

TRACKING EPILEPSY-RELATED GRAY AND WHITE MATTER ATROPHY ACROSS THE ADULT LIFESPAN: A WORLD WIDE ENIGMA STUDY

Judy Chen¹, Alexander Ngo¹, Raúl Rodríguez-Cruces¹, Jessica Royer¹, Maria Eugenia Caligiuri², Antonio Gambardella^{2,3}, Luis Concha⁴, Simon S. Keller^{5,6}, Fernando Cendes⁷, Clarissa L. Yasuda⁷, Leonardo Bonilha⁸, Ezequiel Gleichgerrcht⁸, Niels K. Focke⁹, Martin Domin¹⁰, Felix von Podewils¹¹, Soenke Langner¹², Christian Rummel¹³, Roland Wiest¹³, Pascal Martin¹⁴, Raviteja Kotikalapudi^{14,9}, Terence J. O'Brien^{15,16}, Benjamin Sinclair^{15,16}, Lucy Vivash^{15,16}, Patrick Kwan¹⁶, Patricia M. Desmond¹⁶, Elaine Lui¹⁶, Alice Ballerini¹⁸, Anna Elisabetta Vaudano^{17,18}, Stefano Meletti^{17,18}, Manuela Tondelli^{18,19}, Saud Alhusaini^{20,21}, Colin P. Doherty^{22,23}, Gianpiero L. Cavalleri^{20,23}, Norman Delanty^{20,23}, Reetta Kälviäinen^{24,25}, Graeme D. Jackson²⁶, Magdalena Kowalczyk²⁶, Mario Mascalchi²⁷, Mira Semmelroch²⁶, Rhys H. Thomas²⁸, Hamid Soltanian-Zadeh^{29,30}, Esmaeil Davoodi-Bojd³¹, Junsong Zhang³², Matteo Lenge³³, Renzo Guerrini³³, Emanuele Bartolini³⁵, Khalid Hamandi^{36,37}, Sonya Foley³⁶, Theodor Rüber^{38,39}, Tobias Bauer^{38,39}, Bernd Weber⁴¹, Benoit Caldaïrou⁵⁴, Chantal Depondt⁴², Julie Absil⁴³, Sarah J. A. Carr⁴⁴, Eugenio Abela⁴⁴, Mark P. Richardson⁴⁴, Orrin Devinsky⁴⁵, Mariasavina Severino⁴⁶, Pasquale Striano^{46,47}, Domenico Tortora⁴⁷, Erik Kaestner⁵⁵, Donatello Arienzo⁵⁵, Sjoerd B. Vos^{49,50,51}, Mina Ryten^{49,52}, John S. Duncan^{49,50}, Christopher D. Whelan²⁰, Marian Galovic⁵⁷, Gavin P. Winston^{49,50,58}, Sophia I. Thomopoulos⁵³, Paul M. Thompson⁵³, Sanjay M. Sisodiya^{49,50}, Angelo Labate⁶⁰, Carrie R. McDonald⁵⁵, Lorenzo Caciagli^{49,52}, Neda Bernasconi⁵⁴, Andrea Bernasconi⁵⁴, Sara Larivière^{1,59*}, Dewi Schrader^{56*}, Boris C. Bernhardt^{1*}

¹Multimodal Imaging and Connectome Analysis Laboratory, McConnell Brain Imaging Centre, Montreal Neurological Institute and Hospital, McGill University, Montreal, QC, Canada

²Neuroscience Research Center, University Magna Græcia, Catanzaro, CZ, Italy

³Institute of Neurology, University Magna Græcia, Catanzaro, CZ, Italy

⁴Institute of Neurobiology, Universidad Nacional Autónoma de México, Querétaro, México

⁵Institute of Systems, Molecular and Integrative Biology, University of Liverpool, Liverpool, United Kingdom

⁶Walton Centre NHS Foundation Trust, Liverpool, United Kingdom

⁷Department of Neurology, University of Campinas—UNICAMP, Campinas, São Paulo, Brazil

⁸Department of Neurology, Emory University, Atlanta, GA, USA

⁹Department of Neurology, University of Medicine Göttingen, Göttingen, Germany

¹⁰Institute of Diagnostic Radiology and Neuroradiology, Functional Imaging Unit, University Medicine Greifswald, Greifswald, Germany

¹¹Department of Neurology, University Medicine Greifswald, Epilepsy Center, Greifswald, Germany

¹²Institute of Diagnostic Radiology and Neuroradiology, University Medicine Greifswald, Greifswald, Germany

¹³Support Center for Advanced Neuroimaging (SCAN), University Institute of Diagnostic and Interventional Neuroradiology, University Hospital Bern, Bern, Switzerland

¹⁴Department of Neurology and Epileptology, Hertie Institute for Clinical Brain Research, University of Tübingen, Tübingen, Germany

¹⁵Department of Neuroscience, Central Clinical School, Alfred Hospital, Monash University, Melbourne, Victoria, Australia

¹⁶Departments of Medicine and Radiology, The Royal Melbourne Hospital, The University of Melbourne, Parkville, Victoria, Australia

¹⁷Neurology Unit, OCB Hospital, Azienda Ospedaliera-Universitaria, Modena, Italy

¹⁸Department of Biomedical, Metabolic and Neural Science, University of Modena and Reggio Emilia, Modena, Italy

¹⁹Primary Care Department, Azienda Sanitaria Locale di Modena, Modena, Italy

²⁰Department of Molecular and Cellular Therapeutics, The Royal College of Surgeons in Ireland, Dublin, Ireland

²¹Department of Neurology, Alpert Medical School of Brown University, Providence, RI, USA

²²Department of Neurology, St James' Hospital, Dublin, Ireland

²³FutureNeuro SFI Research Centre, Dublin, Ireland

²⁴Epilepsy Center, Neuro Center, Kuopio University Hospital, Member of the European Reference Network for Rare and Complex Epilepsies EpiCARE, Kuopio, Finland

²⁵Faculty of Health Sciences, School of Medicine, Institute of Clinical Medicine, University of Eastern Finland, Kuopio, Finland

²⁶Florey Institute of Neuroscience and Mental Health, University of Melbourne, Melbourne VIC 3010, Australia

²⁷Neuroradiology Research Program, Meyer Children Hospital of Florence, University of Florence, Florence, Italy

²⁸Transitional and Clinical Research Institute, Newcastle University, Newcastle upon Tyne, UK

²⁹Control and Intelligent Processing Center of Excellence (CIPCE), School of Electrical and Computer Engineering, University of Tehran, Tehran, Iran

³⁰Departments of Research Administration and Radiology, Henry Ford Health System, Detroit, MI, USA

³¹Department of Neurology, Henry Ford Health System, Detroit, MI, USA

³²Cognitive Science Department, Xiamen University, Xiamen, China

³³Child Neurology Unit and Laboratories, Neuroscience Department, Meyer Children's Hospital IRCCS, Florence, Italy

³⁴Functional and Epilepsy Neurosurgery Unit, Neurosurgery Department, Children's Hospital A. Meyer-University of Florence, Italy

³⁵IRCCS Stella Maris Foundation, Pisa, Italy

³⁶Cardiff University Brain Research Imaging Centre (CUBRIC), College of Biomedical and Life Sciences, Cardiff University, Cardiff, UK

³⁷The Welsh Epilepsy Unit, Department of Neurology, University Hospital of Wales, Cardiff, UK

³⁸Department of Neuroradiology, University Hospital Bonn, Bonn, Germany

³⁹Department of Neuroradiology, University Hospital Bonn, Bonn, Germany

40Center for Personalized Translational Epilepsy Research (CePTER), Goethe-University Frankfurt, Frankfurt am Main, Germany
 41Institute of Experimental Epileptology and Cognition Research, University Hospital Bonn, Bonn, Germany
 42Department of Neurology, Hôpital Erasme, Hôpital Universitaire de Bruxelles (HUB), Université Libre de Bruxelles, Brussels, Belgium
 43Department of Radiology, Hôpital Erasme, Université Libre de Bruxelles, Brussels, Belgium
 44School of Neuroscience, Institute of Psychiatry, Psychology and Neuroscience, King's College London, UK
 45Department of Neurology, NYU Grossman School of Medicine, New York, NY, United States
 46IRCCS Istituto Giannina Gaslini, Genova, Italy
 47Department of Neurosciences, Rehabilitation, Ophthalmology, Genetics, Maternal and Child Health, University of Genova, Genova, Italy
 48Department of Neurosciences, Center for Multimodal Imaging and Genetics, University of California San Diego, La Jolla, CA, United States
 49Department of Clinical and Experimental Epilepsy, UCL Queen Square Institute of Neurology, London, UK
 50Chalfont Centre for Epilepsy, Bucks, UK
 51Centre for Medical Image Computing, University College London, London, UK
 52MRI Unit, Epilepsy Society, Chalfont St Peter, Bucks, UK
 53Imaging Genetics Center, Mark & Mary Stevens Institute for Neuroimaging and Informatics, Keck School of Medicine, University of Southern California, Los Angeles, CA, United States
 54Neuroimaging of Epilepsy Laboratory, McConnell Brain Imaging Centre, Montreal Neurological Institute and Hospital, McGill University, Montreal, QC, Canada
 55Department of Radiation Medicine and Applied Sciences, Department of Psychiatry, Center for Multimodal Imaging and Genetics, University of California San Diego, La Jolla, CA, United States
 56BC Children's Hospital, University of British Columbia, Vancouver, BC, Canada
 57Department of Neurology, Clinical Neuroscience Center, University Hospital and University of Zürich, Zürich, Switzerland
 58Department of Medicine, Division of Neurology, Queen's University, Kingston, ON, Canada
 59Center for Brain Circuit Therapeutics, Brigham and Women's Hospital, Harvard University, Boston, MA, USA
 59Neurophysiopathology and Movement Disorders Clinic, Regional Epilepsy Center, University of Messina, Italy

Running head: TBD

**joint senior authors*

Correspondence to:

Boris C. Bernhardt, PhD
 Montreal Neurological Institute (NW-254)
 3801 University Street
 Montreal, Quebec, Canada H3A 2B4
 p: (514) 398-3579
 e: boris.bernhardt@mcgill.ca

Judy Chen, PhD student
 e: judy.chen@mail.mcgill.ca

ABSTRACT

Objectives. Temporal lobe epilepsy (TLE) is commonly associated with mesiotemporal pathology and widespread alterations of grey and white matter structures. Evidence supports a progressive condition although the temporal evolution of TLE is poorly defined. This ENIGMA-Epilepsy study utilized multimodal magnetic resonance imaging (MRI) data to investigate structural alterations in TLE patients across the adult lifespan. We charted both grey and white matter changes and explored the covariance of age-related alterations in both compartments.

Methods. We studied 769 TLE patients and 885 healthy controls across an age range of 17-73 years, from multiple sites. To assess potentially non-linear lifespan changes in TLE, we harmonized data and combined median split assessments with cross-sectional sliding window analyses of grey and white matter age-related changes. Covariance analyses examined the coupling of grey and white matter lifespan curves.

Results. In TLE, age was associated with a robust gray matter thickness/volume decline across a broad cortico-subcortical territory, extending beyond the mesiotemporal disease epicentre. White matter changes were also widespread across multiple tracts with peak effects in temporo-limbic fibers. While changes spanned the adult time window, changes accelerated in cortical thickness, subcortical volume, and fractional anisotropy (all decreased), and mean diffusivity (increased) after age 55 years. Covariance analyses revealed strong limbic associations between white matter tracts and subcortical structures with cortical regions.

Conclusions. This study highlights the profound impact of TLE on lifespan changes in gray and white matter structures, with an acceleration of aging-related processes in later decades of life. Our findings underscore the importance of targeted longitudinal studies focusing on epilepsy-related effects in this age group.

Keywords. Epilepsy, Temporal Lobe Epilepsy, aging, lifespan, atrophy, multimodal

INTRODUCTION

Epilepsy is a common and debilitating neurological disease affecting over 50 million individuals worldwide, of which a substantial proportion suffers from functional deficits and markedly compromised quality of life.¹ Temporal lobe epilepsy (TLE) is the most common pharmaco-resistant epilepsy in adults associated with mesiotemporal pathology.² While traditionally considered a prototypical focal epilepsy, increasing evidence suggests that atrophy and white matter alterations are not limited to the mesiotemporal region, but affect widely distributed grey and white matter systems. Single and multi-site morphometric studies have reported significantly reduced grey matter volume in temporal, frontal, and centroparietal cortices as well as subcortical structures such as the thalamus and amygdala.^{3–5} Moreover, there is a compromise of white matter microstructure and architecture of numerous fiber tracts, with larger effect sizes in limbic and subcortico-cortical systems.⁶ Alterations have been shown to follow structural connectivity patterns between distributed regions and the mesiotemporal epicentre, and is evident in large-scale inter-regional morphometry.^{7,8} As such, TLE is now increasingly recognized as a “network” disorder accompanied by profound changes in cortico-subcortical grey matter morphology and changes in related white matter compartments.^{9–12}

There is evidence that TLE is not a static condition, but shows progression over time. Advanced age is a critical risk factor for cognitive decline, particularly in TLE where the majority of patients older than 55 years have been found to present with memory and language deficits that may meet the criteria for mild cognitive impairment.¹³ In addition to cognitive decline, older TLE patients exhibit accelerated bilateral cortical thinning beyond the mesiotemporal region.^{14,15} A recent meta-analysis of structural magnetic resonance imaging (MRI) studies suggested that TLE is associated with cumulative atrophy exceeding typical aging-related processes, but most constituent studies were limited by the low quality of evidence and heterogeneous approach to controlling for age-related effects.¹⁶ Even fewer studies have investigated progression of effects on white matter despite the role of this compartment in seizure propagation. Despite some evidence for age as a key determinant of global white matter volume changes, aging-related effects on white matter microstructure remain underexplored.¹⁷

Thus, whilst these prior investigation provide some initial evidence for age-modulated changes in brain structure in TLE, findings are limited in several ways. First, most research has focused on gray matter morphology,¹⁶ not addressing white matter changes nor the covariance between grey and white matter alterations. Such studies would be needed to understand aging effects on brain-wide integrity and to identify pathology affecting white and grey matter in epilepsy. Second, research into aging effects in TLE has been limited by focusing on only small cohorts and has been mainly restricted to single sites.^{15,18,19} Such an approach may be limited in terms of reproducibility and generalizability and affected by cohort- and centre-specific idiosyncrasies.³ Third, age is generally considered a confounder for the more prevailing analyses of disease duration or between-group difference effects, and has therefore not been a key focus in assessing its direct relation to MRI measures in TLE in prior work.¹⁶ Here, we address these gaps by leveraging a multi-site TLE cohort aggregated by the ENIGMA-Epilepsy Consortium to map lifespan changes in multimodal MRI measures of grey matter morphology and white matter microstructure.^{4,6} In addition to assessing grey and white matter changes independently, we examined their covariance to quantify the inter-regional and cross-compartment coupling of aging-related processes.

METHODS

Participants. We studied 769 TLE patients (455 females, mean $\pm$ standard deviation [SD] age = 38.2 ± 11.1 years, 430/339 with left/right-sided seizure focus, age of onset = 16.5 ± 12.0 years, disease duration = 21.3 ± 13.5 years) and 885 age- and sex-matched healthy controls (508 females, mean $\pm$ SD age = 36.0 ± 11.7 years) spanning an age range of 17-73 years from the ENIGMA-Epilepsy cohort. Summary clinical and demographic characteristics are presented in **Table S1**. Engel scores of all patients that had undergone surgery ($n=769$) are reported as well. TLE was diagnosed according to standardized International League Against Epilepsy criteria (REF) across 18 recruitment sites.

Multimodal MRI data processing and harmonization. Acquisition and scanner descriptions of both the structural T1-weighted (T1w) MRI and diffusion-tensor imaging (DTI) for neocortical and subcortical regions obtained from all participants are described in previous ENIGMA-Epilepsy studies.^{4,6} Images were independently processed on site using a standardized workflow (T1w: <https://enigma.ini.usc.edu/protocols/imaging-protocols/>, DTI: <https://enigma.ini.usc.edu/protocols/dti-protocols/>, <https://enigma.ini.usc.edu/ongoing/dti-working-group/>)³, which allowed for the measurement of cortical thickness (CT) and subcortical volume (SV) based on T1w according to the Desikan-Killiany atlas, composed of 68 cortical and 16 subcortical regions.²⁰ Diffusion MRI data were processed using TBSS to derive the tensor parameters fractional anisotropy (FA) and mean diffusivity (MD) for 38 unique white-matter tracts referenced from the JHU ICBM-DTI-81 atlas.²¹

Statistical analysis. Analyses were performed in Matlab R2019b (The Mathworks, Natick, MA). To control for potential confounders such as scanner and site variability but to preserve biological covariates (age, sex, and diagnostic group), harmonization via ComBat was performed for DTI and T1w data.^{22,23} Patient measures were z-scored relative to controls and sorted into ipsilateral/contralateral to the seizure focus as per previous studies.^{7,24} We then used the openly accessible BrainStat toolbox for statistical inference and data visualization (<http://brainstat.readthedocs.io>).²⁵

a) Age-bisected cohort analysis. To preserve statistical power of this large dataset, while establishing age-dependent differences between TLE patients and controls, the cohort was first split via its median age across the entire sample (35 years), which also approximates to the median in TLE (37 years) and control (33 years) groups. This resulted in a “young” (< 35 years) and “older” (≥ 35 years) cohort for both patients (n young/old=292/477) and healthy controls ($n=481/405$). Linear models highlighted CT, SV, FA, and MD effect size (Cohen’s d) profiles of young patients as compared to their age-equivalent healthy control cohort, and likewise of old patients and their age-equivalent control cohort. This illustrated the progression of differences between the two age groups. Group-by-age interactions analysis identified regions showing more marked age-dependent changes in TLE patients compared to controls, with participant ages treated categorically to identify only regions with the most robust age-dependent changes and to align with the prior median split analyses. All findings controlled for sex and were corrected for multiple comparisons using the false discovery rate (FDR) procedure.²⁶

b) Sliding age-window and covariance analysis. Age-sliding windows were utilized to stabilize trends and limit the influence of potential outliers (REF). Though various window ranges can be constructed, a range of ± 2 years from an age of interest (AOI) was chosen for presented analyses to best maintain granularity of trajectory perturbations while maximizing the number of patients. Sensitivity analyses were conducted to ensure preservation of trends across different age windows (**Fig. S3**). AOI and subsequent windows begin at 19 and end at 70 years of age; each window is slid incrementally by 1 year, resulting in 52 unique sub-cohorts. For each sub-cohort, structural/microstructural measures were calculated via normally distributed weights based on their age distance to the AOI. Finally, each sliding age-window was z-scored against the healthy control cohort to yield age-associated z-score trajectories across all cortical, subcortical, and white matter tract regions (**Fig. 1B**). Correlations of each region-associated age trajectory across the lifespan for

CT, SV, FA, and MD were also calculated, with the top 10% and bottom 10% highlighted to identify region(s) for which their lifespan trajectories are most closely related.

RESULTS

Gray matter morphometric lifespan analysis

Median split analysis showed an increased rate of ipsilateral reductions in CT and SV in TLE patients over adulthood as compared to changes seen in healthy individuals (**Fig. 1A**). Young TLE patients exhibited bilateral neocortical thinning in the precentral ($d = -0.22$), postcentral ($d = -0.28$), superior parietal ($d = -0.27$), and middle frontal ($d = -0.27$) regions compared to their age-matched healthy counterparts ($p_{FDR} < 0.05$). In the ipsilateral hemisphere, there was additional atrophy in the transverse temporal areas ($d = -0.20$, $p_{FDR} < 0.05$). The differences between patients and controls in the older TLE cohort showed almost identical overlap for atrophic regions relative to the results of the younger cohorts, but with further atrophy of the neocortex, particularly in the ipsilateral hemisphere. Additional areas of atrophy in the older age group included ipsilateral entorhinal ($d = -0.26$, $p_{FDR} < 0.0005$) and parahippocampal regions ($d = -0.23$, $p_{FDR} < 0.005$), as well as further bilateral inferior parietal (d ipsi/contra = $-0.20/-0.18$, $p_{FDR} < 0.05$) and contralateral occipital ($d = -0.23$, $p_{FDR} < 0.005$) regions. While the contralateral hemisphere presented with fewer significant regions, atrophy overall still included a similar pattern as seen in the younger TLE cohort with additional effects in bilateral frontal (superior frontal d ipsi/contra = $-0.23/-0.21$, caudal middle frontal (d ipsi/contra = $-0.24/-0.24$, $p_{FDR} < 0.05$) thinning. In total, 18 (d ipsi/contra = 11/7) cortical regions of interest (ROI) significantly differed in the young TLE cohort, while 26 (d ipsi/contra = 16/10) ROIs significantly differed in the older TLE cohort. Overall, the group-by-age interaction analysis showed that neocortical thinning in the ipsilateral superior parietal, contralateral precuneus, cuneus, and pericalcarine regions was significantly more marked in older patients than younger patients in comparison to their healthy control counterparts ($p_{FDR} < 0.05$).




A | Gray matter analysis of age-bisected cohort**B | Construction of sliding-age windows (gray matter)****C | TLE cortical thickness and subcortical volume sliding-age window trajectories**

Figure 1. Gray matter morphometry via median-split analysis and sliding-age window trajectories in temporal lobe epilepsy (TLE). (A) Median age of 35 was chosen as the division into young and old cohorts. All right-sided temporal lobe epilepsy (TLE) patients were flipped, with results demonstrating significant left hippocampal reductions in both young and old age cohorts, and increasing occipital gray matter atrophy with age as seen in the old cohort. There are strong group by age interaction with significant subcortical regions and some cortical involvement in the parieto-occipital regions. (B) After z-scoring cortical thickness and subcortical volume measures, windows of ± 2 years of the age of interest were chosen, and the mean values for cortical (68 total brain regions) and subcortical regions (12 subcortical gray matter regions and bilateral ventricles and hippocampi) following the Desikan-Killiany atlas were multiplied by normally distributed weights to yield a weighted average. Global weighted-average z-scores were then plotted across age to illustrate lifespan trajectory. (C) In temporal lobe epilepsy (TLE) patients, older age especially past age 55 seems to be correlated with an accelerated and pronounced decline in cortical thickness. The hippocampi and other subcortical regions also show a marked decreased volume with aging.

There were significant ipsilateral hippocampal volume reductions in both young and older TLE cohorts compared to their healthy age counterparts ($d = -0.60$, $p_{FDR} < 0.05e-15$). Young TLE cohorts exhibited

bilateral volume reductions in the thalamus (d ipsi/contra=-0.41/-0.29, $p_{FDR}<0.05$) and pallidum (d ipsi/contra=-0.17/-0.19, $p_{FDR}<0.05$). Older TLE cohorts showed more extensive, bilateral findings ($p_{FDR}<0.05$), with greater volume reductions in the hippocampi (d ipsi/contra=-0.82/-0.67), thalamus (d =-0.29/-0.68), caudate (d =-0.41/-0.31), and ipsilateral amygdala (d =-0.26).

To discern trajectories of neocortical thinning over the lifespan (**Fig. 1B-C**), age-sliding windows were constructed, and cortical thickness was z-scored against controls for each region. Overall, there was a slow but progressive decline in mean cortical thickness from young adulthood to middle age, with mean z-scores decreasing to -1 by age 55 and dropping more steeply to -4 beyond age 60; this pattern was reflected across all cortical regions bilaterally. In contrast, subcortical volumes decreased linearly and less severely (ranging from z-scores of 0 to -2) as age advanced. This pattern remained consistent across all subcortical regions and the hippocampus. In particular, the volume of the ipsilateral hippocampus exhibited the most consistent decline, with substantially decreased z-scores compared to other cortical and subcortical regions. Furthermore, comparative sensitivity analyses of these age-dependent trajectories demonstrated much higher correlation with the TLE-mesial temporal sclerosis (MTS) sub-cohort (CT/SV $r=0.88/0.91$) compared to the TLE-non lesional (NL) sub-cohort (CT/SV $r=0.68/0.70$). There were little differences between the correlations of the left-sided TLE (CT/SV $r=0.77/0.83$) and right-sided TLE (CT/SV $r=0.82/0.86$) sub-cohorts.

White matter microstructure lifespan analysis

Within the age-bisected cohort, young TLE patients exhibited smaller FA decreases compared to older TLE patients against their respective age-matched control cohort with a total of 19/21 significantly different ROIs ($p_{FDR}<0.05$) within the younger TLE cohort, and 21/21 significantly different ROIs ($p_{FDR}<0.05$) within the older TLE cohort. Most white matter tracts within the younger TLE cohort were significantly decreased, particularly in the fornix (FX: $d=-0.37$; fornix stria terminalis [FXST]: d ipsi/contra=-0.29/-0.33 $p_{FDR}<0.05$), genu (GCC: -0.41, $p_{FDR}<0.05$) and body of the corpus callosum (BCC: -0.43, $p_{FDR}<0.05$), bilateral external capsule (EC: d ipsi/contra=-0.45/-0.53, $p_{FDR}<0.05$), and cingulum (cingulate gyrus [CGC]: d ipsi/contra=-0.42/-0.47 ; cingulum adjacent to the hippocampus [CGH]: d ipsi/contra=-0.3/-0.42, $p_{FDR}<0.05$). All white matter tracts were significantly decreased in older TLE patients with even greater decline in FA values within the fornix (FX: $d = -0.59$, $p_{FDR}<0.05$; FXST: d ipsi/contra=-0.45/-0.47, $p_{FDR}<0.05$), genu (GCC: $d=-0.61$, $p_{FDR}<0.05$) and body (BCC: $d=-0.72$, $p_{FDR}<0.05$) of the corpus callosum, and cingulum (CGC: d ipsi/contra=-0.60/-0.61; CGH d ipsi/contra=-0.54/-0.57; $p_{FDR}<0.05$). Overall, there was a clear decrease in FA values between older cohorts and younger cohorts, with significant age-effects across almost all white matter regions, suggesting a weakening of structural integrity of the fibre tracts and cumulative damage with age.

For MD, we observed a low to moderate increase in most tracts in young TLE cohorts when compared to their age-matched controls, totalling 14/21 bilaterally altered ROIs ($p_{FDR}<0.05$). The fornix ($d=0.41$, $p_{FDR}<0.05$), bilateral external capsule (EC: d ipsi/contra=0.31/0.32, $p_{FDR}<0.05$), cingulum adjacent to the hippocampus (CGH: d ipsi/contra=0.33/0.30, $p_{FDR}<0.05$), and the anterior and posterior corona radiata (ACR: d ipsi/contra=0.25/0.23; PCR: d ipsi/contra=0.27/0.26, $p_{FDR}<0.05$) had the highest MD increases within the young TLE cohort. The ipsilateral corticospinal tract was the sole structure that had an unexpected decrease in MD ($d=-0.65$, $p_{FDR}<0.05$). In the older TLE cohort, almost all (20/21 ROIs) white matter structures demonstrated larger increases as compared to the differences seen in the younger TLE cohort ($p_{FDR}<0.05$), with the greatest increases in the corpus callosum (GCC: $d=0.50$; BCC: $d=0.65$, $p_{FDR}<0.05$), corona radiata (ACR: d ipsi/contra=0.37/0.42; SCR: d ipsi/contra = 0.41/0.42; PCR d ipsi/contra=0.46/0.45, $p_{FDR}<0.05$), cingulum adjacent to the hippocampus (CGH: d ipsi/contra=0.40/0.36, $p_{FDR}<0.05$), and the superior fronto-occipital fasciculus (SFO: d ipsi/contra=0.31/0.43, $p_{FDR}<0.05$). The fornix was the only structure that exhibited a higher MD effect size within the younger TLE cohort ($d=0.41$, $p_{FDR}<0.05$) than in the older TLE cohort ($d=0.27$, $p_{FDR}<0.05$). Apart from the fornix, all other regions

exhibited significantly increased effect for the older TLE cohort compared to their age-matched controls. Group-by-age interaction analyses for both FA and MD of the white matter structures showed almost all regions were significantly more increased for MD and decreased for FA in patients than their healthy control counterparts ($p_{\text{FDR}} < 0.05$).

The trajectories of FA measures across age (**Fig. 2**) resembled gray matter trajectories, with a gradual downward trend until age 50, followed by a steeper decline. This general trend occurs across most white matter tracts, such as the limbic, cortical-subcortical, and cortical structures (see Table S2 for a complete list). The splenium, body, and genu of the corpus callosum white matter tracts were more pronounced in their FA decreases. Likewise, MD age trajectories for all white matter tracts gradually increased until age 60, after which MD values showed even greater increases. Most pronounced findings were with structures associated with the limbic, cortical-subcortical, and cortical regions bilaterally. The correlation of the age trajectories with the TLE-MTS subcohort again were again much higher (FA/MD $r=0.86/0.78$) compared to their TLE-NL sub-cohort counterparts (FA/MD $r=0.57/0.50$). Additional correlations calculated between the FA/MD age trajectories showed a much more positive correlation with the right-sided TLE patient cohort (FA/MD $r=0.91/0.86$) as opposed to the left-sided patient cohort (FA/MD $r=0.68/0.75$).




Figure 2. White matter alterations via median-split analysis and sliding-age window trajectories in temporal lobe epilepsy (TLE). (A) Median age of 35 was chosen as the division into young and old cohorts. All right-sided temporal lobe epilepsy (TLE) patients were flipped, with results demonstrating almost all white matter tracts significantly decreased for fractional anisotropy (FA) measures and increased for mean diffusivity (MD) measures for both young and old TLE cohorts against their age-matched controls. (B) Average age trajectories were calculated for ipsilateral and contralateral white matter tracts associated with the corpus callosum, limbic, cortical-subcortical, and cortical regions. All tracts demonstrate an accelerated and pronounced decline in FA measures past age 50 and a similarly pronounced increase in MD measures past age 60.




Figure 3. Sliding age-window structural covariance correlations in temporal lobe epilepsy (TLE). Correlograms (center) showing the correlational of each region-associated age trajectory across the lifespan for CT, SVC, FA, and MD were calculated. The top 10% (*red*) and bottom 10% (*blue*) of connections were preserved to identify regions more closely associated with each gray or white matter measures. Scatter plots demonstrate the age-dependent (*gray*) distribution of z-score values against controls.

Age-mediated structural covariance dynamics

Aside from region-specific correlations to itself, there is a general positive correlation between FA, SV, and CT measures, and a general negative correlation between MD, SV, and CT measures (**F3**). This is further illustrated by the scatter plots of every patient and their z-scores against controls for gray and white matter measures. The top 10% of correlations were calculated for FA by CT, FA by SV, and CT by SV correlograms, while the bottom 10% of correlations were calculated for MD by CT, MD by SV, and MD by FA to reflect their general trend. In the FA by CT correlogram, more than half of FA structures associated with the right caudal anterior cingulate neocortical region belong in the top 10%. Likewise, more than half of all CT structures correlated with the fornix cres/stria terminalis (FXST), ipsilateral and contralateral posterior thalamic radiation (PTR) were found in the top 10%. The ipsilateral posterior thalamic (PTR) radiation was again implicated in the FA by SV correlogram. Finally, in the SV by CT correlogram, the right pallidum had almost all its correlations with the neocortical regions within the top 10% of connections. Similar analyses done within the MD by CT, MD by SV, and MD by FA implicate the MD value of contralateral corticospinal tract (CST) for both FA white matter regions and subcortical volumes. Both ipsilateral and contralateral FXST were also strongly associated across most neocortical and subcortical regions. Covariance patterns between the ipsilateral/ipsilateral, ipsilateral/contralateral, and contralateral/contralateral structures showed concordant trends across hemispheres (**Fig. S4**).

DISCUSSION

Our multi-site cross-sectional study investigated aging trajectories and the correlative interplay of gray and white matter atrophy in pharmaco-resistant TLE patients. Epilepsy exerts an age-dependent effect across the adult lifespan, and an acceleration of aging-related effects in later decades. Benefiting from a large sample size, multiparametric assessment of gray and white matter alterations, and standardized protocols for data extraction and processing from all ENIGMA-Epilepsy sites, our findings illustrate TLE as a temporo-limbic network disorder exhibiting age-mediated decline of gray/white matter structural integrity that accelerates with advanced age (>55 years). Collectively, our findings highlight (i) the influence age exerts on disease effects and (ii) that patients above 55 years of age are more severely affected.

Gray matter atrophy not only became significantly more pronounced in older patients but also extended over a broader anatomical territory with advancing age. Prior literature on cortical thickness, hippocampal volumes, and subcortical structures supports a pattern of diffuse whole brain atrophy in TLE^{4,7,14,16,18} echoing our findings of reductions in the entorhinal, medial parietal, cingulate, and frontal cortices as well as hippocampal and thalamic volumes. Other cross-sectional cohort studies from ENIGMA-Epilepsy have also highlighted broad cortical atrophy and subcortical volume reductions in TLE patients on a group level.^{4,7} However, little has been done to investigate the influence of age on such alterations. Previous evaluations of cortical thinning across the lifespan in healthy individuals have revealed unique patterns of decline specific to neocortical regions; here, entorhinal and temporopolar cortices seemed to increase initially following an inverse U-shape pattern, unlike most other regions for which cortical thickness is highest during childhood.²⁷ Our findings of linearly progressive thinning, particularly for the entorhinal and temporopolar cortices, is in line with cumulative disease effects and complement previous reports of disease progression in epilepsy.¹⁶ Though such preceding studies have assessed disease duration in relation to atrophy to evaluate progression in TLE patients, many either report no correlations between age and gray matter atrophy in healthy control and patient populations despite compelling evidence for declines in normal aging in adulthood,^{14,15,19,28} or identify age as a likely determinant on atrophy rate and volume loss.¹⁷ A meta-analysis reviewing progressive atrophy in TLE has shown that the majority of reports are cross-sectional with narrow age-ranges, small cohort sizes, and consequently, poor age control.¹⁶ Here, direct comparisons of age to atrophy metrics demonstrated age as having increasing influence through later years and may itself be a key biomarker of disease effects. The increased rate of cortical thinning and subcortical volume reductions of TLE patients older than 50 years is similar to previous longitudinal studies demonstrating increased progression and severity of cortical thinning in TLE cohorts,^{14,15} suggestive of an increased vulnerability to cortical damage and potential accelerated age-mediated decline.

In line with prior investigations showing white matter abnormalities in TLE,^{6,29–31} almost all white matter tracts were significantly altered in our young and old TLE cohorts compared to their respective counterparts controls at the group level. Magnitudes of MD increase and FA reduction were highest bilaterally in the fornix, cingulum, and corpus callosum. The alteration of these tracts adds to prior evidence for marked changes of temporo-limbic tracts, given their close associations anatomically and functionally to the mesio-temporal TLE epicentre and thus further emphasizing the network level impact of epilepsy.³² The accelerated structural changes in temporo-limbic tracts provide evidence of secondary damage from the disease, consistent with previous studies on adult TLE that associated MD decreases with age of onset in early-onset TLE patients and seizure frequency in late-onset TLE patients, with elevated MD values primarily in the uncinate and ipsilateral hemisphere.³³ The accelerated microstructural damage of almost all white matter fascicles suggests that whole-scale white matter perturbations beyond atrophy of these lesion-associated anatomical networks can occur, particularly in older TLE patients. Indeed, baseline age has been found as the main determinant of cerebral white matter volume decreases when assessing disease progression.¹⁷ The relationship between white matter atrophy and aging remains inadequately explored; most earlier studies on group level differences or spatial characteristics indicate no correlation between age and white matter metrics as they possess limited cohort sizes of age-matched controls,^{31,34} entirely omit age analyses,³⁵ or report significant associations of duration and age of onset to diffusion metrics after correcting

for age.^{6,29} Simply correcting for age, however, does not rule out potential for age-mediated effects on disease duration or age of onset, as older patients may still show accelerated loss of gray and white matter integrity compared to a younger counterpart despite having been diagnosed more recently. Given the highly correlative characteristics of age, age of onset, and disease duration, future work towards assessing outcomes involving these variables should be prudent to employ methodologies that ensure nuanced insight.

Our evaluation of the concomitant covariance patterns of white and gray matter microstructural change over the lifespan implicate the fornix, posterior thalamic radiation, corticospinal tracts, pallidum, and thalamus as structures most strongly correlated to the global aging trajectories of at least one other metric (CT, SV, FA, or MD), reinforcing the notion of TLE as a limbic network disorder.^{34,36} This aligns with hypothesized axonal degeneration within limbic pathways along the fornix, posterior thalamic radiation, and thalamic structures,^{31,36} and further suggest an interaction between age and epilepsy in network pathology. The functional overlap of these limbic structures to memory formation, specifically the lifespan trajectories of the entorhinal cortex and FA/MD metrics of the fornix, may explain the increased cognitive decline observed in many older individuals with TLE. Even less is known regarding networks associated with the pallidum and corticospinal tracts, though we speculate a relationship to motor pathway activation during seizures.³⁷ More information regarding cognitive, electro-clinical, and socio-demographic data in these participants is needed to further ascertain these relationships in future work, ideally based on prospective and longitudinal investigations.

Continued seizures represent long-term, repeated insults to the brain that have been associated with progressive gray and white matter atrophy.^{38,39} Recent findings also suggest a bidirectional relationship mediated by common vascular risk factors, molecular profiles, and large-scale network changes between epilepsy and vascular-related dementia.^{40–42} One could thus consider age to be the largest risk factor for these neurodegenerative diseases, an equally independent clinical marker of vulnerability to seizures and epilepsy in older patient cohorts. Psychosocial difficulties are also exacerbated in older epilepsy patients, who are increasingly at risk of a psychiatric disorder.⁴³ Furthermore, there have been reports suggesting an association between structural compromise and antiseizure medication which could modulate age-related effects seen here.⁴⁴ The multiple factors likely coalesce to explain the accelerated atrophy we see in older TLE cohorts, but the cross-sectional nature of this present study in evaluating these aging effects lends susceptibility to confounders such as cohort effects and thus prevents establishing causal associations. A significant proportion of our cohort had paediatric age of onset, therefore more comprehensive lifespan analyses would begin around the mean age of onset to fully illustrate gray and white matter trajectories. Stratified sampling and increased recruitment of older age populations given our limited cohort size past age 55 would reinforce the argument for accelerated disease processes linked to aging as opposed to cohort effects. Our analyses did not correct for disease duration as doing so in a cross-sectional study may fail to show discernable differences.¹⁶

Our findings indicate that age, whether acting as a surrogate for disease duration or used to argue for disease progression, is an overlooked yet clinically relevant biomarker in TLE, particularly in patients above the age of 55. The heterogeneity of older patients in educational attainment, socioeconomic status, or other factors that confer protective effects on cortical thinning and cognitive outcomes may necessitate an individualized approach and more comprehensive data to evaluate the true risk baseline age poses for accelerated brain atrophy and potential worsening of symptoms. In this context, future longitudinal work with stratified sampling across the lifespan is recommended to explore eEpilepsy-specific characteristics such as frequency of seizures, alongside with additional contributions of neurological, neurovascular, and psychiatric comorbidities. Longitudinal designs may also help to between address cohort-level health effects such as differences in drug exposure, shifts in treatment, and world-scale events. Ultimately, future work should be directed towards patients-specific longitudinal tracking of network-level trajectories to fully comprehend the extent of disease progression, its potential acceleration in older age, and associations to functionally and clinically relevant outcomes.

ACKNOWLEDGEMENTS

R.R.C. is funded by the Fonds de la Recherche du Québec – Santé (FRQS). L.C. is supported by the Berkeley Fellowship jointly awarded by UCL and Gonville and Caius College, Cambridge, and by Brain Research UK (award 14181). A.B. and N.B. are supported by FRQS and CIHR. B.C.B. acknowledges research support from the NSERC Discovery-1304413, CIHR (FDN-154298, PJT-174995), SickKids Foundation (NI17-039), the Helmholtz International BigBrain Analytics and Learning Laboratory (HIBALL), HBHL, Brain Canada, and the Tier-2 Canada Research Chairs program. G.P.W. acknowledges support from the MRC (G0802012, MR/M00841X/1). C.R.M. acknowledges research support from the NIH/NINDS (R01 NS124585; R01 NS122827). P.S. acknowledges support from the PNRR-MUR-M4C2 PE0000006 Research Program “MNESYS”. IRCCS ‘G. Gaslini’ is a member of ERN-Epicare. S.L. acknowledges funding from the Canadian Institutes of Health Research (CIHR).

SUPPLEMENTARY

Table S1 | Detailed clinico-demographic data of ENIGMA-Epilepsy TLE cohort

Case-control cohort	Age (years)	Sex (F/M)	Age of onset (years)	Disease duration (years)	Engel score (I/II+)
HC (<i>n</i> = 886)	36.0 ± 11.7	508/378	—	—	—
L TLE MTS (<i>n</i> =280)	39.0 ± 10.8	161/119	14.9 ± 11.8*	23.8 ± 13.5*	102/178
R TLE MTS (<i>n</i> =226)	39.8 ± 10.9	130/96	15.1 ± 11.3*	24.4 ± 13.8*	66/160
L TLE non-lesional (<i>n</i> = 150)	36.0 ± 10.5	93/133	19.4 ± 11.4*	16.8 ± 12.0*	23/127
R TLE non-lesional (<i>n</i> =113)	36.5 ± 11.5	71/42	19.9 ± 13.4*	14.8 ± 10.9*	25/88

Age, age of onset, and disease duration are presented in mean ± SD. *Clinical information available: a) age of onset: L-TLE MTS = 8/280, R-TLE MTS = 11/130, L-TLE non-lesional = 8/150, R-TLE non-lesional 7/113; b) disease duration: L-TLE MTS = 32/280, R-TLE MTS = 24/130, L-TLE non-lesional 16/150, R-TLE non-lesional 18/113.

TABLE S2 | White matter tract labels

Group	Tract abbreviation	Full name
Corpus Callosum	SCC	Splenium of the corpus callosum
	BCC	Body of the corpus callosum
	GCC	Genu of the corpus callosum
Limbic	FX	Fornix (column and body)
	FXST	Fornix (cres) / Stria terminalis
	CGH	Cingulum (hippocampus)
	CCG	Cingulum (cingulate gyrus)
	UNC	Uncinate fasciculus
Cortical-subcortical	ALIC	Anterior limb of the internal capsule
	PLIC	Posterior limb of the internal capsule
	RLIC	Retrolenticular part of the internal capsule
	ACR	Anterior corona radiata
	SCR	Superior corona radiata
	PCR	Posterior corona radiata
	PTR	Posterior thalamic radiation
	EC	External capsule
	CST	Corticospinal tract
Cortical	SS	Sagittal stratum
	SLF	Superior longitudinal fasciculus
	SFO	Superior fronto-occipital fasciculus
	IFO	Inferior fronto-occipital fasciculus

FIGURE S3 | Sensitivity analyses with age-sliding window ranges





FIGURE S4 | Covariance of lifespan trajectories separated by ipsilateral and contralateral structures and regions



Bibliography

1. Feigin VL, Nichols E, Alam T, et al. Global, regional, and national burden of neurological disorders, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet Neurol.* Elsevier BV; 2019;18:459–480.
2. Wiebe S, Blume WT, Girvin JP, Eliasziw M. A randomized, controlled trial of surgery for temporal-lobe epilepsy. *N Engl J Med.* Massachusetts Medical Society; 2001;345:311–318.
3. Sisodiya SM, Whelan CD, Hatton SN, et al. The ENIGMA-Epilepsy working group: Mapping disease from large data sets. *Hum Brain Mapp.* Wiley; 2020;43:113–128.
4. Whelan CD, Altmann A, Botía JA, et al. Structural brain abnormalities in the common epilepsies assessed in a worldwide ENIGMA study. *Brain.* Oxford University Press (OUP); 2018;141:391–408.
5. Park B-Y, Larivière S, Rodríguez-Cruces R, et al. Topographic divergence of atypical cortical asymmetry and atrophy patterns in temporal lobe epilepsy. *Brain.* Oxford University Press (OUP); 2022;145:1285–1298.
6. Hatton SN, Huynh KH, Bonilha L, et al. White matter abnormalities across different epilepsy syndromes in adults: an ENIGMA-Epilepsy study. *Brain.* Oxford University Press (OUP); 2020;143:2454–2473.
7. Larivière S, Rodríguez-Cruces R, Royer J, et al. Network-based atrophy modeling in the common epilepsies: A worldwide ENIGMA study. *Sci Adv.* American Association for the Advancement of Science (AAAS); 2020;6:eabc6457.
8. Larivière S, Royer J, Rodríguez-Cruces R, et al. Structural network alterations in focal and generalized epilepsy assessed in a worldwide ENIGMA study follow axes of epilepsy risk gene expression. *Nat Commun.* Springer Science and Business Media LLC; 2022;13:4320.
9. Bernhardt BC, Bonilha L, Gross DW. Network analysis for a network disorder: The emerging role of graph theory in the study of epilepsy. *Epilepsy Behav.* Elsevier BV; 2015;50:162–170.
10. Tavakol S, Royer J, Lowe AJ, et al. Neuroimaging and connectomics of drug-resistant epilepsy at multiple scales: From focal lesions to macroscale networks. *Epilepsia.* Wiley; 2019;60:593–604.
11. Gleichgerricht E, Kocher M, Bonilha L. Connectomics and graph theory analyses: Novel insights into network abnormalities in epilepsy. *Epilepsia.* Wiley; 2015;56:1660–1668.
12. Royer J, Bernhardt BC, Larivière S, et al. Epilepsy and brain network hubs. *Epilepsia.* Wiley; 2022;63:537–550.
13. Reyes A, Kaestner E, Edmonds EC, et al. Diagnosing cognitive disorders in older adults with epilepsy. *Epilepsia.* Wiley; 2021;62:460–471.
14. Galovic M, van Dooren VQH, Postma TS, et al. Progressive cortical thinning in patients with focal epilepsy. *JAMA Neurol.* American Medical Association (AMA); 2019;76:1230–1239.
15. Bernhardt BC, Worsley KJ, Kim H, Evans AC, Bernasconi A, Bernasconi N. Longitudinal and cross-sectional analysis of atrophy in pharmacoresistant temporal lobe epilepsy. *Neurology.* Ovid Technologies (Wolters Kluwer Health); 2009;72:1747–1754.
16. Caciagli L, Bernasconi A, Wiebe S, Koepp MJ, Bernasconi N, Bernhardt BC. A meta-analysis on progressive atrophy in intractable temporal lobe epilepsy. *Neurology.* Ovid Technologies (Wolters Kluwer Health); 2017;89:506–516.
17. Liu RSN, Lemieux L, Bell GS, et al. Cerebral damage in epilepsy: a population-based longitudinal quantitative MRI study. *Epilepsia.* Wiley; 2005;46:1482–1494.
18. Bernhardt BC, Bernasconi N, Concha L, Bernasconi A. Cortical thickness analysis in temporal lobe epilepsy: reproducibility and relation to outcome. *Neurology.* Ovid Technologies (Wolters Kluwer Health); 2010;74:1776–1784.
19. Tasch E, Cendes F, Li LM, Dubeau F, Andermann F, Arnold DL. Neuroimaging evidence of progressive neuronal loss and dysfunction in temporal lobe epilepsy. *Ann Neurol.* Wiley; 1999;45:568–576.
20. Desikan RS, Ségonne F, Fischl B, et al. An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *Neuroimage.* Elsevier BV; 2006;31:968–980.

21. Wakana S, Jiang H, Nagae-Poetscher LM, van Zijl PCM, Mori S. Fiber tract-based atlas of human white matter anatomy. *Radiology. Radiological Society of North America (RSNA)*; 2004;230:77–87.
22. Fortin J-P, Parker D, Tunç B, et al. Harmonization of multi-site diffusion tensor imaging data. *Neuroimage. Elsevier BV*; 2017;161:149–170.
23. Fortin J-P, Cullen N, Sheline YI, et al. Harmonization of cortical thickness measurements across scanners and sites. *Neuroimage. Elsevier BV*; 2018;167:104–120.
24. Liu M, Bernhardt BC, Bernasconi A, Bernasconi N. Gray matter structural compromise is equally distributed in left and right temporal lobe epilepsy. *Hum Brain Mapp. Wiley*; 2016;37:515–524.
25. Larivière S, Bayrak Ş, Vos de Wael R, et al. BrainStat: A toolbox for brain-wide statistics and multimodal feature associations. *Neuroimage. Elsevier BV*; 2023;266:119807.
26. Controlling the False Discovery Rate: AControlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing Practical and Powerful Approach to Multiple Testing.
27. Frangou S, Modabbernia A, Williams SCR, et al. Cortical thickness across the lifespan: Data from 17,075 healthy individuals aged 3-90 years. *Hum Brain Mapp. Wiley*; 2022;43:431–451.
28. Bernasconi N, Natsume J, Bernasconi A. Progression in temporal lobe epilepsy: differential atrophy in mesial temporal structures. *Neurology. Ovid Technologies (Wolters Kluwer Health)*; 2005;65:223–228.
29. Ballerini A, Arienzo D, Stassenko A, et al. Spatial patterns of gray and white matter compromise relate to age of seizure onset in temporal lobe epilepsy. *NeuroImage Clin. Elsevier BV*; 2023;39:103473.
30. Yasuda CL, Pimentel-Silva LR, Beltramini GC, et al. Brain volumes and white matter diffusion across the adult lifespan in temporal lobe epilepsy. *Ann Clin Transl Neurol. Wiley*; 2023;10:1106–1118.
31. Concha L, Kim H, Bernasconi A, Bernhardt BC, Bernasconi N. Spatial patterns of water diffusion along white matter tracts in temporal lobe epilepsy. *Neurology. Ovid Technologies (Wolters Kluwer Health)*; 2012;79:455–462.
32. Bernhardt BC, Kim H, Bernasconi N. Patterns of subregional mesiotemporal disease progression in temporal lobe epilepsy. *Neurology. Ovid Technologies (Wolters Kluwer Health)*; 2013;81:1840–1847.
33. Nagy SA, Horváth R, Perlaki G, et al. Age at onset and seizure frequency affect white matter diffusion coefficient in patients with mesial temporal lobe epilepsy. *Epilepsy Behav. Elsevier BV*; 2016;61:14–20.
34. Concha L, Beaulieu C, Gross DW. Bilateral limbic diffusion abnormalities in unilateral temporal lobe epilepsy. *Ann Neurol. Wiley*; 2005;57:188–196.
35. Coan AC, Appenzeller S, Bonilha L, Li LM, Cendes F. Seizure frequency and lateralization affect progression of atrophy in temporal lobe epilepsy. *Neurology. Ovid Technologies (Wolters Kluwer Health)*; 2009;73:834–842.
36. Bernhardt BC, Worsley KJ, Besson P, et al. Mapping limbic network organization in temporal lobe epilepsy using morphometric correlations: insights on the relation between mesiotemporal connectivity and cortical atrophy. *Neuroimage. Elsevier BV*; 2008;42:515–524.
37. Hsin Y-L, Harnod T, Chang C-S, Peng S-J. Increase in gray matter volume and white matter fractional anisotropy in the motor pathways of patients with secondarily generalized neocortical seizures. *Epilepsy Res. Elsevier BV*; 2017;137:61–68.
38. Coan AC, Campos BM, Yasuda CL, et al. Frequent seizures are associated with a network of gray matter atrophy in temporal lobe epilepsy with or without hippocampal sclerosis. *PLoS One. Public Library of Science (PLoS)*; 2014;9:e85843.
39. Coan AC, Appenzeller S, Bonilha L, Li LM, Cendes F. Seizure frequency and lateralization affect progression of atrophy in temporal lobe epilepsy. *Neurology. Ovid Technologies (Wolters Kluwer Health)*; 2009;73:834–842.
40. Hamed SA. Atherosclerosis in epilepsy: its causes and implications. *Epilepsy Behav. Elsevier BV*; 2014;41:290–296.
41. Tai XY, Koeppe M, Duncan JS, et al. Hyperphosphorylated tau in patients with refractory epilepsy correlates with cognitive decline: a study of temporal lobe resections. *Brain. Oxford University Press (OUP)*; 2016;139:2441–2455.
42. Sen A, Capelli V, Husain M. Cognition and dementia in older patients with epilepsy. *Brain*. 2018;141:1592–1608.

43. Hernández-Ronquillo L, Adams S, Ballendine S, Téllez-Zenteno JF. Epilepsy in an elderly population: Classification, etiology and drug resistance. *Epilepsy Res.* 2018;140:90–94.
44. Pardoe HR, Berg AT, Jackson GD. Sodium valproate use is associated with reduced parietal lobe thickness and brain volume. *Neurology*. Ovid Technologies (Wolters Kluwer Health); 2013;80:1895–1900.