

Short report

Ethnic disparities in time to bipolar disorder diagnosis: retrospective cohort study

Maria Richards-Brown, Violeta Perez-Rodriguez, Emma Butler, Sophie M. Allan, Rashmi Patel, Rebecca Strawbridge, Katherine Beck, Oliver D. Howes, Allan Hunter Young and Sameer Jauhar

Summary

Lithium is a first-line maintenance treatment for bipolar disorder yet delays in diagnosis and lithium initiation remain common. This retrospective cohort study examined whether time to bipolar disorder diagnosis and lithium initiation differ by ethnicity, gender or age among 88 patients at South London and Maudsley National Health Service Foundation Trust. Delay outcomes were non-normally distributed and log-transformed before analysis. No demographic characteristic was significantly associated with time to lithium initiation. Ethnicity was significantly associated with diagnostic delay, with individuals from minority ethnic groups experiencing longer delays than White individuals (median 556 v. 50.5 days; IQR [89.8, 1342] v. [6.25, 222]), which persisted after adjustment for age, gender, substance use and illness severity, and remained robust to outlier removal. Although social deprivation was not accounted

for, these findings suggest possible systematic diagnostic inequities in bipolar disorder, calling for culturally informed diagnostic practices and further investigation in larger, multi-site samples.

Keywords

Bipolar type I or II disorders; lithium; ethnicity; diagnostic delay; treatment.

Copyright and usage

© The Author(s), 2026. Published by Cambridge University Press on behalf of Royal College of Psychiatrists. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

Lithium is considered a first-line maintenance treatment for bipolar disorder, associated with substantial reductions in symptomatology, relapse, hospitalisation and suicide risk.^{1–3} Despite its established efficacy, delays in lithium initiation are common, with a recent analysis of the South London and Maudsley (SLaM) data used in the present study identifying a delay of 1.8 years between first assessment and lithium initiation.⁴ Diagnostic inaccuracy, such as misdiagnosing young women with borderline personality disorder rather than bipolar disorder or misinterpreting differential presentations of affective symptoms between ethnic groups,⁵ is thought to contribute to bipolar disorder diagnosis delays, with estimates ranging from 6 to 10 years.^{6,7} This study investigated whether time to bipolar disorder diagnosis and lithium initiation differ by gender, ethnicity or age.

Method

This sub-analysis of Perez-Rodriguez and colleagues⁴ used equivalent methods and sample (see original paper). The study was approved by the Clinical Records Interactive Search (CRIS) Oversight Committee in November 2020. CRIS has ethical approval from the South Central – Oxford C Research Ethics Committee (REC) as a database for secondary research. The REC reference is: 23/SC/0257. Briefly, this is a retrospective observational cohort study utilising routinely collected data within SLaM. Delay was defined as time from first SLaM assessment to recorded bipolar disorder or lithium initiation.

Delay distributions were assessed using Shapiro–Wilk tests. As outcomes were non-normally distributed, Mann–Whitney *U* tests, Kruskal–Wallis tests and Spearman's correlations examined associations with gender, ethnicity and age, respectively. Ethnicity was subsequently collapsed into White versus non-White categories. Univariate linear regressions were performed using log-transformed delay outcomes, followed by adjusted models incorporating age, gender, in-patient initiation and substance use.

Analyses were repeated after outlier removal, and *post hoc* power was assessed. All analyses were conducted in R (version 4.6.1 for macOS, Posit Software, PBC, Boston, USA, <https://posit.co/downloads>) with $\alpha = 0.05$.

Results

Time to diagnosis was the only outcome significantly associated with ethnicity (Kruskal–Wallis: $\chi^2 = 12.70$, $p = 0.026$, effect size = 0.094). Given small subgroup sizes (White $n = 38$, Black $n = 25$, Asian $n = 9$, Mixed $n = 5$), ethnicity was collapsed into a binary variable (White v. non-White). Although the authors are aware that research should avoid broad ethnic categories that minimise within-group heterogeneity, this collapsing was required to facilitate analyses.⁸ Non-White patients were diagnosed significantly later (median 556 v. 50.5 days, IQR (89.8, 1342) v. (6.25, 222); $\beta = 1.81$, s.e. = 0.51, $t = 3.60$, $p < 0.001$).

In the fully adjusted model (covariates: age, gender, illicit substance use, in-patient/out-patient prescribing), ethnicity remained the sole significant predictor ($\beta = 2.04$, s.e. = 0.54, $t = 3.80$, $p < 0.001$). The model explained 16.7% of variance overall ($R^2 = 0.167$, adjusted $R^2 = 0.117$, $F(5, 82) = 3.30$, $p = 0.009$). A *post hoc* power calculation confirmed 90% power to detect the observed effect ($f^2 = 0.20$) at $\alpha = 0.05$.

Outlier removal ($n = 12$, identified via Cook's distance, leverage and standardised residuals) strengthened the ethnicity effect and the model explained substantially more variance ($\beta = 2.58$, $p < 0.001$; $R^2 = 0.367$). In-patient versus out-patient setting of lithium prescribing also emerged as a significant predictor in the sensitivity model ($\beta = 1.06$, $p = 0.022$), with out-patients experiencing longer diagnostic delays.

Moreover, patients with an initial schizophrenia-spectrum disorder (SSD) diagnosis were disproportionately non-White: 9 of 11 (81.8%) were non-White compared with 41 of 77 (53.2%) without an initial SSD diagnosis. Given the small subgroup, findings are presented descriptively and should be interpreted cautiously.

Table 1 Model statistics from unadjusted and adjusted regressions examining demographic and clinical characteristics associated with time to diagnosis					
Variable	N (%) / mean (s.d.)	Univariate beta		Multivariable beta	
		coefficient log, days (s.e.)	p value	coefficient log, days (s.e.)	p-value
Age (years), mean (s.d.)	34.8 (12)	−0.01 (0.02)	0.68	0.02 (0.023)	0.49
Gender (ref: male), n (%)	27 (30.7)	0.56 (0.58)	0.34	0.90 (0.59)	0.13
Illicit substance use (ref: no substance use), n (%)	17 (19.3)	−0.29 (0.68)	0.67	0.04 (0.68)	0.95
Lithium initiation setting (ref: in-patient), n (%)	60 (68.2)	0.40 (0.58)	0.49	1.06	0.022
Ethnicity (ref: White), n (%)	38 (43.2)	1.81 (0.51)	<0.001	2.04 (0.54)	<0.001

Discussion

This study found that non-White individuals experienced significantly longer delays to bipolar disorder diagnosis than White individuals, an effect that remained after covariate adjustment and outlier removal (Table 1). The modest R^2 values are consistent with the multifactorial nature of diagnostic delay and noise in routinely collected clinical data, thus residual confounding from unmeasured factors cannot be excluded.

General barriers to bipolar disorder diagnosis, including misdiagnosis, healthcare access challenges, stigma and mistrust of services,⁹ may be amplified in non-White individuals. Misinterpretation of affective symptoms within psychotic rather than bipolar frameworks may contribute, alongside documented over-diagnosis of schizophrenia and greater diagnostic instability in Black populations.^{5,10,11} Supporting this possibility, non-White individuals disproportionately received SSD diagnoses before their bipolar disorder diagnosis, suggesting that diagnostic delay may represent a downstream consequence of misdiagnosis.

The absence of an ethnicity effect on time to lithium prescription may indicate that treatment decisions become more equitable once bipolar disorder is diagnosed. However, previous studies report less lithium and more antipsychotic prescribing among people of African ancestry.^{12–14} This null finding may reflect sampling bias, as all participants received lithium. Therefore, future research should expand the sample to individuals who are not administered lithium at all.

Age and gender did not significantly influence delay outcomes. Females nevertheless had descriptively longer delays to bipolar disorder diagnosis ($M=742.38$ days) compared with males ($M=514.93$ days). The predominantly female sample may be limiting the detection of gender effects. Other limitations, including limited sample size, insufficient documentation and broad ethnic classifications, have been reported previously and may obscure within-group heterogeneity.

These findings align with the evidence of under-recognition of bipolar disorder in non-White populations.^{5,10} Clinically, this emphasises the need for culturally informed diagnostic practices and reduced diagnostic overshadowing by psychotic disorder frameworks in non-White populations. Future studies should examine these findings in larger, representative samples while accounting for socioeconomic status.

Lived experience commentary (4th author)

I write as someone from a White British background who has lived with bipolar disorder type 1 throughout my adult life (first manic episode aged 18), with a lithium treatment delay of 14 years (first prescribed lithium aged 32). In the 5 years prior to commencing lithium, I had four manic episodes, but in the 5 years following lithium initiation, just one. My episodes have been devastating in

their impact on my ability to function, maintain relationships and sustain employment. Happily, I am now functioning very well.

I was offered lithium aged 19 but declined due to reluctance to have blood tests. I feel my consultant psychiatrist should have made a stronger case for the drug. A later psychiatrist stated lithium was not a good fit because it would make me feel ‘sluggish’ (it doesn’t). I was only started on lithium after a lengthy hospital admission, where I was given the options of either lithium or electroconvulsive therapy. I responded quickly to lithium and have only had one more minor manic episode since.

Lithium has been absolutely transformative for me and I only wish I had been prescribed it earlier. I’m sure it has also saved the National Health Service (NHS) significant costs. I commend this research into inequalities in lithium initiation, and while my story does not speak to the ethnic inequalities found in this paper, they are no doubt worthy of further exploration.

Maria Richards-Brown , Department of Psychosis Studies, King’s College London Institute of Psychiatry Psychology & Neuroscience, UK; and South London and Maudsley Mental Health NHS Trust, London, UK; **Violeta Perez-Rodriguez**, Department of Psychosis Studies, King’s College London Institute of Psychiatry Psychology & Neuroscience, UK; and South London and Maudsley Mental Health NHS Trust, London, UK; **Emma Butler** , Department of Psychiatry, University of Cambridge, UK; and Cambridgeshire and Peterborough Mental Health Partnership NHS Trust, Cambridge, UK; **Sophie M. Allan** , Department of Psychology, Clinical Psychology & Psychological Therapies, University of Cambridge, UK; and Department of Health Sciences, University of East Anglia, UK; **Rashmi Patel**, Department of Psychiatry, University of Cambridge, UK; and Department of Psychological Medicine, King’s College London Institute of Psychiatry Psychology & Neuroscience, UK; **Rebecca Strawbridge** , Department of Psychological Medicine, King’s College London Institute of Psychiatry Psychology & Neuroscience, UK; **Katherine Beck** , Department of Psychosis Studies, King’s College London Institute of Psychiatry Psychology & Neuroscience, UK; and South London and Maudsley Mental Health NHS Trust, London, UK; **Oliver D. Howes** , Department of Psychosis Studies, King’s College London Institute of Psychiatry Psychology & Neuroscience, UK; and South London and Maudsley Mental Health NHS Trust, London, UK; **Allan Hunter Young** , Department of Brain Sciences, Imperial College London, UK; **Sameer Jauhar** , Department of Brain Sciences, Imperial College London, UK

Correspondence: Maria Richards-Brown. Email: maria.richards-brown@kcl.ac.uk
First received 12 Aug 2026 UTC, final revision 3 Sep 2026 UTC, accepted 14 Sep 2026 UTC

Data availability

The data that support the findings of this study are available on request from the corresponding author (M.R.-B.).

Acknowledgement

No artificial intelligence tools have been used in the research or writing process for this manuscript.

Author contributions

M.R.-B. contributed to conceptualisation, methodology, formal analysis, project administration, visualisation, writing – original draft preparation and writing – review and editing. V.P.R. contributed to conceptualisation, data curation, methodology, investigation, project administration, resources and writing – review and editing. E.B. contributed to conceptualisation, methodology, formal analysis and writing – review and editing. S.M.A. contributed to methodology and writing – original draft preparation. R.P. contributed to methodology, resources, validation, visualisation and writing – review and editing. R.S. contributed to methodology, validation and writing – review and editing. K.B. contributed to supervision and writing – review

and editing. A.H.Y. contributed to funding acquisition, resources, supervision and writing – review and editing. O.D.H. contributed to funding acquisition, resources, supervision and writing – review and editing. S.J. contributed to funding acquisition, resources, supervision and writing – review and editing.

Funding

The authors disclosed receipt of the following financial support for the research, authorship and/or publication of this article. This paper represents independent research funded by the National Institute for Health and Care Research (NIHR) Maudsley Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London. V.P.R.'s salary is funded by the Rosetrees Trust. K.B. is supported by the Early Intervention Mission study (IRAS Ref: 333754), NIHR Mental Health Translational Research Collaboration Mission, hosted by the NIHR Oxford Health Biomedical Research Centre (NIHR203316) and the Rosetrees Trust. R.P. has received grant funding from the National Institute for Health and Care Research (NIHR301690). O.D.H. has received funding by Medical Research Council-UK (no. MC_A656_5QD30_2135), Maudsley Charity (no. 666) and Wellcome Trust (no. 094849/Z/10/Z) grants to O.D.H. and the NIHR Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London.

Declaration of interest

E.B. currently holds an unpaid secondment with Akviva Health, but has no financial relationship with the company and the duties are distinct from the included work. R.P. has participated in scientific advisory boards for Boehringer Ingelheim and Teva, has received grant funding from Janssen and has received consulting fees from Akviva Health, Boehringer Ingelheim, Bristol Myers Squibb, Clinilabs, Columbia Data Analytics, Eisai, Holmusk, Otsuka, Social Finance, Supernus and Teva. R.S. has received honoraria for educational lectures from Janssen. K.B. provides consultancy services to Bristol Myers Squibb. O.D.H. has received investigator-initiated research funding from and/or participated in advisory/speaker meetings organised by Alkermes, Angelini, Autifony, Biogen, Boehringer Ingelheim, Bristol Myers Squibb, Eli Lilly, Elysium, Heptares, Global Medical Education, Invivo, Janssen, Karuna, Lundbeck, Merck, Minerva, Neurocrine, Ontrack/ Pangea, Otsuka, Ovid, Pasithea, QCell, Sunovion, Recordati, Roche, Rovi, Teva and Viartis/Mylan. He was previously a part-time employee of Lundbeck A/S. O.D.H. has a patent for the use of dopaminergic imaging. The views expressed are those of the author(s) and not necessarily those of H Lundbeck A/S, the NHS/NIHR or the Department of Health. A.H.Y. is employed by Imperial College London and is an honorary consultant at South London and Maudsley NHS Foundation Trust (NHS UK). A.H.Y. is the editor of *Journal of Psychopharmacology* and deputy editor of *BJPsych Open*. A.H.Y. gives paid lectures and serves on advisory boards for the following companies with drugs used in affective and related disorders: Flow Neuroscience, Novartis, Roche, Janssen, Takeda, Noema pharma, Compass, AstraZeneca, Boehringer Ingelheim, Eli Lilly, LivaNova, Lundbeck, Sunovion, Servier, Allegan, Bionomics, Sumitomo Dainippon Pharma, Sage, Neurocentrx and Otsuka. A.H.Y. is principal investigator in the Restore-Life Vagus Nerve Stimulation registry study funded by LivaNova. A.H.Y. is principal investigator on ESKETINTRD3004: 'An Open-label, Long-term, Safety and Efficacy Study of Intranasal Esketamine in Treatment-resistant Depression.' A.H.Y. is principal investigator on 'The Effects of Psilocybin on Cognitive Function in Healthy Participants'. A.H.Y. is principal investigator on 'The Safety and Efficacy of Psilocybin in Participants with Treatment-Resistant Depression (P-TRD)'. A.H.Y. is principal investigator on 'A Double-Blind, Randomized, Parallel-Group Study with Quetiapine Extended Release as Comparator to Evaluate the Efficacy and Safety of Seltorexant 20 mg as Adjunctive Therapy to Antidepressants in Adult and Elderly Patients with Major Depressive Disorder with Insomnia Symptoms Who Have Responded Inadequately to Antidepressant Therapy.' (Janssen). A.H.Y. is principal investigator on 'An Open-label, Long-term, Safety and Efficacy Study of Aticaprant as Adjunctive Therapy in Adult and Elderly Participants with Major Depressive Disorder (MDD)'. (Janssen). A.H.Y. is principal investigator on 'A Randomized, Double-blind, Multicentre, Parallel-group, Placebo-controlled Study to Evaluate the Efficacy, Safety, and Tolerability of Aticaprant 10 mg as Adjunctive Therapy in Adult Participants with Major Depressive Disorder (MDD) with Moderate-to-severe Anhedonia and Inadequate Response to Current Antidepressant Therapy'. A.H.Y. is principal investigator on 'A Study of Disease Characteristics and Real-life Standard of Care Effectiveness in Patients with Major Depressive Disorder (MDD) With Anhedonia and Inadequate Response to Current Antidepressant Therapy Including an SSRI or SNRI.' (Janssen). A.H.Y. is the UK chief investigator for Compass; COMP006 & COMP007 studies (Compass). A.H.Y. is the UK chief investigator for Novartis MDD study MJ821A12201. A.H.Y. has no shareholdings in pharmaceutical companies. Grant funding (past and present): National Institute of Mental Health, USA; Canadian Institutes of Health Research, Canada; National Alliance for Research on Schizophrenia and Depression, USA; Stanley Medical Research Institute, USA; Medical Research Council, UK; Wellcome Trust, UK; Royal College of Physicians of Edinburgh; British Medical Association, UK; University of British Columbia – Vancouver General Hospital Foundation, Canada; Women's Economic Development Council, Canada;

Coast Capital Savings Depression Research Fund, Canada; Medical Services Foundation of British Columbia, Canada; NIHR, UK; Janssen (UK) and European Union Horizon 2020. S.J. has received personal fees or honoraria from Recordati, LB Pharmaceuticals, Boehringer Ingelheim, Accord Pharmaceuticals, Lundbeck, Janssen and Sunovion. He has served on an advisory board for Boehringer Ingelheim and has undertaken consultancy work for LB Pharmaceuticals. He has received personal fees from the Wellcome Trust and has served on panels or undertaken non-financial professional work for the Wellcome Trust, the National Institute for Health and Care Excellence, the British Association for Psychopharmacology and the Royal College of Psychiatrists. All interests were outside the submitted work. It should be noted that R.S., A.H.Y. and S.J. are members of the *BJPsych Open* Editorial Board; however, they did not take part in the review or decision-making process for this paper. All other authors (M.R.-B., V.P.R. and S.M.A.) had no declarations.

References

- 1 Berk M, Daglas R, Dandash O, Yücel M, Henry L, Hallam K, et al. Quetiapine v. lithium in the maintenance phase following a first episode of mania: randomised controlled trial. *Br J Psychiatry* 2017; **210**: 413–21.
- 2 Del Matto L, Muscas M, Murru A, Verdolini N, Anmella G, Fico G, et al. Lithium and suicide prevention in mood disorders and in the general population: a systematic review. *Neurosci Biobehav Rev* 2020; **116**: 142–53.
- 3 Severus E, Taylor MJ, Sauer C, Pfennig A, Ritter P, Bauer M, et al. Lithium for prevention of mood episodes in bipolar disorders: systematic review and meta-analysis. *Int J Bipolar Disord* 2014; **2**: 15.
- 4 Perez-Rodriguez V, Nguyen T, Beck K, Butler R, Strawbridge R, et al. Time to lithium: a case register study of lithium initiation in bipolar disorder. *J Psychopharmacol* 2026; **40**: 945–50.
- 5 Haeri S, Williams J, Kopeykina I, Johnson J, Newmark A, Cohen L, et al. Disparities in diagnosis of bipolar disorder in individuals of African and European descent: a review. *J Psychiatr Pract* 2011; **17**: 394–403.
- 6 Scott J, Graham A, Yung A, Morgan C, Bellivier F, Etain B. A systematic review and meta-analysis of delayed help-seeking, delayed diagnosis and duration of untreated illness in bipolar disorders. *Acta Psychiatr Scand* 2022; **146**: 389–405.
- 7 Drancourt N, Etain B, Lajnef M, Henry C, Raust A, Cochet B, et al. Duration of untreated bipolar disorder: missed opportunities on the long road to optimal treatment. *Acta Psychiatr Scand* 2013; **127**: 136–44.
- 8 Sharghi S, Khalatbari S, Laird A, Lapidus J, Enders FT, Meinzen-Derr J, et al. Race, ethnicity, and considerations for data collection and analysis in research studies. *J Clin Transl Sci* 2024; **8**: e182.
- 9 Mehra N, Bhatia A, Ayis S, Kordowicz M, Hack-Polay D, Akilan V, et al. *Reasons for Diagnostic Delays in Bipolar Disorder: Systematic Review and Narrative Synthesis*. Qeios, 2024 (<https://www.qeios.com/read/BGTD5K>).
- 10 Akinhanmi MO, Biernacka JM, Strakowski SM, McElroy SL, Balls Berry JE, Merikangas KR, et al. Racial disparities in bipolar disorder treatment and research: a call to action. *Bipolar Disord* 2018; **20**: 506–14.
- 11 Barry S, Jarskog LF, Xia K, Torpurni RS, Wu X, Zeng X. Racial disparities in clozapine prescription patterns among patients with schizophrenia. *Psychiatr Serv* 2024; **75**: 733–9.
- 12 Akinhanmi M, El-Amin S, Balls-Berry JE, Vallender EJ, Ladner M, Geske J, et al. Decreased core symptoms of mania and utilization of lithium/mood stabilizing anticonvulsants in U.S. bipolar I patients of African vs European ancestry. *J Affect Disord* 2020; **260**: 361–5.
- 13 Johnson KR, Johnson SL. Inadequate treatment of black Americans with bipolar disorder. *Psychiatr Serv* 2014; **65**: 255–8.
- 14 Taylor-Desir MJ, Sanchez-Ruiz JA, Vallender EJ, Singh B, Meagher KM, Frye MA. Persistence of lithium underutilization in patients of African ancestry with bipolar disorder underuse of lithium in black patients. *Gen Psychopharmacol Bull* 2025; **55**: 47–63.