

This document is confidential and is proprietary to the American Chemical Society and its authors. Do not copy or disclose without written permission. If you have received this item in error, notify the sender and delete all copies.

The muscle metabolome differs between healthy and frail older adults

Journal:	<i>Journal of Proteome Research</i>
Manuscript ID	pr-2015-00840a.R2
Manuscript Type:	Article
Date Submitted by the Author:	27-Dec-2015
Complete List of Authors:	Fazelzadeh, Parastoo; Wageningen University, Human Nutrition Hangelbroek, Roland; Wageningen University, Human Nutrition Tieland, Michael; Wageningen University, Human Nutrition de Groot, Lisette; Wageningen University, Human Nutrition Verdijk, Lex; Maastricht University, Human Movement Sciences van Loon, Luc; Maastricht University, Human Movement Sciences Smilde, Age; Swammerdam Institute for Life Sciences, Alves, Rodrigo; Leiden University, Analytical BioSciences Vervoort, Jacques; Wageningen University, Laboratory of Biochemistry Muller, Michael; University of East Anglia, Norwich, Institute of Food Research van Duynhoven, John; Wageningen University, Laboratory of Biophysics Boekschoten, Mark; Wageningen University, Human Nutrition

SCHOLARONE™
Manuscripts

The muscle metabolome differs between healthy and frail older adults

Parastoo Fazelzadeh^{§□=}, Roland W.J. Hangelbroek^{§□=}, Michael Tieland^{§□}, Lisette CPGM de Groot^{§□}, Lex B Verdijk^{§Δ}, Luc JC van Loon^{§Δ}, Age K. Smilde[‡], Rodrigo DAM Alves^{‡□}, Jacques Vervoort^φ, Michael Müller⁺, John P.M. van Duynhoven^{□†}, Mark V Boekschoten^{§□}*

[§]Top Institute Food and Nutrition, Wageningen, The Netherlands

[□] Nutrition, Metabolism and Genomics group, Division of Human Nutrition, Wageningen University, Wageningen, The Netherlands

^ΔDepartment of Human Movement Sciences, NUTRIM School of Nutrition and Translational Research in Metabolism, Maastricht University, Maastricht, the Netherlands

[‡] Biosystems Data Analysis group, University of Amsterdam, Amsterdam, The Netherlands

[‡] Department of Analytical Biosciences, Leiden University, Leiden, The Netherlands

^φLaboratory of Biochemistry, Wageningen University, Wageningen, The Netherlands

[†]Laboratory of Biophysics, Wageningen University, Wageningen, The Netherlands

⁺Norwich Medical School, University of East Anglia, United Kingdom

[□]Netherlands Metabolomics Centre, Leiden, The Netherlands

⁼These authors contributed equally to this study.

ABSTRACT:

Populations around the world are aging rapidly. Age-related loss of physiological functions negatively affects quality of life. A major contributor to the frailty syndrome of aging is loss of skeletal muscle. In this study we assessed the skeletal muscle biopsy metabolome of healthy young, healthy older and frail older subjects to determine the effect of age and frailty on the metabolic signature of skeletal muscle tissue. In addition, the effects of prolonged whole-body resistance-type exercise training on the muscle metabolome of older subjects were examined. The baseline metabolome was measured in muscle biopsies collected from 30 young, 66 healthy older subjects and 43 frail older subjects. Follow-up samples from frail older (24 samples) and healthy older subjects (38 samples) were collected after 6 months of prolonged resistance-type exercise training. Young subjects were included as a reference group. Primary differences in skeletal muscle metabolite levels between young and healthy older subjects were related to mitochondrial function, muscle fiber type, and tissue turnover. Similar differences were observed when comparing frail older subjects with healthy older subjects at baseline. Prolonged resistance-type exercise training resulted in an adaptive response of amino acid metabolism, especially reflected in branched chain amino acids and genes related to tissue remodeling. The effect of exercise training on branched-chain amino acid-derived acylcarnitines in older subjects points to a downward shift in branched-chain amino acid catabolism upon training. We observed only modest correlations between muscle and plasma metabolite levels, which pleads against the use of plasma metabolites as a direct read-out of muscle metabolism and stresses the need for direct assessment of metabolites in muscle tissue biopsies.

KEYWORDS: muscle biopsy, frailty, aging, tissue remodeling

INTRODUCTION

Populations around the world are aging rapidly. It has been estimated that people older than 60 years of age will make up 22% of the world population, whereas people older than 80 years of age will account for 4.4% of the world population in 2050¹. Age-related loss of physiological functions compromises independence at older age. A major contributor to the frailty syndrome of aging is skeletal muscle loss, which can lead to increased disability in the older population. In most people muscle mass and strength start to decline around the age of 35 with more progressive muscle loss observed after the age of 65²⁻⁶.

Exercise training has been applied to counter the effects muscle mass loss, such as loss of strength^{7, 8}. We recently demonstrated that prolonged resistance-type exercise training partially shifts the skeletal muscle transcriptome of older subjects toward an expression pattern observed in muscle tissue of young subjects, with changes in gene expression related to vascularisation, tissue remodelling and glucose metabolism⁹. The transcriptome analysis also revealed substantial differences between healthy young men, healthy older subjects and frail older subjects before any intervention was undertaken. In particular, genes related to mitochondrial function had lower expression levels in older subjects compared to young subjects.

The age-related differences in gene expression level in muscle are expected to be reflected in local levels of metabolites. However, limited information is available on the effect of aging on muscle metabolite levels. A recent study in mice showed that aging affects muscle glucose and fatty acid metabolism¹⁰, whereas a study in rats indicated that aging leads to muscle group-specific perturbations in lipid and glucose metabolism consistent with mitochondrial dysfunction¹¹. In humans, it was found that changes in lipid content and oxidative activity in skeletal muscle during aging are related to a shift in muscle fiber type¹². Interestingly, exercise training led to reprogramming of mitochondrial function and intermediary

1
2
3 metabolism in insulin-insensitive obese subjects ¹³. In both human studies, the metabolic
4
5 profiling platforms that were used had limited coverage and focused on specific sub-
6
7 metabolomes. A major bottleneck in achieving extended coverage has been the limited
8
9 amount of muscle tissue material that can be sampled from human subjects.
10

11 On the basis of the observed shift in transcriptome profiles after training we hypothesized that
12
13 a similar shift towards the young phenotype occurs in the muscle metabolome after resistance-
14
15 type exercise training in older subjects. The recent development of a comprehensive targeted
16
17 metabolic profiling platform optimized and validated for small muscle biopsies makes it
18
19 possible to elaborately verify this hypothesis ¹⁴. Here we aim to investigate the effect of aging
20
21 and frailty on the skeletal muscle metabolome. In addition, we examined the impact of
22
23 prolonged resistance-type exercise training on the metabolome of frail and healthy older
24
25 subjects.
26
27
28
29
30
31
32

33 MATERIALS AND METHODS

34 Experimental design

35
36
37
38
39 Muscle biopsies (*m. vastus lateralis*), plasma and serum samples were collected from frail
40
41 older ^{15, 16}, healthy older ^{17, 18} and young subjects ¹⁹. The frail group included both frail and
42
43 pre-frail subjects and are hereafter referred to as frail. Medical history of all subjects was
44
45 evaluated by medical questionnaires, which were analyzed by a physician. Subjects who were
46
47 unable to participate in the exercise program due to pain were excluded prior to starting the
48
49 intervention. Included subjects who showed severe discomfort during the training sessions
50
51 were excluded from further participation in the study. Baseline metabolite levels were
52
53 measured in skeletal muscle tissue of 30 young, 66 healthy and 43 frail older subjects (Table
54
55
56
57 1). We also measured metabolites in plasma and serum for 50 young, 76 healthy and 62 frail
58
59
60

1
2
3 older subjects (Supporting Information Table S1). The transcriptome of the muscle samples
4
5 was measured in an earlier study⁹.
6

7 Baseline samples from healthy young male subjects served as reference and were derived
8
9 from several studies performed within our group, in which exactly the same techniques and
10
11 processing were used for sample collection. Samples from the frail and healthy older subjects
12
13 were obtained from two study centers. More details of the studies can be found in the papers
14
15 published previously^{15, 18}. In addition, muscle biopsies were obtained after 6 months of
16
17 resistance-type exercise training for 38 healthy and 24 frail older subjects. All muscle biopsies
18
19 and blood samples were obtained in the morning after an overnight fast. Subjects ate a
20
21 standardized meal the evening before sample collection. Subjects refrained from doing any
22
23 strenuous physical activity for 3 days prior to muscle biopsy collection.
24
25
26

27
28 Fried criteria²⁰ were used to assess the frailty in older subjects, which takes into account
29
30 unintentional weight loss, weakness, self-reported exhaustion, slow walking speed, and low
31
32 physical activity. Based on the above mentioned criteria, the healthy older subjects were not
33
34 considered frail or pre-frail at the start of the intervention study²¹.
35
36

37 In our study population, all older subjects regardless of their health status (frail or healthy)
38
39 improved in muscle performance following 6 months of resistance-type exercise training
40
41 (Supporting Information Figure S1 Table S2), as illustrated by significantly increased leg
42
43 extension and leg press strength after training (P -value < 0.01)^{9, 15, 21}. Both healthy older and
44
45 frail older subjects followed similar progressive full-body resistance type exercise training. In
46
47 brief, the training consisted of a 5 minute warm-up on a cycle ergometer, followed by 4 sets
48
49 on the leg-press and leg-extension machines. In addition, 3 sets of chest press, lat pulldown,
50
51 pec-dec and vertical row machines were performed (Technogym, Rotterdam, The
52
53 Netherlands). The healthy older subjects trained 3 times per week and frail subjects trained 2
54
55 times per week. Moreover, subjects received a protein or control supplement during the study.
56
57
58
59
60

1
2
3 Healthy older subjects took 15 gram milk protein or control drink at breakfast and frail older
4
5 subjects took a similar 15 gram drink (milk protein or control) at breakfast and lunch every
6
7 day throughout the entire 6 months intervention. Full details can be found in the earlier papers
8
9 on the phenotypical impact of training on our study population ^{15, 21}. All studies were
10
11 approved by The Medical Ethics Committee of either Wageningen University or Maastricht
12
13 University and comply with the Declaration of Helsinki.
14
15

16 17 18 **Metabolomics analysis of circulating metabolites**

19
20 Amino acids and biogenic amines were derivatized (Acc-Tag) in 5 μ L aliquots of plasma.
21
22 Samples were analyzed using an ACQUITY UPLC system with autosampler (Waters, Etten-
23
24 Leur, The Netherlands) coupled with a Xevo Tandem quadrupole mass spectrometer (Waters)
25
26 operated using QuanLynx data acquisition software (version 4.1; Waters). An Accq-Tag Ultra
27
28 column (Waters) was used. The Xevo TQ was used in the positive-ion electrospray mode and
29
30 all analytes were monitored in Multiple Reaction Monitoring (MRM) using nominal mass
31
32 resolution. Acquired data were evaluated using TargetLynx software (Waters), by integration
33
34 of assigned MRM peaks and normalization using proper internal standards²²,
35
36

37
38 Acylcarnitines, trimethylamine-N-oxide, choline, betaine, deoxycarnitine and carnitine were
39
40 analyzed in 5 μ L plasma, spiked with an internal standard, using a UPLC-MS/MS. Also here
41
42 an Accq-Tag Ultra column was used. The Xevo TQ was used in the positive-ion electrospray
43
44 mode and all analytes were monitored in MRM using nominal mass resolution. In-house
45
46 developed algorithms²³ were applied using the pooled QC samples to compensate for shifts in
47
48 the sensitivity of the mass spectrometer over the batch.
49
50

51
52 Organic acids were measured by GC-MS using 50 μ L of plasma sample prepared using a two-
53
54 step derivatization procedure with subsequent oximation using methoxyamine hydrochloride
55
56 (MeOX) and silylation using N-Methyl-N-(trimethylsilyl)trifluoroacetamide (MSTFA).
57
58 Samples were measured on an Agilent GC (7890A) coupled to Agilent Quadrupole-MS with
59
60

1
2
3 EI source (Agilent MSD 5975C). Separation was performed using a HP-5MS column (30 m x
4 0.25 m x 0.25 μm ; Agilent). The raw data were pre-processed using Agilent MassHunter
5 Quantitative Analysis software for GC-MS (Agilent, Version B.04.00), and quantitation of
6
7 metabolite response was calculated as the peak area ratios of the target analyte to the
8
9 respective internal standard. In-house developed algorithms were applied using the pooled QC
10
11 samples to compensate for shifts in the sensitivity of the mass spectrometer over the batch.
12
13
14 Serum metabolite concentrations determined by NMR were measured as described by
15
16 Mihaleva et al ²⁴. In short, serum samples were ultrafiltrated and automated quantum
17
18 mechanical line shape fitting of ¹H NMR spectra was performed using PERCH.
19
20
21
22
23
24

25 **In tissue metabolome analysis**

26
27 Metabolites were extracted from 10 mg of wet muscle tissue. This tissue was further
28
29 lyophilized and weighted to determine the dry tissue mass. After pulverizing the tissue,
30
31 metabolites were extracted using methanol/chloroform/water (MCW). The extraction method
32
33 used in this study has extensively been described and characterized elsewhere ¹⁴.
34
35

36 Amines, acylcarnitines and oxylipins were measured using the platforms also used for
37
38 measurement of these metabolites in plasma. The validation of these methods for human
39
40 tissue biopsies is described in detail elsewhere ²³. In-house developed algorithms were applied
41
42 to compensate for shifts in the sensitivity of the mass spectrometer over multiple batches of
43
44 measurements ²³. The metabolite response was determined by the peak area ratio of the target
45
46 analyte to the appropriate internal standard. These response ratios were used in the subsequent
47
48 data analysis. ATP, ADP, creatine and phosphocreatine were determined using established
49
50 enzymatic assays. For the first 3 metabolites, commercially available fluorimetric assay kits
51
52 were used (BioVision cat.# K354-100, K355-100 and K635-100) following the
53
54 manufacturer's instructions. Phosphocreatine was measured according to a colorimetric assay
55
56
57
58
59
60

1
2
3 kit protocol described by Szas et al.²⁵. In total 96 metabolites including amine, acylcarnitines,
4
5 organic acids, oxylipins and a number of nucleotides were measured.
6
7

8 **Statistical analysis**

9
10 Statistical analysis was performed on log-transformed data. We used analysis of variance
11
12 (ANOVA) for between group comparisons at baseline. *P*-value <0.05 was considered
13
14 statistically significant. We used linear mixed models for assessment of the training effect.
15
16 Our model included exercise training, subject, sex, protein supplementation and within
17
18 subject correlation. Analyses of the training effect was performed separately for frail and
19
20 healthy older subjects due to differences in training frequency. To summarize acylcarnitines
21
22 into one single metabolite (eigenmetabolite), the Non-linear Iterative partial least squares
23
24 (NIPALS) algorithm ²⁶ of the mixOmics R library was used to calculate the Singular Value
25
26 Decomposition (SVD) of acylcarnitines. To integrate metabolomics and transcriptomics data
27
28 the multilevel sparse partial least squares (sPLS) module from the mixOmics R library was
29
30 used ²⁷. We applied canonical correlation (cut-off => 0.8), which highlights the strongest
31
32 correlations between the two data sets. FactoMineR was used to perform principal component
33
34 analysis (PCA) ²⁸. All analyses were done using R (version 3.02).
35
36
37

38
39 ASCA (ANOVA Simultaneous Component Analysis) was performed to determine global
40
41 differences on metabolites. ASCA is a multivariate method that partitions variation in the data
42
43 and enables to interpret these partitions by simultaneous component analysis ²⁹. Analysis was
44
45 performed separately for the frail and healthy older subjects, using delta values of each
46
47 metabolite for each individual (value after training - value before training) with supplement,
48
49 sex and their interaction in the model as factors. Analysis was done under Matlab (version
50
51 R2012a).
52
53
54
55
56
57
58
59
60

RESULTS

Baseline comparisons between healthy older, frail older and young subjects

Comparison of the plasma metabolite profiles of young and healthy older subjects showed marked differences in levels of many metabolites between the groups. Analysis of variance (ANOVA) showed that differences in plasma metabolite levels between older and young subjects were mainly accounted for by the amino acids and acylcarnitines. These differences were in line with previous observations of age-related effects on the plasma metabolome^{10, 30, 31}.

Figure 1 shows a principal component analysis (PCA) plot of the biogenic amine profiles of the muscle biopsies obtained from young, healthy older and frail older subjects, before and after exercise. We observed clear age-related differences in the biogenic amine profiles of skeletal muscle, as well as a difference between healthy and frail older subjects. The PCA plot also revealed an effect of prolonged resistance-type exercise training on amine concentrations in muscle tissue. PCA revealed a similar effect for organic acids in muscle (Supporting Information Figure S2).

ANOVA yielded a series of skeletal muscle metabolites that differ significantly between healthy older subjects and young subjects. Many of these metabolites are amino acids and organic acids (Table 2). The outcome of ANOVA modeling of muscle metabolites in healthy vs. frail older subjects is presented in Table 3.

Effect of prolonged resistance-type exercise training in frail and healthy older subjects

In the PCA plot in Figure 1 we can observe that prolonged resistance-type exercise training has an effect on the biogenic amine profile of skeletal muscle tissue of frail and healthy older subjects. The trends in the PCA plot suggest that upon training both healthy and frail older subjects shift towards a younger phenotype (see also the plot for the average value of isoleucine in Supporting information Figure S3,a). The variation in this PCA plot may not

1
2
3 only be due to phenotype (young, healthy and frail old) and exercise, but also to sex and
4
5 protein supplementation. Accordingly, we performed ANOVA Simultaneous Component
6
7 Analysis (ASCA) to account for these different sources of biological variation. We observed
8
9 that prolonged resistance-type exercise training had a weak effect on muscle tissue metabolite
10
11 levels. No significant interaction of protein supplementation with training in both frail and
12
13 healthy older subjects could be observed. Interaction of sex with training was not significant
14
15 in healthy older subjects, but was significant in frail older subjects (P -value=0.03).

16
17
18 Next we used linear mixed models on univariate metabolite levels to account for phenotype,
19
20 exercise, protein supplementation and sex as sources of biological variation. We observed that
21
22 prolonged resistance-type exercise training affected muscle levels of acylcarnitines in both the
23
24 healthy older and frail older subjects (Table 4 and 5, respectively). These include
25
26 acylcarnitines that are produced during oxidation of branched chain amino acids (propionyl
27
28 (C3), methylmalonyl (C4-DC), and isovaleryl (C5) acylcarnitines) and acylcarnitines that are
29
30 produced during fatty acid oxidation (C6-C20 acylcarnitines)³². Because the exercise effect
31
32 on several acylcarnitines was nearly significant (P -value >0.05), we used the singular value
33
34 decomposition to summarize the levels of fatty acid derived and amino acid derived
35
36 acylcarnitines (denoted as FAAC and AAAC, respectively in Figure 4 and 5). Subsequently,
37
38 we also performed linear mixed model analysis to determine the effect of training on the fatty
39
40 acid derived and amino acid derived acylcarnitines. The amino acid derived acylcarnitines
41
42 levels were significantly decreased after training in both healthy and frail older subjects (see
43
44 example of the effects for propionylcarnitine in Supporting information Figure S3,b).

45
46
47 We used multilevel sPLS to integrate changes in the muscle transcriptome and muscle
48
49 metabolome after training. Here the goal was to investigate the interaction of two matched
50
51 datasets and the selection of subsets of either positively or negatively correlated variables
52
53 across all subjects. Only those genes were selected that were significantly changed by training
54
55
56
57
58
59
60

1
2
3 (FDR<0.05) in both frail and healthy older subjects. The resulting correlation network in
4
5 Figure 2 shows that especially branched chain amino acids correlate with genes related to
6
7 connective tissue/extracellular matrix such as collagen, laminin and Secreted Protein, Acidic,
8
9 Cysteine-Rich (SPARC) .
10

11 12 **Correlation between muscle and plasma metabolites levels**

13
14 Next we investigated to what extent baseline plasma metabolite levels reflect muscle
15
16 metabolite levels. To that end we first constructed separate correlation heatmaps for muscle
17
18 metabolite levels. Figure 3a-b shows that metabolites that belong to the same group of metabolites (e.g. amino acids, acylcarnitines) are correlated to
19
20 each other, which is observed for both plasma and muscle. However, the heatmap of the
21
22 correlation between muscle and plasma metabolites (Figure 3c) showed only minor to
23
24 moderate correlations (Pearson correlation between 0.3 and 0.5). The correlation networks
25
26 (Figure 3d) revealed the strongest correlations for 3-hydroxybutyric acid, 4-hydroxyproline,
27
28 proline, branched chain amino acids and several acylcarnitines. Correlations between serum
29
30 and muscle metabolites are presented in Supporting Information Figure S4. The results are
31
32 generally similar to the correlations observed between plasma and muscle metabolites.
33
34
35
36
37
38
39
40

41 **DISCUSSION**

42 43 44 **Comprehensive metabolic profiling of muscle biopsies: differences with age and effect of** 45 46 **resistance-type exercise training in a heterogeneous study population** 47

48
49
50 Comprehensive metabolic profiling of muscle biopsies revealed pronounced differences
51
52 between the muscle biopsy metabolomes of healthy young, healthy older and frail older
53
54 subjects. Furthermore, distinct effects of prolonged resistance-type exercise training could be
55
56 observed in both groups of older subjects..
57
58
59
60

1
2
3 A key asset of our analysis was the deployment of five robust profiling platforms using a
4 single and thoroughly validated muscle-biopsy extraction procedure ¹⁴. The analytical
5 variation in these profiles was around 20-30%, which was smaller than the biological
6 variation between the muscle biopsies. In order to separate the sources of biological
7 variation in our heterogeneous study population, we used univariate linear mixed models. Interestingly,
8 the metabolic effects of supplementation and differences between males and females were
9 relatively minor compared to the differences between old and young and between healthy and
10 frail. The effects were also minor compared to the response to prolonged resistance-type
11 exercise training. A multivariate approach (ASCA) did not show any significant effects of
12 training, probably because not all sources of biological variation could be adequately
13 accounted for ³³. In the following discussion we focus on the univariate linear mixed model
14 approach since it allows us to more precisely account for protein supplementation and sex in
15 our study population.
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31

32 **Baseline comparisons**

33
34 Interestingly, in muscle biopsies baseline levels of metabolites from the tricarboxylic acid
35 cycle (succinic acid, fumaric acid, 2-ketoglutaric acid) were lower in the healthy older
36 subjects compared to young subjects. These differences were accompanied by lower levels of
37 ATP, ADP, branched chain amino acids and acylcarnitines in the healthy older subjects,
38 suggesting impaired mitochondrial function or a lower number of mitochondria in the muscle
39 of the older subjects, which in turn may be a consequence of the lower habitual physical
40 activity of the older subjects. These observations are in line with data on the muscle
41 transcriptomic profiles of these subjects, which show decreased expression of genes related to
42 mitochondrial function and oxidative phosphorylation in the older subjects compared to
43 young subjects, with the lowest expression observed in the frail older subjects ⁹. These
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 findings are in agreement with other studies^{10, 34-38}, which also showed a down-regulation of
4
5 the mitochondrial electron-transport chain pathway in older as compared to young subjects.
6

7
8 One of our findings is that levels of 4-hydroxyproline and proline are lower in the healthy
9
10 older subjects compared to the young subjects. Both of these amino acids have been
11
12 associated with collagen turnover^{39, 40}. Muscle levels of two precursors of proline, ornithine
13
14 and arginine, are higher in the older subjects, which together with lower levels of 4-
15
16 hydroxyproline and proline may be due to dysfunction of the mitochondrial ornithine
17
18 aminotransferase⁴¹, leading to accumulation of arginine and ornithine. This notion is in line
19
20 with slightly lower expression of genes related to tissue remodeling, such as collagen in the
21
22 older subjects compared to the young subjects⁹.
23

24
25 We observed higher levels of β -isoamino butyric acid (BAIBA) in muscle tissue of healthy
26
27 older subjects compared to young subjects. This metabolite has been suggested to be
28
29 produced upon exercise by expression of PGC-1 α and has been proposed as a myokine
30
31 stimulating browning of white adipose tissue and hepatic β -oxidation⁴². In our study
32
33 population we observed lower expression of PGC-1 α target genes in healthy older subjects
34
35 compared to young subjects, and in frail older subjects compared to healthy older subjects⁹.
36
37 The activity of PGC-1 α is modulated post-translationally by NAD⁺ dependent deacetylation
38
39 by SIRT1 (Figure 4-b)⁴³. Since NAD⁺ levels decrease with age⁴⁴, we expect a concurrent
40
41 downregulation of PGC-1 α ⁴⁵. However, lower levels of BAIBA were only observed when
42
43 comparing frail to healthy older subjects, whereas BAIBA levels were higher in healthy older
44
45 subjects compared to young subjects. Assuming that aging lead to down-regulation of PGC-
46
47 1 α , the observed changes in baseline muscle BAIBA levels are not consistent with BAIBA
48
49 serving as a marker for PGC-1 α activity. Thus, we cannot confirm the notion that BAIBA acts
50
51 as a PGC-1 α induced myokine. A possible explanation for the discrepancy is that in our study
52
53
54
55
56
57
58
59
60

1
2
3 differences in PGC-1 α expression are caused by age and frailty-dependent processes, whereas
4
5 previously described PGC-1 α -mediated effects on BAIBA were caused by acute exercise.
6

7
8 Two polyamines, spermine and spermidine, were found to be significantly different between
9
10 the frail in comparison to the healthy older subjects. Polyamines are involved in tissue
11
12 regeneration and cell proliferation and differences are associated with both exercise and
13
14 muscle pathology⁴⁶⁻⁴⁸. The genes directly involved in the polyamine pathway are however
15
16 not differentially expressed between frail and healthy older subjects. Hence the observed
17
18 differences in polyamine levels between the frail and healthy older subjects are more likely to
19
20 be attributed to effects at the level of enzyme activity or metabolite transport. Previous studies
21
22 have shown that perturbations in polyamine metabolism are associated with neuromuscular
23
24 disorders^{49, 50}. On the transcriptome level we indeed found indications of neuromuscular
25
26 perturbations in the frail older subjects⁹. In addition, increased spermine levels were recently
27
28 linked to skeletal muscle atrophy⁵¹. However, we observed lower levels of spermine in frail
29
30 subjects in comparison to healthy older subjects, even though frail older subjects generally
31
32 have less skeletal muscle and are likely to exhibit more extensive muscle atrophy. Ost *et al.*
33
34 recently reported that spermidine is increased in the skeletal muscle of mice overexpressing
35
36 uncoupling protein 1. The authors proposed that this might be an adaptive response to cope
37
38 with the additional oxidative stress⁵². This would suggest that also in the frail elderly the
39
40 increase in spermidine is a response to reduced oxidative capacity and increase in oxidative
41
42 stress.
43
44
45
46
47

48
49 The level of carnosine is decreased in healthy older subjects in comparison to young subjects,
50
51 and in frail in comparison to healthy older subjects. Carnosine is an abundant metabolite in
52
53 muscle where it plays an important role in intracellular pH buffering⁵³. Carnosine has also
54
55 been associated with chelation of metal ions and antioxidant activity⁵⁴. Furthermore,
56
57 carnosine levels are higher in type II muscle fibers compared to type I. A likely explanation
58
59
60

1
2
3 for the significantly lower levels of carnosine in healthy and frail older subjects is therefore a
4
5 decrease of the muscle fiber II/I ratio with respectively age and lack of exercise⁵⁵.
6

7
8 Several oxylipins derived from linoleic acid (LA) and α -linoleic acid (ALA) occur at higher
9
10 levels in the muscle of the healthy older subjects compared to young subjects. On the other
11
12 hand, levels of metabolites derived from the Δ -6 desaturase product dihomo- γ -linoleic acid
13
14 (DGLA) are lower in healthy older subjects. We postulate that due to reduced Δ -6 desaturase
15
16 activity linoleic acid and α -linoleic acid accumulate in the muscle of the healthy older
17
18 subjects, whereas downstream Δ -6 desaturase product dihomo- γ -linoleic acid products are
19
20 depleted⁵⁶.
21
22

23 24 **Effect of prolonged resistance-type exercise training**

25
26 We compared the effect of prolonged resistance-type exercise training in both healthy and
27
28 frail older subjects on all measured metabolites using multilevel sPLS. There was a profound
29
30 correlation between the adaptive response to training between the transcriptome and amino
31
32 acids in the muscle metabolome (canonical correlations between 0.7 and 0.8). Particularly
33
34 high correlations were observed between expression changes of extracellular matrix genes and
35
36 amino acids. Although it is unlikely that there is a direct link between expression of these
37
38 genes and levels of these metabolites, it does imply that these changes in amino acid levels are
39
40 part of the adaptive response to resistance-type exercise training.
41
42
43

44
45 At the metabolite level, the most striking effects of resistance-type exercise training in frail
46
47 and healthy older subjects were observed for the C3 (propionyl) and C5 (isovaleryl) muscle
48
49 acylcarnitines which are derived from branched chain amino acids. After training, the amino
50
51 acid derived acylcarnitines showed a significant decrease both in the healthy and frail older
52
53 subjects, accompanied by an increase of branched chain amino acids. A likely explanation is
54
55 that the flux-determining mitochondrial branched chain α -keto acid hydrogenase (BCKDH)
56
57
58
59
60

1
2
3 complex ⁵⁷ has a compromised response to prolonged resistance-type exercise training. As is
4
5 schematically depicted in Figure 4-a, the BCKDH complex can respond to exercise via
6
7 different mechanisms. PGC-1 α is a known activator of BCKDH, but training did not have an
8
9 effect on its gene expression in our study. Exercise is known to increase NAD⁺ levels ⁴⁵ and
10
11 could thus activate PGC-1 α in a post-translational manner via SIRT1. Apparently this
12
13 mechanism is also not activated by training in the older subjects. These effects are specific for
14
15 branched chain amino acids oxidation and no significant effects on fatty acid derived
16
17 acylcarnitines were found. A decrease in branched chain amino acids oxidation may stimulate
18
19 mTOR related pathways activation and protein synthesis or attenuate muscle protein
20
21 breakdown^{58, 59}, which could be beneficial for older subjects. As this mechanism occurs at
22
23 enzymatic level, the available metabolomics and transcriptomics data can however not
24
25 confirm this hypothesis and in future studies proteomics analysis would be called for.
26
27
28
29

30 **Correlation of plasma and muscle metabolome**

31
32
33 The weak correlations between plasma and muscle metabolite levels indicate that plasma
34
35 levels only partially reflect muscle metabolism, even though skeletal muscle is one of the
36
37 largest metabolically active tissues in the human body. This seems to suggest that these
38
39 metabolites are also produced by other metabolic compartments. In a recent study, lack of
40
41 correlation between acylcarnitine levels in plasma and tissues in mice was attributed to
42
43 differences in turnover in plasma and muscle compartments, and contribution of other
44
45 compartments than muscle to plasma acylcarnitine levels ⁶⁰. The same rationale very likely
46
47 also applies to amino acids involved in collagen metabolism (proline, 4-hydroxyproline),
48
49 which can also be formed in bone. Adipose tissue is also a metabolically active compartment
50
51 with regard to branched chain amino acids besides muscle ⁶¹, and this will likely weaken
52
53 plasma-muscle level correlations. As a consequence, correlations between plasma metabolites
54
55
56
57
58
59
60

1
2
3 and muscle metabolites are too modest to support their use as direct read-outs of muscle
4
5 metabolism^{62, 63}.
6
7
8
9

10 **Conclusion**

11
12
13 The major differences in muscle metabolome of healthy older and young subjects relate to
14
15 mitochondrial function, fiber-type composition, and tissue turnover. Similar differences were
16
17 observed when comparing frail older subjects to healthy older subjects. Prolonged resistance-
18
19 type exercise training resulted in a correlative adaptive response of amino acids and genes
20
21 responsible for tissue remodeling. The effect of exercise on amino acid derived acylcarnitines
22
23 in healthy and frail older subjects points towards decreased branched chain amino acids
24
25 catabolism, likely due to attenuated activation of the flux-determining mitochondrial branched
26
27 chain α -keto acid hydrogenase complex in older subjects. Only modest correlations between
28
29 muscle metabolite and plasma levels were found, which prohibits the use of the latter as read-
30
31 outs of muscle metabolism.
32
33
34
35
36

37 **AUTHOR INFORMATION**

38 39 40 **Corresponding author**

41
42
43
44 *E mail: john.vanduyhoven@wur.nl. Phone: +31(0)317 482044.
45
46

47 **Acknowledgements**

48
49 This research was financially supported by Top Institute Food and Nutrition, Wageningen,
50
51 The Netherlands and the Netherland Metabolomics Centre (NMC), part of the Netherlands
52
53 Genomics Initiative (NGI). TI Food and Nutrition is a public-private partnership on
54
55 precompetitive research in food and nutrition. The public partners are responsible for the
56
57
58
59
60

1
2
3 study design, data collection and analysis, decision to publish, and preparation of the
4
5 manuscript. The private partners have contributed to the project through regular discussion.
6
7 The authors gratefully acknowledge the staff of the NMC Demonstration and Competency
8
9 Centre for performing the expert metabolic profiling work on plasma and muscle biopsies.
10
11 We also thank Dr.Velitchka V. Mihaleva for expert deconvolution of ^1H NMR spectra of
12
13 serum.
14
15
16
17
18

19 ASSOCIATED CONTENT

20 21 Supporting Information

22
23 **Figure S1.** Change of leg extension 1RM after resistance type exercise training. **Figure S2.**
24
25 PCA plot of metabolites detected by the organic acid platform in muscle tissue. **Figure S3.**
26
27 Group means with 95 percent confidence intervals for isoleucine and propionylcarnitine.
28
29 **Figure S4.** Pearson correlation heatmap between muscle and serum metabolites. **Table S1.**
30
31 Characteristics of subjects of which plasma samples were studied. **Table S2.** Characteristics
32
33 of the older subjects of which samples were obtained before and after training.
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

1. PD, U. N., Department of Economic and Social Affairs. Population Ageing and Development. New York. **September 2012**.
2. Frontera, W. R.; Hughes, V. A.; Fielding, R. A.; Fiatarone, M. A.; Evans, W. J.; Roubenoff, R., *Ageing of skeletal muscle: a 12-yr longitudinal study*. 2000; Vol. 88, p 1321-1326.
3. Goodpaster, B. H.; Park, S. W.; Harris, T. B.; Kritchevsky, S. B.; Nevitt, M.; Schwartz, A. V.; Simonsick, E. M.; Tyllavsky, F. A.; Visser, M.; Newman, A. B., The loss of skeletal muscle strength, mass, and quality in older adults: the health, aging and body composition study. *J Gerontol A Biol Sci Med Sci* **2006**, 61, (10), 1059-64.
4. Sayer, A. A.; Robinson, S. M.; Patel, H. P.; Shavlakadze, T.; Cooper, C.; Grounds, M. D., New horizons in the pathogenesis, diagnosis and management of sarcopenia. *Age Ageing* **2013**, 42, (2), 145-50.
5. Sayer, A. A.; Syddall, H.; Martin, H.; Patel, H.; Baylis, D.; Cooper, C., The developmental origins of sarcopenia. *J Nutr Health Aging* **2008**, 12, (7), 427-32.
6. Sayer, A. A.; Syddall, H. E.; Martin, H. J.; Dennison, E. M.; Anderson, F. H.; Cooper, C., Falls, sarcopenia, and growth in early life: findings from the Hertfordshire cohort study. *Am J Epidemiol* **2006**, 164, (7), 665-71.
7. Snijders, T.; Verdijk, L. B.; van Loon, L. J., The impact of sarcopenia and exercise training on skeletal muscle satellite cells. *Ageing Res Rev* **2009**, 8, (4), 328-38.
8. Morley, J. E., Editorial: Sarcopenia Revisited. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences* **2003**, 58, (10), M909-M910.
9. Hangelbroek, R. W. J.; Fazelzadeh, P.; Tieland, M.; Boekschoten, M. V.; Hooiveld, G. J. E. J.; van Duynhoven, J. P. M.; Timmons, J. A.; Verdijk, L. B.; van Loon, L. J. C.; de Groot, L. C. P. G. M.; Müller, M., Expression of protocadherin gamma in skeletal muscle is associated with muscle weakness and age. *In press* **2015**.
10. Houtkooper, R. H.; Argmann, C.; Houten, S. M.; Canto, C.; Jenning, E. H.; Andreux, P. A.; Thomas, C.; Doenlen, R.; Schoonjans, K.; Auwerx, J., The metabolic footprint of aging in mice. *Sci Rep* **2011**, 1, 134.
11. Garvey, S. M.; Dugle, J. E.; Kennedy, A. D.; McDunn, J. E.; Kline, W.; Guo, L.; Guttridge, D. C.; Pereira, S. L.; Edens, N. K., Metabolomic profiling reveals severe skeletal muscle group-specific perturbations of metabolism in aged FBN rats. *Biogerontology* **2014**, 15, (3), 217-32.
12. Gueugneau, M.; Coudy-Gandilhon, C.; Théron, L.; Meunier, B.; Barboiron, C.; Combaret, L.; Taillandier, D.; Polge, C.; Attaix, D.; Picard, B.; Verney, J.; Roche, F.; Féasson, L.; Barthélémy, J.-C.; Béchet, D., Skeletal Muscle Lipid Content and Oxidative Activity in Relation to Muscle Fiber Type in Aging and Metabolic Syndrome. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences* **2014**.
13. Huffman, K. M.; Koves, T. R.; Hubal, M. J.; Abouassi, H.; Beri, N.; Bateman, L. A.; Stevens, R. D.; Ilkayeva, O. R.; Hoffman, E. P.; Muoio, D. M.; Kraus, W. E., Metabolite signatures of exercise training in human skeletal muscle relate to mitochondrial remodelling and cardiometabolic fitness. *Diabetologia* **2014**, 57, (11), 2282-95.
14. Alves, R. A. M.; Dane, A.; Harms, A.; Strassburg, K.; Seifar, R.; Verdijk, L.; Kersten, S.; Berger, R.; Hankemeier, T.; Vreeken, R., Global profiling of the muscle metabolome: method optimization, validation and application to determine exercise-induced metabolic effects. *Metabolomics* **2015**, 11, (2), 271-285.
15. Tieland, M.; Dirks, M. L.; van der Zwaluw, N.; Verdijk, L. B.; van de Rest, O.; de Groot, L. C.; van Loon, L. J., Protein supplementation increases muscle mass gain during

1
2
3 prolonged resistance-type exercise training in frail elderly people: a randomized, double-
4 blind, placebo-controlled trial. *J Am Med Dir Assoc* **2012**, 13, (8), 713-9.

5 16. Tieland, M.; van de Rest, O.; Dirks, M. L.; van der Zwaluw, N.; Mensink, M.; van
6 Loon, L. J.; de Groot, L. C., Protein supplementation improves physical performance in frail
7 elderly people: a randomized, double-blind, placebo-controlled trial. *J Am Med Dir Assoc*
8 **2012**, 13, (8), 720-6.

9 17. Wall, B. T.; Hamer, H. M.; de Lange, A.; Kiskini, A.; Groen, B. B.; Senden, J. M.;
10 Gijsen, A. P.; Verdijk, L. B.; van Loon, L. J., Leucine co-ingestion improves post-prandial
11 muscle protein accretion in elderly men. *Clin Nutr* **2013**, 32, (3), 412-9.

12 18. Leenders, M.; Verdijk, L. B.; van der Hoeven, L.; van Kranenburg, J.; Nilwik, R.; van
13 Loon, L. J., Elderly men and women benefit equally from prolonged resistance-type exercise
14 training. *J Gerontol A Biol Sci Med Sci* **2013**, 68, (7), 769-79.

15 19. Gorissen, S. H.; Burd, N. A.; Hamer, H. M.; Gijsen, A. P.; Groen, B. B.; van Loon, L.
16 J., Carbohydrate coingestion delays dietary protein digestion and absorption but does not
17 modulate postprandial muscle protein accretion. *J Clin Endocrinol Metab* **2014**, 99, (6), 2250-
18 8.

19 20. Fried, L. P.; Tangen, C. M.; Walston, J.; Newman, A. B.; Hirsch, C.; Gottdiener, J.;
20 Seeman, T.; Tracy, R.; Kop, W. J.; Burke, G.; McBurnie, M. A., Frailty in Older Adults:
21 Evidence for a Phenotype. *The Journals of Gerontology Series A: Biological Sciences and*
22 *Medical Sciences* **2001**, 56, (3), M146-M157.

23 21. Leenders, M.; Verdijk, L. B.; Van der Hoeven, L.; Van Kranenburg, J.; Nilwik, R.;
24 Wodzig, W. K.; Senden, J. M.; Keizer, H. A.; Van Loon, L. J., Protein supplementation
25 during resistance-type exercise training in the elderly. *Med Sci Sports Exerc* **2013**, 45, (3),
26 542-52.

27 22. Noga, M. J.; Dane, A.; Shi, S.; Attali, A.; van Aken, H.; Suidgeest, E.; Tuinstra, T.;
28 Muilwijk, B.; Coulier, L.; Luiders, T.; Reijmers, T. H.; Vreeken, R. J.; Hankemeier, T.,
29 Metabolomics of cerebrospinal fluid reveals changes in the central nervous system
30 metabolism in a rat model of multiple sclerosis. *Metabolomics* **2012**, 8, (2), 253-263.

31 23. van der Kloet, F. M.; Bobeldijk, I.; Verheij, E. R.; Jellema, R. H., Analytical Error
32 Reduction Using Single Point Calibration for Accurate and Precise Metabolomic
33 Phenotyping. *Journal of Proteome Research* **2009**, 8, (11), 5132-5141.

34 24. Mihaleva, V. V.; Korhonen, S. P.; van Duynhoven, J.; Niemitz, M.; Vervoort, J.;
35 Jacobs, D. M., Automated quantum mechanical total line shape fitting model for quantitative
36 NMR-based profiling of human serum metabolites. *Anal Bioanal Chem* **2014**, 406, (13),
37 3091-102.

38 25. Szasz, G., Reaction-rate method for gamma-glutamyltransferase activity in serum.
39 *Clin Chem* **1976**, 22, (12), 2051-5.

40 26. Wold, H., *Path models with latent variables: The NIPALS approach*. Acad. Press:
41 1975.

42 27. Lê Cao, K.-A.; González, I.; Déjean, S., integrOmics: an R package to unravel
43 relationships between two omics datasets. *Bioinformatics* **2009**, 25, (21), 2855-2856.

44 28. Sébastien Lê, J. J., François Husson, FactoMineR: An R Package for Multivariate
45 Analysis. *Journal of Statistical Software* **2008**.

46 29. Smilde, A. K.; Jansen, J. J.; Hoefsloot, H. C. J.; Lamers, R.-J. A. N.; van der Greef, J.;
47 Timmerman, M. E., ANOVA-simultaneous component analysis (ASCA): a new tool for
48 analyzing designed metabolomics data. *Bioinformatics* **2005**, 21, (13), 3043-3048.

49 30. Berger, A.; Milgram, E.; Mitchell, M.; Lawton, K.; Hanson, R.; Kalhan, S.; Milburn,
50 M., The metabolomics of aging. *Faseb Journal* **2007**, 21, (6), A1040-A1040.

- 1
2
3 31. Lawton, K. A.; Berger, A.; Mitchell, M.; Milgram, K. E.; Evans, A. M.; Guo, L.;
4 Hanson, R. W.; Kalhan, S. C.; Ryals, J. A.; Milburn, M. V., Analysis of the adult human
5 plasma metabolome. *Pharmacogenomics* **2008**, 9, (4), 383-397.
- 6 32. Laferrere, B.; Reilly, D.; Arias, S.; Swerdlow, N.; Gorroochurn, P.; Bawa, B.; Bose,
7 M.; Teixeira, J.; Stevens, R. D.; Wenner, B. R.; Bain, J. R.; Muehlbauer, M. J.; Haqq, A.;
8 Lien, L.; Shah, S. H.; Svetkey, L. P.; Newgard, C. B., Differential metabolic impact of gastric
9 bypass surgery versus dietary intervention in obese diabetic subjects despite identical weight
10 loss. *Sci Transl Med* **2011**, 3, (80), 80re2.
- 11 33. Saccenti, E.; Hoefsloot, H. J.; Smilde, A.; Westerhuis, J.; Hendriks, M. W. B.,
12 Reflections on univariate and multivariate analysis of metabolomics data. *Metabolomics*
13 **2014**, 10, (3), 361-374.
- 14 34. Balagopal, P.; Rooyackers, O. E.; Adey, D. B.; Ades, P. A.; Nair, K. S., Effects of
15 aging on in vivo synthesis of skeletal muscle myosin heavy-chain and sarcoplasmic protein in
16 humans. *Am J Physiol* **1997**, 273, (4 Pt 1), E790-800.
- 17 35. Short, K. R.; Vittone, J. L.; Bigelow, M. L.; Proctor, D. N.; Nair, K. S., Age and
18 aerobic exercise training effects on whole body and muscle protein metabolism. *Am J Physiol*
19 *Endocrinol Metab* **2004**, 286, (1), E92-101.
- 20 36. Yarasheski, K. E.; Welle, S.; Nair, K. S., Muscle protein synthesis in younger and
21 older men. *JAMA* **2002**, 287, (3), 317-8.
- 22 37. Yarasheski, K. E.; Zachwieja, J. J.; Bier, D. M., Acute effects of resistance exercise on
23 muscle protein synthesis rate in young and elderly men and women. *Am J Physiol* **1993**, 265,
24 (2 Pt 1), E210-4.
- 25 38. Rooyackers, O. E.; Adey, D. B.; Ades, P. A.; Nair, K. S., Effect of age on in vivo rates
26 of mitochondrial protein synthesis in human skeletal muscle. *Proc Natl Acad Sci U S A* **1996**,
27 93, (26), 15364-9.
- 28 39. Kjaer, M., Role of extracellular matrix in adaptation of tendon and skeletal muscle to
29 mechanical loading. *Physiol Rev* **2004**, 84, (2), 649-98.
- 30 40. Gosselin, L. E., Skeletal Muscle Collagen: Age, Injury and Disease. In *Sarcopenia-
31 Age-Related Muscle Wasting and Weakness*, Springer: 2011; pp 159-172.
- 32 41. Morris, S. M., Jr., Arginine metabolism: boundaries of our knowledge. *J Nutr* **2007**,
33 137, (6 Suppl 2), 1602s-1609s.
- 34 42. Roberts, L. D.; Bostrom, P.; O'Sullivan, J. F.; Schinzel, R. T.; Lewis, G. D.; Dejam,
35 A.; Lee, Y. K.; Palma, M. J.; Calhoun, S.; Georgiadi, A.; Chen, M. H.; Ramachandran, V. S.;
36 Larson, M. G.; Bouchard, C.; Rankinen, T.; Souza, A. L.; Clish, C. B.; Wang, T. J.; Estall, J.
37 L.; Soukas, A. A.; Cowan, C. A.; Spiegelman, B. M.; Gerszten, R. E., beta-Aminoisobutyric
38 acid induces browning of white fat and hepatic beta-oxidation and is inversely correlated with
39 cardiometabolic risk factors. *Cell Metab* **2014**, 19, (1), 96-108.
- 40 43. Gomes, A. P.; Price, N. L.; Ling, A. J.; Moslehi, J. J.; Montgomery, M. K.; Rajman,
41 L.; White, J. P.; Teodoro, J. S.; Wrann, C. D.; Hubbard, B. P.; Mercken, E. M.; Palmeira, C.
42 M.; de Cabo, R.; Rolo, A. P.; Turner, N.; Bell, E. L.; Sinclair, D. A., Declining NAD(+) induces
43 a pseudohypoxic state disrupting nuclear-mitochondrial communication during aging.
44 *Cell* **2013**, 155, (7), 1624-38.
- 45 44. Imai, S.; Guarente, L., NAD⁺ and sirtuins in aging and disease. *Trends Cell Biol* **2014**,
46 24, (8), 464-71.
- 47 45. Overmyer, K. A.; Evans, C. R.; Qi, N. R.; Minogue, C. E.; Carson, J. J.; Chermiside-
48 Scabbo, C. J.; Koch, L. G.; Britton, S. L.; Pagliarini, D. J.; Coon, J. J.; Burant, C. F., Maximal
49 oxidative capacity during exercise is associated with skeletal muscle fuel selection and
50 dynamic changes in mitochondrial protein acetylation. *Cell Metab* **2015**, 21, (3), 468-78.
- 51 46. Pegg, A. E.; Mccann, P. P., Polyamine Metabolism and Function. *American Journal of*
52 *Physiology* **1982**, 243, (5), C212-C221.
- 53
54
55
56
57
58
59
60

- 1
2
3 47. Turchanowa, L.; Rogozkin, V. A.; Milovic, V.; Feldkoren, B. I.; Caspary, W. F.;
4 Stein, J., Influence of physical exercise on polyamine synthesis in the rat skeletal muscle.
5 *European Journal of Clinical Investigation* **2000**, *30*, (1), 72-78.
- 6 48. Lee, N. K. L.; Maclean, H. E., Polyamines, Androgens, and Skeletal Muscle
7 Hypertrophy. *Journal of Cellular Physiology* **2011**, *226*, (6), 1453-1460.
- 8 49. Kaminska, A. M.; Stern, L. Z.; Russell, D. H., Altered Muscle Polyamine Levels in
9 Human Neuromuscular Diseases. *Annals of Neurology* **1981**, *9*, (6), 605-607.
- 10 50. Kaminska, A. M.; Stern, L. Z.; Russell, D. H., Polyamine Accumulation in Normal
11 and Denervated Neonatal Muscle. *Experimental Neurology* **1981**, *72*, (3), 612-618.
- 12 51. Bongers, K. S.; Fox, D. K.; Kunkel, S. D.; Stebounova, L. V.; Murry, D. J.; Pufall, M.
13 A.; Ebert, S. M.; Dyle, M. C.; Bullard, S. A.; Dierdorff, J. M.; Adams, C. M., Spermine
14 oxidase maintains basal skeletal muscle gene expression and fiber size and is strongly
15 repressed by conditions that cause skeletal muscle atrophy. *Am J Physiol Endocrinol Metab*
16 **2015**, *308*, (2), E144-58.
- 17 52. Ost, M.; Keipert, S.; van Schothorst, E. M.; Donner, V.; van der Stelt, I.; Kipp, A. P.;
18 Petzke, K. J.; Jove, M.; Pamplona, R.; Portero-Otin, M.; Keijer, J.; Klaus, S., Muscle
19 mitohormesis promotes cellular survival via serine/glycine pathway flux. *Faseb j* **2015**, *29*,
20 (4), 1314-28.
- 21 53. Sale, C.; Artioli, G. G.; Gualano, B.; Saunders, B.; Hobson, R. M.; Harris, R. C.,
22 Carnosine: from exercise performance to health. *Amino Acids* **2013**, *44*, (6), 1477-91.
- 23 54. Boldyrev, A. A.; Aldini, G.; Derave, W., Physiology and pathophysiology of
24 carnosine. *Physiol Rev* **2013**, *93*, (4), 1803-45.
- 25 55. Harris, R. C.; Wise, J. A.; Price, K. A.; Kim, H. J.; Kim, C. K.; Sale, C., Determinants
26 of muscle carnosine content. *Amino Acids* **2012**, *43*, (1), 5-12.
- 27 56. Kalish, B. T.; Fallon, E. M.; Puder, M., A tutorial on fatty acid biology. *JPEN J*
28 *Parenter Enteral Nutr* **2012**, *36*, (4), 380-8.
- 29 57. Shimomura, Y.; Murakami, T.; Nakai, N.; Nagasaki, M.; Harris, R. A., Exercise
30 promotes BCAA catabolism: effects of BCAA supplementation on skeletal muscle during
31 exercise. *J Nutr* **2004**, *134*, (6 Suppl), 1583s-1587s.
- 32 58. Um, S. H.; D'Alessio, D.; Thomas, G., Nutrient overload, insulin resistance, and
33 ribosomal protein S6 kinase 1, S6K1. *Cell Metab* **2006**, *3*, (6), 393-402.
- 34 59. Newgard, C. B., Interplay between lipids and branched-chain amino acids in
35 development of insulin resistance. *Cell Metab* **2012**, *15*, (5), 606-14.
- 36 60. Schooneman, M. G.; Achterkamp, N.; Argmann, C. A.; Soeters, M. R.; Houten, S. M.,
37 Plasma acylcarnitines inadequately reflect tissue acylcarnitine metabolism. *Biochim Biophys*
38 *Acta* **2014**, *1841*, (7), 987-94.
- 39 61. Herman, M. A.; She, P.; Peroni, O. D.; Lynch, C. J.; Kahn, B. B., Adipose tissue
40 branched chain amino acid (BCAA) metabolism modulates circulating BCAA levels. *J Biol*
41 *Chem* **2010**, *285*, (15), 11348-56.
- 42 62. Brosnan, J. T., Glutamate, at the Interface between Amino Acid and Carbohydrate
43 Metabolism. *The Journal of Nutrition* **2000**, *130*, (4), 988.
- 44 63. van de Poll, M. C.; Siroen, M. P.; van Leeuwen, P. A.; Soeters, P. B.; Melis, G. C.;
45 Boelens, P. G.; Deutz, N. E.; Dejong, C. H., Interorgan amino acid exchange in humans:
46 consequences for arginine and citrulline metabolism. *The American Journal of Clinical*
47 *Nutrition* **2007**, *85*, (1), 167-172.
- 48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

TABLES

Table 1. Characteristics of subjects of which skeletal muscle tissue biopsies were studied

	Young	Healthy older	Frail older
N (male / female)	30/0	47/19	25/18
Age (years)	21.7 ± 2.5	71.7 ± 5.2	77.5 ± 8.0
Height (m)	1.83 ± 0.06	1.72 ± 0.08	1.67 ± 0.09
Weight (kg)	76.7 ± 11.8	75.9 ± 13.3	77.5 ± 11.1
BMI ^a (kg / m ²)	22.6 ± 2.7	25.5 ± 3.0	27.5 ± 3.7
Body Fat (%)	14.9 ± 4.9	24.5 ± 5.6	32.1 ± 8.8

Data was presented as mean ± SD. a: body mass index,

Table 2. Muscle metabolites that are significantly different between healthy older and young males ^a

Metabolite	<i>P</i> -value	FC ^b (Older/Young)
<i>TCA Cycle</i>		
Succinic acid	0.02	0.76
2-ketoglutaric acid	0.03	0.76
Fumaric acid	0.04	0.82
Lactic acid	0.05	0.69
<i>Energy</i>		
ATP	<0.01	0.75
ADP	0.01	0.88
<i>Branched chain amino acids</i>		
Valine	<0.01	0.81
Leucine	<0.01	0.81
Isoleucine	0.03	0.84
<i>Acylcarnitines</i>		
Acetylcarnitine (C2)	<0.01	0.49
Malonylcarnitine (C3-DC)	<0.01	0.46
<i>Intracellular buffering</i>		
Carnosine	<0.01	0.7
<i>Arginine, Proline Pathway</i>		
Ornithine	<0.01	1.55
Arginine	<0.01	1.34
4-hydroxy-proline	0.01	0.69
Proline	0.02	0.84
Glycylglycine	0.05	0.87
Methionine	<0.01	0.8
<i>Other amino acids</i>		
Lysine	<0.01	1.44
Aspartic acid	<0.01	1.45
<i>Oxylipins</i>		
LA (LOX)		
9-HODE	<0.01	1.4
13-HODE	0.01	1.37
13-KODE	0.03	1.3
LA(CYP450)		
9,10-EpOME	0.04	1.39
ALA (LOX)		
9-HOTrE	0.01	1.71
DGLA (LOX)		
15S-HETrE	0.02	0.83
8-HETrE	0.03	0.85
<i>Aminobutyric acids</i>		

β amino isobutyric acid	<0.01	1.61
α -aminobutyric acid	0.01	0.82

a: Metabolites are presented that significantly (P -value ≤ 0.05) differ between young and old subjects according to univariate ANOVA models. b: Fold Change

Table 3. Muscle metabolites that are significantly different between frail and healthy older subjects.^a

Metabolite	Group	Sex	Interaction	FC ^b (Frail/Healthy)
<i>TCA Cycle</i>				
Citric acid	<0.01	NS ^c	NS	0.54
<i>Acylcarnitines</i>				
Isovalerylcarnitine (C5)	<0.01	NS	NS	0.42
Octenoylcarnitine (C8)	0.03	0.03	NS	0.77
Malonylcarnitine(C3-DC)	0.02	NS	NS	0.77
Carnitine (C0)	0.01	NS	NS	0.75
<i>Intracellular buffering</i>				
Carnosine	0.01	NS	NS	0.8
<i>Oxylipins</i>				
LA (CYP450)				
12,13DiHOME	0.04	NS	0.03	1.18
DGLA (LOX)				
8HETrE	0.03	NS	NS	0.81
15SHETrE	<0.01	NS	NS	0.77
<i>Polyamine metabolism</i>				
Spermidine	0.01	0.02	NS	1.24
Spermine	0.04	NS	NS	0.9
<i>Other amino acids</i>				
Histidine	<0.01	NS	NS	0.79
Asparagine	0.01	NS	NS	0.81
Taurine	0.01	NS	NS	0.79
Serine	0.01	NS	NS	0.86
Glycine	0.02	NS	NS	0.81
oacetylserine	0.02	NS	NS	0.9
Homoserine	0.02	NS	NS	0.85
Tyrosine	0.02	NS	NS	0.83
Tryptophan	0.02	0.04	NS	0.83
Methionine	0.02	NS	NS	0.83
Glutamine	0.03	NS	NS	0.82
Pyroglutamic acid	0.03	NS	NS	0.83

Glutamic acid	0.04	NS	NS	0.82
Glycylglycine	<0.01	NS	NS	0.77
<hr/>				
<i>Aminobutyric acids</i>				
β amino isobutyric acid	0.05	NS	NS	0.79

a: Metabolites are presented that significantly (P -value ≤ 0.05) differ between frail and healthy older subjects according to a univariate ANOVA models that accounted for Group effect (Frail vs. Healthy), Sex effect and Group-Sex interaction; Significance of Group (Frail vs. Healthy older subjects) effects and Group-Sex interactions has been indicated; b :fold change; c: not significant

Table 4. Training effect on muscle metabolites in healthy older subjects ^a

Metabolite	Training	FC(post/pre training) ^b
Pipecolic acid	0.002	1.64
Isovalerylcarnitine (C5)	0.005	0.56
Linoleylcarnitine (C18:2)	0.01	0.61
Oleylcarnitine (C18:1)	0.01	0.7
Propionylcarnitine (C3)	0.01	0.73
Palmitoylcarnitine (C16)	0.02	0.75
11.12.EpETrE	0.03	1.26
Tetradecenoylcarnitine (C14:1)	0.03	0.47
AAAC ^c	0.02	0.77
FAAC ^d	NS	0.96

a: Metabolites are presented that significantly (P -value ≤ 0.05) differ pre- and post-training in healthy older subjects according to univariate linear mixed models. We note that we constructed also linear mixed models that account for other factors and their interactions, but these were not found to be significant b: fold change, c,d: Single Value Decomposition (SVD) were calculated for amino acid and fatty acid acylcarnitines, denoted as AAAC and FAAC, respectively.

Table 5. Training effect on muscle metabolites in frail older subjects ^a

Metabolite	Training	Supplement	Sex	Interaction	FC(post/pre training) ^b
Propionylcarnitine (C3)	<0.01	NS ^c	NS	NS	0.75
Glucose	<0.01	NS	<0.01	NS	1.35
Lactic acid	0.01	NS	<0.01	NS	1.55
Tetradecenoylcarnitine (C14:1)	0.03	NS	NS	NS	2.00
Methionine	0.04	<0.01	NS	NS	1.22
Tryptophan	0.04	NS	NS	NS	1.21
β Alanine	0.05	NS	NS	0.01	0.92
Isoleucine	0.05	<0.01	0.02	NS	1.21
Myristoylcarnitine (C14)	0.05	NS	NS	NS	1.73
AAAC ^d	0.01	NS	NS	NS	0.76
FAAC ^e	NS	NS	NS	NS	1.95

a: Metabolites are presented that significantly (P -value ≤ 0.05) differ pre- and post-training in frail older subjects according to univariate linear mixed models that account for Supplement, Sex and (Training and Supplement) Interaction. We note that also linear mixed models were constructed that account for other interactions, but these were not found to be significant; b: fold change, c: Not Significant, d & e: Single Value Decomposition (SVD) were calculated for amino acid and fatty acid acylcarnitines (AAAC and FAAC, respectively).

FIGURE LEGENDS

Figure 1. Principal component analysis (PCA) plot of biogenic amines detected in muscle biopsies. To visualize whether groups are significantly different from each other, confidence ellipses (95% Confidence Interval) were drawn around them.

Figure 2. Correlation network of muscle metabolites and genes. Only significantly changed genes were selected (FDR<0.05). Metabolite canonical correlation cutoff ≤ 0.80 . Circle: gene, rectangle: metabolite, Green : positive correlation.

Figure 3. Pearson correlation heatmap of muscle to muscle metabolites (a), plasma to plasma metabolites (b), muscle to plasma metabolites (c). (d) Pearson correlation network of the most strongly correlated muscle and plasma metabolites. Red and blue indicate positive and negative correlations, respectively. Thick lines: correlation ~ 0.5 , thin lines: $0.3 < \text{correlation} < 0.5$. Pink nodes: muscle tissue (T) metabolites and yellow nodes: plasma (P) metabolites,

Figure 4. (a) Schematic representation of mitochondrial oxidation of free fatty acids (FFA) and branched chain amino acids (BCAA). Arrows indicate effect of prolonged resistance-type exercise training on older subjects (healthy and frail) on BCAA (increase) and acylcarnitines (decrease) as well as the proposed (dashed-arrows) downregulation of the branched chain α -keto acid hydrogenase (BCKDH) complex. (b) Schematic representation of age-related NAD^+ dependent acetylation of PGC1 α . Dashed arrows indicate NAD^+ and SIRT1 dependent downregulation of PGC-1 α and β -isoamino butyric acid (BAIBA).

Figure 1.






Figure 2.



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 3.








Figure 4.



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

