

1
2
3
4
5 The latent structure of Acute Stress Disorder symptoms in trauma-exposed children and adolescents.
6

7 Anna McKinnon¹ PhD

8 Macquarie University

9
10 Richard Meiser-Stedman PhD

11 University of East Anglia

12 Peter Watson PhD

13 Medical Research Council Cognition and Brain Sciences Unit

14 Clare Dixon MSc

15 University of Bath

16 Nancy Kassam-Adams PhD

17 The Children's Hospital of Philadelphia

18 Anke Ehlers PhD

19 University of Oxford

20 Flaura Winston PhD

21 The Children's Hospital of Philadelphia

22 Patrick Smith PhD

23 Kings College London

24 William Yule PhD

25 Kings College London

26 Tim Dalgleish PhD

27 Medical Research Council Cognition and Brain Sciences Unit

28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47 Address of work completion: Medical Research Council Cognition and Brain Sciences Unit

48
49
50
51 Keywords: Acute Stress Disorder, DSM-5, factor analysis, children, Post-traumatic Stress Disorder
52 Text word count²: 6684 words; document word count: 7132 words
53

54
55 ¹ Corresponding author: Dr Anna McKinnon; work address: Centre for Emotional Health, Department
56 of Psychology, Macquarie University, NSW 2109, Australia; ph: +61 (0)2 9850 9163; fax: +61 (02)
57 9850 9911; email: anna.mckinnon@mq.edu.au.
58
59
60

Abstract

Background. The revision of Acute Stress Disorder (ASD) in the DSM-5 (*DSM-5*, 2013) proposes a cluster-free model of ASD symptoms in both adults and youth. Published evaluations of competing models of ASD clustering in youth have rarely been examined. **Methods** We used Confirmatory Factor Analysis (combined with multi-group invariance tests) to explore the latent structure of ASD symptoms in a trauma-exposed sample of children and young people (N=594). The DSM-5 structure was compared with the previous DSM-IV conceptualisation (4-factor), and two alternative models proposed in the literature (3-factor; 5-factor). Model fit was examined using goodness-of fit-indices. We also established DSM-5 ASD prevalence rates relative to DSM-IV ASD, and the ability of these models to classify children impaired by their symptoms. **Results.** Based on both the Bayesian Information Criterion (BIC), the inter-factor correlations and invariance testing, the 3-factor model best accounted for the profile of ASD symptoms. DSM-5 ASD led to slightly higher prevalence rates than DSM-IV ASD and performed similarly to DSM-IV with respect to categorising children impaired by their symptoms. Modifying the DSM-5 ASD algorithm to a 3+ or 4+ symptom requirement was the strongest predictor of impairment. **Conclusions.** These findings suggest that a uni-factorial general distress model is not the optimal model of capturing the latent structure of ASD symptom profiles in youth and that modifying the current DSM-5 9+ symptom algorithm could potentially lead to a more developmentally sensitive conceptualisation.

Introduction

While it is common for children to display stress responses in the first few weeks following a trauma, only a minority will go on to develop Posttraumatic Stress Disorder (PTSD) (Bryant et al., 2004; Dalgleish et al., 2008; Kassam-Adams & Winston, 2004; Meiser-Stedman et al., 2008). This has prompted debate about the clinical utility of classifying these early post-trauma responses using an Acute Stress Disorder (ASD) diagnosis (Bryant, 2011). First introduced in the DSM-IV (American Psychiatric Association [APA], 1994) to identify individuals likely to develop subsequent PTSD (only diagnosable 1-month post-trauma), symptoms of ASD were arranged into re-experiencing, avoidance, and arousal clusters, mirroring the structure of DSM-IV PTSD closely. DSM-IV ASD also included a dissociation cluster, based on research indicating that dissociation was a prognostic indicator of later pathology (Bryant et al., 2011). This DSM-IV conceptualisation operationalises the idea that the distinctive symptom clusters of later PTSD actually develop within the first two weeks of a trauma, and, in addition, that dissociation is a distinctive feature of those that will later develop chronic PTSD. However, this ASD model has poor predictive validity, and the overly restrictive nature of the dissociation criterion means that adults and children can miss out on an ASD diagnosis despite having clinically significant problems in other symptom domains (Harvey & Bryant, 1998; Meiser-Stedman, Yule, Smith, Glucksman, & Dalgleish, 2005). Six confirmatory factor analytic tests (Armour, Shevlin, & Elkit, 2011; Brooks et al., 2008; Hansen, Armour, & Elklit, 2012; Hansen, Lasgaard, & Elklit, 2013; Wang, Li, Shi, Zhang, & Shen, 2010) carried out in adults have evaluated the DSM-IV structure (see Table 1) of ASD compared to alternative conceptualisations derived from the PTSD literature (with the addition of a dissociation criterion), showing it is the best fitting model in only two of these six studies (Brooks et al., 2008; Hansen et al., 2013) (Refer to Supplementary Table S1 for a summary of study findings).

Based on these sets of findings, there was a radical overhaul of ASD in the DSM-5 (APA, 2013). Along with minor symptom changes and the revision of Criterion A describing the traumatic event, the DSM-IV requirement of at least one symptom from a set of distinct symptom clusters was removed (although 5 distinct clusters were retained as an organising principal for the different symptoms). This addressed the potentially unhelpful requirement for dissociation to be mandatory for a positive diagnosis. Instead, for a diagnosis of ASD, the DSM-5 specifies experiencing nine

1
2
3
4 symptoms from a single list of 14 (down from the DSM-IV list of 17 symptoms) to be present. The
5
6 DSM-5 ASD thus posits a unifactorial 'general distress model'. The rationale behind this change was
7
8 that no compelling support for any given model of the ASD symptom structure had clearly emerged
9
10 from factor analytic investigations and consequently the structure of early traumatic stress responses
11
12 was considered to be too variable for distinct and reliable clustering patterns to occur (Bryant et al.,
13
14 2011). The advantage of a 'general distress' model, it was proposed, is the potential to better identify
15
16 individuals at risk of developing a *range* of psychological disorders in the future, not just PTSD (Bryant
17
18 et al., 2011).
19

20 The removal of diagnostic clustering in ASD was a bold move. There are significant merits to
21
22 elucidating a valid and reliable ASD clustering system. First, there are concerns that ASD does not
23
24 represent a distinct disorder from PTSD (Brewin, Andrews, & Rose, 2003). Examining the validity of
25
26 putative ASD symptom clusters, in particular the unique dissociation symptoms, is critical to the case
27
28 for ASD as a distinct disorder. This argument extends beyond ASD and PTSD. Many of the symptoms
29
30 of both disorders overlap with other mood and anxiety syndromes calling some to question the validity
31
32 of stress disorders as a separable diagnostic entity (Brady, Killeen, Brewerton, & Lucerini, 2000).
33
34 Evaluating the validity of symptom clusters unique to the stress disorders (notably the re-experiencing
35
36 symptoms) is critical in establishing the validity of this class of disorder. Secondly, elucidating
37
38 potential symptom clusters can improve our understanding of the aetiology and maintenance of ASD
39
40 and of later PTSD, by identifying risk factors for specific clusters/factors of symptoms, and revealing
41
42 relationships between ASD factors and ongoing impairment, including later PTSD. Third, there may
43
44 be underlying cognitive or biological mechanisms mechanisms that only relate to a particular group of
45
46 symptoms and clustering allows these to be better identified (Frances & Widiger, 2012). Such
47
48 relationships might not be detectable if clustering were removed, making it difficult to refine the
49
50 specific pathways necessary for the development of treatment models that target these underlying
51
52 processes. Fourth, clusters make it easier for new theoretical approaches such as the Research
53
54 Domains Criteria (RDoC; Insel et al., 2010) and other trans-diagnostic approaches to evaluate
55
56 whether the psychological constructs/outcomes (e.g., intrusive memories) they aim to evaluate are
57
58 consistent or different across disparate disorders. Finally, a cluster-free diagnostic algorithm has
59
60 potential implications for the prevalence rates of the disorder, relative to a cluster-based algorithm

1
2
3
4 (Magruder & Calderone, 2000; Mullins-Sweatt, Lengel, & DeShong, 2016), with important implications
5
6 for the provision of clinical care. It may be that particular clusters have a stronger relationship to
7
8 clinical impairment and a cluster-free approach impedes greater understanding of such potential
9 relationships.

10
11
12 The body of empirical, modelling and theoretical work around the diagnosis of ASD has to
13 date focused almost entirely on the conceptualisation of ASD in adults with little corresponding
14 analysis of the diagnosis in youth. Reflecting this, the DSM-5 draws no distinctions across the age
15 range in terms of ASD. However, there are several putative issues concerning the application of the
16 DSM-5 ASD model to youth. Firstly, the model contains no developmental adaptations in contrast to
17 the DSM-5 conceptualisation of PTSD and to a growing body of work suggesting that stress disorders
18 present differently in adults relative to not just young children but also older children and adolescents
19 (Meiser-Stedman et al., 2008). In support of this, research indicates the tendency to express
20 individual PTSD symptoms is impacted by age (Chen, Lin, Tseng, & Wu, 2002). Secondly, question
21 marks have been raised about whether the ≥ 9 symptom requirement would have greater clinical utility
22 by reducing the requirement to 3-4 symptoms (Kassam-Adams et al., 2012). Thirdly, unlike in adults,
23 there has been almost no examination of the factor structure of ASD symptoms in youth with the
24 exception of a study of older Filipino adolescents exposed to a flash flood, which supported the 5-
25 factor dysphoria model (Mordeno & Cue, 2015). Taken together, the case against a clustered
26 approach is less clear for traumatised youth and a proper evaluation of symptom clustering
27 immediately post-trauma in younger populations is required.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

43 Given this, the present study had four inter-related aims. The first was to carry out a CFA of
44 ASD symptoms in youth comparing the unifactorial, cluster-free DSM-5 approach with other models
45 derived from the adult field (Table 1). To do this, we carried out a CFA of the 17 DSM-IV ASD
46 symptoms (these include the 14 DSM-5 symptoms but allow us to also test the DSM-IV model) in
47 youth aged 6-17 years, who had experienced a discrete one-off trauma, using pooled data from 4
48 sites. Data-pooling is advantageous over single site CFA's and standard review approaches based on
49 mean data (e.g., meta-analysis, systematic reviews) as robust statistical techniques can be used to
50 assess the potential influence of moderators (i.e., recruitment site, trauma type) of factor structure.
51
52
53
54
55
56

57 Secondly, we compared ASD prevalence rates in our sample for the current DSM-5 diagnosis relative
58
59
60

1
2
3
4 to the previous DSM-IV diagnosis. In youth, between 5% and 25% of children and young people
5
6 suffer from ASD according to DSM-IV (Bryant et al., 2004; Dalgleish et al., 2008; Kassam-Adams &
7
8 Winston, 2004; Meiser-Stedman et al., 2008). Preliminary examinations of the DSM-5 algorithm in
9
10 youth suggests that the new DSM-5 model would lead to substantially lower prevalence in children
11
12 (Kassam-Adams et al., 2012), but this finding must be replicated. Thirdly, we investigated whether the
13
14 unifactorial 'general distress' model was a better predictor of clinical impairment relative to the DSM-
15
16 IV model. Finally, following from the work of Kassam-Adams et al. (2012), we investigated whether
17
18 the DSM-5 requirement of 9+ symptoms was the best predictor of impairment relative to lower
19
20 symptom thresholds.

21 Method

22
23
24 **Sample:** Data for 594 children and young people who had been exposed to a Criterion A discrete
25
26 stressor within the previous 4 weeks (Mean age=12.55 years, SD=2.99, Range=6-17 years) were
27
28 pooled from centres in East Anglia (EA) (n=189, 8-16 years), Oxford (n=65, 6-17 years), London
29
30 (Sample 1: n=59, 7-10 yrs; Sample 2: n=92, 10-16 years) and Philadelphia (n=189, 6-17 years).
31
32 Children were recruited from either an inpatient setting or emergency department. Each study
33
34 received ethics approval from the local recruitment site. Informed consent was obtained from adult
35
36 carers/parents and assent from young people.

37
38 With the exception of EA, details of recruitment flow are published elsewhere (Bryant et al., 2004;
39
40 Dalgleish et al., 2008; Kassam-Adams & Winston, 2004; Meiser-Stedman et al., 2008). In EA the
41
42 inclusion criteria were in line with the other centres as follows: any event that involved the threat of
43
44 death, severe injury, or threat to bodily integrity, or witnessing such an event (typically road traffic
45
46 collisions [RTCs] and assaults). The exclusion criteria were: intellectual disability; assaults by the
47
48 young person's caregiver or close relative; being unconscious for >15 minutes; not being fluent in
49
50 English; ongoing exposure to threat; history of organic brain damage; and significant risk of self-harm.
51
52 A member of the clinical care team at the hospital identified cases from medical records and invited
53
54 families to participate by letter (opt-out consent).

55
56 The characteristics of the final sample are presented in Table 2. The Oxford and Philadelphia
57
58 samples recruited children and young people involved in RTC's. The EA sample included victims of
59
60

1
2
3
4 assaults¹, RTC's and accidental injuries and the London sample comprised assault and RTC victims.

5
6 The majority of the final sample had experienced RTC's (n=441), followed by assaults (n=87) and

7
8 then accidental injuries (n=66). The proportions of children endorsing at least one symptom from
9 individual clusters ranged from 52% for the avoidance cluster to 76% for the dissociation cluster.

10
11 **Measures:** In line with previous approaches (Meiser-Stedman et al., 2008), we pooled data across
12 different DSM-IV ASD instruments. DSM-IV ASD symptoms were indexed using well-validated
13 measures administered 2-4 weeks post-trauma². Measures (described for each site below) were
14 obtained via home interviews except at the EA site where measures were obtained over the phone.

15
16 *Philadelphia.* The Children's Acute Stress Questionnaire (CASQ; Kassam-Adams, 2006) is a 25-
17 item self-report instrument with good internal consistency ($\alpha=.86$), test-retest reliability ($r=.76$ for the
18 total subscale and $r=.59-.68$ for individual subscales), and convergent validity ($r=.77$ with the Child
19 and Adolescent Trauma Symptom Scale; March, 1999).

20
21 All UK sites used structured clinical interviews to assess symptoms. As there are no validated
22 interviews of ASD in children and young people, researchers added developmentally appropriate
23 dissociation items derived from the adult ASD literature to existing PTSD interviews, in line with
24 previous studies (Bryant et al., 2004; Dalgleish et al., 2008; Kassam-Adams & Winston, 2004; Meiser-
25 Stedman et al., 2008).

26
27 *EA.* The Children's Posttraumatic Stress Disorders Inventory (CPTSD-I; Saigh et al., 2000) is a
28 structured interview for assessing PTSD in children and adolescents that has excellent internal
29 consistency, test-retest reliability, and inter-rater reliability (Saigh et al., 2000; Yasik et al., 2001).

30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46 ³ The traumas that precipitated the development of ASD needed to be one-off in nature, for example
47 one-off assaults as opposed to repeated domestic violence or physical abuse. This was either
48 determined via the initial screening carried out with parents in the hospital or via the examination of
49 medical records.

50
51 ² Data collected across all four sites captured the full range of DSM-IV and DSM-5 ASD symptoms.
52 Changes to ASD in DSM-5 are as follows: DSM-IV B1 'emotional numbing' has been refined to
53 describe absence of positive emotional experiences only in DSM-5 (B5); DSM-IV B2 'loss of
54 awareness', B3 'depersonalization' and B4 'derealization' are grouped under DSM-5 B6 'altered
55 sense of reality', and; DSM-IV symptoms c) psychological distress to reminders and e) physiological
56 reactivity are grouped under DSM-5 B4.

1
2
3
4 *London.* Sample 1 ($n=92$ cases) completed The Anxiety Disorders Interview Schedule - Child
5
6 Version (ADIS-C; Silverman & Albano, 1996). The ADIS-C contains 27 items measuring symptoms of
7
8 PTSD in addition to a single item assessing impairment. Sample 2 ($n=59$) completed the Clinician
9 Administered PTSD Scale—Child and Adolescent Version (CAPS-CA; Nader, 2002), which indexes
10 both the frequency and intensity of PTSD symptoms.
11

12
13 *Oxford.* A combination of structured clinical interview and self-report measures determined ASD
14 diagnosis. The widely established Children's Impact of Event Scale (IES-8; Dyregrov & Yule, 1995),
15 an 8-item self-report measure of intrusion and avoidance symptoms, was used to assess symptoms
16 on a 4-point scale (Not at all=0; Rarely=1; Sometimes=3; Often=5). The Child Post-Traumatic Stress
17 Research Index (CPTSD-RI; Pynoos et al., 1987), a systematic clinical assessment of PTSD with
18 widely established psychometric properties, was administered to cover items not contained in the IES-
19 8 (i.e., arousal items).
20
21
22
23
24
25
26
27

28 *Impairment ratings.* Positive categorical impairment ratings were calculated according to whether
29 the young person endorsed problems in at least one ASD Criterion F area of functioning (e.g., school,
30 family, and social) on each of the instruments described above.
31
32

33 **Data analysis:**

34 *Confirmatory Factor Analysis*

35
36
37 Four CFA models were specified (Table 1). Due to the dichotomous nature of the items,
38 tetrachoric inter-item correlations were estimated, and covariate adjustment then made within EQS
39 v6.1 software (Bentler, 2006). A preliminary factor analysis using direct oblimin rotation was carried
40 out to obtain an estimate of the size of the factor loadings for marker variables in need of scaling in
41 the subsequent analysis (Loehlin, 1992). Mardia's multivariate kurtosis coefficient as recommended
42 by, for example, Raykov and Marcoulides (2006) and its normalised estimate were used to assess
43 whether the multivariate distribution of all the observed variables deviated from normality. This
44 statistic, and the univariate item skew of >1.5 , suggested non-normality in the items. Robust
45 maximum likelihood estimation was therefore used to fit the factor models in this sample of
46 intermediate size (Lee, Poon & Bentler, 1995). These estimates also perform better than uncorrected
47 statistics where the normal distribution assumption is false and better than a distribution-free method
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 in all but the largest samples (Chou, Bentler, & Satorra, 1991; Hu, Bentler, & Kano, 1992). The
5
6 tetrachoric correlation is recommended as a measure of association between pairs of categorical
7
8 variables (Tabachnick & Fidell, 2007). The Satorra-Bentler scaled χ^2 goodness of fit test (Satorra &
9
10 Bentler, 1994) was used to index the goodness of fit for each model. The test is sensitive to large
11
12 sample sizes (Tabachnick & Fidell, 2007) and a cut-off of a χ^2 df of ≤ 3 indicates an acceptable fit.

13
14 Goodness of fit indices as recommended by Bentler (2007) were used: the Comparative Fit
15
16 Index (CFI); Tucker-Lewis Index (TLI); and the root mean square error of approximation (RMSEA)
17
18 with its 90% confidence interval (Moschopoulos & Canada, 1984). Better fitting models are denoted
19
20 by higher CFI and TLI, with 0.90 representing a good fit, and 0.95 an excellent fit (Kline, 2005). Better
21
22 fitting models are indicated by lower RMSEA scores. RMSEA values of ≤ 0.05 , and 90% confidence
23
24 intervals whose lower bound contains, or is very close to, zero, and whose upper bound is ≤ 0.08 , are
25
26 thought to indicate a close fit, 0.05-0.08 a fair fit, and 0.08-0.10 a marginal fit by one standard
27
28 deviation (Browne & Cudeck, 1992). Multiple fit indices assessed model fit as fit indices are heavily
29
30 influenced by sample size, model parameters, and data-normality (Bentler, 2007). A good model
31
32 would meet at least two fit criteria, and meeting three fit criteria would only be considered necessary
33
34 according to a stringent criterion (Schermelleh-Engel, Moosbrugger, & Müller, 2003). Schwarz's
35
36 Bayesian information criterion model (BIC) combines goodness of fit with the number of model
37
38 parameters needed to allow for model comparisons. BIC scores that are ≥ 10 points lower than the
39
40 next lowest model are evidence for the superiority of one model (Raftery, 1995). Factor loadings of
41
42 $\geq .30$ are needed for an item to be considered of practical significance (Hair et al., 2006).

43
44 MIMIC modelling was used to explore the moderating effects of recruitment site (Brown,
45
46 2006). A series of dummy variables was created, which were then specified in the structural equation
47
48 for each of the items. To determine whether interpersonal (i.e., assault; n=87 cases) and non-
49
50 interpersonal (i.e., RTC and accidental injuries; n=507) experiences led to a distinct profile of PTSD
51
52 symptoms (cf. Shevlin & Elklit, 2011), a multiple-sample group analysis (Bentler, 2006) was carried
53
54 out in which tetrachoric correlation matrices were specified. In each analysis, factors that held
55
56 univariate correlations to any ASD items were also co-varied; in this case age and sub-ASD (whether
57
58 individuals met all of the DSM-IV ASD criteria except dissociation—a useful index of clinical status
59
60

given the problems with the dissociation criterion¹) were removed by specifying them in the structural equation for each item.

Results

The latent structure of ASD in children and young people

Table 3 presents the fit indices for the four models for the full sample. The χ^2 statistics for each model were significant, and all but the 1-factor DSM-5 model met the pre-requisite of a χ^2 : *df* of ≤ 3 . The 1-factor DSM-5 model was a good fit to the data ($\geq .90$) according to the CFI but not according to the TLI. The 3-, 4- (DSM-IV) and 5-factor models were all an excellent fit to the data according to both fit indices (CFI and TLI $\geq .95$). According to RMSEA scores, the 1-factor DSM-5 model was a fair fit (i.e., RMSEA $\leq .08$) whereas the 3- 4- and 5-factor models were all a close fit of the data (RMSEA $\leq .05$). The lower bound confidence intervals within the RMSEA were also close to 0 for the three more complex models, indicating a higher level of precision. The model BIC highlighted the 3-factor structure as the preferred model, in that this model was 17 points lower than the next lowest model. Inter-factor correlations for the three factor model ranged between $r = .50$ ($p < .001$) and $.80$ ($p < .001$).

We sought to determine the generalizability of this best-supported 3-factor model across: (a) recruitment site; and, (b) trauma type. Similar results were obtained when site differences were taken into account. The three factor model had the lowest BIC (by over 100) and an acceptable fit according to both CFI ($\geq .90$) and RMSEA ($\leq .08$) although not by TLI ($= .85 < .90$) or χ^2 : *df* ($= 3.3 > 3$). The multi-sample CFA assessing differences in model fit across trauma type also showed the three factor model had the lowest BIC (by over 100), a χ^2 : *df* of ≤ 3 and good fits using CFI ($\geq .90$), RMSEA ($\leq .05$) and TLI ($\geq .90$).

The factor loadings for the 3-factor model in the full sample are presented in Table 4. The pre-requisite of a factor loading of $\geq .30$ was met for all items aside from amnesia for the trauma (DSM-5 B7), which poorly loaded onto all other factors².

Prevalence rates

DSM-5 ASD (9+ symptoms) led to a .3% increase in prevalence of ASD relative to DSM-IV, with rates of 13.6% and 13.3% found for the two models respectively.

¹ Prevalence of Sub-ASD varied between 10% and 49% across groups¹⁰

² This pattern of results for factor loadings was also replicated in group invariance tests addressing recruitment site and trauma type.

Relationship to impairment

Two hundred and nine (35.2%) young people met the impairment criterion (i.e., whether the young person endorsed problems in at least one Criterion F area of functioning). Table 5 presents sensitivity, specificity, positive predictive value (PPV), negative predictive value (NPV), for the percentage of young people correctly classified as suffering impairment using for the DSM-IV and DSM-5 models. Please see Supplementary Table S2 for a similar table documenting the results of alternate factor models.

The uni-factorial DSM-5 ASD model of 9+ symptoms and DSM-IV were remarkably similar, with DSM-5 only marginally outperforming the DSM-IV model in terms of sensitivity and numbers of cases correctly identified, although neither was strong. Reducing the DSM-5 symptom requirements to either 3+ or 4+ (cf. Kassam-Adams et al., 2012) symptoms improved the sensitivity of both models by >46 percentage points. However, this improvement was off-set by low specificity and elevated ASD prevalence rates.

Discussion

In the present study, we aimed to evaluate the latent structure of ASD symptoms in children and young people following single incident traumatic events and compare DSM-5 and DSM-IV prevalence rates and their ability to categorise children impaired by their symptoms. Our findings provided mixed support for the new diagnosis in children and young people. Our exploratory examination of the 3-factor model showed that it offered the optimal account of symptom clustering. The findings showed that the DSM-5 'general distress' model was a good fit for the data according to two of three fit indices, whereas the 3- and 4- and 5-factor models were excellent fits for the data according to all three fit indices. DSM-5 and DSM-IV models led to very similar prevalence rates and were both poor models for categorising children's level of impairment. The sensitivity of the uni-factorial model was improved by lowering the symptom threshold from 9+ (DSM-5) to 3+/4+ symptoms (cf. Kassam-Adams et al., 2012).

Overall, the preferred model from a structural perspective was the 3-factor model, with re-experiencing and arousal symptoms clustered together, but distinct from avoidance and dissociation. It is important to note that this consistent factor structure was upheld after adjusting for, age, sub-ASD and site differences and comparisons across trauma type. The 3-factor model was previously shown to meet the requisite requirements of a good fitting model in three studies of ASD in adults

1
2
3
4 (Edmondson et al., 2010; Hansen et al., 2013; Wang et al., 2010), but this model was not evaluated in
5
6 the only adolescent study on this issue (Mordeno & Cue, 2015). The findings challenge the cluster-
7
8 free approach of ASD in the DSM-5, as clustering is clearly present acutely following traumatic
9
10 experiences. The good fit of adult post-traumatic stress models on the whole suggests that early
11
12 responses to trauma are perhaps not dissimilar from chronic responses in children. Theoretically, this
13
14 raises important further questions. If the symptom structure of ASD is similar to PTSD, we might
15
16 surmise that there is no evidence for an early “general distress” syndrome, but rather a distinctive
17
18 early post-traumatic stress response that proceeds (without fundamental change in symptom
19
20 structure) to PTSD in some cases, but remits naturally over time without intervention in others
21
22 (Kassam-Adams et al. 2004; Meiser-Stedman et al., 2005, 2008). However, this is speculative as a
23
24 two-factor PTSD model (i.e., avoidance, re-experiencing+arousal) has rarely been examined in the
25
26 child literature. Future CFA studies carried out in youth should test this model to determine the
27
28 continuities/discontinuities in the structure of acute and chronic trauma responses in youth. In
29
30 particular, future research must consider whether dissociation symptoms are an essential feature of
31
32 the ASD diagnosis, whether there is any benefit if a more liberal number of symptoms are endorsed in
33
34 competing models with a stronger factor structure, or whether a dissociative subtype of ASD should
35
36 be introduced.

37 Examination of patterns of prevalence and discriminant validity were informative. Our results
38
39 indicate that the adoption of the DSM-5 model does not have a negative impact on the detection of
40
41 children impaired by their symptoms relative to the other competing models (with superior patterns of
42
43 clustering). Prevalence rates for both models were within the ranges suggested by previous research
44
45 of ASD in youth (Bryant et al., 2004; Dalgleish et al., 2008; Kassam-Adams & Winston, 2004; Meiser-
46
47 Stedman et al., 2008). Replicating previous research (Kassam-Adams et al., 2012), the DSM-5 cluster
48
49 free model was more closely associated with clinical impairment when the developmental changes
50
51 suggested by Kassam-Adams et al. (2012) of 3+ or 4+ symptom cut-offs were suggested, although
52
53 making such a change to the DSM would need to be carefully considered as our data indicate these
54
55 lower cut-offs would likely lead to a higher rates of false positives as well as markedly increasing
56
57 prevalence rates. This might then increase the provision of treatment to children whose symptoms are
58
59 in fact likely to abate over time naturally, directing valuable resources away from children potentially in
60
61 greater need of help (Magruder & Calderone, 2000; Mullins-Sweatt, Lengel, & DeShong, 2016). To

1
2
3
4 explore this position further, prospective studies investigating the impact of factor structure on
5
6 recovery trajectories and relapse rates in youth must be carried out.
7

8 Some aspects of the present methodology merit comment. The data here were collected prior to
9 the publication of the DSM-5 and we therefore have no pure measure of the new B5 emotional
10 numbing symptom that refers to an absence of positive affect only (instead we used an item asking
11 people to indicate their recent experience of positive or negative affect). In saying that, the addition of
12 this new symptom would not have influenced the results of the CFA analyses that suggest that
13 cluster-free approaches are the weakest of each of the models. We also pooled data across four sites
14 that use different instruments on the basis that DSM diagnoses are universal and different diagnostic
15 instruments are designed to yield the same underlying constructs. However, the low numbers in some
16 groups (e.g., Oxford, interpersonal-trauma) was a limitation. Furthermore, the fit of the 3-factor model
17 was reduced when controlling for site, although it is important to highlight that it still met the minimum
18 fit requirement of a 'good' model (Schermelleh-Engel et al., 2003). Prevalence rates of ASD varied
19 widely across site. Prevalence rates of ASD varied widely across site. These differences are likely
20 due to a number of reasons, including differing age distributions, different countries within which the
21 studies were conducted, different distributions of trauma types recruited, and different research
22 teams. Power limitations preclude a satisfactory examination of these factors via sub-analyses.
23
24
25
26
27
28
29
30
31
32
33
34
35

36 To our knowledge, this is the first study to show that the 'uni-factorial' DSM-5 model is not the
37 optimal account of ASD symptom structure in youth. With the publication of DSM-5 in 2013, and its
38 cluster-free approach to ASD, there is a significant need to continue to more fully explore the impact
39 of nosological models of ASD in youth on theory and clinical practice.
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Key Points

- The latent structure of Acute Stress Disorder (ASD) symptoms has never before been investigated in a sample of youth following ED attendance or hospital admission.
- The results of this Confirmatory Factor Analysis (CFA) investigation showed that a “cluster free” model of ASD, the model adopted for the DSM-5 (i.e. a “general distress” conceptualisation of early responses to trauma), was not supported in youth; instead a 3-factor model comprising dissociation, re-experiencing/arousal and avoidance dimensions was found to be the preferred model.
- This study shows that the pattern of symptom clustering directly opposes the ‘general-distress’ model of ASD in DSM-5 and supports a continuity between the structure of acute stress reactions and chronic post-traumatic stress (as specified in the PTSD diagnosis).
- The proportion of children that will develop DSM-5 ASD following exposure to single incident trauma is approximately 13.6 percent.
- Reducing the DSM-5 ASD algorithm to either 3 or 4 symptoms improved the sensitivity of DSM-5 ASD to detect children impaired by their symptoms.

Acknowledgements

All authors declare they have no biomedical financial interests or potential conflicts of interest.

Professor Dagleish and Dr Watson are supported by the UK Medical Research Council (MRC). Dr Meiser-Stedman, Ms Dixon & Dr McKinnon were supported by an MRC Clinician Scientist Fellowship (G0802821) awarded to Dr Meiser-Stedman, supporting data collection at the Cambridge site. The London site was funded by an MRC Research Studentship (G78/6730) and a Peggy Pollak Research Fellowship in Developmental Psychiatry, by the Psychiatry Research Trust to Dr Meiser-Stedman. The Philadelphia site was supported by a research grant to Drs. Winston and Kassam-Adams from the Maternal and Child Health Bureau, US Health Resources and Services Administration. Professor Anke Ehlers and the Oxford study were funded by the Wellcome Trust.

Table 1. Model specifications for alternative factor models of ASD.

DSM-5 ASD symptoms	Model 1	Model 2	Model 3	Model 4
	DSM-5 1-factor	Alternate 3-factor	3 DSM- IV	Alternate 5-factor
B1. Intrusive memories, thoughts	ASD	Re-ex/Ar	Re-ex	Re-ex
B2. Nightmares	ASD	Re-ex/Ar	Re-ex	Re-ex
B3. Flashbacks	ASD	Re-ex/Ar	Re-ex	Re-ex
B4a. Psychological distress to reminders ¹	ASD	Re-ex/Ar	Re-ex	Re-ex
B4b. Physiological reactivity ¹	ASD	Re-ex/Ar	Re-ex	Re-ex
B5. Emotional numbing ²	ASD	Diss	Diss	Diss
B6a. Altered sense of reality: loss of awareness ³	ASD	Diss	Diss	Diss
B6b. Altered sense of reality: Derealisation ³	ASD	Diss	Diss	Diss
B6c. Altered sense of reality: Depersonalisation ³	ASD	Diss	Diss	Diss
B7. Amnesia	ASD	Diss	Diss	Diss
B8. Avoid thinking/conversations	ASD	Av	Av	Av
B9. Avoid places/things/people	ASD	Av	Av	Av
B10. Difficulty sleeping	ASD	Re-ex/Ar	Ar	Dys-Ar
B11. Irritability	ASD	Re-ex/Ar	Ar	Dys-Ar
B12. Difficulty concentrating	ASD	Re-ex/Ar	Ar	Dys-Ar
B13. Hyper-vigilance	ASD	Re-ex/Ar	Ar	Anx-Ar
B14. Startle	ASD	Re-ex/Ar	Ar	Anx-ar

¹ DSM-IV symptoms (C: psychological distress to reminders; E: Physiological reactivity) are grouped under the one symptom in DSM-5 (DSM-5 B4).

² DSM-IV emotional numbing (B1) is reworded in DSM-5 to describe absence of positive emotional experiences (DSM-5 B5).

³ DSM-IV reduction in awareness of surroundings (B2), derealization (B3), and depersonalization (B4) are grouped under DSM-5 Altered Sense of Reality (B6).

Note. Diss=dissociation; Av=avoidance; Ar=arousal; Re-ex=re-experiencing; Dys-Ar=dysphoric arousal; Anx-ar=anxious arousal; ASD=Acute Stress Disorder.

Table 2. Demographic characteristics and trauma related characteristics for the 4 sites.

	Descriptives				Total
	East Anglia	London	Oxford	Philadelphia	
<i>Demographics</i>					
n	32% (n=189)	25% (n=151)	11% (n=65)	32% (n=189)	100% (n=594)
Age (M, SD)	14.17 (2.89)	11.94 (3.00)	12.67 (2.68)	11.39 (2.41)	12.56 (2.99)
6-8 yrs	5.8% (n=11)	19% (n=28)	11% (n=7)	14% (n=27)	12% (n=73)
9-12 yrs	30% (n=56)	41% (n=62)	42% (n=27)	55% (n=103)	42% (n=248)
13-16 yrs	65% (n=122)	40% (n=61)	48% (n=31)	31% (n=59)	46% (n=273)
Male gender	56% (n=107)	58% (n=88)	58% (n=38)	76% (n=144)	64% (n=377)
<i>Trauma type</i>					
Assault	19% (n=35)	34% (n=52)	0% (n=0)	0% (n=0)	15% (n=87)
RTC	47% (n= 88)	66% (n=99)	100% (n=65)	100% (n=189)	74% (n=441)
Accidental injury	35% (n=66)	0% (n=0)	0% (n=0)	0% (n=0)	11% (n=66)
Full DSM-5 ASD met	13%(n=25)	23% (n=35)	28% (n=12)	5% (n=9)	14% (n=103)

Table 3. Fit Indices for the four alternative ASD models (N=594) in the pooled sample.

Item Models	Satorra-Bentler $\chi^2(\text{df})^1$	p	BIC ²	CFI ³	90% CI RMSEA ⁴	TLI ⁵
1 Factor	$\chi^2(118)=391.78$	<.001	-361.87	0.90	0.063; 0.056, 0.069	0.89
3 Factors⁶	$\chi^2(116)=234.09$	<.001	-506.79	0.96	0.041; 0.034, 0.049	0.95
4 Factors	$\chi^2(113)=231.10$	<.001	-490.61	0.96	0.042; 0.034, 0.050	0.95
5 Factors	$\chi^2(109)=207.14$	<.001	-489.03	0.97	0.039; 0.031, 0.047	0.96

Note

¹ Satorra Bentler χ^2 ; ² Bayesian Information Criterion; ³ Comparative Fit Index; ⁴ Root Mean Square Error of Approximation; ⁵ Tucker Lewis Index; ⁶ Models in **bold** indicate the best fitting model.

Table 4. Standardized factor loadings (standard errors) for the ASD 3-factor model in the pooled sample (N=594).

DsM-5 ASD symptoms	Dissociation	Re-experiencing/ Arousal	Avoidance
B1. Intrusive memories, thoughts		0.83 (0.04)	
B2. Nightmares		0.69 (0.05)	
B3. Flashbacks		0.82 (0.06)	
B4a. Distress to reminders ¹		0.71 (0.05)	
B4b. Physiological reactivity ¹		0.75 (0.05)	
B5. Emotional numbing ²	0.72 (0.05)		
B6a. loss of awareness ³	0.70 (0.05)		
B6b. Derealisation ³	0.87 (0.06)		
B6c. Depersonalisation ³	0.75 (0.05)		
B7. Amnesia	0.16 (0.06)		
B8. Avoid thinking/conversations			0.83 (0.05)
B9. Avoid places/things/people			0.85 (0.05)
B10. Difficulty sleeping		0.70 (0.05)	
B11. Irritability		0.63 (0.05)	
B12 Difficulty concentrating		0.73 (0.05)	
B13. Hyper-vigilance		0.75 (0.05)	
B14. Startle		0.77 (0.04)	

¹ DSM-IV symptoms (C: psychological distress to reminders; E: Physiological reactivity) are grouped under DSM-5 B4.

² DSM-IV emotional numbing (B1) is reworded in DSM-5 to describe absence of positive emotional experiences (DSM-5 B5).

³ DSM-IV reduction in awareness of surroundings (B2), derealization (B3), and depersonalization (B4) are grouped under DSM-5 B6.

Table 5. Performance of different symptom requirements per acute stress models to predict concurrent ratings of impairment (N=594).

Model	Cluster	Frequency symptom/diagnosis ¹	Sensitivity	Specificity	PPV	NPV	% correctly classified	% ASD diagnosis
Model 1: DSM-5	One-factor: [3+] ²	194 (32.7%)	92.82	49.35	49.87	92.68	64.6	33.0%
Model 1: DSM-5	One-factor: [4+] ³	178 (30.0%)	85.17	63.64	55.97	88.77	71.2	30.0%
Model 1: DSM-5	One-factor: [9+] ⁴	103 (17.3%)	38.77	94.29	78.64	73.93	74.7	13.6%
Model 3: DSM-IV	Four factor DSM-IV ⁵	103 (17.3%)	37.80	93.77	76.70	73.52	74.1	13.3%

Note. NPV=negative predictive value; PPV=positive predictive value

¹ The number of cases meeting the frequency requirement per symptom cluster and diagnosis (i.e., without meeting impairment).

² 3+ symptoms from the DSM-5 list of 14 symptoms

³ 4+ symptoms from the DSM-5 list of 14 symptoms

⁴ 9+ symptoms from the DSM-5 list of 14 symptoms

⁵

References

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- Armour, C., Shevlin, M., & Elkit, A. (2013). The latent structure of Acute Stress Disorder: A post-traumatic stress disorder approach. *Psychological trauma: theory, research, practice, and policy*, 5, 18
- Bentler, P. M. (2006). *EQS 6 structural equations program manual*. Encino, CA: Multivariate software Inc.
- Bentler, P. M. (2007). On tests and indices for evaluating structural models. *Personality and Individual Differences*, 42, 825-829.
- Brady, K. T., Killeen, T. K., Brewerton, T., & Lucerini, S. (2000). Comorbidity of psychiatric disorders and posttraumatic stress disorder. *Journal of Clinical Psychiatry*, 61, 22-32.
- Brewin, C. R., Andrews, B., & Rose, S. (2003). Diagnostic overlap between acute stress disorder and PTSD in victims of violent crime. *American Journal of Psychiatry*, 160, 783-785. doi: 10.1176/appi.ajp.160.4.783
- Brooks, R., Silove, D., Bryant, R., O'Donnell, M., Creamer, M., & McFarlane, A. (2008). A confirmatory factor analysis of the Acute Stress Disorder Interview. *Journal of Traumatic Stress*, 21, 352-355.
- Brown, T. A. (2006). *Confirmatory factor analysis for applied research*. New York: The Guilford Press.
- Browne, M. W., & Cudeck, R. (1992). Alternative ways of assessing model fit. *Sociological Methods & Research*, 21, 230-258.
- Bryant, R. A. (2011). Acute Stress Disorder as a Predictor of Posttraumatic Stress Disorder: A Systematic Review. *Journal of Clinical Psychiatry*, 72, 233-239.
- Bryant, B., Mayou, R., Wiggs, L., Ehlers, A., & Stores, G. (2004). Psychological consequences of road traffic accidents for children and their mothers. *Psychological Medicine*, 34, 335-346.
- Chen, S. H., Lin, Y. H., Tseng, H. M., & Wu, Y. C. (2002). Posttraumatic stress reactions in children and adolescents one year after the 1999 Taiwan Chi-Chi Earthquake. *Journal of the Chinese Institute of Engineers*, 25, 597-608. doi: 10.1080/02533839.2002.9670734

- 1
2
3
4 Chou, C. P., Bentler, P. M., & Satorra, A. (1991). Scaled test statistics and robust standard errors for
5
6 nonnormal data in covariance structure analysis: A Monte-Carlo study. *British Journal of*
7
8 *Mathematical & Statistical Psychology*, 44, 347-357.
- 9 Dagleish, T., Meiser-Stedman, R., Kassam-Adams, N., Ehlers, A., Winston, F., Smith, P., . . . Yule,
10
11 W. (2008). Predictive validity of acute stress disorder in children and adolescents. *British*
12
13 *Journal of Psychiatry*, 192, 392-393.
- 14
15 *Diagnostic and Statistical Manual of Mental Disorders, 4th edition.* (1994). Washington, DC: American
16
17 Psychiatric Association.
- 18
19 *Diagnostic and Statistical Manual of Mental Disorders, 5th edition.* (2013). Washington, DC: American
20
21 Psychiatric Association.
- 22 Dyregrov, A., & Yule, W. (1995). *The Development of the UNICEF Screening Battery.* Paper
23
24 presented at the Fourth European Conference on Traumatic Stress Paris.
- 25
26 Edmondson, D., Mills, M. A., & Park, C. L. (2010). Factor Structure of the Acute Stress Disorder Scale
27
28 in a Sample of Hurricane Katrina Evacuees. *Psychological Assessment*, 22, 269-278.
- 29
30 Frances, A. J., & Widiger, T. (2012). Psychiatric Diagnosis: Lessons from the DSM-IV Past
31
32 and Cautions for the DSM-5 Future. In S. NolenHoeksema (Ed.), *Annual Review of*
33
34 *Clinical Psychology, Vol 8* (Vol. 8, pp. 109-130). Palo Alto: Annual Reviews.
- 35
36 Hair, J. F. J., Black, W. C., Babin, B. J., Anderson, R. E., & Tatham, R. L. (2006). *Multivariate data*
37
38 *analysis* (6th ed.). New Jersey: Prentice Hall.
- 39
40 Hansen, M., Armour, C., & Elklit, A. (2012). Assessing a dysphoric arousal model of acute stress
41
42 disorder symptoms in a clinical sample of rape and bank robbery victims. *European Journal of*
43
44 *Psychotraumatology*, 3.
- 45
46 Hansen, M., Lasgaard, M., & Elklit, A. (2013). The latent factor structure of acute stress disorder
47
48 following bank robbery: Testing alternative models in light of the pending DSM-5. *British*
49
50 *Journal of Clinical Psychology*, 52, 82-91
- 51
52 Harvey, A. G., & Bryant, R. A. (1998). The relationship between acute stress disorder and
53
54 posttraumatic stress disorder: A prospective evaluation of motor vehicle accident survivors.
55
56 *Journal of Consulting and Clinical Psychology*, 66, 507-512.
- 57
58 Hu, L. T., Bentler, P. M., & Kano, Y. (1992). Can test statistics in covariance structure-anlysis be
59
60 trusted? *Psychological Bulletin*, 112, 351-362.

- 1
2
3
4 Kassam-Adams, N. (2006). The acute stress checklist for children (ASC-Kids): Development of a child
5
6 self-report measure. *Journal of Traumatic Stress, 19*, 129-139.
7
- 8 Kassam-Adams, N., Palmieri, P. A., Rork, K., Delahanty, D. L., Kenardy, J., Kohser, K. L., . . .
9
10 McGrath, C. (2012). Acute Stress Symptoms in children: Results from an International Data
11
12 Archive. *Journal of the American Academy of Child and Adolescent Psychiatry, 51*, 812-820.
13
- 14 Kassam-Adams, N., & Winston, F. K. (2004). Predicting child PTSD: The relationship between acute
15
16 stress disorder and PTSD in injured children. *Journal of the American Academy of Child and*
17
18 *Adolescent Psychiatry, 43*, 403-411.
19
- 20 Kline, R. B. (2005). *Principles and practice of structural equation modeling*. New York: Guilford Press.
- 21 Lee, S. Y., Poon, W. Y., & Bentler, P. M. (1995). A two-stage estimation of structural equation models
22
23 with continuous and polytomous variables. *British Journal of Mathematical & Statistical*
24
25 *Psychology, 48*, 339-358.
26
- 27 Loehlin, J. C. (1992). *Latent variable models an introduction to factor, path, and structural analysis*
28
29 (2nd ed.). Hillsdale, NJ: Lawrence Erlbaum.
- 30 March, J. (1999). Assessment of pediatric Post-traumatic stress disorder. In P. Saigh & D. Bremner
31
32 (Eds.), *Post-traumatic stress disorder* Washington, DC: American Psychological Press.
- 33 Meiser-Stedman, R., Smith, P., Glucksman, E., Yule, W., & Dalgleish, T. (2008). The posttraumatic
34
35 stress disorder diagnosis in preschool- and elementary school-age children exposed to motor
36
37 vehicle accidents. *American Journal of Psychiatry, 165*, 1326-1337.
38
- 39 Meiser-Stedman, R., Yule, W., Smith, P., Glucksman, E., & Dalgleish, T. (2005). Acute stress disorder
40
41 and posttraumatic stress disorder in children and adolescents involved in assaults or motor
42
43 vehicle accidents. *American Journal of Psychiatry, 162*, 1381-1383.
44
- 45 Magruder, K. M., & Calderone, G. E. (2000). Public health consequences of different thresholds for
46
47 the diagnosis of mental disorders. *Comprehensive Psychiatry, 41*, 14-18. doi: 10.1016/s0010-
48
49 440x(00)80003-6.
50
- 51 Mullins-Sweatt, S. N., Lengel, G. J., & DeShong, H. L. (2016). The Importance of Considering Clinical
52
53 Utility in the Construction of a Diagnostic Manual. In T. D. Cannon & T. Widiger (Eds.), *Annual*
54
55 *Review of Clinical Psychology, Vol 12* (Vol. 12, pp. 133-155).
56
- 57 Mordeno, I.G. & Cue, M.P. (2015). Examining the latent structure of Acute Stress Disorder symptoms
58
59 in Filipino-vctims of a flashflood disaster. *Community Mental Health Journal, 51*, 613-620.
60

- 1
2
3
4 Moschopoulos, P. G., & Canada, W. B. (1984). The distribution function of a linear combination of chi-
5 squares. *Computers & Mathematics with Applications*, 10(4-5), 383-386.
6
7
8 Nader, K., Kriegler, J.A., Blake, D.D., Pynoos, R.S., Newman, E., & Weather, F.W. (1996). Clinician
9 Administered PTSD Scale, Child and Adolescent Version. White River Junction, VT: National
10 Center for PTSD.
11
12 Pynoos RS, Frederick C, Nader K, et al. (1987) Life threat and posttraumatic stress in school-age
13 children. *Archives of General Psychiatry*, 44, 1057-1063.
14
15 Raftery, A. E. (1995). Bayesian model selection in social research. *Sociological Methodology*, 25, 111-
16 163.
17
18
19 Raykov, T., & Marcoulides, G. A. (2006). On multilevel model reliability estimation from the
20 perspective of structural equation modeling. *Structural Equation Modeling-a Multidisciplinary*
21 *Journal*, 13, 130-141.
22
23
24 Saigh, P. A., Yasik, A. E., Oberfield, R. A., Green, B. L., Halamandaris, P. V., Rubenstein, H., . . .
25
26 McHugh, M. (2000). The children's PTSD Inventory: Development and reliability. *Journal of*
27 *Traumatic Stress*, 13, 369-380.
28
29
30 Satorra, A., & Bentler, P. M. (1994). Corrections to test statistics and standard errors in covariance
31 structure analysis. In A. von Eye & C. C. Clogg (Eds.), *Latent variables analysis: applications*
32 *for developmental research* (pp. 399-419). Thousand Oaks, CA: Sage.
33
34
35 Schermelleh-Engel K, Moosbrugger H, & Müller H. Evaluating the fit of structural equation models:
36 Tests of significance and descriptive goodness-of-fit measures. *Methods of Psychological*
37 *Research Online*. 2003;8:23-74.
38
39
40 Shevlin, M., & Elklit, A. (2011). The latent structure of Posttraumatic Stress Disorder: Different models
41 or different populations? *Journal of Abnormal Psychology*, 121, 610-615.
42
43 Silverman, W. K., & Albano, A. M. (1996). *Anxiety Disorder Interview Schedule for DSM-IV: Child and*
44 *Parent Interview Schedule*. San Antonio, TX: The Psychological Corporation.
45
46
47
48 Tabachnick, B. G., & Fidell, L. S. (Eds.). (2007). *Using multivariate statistics* (5th ed.). Boston:
49 Pearson International Edition.
50
51
52
53 Wang, L., Li, Z. Q., Shi, Z. B. A., Zhang, Y. Q., & Shen, J. H. (2010). Factor structure of acute stress
54 disorder symptoms in Chinese earthquake victims: A confirmatory factor analysis of the acute
55 stress disorder scale. *Personality and Individual Differences*, 48, 798-802.
56
57
58
59
60

1
2
3
4 Yasik, A. E., Saigh, P. A., Oberfield, R. A., Green, B., Halamandaris, P., & McHugh, M. (2001). The
5
6 validity of the children's PTSD Inventory. *Journal of Traumatic Stress, 14*, 81-94.
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Supplementary Table 1. Confirmatory Factor Analytic studies examining the structure of Acute Stress Disorder symptoms in adults.

Study	Sample	Models tested ¹	CFI ²	RMSEA ³
Brooks et al. (2008)	N=587 level 1 trauma patients	Dissociation, re-experiencing, arousal, avoidance	.97*	.036*
		<i>Dissociation^{1st order}, re-experiencing^{1st order}, arousal^{1st order}, avoidance^{1st order}, general distress^{2nd order}</i>	.97*	.032*
Edmondson et al. (2010)	N=132 Hurricane Katrina evacuees.	Dissociation, re-experiencing, arousal, avoidance	.86	.09
		<i>Dissociation, re-experiencing/arousal, avoidance</i>	.91*	.07*
		<i>Dissociation^{1st order}, re-experiencing^{1st order}, arousal^{1st order}, avoidance^{1st order}, general distress^{2nd order}</i>	.92*	.07*
		Dissociation^{1st order}, re-experiencing^{2nd order}, avoidance^{2nd order}, arousal^{2nd order}	.93*	.07*
Wang et al. (2010)	N=353 Chinese earthquake victims	Dissociation, re-experiencing, arousal, avoidance	.99*	.038*
		<i>Dissociation, re-experiencing/arousal, avoidance</i>	.99*	.039*
		<i>Dissociation, dissociative amnesia, acute posttraumatic stress reactions</i>	.99*	.042*
		<i>Dissociation^{1st order}, re-experiencing^{1st order}, arousal^{1st order}, avoidance^{1st order}, general distress^{2nd order}</i>	.99*	.039*
Hansen et al. (2012)	N=404 rape and bank robbery with ASD.	ASD	.59	.085
		Dissociation, re-experiencing /arousal, avoidance	.74	.069*
		Dissociation, re-experiencing, arousal, avoidance	.74	.069*
		Dissociation, re-experiencing, dysphoria, arousal, avoidance	.81	.061*
Armour et al. (2013)	N=380 Danish rape victims	ASD	.67	.075*
		Dissociation, re-experiencing/arousal, avoidance	.80	.058*
		Dissociation, re-experiencing, arousal, avoidance	.80	.058*
		Dissociation, re-experiencing, dysphoria,	.81	.058*

Study	Sample	Models tested ¹	CFI ²	RMSEA ³
		arousal, avoidance		
		Dissociation, re-experiencing, numbing, avoidance, arousal	.79	.06*
Hansen et al. (2013)	N=450 bank robbery victims.	ASD	.96*	.089
		<i>Dissociation^{1st order}, re-experiencing^{2nd order}, avoidance^{2nd order}, arousal^{2nd order}</i>	.97*	.076*
		<i>Dissociation, re-experiencing /arousal, avoidance</i>	.98*	.070*
		Dissociation, re-experiencing, arousal, avoidance	.98*	.068*

Note

¹ Models *italicised* either had a good or excellent fit of the data according to the χ^2 in addition to CFI and RMSEA fit indices. Models in **bold** are the best fitting model.

²Comparative Fit Index: The ratio of the difference between the χ^2 for the fitted model and the null model divided by the χ^2 for the null model with $\geq .90$ =good fit and $\geq .95$ =excellent fit. CFI's meeting either criteria are marked by an asterisk.

³Root Mean Square Error of approximation: The amount of unexplained variance left by the models with $\leq .05$ suggesting a close fit and $\leq .08$ suggesting a fair fit. CFI's meeting either criteria are marked by an asterisk.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Supplementary Table 2. Performance of different symptom requirements per 3- and 5-factor models to predict concurrent ratings of impairment (N=594).

Cluster	Frequency symptom/diagnosis ¹	Sensitivity	Specificity	PPV	NPV	% correctly classified	% ASD diagnosis
3-factor ²	117	41.63	92.21	74.36	74.42	74.41	14.65%
5-factor ³	69	33.01	100	100	73.3	76.43	4.88%

Note. NPV=negative predictive value; PPV=positive predictive value

¹ The number of cases meeting the frequency requirement per symptom cluster and diagnosis (i.e., without meeting impairment).

² 3+ dissociation, 2+ re-experiencing/arousal, 1+ avoidance.

³ 3+ dissociation, 1+ re-experiencing, 1+ avoidance, 1+ arousal.