

Transcriptional and translational adaptation to aerobic nitrate anabolism in the denitrifier *Paracoccus denitrificans*

Victor M. Luque-Almagro¹, Isabel Manso¹, Matthew J. Sullivan², Gary Rowley², Stuart J. Ferguson³, Conrado Moreno-Vivián¹, David J. Richardson², Andrew J. Gates², M. Dolores Roldán^{1*}

¹*Departamento de Bioquímica y Biología Molecular, Universidad de Córdoba, Edificio Severo Ochoa, 1ª planta, Campus de Rabanales, Córdoba, 14071, Spain.*

²*School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich, NR4 7TJ, UK.*

³*Department of Biochemistry, University of Oxford, South Parks Road, Oxford, OX1 3QU, UK.*

* Corresponding author address:

Departamento de Bioquímica y Biología Molecular

Universidad de Córdoba

Córdoba 14071

Spain

Tel: +34 957218318

Fax: +34 957218592

E-mail: bb2rorum@uco.es

Running title: Regulation of nitrate and nitrite assimilation in *Paracoccus denitrificans*

Keywords: Global nitrogen regulator Ntr, nitrate assimilation, nitrate reductase, nitrate/nitrite sensor protein, nitrogen regulation, *Paracoccus*.

SUMMARY STATEMENT

Transcriptomic, proteomic and mutational analyses of the nitrate assimilatory system in *P. denitrificans* PD1222 reveals that nitrate exerts a very tight regulation at the transcriptional and translational levels through the NtrBC and the NasTS systems.

AUTHOR CONTRIBUTION

Matthew J. Sullivan (MJS) and Gary Rowley (GR) contributed to the microarray construction under the supervision of David J. Richardson (DJR), Stuart J. Ferguson (SJF) and Andrew J. Gates (AJG). Proteomic analysis and generation of mutant strains were carried out by Victor M. Luque-Almagro (VLA) under the supervision of Conrado Moreno-Vivián (CMV) and M. Dolores Roldán (MDR). RT-PCR and qPCR was performed by Isabel Manso (IM) under the supervision of CMV, MDR and AJG. The final version of this manuscript has been written by MDR and discussed and revised in agreement with DJR, CMV and SJF.

1 **ABSTRACT**

2
3 Transcriptional adaptation to nitrate-dependent anabolism by *Paracoccus*
4 *denitrificans* PD1222 was studied. A total of 74 genes were induced in cells grown
5 with nitrate as N-source compared to ammonium, including *nasTSABGHC* and
6 *ntrBC* genes. The *nasT* and *nasS* genes were cotranscribed, although *nasT* was more
7 strongly induced by nitrate than *nasS*. The *nasABGHC* genes constituted a
8 transcriptional unit, which is preceded by a non-coding region containing hairpin
9 structures involved in transcription termination. The *nasTS* and *nasABGHC*
10 transcripts were detected at similar levels with nitrate or glutamate as N-source,
11 but *nasABGHC* transcript was undetectable in ammonium-grown cells. The nitrite
12 reductase NasG subunit was detected by 2D-PAGE in cytoplasmic fractions from
13 nitrate-grown cells, but it was not present when either ammonium or glutamate
14 was used as N-source. The *nasT* mutant lacked both *nasABGHC* transcript and
15 NADH-dependent nitrate reductase activity. On the contrary, the *nasS* mutant
16 showed similar levels of the *nasABGHC* transcript to the wild-type strain and
17 displayed NasG protein and NADH-nitrate reductase activity with all N-sources
18 tested, except with ammonium. Ammonium repression of *nasABGHC* was
19 dependent on the Ntr system. The *ntrBC* and *ntrYX* genes were expressed at low
20 levels regardless of the nitrogen source supporting growth. Mutational analysis of
21 the *ntrBCYX* genes indicated that while *ntrBC* genes are required for nitrate
22 assimilation, *ntrYX* genes can only partially restore growth on nitrate in absence of
23 *ntrBC* genes. The existence of a regulation mechanism for nitrate assimilation in *P.*
24 *denitrificans*, by which nitrate induction operates at both transcriptional and
25 translational levels, is proposed.
26

1 INTRODUCTION

2
3 The soil denitrifier *Paracoccus denitrificans* PD1222 makes multiple metabolic uses of
4 nitrate: (i) as a respiratory electron acceptor to support anaerobic growth, catalysed by a
5 membrane-bound nitrate reductase (Nar); (ii) as an electron sink to dispose of excess
6 reductant during aerobic metabolism of highly reduced carbon substrates, catalysed by
7 the periplasmic nitrate reductase (Nap); and (iii) as nitrogen source for anabolism
8 during both oxic and anoxic growth. The first two processes have been widely studied
9 [1], but the biochemistry and regulation of the third process, nitrate assimilation, has
10 received much less attention. The nitrate assimilation system (Nas) is encoded by the
11 *nasABGHC* gene cluster, which contains a nitrate transporter (NasA), an NADH-
12 dependent nitrite reductase (NasB), a small Rieske-type protein required for both nitrate
13 and nitrite reduction (NasG), a nitrite transporter (NasH), and a nitrate reductase
14 (NasC), which receives electrons from NADH *via* NasB and NasG [2]. The NasB and
15 NasG proteins are essential for growth with nitrate or nitrite as sole nitrogen source
16 under both aerobic and anaerobic conditions. However, the NasA and NasC proteins are
17 required for nitrate import and reduction only under aerobic conditions because of a
18 functional overlap with biochemical components of the respiratory nitrate reductase
19 system that are synthesised under anaerobic conditions [2,3].

20 Nitrate assimilation is regulated in response to nitrate by a two-component
21 system, encoded by the *nasTS* genes, which are located directly upstream from the *nasA*
22 gene. NasS acts as nitrate/nitrite sensor whereas NasT has an ANTAR (AmiR and NasR
23 transcription antitermination regulators) domain for acting as a transcriptional
24 antiterminator [4,5]. In the absence of nitrate or nitrite, NasS and NasT form an inactive
25 tetrameric complex (two units of each), leading to the premature termination of the
26 *nasABGHC* gene transcription. In the presence of nitrate or nitrite, the NasTS complex
27 becomes dissociated, and thereby NasT allows expression of *nas* genes [5]. Recently,
28 the *P. denitrificans* PD1222 whole genome has been analyzed by using the Quadparser
29 programme, revealing the existence of a guanine-rich region located upstream of the
30 *nasT* gene that forms a canonical G-quadruplex structure. Stabilization of this secondary
31 structure in DNA has been suggested to act as a negative regulator for nitrate-dependent
32 growth [6].

33 Nitrate assimilation is a widespread metabolic capacity in proteobacteria that
34 usually is controlled at the transcriptional level by nitrate and nitrite induction and by
35 ammonium repression [7,8]. In cyanobacteria, the catabolite activator protein (CAP)
36 family transcription factor NtcA represses the nitrate reductase genes in the presence of
37 ammonium, whereas it activates transcription of these genes at a high C/N ratio
38 (nitrogen depletion), reflected by high 2-oxoglutarate levels [9-11]. In some
39 cyanobacteria, the LysR family transcription regulator NtcB is required for the
40 nitrate/nitrite-dependent induction of the nitrate reductase gene [11]. In Gram positive
41 bacteria like *Bacillus subtilis* and *Streptomyces coelicolor* the TnrA and GlnR regulators
42 respond to nitrogen starvation [12,13]. In *Klebsiella oxytoca*, expression of the nitrate
43 assimilation genes is activated under low nitrogen conditions through the global
44 nitrogen regulatory Ntr system, including NtrA, NtrB and NtrC proteins, and by
45 nitrate/nitrite induction through NasR, a transcription antitermination protein that also
46 binds to nitrate [14,15]. The crystal structure of the *Klebsiella oxytoca* NasR protein has
47 been solved; it is a dimer with a large N-terminal nitrate and nitrite-sensor (NIT)
48 domain and a C-terminal ANTAR domain necessary for specific binding to leader

1 mRNA. The NIT domain binds nitrate and nitrite between two conserved arginine
2 residues located on adjacent helices [16]. In *Azotobacter vinelandii* and *Pseudomonas*
3 *putida*, as well as in *P. denitrificans*, two different proteins NasT and NasS act as
4 transcriptional antiterminator and nitrate sensor, respectively [17-20].

5 The NtrBC two-component system has been extensively characterized in enteric
6 bacteria [21]. NtrB is a sensor kinase that autophosphorylates on a histidine residue
7 under low nitrogen concentrations and transfers a phosphoryl group to the NtrC
8 response regulator protein on a specific aspartate residue [22]. Phosphorylated NtrC acts
9 as a transcriptional activator that oligomerizes on the DNA template with ATPase
10 activity [23]. The NtrC members are usually dependent on the σ^{54} factor (NtrA) and
11 they are involved in transcription of genes related to nitrogen metabolism, such as the
12 glutamine synthetase *glnA* gene. However, in *Rhodobacter capsulatus* a regulatory two-
13 component NtrBC system has been described in which the NtrC component is not σ^{54} -
14 dependent to activate transcription of *nifA1* and *nifA2* genes, which code for
15 transcriptional activators that induce nitrogen fixation gene expression, and the *glnB*
16 gene that is a negative regulator of the *R. capsulatus* NtrBC system under nitrogen
17 excess [24].

18 The NtrYX system is also a two-component regulatory system with similarity to
19 the sensor/kinase NtrB and the regulatory protein NtrC, respectively. This system has
20 been investigated in diazotrophs with a proposed role in nitrogen assimilation. In
21 *Azorhizobium caulinodans*, a mutant in the *ntrX* gene was found to be defective in using
22 nitrate as nitrogen source, and showed also a reduced *nifA* expression under nitrogen
23 fixation conditions with a disturbed symbiotic phenotype. In the *ntrC* mutant strain,
24 expression of the *ntrYX* operon was derepressed in the presence of nitrate, suggesting an
25 interaction between both NtrBC and NtrYX systems [25]. The NtrBC system in
26 *Azospirillum brasilense* is involved in regulation of nitrate assimilation, ammonium
27 transport, and nitrogenase switch-off by ammonium. The NtrYX system may be
28 involved in nitrate utilization through a possible substitution of the NtrBC system by the
29 NtrYX sensor-regulator pair [26,27]. *Herbaspirillum seropedicae* is a diazotrophic β -
30 proteobacterium with both NtrBC and NtrYX systems displaying a role in regulation of
31 nitrate assimilation [28]. The photosynthetic bacterium *Rhodobacter capsulatus* also has
32 both NtrBC and NtrYX systems, with a function of the NtrBC system in urea
33 assimilation and nitrogen fixation, but with an unclear physiological function of the
34 NtrYX system [29]. The NtrYX system of *Brucella* spp. is involved in redox sensing
35 and regulation of denitrification genes. Expression of *narGHIJK*, *nirKV*, *norBCDEFQ*
36 and *nosDFLRXYZ* genes are down-regulated in an *ntrY* mutant strain under aerobic and
37 microaerobic conditions, with a marked down-regulation of the *nir*, *nor* and *nos* genes
38 under microaerobic conditions. The *Brucella* spp. NtrY protein contains one haem
39 group for sensing the oxygen status in the cell and shows its maximal activity as an
40 autohistidine kinase in the ferrous state under low oxygen tension [30,31].

41 Analysis of the whole genome sequence of *P. denitrificans* reveals the presence
42 of genes encoding NtrBC and NtrYX proteins. In this work, we explore the global
43 transcriptomic changes that underpin the transition from ammonium-dependent to
44 nitrate-dependent aerobic growth, including the relationships within the different two-
45 component regulatory systems NasTS, NtrBC and NtrYX. An additional mechanism to
46 the transcriptional regulation of nitrate assimilation, based on a nitrate-dependent
47 translation control of the *P. denitrificans* PD1222 Nas proteins, is also proposed.

1 MATERIALS AND METHODS

2

3 **Bacterial strains, media and growth conditions**

4 All strains used in this study are listed in Table S1. *P. denitrificans* PD1222 was grown
5 at 30 °C in Luria-Bertani (LB) medium [32] or defined mineral salts medium as
6 previously described [33], containing 50 mM succinate as carbon source. Ammonium
7 chloride, potassium nitrate, potassium nitrite or L-glutamate were used as nitrogen
8 source (10 mM each), as stated in the text. Bacteria were cultured in 250 mL flasks
9 containing 50 mL of medium that were rotated at 200 rpm. *E. coli* strains were grown at
10 37 °C in LB medium. Cell growth was followed measuring the absorbance of the
11 cultures at 600 nm (A_{600}) or by protein determination [34]. Antibiotic supplements were
12 used at the following concentrations ($\mu\text{g}\cdot\text{mL}^{-1}$): ampicillin (Amp), 100; kanamycin
13 (Km), 25; rifampicin (Rif), 100; spectinomycin (Sp), 25; streptomycin (Sm), 60;
14 tetracycline (Tet), 10; chloramphenicol (Cm), 50.

15

16 **Analytical methods**

17 Upon reaching exponential growth phase, a 50 mL culture volume was subject to cell
18 fractionation. Cells were harvested, washed twice with 50 mM Tris-HCl (pH 8.0) and
19 re-suspended in 10 mM Tris-HCl (pH 8.0), 500 mM sucrose and 3 mM EDTA.
20 Lysozyme (chicken egg white, EC 3.2.1.17) at $0.2\text{ mg}\cdot\text{mL}^{-1}$ final concentration and a
21 few grains of DNase I (bovine pancreas, EC 3.1.21.1) were added. This mixture was
22 incubated with shaking for 30 minutes at 30 °C. The sphaeroplasts formed were
23 harvested by centrifugation ($13000\times g$) for 15 minutes at 4 °C and the periplasm was
24 recovered. Sphaeroplasts were re-suspended in a volume of 10 mL 100 mM Tris-HCl
25 (pH 8.0) and disrupted by sonication, and cytoplasmic and membrane fractions were
26 separated by ultracentrifugation ($40000\times g$). Assimilatory NADH-dependent nitrate
27 reductase activity was assayed in cytoplasmic fractions in the presence of NADH as
28 electron donor by measuring the nitrite formed from nitrate [35]. β -galactosidase
29 activity was determined spectrophotometrically as previously described [36]. Protein
30 concentration was measured by using a Bradford protein assay kit (Bio-Rad, UK) with a
31 BSA standard (Fraction V, Sigma, UK).

32

33 **Microarray analysis of *P. denitrificans***

34 *P. denitrificans* genomic DNA was isolated using a Genomic DNA extracting kit and
35 100/G columns (Qiagen) from 10 mL exponential phase cells according with the
36 specification of the manufacturer. For RNA extractions, 30 mL of early-exponential
37 phase cells ($A_{600} \sim 0.4$) was added to 12 mL of ice-cold 95% ethanol/5% phenol (v/v)
38 solution, and incubated on ice for 30 minutes to prevent RNA degradation. Cells were
39 then pelleted and stored at -80 °C until RNA was extracted by using SV Total RNA
40 isolation kit (Promega). Trace DNA contamination was removed by treatment with
41 Turbo DNA-*free*TM DNase (Ambion), and this was confirmed by PCR amplification of
42 RNA samples using MyFiTM DNA polymerase (Bioline). Nucleic acids were quantified
43 spectrophotometrically in a Nanodrop 2000 (Thermo Scientific), and integrity of RNA
44 samples was analyzed using an ExperionTM Automated Electrophoresis platform
45 (BioRad) using RNA StdSens chips (BioRad). All standard protocols were carried out
46 according with the instructions of the manufacturers.

47 For labelling and hybridisation of microarray slides, total RNA (10 μg) from
48 three independent bacterial cultures were reverse-transcribed to cDNA using

1 AffinityScript multiple temperature reverse transcriptase (Agilent), and fluorescently
2 labelled using random primers (Invitrogen) to incorporate Cy5-dCTP (Amersham). *P.*
3 *denitrificans* genomic DNA (10 µg) was labelled with Cy3-dCTP (Amersham) using a
4 Bioprime[®] DNA labelling system (Invitrogen), prior to mixing (1:5) with labelled
5 cDNA and hybridised to custom-designed 4 x 44K oligonucleotide array slides
6 (Agilent). Hybridisation buffer (50 mM morpholine-4-ethanesulfonic acid pH 6.5, 1 M
7 NaCl, 20% w/v formamide, 20 mM EDTA, 1% w/v, Triton-X-100) mixed with Cy5-
8 and Cy3-dCTP labelled nucleic acids were loaded onto the GASKET slide prior to
9 placing the microarray slide in contact with the hybridization mix, which were sealed in
10 a tight chamber and incubated at 55 °C in a rotary hybridisation oven at 8 rpm, for
11 approximately 60 hours in the dark. Following hybridizations, slides were removed and
12 washed for 5 minutes in a microscope-slide chamber using a solution of 6 x SSPE (0.2
13 M phosphate buffer, 2.98 M NaCl, 20 mM EDTA, pH 7.4) supplemented with 0.005%
14 N-lauryl-sarcosine, followed by 5 minutes in a solution of 0.6 x SSPE supplemented
15 with 0.18% polyethylene glycol 200. Slides were then dried by centrifugation for 30
16 seconds.

17 For analyses and interpretation, microarray slides were scanned using a scanner
18 (GenePix 4000A, Axon Instruments) with excitation wavelengths of 532 nm and 635
19 nm. Fluorescent spots and background intensities were quantified using GenePix Pro
20 software (Axon Instruments) and filtered to omit those with a reference signal lower
21 than two standard-deviations from the background intensity in further analyses. Signal
22 intensities were corrected by subtracting the background and the red/green (Cy5/Cy3)
23 ratios. All datasets were normalised using Batch Anti-Banana Algorithm in R (BABAR)
24 which uses cyclic loess to normalise across the complete dataset [37] and analyzed
25 using Gene Spring 7.3 (Agilent) to filter genes that were differentially expressed ≥ 2 -
26 fold with significance of $\geq 95\%$ across three independent cultures. These data are
27 represented in Figure 1, where \log_2 of normalised expression values are shown as a
28 heat-map. Microarrays were validated by qPCR with oligonucleotide primers that
29 annealed to internal regions of the *nas* genes (Table S2), as described below. Primers
30 were designed using Primer³ Plus software [38], to amplify products between 100 and
31 150 bp, with a T_m of about 60 °C and used at a final concentration of 0.4 µM. The
32 relative fold-change values were calculated by using amplification efficiencies, as
33 described previously [39].

34

35 **Proteomic analysis**

36 Two-dimensional gels electrophoresis (2D-PAGE) was performed with sample
37 preparations that were obtained from *P. denitrificans* cells grown to mid-log phase in
38 minimal medium with different nitrogen sources as previously described [2].
39 Subcellular fractionation was carried as described above, and isolated cytoplasmic
40 fractions containing 250 µg protein were used to rehydrate 11 cm strips (Immobiline
41 DryStrips with the appropriate pH range; Amersham Biosciences) for 12 h. Isoelectric
42 focusing was carried out in an IPGphor (Pharmacia) until 20000 volt-hours were
43 reached. After isoelectric focusing, strips were equilibrated as previously described [40]
44 and applied to 12% (v/v) polyacrylamide gels. Second dimension SDS-PAGE was
45 performed by using the Hoefer SE600 system (Amersham Biosciences) and gels were
46 stained using the Brilliant Blue G-colloidal concentrate (Sigma) and scanned with a
47 Molecular Image FX (Bio-Rad). Triplicate 2D-PAGE separations were generated for
48 each sample. Protein identification was carried out in the UCO-SCAI Proteomic Centre,

1 University of Córdoba (Spain), a member of ProteoRed network. Protein spots of
2 interest were excised automatically in a ProPic station (Genomic Solutions, UK) and
3 samples were automatically digested with trypsin according with standard protocols in a
4 ProGest station (Genomic Solutions), and analyzed in a 4700 Proteomics Analyzer
5 MALDI-TOF/TOF mass spectrometer (Applied Biosystems), in the m/z range 800 to
6 4000, with an accelerating voltage of 20 kV, in reflectron mode and with delayed
7 extraction set to “on” and an elapsed time of 120 ns. Proteins were identified by peptide
8 mass fingerprinting (PMF) and confirmed by MS/MS analysis of the three most
9 abundant peptide ions. MASCOT searching engine (Matrixscience, UK) was used for
10 protein identification over the non-redundant NCBI database of proteins. The
11 confidence in the peptide mass fingerprinting matches ($p < 0.05$) was based on the
12 MOWSE score (higher than 65) and CI > 99.8%, and confirmed by the accurate
13 overlapping of the matched peptides with the major peaks of the mass spectrum.

14 15 **Routine DNA manipulations and site-directed mutagenesis of *nasA* leader** 16 **sequence**

17 Genomic and plasmid DNA were routinely isolated and purified using the Wizard®
18 Genomic DNA purification kit (Promega, USA) and the Qiagen plasmid kit (Qiagen,
19 Germany), respectively. Custom oligonucleotide primers, listed in Table S2, were
20 supplied by Invitrogen (Paisley, UK) and the polymerase chain reaction (PCR) was
21 performed using the Expand High Fidelity PCR system (Roche, Switzerland) with 5%
22 DMSO added as standard.

23 In the mutational analysis carried out *in trans* of the *nasA* leader region, a *nasA*
24 promoter transcriptional fusion was constructed by using an intergenic *nasS-nasA* 392-
25 bp fragment amplified by PCR with oligonucleotides FA1_A/RA1, which were located
26 at the 3'-end of *nasS* and at the 5'-end of *nasA*, respectively. This fragment was cloned
27 within the pSparkII vector and subcloned as *SphI-PstI* in the promoter probe vector
28 pMP220, generating pMP220-*PnasA* (Table S1). Different mutated versions of *PnasA-*
29 *lacZ* were also constructed by using oligonucleotides containing substitutions of three
30 bases for two hairpins, HI or HII, identified in the *nasA* gene leader region (HI 137 bp
31 and HII 91 bp upstream 5'-end *nasA* gene, respectively). Oligonucleotides FA1_B/RHI
32 and FHII/RA1 were used for mutagenesis of hairpins I or II, respectively, and the PCR
33 products amplified with these oligonucleotides were cloned into the pSparkII vector.
34 The pMP220-*PnasA* vector was digested with *PstI/AscI* and used to clone the mutated
35 fragment of the hairpin I previously liberated from pSparkII with *PstI/AscI* restriction
36 enzymes, obtaining the pMP220-*PnasA1* construct. The pMP220 plasmid is a non-
37 integrative vector. For mutagenesis of hairpin II, the corresponding mutated fragment
38 was liberated from pSparkII with *AscI/SphI* and cloned into pMP220-*PnasA* previously
39 digested with *AscI/SphI*, obtaining pMP220-*PnasA2* plasmid. All constructs were
40 sequenced (UCO-SCAI, University of Córdoba) to confirm the native or the mutated
41 sequences and the final constructs were introduced into *P. denitrificans* PD1222 by
42 triparental mating [2].

43 In the mutational analysis carried out *in cis* of the ANTAR region (hairpin I), the
44 DNA regions upstream and downstream this secondary structure were amplified by
45 PCR with oligonucleotide pairs AIF1/AIR1a and AIF2/AIR2, respectively. These two
46 fragments were cloned separately into the pSparkII cloning vector. The upstream
47 fragment (524 bp) was subcloned as *EcoRI/SalI* into pK18mobsacB, a kanamycin-
48 resistant suicide vector in *Paracoccus denitrificans*. The downstream fragment (578 bp)
49 was subcloned as *SalI/HindIII* into the previous construction. The final construction

1 contained a 35 bp sequence corresponding to the multicloning site of the pSparkII
2 vector replacing the native hairpin I sequence (37 bp). This plasmid was used for
3 triparental mating with the receptor strain *Paracoccus denitrificans* and the helper strain
4 *E. coli* containing the pRK2013 plasmid. First transconjugant selection was carried out
5 in LB media with spectinomycin and kanamycin. After that, a sucrose selection for
6 double-crossover events was performed. Finally, mutation was confirmed by PCR and
7 enzyme restriction analyses. The wild-type *nasA* leader region with hairpin I was
8 amplified with oligonucleotides FA1_A/RA1. This wild-type fragment was not digested
9 by *SalI*, whereas the fragment amplified from the mutant was digested by this restriction
10 enzyme because it contains the *SalI* recognition sequence of the pSparkI multicloning
11 site.

12

13 **Site-directed mutagenesis of the NtrC binding sequence**

14 The native *nasT* promoter, as well as a mutated version affected in the putative NtrC
15 binding site located upstream of *nasT* gene, were used to generate transcriptional
16 fusions by PCR-driven overlap extension. Initial PCRs were carried out with
17 oligonucleotides PTF/PTNtrR and PTR/PTNtrF for the native construct, and
18 PTF/PTNtrR_Mut and PTR/PTNtrF_Mut for the mutated version. Internal primers
19 (PTNtr) were used to generate overlapping and complementary 3'-ends on the
20 intermediate segments. In the case of the mutated *nasT* promoter, the PTNtr primers
21 were also used to introduce the mutated bases. Overlapping strands hybridize at this
22 region in a subsequent PCR and were extended to generate the full-length product (619-
23 bp) amplified by flanking primers that include restriction enzyme sites for cloning as
24 *PstI/SphI* into the promoter probe vector pMP220. The final constructs for the analysis
25 of the native *nasT* promoter and the NtrC-binding site mutated version of *nasT*
26 promoter were called pMP220-*PnasT* and pMP220-*PnasT1*, respectively. All plasmids
27 were checked by sequencing (UCO-SCAI, University of Córdoba) and the final
28 constructs were introduced into *P. denitrificans* PD1222 by triparental mating [2].
29

30

30 **RT-PCR and qPCR reactions**

31 *P. denitrificans* wild-type and mutant strains were cultured in minimal medium with
32 succinate as carbon source and different nitrogen sources. Cells were harvested ($A_{600} \sim$
33 0.4) and washed in TEG buffer with 25 mM Tris-HCl (pH 8.0), 1% glucose and 10 mM
34 EDTA. RNA isolations were performed following the Qiagen RNA extraction kit
35 (RNeasy midi kit). DNase incubation was carried out in the column with RNase-free
36 DNase set (Qiagen) and an additional post-column treatment was required with DNase I
37 (Ambion). The concentration and purity of the RNA samples were measured in a
38 ND1000 spectrophotometer (Nanodrop Technologies). Synthesis of total cDNA was
39 achieved in 20 μ L final volume, containing: 500 ng RNA, 0.7 mM dNTPs, 200 U
40 SuperScript II Reverse Transcriptase (Invitrogen) and 3.75 μ M random hexamers
41 (Applied Biosystems). Samples were initially heated at 65 °C for 5 min and then
42 incubated at 42 °C for 50 min, followed by incubation at 70 °C for 15 min. To carry out
43 PCR reactions, 2 μ L of each cDNA were initially heated at 98 °C for 2 min, followed by
44 30 cycles of amplification: 98 °C, 30 s; 60 °C, 30 s and 69 °C, 1 min. Polymerase
45 extension reactions were completed by an additional incubation at 69 °C for 10 min. For
46 real-time assays, the cDNA was purified using Favorprep Gel/PCR purification kit
47 (Favorgen) and the concentration was measured using a ND1000 spectrophotometer.
48 The iQ5 Multicolor Real-Time PCR Detection System (Bio-Rad) was used in a 25 μ L
49 reaction (final volume), containing 2 μ L diluted cDNA (12.5, 2.5 and 0.5 ng), 0.2 μ M of

1 each primer (Table S2), and 12.5 μ L iQ SYBR Green Supermix (Bio-Rad). Target
2 cDNAs and reference samples were amplified three times in separate PCR reactions.
3 Samples were initially denatured by heating at 95 $^{\circ}$ C for 3 min, followed by 40 cycles of
4 amplification (95 $^{\circ}$ C, 30 s; test annealing temperature, 60 $^{\circ}$ C, 30 s; elongation and signal
5 acquisition, 72 $^{\circ}$ C, 30 s). For relative quantification of the fluorescence values, a
6 calibration curve was made using dilution series from 100-0.001 ng of *P. denitrificans*
7 genomic DNA sample. Represented data were normalized by using the *rpoB* and the
8 *dnaN* genes as housekeeping (Table S2). Primer design and relative fold-change values
9 calculation were carried out as previously described [38,39].

10

11 **Generation of *ntrB* and *ntrY* mutant strains of *P. denitrificans***

12 Mutant strains of *P. denitrificans* were generated by significant allelic replacement with
13 specific antibiotic resistance markers. The *ntrB* mutant (*ntrB* Δ ::Sm) was generated by
14 PCR amplification of the 3'-end (682 bp) and 5'-end (533 bp) of the *ntrB* gene with the
15 respective set of primers ntrB1/ntrB2 and ntrB3/ntrB4 (Table S2). The PCR products
16 were cloned within the pGEM-T Easy vector and then assembled within pUC18 vector
17 (Table S1). A unique *Bam*HI restriction site was generated at the interface of the 3'-end
18 and the 5'-end of *ntrB* gene to generate a 685 bp deletion and to insert a streptomycin
19 resistance cassette from the pSRA2 vector (Table S1). A single *Eco*RI fragment
20 (*ntrB* Δ ::Sm) was transferred to pSUP202 to produce the final mobilizable vector.

21 For the *ntrY* mutant strain (*ntrY* Δ ::Km), fragments containing the 3'-end (510
22 bp) and 5'-end (717 bp) of *ntrY* were produced by PCR using the respective primer
23 pairs ntrY1/ntrY2 and ntrY3/ntrY4 (Table S2) and cloned within the pGEM-T vector
24 (Table S1). The upstream region was screened by restriction digestion and selected by
25 using an *Eco*RI recognition site and the downstream region of *ntrY* was introduced as an
26 *Aat*II-*Bam*HI fragment, yielding a unique *Bam*HI restriction site between both regions,
27 generating a 1035 bp deletion. A kanamycin resistance cassette obtained from the
28 pSRA2 vector (Table S1) was inserted as a *Bam*HI fragment between these regions of
29 *ntrY*. Finally, the *Aat*II-*Eco*RI fragment (*ntrY* Δ ::Km) was cloned into the mobilizable
30 vector pSUP202. This construct was transferred to either the wild-type strain or the *ntrB*
31 mutant to generate the single *ntrY* or the double NtrB/NtrY mutant, respectively.
32 Plasmids were analyzed by DNA-sequencing (UCO-SCAI, University of Córdoba) and
33 mutants were checked by PCR by using primer pairs that amplify for a product
34 containing the deletion-insertion mutation. Mutants showed bands higher than those
35 presented in the wild type because the inserted resistance cassettes were larger than the
36 deleted fragments, confirming allelic exchange by double-crossover mechanism.

37

1 RESULTS

2 3 **Transcriptomic differences between bacteria cultured aerobically under nitrate-** 4 **versus ammonium-dependent anabolism**

5 To investigate the underlying biochemical adaptation to nitrate assimilation compared
6 to ammonium-dependent anabolism, transcriptomic analysis was undertaken through
7 microarray analysis of RNA isolated from *P. denitrificans* cells grown in batch culture
8 with 20 mM succinate and either 10 mM nitrate or 10 mM ammonium as the sole N-
9 source. Of the 5077 protein-coding genes that are present in the genome of *P.*
10 *denitrificans* PD1222, only 83 genes (1.64 % of the whole genome) displayed at least 2-
11 fold expression differences at a 95% significance level between the two growth
12 conditions tested. Among these, 74 genes were induced during nitrate-dependent
13 growth, while 9 genes were down-regulated (Fig. 1A).

14 The genes induced when nitrate was the only nitrogen source belonged to 30
15 gene clusters, including among others, the *gltB* and *gltD* genes encoding the small and
16 large subunits of the glutamate synthase (GOGAT), the *amtB* gene that codes for a high-
17 affinity ammonium transporter, the *ntrB* and *ntrC* genes that code for the general
18 nitrogen regulatory Ntr system, the *glnB* gene that codes for the nitrogen regulatory
19 protein PII, the *glnA* gene encoding the glutamine synthetase (GS), the *bztA-D* gene
20 cluster involved in glutamate/aspartate transport, the *ureA-GJ* gene cluster for urea
21 metabolism, the *urtA-E* gene cluster for urea transport, the *potDF-I* gene cluster
22 involved in polyamine transport, the *braC-G* gene cluster involved in branched-chain
23 amino acids transport, the *dctPQM* genes encoding a TRAP dicarboxylate transporter
24 system, the respiratory nitrate reductase *narGJ* genes, and the *nasTSABGHC* gene
25 cluster for nitrate/nitrite assimilation (Fig. 1A). The genes down-regulated during
26 growth with nitrate included the *gdhA* gene that encodes the glutamate dehydrogenase
27 (Fig. 1A). Expression of a selection of these genes was also quantified by qPCR and
28 corroborated well with the microarray data (Table 1). The genes for which expression
29 was highly increased on nitrate included *nasA*, *nasH*, *narG*, *narJ*, *nasT*, *nasS*, *urtC*,
30 *amtB*, *ureD*, *glnA*, *glnB*, *gltD*, and *ntrC*. The *nasT* gene was more strongly induced by
31 nitrate than the *nasS* gene (Table 1).

32 Qualitative proteomic analysis by 2D-PAGE of soluble fractions of *P.*
33 *denitrificans* cells grown with nitrate or ammonium as the sole N-source confirmed that
34 a significant number of the key changes in the transcriptome could be also indentified
35 in the proteome (Fig. 1B). The up-regulated proteins for which higher levels of
36 synthesis in the presence of nitrate could be confirmed included the nitrogen regulatory
37 protein GlnB (PII), the periplasmic component of the urea ABC-type transporter (UrtA),
38 the polyamine transporter component (PotD), the glutamate/aspartate transporter
39 component (BztA), and two proteins encoded by the *nas* gene cluster of *P. denitrificans*
40 (NasG and NasB). A decreased synthesis of the glutamate dehydrogenase (GdhA) was
41 observed in the nitrate grown cells compared to the ammonium-grown cells (Fig. 1B).

42 43 **Analysis of the expression of the *nasTS* and *nasABGHC* transcripts**

44 The transcriptome differences between nitrate- and ammonium-anabolising cells
45 suggest a general adaptation to metabolism of secondary (non-ammonium) nitrogen
46 sources, nitrogen scavenging and the need to search for organic carbon to provide the
47 reductant for nitrate assimilation. Alongside this is a more specific adaptation to nitrate
48 anabolism through expression of the *nasABGHC* genes. Therefore, the control of the

1 *nas* gene expression was investigated. The regulatory pattern for *P. denitrificans nas*
2 gene expression was analyzed at the transcriptional level by RT-PCR analysis with
3 RNA from cells grown with nitrate, glutamate, ammonium, or ammonium plus nitrate
4 as N-source. Primer pairs were designed to explore transcription across the 3'- and the
5 5'-ends of different *nas* gene boundaries (Fig. 2A). The PCR products were detected
6 across the *nasTS*, *nasAB*, *nasBG*, *nasGH*, *nasHC*, and *nasGHC* boundaries (Fig. 2B).
7 However, PCR product was undetectable when the primers located at the 3'-end of the
8 *nasS* gene and the 5'-end of the *nasA* gene were used. These results suggest that the
9 *nasTSABGHC* gene cluster comprises two different transcriptional units, one
10 corresponding to the regulatory *nasTS* genes and the other constituted by the structural
11 *nasABGHC* genes.

12 The RT-PCR analysis also indicated that the *nasA-C* transcript is found in cells
13 grown with either nitrate or glutamate as sole N-source, but not when ammonium was
14 present (Fig. 2). However, a high glutamine concentration (5 mM) or a 2-
15 oxoglutarate/glutamine 5 mM/5 mM ratio did not repress expression of the *nas* genes.
16 The *nasA* gene expression, determined by RT-qPCR, was similar in cells grown with
17 glutamate or glutamate plus nitrate (Fig. 2C). The *nasTS* transcript was detected in cells
18 grown with all nitrogen sources tested, although only at very low levels in the presence
19 of ammonium (Fig. 2C).

20 A proteomic approach was applied to confirm the presence of polypeptides
21 encoded by the *nas* genes in cytoplasmic fractions of *P. denitrificans* cells grown with
22 different nitrogen sources. The Rieske-type protein NasG required for both nitrate and
23 nitrite reductase activities (Fig. 3A) and the catalytic subunit of the nitrite reductase
24 NasB (not shown) were only detected when nitrate was present in the media and
25 ammonium was absent. NADH-dependent nitrate reductase activity was determined in
26 the cytoplasmic fraction from cells grown with different nitrogen sources. In accordance
27 with the proteomic analysis, the nitrate reductase activity was detected only in presence
28 of nitrate and absence of ammonium (Fig. 3A). It is notable that although the
29 *nasABGHC* transcript was detected in cells grown with glutamate in absence of nitrate
30 (Fig. 2B), the NasG and NasB proteins and the nitrate reductase activity were not
31 detectable under these conditions (Fig. 3A).

32 33 **Analysis of *nas* gene expression in the *nasT* and *nasS* mutants**

34 It has been previously described that a *nasT* mutant of *P. denitrificans* is unable to grow
35 with nitrate as the sole nitrogen source, whereas a *P. denitrificans nasS* mutant grows
36 with nitrate as the sole N-source with similar rate and yield to the wild-type strain [5].
37 To analyze the regulatory effect of nitrate in the *nasS* and *nasT* mutants, a comparison
38 of *nas* gene expression between cells grown with glutamate versus cells grown with
39 glutamate plus nitrate was performed. RT-qPCR analysis showed that the *nasT* mutant
40 lacks the *nasABGHC* transcript (Fig. 3B). Accordingly, 2D-PAGE revealed that this
41 mutant strain is also devoid of the NasG polypeptide (Fig. 3C). It is noticeable that the
42 respiratory nitrate reductase *narG* and *narJ* genes were up-regulated in the *nasT* mutant
43 strain, whereas genes coding for glutamine synthetase (*glnA*) and PII regulatory protein
44 (*glnB*) were down-regulated (Table 2), suggesting that NasT might also serve as part of
45 a more complex global regulatory system.

46
47 In the *P. denitrificans nasS* mutant, the *nasABGHC* transcript was detected in all
48 nitrogen sources, except when ammonium was used as N-source, similarly to the wild-
49 type strain (Fig. 3B). The NasG polypeptide and the nitrate reductase activity were

1 detected in the *nasS* mutant strain grown with glutamate both in presence and absence
2 of nitrate (Fig. 3C), whereas in the wild-type strain the NasG polypeptide and the nitrate
3 reductase activity were detected only in the presence of nitrate (Fig. 3A). In addition,
4 RT-qPCR analysis revealed that the *nasS* mutant strain showed significantly increased
5 levels of the *narG* and *gdhA* genes in response to nitrate (Table 2).

7 **Identification of putative *cis*-regulatory RNA secondary structures in the *nasA*** 8 **leader region**

9 Between the regulatory *nasTS* genes and the structural *nasABGHC* genes lies a ~200 bp
10 non-coding region with putative hairpin structures that could lead to transcription
11 termination (Fig. 4A). Secondary structures in the RNA leader sequence of *nasA* are
12 thought to terminate transcription [14,15], but binding of NasT could allow
13 transcription antitermination. Putative hairpin structures that could be a target for the
14 transcriptional antiterminator NasT in the *nasA* leader region were identified by using
15 the RNAfold web server <http://rna.tbi.univie.ac.at/cgi-bin/RNAfold.cgi>. To explore the
16 importance of this region, several *lacZ* transcriptional fusions were constructed and
17 expressed in *P. denitrificans* cells grown with either nitrate or ammonium as sole N-
18 source. In cells grown under nitrate anabolizing conditions, the mutation of the HI
19 hairpin resulted in a severe deficiency in the transcriptional activity. However,
20 transcriptional activity was relatively unaffected in the HII mutation (Fig. 4B). In cells
21 grown with ammonium the transcriptional activity was 100-fold lower than in nitrate-
22 assimilating cells, but there was a two-fold increase in cells carrying the HII mutation.
23 These results suggest that the HI hairpin is important for NasT binding, and that HII
24 structure may contribute to transcription attenuation. To corroborate the importance of
25 HI, a specific chromosomal mutation *in cis* of the putative NasT binding region was
26 introduced. This chromosomal mutant lost the parental capacity to grow with nitrate as
27 the sole nitrogen source (Fig. 4C). Controls established that the chromosomal HI mutant
28 strain showed a similar growth to the parental strain with ammonium as nitrogen source.

30 **The contribution of *ntrBCYX* to the regulation of nitrate assimilation**

31 In addition to *nasTS* genes, the *ntrBC* genes were also up-regulated during nitrate-
32 dependent anabolism compared to cells grown with ammonium (Fig. 1A, Table 1). This
33 suggests an integrated regulation of nitrate metabolism between these two systems. The
34 *ntrBC* genes are clustered together with the *ntrYX* genes. The *ntrB* gene encodes a
35 signal transduction protein (386 residues) with histidine kinase activity involved in
36 nitrogen metabolism, and the *ntrC* gene codes for a Fis-like σ^{54} -dependent regulator
37 receiver protein (486 residues). Similarly, the *ntrY* gene codes for a multi-sensor signal
38 transduction regulator (709 residues) with histidine kinase activity and the *ntrX* gene
39 encodes a σ^{54} -dependent regulatory protein (463 residues). A non-coding region of 217
40 bp lies between *ntrC* and *ntrY* genes, whereas *ntrY* and *ntrX* genes overlap by 3 bp.
41 NtrB and NtrC homologues have been found in many different microorganisms (Fig.
42 S1). Although the NtrYX system is not widely distributed among bacteria, the *P.*
43 *denitrificans* NtrY protein showed the highest identity (41%) with *Azorhizobium*
44 *caulinodans* NtrY, whereas the *P. denitrificans* NtrX protein showed the highest
45 identity (56%) with its homologues in *Azorhizobium caulinodans* and *Azospirillum*
46 *brasilense* (Fig. S2).

47 Isolation and characterization of single *ntrB* and *ntrY* mutants and double
48 *ntrB/ntrY* mutant of *P. denitrificans* were carried out. All mutant strains displayed
49 similar growth rates and yields to that shown by the wild-type strain when ammonium

1 was used as nitrogen source. However, the *ntrY* mutant was only slightly affected on its
2 growth on nitrate, whereas the *ntrB* mutant showed a longer lag-phase than the wild-
3 type, and the double *ntrB/ntrY* mutant was unable to grow with nitrate (Fig. 5A).

4 Expression of the *nasT*, *nasS*, *ntrB*, *ntrY* regulatory genes and the structural *nasA*
5 gene was analyzed by RT-qPCR in the *ntrB* and *ntrY* mutants and the wild-type strain
6 of *P. denitrificans*. In the *ntrB* mutant *nasTS* and *nasA* were down-regulated (Fig. 5B),
7 whereas in the *ntrY* mutant all these genes were expressed similarly to the wild-type
8 strain (not shown). The assimilatory NADH-dependent nitrate reductase activity was
9 determined in cytoplasmic fractions from wild-type, *ntrB* and *ntrY* mutants grown with
10 nitrate, when all strains reached similar biomass (protein concentration about 150 mg
11 mL⁻¹). The *ntrY* mutant showed similar levels of activity to that presented by the wild-
12 type, about 4 nmol NO₂⁻ formed min⁻¹ mg⁻¹, whereas the *ntrB* mutant only presented an
13 activity of about 1 nmol NO₂⁻ formed min⁻¹ mg⁻¹.

14 Transcriptional fusions to the β-galactosidase-encoding gene *lacZ* were carried
15 out in the *nasT* leader sequence and expressed in the wild-type, *ntrB* and *ntrY* mutant
16 strains of *P. denitrificans*. The β-galactosidase activities in the *ntrY* mutant were very
17 similar to those found in the wild-type strain (Table 3). However, β-galactosidase
18 activities in the *P. denitrificans ntrB* mutant were much lower than those described for
19 the wild-type strain (Table 3).

20
21 These data strongly suggest a role for NtrBC in the regulation of nitrate
22 assimilation at the level of expression of *nasT* and so the possibility of a *cis*-acting
23 binding site for NtrC was explored. The RegPrecise program for collection of manually
24 curated inferences of regions in prokaryotic genomes identified a very well conserved
25 putative NtrC binding site (5'- (T/C)GC(C/A)NNNNTTNNT(G/A)GCA-3') in the
26 leader regions of a number of the different genes in *P. denitrificans*, including the
27 *glnBA*, *amtB*, and *nasTS* genes. These genes were upregulated during nitrate
28 assimilation, as revealed by the transcriptomic analyses (Fig. 1A, Table 1). Several base
29 pairs were mutated in the putative NtrC binding site located in the *nasT* leader
30 sequence, resulting in the loss of the up-regulation observed under nitrate- or glutamate-
31 dependent growth conditions (Table 3). These results highlight the importance of this
32 sequence motif in the regulation of nitrate assimilation.

33
34

1 DISCUSSION

2
3 The soil denitrifier *P. denitrificans* can use nitrate as the sole N-source by an
4 assimilatory nitrate reduction system. The process of nitrate assimilation in this
5 bacterium has been comprehensively investigated by comparative transcriptomic and
6 proteomic analyses in cells grown under nitrate- or ammonium-assimilating conditions
7 (Figs. 1, 3 and S3). Comparing nitrate- to ammonium-grown cells, a key metabolic
8 change was the decrease in the expression of glutamate dehydrogenase, the enzyme
9 related with the low affinity ammonium assimilation pathway. This was accompanied
10 by induction of components for organic nitrogen scavenging, urea uptake and
11 utilisation, high affinity ammonium uptake and assimilation, regulatory and structural
12 components of inorganic nitrate and nitrite assimilation, and scavenging of
13 dicarboxylate organic acids to provide the additional reducing equivalents required for
14 nitrate assimilation (eight electrons per nitrate ion converted into ammonia).

15 In *P. denitrificans*, the regulatory *nasTS* genes are located at the 5'-end of the *P.*
16 *denitrificans nas* gene cluster for nitrate assimilation (Fig. 2A). RT-PCR and qPCR
17 analyses of the *P. denitrificans nasTS* and *nasABGHC* genes demonstrate that these
18 genes are transcribed as separated transcriptional units. The existence of two
19 independent transcriptional units in the *P. denitrificans nas* region is also supported by
20 the presence of a non-coding region between the regulatory *nasTS* genes and the
21 structural *nasABGHC* genes, with two putative hairpins (HI and HII) that could lead to
22 secondary structures in this mRNA region (Fig. 4). These two transcriptional units show
23 different regulation, since expression of the regulatory *nasT* and *nasS* genes was
24 observed in all N-sources tested, including ammonium, although in general at very low
25 levels, whereas *nasABGHC* transcript was detected either with nitrate or glutamate, but
26 it was undetectable with ammonium independently of the presence or absence of nitrate
27 (Fig. 2). However, although the *nasABGHC* transcript was present in wild-type cells
28 grown with glutamate as the sole N-source, the Rieske-type NasG protein and the large
29 subunit of the assimilatory nitrite reductase NasB, required for both nitrate and nitrite
30 reductase activities, were not detected by 2D-PAGE and these proteins were only
31 observed in the presence of nitrate. Accordingly, the assimilatory NADH-dependent
32 nitrate reductase activity was not detected when glutamate was used as the sole nitrogen
33 source (Fig. 3A). These results suggest that nitrate is not an obligate inducer of the
34 *nasABGHC* gene expression, and that it may also play a positive role at the translational
35 level.

36 Mutational analysis of the *nasT* gene of *P. denitrificans* reveals that NasT could
37 act as transcription antiterminator because the *nasT* mutant strain is unable to grow with
38 nitrate as the sole N-source and lacks both *nasABGHC* transcript and NADH-nitrate
39 reductase activity (Fig. 3BC). Also, the strain harbouring the chromosomal mutation in
40 a putative *cis*-acting NasT binding site (hairpin I) was unable to grow with nitrate (Fig.
41 4C). Therefore, in this work we have identified the RNA region where NasT is likely to
42 bind as a transcriptional antiterminator. On the contrary, the *nasS* mutant strain of *P.*
43 *denitrificans* is capable to use nitrate as the sole N-source, similarly to the wild-type
44 strain, but shows a deregulated nitrate reductase activity that can be detected
45 independently of the presence or absence of nitrate (Fig. 3BC). In addition, assimilatory
46 nitrate reductase activity in the *P. denitrificans nasS* mutant is abolished in the presence
47 of ammonium, as previously described in the wild-type strain [5]. Expression of the
48 *nasT* gene is higher than *nasS* gene expression (Tables 1 and 2), probably because

1 strong secondary structures present at the 5'-end of the *nasS* gene might provoke RNA
2 polymerase to be occasionally released without completing the whole transcript.

3 *P. denitrificans* NasS and NasT proteins expressed in *E. coli* have been purified
4 as a tetrameric complex in the absence of nitrate or nitrite, but in addition to the NasT-
5 NasS complex, free NasT protein can be also detected [5]. Therefore, in the absence of
6 nitrate, NasS and NasT form a protein-protein complex, probably with two units of each
7 protein, which limits binding of NasT to the *nas* mRNA to act as transcription
8 antiterminator. However, in the presence of nitrate, NasS clamps the oxyanion,
9 changing conformation and dissociates from NasT, thus increasing the size of the free
10 NasT pool available to serve for *nasABGHC* transcription antitermination [5]. However,
11 the elevated *nasT* gene expression over the *nasS* gene could lead to an excess of NasT
12 protein over the NasS protein. This unbalance implies that there is always enough free
13 NasT to allow transcription of *nasABGHC* genes in the absence of both nitrate and
14 ammonium (i.e., glutamate as N-source), suggesting additional global regulatory roles.
15 In the *P. denitrificans nasT* mutant the structural *nasABGHC* transcript was absent in all
16 N-sources tested (Fig. 3B). This is consistent with a role for NasT as transcription
17 antiterminator and with the phenotype of the *P. denitrificans nasT* mutant, which is
18 unable to grow with nitrate or nitrite as the sole N-source. In addition, up-regulation of
19 *narG* and *narJ* genes and down-regulation of *glnAB* genes in the *nasT* mutant also
20 indicated that NasT may be involved in other regulatory processes. Recently, it has been
21 suggested in the soybean endosymbiont *Bradyrhizobium japonicum* that *nasST* genes
22 regulate respiratory nitrous oxide and periplasmic nitrate reductases [41]. Therefore,
23 additional targets for NasT may exist in *P. denitrificans*, which could function as
24 negative regulators of *nar* gene expression.

25 In the *nasS* mutant the *nasABGHC* gene expression is no longer under nitrate
26 control because in the absence of the NasS protein, all the NasT pool is free to serve as
27 transcription antiterminator under all growth conditions (Fig. 3BC). Thus, in contrast to
28 the wild-type strain, the *nasS* mutant of *P. denitrificans* displayed in 2D-PAGE gels the
29 assimilatory nitrite reductase NasB and the Rieske-type NasG polypeptides when
30 glutamate was used as nitrogen source, independently of the presence or absence of
31 nitrate (Fig. 3C). These results corroborate that nitrate exerts a regulatory control at
32 post-transcriptional (translational) level in addition to the role of NasTS in the
33 transcription antitermination.

34 The phenotypes of the *ntrB*, *ntrY* and *ntrB/ntrY* mutants of *P. denitrificans*
35 suggest that the NtrBC system is mainly required for nitrate assimilation, while the
36 NtrYX system only has a slight contribution (Fig. 5A). The regulatory *nasTS* genes and
37 the structural *nasA* gene are down-regulated in the *ntrB* mutant of *P. denitrificans* (Fig.
38 5B), suggesting that these genes are under the control of the NtrBC system. This idea
39 has been corroborated by using the *lacZ* gene transcriptionally fused to native and
40 mutated *nasT* leader sequence (Table 3). These results indicate that, in the absence of
41 ammonium, phosphorylated NtrC could bind to the *nasT* promoter region, where a
42 consensus NtrC-binding site has been identified. The absence of the *nasABGHC*
43 transcript in cells grown with ammonium also suggests that NtrC or NrtX may bind to
44 an, as yet unidentified, site in the *nasA* leader region. According with the RegPrecise
45 program, very well conserved putative NtrC binding sites are found in the promoter
46 regions of the *nasT* and *ntrBC* genes, suggesting that these two gene clusters are under
47 the regulation of the general nitrogen control.

1 The increased synthesis of *P. denitrificans nasABGHC* genes reflects a system-
2 specific response to ammonium-limited growth in which nitrate is initially the sole
3 available N-source [2]. This bacterium also uses nitrate as terminal electron acceptor by
4 a membrane-bound nitrate reductase [1]. This respiratory nitrate reductase (Nar) can
5 substitute for NasC (assimilatory nitrate reductase) during anaerobic growth. The
6 present study shows that transcription of *nasTS* occurs at a low level in the presence of
7 ammonium, whereas transcription of *nasABGHC* genes is fully repressed in cells grown
8 with ammonium (Fig. 2). In *K. oxytoca*, *nasR* expression is sensitive to ammonium
9 since it is induced by the NtrBC system when ammonium is absent [14]. In *Azotobacter*
10 *vinelandii* expression of *nasST* genes is also sensitive to ammonium, although is not
11 under the control of the Ntr system [17]. Nevertheless, the capacity for nitrate
12 assimilation is lost in several *ntr* mutants [19]. Recently, it has been demonstrated that
13 NtrBC and NasST co-regulate the *nasAB* genes required for nitrate/nitrite uptake and
14 reduction in *A. vinelandii* [20]. Bioinformatic analysis has revealed that the two-
15 component NasTS system is a regulatory mechanism for nitrate assimilation that is
16 widely spread among bacteria. Particularly, NasT is mainly present in α -proteobacteria
17 like Rhizobiales and Rhodobacterales, and β -proteobacteria like Burkholderiales [5].
18 NasTS is involved in the regulation of the assimilatory nitrate/nitrite reductase genes in
19 response to nitrate or nitrite in *Azotobacter vinelandii*, *Pseudomonas putida* and
20 *Rhodobacter capsulatus* [17,18,42]. However, in *Azotobacter vinelandii* these genes are
21 arranged in the *nasST* orientation, instead of the *nasTS* organization found in *P.*
22 *denitrificans*, and they do not cluster together with the *nas* genes encoding the
23 nitrate/nitrite uptake and reduction system [43]. The NasS protein is homologous to
24 proteins which belong to the ATP-dependent nitrate transport systems found in
25 cyanobacteria, although NasS lacks the signal peptide for periplasmic translocation.
26 This is consistent with the role of NasS as a nitrate/nitrite sensor rather than acting as
27 the periplasmic component of the transport system for these oxyanions. The NasT
28 protein is a regulator with the ANTAR binding domain characteristic of transcription
29 antiterminators like the *Klebsiella oxytoca* NasR protein [4,14,16]. However, the *nasTS*
30 genes are absent in the *nas* cluster of *Klebsiella*, in which they are functionally replaced
31 by *nasR*, a gene mainly found in γ -proteobacteria. It has been proposed that NasR is
32 both a nitrate/nitrite sensor and a transcription antiterminator. A hairpin structure in
33 mRNA upstream from the *K. oxytoca nas* structural genes causes early termination of
34 transcription [14]. It has been suggested that in the presence of nitrate or nitrite NasR
35 binds to the mRNA transcript preventing hairpin formation. Additional novelty on the
36 regulatory mechanisms for nitrate assimilation was found in *P. denitrificans* because
37 repression of structural *nas* genes by ammonium could be directly exerted by
38 ammonium itself rather than by glutamine or the glutamine/2-oxoglutarate ratio. Under
39 low nitrogen conditions (absence of ammonium), the NtrB and NtrY proteins become
40 active to phosphorylate their respective partners, NtrC or NtrX, which in turn, become
41 active for binding to promoter regions of the regulated genes. In *R. capsulatus* the
42 histidine kinases NtrB and NtrY can substitute for each other as phosphodonors towards
43 the response regulator NtrC [29]. Also, a cross-talk between the NtrB/C and NtrY/X
44 sensor/regulator pairs has been suggested in *A. brasilense* during nitrate-dependent
45 growth [26]. However, NtrY protein is much larger than NtrB suggesting that they may
46 play different functions. The NtrBC system is widespread among microorganism and
47 displays known functions, like the control of genes for glutamine synthetase, amino
48 acids transport, nitrate and nitrite assimilation, nitrogen fixation and, also for the
49 expression of other nitrogen regulation genes. Although the role of the NtrBC system
50 seems to be more related to nitrogen metabolism, in *A. brasilense* there is also a link

1 between this two-component regulatory system and polyhydroxybutyrate production
2 [44]. To summarize, Fig. 6 represents the integration of mutational, transcriptomic and
3 proteomic data to establish a novel regulation model of the nitrate/nitrite assimilation
4 process in *P. denitrificans*, in which nitrate exerts a transcriptional and translational
5 control that has no precedents in the literature as far as we know. In this model three
6 conditions have been considered: i) presence of ammonium, in which the *nasABGHC*
7 genes are repressed and there is a very low expression of the *nasTS* genes; ii) absence of
8 both ammonium and nitrate (i.e. glutamate as N-source), in which *nasABGHC* and
9 *nasTS* genes are expressed, but Nas polypeptides and nitrate reductase activity
10 are not detected; and iii) absence of ammonium and presence of nitrate, in which
11 *nasABGHC* and *nasTS* genes are expressed and Nas proteins are synthesized, leading to
12 an active nitrate assimilation system. Our experimental data supports the existence of a
13 regulatory cascade with three levels of control. At level 1, the phosphorylated NtrC
14 protein (and probably NtrX in a lesser extent) interacts with the *nas* promoter to activate
15 *nas* gene expression in absence of ammonium, as described in other bacteria (general
16 nitrogen control). Accordingly, a conserved NtrC-binding sequence has been found in
17 the promoter region of the *nasT* gene. At level 2, the NasTS system controls expression
18 of the *nasABGHC* genes by transcription antitermination. In the absence of nitrate, an
19 inactive NasTS complex is formed but there is enough free NasT protein to allow
20 transcription antitermination for the synthesis of a complete *nasABGHC* transcript,
21 which is detectable in cells grown with glutamate. This NasT/NasS unbalance is
22 probably a consequence of the high level of expression of the *nasT* gene compared to
23 the *nasS* gene. At level 3, translation of the Nas proteins only occurs in the presence of
24 nitrate. This regulation at the translational level may occur when NasS binds to nitrate
25 and the NasTS complex dissociates, leading to an increase of free NasT protein that, in
26 turn, may lead to up-regulation of systems involved in post-translational regulation.

27 This is a novel regulatory mechanism, but bioinformatics analyses may suggest
28 that NasTS regulation is emerging in a wide number of bacteria, as previously reviewed
29 [8]. As the *P. denitrificans* Nas system is very well characterized [2,5], this work may
30 now lead to comparative studies in other bacteria, in which the NasTS system regulated
31 nitrate assimilation and even other processes like nitrous oxide reduction, as recently
32 described in *Bradyrhizobium japonicum* [41].

ACKNOWLEDGEMENTS

This work was supported by Ministerio de Ciencia y Tecnología (Grant BIO2015-64311-R) and Junta de Andalucía (Grant CVI-7560), Spain. DJR was a Royal Society and Wolfson Foundation for Merit Award Fellow and thank the Biotechnology and Biological Sciences Research Council (Grants BB/EO21999/1 and BB/D523019/1). This work was also funded by the Biotechnology and Biological Sciences Research Council (Grant BB/M00256X/1). VML-A was recipient of a postdoctoral fellowship from the Junta de Andalucía, Spain.

ABBREVIATIONS

ANTAR, Ami and NasR transcription antitermination regulator; Cy5-dCTP and Cy3-dCTP, deoxycytidine triphosphate containing either Cy5 or Cy3 dyes; dNTPs, deoxyribose-nucleoside triphosphates; EDTA, ethylenediaminetetraacetic acid; MALDI-TOF/TOF, matrix-assisted laser desorption/ionization (time of flight); NADH, nicotinamide adenine dinucleotide; NIT, N-terminal nitrate/nitrite sensor domain; Ntr, nitrogen regulator; RT-PCR, reverse transcription polymerase chain reaction; RT-qPCR, quantitative reverse transcription polymerase chain reaction; 2D-PAGE, two-dimensional polyacrylamide gel electrophoresis.

Table 1. Analysis by RT-qPCR of nitrate-induced genes detected in the microarrays of *P. denitrificans*.

Gene	$\text{NO}_3^-/\text{NH}_4^+$
Pden_0490 <i>gltD</i>	2.8 ± 0.5
Pden_0794 <i>bztB</i>	1.9 ± 0.4
Pden_1212 <i>ureD</i>	5.1 ± 1.8
Pden_2032 <i>amtB</i>	5.3 ± 1.7
Pden_2151 <i>potD</i>	2.6 ± 0.2
Pden_4019 <i>urtC</i>	6.2 ± 0.4
Pden_4129 <i>ntrC</i>	4.6 ± 0.5
Pden_4130 <i>ntrB</i>	1.6 ± 0.2
Pden_4234 <i>narJ</i>	24.3 ± 11.1
Pden_4236 <i>narG</i>	100.5 ± 23.3
Pden_4450 <i>nasH</i>	185.7 ± 9.8
Pden_4453 <i>nasA</i>	385.0 ± 11.9
Pden_4454 <i>nasS</i>	7.7 ± 1.5
Pden_4455 <i>nasT</i>	23.3 ± 6.3
Pden_4461 <i>glnB</i>	2.6 ± 0.6
Pden_4462 <i>glnA</i>	3.4 ± 0.7

Data correspond to the mean ± standard deviation of the nitrate/ammonium gene expression ratio.

Table 2. Analysis by RT-qPCR of nitrate-induced genes in the wild-type, *nasT* and *nasS* mutant strains of *P. denitrificans*.

Gene	Wild-type Glu+NO₃⁻/Glu	<i>nasT</i> mutant Glu+NO₃⁻/Glu	<i>nasS</i> mutant Glu+NO₃⁻/Glu
Pden_0490 <i>gltD</i>	0.6 ± 0.1	0.7 ± 0.1	2.6 ± 0.2
Pden_0794 <i>bztB</i>	2.2 ± 1.6	0.7 ± 0.4	-
Pden_1212 <i>ureD</i>	1.1 ± 0.4	1.0 ± 0.1	1.8 ± 0.1
Pden_2032 <i>amtB</i>	0.4 ± 0.0	0.6 ± 0.1	0.3 ± 0.1
Pden_2151 <i>potD</i>	0.5 ± 0.1	0.6 ± 0.1	-
Pden_4019 <i>urtC</i>	0.5 ± 0.1	0.5 ± 0.1	-
Pden_4129 <i>ntrC</i>	0.4 ± 0.1	0.5 ± 0.1	0.8 ± 0.2
Pden_4130 <i>ntrB</i>	0.7 ± 0.1	0.6 ± 0.2	0.6 ± 0.1
Pden_4234 <i>narJ</i>	7.7 ± 1.2	57.0 ± 14.4	-
Pden_4236 <i>narG</i>	2.5 ± 0.2	67.5 ± 12.2	173.8 ± 12.3
Pden_4450 <i>nasH</i>	0.6 ± 0.2	0.9 ± 0.1	0.2 ± 0.1
Pden_4453 <i>nasA</i>	0.9 ± 0.3	0.3 ± 0.1	0.2 ± 0.1
Pden_4454 <i>nasS</i>	1.0 ± 0.3	1.4 ± 0.1	0.0 ± 0.0
Pden_4455 <i>nasT</i>	2.0 ± 0.4	0.0 ± 0.0	1.8 ± 0.1
Pden_4461 <i>glnB</i>	2.0 ± 0.2	0.4 ± 0.1	1.0 ± 0.1
Pden_4462 <i>glnA</i>	1.7 ± 0.1	0.4 ± 0.1	1.2 ± 0.1
Pden_3872 <i>gdhA</i>	2.0 ± 0.3	1.1 ± 0.2	11.4 ± 1.0

Data correspond to the mean ± standard deviation of the glutamate plus nitrate/glutamate gene expression ratio.

Table 3. Analysis of the *nasT* promoter by *lacZ* (β -galactosidase activity) gene fusion in *P. denitrificans* wild-type and *ntrB* and *ntrY* mutant strains.

Strain/construct	N-source			
	NH ₄ ⁺	NO ₃ ⁻	Glu	Glu + NO ₃ ⁻
WT/PnasT (native)	71 ± 5	14500 ± 143	11000 ± 120	15000 ± 152
WT/PnasT (mutated)*	50 ± 3	220 ± 14	53 ± 5	600 ± 25
<i>ntrY</i> / PnasT (native)	n.d.	n.d.	11500 ± 115	15200 ± 135
<i>ntrB</i> / PnasT (native)	n.d.	n.d.	10 ± 1	120 ± 4

* The base substitutions are specified in Table S2; n.d.: not determined.

REFERENCES

1. Richardson, D. J., Berks, B. C., Rusell, D. A., Spiro, S., and Taylor, C. J. (2001) Functional, biochemical and genetic diversity of prokaryotic nitrate reductases. *Cell. Mol. Life Sci.* **58**, 165-178
2. Gates, A. J., Luque-Almagro, V. M., Goddard, A. D., Ferguson, S. J., Roldán, M. D., and Richardson, D. J. (2011) A composite biochemical system for bacterial nitrate and nitrite assimilation as exemplified by *Paracoccus denitrificans*. *Biochem. J.* **435**, 743-753
3. Goddard, A. D., Bali, S., Mavridou, D. A., Luque-Almagro, V. M., Gates, A. J., Roldán, M. D., Newstead, S., Richardson, D. J., and Ferguson, S. J. (2017). The *Paracoccus denitrificans* NarK-like nitrate and nitrite transporters-probing nitrate uptake and nitrate/nitrite exchange mechanisms. *Mol. Microbiol.* **103**, 117-133
4. Shu, C. J., and Zhulin, I. B. (2002) ANTAR: an RNA-binding domain in transcription antitermination regulatory proteins. *Trends Biochem. Sci.* **27**, 3-5
5. Luque-Almagro, V. M., Lyall, V. J., Ferguson, S. J., Roldán, M. D., Richardson, D. J., and Gates, A. (2013) Nitrogen oxyanion-dependent dissociation of a two-component complex that regulates bacterial nitrate assimilation. *J. Biol. Chem.* **288**, 29692-29702
6. Waller, Z. A. E., Pinchbeck, B. J., Buguth, B. S., Meadows, T. J., Richardson, D. J., and Gates, A. J. (2016). Control of bacterial nitrate assimilation by stabilization of G-quadruplex DNA. *Chem. Commun.* **52**, 13511-13514
7. Moreno-Vivián, C., and Flores, E. (2007) Nitrate assimilation in Bacteria. In *Biology of the Nitrogen Cycle*. Bothe, H., S. J. Ferguson, W. E. and Newton (eds). Amsterdam: Elsevier, pp.263-282
8. Luque-Almagro, V. M., Gates, A. J., Moreno-Vivián, C., Ferguson, S. J., Richardson, D. J., and Roldán, M. D. (2011) Bacterial nitrate assimilation: gene distribution and regulation. *Biochem. Soc. Trans.* **39**, 1838-1843
9. Paz-Yepes J., Flores, E., and Herrero, A. (2003) Transcriptional effects of the signal transduction protein P_{II} (*glnB* gene product) on NtcA-dependent genes in *Synechococcus* sp. PCC 7942. *FEBS Lett.* **543**, 42-46
10. Forchhammer, K. (2004) Global carbon/nitrogen control by pII signal transduction in cyanobacteria: from signals to targets. *FEMS Microbiol. Rev.* **28**, 319-333
11. Flores, E., Frias, J. E., Rubio L. M., and Herrero, A. (2005) Photosynthetic nitrate assimilation in cyanobacteria. *Photosynth. Res.* **83**, 117-133
12. Kayumov, A., Heinrich, A., Sharipova, M., Lljinskaya, O., and Forchhammer, K. (2008) Inactivation of general transcription factor TnrA in *Bacillus subtilis* by proteolysis. *Microbiology* **154**, 2348-2355
13. Tiffert, Y., Supra, P., Wurm, R., Wohlleben, W., Wagner, R., and Reuther, J. (2008) The *Streptomyces coelicolor* GlnR regulon: identification of new GlnR targets and evidence for a central role of GlnR in nitrogen metabolism in actinomycetes. *Mol. Microbiol.* **67**, 861-880
14. Lin, J. T., and Stewart, V. (1996) Nitrate and nitrite-mediated transcription antitermination control of *nasF* (nitrate assimilation) operon expression in *Klebsiella pneumoniae* M5a1. *J. Mol. Biol.* **256**, 423-435
15. Chai, W., and Stewart, V. (1999) RNA sequence requirements for NasR-mediated, nitrate-responsive transcription antitermination of the *Klebsiella oxytoca* M5a1 *nasF* operon leader. *J. Mol. Biol.* **292**, 203-216

16. Boudes, M., Lazar, N., Graille, M., Durand, D., Gaidenko, T. A., Steward, V., and Tilbeurgh, H. (2012) The structure of the NasR transcription antiterminator reveals a one-component system with a NIT nitrate receptor coupled to an ANTAR RNA-binding effector. *Mol. Microbiol.* **85**, 431-444
17. Gutiérrez, J. C., Ramos, F., Ortner, L., and Tortolero, M. (1995) *nasST*, two genes involved in the induction of the assimilatory nitrate-nitrite reductase operon (*nasAB*) of *Azotobacter vinelandii*. *Mol. Microbiol.* **18**, 579-591
18. Caballero, A., Esteve-Núñez, A., Zylstra, G. J., and Ramos, J. L. (2005) Assimilation of nitrogen from nitrite and trinitrotoluene in *Pseudomonas putida* JLR11. *J. Bacteriol.* **187**, 396-399
19. Toukdarian, A., and Kennedy, C. (1986) Regulation of nitrogen metabolism in *Azotobacter vinelandii*: isolation of *ntr* and *glnA* genes and construction of *ntr* mutants. *EMBO J.* **5**, 399-407
20. Wang, B., Pierson III, L. S., Rensing, C., Gunatilaka, M. K., and Kennedy, C. (2012) NasT-mediated antitermination plays an essential role in the regulation of the assimilatory nitrate reductase operon in *Azotobacter vinelandii*. *Appl. Environ. Microbiol.* **78**, 6558-6567
21. Swanson, R. V., Alex, L. A., and Simon, M. I. (1994) Histidine and aspartate phosphorylation: two-component systems and the limits of homology. *Trends Biochem. Sci.* **19**, 485-90
22. Sanders, D. A., Gillice-Castro, B. L., Burlingame, A. L., and Koshland, D. E., Jr. (1992) Phosphorylation site of NtrC, a protein phosphatase whose covalent intermediate activates transcription. *J. Bacteriol.* **174**, 5117-5122
23. Chen, P., and Reitzer, L. J. (1996) Active contribution of two domains to cooperative DNA binding of the enhancer-binding protein regulator I (NtrC) of *Escherichia coli*: stimulation by phosphorylation and the binding of ATP. *J. Bacteriol.* **177**, 2490-2496
24. Cullen, P. J., Bowman, W. C., and Kranz, R. G. (1996) In vitro reconstitution and characterization of the *Rhodobacter capsulatus* NtrB and NtrC two-component system. *J. Biol. Chem.* **271**, 6530-6536
25. Pawlowsky, K., Klosse, U., and the Bruijn, F. J. (1991). Characterization of a novel *Azorhizobium caulinodans* ORS571 two-component regulatory system, NtrY/NtrX, involved in nitrogen fixation metabolism. *Mol. Gen. Genet.* **231**, 124-138
26. Ishida, M. L., Assumpção, M. C., Machado, H. B., Benelli, E. M., de Souza, E. M., and Pedrosa, F. O. (2002) Identification and characterization of the two-component NtrY/NtrX regulatory system in *Azospirillum brasilense*. *Braz. J. Med. Biol. Res.* **35**, 651-661
27. Assumpção, M. C., de Souza, E. M., Yates, M. G., de Oliveira-Pedrosa, F., and Machado-Benelli, E. (2007) Purification and characterization of *Azospirillum brasilense* N-truncated NtrX protein. *Prot. Express. Purif.* **53**, 302-308
28. Bonato, P., Alves, L. R., Osaki, J. H., Rigo, L. U., Pedrosa, F. O., de Souza, E. M., Zhang, N., Schumacher, J., Buck, M., Wasseem, R., and Chubatsu1, L. S. (2016). The NtrY/NtrX two-component system is involved in controlling nitrate assimilation in *Herbaspirillum seropedicae* strain SmR1. *FEBS J.* **283**, 3919-3930
29. Drepper, T., Wiethaus, J., Giaourakis, D., Groß, S., Schubert, B., Vogt, M., Wiencek, Y., McEwan, A. G., and Masepohl, B. (2006) Cross-talk towards response regulator NtrC controlling nitrogen metabolism in *Rhodobacter capsulatus*. *FEMS Microbiol. Lett.* **258**, 210-256

30. Carrica, M. C., Fernandez, I., Martí, M. A., Paris, G., and Goldbaum, F. A. (2012) The NtrY/X two-component system of *Brucella* spp. acts as a redox sensor and regulates the expression of nitrogen respiration enzymes. *Mol. Microbiol.* **85**, 39-50
31. Roop II, R. M., and Caswell, C. C. (2012) Redox-responsive regulation of denitrification genes in *Brucella*. *Mol. Microbiol.* **85**, 5-7
32. Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989) Molecular cloning. In A laboratory manual, 2nd edn. New York: Cold Spring Harbor Laboratory, Cold Spring Harbor
33. Robertson, L. A., and Kuenen, J. G. (1983) *Thiosphaera pantotropha* gen. nov. sp. nov., a facultatively anaerobic, facultatively autotrophic sulphur bacterium. *J. Gen. Microbiol.* **129**, 2847-2855
34. Shakir, F. K., Audilet, D., Drake, A. J. III, and Shakir, K. M. M. (1994) A rapid protein determination by modification of the Lowry procedure. *Anal. Biochem.* **216**, 232-233
35. Snell, F. D., and Snell, C. T. (1949) Nitrites. In *Colorimetric methods of analysis*. Snell, F. D., and C. T. Snell (eds). New York: van Nostrand, pp.802-807
36. Miller, J. M. (1972) In *Experiments in molecular genetics*. Miller, J. (ed). New York: Cold Spring Harbor Laboratory Press, pp.171-224
37. Alston, M. J., Seers, J., Hinton, J. C., and Lucchini, S. (2010) BABAR: an R package to simplify the normalisation of common reference design microarray-based transcriptomic datasets. *BMC Bioinformatics* **11**, 73
38. Untergasser, A., Nijveen, H., Rao, X., Bisseling, T., Geurts, R., and Leunissen, J. A. (2007) Primer3Plus, an enhanced web interface to Primer3. *Nucleic Acids Res* **35**, W71-74
39. Pfaffl, M. W. (2001) A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res.* **29**, E45
40. Luque-Almagro, V. M., Huertas, M. J., Roldán M. D., Moreno-Vivián, C., Martínez-Luque, M., Blasco, R., and Castillo, F. (2007) The cyanotrophic bacterium *Pseudomonas pseudoalcaligenes* CECT5344 responds to cyanide by defence mechanism against iron deprivation, oxidative damage and nitrogen stress. *Environ. Microbiol.* **9**, 1541-1549
41. Sánchez, C., Itakura, M., Okubo, T., Matsumoto, T., Yoshikawa, H., Gotoh, A., Hidaka, M., Uchida, T., and Minamisawa, K. (2014) The nitrate-sensing NasST system regulates nitrous oxide reductase and periplasmic nitrate reductase in *Bradyrhizobium japonicum*. *Environ. Microbiol.* **16**, 3263-3274
42. Pino, C., Olmo-Mira, M. F., Cabello, P., Martínez-Luque, M., Castillo, F., Roldán, M. D., and Moreno-Vivián, C. (2006) The assimilatory nitrate reduction system of the phototrophic bacterium *Rhodobacter capsulatus* E1F1. *Biochem. Soc. Trans.* **34**, 127-129
43. Wang, B., Rensing, C., Pierson III, L. S., Zhao, H., and Kennedy, C. (2014) Translational coupling of *nasST* expression in *Azotobacter vinelandii* prevents overexpression of the *nasT* gene. *FEMS Microbiol. Lett.* **361**, 123-130
44. Sun, J., Peng, X., van Impe, J., and Vanderleyden J. (2000) The *ntrB* and *ntrC* genes are involved in the regulation of poly-3-hydroxybutyrate biosynthesis by ammonia in *Azospirillum brasilense* Sp7. *Appl. Environ. Microbiol.* **66**, 11-117

FIGURE LEGENDS

Figure 1. Transcriptomic and proteomic analysis of aerobically cells grown with nitrate or ammonium as nitrogen source. (A) Heat map from DNA microarrays analyses of *P. denitrificans* PD1222 grown aerobically with either ammonium or nitrate as the sole nitrogen source. Colours indicate \log_2 of normalised expression values in either condition. Genes were selected based on at least 2-fold change in expression with p -values ≤ 0.05 . A total of 74 genes were up-regulated more than 2-fold under nitrate assimilation conditions, whereas nine genes were down-regulated more than 2-fold under these conditions. (B) 2D-PAGE analysis of *P. denitrificans* PD1222 soluble fractions from cells grown aerobically with ammonium or nitrate (10 mM each) as the sole nitrogen source. Spots present or absent in gels are highlighted by black or white triangles, respectively. The presented gels are representative of three independent replicants.

Figure 2. Analysis by RT-PCR and qPCR of the *P. denitrificans* nitrate assimilation *nas* genes. (A) The *nas* gene cluster of *P. denitrificans* and the location of the oligonucleotides used in this study are shown. (B) RT-PCR reactions with RNA from the wild-type strain of *P. denitrificans* grown in the presence of different N-sources. To isolate RNA, cells were cultured in minimal medium with different N-sources as described in Materials and Methods section and harvested at an A_{600} of about 0.4. The *rpoB* gene was used as housekeeping. -/+, without/with reverse transcriptase (C) RT-qPCR analysis of the *nasT*, *nasS* and *nasA* genes from cells grown with different nitrogen sources. Error bars represent standard deviation calculated from the results of three independent experiments.

Figure 3. Characterization of the wild-type strain and the *nasT* and *nasS* mutants of *P. denitrificans* with different nitrogen sources. (A) NADH-dependent nitrate reductase activity and NasG polypeptide detection by 2D-PAGE analysis in the wild-type strain of *P. denitrificans*. The 2D-PAGE analyses were carried out in cytoplasmic fraction from cells grown with different nitrogen sources obtained by subcellular fractionations as indicated in Materials and Methods. The different nitrogen sources were nitrate, ammonium, ammonium plus nitrate, glutamate or glutamate plus nitrate (10 mM each). Isoelectric focusing was performed from IPG strips range 4-7 and second dimension was carried out onto 12% polyacrylamide gels. The nitrate reductase activity (NR) was assayed in cytoplasmic fractions and was undetectable with glutamate or ammonium as N-source, and expressed as $\text{nmol NO}_2^- \text{ formed min}^{-1} \text{ mg}^{-1}$ (n.d, not detected). (B) RT-PCR analysis of structural *nas* genes in the wild-type and *nasT* and *nasS* mutant strains of *P. denitrificans*. To isolate RNA, wild-type and *nasT* and *nasS* strains were cultured in minimal medium with different N-sources as described in Materials and Methods section and harvested at an A_{600} of about 0.4. The *rpoB* gene was used as housekeeping. -/+, without/with reverse transcriptase. (C) NasG polypeptide detection by 2D-PAGE analysis and NADH-nitrate reductase activity in cytoplasmic fractions from *nasT* and *nasS* mutant strains of *P. denitrificans*. The 2D-PAGE analysis and the NR activity assays were performed as indicated in (A) for the wild-type strain.


Figure 4. Analysis of the *nasA* leader region of *P. denitrificans*. (A) Intergenic *nasS-nasA* region from *P. denitrificans*. (B) Expression of the transcriptional $P_{nasA-lacZ}$ fusions and effect of mutations within the leader sequence on *nasA-lacZ* expression in cells grown with ammonium or nitrate (10 mM each). The β -galactosidase activity was

measured when the A_{600} was about 1.0 and it was represented as Miller units (M.U.). Control represents the wild type (native) construction from -321 to +71 nucleotides of the *nasS-nasA* intergenic region. HI and HII represent the $P_{nasA}::lacZ$ constructions with 3 bp mutations in each hairpin. (C) Growth curves with nitrate as sole nitrogen source of the *P. denitrificans* wild-type strain and the mutants in the *nasT* gene and the *cis*-acting NasT binding region (hairpin I, HI).

Figure 5. Characterization of the *ntrB*, *ntrY* and double *ntrB/ntrY* mutant strains of *P. denitrificans* in media with nitrate. (A) Growth-curves of *P. denitrificans* wild-type strain (●), and *ntrB* (■), *ntrY* (▲) and double *ntrB/ntrY* (◆) mutants with 10 mM nitrate as nitrogen source. (B) RT-qPCR analysis of the regulatory genes *nasT*, *nasS*, *ntrB* and *ntrY* and the structural *nasA* gene in the wild-type strains and the *ntrB* mutant strains of *P. denitrificans* grown with 10 mM nitrate as N-source. The oligonucleotides used in the qPCR analysis for the *ntrB* and *ntrY* genes hybridized upstream the mutation site. Represented data were normalized by using the *rpoB* and *dnaN* genes as housekeeping (Table S2). Error bars represent standard deviation calculated from the results of three independent experiments.

Figure 6. Proposed model of the nitrate assimilation regulation in *P. denitrificans*. The model describes the regulatory cascade controlling expression of nitrate assimilation *nas* genes by the *ntrBCYX* gene cluster in response to ammonium availability and by the *nasTS* genes for nitrate control. Levels 1 and 2 include regulatory mechanisms that control transcription of the *nas* genes, whereas level 3 involves a control on the translation of the Nas proteins. A circle with a plus symbol means activation of the transcription and a circle with the capital letter A indicates transcription antitermination.

A



B




Figure 1




Figure 2






Figure 3








Figure 4

A**B****Figure 5**




Figure 6

Table S1. Bacterial strains and plasmids used in this study.

Strain or plasmid	Genotype and description	Source or reference
<i>Paracoccus denitrificans</i>		
PD1222		
Wild-type	Sp ^R , Rif ^R , enhanced conjugation frequencies	[1]
<i>nasT</i>	<i>nasT</i> Δ::Sm ^R	[2]
<i>nasS</i>	<i>nasS</i> Δ::Km ^R	[2]
<i>ntrB</i>	<i>ntrB</i> Δ::Sm ^R	This work
<i>ntrY</i>	<i>ntrY</i> Δ::Km ^R	This work
<i>ntrB/ntrY</i>	<i>ntrB</i> Δ::Sm ^R / <i>ntrY</i> Δ::Km ^R	This work
HI (hairpin I)	Unmarked mutant	This work
<i>Escherichia coli</i>		
DH5α	<i>deoR endA1 gyrA96 hsdR17</i> (r _k ⁻ m _k ⁺) <i>recA1 relA1 supE44 thi-1</i> Δ(<i>lacZYA-argFV169</i>) Ø80 δ <i>lacZ</i> ΔM15 F ⁻ λ ⁻	[3]
pRK2013	Km ^R , Tra ⁺ (pRK2013 encoded) helper strain	[4]
Plasmids		
pUC18	Amp ^R , cloning vector	[5]
pGEM-Teasy	Amp ^R , cloning vector	Promega (USA)
pGEM-T	Amp ^R , cloning vector	Promega (USA)
pSparkII	Amp ^R , cloning vector	Canvax (Spain)
pSUP202	Amp ^R , Tet ^R , Cm ^R , Mob ⁻ , mobilizable suicide vector	[6]
pSUP202*	pSUP202::Km-ΔTet, mobilizable suicide vector	[7]
pSUP2021	pSUP202::Tn5, source of Km ^R cassette	[6]
pK18 <i>mobsacB</i>	Gene knockout vector, Km ^R	[8]
pSRA2	Amp ^R , Sp ^R , Sm ^R , source of Sm ^R cassette	[9]
pMP220	Promoter probe vector, <i>lacZ</i> gene, IncP1 Tc ^R	[10]
pMP220- <i>PnasA</i>	pMP220-based transcriptional fusion spanning the intergenic region between <i>nasS</i> and <i>nasA</i> (Tet ^R)	This work
pMP220- <i>PnasA1</i>	pMP220-based transcriptional fusion spanning the intergenic region between <i>nasS</i> and <i>nasA</i> containing the hairpin I mutated (Tet ^R)	This work
pMP220- <i>PnasA2</i>	pMP220-based transcriptional fusion spanning the intergenic region between <i>nasS</i> and <i>nasA</i> containing the hairpin II mutated (Tet ^R)	This work
pMP220- <i>PnasT</i>	pMP220-based transcriptional fusion spanning the <i>nasT</i> leader region (Tet ^R)	This work
pMP220- <i>PnasT1</i>	pMP220-based transcriptional fusion spanning the <i>nasT</i> leader region and containing the	This work

	NtrC-binding site mutated (Tet ^R)	
--	---	--

Table S2. Oligonucleotide primers used in this study.

Primer	Sequence (5' → 3') ^a	Used for
ntrB1-HB (<i>HindIII</i> - <i>Bam</i> HI)	CCCAAGCTTACGAATTCTCGCCAATGGCGACGTGGTGGA TGCGGCCTCGC	<i>ntrB</i> mutagenesis
ntrB2-B (<i>Bam</i> HI)	CGGGATCCAGGATCGCGCCGAGCCGGGGCACGATG	<i>ntrB</i> mutagenesis
ntrB3-B (<i>Bam</i> HI)	CCCTGCGGATCCAGATCGAGGACGACGGCCC	<i>ntrB</i> mutagenesis
ntrB4-K (<i>Kpn</i> I)	ACGGGGTACCCGAAGGGCTTGGGCAGGTAGTCGAAGGC	<i>ntrB</i> mutagenesis
ntrY1-A (<i>Aat</i> II)	GATGACGTCGCGAGAGTCGCTGAACGGCCACAGAA	<i>ntrY</i> mutagenesis
ntrY2-B (<i>Bam</i> HI)	CGCGGATCCCGAGGTGCAAACACCTGCTGCACCC	<i>ntrY</i> mutagenesis
ntrY3-B (<i>Bam</i> HI)	GCCGGATCCCTGGAGGGCTATGTGGTGGCGCTGGA	<i>ntrY</i> mutagenesis
ntrY4-E (<i>Eco</i> RI)	GCCGAATTCCTCGGGGGCATCGGTCAGCATCAGGC	<i>ntrY</i> mutagenesis
PTF	CGCCACGCCGCATAATGGTCGAAGGGATAGT	<i>nasT</i> :: <i>lacZ</i> fusion
PTR	GCGAGGGATTGGCCAGGTCGATCAGCA	<i>nasT</i> :: <i>lacZ</i> fusion
PTNtrR	GCGGCGCTGCCCATGGTTCGGTAGGCG	<i>nasT</i> :: <i>lacZ</i> fusion
PTNtrR_Mut	GCGGCGCT AT (GC) CC (A)TGGTTCGGTAGGCG	Mutated <i>nasT</i> promoter:: <i>lacZ</i> fusion
PTNtrF	CGCCTACCGAACCATGGGCAGCGCCGC	<i>nasT</i> :: <i>lacZ</i> fusion
PTNtrF_Mut	CGCCTACCGAACC AG (T)GG AT (CG)AGCGCCGC	Mutated and native <i>nasT</i> promoter:: <i>lacZ</i> fusion
FA1_A	CGACGGCACCATCTTCGA	<i>nasA</i> :: <i>lacZ</i> fusion
RA1	GTAAGGCGAAGGTCGAGAG	Mutated <i>nasA</i> promoter:: <i>lacZ</i> fusion
RA1-S (<i>Sph</i> I)	TAGCATGCGTAAAGGCGAAGGTCGAGAG	Mutated <i>nasA</i> promoter :: <i>lacZ</i> fusion
FA1_B-P (<i>Pst</i> I)	ATCTGCAGACGGCACCATCTTCGAACCACC	Mutated <i>nasA</i> promoter (hairpin I):: <i>lacZ</i> fusion
RHI-A (<i>Asc</i> I)	GCCGGTGGCGCGCCGTT CG (GC) GG (C)CGGGCCAT	Mutated <i>nasA</i> promoter (haipin I):: <i>lacZ</i> fusion

FHII-A (<i>AscI</i>)	AACGGCGCGCCACCGGCCGCTCGCAATGATGCGGCAC GGATA(T)CGG(CC)GCGA	Mutated <i>nasA</i> promoter (hairpin II):: <i>lacZ</i> fusion
AIF1 (<i>EcoRI</i>)	AAGAATTCGCGAATATCTGGACCTGCCGGCGGAA	Unmarked hairpin I mutant
AIR1a	ATGGGATCAGACTGGGTCAGGACGCA	Unmarked hairpin I mutant
AIF2 (<i>SalI</i>)	AAGTCGACCGCTCGCAATGATGCGGCACGGATTC	Unmarked hairpin I mutant
AIR2 (<i>HindIII</i>)	TTAAGCTTCATTGCCGGTGCCGAAGATGCCCAAC	Unmarked hairpin I mutant
nasTR	GATGCGGCCATCGCACGCTTTCACA	RT-PCR
nasSF	GCGGCGTCGATCAGCGGCACATAGC	RT-PCR
nasSR	GCGATGGCGCATTTCCGCACCGATCTTACC	RT-PCR
nasAF	ATGCGTTGCTGATGGCGAAGGCTGTCCGAAA	RT-PCR
nasAR	CTGGGCAAGGCGGCCGTCTACAAGCACAT	RT-PCR
nasBF	CCCGACAGGACCGGCGACAGCATCAG	RT-PCR
nasBR	TGGTCTGCGTCGACAGCGGCTACCAGATCAG	RT-PCR
nasGF	GCGGTGCCCCGAGTTCATGTGCAACACCCAGTT	RT-PCR
nasGR	CTGCCCCGCTGCACAACCTGGGTGTTGACATGA	RT-PCR
nasHF	GCGAAATTGCCGAAAAGACCAGGCCCCAGTT	RT-PCR
nasHR	ATCCTGATCTTCTTCTACATGGGTTTCGAGCATT	RT-PCR
nasCF	CGTCGCCAGCACGCCGAGCCCACGCCGCAATA	RT-PCR
RT-RpoB5'	TTTGTGCGGAAAGACATCATCAACGAGCAGAC	RT-PCR (housekeeping)
RT-RpoB3'	TGGGTATCCGGCGCATCCAGGTCAAGG	RT-PCR (housekeeping)
RT-A5'	GATGTTCTGGACCGCTCGGATGAG	qPCR (<i>nasT</i> gene)
RT-A3'	GGAACATGTGAAAGCGTGCGATGGC	qPCR (<i>nasT</i> gene)
RT-K5'	GGACGAGCCCCGAGAACCGC	qPCR (<i>nasS</i> gene)
RT-K3'	CGGCGTGGAAGGCGATGAA	qPCR (<i>nasS</i> gene)
RT-C5'	GCGGTGATGTATGCGACCTTCATCG	qPCR (<i>nasA</i> gene)
RT-C3'	GCTGGGGATGTGCTTGTAGACGGC	qPCR (<i>nasA</i> gene)


RT-H5'	GCGACGTTTCATCCGACCCATTACG	qPCR (<i>rpoB</i> -housekeeping)
RT-H3'	TGCCCTCGACCACCTTGCGGTAGG	qPCR (<i>rpoB</i> -housekeeping)
dnaN-1F'	CATGTCGTGGGTCAGCATAC	qPCR (<i>dnaN</i> -housekeeping)
dnaN-1R'	CTCGCGACCATGCATATAGA	qPCR (<i>dnaN</i> -housekeeping)
ntrB10	GCCATCATGGGCGAGGTTTTCGAGC	qPCR
ntrB11	CGGGGCGGGATTCCAGTGCGAT	qPCR
ntrY4	CGGTGAATATCGGGCTGGAGGGCTGG	qPCR
ntrY5	TGGCGTCGTTGGTCAAGTCTCGGCG	qPCR
nasT2F	TCGATCGTCGTCATCGAAT	qPCR to verify microarrays
nasT2R	GGTTTCTTCCGAGATGATGC	qPCR to verify microarrays
nasS2F	CTGGACCTGGTGATGTTCT	qPCR to verify microarrays
nasS2R	ATCGCGAAAGTCGAAATCAT	qPCR to verify microarrays
nasA1F	CCTGACGCAGGTCTATGGTT	qPCR to verify microarrays
nasA1R	GACGATGAAGGTCGCATACA	qPCR to verify microarrays
nasB2F	TCTGGACCGGCCCTATAAAT	qPCR to verify microarrays
nasB2R	CGATAGACCGACTGGCTGAT	qPCR to verify microarrays
nasG1F	GTCTTTGCGCTGGACGAC	qPCR to verify microarrays
nasG1R	ATGTCGAACACCCAGTTGTG	qPCR to verify microarrays
nasH1F	TCATGTGCAACTGGATGGTT	qPCR to verify microarrays
nasH1R	CATGTTACGATGGAATGCT	qPCR to verify microarrays
nasC1F	CTATGTCGCCAACAAAGCTGA	qPCR to verify microarrays

nasC1R	CCAGCTCCAGATCCTCGTAG	qPCR to verify microarrays
ntrB10	GCCATCATGGGCGACGGTTTTCGAGC	qPCR to verify microarrays
ntrB11	CGGGGCGGGATTCCAGTGCGAT	qPCR to verify microarrays
ntrC1F	ACCGCTATTCGACCTGCAT	qPCR to verify microarrays
ntrC1R	GGTTTATGCCCAGAAGATCG	qPCR to verify microarrays
GIDH1F	ACGGCAACACCTATTCACC	qPCR to verify microarrays
GIDH1R	ATCCTGCACCTTCTCGATG	qPCR to verify microarrays

^aEnzyme restriction sites are underlined. Bases changed are marked in bold and italics

and native bases are in brackets.

A



B




Figure S1

A

NtrY Pd 1 -MAGTLSKGTWERVARLRRRLRYRN-----TATLGLALLGPALAGLTFVAMGPFADSTSGG-TVLRLLVLLADLLYLIIVLTGLVVARMAQ 82
NtrY Ab 1 -MSPTTPEVTVPLWQQFLRWAARVG---LAKRLAFALSLAALVAGPATYALTESAPGETNPRVTWLLTLDLALLLLLVGLIARRIVY 86
NtrY Ac 1 -MTQAAPDQASDNGPMTPSGSSFG----LFAP--AVVLLALISALATFLIILMGLTPVVVPTH-QVVISVLLVNAAAVLLSAMVGRRIWR 81
NtrY Ba 1 -MNRANLTTDMNGTRQSSQDEGQEGRRLLALPGIITVVSALVTASFSPAILIGITPTIPDR-TVTLALVINIVALILFLILLICREVIYR 88
NtrY Cc 1 -MGGGYALAVILITVT-----GVALASSPPRTDSIGAASTVILVVLGPNLVLILGVATIVGLRLYE 59
NtrY Sm 1 MVDGMALPLGTEDGVTAADR-----RASFALPGLMLATGALICATLSLLVLLGLTPIRPER-NIVACAGINGLFFVGLIYLITAREIFR 84
NtrY Ml 1 -----MLALPGVVAVVGALVMAAISFTIILVGATPIAPDA-RTTWALIALNAAFVFLMALVGREVHR 61

NtrY Pd 83 TVAAFRKSAAGSRILHFLVGVFATIALVETVVLVAFAGLTVNIGLEWFSNRVQVQVSTSLAAEAETODEHRRDLTNDAKLAGATQAA 172
NtrY Ab 87 LWIGRRRLGAGSQMVRLLVAVFSLLAVABAIIMAFISTVFFYVGVQSWSERVAVNESLAVASAMLHEHQNRADALAMANDLQEA 176
NtrY Ac 82 IAKARARGRAARHFRIVGLFAVVSVEAILVAVVAVSLTLDRGLDRWBSMRTQEIIVASSVVAQTVREHALNIRGDIILAMSADLTRLK 171
NtrY Ba 89 IITARRLLGKAASRIHFRIVALFSLIAAIEAIVVAVIVASVTLNIGLDRWBDQNTDIISSSQNFYTAQLQETALNLQSTSYSLMQDDAQR 178
NtrY Cc 60 LIDAR-ASDAGARHLHFRVGLFSLAAVAFAVIVLFFGVLVNRGVDGWBSERVQTVVGNSAKVANSVVEQQKNYI SEHIGPMAASLNQAA 148
NtrY Sm 85 LLRARSKGRPAARHFRIVALFSLIATLFAILLVAVIFASITLVDGLDRWBSLRTQAVIRSSLNVAQAQVLENASYLQGGQTVSMANDLERNR 174
NtrY Ml 62 TVMARRHGKAASRIHFRIVAMPALVAALIAIMVAVI IASTITLDTGLDRWBSLRTQAVIRSSLSIADAVQENARNLQGTTLMSAYDLDASR 151

NtrY Pd 173 RQNPLMDGDELRLLLGQQAALIQRLREBYVID-GRGTIRARGERSYQFWYEEPSVQFDEAATRGLVLIEDWQNEFRALVPLVPLADR 261
NtrY Ab 177 ARLASDPERFEQVWATQ---AMLRALSBAIVFNGTGAIIVARSGYTFALEFPDPI PDDKLATARRGEVAMIVSENDDRVRLVRLDRFADT 263
NtrY Ac 172 SVYEGDRSRFNILLTAQ---AALRNLPQAMLIIR-RDLSVVERANVNI GREFIVPANLAIGDADPDQVPVYLPNDADYAAVPLKDYDDL 257
NtrY Ba 179 AYSLDRNGFITQLMTLQ--VRGRALLCAFVLR-DNGDGVVQSDTGMESHLPPTKDSLEKAPDGKPVII PPGNTNFVGFALIKMREIPDM 264
NtrY Cc 149 PALAQSPVAFGHFLAGLTQ---DNGFSAPYVLD-RDGRILARTEAEGAPPFLAPPASFDWADGGEASAQRFVSTDLFRTLKLGKDFDG 234
NtrY Sm 175 QYSLDRDTGFVFLMTRQ---ARGRMGLCAFVLR-ADGSAALQANISTDRPLPAI PQDALKSTIAGQPTLI PPGVNTLVGAVIIPLENIPGT 260
NtrY Ml 152 SLYGLDRTGFLDLLNKE---AVGRSLAHAAALIK-PDGSFVMSAQTDADFAMPEPPEGVSVSATDGPVLI EPTRNIMGAVIKGREIEGL 237

NtrY Pd 262 VLYVTRDVGDSLLGLDDTRQTVGDYQRLEQERGRVLPFESLLYLGFAPLLVARFMLGLWFADRLSREI GRFAEASECYGECNIDFQIP 351
NtrY Ab 264 VLYVGRMVEPRVLSHMASAEGAVREFGALESQRGSLQITFTLIFLCPVALLLLAFAVWAGLIFATRLVRFISALIGADRVRACQIDTVRVT 353
NtrY Ac 258 VLYVARDLDPVIGYLYKTTQETLADYRSLEERRPFGVQVAGALMYAVITLIVLLSAVVLGGLNFSKWLVAPIRRMSADHVAECNLDVVRVP 347
NtrY Ba 265 VLYTVRSVPPQILDANLNMANARRYQEMDANRIPTOIAGALYFGLTILVLLSAIWTGIAVADRVRHRLIGASDDVAACNLDVSV 354
NtrY Cc 235 VLYVVRPEKGINHLIETQDALVSYRDAERSGRIOAIFGLSYLETALVVLVPAIHWGIAAANSIAGVAGIIGASAGRVSGCQIDARVE 324
NtrY Sm 261 VLYTVRNVDPVEMSRMLMEENTAEYKTLAEAGRTSLQIAGVLYIGFAPLIVLLAIAIWAIAVADRIVRPHRQIIGASAGRVVASCNLDVVRVP 350
NtrY Ml 238 VLYTIRLVDPPEVIKARQIVRSNTDEYRNLEDNRRTSQAVALPYLSLTIITILSAIWTGIAVADRVRHRLIGADEVATCNLDVAVP 327

NtrY Pd 352 APDTGDEIQTIGESFARNMTROLKCRQELIVESYRAADDORRLFDSVLSVTVAGVIGLDAAGEIDFLNRSATRLLGL-DPATAHD-RITSE 439
NtrY Ab 354 ERPAEDDLLALSRFNRMTTELESCHALLSANRLIDSRFRFTETVLSGVSAGRDGLDAEGRTLSKFSAAARLLGVKDAESLIG-MRLAE 442
NtrY Ac 348 IYRAGDDLASAETFNKMTHELRSCEALITARQIDISRRFRTEAVLSGVGAGVIGLDSQERTILNRSARLLGLSEVEALH-RHIAE 435
NtrY Ba 355 VRSSDGDVAGLSGTFNMMVAELKSCRNELISAKQIDIBERFSEAVLSGVTVAGVIGVIGLDSISILNRSAEHMFQSDDAVAG-KSATS 442
NtrY Cc 325 VERGPEEIRASNAFNMMTSELQLQAAKKAASLDAESRRQFIETVLSGVSAAGVVSLLDDRKGISAVNRRRAVALLGLPDDALGVD--HIA 411
NtrY Sm 351 VRAVDGVDGNRSRTFNKMWSEIRTOQDQLVAKDEVQRRRFTIETAVLSGVTAIVIGVGRDRRITIFNPSSEGLMKKAAPELIG-ANLSD 438
NtrY Ml 328 VRSSDGDVASIGDIFNKMILLEKSCRNELISAKDLIDERFSEAVLACVTVAGVIGVDPYGIIVTVNRSAESMLAISASAALG-QNLSA 415

NtrY Pd 440 AVPEFAPLPERLAQSVNESVQDEIRLITREGRV-ESLLVLRMAIRR-GAAGGLEGYVALDDVTDLVS-AQRMAAGDVARVREH-EIKNP 524
NtrY Ab 443 LVPEMGGLEAEAGPRGLVVDQIKIRRECDTTLTLVLRISTE-GRGSGMMRGYVITREDEHRTVSFAQRKAARARRPIAASPTRVKNP 531
NtrY Ac 436 VVPETAGLLEAEHARQSVQGNITLITRECRE-RVFAVVRVTE--QSPAEAHGMYVITDDITELIS-AQRSAWADVARRIAH-EIKNP 519
NtrY Ba 442 IAPEVGGQAFVARATGRTVHREQVSMARCGGS-RSFNQVQVTE--DAESEDHSYVITVDDITDLVQ-AQRSSAWADVARRIAH-EIKNP 526
NtrY Cc 413 LAPEFETVMTLSSESREP-TDVEVDIMREGCT-----RRLRVR--AAGHFAEGIVLITDDITRLVA-AQRNPAWKDVARVIAH-EIKNP 490
NtrY Sm 439 VAPEIEAVLVEAESRYRNDYRKQINIMRGCTE-RTMNQVQVRE--EGDESHGSYVITVDDITDLVI-AQRSTAWADVARRIAH-EIKNP 522
NtrY Ml 416 IIPHVRGVFEIGRQSGKPVYREQVFFRAGTE-RTFNQVITIEAGDDGSEKSYVITVDDITDLVQ-AQRSSAWADVARRIAH-EIKNP 501

NtrY Pd 525 LTPIQLSAERLKRKFGPIAG-EEREALQYTEVITRQVNDLRRIVDFEFSRFAFMPPEPRKETDI AKLLRDAELMQRDALQ--GALVSDIP 611
NtrY Ab 532 LTPIQLSAEPCAASTLKEIT-SDTEVFTMCTDITRQVVDLIRRMVDEFSAFARMPQPMVKPCNLDLVRQAVFLQSSAHAGIKFDMALP 620
NtrY Ac 520 LTPIQLSAERLKRKFGRHVT-QDREIFDQCDDTITRQVVDLIRRMVDEFSAFARMPKPVVDSQDMSEIIRQTVFLMRVGHV-EVVFDSPEV 607
NtrY Ba 527 LTPIQLSAERIRRRYGVKIT-EDREVFDQCDDTITRQVVDLIRRMVDEFSAFARMPKPEMRHMDLREALREASFLIEVSRSDIHFDNDFG 614
NtrY Cc 491 LTPIQLSAERLRRYKKEIA-SDLETDFRCDTITRQVVDLIRRMVDEFSAFARMPAERFEPNSLITMLRQAVFAQRFQDAIEVRLPEEPG 579
NtrY Sm 523 LTPIQLSAERLRRYKQIDQEDRAVFDQCDDTITRQVVDLIRRMVDEFSAFARMPKPTKEKSDLRALIKDAVFLREMGNS-HINFVRDFG 611
NtrY Ml 502 LTPIQLSAERLRRYKGVKIT-EDREVFDQCDDTITRQVVDLIRRMVDEFSAFARMPKPEMKAIDLRESLREASFLIEVSRA-DITFERIFG 589

NtrY Pd 612 DEPVI VDA DAGMMRQVFTNLLKNAGEBAIDELRENPPED--WTPQVHVKLVRVAPDSVAITRIDNGGGLP-DDRTRLEPEYVITMKPGGTGLG 698
NtrY Ab 621 QGPLTVQDSRQISCALTNLLKNADATEGRPPPAEGETLPPGHVAVIRVEADAERIAMIIDNCKGLPTEERDRLTEPVVITRAKGTGLG 710
NtrY Ac 608 P-AMPARDRLRLVSCALTNLLKNABABAEAVPPDVRG---QGRIRVSANRVGEDLVEDIIDNGTGLPQESRNLLEPEYVITREKGTGLG 692
NtrY Ba 615 SEKLGSRFDRSLICAFGNVKNASBAIDAVAKEERG---EGHIRIHAYRADGQLVVDVIDNCKGLPDDDRQKLEPEYVITREKGTGLG 700
NtrY Cc 580 DGDVWITSDERMVQCALTNLLKNAGEAVGARRANEPEL---QGRIATLVCEGDELCTVVEDNCGVLPKDRDRLTEPVVITREKGTGLG 666
NtrY Sm 612 DEPLEGQDGRMLCQAFGNVKNABABAEAVPAGTSRG-APTVVIRSRDDTTGRFVVDVIDNCKGLPTEENRRIIEPEYVITMRDKGTGLG 700
NtrY Ml 590 NEPLKGTDRSRLICAFGNVKNABABADGLEQKDGSD---HGIIRIQAGRQNGAIRIDVIDNCKGLPRENRQRLLEPEYVITREKGTGLG 675

NtrY Pd 699 LAIVKKIIVEDHGGSLMIDAPE---GRGMAEIRLPRE-----RQPVRLSARRHKHEQETLK----- 752
NtrY Ab 711 LAIVKKIIVEDHGGVLTLEDREG---GGRVGLVVIQHIIPASG-----TAAGDAPGGVGTPTAETGEKRRHAHGA----- 777
NtrY Ac 693 LAIVGKIIVEDHGGSIENDAPE---GRGAVIRLTLKAEQKAEPTDASTKATGAAATPAAPASAMARDAADASAARGKNERT----- 771
NtrY Ba 701 LAIVRKIIVEDHGGNLEHDAPADFHAGRGAMIRMVFPED-----PPIPSDASDNGEKHTAGQVI----- 760
NtrY Cc 667 LAIVKRIVEDHGGSLAVDARE---PPGARVVMKFP-----TTARLPVAAQSGVEEMI----- 716
NtrY Sm 701 LAIVKKIIVEDHGGDLLEDAPPDFDGGAGAMIRVVLP-----PAGETGGEDNLKDKGNTNGG----- 757
NtrY Ml 676 LAIVKKIIVEDHGGLELEDAPADFHGGRGAMISITLPLA-----AVTPLRGEKTEHERETEKVNGV----- 738

B

```

**                               *                               *                               *
NtrX Pd 1  -MSDILLVDDBRDIRELISDILRDECFSTRLEANSDCVVELNAAEPATMI LDIWLKDKSKMDGIDHDKQVKRNNPDVPIVHISGHGNIET 89
NtrX Ab 1  MAHDILLVDDBADIRMLIAGILNDEGMKTRERADADQAFRAQVSARRPESVVLVDIWLQGSRIIDGLQIIEQLMRDHRNLPVIMISGHGNIET 90
NtrX Ac 1  MAHDILLVDDBEDISGLVAGILEDEGYARTRADADGALAEIARRFNIFLDIWLQGSRIIDGLLELDIIKREHPEVPVVMISGHGNIET 90
NtrX Ba 1  MAADILLVDDBDIRDVLVAGILSDEGHETRTAFDADSALBAINDRAPRFVLDIWLQGSRIIDGLALDEIKKQHPPELVPVIMISGHGNIET 90
NtrX Cc 1  MSADVLLVDDBADIRDVLVAGILEDEGYAVRTRADSDGALAIRARKEPALLVLDIWLQGSRIIDGLLELDIMVKALDADLPVIMISGHGNIET 90
NtrX Sm 1  MAADILLVDDBEDIRELVAGILSDEGHETRTAFDADSALBAINDRVPRFVLDIWLQGSRIIDGLALDEIKNRHPDLPVIMISGHGNIET 90
NtrX M1 1  MASDILLVDDBEDIRELVAGILSDEGHETRTAFDADSALBAIADRPRFVLDIWLQGSRIIDGLALDEIKTMHPTLPVIMISGHGNIET 90

* **                               *                               *                               *
NtrX Pd 90 AVAAIKQGGAYDFIEKPKENITQLLVVIRRAEMETARTRRENATIKRGEARAAEMIGNSAPFRRIEQLDKVAKSNGRVMLTICEFGKGIAP 179
NtrX Ab 90 AVSAIKIGAYDFIEKPKKARLLLLMVDRAIEAARIKRENEELKLRAGGEVELIGRSTAVNHVRQSIIEKVSPTGSRVLIITCPAGSGKGVVA 180
NtrX Ac 90 AVAAIKRGGAYDFIEKPKENARLVVITERALELTLIRREVRRELKQLT-QPHTMVGRSSVIQQLRATVDRVGTNSRILIVGSPSGKGLITA 179
NtrX Ba 90 AVSAIKRGGAYDFIEKPKKARLLLVABRALETSILKREVSDDLKRTGDQLELVGTSLAMNQLRQTIERVAPTNSRIMITCPAGSGKGLIVA 180
NtrX Cc 90 AVSAIKRGGAYDFIEKPKKARLLLLIVERALEAAGLRRENRRLRQAQLSPDGLICKSAPAQALRQLLLKVPANSRVLSVCPGSGKGLIVA 180
NtrX Sm 90 AVSAIKRGGAYDFIEKPKKARLLLLIABRALENSILKRENSLRKSGDPVELICTSVAVSQLRQMIKVVAPTNSRIMITCPAGSGKGLIVA 180
NtrX M1 90 AVSAIKRGGAYDFIEKPKKARLLLLIABRALETSILKREVSDDLKQRSGETFDLIGMSSAMSQLRQTIERVAPTNSRVMITCPAGSGKGLIVA 180

+
NtrX Pd 180 RYVHAQSPRAREFVTVPCITLIEFEHMEBEVLFGEETAERICIEP--LLIFQAHGCVIYHDEVADMFLCTCPKILRVLTGECQFCFRAGCTDKV 267
NtrX Ab 181 RLIIHARSRRAGGPFVGLNQTMRPDRLEMELEFGIEAGVDCGGRKICITFQAHGCVIYHDEVADMFLCTCGKIVRAIQOCVFRVGGGQRV 270
NtrX Ac 180 RMIHAASARAQGFVVINAPATITPERLEYLELFGIEE-GEGRERHRGALBEAHGCVIYHDEVADMFLCTGRVIRVIVGECVFRVGGGQRV 268
NtrX Ba 181 RAIHAQSSRRANGPFVTVNAATITPERMELELFGIEE--MDGGERKVCALBEAHGCVIYHDEVADMFLCTGNKILRVIVGECVFRVGGGQRV 268
NtrX Cc 181 RLIIHARSRRAGGPFVAVSAPGMAPERLDVLELFGIEE-EGGRPRKICVFRRAHGGCVIYHDEVADMFLCTGRVIRVIVGECVFRVGGGQRV 269
NtrX Sm 181 RMIHRSKARANGPFVALNAPATITPERMELELFGIEE-TTCQPRRTCALBEAHGCVIYHDEVADMFLCTGNKILRVIVGECVFRVGGGQRV 269
NtrX M1 181 RAIHLSARKGAPVTLSPANTITPERMELELFGIEE--SNVVERKVCALBEAHGCVIYHDEVADMFLCTGNKILRVIVGECVFRVGGGQRV 268

NtrX Pd 268 RVDLRFVSSNTRDIAABEIAAGRFRQELVLRNINNVVAVVPSLABRRDDIPMLATHITELFHASCCLPQRALPEETSALCAMNWPGNVROL 357
NtrX Ab 271 EVDVRFVVATSNRDIQABIDQCRFRQDLFYRLAVVIRVPSLABRRDDIPMLARHFMQRSAEAAELPARDFGEDAMAALQAYDWPGNVROL 360
NtrX Ac 269 RVDVRFVSSNTRDIAACRFREDLYHRLSVMVIRVPSLABRRDDIPDLVDFEIDLISQTTCLQRKRVGEDAMAVLQSHDWPGNVROL 358
NtrX Ba 269 KVDVRFVSSNTRDIAAGRFREDLYHRLSVMVIRVPSLABRRDDIPSLVEFRMKQIAEQACIKFRKIGPDAMAVLCAHSWPGNVROL 358
NtrX Cc 270 QVDVRFVSSNTRDIAAGRFRREDLYHRLSVMVIRVPSLABRRDDIPELINVFVERISEATCLARRRIGEDALAVLQVQAWPGNVROL 359
NtrX Sm 270 KVDVRFVSSNTRDIAAGRFREDLYHRLSVMVIRVPSLABRRDDIPFLVDMFMRQVSEQACIRERKIGEDALAVLCAHDPGNVROL 359
NtrX M1 269 KVDVRFVSSNTRDIAAGRFREDLYHRLSVMVIRVPSLABRRDDIPFLVDMFMRQVSEQACIKFRKIGEDALAVLCAHDPGNVROL 358

NtrX Pd 358 RNVIERVILLGDGT--GPIQPSLEESQNGS-AGTSEALALGPQITSLALREARELFEREYLLACINRFGGNISRFAEIVGMERSALHRK 443
NtrX Ab 361 RNVVDWLLIMAQGD-PKEPIRADQPPPEIGAITPTVLKWDKGGEIMGLLEAREAREVFEREYLLACVIRFGGNISRFAEIVGMERSALHRK 449
NtrX Ac 359 RNNVERLLILAGGD-PDAEVTASMLPPDVGALVPTLPNGNGGHEMLGLLEAREAREVFEREYLLACINRFGGNISRFAEIVGMERSALHRK 447
NtrX Ba 359 RNNVERMLILTRGDDDELVTADLIPAEIGDTLPAPT-ESDQHIMALLEAREAREVFEREYLLACINRFGGNISRFAEIVGMERSALHRK 447
NtrX Cc 360 RNNVERMLILASGE-PGDVITAEMLSGAEQPSAGNAGA-IGAERI IALEAREAREVFEREYLLACVIRFGGNISRFAEIVGMERSALHRK 447
NtrX Sm 360 RNNIERLLILARSQDPDTPITADMLPNEVGDTLPKISA-QGDQHIMTLEAREAREVFEREYLLACINRFGGNISRFAEIVGMERSALHRK 448
NtrX M1 359 RNNVERMLILARGDDVDAPTADLIPSEIGDVMPTPN-QSDQHIMALLEAREAREVFEREYLLACINRFGGNISRFAEIVGMERSALHRK 447

NtrX Pd 444 LKSLGVVGGMRVEEEMGK 462
NtrX Ab 450 LKSLGVHGEKGLFVE-- 466
NtrX Ac 448 LKALGVG----- 454
NtrX Ba 448 LKSLGV----- 453
NtrX Cc 448 LKSLGVSGARGDEEE---- 462
NtrX Sm 449 LKSLGV----- 454
NtrX M1 448 LKSLGV----- 453
```

Figure S2




Figure S3

FIGURE LEGENDS

Figure S1. Phylogenetic trees with NtrB (A) and NtrC (B) homologues. The evolutionary history was inferred using the Neighbor-Joining method [11]. The bootstrap consensus tree was inferred from 3000 replicates [12]. Branches corresponding to partitions reproduced in less than 50% bootstrap replicates are collapsed. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (3000 replicates) are shown next to the branches [12]. The evolutionary distances were computed using the p-distance method and are in the units of the number of amino acid differences per site. The analysis involved 18 amino acid sequences. All positions containing gaps and missing data were eliminated. There were a total of 330 positions in the final dataset. Evolutionary analyses were conducted in MEGA5 [13]. The *P. denitrificans* sequences are highlighted by arrows.

Figure S2. Sequence alignment of the two-component regulatory system NtrY (A) and NtrX (B). Alignment was performed using ClustalW. Identical and similar residues are shaded in black and grey respectively. Regions of selected sequences are highlighted by boxes (located below their respective sequences) and show significant matches with the protein family database (Pfam-A, <http://pfam.sanger.ac.uk/>) for the following domains: A. HAMP linker domain (blue box), PAS domain involved in many signaling proteins where they are used as a signal sensor domain (red box), phospho-acceptor domain (green box), ATPase domain of histidine kinases (black box). B. Response regulator receiver domain (blue box), sigma54 interaction domain (red box); and helix-turn-helix (HTH) structural motif capable of binding DNA (green box). Asterisks define the position of conserved residues at the active site, triangles indicate the phosphorylation site, crosses define the ATP binding site, and points indicate the heme binding pocket. Pd, *Paracoccus denitrificans* PD1222 (NtrY, gi|119377428 and NtrX, gi|119377427); Ab, *Azospirillum brasilense* Sp245 (NtrY, gi|20451353 and NtrX, gi|119377427); Ac, *Azorhizobium caulinodans* ORS 571 (NtrY, gi|38737 and NtrX, gi|38738); Ba, *Brucella melitensis* biovar Abortus 2308 (NtrY, gi|123754544 and NtrX, gi|82699956); Cc, *Caulobacter crescentus* CB15 (NtrY, gi|16125986 and NtrX, gi|16125987); Sm, *Sinorhizobium meliloti* SM11 (NtrY, gi|336033438 and NtrX,

gi|336033437); Ml, *Mesorhizobium loti* MAFF303099 (NtrY, gi|14021375 and NtrX, gi|14021376).

Figure S3. Schematic overview of the nitrate-regulated systems in *Paracoccus denitrificans*. Transcriptomic data reveal systems that are up-regulated (\uparrow) or down-regulated (\downarrow) by nitrate. The fold changes are indicated. Proteomic analysis also corroborated these results.

REFERENCES

1. de Vries, G. E., Harms, N., Hoogendijk, J., and Stouthamer, A.-H. (1989) Isolation and characterization of *Paracoccus denitrificans* mutants with increased conjugation frequencies and pleiotropic loss of (nGATCn) DNA-modifying property. *Arch. Microbiol.* **152**, 52-57
2. Luque-Almagro, V. M., Lyall, V. J., Ferguson, S. J., Roldán, M. D., Richardson, D. J., and Gates, A. J. (2013) Nitrogen oxyanion-dependent dissociation of a two-component complex that regulates bacterial nitrate assimilation. *J. Biol. Chem.* **288**, 29692–29702
3. Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989) Molecular cloning. In *A laboratory manual*, 2nd edn. New York: Cold Spring Harbor Laboratory, Cold Spring Harbor
4. Figurski, D. H., and Helinski, D. R. (1979) Replication of an origin-containing derivative of plasmid RK2 dependent on a plasmid function provided in trans. *Proc. Natl. Acad. Sci. USA* **76**, 1648–1652
5. Yanisch-Perron, C., Vieira, J., Messing, J. (1985) Improved M13 phage cloning vectors and host strains: nucleotide sequencing of the M13mp18 and pUC9 vectors. *Gene* **33**, 103-119
6. Simon, R., Priefer, U., and Pühler, A. (1983) A broad host range mobilization system for *in vivo* genetic engineering. *Biotechnology* **1**, 784–791
7. Gates A. J., Luque-Almagro, V. M., Goddard, A. D., Ferguson, S. J., Roldán, M. D., and Richardson, D. J. (2011) A composite biochemical system for bacterial nitrate and nitrite assimilation as exemplified by *Paracoccus denitrificans*. *Biochem. J.* **435**, 743-753
8. Schafer A., Tauch A., Jager W., Kalinowski J., Thierbach G., Pühler A. (1994) Small mobilizable multi-purpose cloning vectors derived from the *Escherichia coli* plasmids pK18 and pK19: selection of defined deletions in the chromosome of *Corynebacterium glutamicum*. *Gene* **145**, 69-73
9. Frigaard, N. U., Li, H., Milks, K. J., and Bryant, D. A. (2004) Nine mutants of *Chlorobium tepidum* each unable to synthesize a different chlorosome protein still assemble functional chlorosomes. *J. Bacteriol.* **186**, 646–653

10. Spaink, H. P., Wijffelman, C. A., Pees, E., Okker, R. J. H., and Lugtenberg B. J. J. (1987) *Rhizobium* nodulation gene *nodD* as a determinant of host specificity. *Nature* **328**, 337-340
11. Saitou N. and Nei M. (1987) The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* **4**, 406-425
12. Felsenstein J. (1985) Confidence limits on phylogenies: An approach using the bootstrap. *Evolution* **39**, 783-791
13. Tamura K., Peterson D., Peterson N., Stecher G., Nei M., and Kumar S. (2011) MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* **28**, 2731-2739