



**Identification of selective protein-protein interaction inhibitors using efficient in silico peptide-directed ligand design.**

Journal:	<i>Chemical Science</i>
Manuscript ID	SC-EDG-01-2019-000059
Manuscript Type:	Edge Article
Date Submitted by the Author:	04-Jan-2019
Complete List of Authors:	Beekman, Andrew; University of East Anglia, School of Pharmacy Cominetti, Marco Maurizio Daniele; University of East Anglia, School of Pharmacy Walpole, Samuel; University of East Anglia, Pharmacy Prabhu, Saurabh; University of East Anglia Faculty of Medicine and Health Sciences O'Connell, Maria; University of East Anglia, Pharmacy Angulo, Jesus; University of East Anglia, School of Pharmacy Searcey, Mark; University of East Anglia, School of Pharmacy



Journal Name

ARTICLE

## Identification of selective protein-protein interaction inhibitors using efficient *in silico* peptide-directed ligand design.

Andrew M. Beekman,<sup>\*a</sup> Marco M. D. Cominetti,<sup>a</sup> Samuel J. Walpole,<sup>a</sup> Saurabh Prabhu,<sup>a</sup> Maria A. O'Connell,<sup>a</sup> Jesus Angulo<sup>a</sup> and Mark Searcey<sup>\*a,b</sup>

Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

The development of protein-protein interaction (PPI) inhibitors with therapeutic value is of increasing importance as the first clinical agent has now been approved, but PPIs remain difficult targets for the development of small molecule ligands. This article describes a highly efficient approach to the development of inhibitors of the p53/*hDMX* or *hDM2* interaction that involves the design of small molecules *in silico* based upon a peptide/protein structure. The process for molecule design, starting from a virtual library of just over 1200 fragments, led to the eventual synthesis of twenty compounds, of which ten bound to either *hDM2*, *hDMX* or both in vitro binding assays. This 50% success rate is extremely efficient compared to traditional high throughput screening. The identification of two selective *hDMX* inhibitors from twenty compounds highlights this efficiency as, to date, only two other *hDMX*-selective agents exist in the literature. Preliminary biological studies show that 20% of the compounds identified have cellular activity and activate downstream pathways associated with p53 activation.

### Introduction

Targeting the *hDM2* (human double minute 2) p53 protein-protein interaction (PPI) has been a paradigm for PPI inhibitor development in cancer since the discovery of nutlin-3, which is currently undergoing clinical trials.<sup>1,2</sup> Although there are many studies of p53-*hDM2* interaction inhibitors ranging from natural products to designed small molecules, the scope of this promising strategy has been severely limited by the activity of *hDMX*.<sup>3</sup> Despite the highly homologous nature of *hDMX* to *hDM2* the two proteins do not perform redundant roles.<sup>4,5</sup> *hDMX* can prevent p53 activity by sequestering it, but *hDMX* also enhances the ligase activity of *hDM2* towards p53 by forming *hDM2/hDMX* dimers.<sup>6</sup> The overexpression of one or both of these proteins results in the loss of p53 activity in cells.<sup>7</sup> Therefore, dual *hDM2/hDMX* inhibitors have become highly desirable.

The development of *hDMX* selective molecules is also of significant importance. It has been shown that direct inhibition of *hDM2* leads to p53 dependent toxicity in healthy cells.<sup>8,9</sup> Additionally, the non-redundant roles and common mutual exclusivity in cancers suggest selective compounds are useful.<sup>10</sup> There are currently no small molecules that modulate

*hDMX/p53* and exhibit both cell and animal efficacy.<sup>11</sup> Two compounds have been highlighted which modulate *hDMX/p53* selectively, but are unsuitable for clinical development.<sup>8,9,12</sup> Therefore, there is still a demand for chemical probes or leads that are selective for *hDM2*, selective for *hDMX* or dual *hDM2/hDMX* inhibitors. The p53 binding domains of *hDM2* and *hDMX* display high structural homology, but differences exist which could be exploited.<sup>7,10</sup>

Several approaches to the identification of inhibitors of the p53-*hDM2* and p53-*hDMX* interaction have been described. Natural products such as chlorofusin can be utilised as starting points for design.<sup>13</sup> Oligobenzamide analogues that mimic the interactions of the helical peptide have been described<sup>14,15</sup> and helical peptides have been stabilised and shown to have cellular activity through the introduction of stapling groups.<sup>16</sup> Both high throughput screening and computational design against the *hDM2* target have had varying degrees of success with hit rates for screening of around 0.02%.

We believe these challenging PPIs are excellent targets for the exemplification of *in silico* peptide directed binding, a new methodology for identifying selective PPI modulators. Using copper(I) catalysed azide-alkyne cycloaddition (CuAAC) click chemistry and taking a small commercially available library of 896 alkynes and 214 azides would require the synthesis of 191,744 compounds to screen all possible compounds. Using these compounds as fragments for screening would need high resolution NMR with <sup>15</sup>N-labelled protein or high throughput crystallography, relatively specialised techniques. Our approach has been to exploit the peptidic dual inhibitor Ac-Phe-Met-Aib-Pmp-6-Cl-Trp-Glu-AC<sub>3</sub>C-Leu-NH<sub>2</sub> (**1**) to develop small molecule

<sup>a</sup> School of Pharmacy, University of East Anglia, Norwich Research Park, Norwich, Norfolk, NR47TJ, United Kingdom

<sup>b</sup> School of Chemistry, University of East Anglia, Norwich Research Park, Norwich, Norfolk, NR47TJ, United Kingdom.

† E-mail: A.Beekman@uea.ac.uk, M.Searcey@uea.ac.uk.

Electronic Supplementary Information (ESI) available: Experimental procedures, spectra and processed biophysical assay data. See DOI: 10.1039/x0xx00000x

probes that target *hDM2*<sup>17</sup> and *hDMX*.<sup>18</sup> Briefly, the crystal structure of **1** bound to both proteins is known (*hDMX*: PDB ID 2GV2, *hDM2*: PDB ID 3FEA). The sequence of the high affinity peptide binder is used to generate two smaller peptides with reactive azide and alkyne terminals (Figure 1).<sup>19</sup> *In silico* screening of the small molecule fragments then allows the identification of peptide/small molecule hybrids with restored affinity for the target site. The restoration of binding affinity implies that the small molecule fragment in some way emulates the peptide section it has replaced. The small molecule portion of the hybrid hits are then combined through *in silico* click chemistry and rescreened to identify potential small molecules with high affinity for the target site. Computational modelling is used to perform the entire peptide directed binding process, identifying small molecules to be prepared. This use of virtual design in peptide directed binding further improves the rapid and economic nature of this process, identifying compounds not highlighted by experimental peptide directed binding.<sup>19</sup> Herein, we demonstrate that this approach led to the synthesis of twenty small molecules of which ten bound to *hDM2* or *hDMX*, representing a 50% success rate. Furthermore, one compound was selective for *hDMX* both in vitro and in tissue culture, outperforming previously reported selective modulators.

## Results

### *In silico* peptide directed binding identifies candidate small molecule *hDM2/X* binders

Peptide **1** binds with high affinity to both *hDM2* and *hDMX*, which allows its exploitation to develop small molecule dual inhibitors as both chemical probes and potential therapeutic lead molecules. In order to identify small molecules to be synthesised and evaluated against *hDM2* and *hDMX* the crystal structures for the peptide 8-mer **1** bound to *hDM2* (PDB ID: 2GV2)<sup>17</sup> (Figure 2A) and *hDMX* (PDB ID: 3FEA)<sup>18</sup> were edited to identify small molecule fragments that emulate a section of the 8 mer (Figure 2B and 2C). The peptide was split, Phe-Met-Aib-Pmp-propargyl glycine and azidoacetamide-6-Cl-Trp-Glu-*Ac*<sub>3</sub>-Leu (Figure 1B), allowing two key binding residues to be present in both smaller peptides (Phe and Pmp, 6-Cl-Trp and Leu). The site left vacant by the removed peptide fragment represents a more tractable binding site for a small molecule. Covalent docking was then performed simulating the Huisgen 1,3-dipolar cycloaddition (see SI, pg 6), with the commercially available library of alkynes or azides (SI, pg 12). The semi-peptide was held in place with flexibility in the reactive terminal and the final residue, allowing for the possibility that the added small molecule fragment is held in the correct orientation for binding. The results were scored and ranked<sup>20</sup> and the top ten azides and top ten alkynes that were synthetically viable were used to generate one hundred virtual triazoles. The generated triazoles were then induced-fit docked to the protein, scored and ranked




Figure 1. Concept of peptide directed binding utilised in this work with *in silico* methods to improve the rapid and economic nature of finding new PPI modulators. Peptide **1** binds to *hDM2/X*, and is separated in to two smaller peptides with no appreciable binding despite having two key residues each. Azide-alkyne cycloaddition (AAC) identifies small molecule fragments which restore binding affinity. The identified small molecule fragments can be combined to generate a small molecule substitute for the initial peptide.

to identify the top ten small molecules that bound in silico with high affinity to each protein and that would be synthesised (Figure 2D).<sup>21</sup> The identified triazole small molecules are likely to bind with the same binding elements as identified by the hybrid, but may sit in a slightly altered orientation to allow for tighter binding. This process was carried out for both *hDM2* and *hDMX* and as such twenty small molecule triazoles were synthesised following published procedures with CuSO<sub>4</sub> and sodium ascorbate in a <sup>1</sup>BuOH/H<sub>2</sub>O mix, followed by purification by reverse phase HPLC (See SI pg. 9 for all synthesised compounds).<sup>13</sup>











### Identified small molecules selectively modulate the p53-*hDM2/X* PPIs

It was expected that some of these molecules would display selectivity for the target used in their design, but inevitably dual modulators would be identified due to the homogeneity of the *hDM2/X* binding sites and the use of the same peptide **1**. The synthesised compounds were first analysed with competitive fluorescence anisotropy (FA) assays. A FAM tagged p4 peptide<sup>22, 23</sup> was used as the fluorescently tagged peptide for both *hDM2*, reported previously,<sup>13</sup> and *hDMX*. The wild-type p53 peptide (residues 15-27) was employed as a positive control and an indication that the FA assays were performing adequately. Of the twenty compounds synthesised, 50% demonstrated binding in the FA assays to either *hDM2* or *hDMX*, with IC<sub>50</sub> values less than 100 μM (Table 1). Strikingly, one compound displayed significant selectivity for *hDMX* over *hDM2* (**5**), while overall 40% of compounds designed for *hDMX* modulated the *hDMX* interaction (**2, 3, 4, 5**). Two of these compounds, **2** and **3**, had

## Journal Name

## ARTICLE

Table 1. IC<sub>50</sub> values for inhibition of the binding of FAM-p4 to *hDM2* and *hDMX* of small molecules.<sup>[a]</sup> A green background indicates the compound was designed for and was selective to the same protein. Yellow indicates the compound is a dual inhibitor. A red background indicates the compound was designed for one protein but was selective for the other protein.

Structure	<i>hDMX</i>	<i>hDM2</i>	Structure	<i>hDMX</i>	<i>hDM2</i>
	FA IC <sub>50</sub> (μM) [95% CI]	FA IC <sub>50</sub> (μM) [95% CI]		FA IC <sub>50</sub> (μM) [95% CI]	FA IC <sub>50</sub> (μM) [95% CI]
	0.013 [0.003, 0.046]	0.24 [0.16, 0.36]		>100	2.46 [1.01, 4.70]
	0.019 [0.005, 0.071]	0.10 [0.03, 0.76]		>100	5.77 [4.00, 8.37]
	1.28 [0.64, 2.52]	8.00 [6.12, 20.06]		>100	12.42 [8.61, 21.23]
	1.49 [1.06, 2.10]	>100		10.97 [8.28, 19.66]	22.44 [17.92, 29.60]
	0.97 [0.79, 1.44]	1.74 [0.89, 3.38]		1.17 [1.02, 4.20]	>100

[a]IC<sub>50</sub> values determined by non-linear regression of at least three independent experiments (see SI, pg 10). Errors are 95% confidence intervals (CI). Fmoc, 9-fluorenylmethylcarbonyl

an IC<sub>50</sub> in the tens of nanomolar range. Of the compounds designed for *hDM2*, 50% demonstrated the ability to modulate the *hDM2* PPI (**6**, **7**, **8**, **9**, **10**), with three showing selectivity for *hDM2* over *hDMX* (**7**, **8**, **9**). Interestingly, one of the compounds designed for *hDM2* (**11**) had affinity for *hDMX* but no appreciable binding to *hDM2*. To confirm selectivity, these compounds were analysed in other PPI FA assays routinely utilised in our lab. Pleasingly, compounds **2-11** demonstrated no appreciable binding at 100 μM towards the PPIs Mcl-1/Noxa, Bcl-2/Bid<sup>19</sup> and Nrf2/Keap1,<sup>24</sup> despite similarities to compounds reported to bind to these PPIs.

The compounds that exhibited binding in the FA assay were validated with differential scanning fluorimetry (DSF).<sup>25</sup> Compounds displayed the ability to increase the melting temperature of proteins compared to a vehicle control (see SI, pg. 41).<sup>26</sup> Thermal stability was demonstrated using the

hydrophobic dye Sypro orange, mimicking the observed FA assay results (see SI, pg. 40).

#### Use of STD NMR indicates that compounds bind in the p53 binding site of *hDM2/X*

To provide structural details of the binding of these compounds in the p53 binding site, we employed an STD NMR initial growth-rates approach to identify the binding mode and the interactions with the binding grooves of *hDM2* and *hDMX*.<sup>27-29</sup> Compound **4** was chosen for analysis by STD NMR because of its solubility in aqueous buffer and its relatively weak binding to both *hDM2* and *hDMX*, allowing the binding kinetics to fall within the fast-exchange conditions necessary for STD NMR. This criteria precludes tighter binders (**2** & **3**) from being analysed with this technique. Compound **4** displayed STD

signals in the presence of either *hDM2* or *hDMX*, indicating binding to both proteins.




Figure 2. Examples of competitive-displacement assays of small molecules towards a selection of PPIs. Compounds **5** and **10** used to exemplify the titration of compounds towards the *hDMX/p53*, *hDM2/p53*, *Mcl-1/NOXA*, *Bcl-2/Bid* and *Nrf2/Keap1* interactions. Dose response curves can be seen for **5** towards *hDMX*, and for **10** towards *hDMX* and *hDM2*, with no competitive binding towards any other PPIs observed.

Additionally, the ligand binding epitope of compound **4** was mapped using the STD-NMR initial growth rates method (SI, pg 15). In order to gain some structural information docking calculations were used to identify a binding pose. We were able to validate the binding pose within the *hDM2* binding groove using CORCEMA-ST,<sup>30</sup> which calculates theoretical STD intensities based on a given 3D model complex. The NOE R-factor between the experimental and theoretical STD intensities was 0.17, indicating good agreement. Interestingly, preliminary CORCEMA-ST trials of **4** binding to *hDMX* suggested the formation of oligomers. Dimerization induced by structurally similar compounds has been observed previously,<sup>8</sup> and suggests a possible selective binding mode which could be exploited for *hDMX* modulation. Figure 3 displays the binding pose of **4** to *hDM2* demonstrating tight binding of the fluorinated aryl ring in the pocket vacated by chlorotryptophan of **1**. This STD/CORCEMA-ST validated binding pose, along with the binding affinities determined by the competitive fluorescence anisotropy assay indicate these compounds are binding in the

*p53* binding groove. The identified binding epitope places the fluorine atom in the same pocket as that occupied by the chlorine atom of the Cl-Trp in **1**. This same pocket also exists in the *hDMX* binding site and may explain the improved binding of the fluoroaryl group (**2**, **10** & **11** compared to **S8**, **S2** & **S9**).




Figure 3. Binding pose of **4** (purple) to *hDM2* in the *p53* binding site determined by CORCEMA-ST, with an overlay of peptide **1** (orange) showing the binding residues of Leu and Cl-Trp. The surface of *hDM2* is cut along the plane of the centroid of the aromatic rings of compound **4**.

#### Compounds inhibit cancer cell growth in a selective fashion

Compounds that had activity in the FA assay were examined for their ability to inhibit cell growth and affect metabolism in an array of cancer cell lines that demonstrate varying expressions of the *p53* and *hDM2/X* proteins. The human cancer cell lines A549,<sup>31</sup> HCT116<sup>3,31</sup> and MCF-7<sup>31</sup> express both *hDM2* and *hDMX*, SJS-1 is *hDM2* dependent<sup>11</sup> and JEG3 is *hDMX* dependent.<sup>3</sup> All cell lines express wild type *p53*. Gratifyingly, 10% of the prepared compounds demonstrated the ability to inhibit cell growth. Compound **10**, which was shown to bind to both *hDM2* and *hDMX*, had activity towards cell lines that are dependent on *hDMX*, dependent on *hDM2* or dependent on both, as expected (Table 2). Compound **5**, which showed selectivity for *hDMX* over *hDM2*, demonstrated activity towards the *hDMX* dependent JEG3, but not towards the *hDM2* dependent SJS-1, a further suggestion of selectivity of **5** for *hDMX*. The remaining compounds did not demonstrate activity in the MTS assay, perhaps due to issues with cell penetration. Alterations to the in silico screening process are currently underway to increase the output of compounds with desirable characteristics, such as cell permeability.

Table 2. Cell growth inhibition of compounds which demonstrated activity towards cell line which are dependent on *hDM2*, *hDMX* or both, demonstrated by the MTS assay.<sup>[a]</sup>

	SJS-1 IC <sub>50</sub> (µM)	A549 IC <sub>50</sub> (µM)	HCT116 IC <sub>50</sub> (µM)	MCF-7 IC <sub>50</sub> (µM)	JEG3 IC <sub>50</sub> (µM)
<b>5</b>	>100	16.2 ± 0.06	20.5 ± 0.07	29.8 ± 0.13	35.5 ± 0.10
<b>10</b>	30.0 ± 2.29	65.7 ± 0.22	>100	>100	38.1 ± 0.03

[a] IC<sub>50</sub> values determined by non-linear regression of at least three experiments (see SI, pg 13). Errors are the transformed greater extreme of the standard error.




Figure 4. A) A549 and JEG-3 cells were treated with vehicle (DMSO), positive control (staurosporine, STS), **5** and **10** at 100 μM and 50 μM for 6 hr, 4 hr and 2 hr at 37 °C. Treatment with Caspase-Glo 3/7 reagent (Promega) allowed for the monitoring of the cleavage of pro-luminescent substrate by caspase 3/7. Data are mean ± SEM for at least n = 3. RLU – relative luminescent units. STS – staurosporine. B) JEG-3 cells were treated with vehicle (DMSO), **5** and **10** at 100 μM for 6 hr at 37 °C, and transcriptional upregulation of p53, hDM2, p21 and MIC-1 was evaluated by qPCR analysis. 18S was used as a control. Y-axis is the mRNA-fold induction vs the vehicle control (DMSO). All *p* values were less than 0.05 when compared to the vehicle.

#### Compounds induce events downstream of p53 activation

To ascertain whether compounds **5** and **10** were inducing p53 related cell death we examined the induction of apoptosis with a caspase 3/7 activation assay. A549 and JEG-3 cells were treated with vehicle (DMSO), positive control (staurosporine), **5** and **10**. Cleavage of the DEVD sequence was monitored to observe caspase 3/7 activity. Activation of caspase 3/7 was observed for both **5** and **10** in line with the observed IC<sub>50</sub> values (Figure 4A). The transcriptional activation of targets downstream of p53 was examined with quantitative PCR (qPCR).<sup>3</sup> JEG-3 cells were treated with vehicle control, **5** and **10** for 6 hr at 37 °C, followed by RNA isolation, reverse transcription, and qPCR analysis of the obtained cDNA using the primers of downstream p53 targets p53, hDM2, p21 and MIC-1. Both compounds **5** and **10** significantly increased the expression of each of the target genes when compared to the vehicle control (Figure 4B).

Virtual peptide directed binding was employed to identify new modulators of p53/hDM2 and p53/hDMX. This use of computational peptide directed binding represents a complementary method to identify PPI modulators in a rapid and economic manner. Twenty compounds were prepared, ten designed for hDM2 and ten designed for hDMX, and 50% of the compounds were found to modulate the PPI. This hit rate is a significant improvement compared to recent studies which utilise a chemical array (0.04%)<sup>32</sup> and high-throughput screen (0.07%).<sup>33</sup> Applying peptide directed binding to the p53-hDM2/X PPIs has identified new scaffolds for modulating these PPIs, which have excellent lead like features.<sup>34</sup> Importantly, this process has identified selective inhibitors for both hDM2 and hDMX, despite the high structural homology of the binding sites of these proteins. Only two small molecules have been previously reported which selectively modulate the p53-hDMX PPI, highlighting the power of this technique. Virtual peptide

directed binding has identified compounds that interact with the p53 binding site of hDM2 and hDMX, supported by activity in competitive fluorescence anisotropy assays and, more significantly, quantitative STD NMR analysis. A subset of the identified modulators was shown to have activity towards cancer cell lines that are known to have dependence on hDM2, hDMX or both, and these compounds were shown to induce p53 activation downstream events.

#### Discussion

The deployment of *in silico* peptide directed binding has demonstrated an extraordinarily high hit rate for new PPI small molecule modulators of the p53 hDM2/X interaction. Perhaps more interestingly, this process has identified new binding motifs for these PPIs, as well as highlighting selective scaffolds for hDMX, a notoriously challenging task.<sup>35, 36</sup> The nature of peptide directed binding provides structure activity relationship information without producing specific modifications. The fluoroaryl group shows enhanced binding to hDMX, and the 2,4-difluoroaryl compound may offer selectivity for hDMX (i.e. **5** and **11**). The Fmoc azoalanine drives binding to hDM2, a phenomenon we have shown previously,<sup>13</sup> but does not guarantee binding (**S1** – see SI, pg. 9). The presence of the sugar moiety imparts potent binding, but again does not guarantee it (**S8**). Despite the FP and DSF results supporting tight binding, the hydrophilic sugar moiety raises concerns about compounds **2** and **3** as leads, as the p53 binding site is known to be hydrophobic. Docking suggested the polar sugar occupying the Phe binding pocket of **1** (See SI, pg 14), but docking results should be interpreted with caution. Modifying the sugar with nonpolar protecting groups reduces potency, but affinity is still observed (i.e. **10** and **11**), suggesting key binding moieties may still be available. Investigation of **2** and **3** as leads is ongoing. The

*in silico* calculations highlighted the zidovudine structure (azide section of **9**) as a likely potent binder, but experimentally resulted in a poor hit rate, perhaps highlighting modifications to be made in compound selection process or docking calculations (see SI, pg 9).

The compounds identified which selectively modulate the p53-*hDMX* interaction outperform the two selective molecules reported in the literature. The first known selective binder of *hDMX* – SJ-172550 – was demonstrated to bind with less affinity to *hDMX* than p53. Both **5** and **11** demonstrate greater binding affinity to *hDMX* than p53 (SI, pg 10).<sup>35</sup> Additionally, the cellular activity of these compounds is reported in the millimolar range, with our compounds demonstrating at least two orders of magnitude improvement. SJ-172550 is also known to not be a competitive binder, but rather induces a conformational change through covalent bond formation, which does not exist in a reducing environment.<sup>37</sup> Unfortunately, this limits the value of SJ-172550 as a lead candidate.

CTX1, the second small molecule shown to have selectivity for *hDMX*, has not been validated in *in vitro* protein assays, with this compound being identified through a cellular screen.<sup>9</sup> CTX1 demonstrated IC<sub>50</sub> values in the tens to hundreds of micromolar range. Compound **5** possesses slightly improved potency, but has other important advantages. CTX1 is an acridine based molecule, known to bind to DNA, and act as a PAIN molecule. This characteristic makes CTX1 unusable in many light based assays, such as fluorescence, due to interference. It is also unknown whether CTX1 is a competitive inhibitor of p53 or acts through some other mechanism (some suggestion the CTX1 mode of action overlaps with the action of 9-aminoacridine),<sup>9</sup> making its use as a chemical tool limited.

SJ-172550 was found as one of three hits in a library of 295,848 compounds.<sup>35</sup> CTX1 came from a screen of over 20,000 compounds.<sup>9</sup> In this instance we have prepared twenty molecules and identified two selective p53-*hDMX* modulators. The true value of the significant advance in the improvement of success rate we have demonstrated here is the power of peptide directed binding to allow researchers in academia and the pharmaceutical industry, chemistry and biology to quickly and cheaply develop modulators for the protein-protein interaction they wish to target. The need for enormous libraries of molecules and expensive high throughput screening facilities is not necessary for the identification of small molecule modulators, as has been the case for so many challenging targets.

Notably, the compounds prepared are unoptimised but still display strong activity in the fluorescence anisotropy assay and low micromolar cellular activity, highlighting the rapid ability of peptide directed binding to identify selective chemical probes.

## Conclusions

This work has exemplified the power of peptide directed binding to rapidly identify small molecule PPI modulators, at a high hit rate, that show selectivity for challenging targets. This method complements experimental peptide directed binding

and is expected to be readily applied to other PPIs such as 14-3-3/tau, bromodomains or Nrf2/Keap1.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

SJW is funded by the Biotechnology and Biological Sciences Research Council (BBSRC) Doctoral Training Partnership PhD studentship. SP is funded by BBSRC grant no BB/N008448/1. We acknowledge the EPSRC UK National Mass Spectrometry Facility at Swansea University. We thank Professor Aart Jochemsen (Leiden) for the kind gift of *hDMX*.

## Notes and references

1. L. T. Vassilev, *J. Med. Chem.*, 2005, **48**, 4491-4499.
2. V. Tisato, R. Voltan, A. Gonelli, P. Secchiero and G. Zauli, *J. Hematol. Oncol.*, 2017, **10**, 133.
3. F. Bernal, M. Wade, M. Godes, T. N. Davis, D. G. Whitehead, A. L. Kung, G. M. Wahl and L. D. Walensky, *Cancer Cell*, 2010, **18**, 411-422.
4. V. Böttger, A. Böttger, C. Garcia Echeverria, Y. F. M. Ramos, A. J. van der Eb, A. G. Jochemsen and D. P. Lane, *Oncogene*, 1999, **18**, 189.
5. F. Toledo and G. M. Wahl, *Int. J. Biochem. Cell Bio.*, 2007, **39**, 1476-1482.
6. A. Shvarts, W. T. Steegenga, N. Riteco, T. Laar, P. Dekker, M. Bazuine, R. C. Ham, W. H. v. Oordt, G. Hateboer, A. J. Eb and A. G. Jochemsen, *EMBO J.*, 1996, **15**, 5349-5357.
7. Y. S. Chang, B. Graves, V. Guerlavais, C. Tovar, K. Packman, K.-H. To, K. A. Olson, K. Kesavan, P. Gangurde, A. Mukherjee, T. Baker, K. Darlak, C. Elkin, Z. Filipovic, F. Z. Qureshi, H. Cai, P. Berry, E. Feyfant, X. E. Shi, J. Horstick, D. A. Annis, A. M. Manning, N. Fotouhi, H. Nash, L. T. Vassilev and T. K. Sawyer, *Proc. Nat. Acad. Sci.*, 2013, **110**, E3445-E3454.
8. B. Graves, T. Thompson, M. Xia, C. Janson, C. Lukacs, D. Deo, P. Di Lello, D. Fry, C. Garvie, K. S. Huang, L. Gao, C. Tovar, A. Lovey, J. Wanner and L. T. Vassilev, *Proc. Nat. Acad. Sci.*, 2012, **109**, 11788-11793.
9. G. Karan, H. Wang, A. Chakrabarti, S. Karan, Z. Liu, Z. Xia, M. Gundluru, S. Moreton, Y. Sauntharajah, M. W. Jackson, M. K. Agarwal and D. N. Wald, *Mol. Can. Ther.*, 2016, **15**, 574-582.
10. S. Datta, M. E. Bucks, D. Koley, P. X. Lim and S. N. Savinov, *Bioorg. Med. Chem.*, 2010, **18**, 6099-6108.
11. F. Wachter, A. M. Morgan, M. Godes, R. Mourtada, G. H. Bird and L. D. Walensky, *Oncogene*, 2017, **36**, 2184.
12. G. M. Popowicz, A. Dömling and T. A. Holak, *Angew. Chem. Int. Ed.*, 2011, **50**, 2680-2688.
13. M. M. D. Cominetti, S. A. Goffin, E. Raffel, K. D. Turner, J. C. Ramoutar, M. A. O'Connell, L. A. Howell and M. Searcey, *Bioorg. Med. Chem. Lett.*, 2015, **25**, 4878-4880.
14. J. P. Plante, T. Burnley, B. Malkova, M. E. Webb, S. L. Warriner, T. A. Edwards and A. J. Wilson, *Chem. Commun.*, 2009, DOI: 10.1039/B908207G, 5091-5093.

15. A. Shaginian, L. R. Whitby, S. Hong, I. Hwang, B. Farooqi, M. Searcey, J. Chen, P. K. Vogt and D. L. Boger, *J. Am. Chem. Soc.*, 2009, **131**, 5564-5572.
16. F. Bernal, A. F. Tyler, S. J. Korsmeyer, L. D. Walensky and G. L. Verdine, *J. Am. Chem. Soc.*, 2007, **129**, 2456-2457.
17. K. Sakurai, C. Schubert and D. Kahne, *J. Am. Chem. Soc.*, 2006, **128**, 11000-11001.
18. J. Kallen, A. Goepfert, A. Blechschmidt, A. Izaac, M. Geiser, G. Tavares, P. Ramage, P. Furet, K. Masuya and J. Lisztwan, *J. Bio. Chem.*, 2009, **284**, 8812-8821.
19. A. M. Beekman, M. A. O'Connell and L. A. Howell, *Angew. Chem. Int. Ed.*, 2017, **56**, 10446-10450.
20. K. Zhu, K. W. Borrelli, J. R. Greenwood, T. Day, R. Abel, R. S. Farid and E. Harder, *J. Chem. Inf. Model.*, 2014, **54**, 1932-1940.
21. W. Sherman, H. S. Beard and R. Farid, *Chem. Biol. Drug Des.*, 2006, **67**, 83-84.
22. B. Hu, D. M. Gilkes and J. Chen, *Cancer Res.*, 2007, **67**, 8810-8817.
23. A. Czarna, G. M. Popowicz, A. Pecak, S. Wolf, G. Dubin and T. A. Holak, *Cell Cycle*, 2009, **8**, 1176-1184.
24. R. J. Steel, M. A. O'Connell and M. Searcey, *Bioorg. Med. Chem. Lett.*, 2018, **28**, 2728-2731.
25. F. H. Niesen, H. Berglund and M. Vedadi, *Nat. Protocols*, 2007, **2**, 2212.
26. A. Madhumalar, L. H. Jun, C. J. Brown, D. P. Lane and C. S. Verma, *Cell Cycle*, 2009, **8**, 2828-2836.
27. J. Angulo, S. A. Goffin, D. Gandhi, M. Searcey and L. A. Howell, *Chem. Eur. J.*, 2016, **22**, 5858-5862.
28. B. Bertrand, J. Fernandez Cestau, J. Angulo, M. M. D. Cominetti, Z. A. E. Waller, M. Searcey, M. A. O'Connell and M. Bochmann, *Inorg. Chem.*, 2017, **56**, 5728-5740.
29. J. Angulo, I. Díaz, J. J. Reina, G. Tabarani, F. Fieschi, J. Rojo and P. M. Nieto, *ChemBioChem*, 2008, **9**, 2225-2227.
30. V. Jayalakshmi and N. R. Krishna, *J. Magn. Reson.*, 2002, **155**, 106-118.
31. M. Hirose, K. Yamato, S. Endo, R. Saito, T. Ueno, S. Hirai, H. Suzuki, M. Abei, Y. Natori and I. Hyodo, *Oncoscience*, 2014, **1**, 830-843.
32. T. Noguchi, S. Oishi, K. Honda, Y. Kondoh, T. Saito, T. Kubo, M. Kaneda, H. Ohno, H. Osada and N. Fujii, *Bioorg. Med. Chem. Lett.*, 2013, **23**, 3802-3805.
33. L. Gu, H. Zhang, T. Liu, S. Zhou, Y. Du, J. Xiong, S. Yi, C.-K. Qu, H. Fu and M. Zhou, *Cancer Cell*, 2016, **30**, 623-636.
34. D. J. Foley, A. Nelson and S. P. Marsden, *Angew. Chem. Int. Ed.*, 2016, **55**, 13650-13657.
35. D. Reed, Y. Shen, A. A. Shelat, L. A. Arnold, A. M. Ferreira, F. Zhu, N. Mills, D. C. Smithson, C. A. Regni, D. Bashford, S. A. Cicero, B. A. Schulman, A. G. Jochemsen, R. K. Guy and M. A. Dyer, *J. Bio. Chem.*, 2010, **285**, 10786-10796.
36. M. Wade, Y. C. Li and G. M. Wahl, *Nat. Rev. Cancer*, 2013, **13**, 83.
37. M. Bista, D. Smithson, A. Pecak, G. Salinas, K. Pustelny, J. Min, A. Pirog, K. Finch, M. Zdzalik, B. Waddell, B. Wladyka, S. Kedracka-Krok, M. A. Dyer, G. Dubin and R. K. Guy, *PLOS One*, 2012, **7**, e37518.