

Assistive technology and telecare; Assessment and practice for people with dementia. (Forsyth)

**Title:** Assessment of need and practice for assistive technology and telecare for people with dementia - the ATTILA (Assistive Technology and Telecare to maintain Independent Living At home for people with dementia) Trial.

Kirsty Forsyth<sup>1\*</sup>, Catherine Henderson<sup>2</sup>, Lucy Davis<sup>3</sup>, Anusua Singh Roy<sup>1</sup>, Barbara Dunk<sup>4</sup>, Eleanor Curnow<sup>1</sup>, Rebecca Gathercole<sup>5</sup>, Natalie Lam<sup>3</sup>, Emma Harper<sup>3</sup>, Iracema Leroi<sup>6</sup>, John Woolham<sup>7</sup>, Chris Fox<sup>8</sup>, John O'Brien<sup>9</sup>, Andrew Bateman<sup>10</sup>, Fiona Poland<sup>11</sup>, Peter Bentham<sup>12</sup>, Alistair Burns<sup>6</sup>, Anna Davis<sup>13</sup>, Richard Gray<sup>3</sup>, Rosie Bradley<sup>3</sup>, Martin Knapp<sup>5</sup>, Stanton Newman<sup>13</sup>, Rupert McShane<sup>14</sup>, Craig Ritchie<sup>15</sup>, Emma Talbot<sup>16</sup>, Emma Hooper<sup>17</sup>, Rachel Winson<sup>18</sup>, Bethany Scutt<sup>5</sup>, Victoria Ordonez<sup>18</sup>, Samantha Nunn<sup>18</sup>, Grace Lavelle<sup>5</sup>, and Robert Howard<sup>19</sup>

<sup>1</sup> School of Health Sciences, Queen Margaret University, Edinburgh. EH21 6UU. UK

<sup>2</sup>Personal Social Services Research Unit, London School of Economics and Political Science, Houghton Street, London. WC2A 2AE. UK

<sup>3</sup> Clinical Trial Service Unit, Richard Doll Building, Old Road Campus, Roosevelt Drive, Oxford. OX3 7LF. UK.

<sup>4</sup>South London and Maudsley NHS Foundation Trust, 115 Denmark Hill, The Maudsley SE5 8AZ. UK.

<sup>5</sup>Department of Old Age Psychiatry, Institute of Psychiatry, Box 070, De Crespigny Park, London SE5 8AF. UK

<sup>6</sup>School of Community Based Medicine, University of Manchester, Jean McFarlane Building, 3<sup>rd</sup> Floor, Oxford Road, Manchester. M13 9PL. UK.

<sup>7</sup> Social Care Workforce Research Unit, King's College London, 22 Kingsway, London. WC2B 6LE. UK.

<sup>8</sup>School of Medicine, Health Policy and Practice, University of East Anglia, Norwich, Norfolk. NR4 7TJ. UK.

<sup>9</sup>Department of Psychiatry, University of Cambridge, Level E4, Box 189, Addenbrooke's Hospital, Hills Road, Cambridge. CB2 0QQ. UK.

<sup>10</sup>Oliver Zangwill Centre for Neuropsychological Rehabilitation, Princess of Wales Hospital, Ely. CB6 1DN. UK.

<sup>11</sup>School of Allied Health Professionals, University of East Anglia, Norwich, Norfolk. NR7 4TJ. UK.

<sup>12</sup> The Barberry Centre, 25 Vincent Drive, Birmingham. B15 2FG. UK.

<sup>13</sup>School of Community and Health Science, City University, London. EC1V 0BH. UK.

<sup>14</sup>Oxford Health NHS Foundation Trust, Warneford Hospital, Old Road, Headington, Oxford. OX3 7JX. UK.

<sup>15</sup> Centre for Clinical Brain Sciences, University of Edinburgh, Chancellor's Building, 49 Little France Crescent, Edinburgh. EH16 4SB. UK.

<sup>16</sup> Norfolk and Suffolk NHS Foundation Trust, Room S12. Haymills House, Station Road East, Stowmarket, Suffolk IP14 1RF. UK.

Assistive technology and telecare; Assessment and practice for people with dementia. (Forsyth)

<sup>17</sup> Lantern Centre, Vicarage Lane, Fulwood, Preston PR2 8DW. UK

<sup>18</sup> Cambridgeshire Community Services NHS Trust, Oliver Zangwill Centre, Ely CB61DN. UK

<sup>19</sup> University College London, 6<sup>th</sup> Floor Maple House, 149 Tottenham Court Road, London W1T 7NF.  
UK

\*Corresponding author: Tel. +441314740000. Email address: kforsyth@qmu.ac.uk

## **Abstract**

**INTRODUCTION:** To define current assistive technology and telecare (ATT<sup>1</sup>) practice for people with dementia living at home.

**METHODS:** Randomised controlled trial (N=495) of ATT assessment and ATT installation intervention, compared to control (restricted ATT package). ATT assessment and installation data were collected. Qualitative work identified value networks delivering ATT; established an ATT assessment standard.

**RESULTS:** ATT was delivered by public and not-for-profit telecare networks. ATT assessments showed 52% fidelity to the ATT assessment standard. Areas of assessment most frequently leading to identifying ATT need were daily activities (93%), memory (89%) and problem solving (83%). ATT needs and recommendations were weakly correlated ( $\tau = 0.242$ ;  $P < .000$ ), with ATT recommendations and installations moderately correlated ( $\tau = -0.470$ ;  $P < .000$ ). Half (53%) of recommended technology was not installed. Safety concerns motivated 38% of installations.

**DISCUSSION:** Assessment recommendations were routinely disregarded at the point of installation. ATT was commonly recommended for safety and seldom for supporting leisure.

**Keywords:** Dementia, assistive technology, telecare, ATTILA, assessment of need, community-dwelling.

---

<sup>1</sup> Abbreviations: ATT – Assistive Technology and Telecare, ATTILA – Assistive Technology and Telecare to maintain Independent Living At home for people with dementia, TIDieR – Template for Intervention Description and Replication, SMMSE – Standardised Mini-Mental State Examination, CASSR – Councils with Adult Social Service Responsibilities, NFP- Not for Profit

## **1. Background**

Approximately 46.8 million people are living with dementia worldwide [1]. As the disease progresses, it can be a challenge for people with dementia to live safely in their own homes [2]. Assistive technology and telecare (ATT) offer a means of managing the risks facing older people with dementia who wish to remain living independently at home. The first use of electronic ATT in the UK was to provide support for people with dementia and their carers [3]. Within a decade, interest in ATT prompted the development of a Department of Health strategy [4] and, increasingly, ATT moved into the mainstream. However, as interest in ATT has increased, the specific focus on its application for those living with dementia has diminished [5]. Early studies of ATT for dementia highlighted the importance of assessment [6] to construct a sufficiently detailed picture of an individual's life to enable assistive technology to meet needs and maximise good outcomes [7, 8]. However, to date, there is scant evidence on the effectiveness of usual ATT provision and associated practices for people with dementia. The ATTILA (Assistive Technology and Telecare to maintain Independent Living At Home for People with Dementia) trial was a pragmatic randomised controlled trial, comparing outcomes for people with dementia receiving ATT and those receiving equivalent community services but not ATT [9].

### **1.1 Aim**

A detailed exploration of the intervention under investigation is needed to give insight into the fidelity of the intervention and to allow for replication [10]. We aimed to provide an investigation of routine ATT practice and the systems in place to deliver ATT for people with dementia.

## **2. Method**

We adhered to the Template for Intervention Description and Replication (TIDieR) [11] in describing the components of the ATTILA intervention, in terms of what happened, who was involved, how, where and when did the intervention happen, how much was provided and whether it was tailored to participants.

### **2.1 Assistive Technology and Telecare (ATT)**

The ATT intervention was defined for the purposes of the ATTILA Trial as a two-stage process:

1. an ATT assessment, with subsequent ATT recommendation(s),
2. the installation of ATT devices alongside monitoring services as appropriate.

ATT devices included simple, battery operated, stand-alone technologies and/or telecare i.e., a range of devices and sensors, which communicate and relay messages to an external call centre where an appropriate response is arranged. Participants were randomised to receive an ATT assessment followed by either installation of ATT as indicated by the assessment (intervention arm), or an ATT package

Assistive technology and telecare; Assessment and practice for people with dementia. (Forsyth)

restricted to only non-monitored smoke & carbon monoxide detectors & pendant alarm (control arm) [9].

## 2.2 Participants

People with either a diagnosis of dementia (early or late onset) or evidence of cognitive difficulties suggesting presence of dementia were eligible for ATTILA. Those lacking capacity could participate with the consent of a personal consultee. Additionally, participants had to meet eligibility criteria for access to social care services, live in an ordinary community dwelling (including sheltered accommodation) and have a working telephone line at home. Exclusions from the study included having already received ATT (excluding smoke, carbon monoxide or pendant alarms); having previously received ATT and not used it; or having a perceived urgent need for a package of care [9].

## 2.3 ATT Delivery Systems

To describe the delivery systems for ATT deployment, interviews were conducted by CH with key informants from local authority operational/commissioning teams and telecare monitoring centre managers in the seven sites from which the majority of trial participants were recruited (N=484). Invitations were sent to 21 potential key informants, resulting in 14 interviews covering six sites (no key informants were available for interview in one site) between June and September 2016. Interviews were not recorded but written notes were taken; interviewees were also asked for supporting documentation that might help to understand the policies and procedures in relation to ATT deployment. Data were also collected on ATT assessment and delivery processes via pro-formas completed by local researchers in 2015 and via a follow-up desk-based search in 2017. Data were examined using NVIVO qualitative data analysis software (Version 11) by CH. Data were first structured into five production stages within a Framework Analysis [12]: assessment, equipment procurement/ordering, installation, call monitoring and response to sensor activations. To identify commonalities in local systems for delivering ATT to trial participants, we took an approach based on value network role analysis [13, 14]. Production inputs and processes observed in each site were mapped onto value network frameworks.

## 2.4 Baseline Participant Characteristics

Local trial researchers collected data on characteristics of participants (age, gender, ethnic background, marital status, living situation (living with another, living alone) and carers (ethnic background, marital status, carer). They completed Standardised Mini-Mental State Examinations (SMMSE) [15] with participants, and rated participants' risk of wandering (low, moderate, high), and safety risk within the home (low, moderate, high).

## 2.5 Local baseline ATT assessments/recommendations

We assumed that Social Services Departments in each ATTILA site had distilled local and national guidelines on best practice in ATT assessment when constructing local assessment templates. To establish a practice standard for ATT assessments in the ATTILA sites, ATT assessment templates and guidance were sourced from each site between August 2013 and August 2016. Sites were asked to resend documentation if there were changes during the lifetime of the study; as a result, two sets of new documentation were submitted. Framework analysis [12] to identify common assessment themes across sites was applied to this documentation (by KF), using the Model of Human Occupation Screening Tool (MOHOST) [16]. The Model of Human Occupation Screening Tool is designed to detail people's values, insight, interests, routines, communication, cognitive and physical skills, and physical and social environment to gain a detailed picture of an individual's life. The resultant ATT assessment standard consisted of a set of 14 ATT assessment areas (see supplementary materials 1). A four point scale was developed for each assessment area within the ATT assessment standard, where 4=no risk when doing daily activity, 3=mostly risk free when doing daily activity, 2= some risk when doing daily activity, and 1=significant multiple risks when doing daily activity. Specific definitions were developed for rating each assessment area (see supplementary materials 2). ATT needs were identified when an assessment area received a rating of 1 (significant multiple risks when doing daily activity) or 2 (some risk when doing daily activity).

Locally completed ATT assessments for each participant were reviewed against the ATT assessment standard to assess whether these addressed the ATT assessment areas identified by the templates across ATTILA sites. Fidelity to this standard was determined by two trial practitioners with experience in dementia care and ATT assessment (KF, EC), who independently classified the content of each locally completed ATT assessment against the ATT assessment standard and assigned risk-ratings. They then reviewed ratings together and resolved discrepancies.

There is no recognised taxonomy of ATT for people with dementia, therefore, a taxonomy was developed in collaboration with Trent Dementia Services Development Centre and the "atdementia" initiative ([www.atdementia.org.uk](http://www.atdementia.org.uk)), an independent online ATT resource. This taxonomy was then developed into two identical Technology Checklist forms (one for recommended ATT and one for installed ATT) which covered the following ATT functions: (a) reminder or prompting devices; (b) devices to support safety; (c) safer walking technologies; (d) communication devices and (e) devices that support meaningful use of leisure time; (f) monitoring and response information. The form also recorded data about which type of assessor had assessed for ATT (ATT assessor, health or social care professional, other), method of assessment (in-person at-home, in-person not-at-home, telephone assessment, using case notes, other), whether ATT was monitored (yes/no), who would respond to ATT alerts (direct to responder or via a call centre). Two trial practitioners with experience in dementia care

Assistive technology and telecare; Assessment and practice for people with dementia. (Forsyth)

and ATT assessment (KF, BD) collaboratively classified each device recommended within the locally completed ATT needs assessment using the technology checklist (for recommended ATT).

## 2.6 Trial Instruments (week 12-104)

Local trial researchers administered the Technology Checklist (for installed ATT) during home visits at weeks 12, 24, 52 and 104.

## 2.7 Statistical analyses

Categorical data were summarised in percentages and numbers of observations. Correlations between count variables were tested using non-parametric methods (Kendall Tau Rank Correlation Coefficient  $\tau$ ). The Kruskal-Wallis test was used to assess if there were statistically significant differences between multiple groups for outcomes. Friedman's test was used to determine significance of change over time in the count variables. In the case of categorical variables, differences between observed and expected frequencies were tested using Pearson's chi square test for independence, or alternatively Fisher's exact test, when the assumption of minimum expected cell count in contingency tables was not met [17]. SMMSE scores were categorised into stages of dementia [18] for the purposes of analysis (30= no dementia, 26-29= questionable dementia, 21-25= mild dementia, 11-20= moderate dementia, 0-10=severe dementia). Effective tailoring of the intervention was described through the strength of the correlation [19] between ATT needs and ATT recommendations at baseline and between ATT recommendations and ATT installation by 24 weeks. We also compared the ATT recommended in the baseline assessment with subsequent installations for each participant in the intervention arm up to 24 weeks. Any installation after 24 weeks was considered unrelated to the baseline ATT assessment.

## 2.8 Ethical approval

The ATTILA trial was approved by the NHS Health Research Authority National Research Ethics Committee (REC reference number 12/LO/186) and is registered with the ISRCTN (<http://www.controlled-trials.com/ISRCTN86537017>).

## 3. Results

In total, 495 people were randomised into the ATTILA trial between August 2013 and October 2016 (247 control, 248 intervention). The majority of participants were >80 years old, female, widowed, white British, did not live alone, had moderate dementia, were at low risk for wandering and of being considered a safety risk within the home. Most caregivers were aged over 65 years, married and spouses or daughters of participants (Table 1). Of the 451 documented baseline ATT assessments available, 413 contained an ATT recommendation. Of the 248 participants recruited to the intervention arm, data from 209 participants was available for analysis of ATT installations (Fig. 1).

### 3.1 Value networks:

Assistive technology and telecare; Assessment and practice for people with dementia. (Forsyth)

Networks delivering services (offering value) to ATTILA participants were classified into two types (Supplementary Materials 3a and 3b). Firstly, ‘public telecare provider networks’ (N=4), where two assessor roles were identified: the ATT assessor, and the authorised (or trusted) assessor (Health or Social Care Professional). ATT assessors were employed by public agencies (NHS or Councils with Adults Social Services Responsibilities - CASSR); their primary role was to assess for a full range of ATT devices (‘networked’ – monitored by a telecare call centre or unpaid carer; or ‘standalone’). Authorised assessors could offer first-generation telecare (pendant-only systems) or straightforward ATT (for instance, adding on an additional sensor or providing a memo minder) depending on their level of experience and local permissions; they performed ATT assessment as a secondary role. In these networks, most or all of the ATT infrastructure for procurement, installation, stock control and maintenance of ATT devices fell to units within the CASSR. Secondly, there were ‘not-for-profit provider networks’ (NFP) (N=3). Three assessor roles were identified across these ‘NFP telecare networks’. Telecare assessors working for NFP telecare providers assessed for AT that was networked to providers’ call-monitoring centres. Assessment for standalone AT fell to assessors within the CASSR. A ‘social care ATT assessor’ role was also identified: these assessors could assess for ATT (networked/standalone) and work with a choice of suppliers to procure and arrange installation of ATT devices. Private companies offered combinations of procurement and stock control, installation and maintenance services to the NFP telecare providers.

### 3.2 Locally completed ATT assessments/recommendations at baseline:

The local ATT assessment fidelity with the ATT assessment standard was 52% (7.2 assessment areas were addressed per assessment) (Table 2). Of ATT assessments reviewed, 99/451 (22%) addressed 0-2 areas of assessment. There was higher fidelity to assessment areas relating to “mobility” (74%), “social support” (72%), “daily activity” (71%), and “memory” (71%). Fidelity varied across sites: the mean number of assessment areas addressed ranged from 2-13 per site ( $P<.000$ ) with public telecare providers addressing more assessment areas than not-for-profit telecare providers ( $P<.026$ ). HSC professionals addressed more assessment areas than ATT assessors ( $P<.046$ ). Fidelity varied across assessment methods ( $P<.000$ ), with the in-person at-home assessment method addressing more assessment areas than in-person but not-at-home ( $P=.003$ ), telephone assessment ( $p<.000$ ), and case notes methods ( $P=.003$ ). Women had more assessment areas addressed than men ( $P=.027$ ). More assessment areas were addressed for participants at medium risk of wandering than for participants at low risk of wandering ( $P=.028$ ).

In total, 60% of assessment responses identified an ATT need, with 4.4 ATT needs (range 0-12) identified per participant (Table 2). The mean number of ATT needs identified varied, ranging from 2-6 per site ( $P<.000$ ). Areas of concern most frequently identified as triggering the need for ATT were:

daily activities (93%), memory (89%), and problem solving (83%). HSC professionals identified more ATT needs than did ATT assessors ( $P=.047$ ). More ATT needs were identified by in-person-home than by telephone assessment methods ( $P<.000$ ). There was no significant difference between ATT needs in men and women ( $P=.337$ ). The number of ATT needs identified for each participant differed depending on the levels of wandering risk ( $P=.005$ ), with medium risk of wandering associated with more ATT needs than in the case of low risk of wandering ( $P=.016$ ). ATT needs varied by category of SMMSE score ( $P<.000$ ): participants with severe dementia had more ATT needs than those with mild ( $P<.000$ ), moderate ( $P=.002$ ) and questionable dementia ( $P<.000$ ).

A documented ATT recommendation was given for 413 participants with 1090 ATT devices recommended at baseline, with a mean of three devices (range 1-14 devices). One or two ATT devices were recommended for 57% (235) of participants. The correlation between the ATT needs and ATT recommendations identified in local ATT assessments was weak ( $\tau=0.242$ ;  $P<.000$ ). Most recommendations were for safety-related devices (59%; 644/1090), followed by reminder/prompting devices (25%; 269/1090). Sixty-two percent (673/1090) of recommended ATT devices required monitoring; and 67% (353/526) of monitored devices with an identified responder required a formal (call centre) response.

### 3.3 Recommendations and Installations at follow up (intervention arm only):

By 24 weeks, a mean of 3.5 devices were recommended for participants in the intervention arm. Of the ATT devices recommended, 53% (306/572) were not installed. However, 62% (438/704) of the ATT devices installed had not been recommended in the baseline assessment (Table 3). There was a moderate negative correlation between number of recommendations and number of installations per participant per ATT category ( $\tau =-0.470$ ; both  $P<.000$ ).

### 3.4 Week 12-104, ATT installed (intervention arm only):

By week 104, there were 888 ATT devices installed for 209 participants in the intervention arm, a mean of 4.2 devices per participant (range 1-15). Of devices installed for intervention participants (Table 4) 42% (374/888) of those involved the types of technology provided to control arm participants (e.g., non-monitored smoke detectors). Installations decreased over time ( $P<.031$ ) with 79% (704/888) of ATT installed by week 24. Intervention participants' ATT devices were most frequently installed for safety reasons (38%) or for reminder/prompting (18%). ATT assessors were most frequently identified as having assessed for the installed devices (32%), followed by HSC professionals (20%) but 40% of assessors' backgrounds were unknown. While 41% of installations followed an in-person home visit (41%), in many cases the participants could not report the method of assessment (42%). Nearly half

(47%) of the ATT devices installed required monitoring; 38% of monitored devices were networked to a call centre (so that any alerts would receive an initial response from paid services).

Results have been summarised using the TIDieR framework in Table 5.

#### **4. Discussion**

Providing ATT for people with dementia is a complex intervention [10] which includes ATT assessment, recommendations, installation, monitoring and response. Above all, a comprehensive assessment is critical to tailoring ATT to individual users' needs [6, 7]. Limiting the amount of assessment information available may reduce a provider's ability to personalise the technology to individual circumstances. Effective tailoring is needed if the technology is to be used and integrated into the person's life [7, 8]. Examining the ATTILA sites' local ATT assessment documentation, we found 14 common assessment areas. An ATT assessment covering all 14 areas would provide a range of information on the person's values, insight, interests, routines, communication, cognitive and physical skills, and physical and social environment. However, we found that only half (52%) of these assessment areas were documented by assessors, with 60% of responses indicating an ATT need. One explanation for limited coverage of the 14 assessment areas in many assessments could be that assessors did address all the assessment areas but only documented the areas of greatest concern. On the other hand, confidence that assessment processes were comprehensive will be tempered by the finding that 22% of participants' assessments consisted of 0-2 assessment areas addressed (typically in the form of one-line emails). Local authorities facing severe budgetary pressures may be tempted to take short-cuts and allow ATT to be installed without a full assessment [7]; our research provides some evidence that this did occur. Furthermore, local ATT assessments tended to be narrowly focussed on certain areas of the assessment, particularly the area of safety.

HSC professionals carried out the bulk of baseline assessments for ATTILA participants. However, more installations were at the request of ATT assessors than HSC professionals across the 12-to-104 week follow-ups. The title 'ATT assessor' suggests a worker with specialist skills; however ATT assessors within our study were from a variety of backgrounds, with and without professional qualifications, including personnel with wider job remits than ATT and personnel working solely in ATT teams. ATT assessors demonstrated lower fidelity to the ATT assessment standard than HSC professionals. A recent study [20] suggested that ATT training within CASSRS for people holding responsibility for ATT deployment was brief, 'product based' and seen as a marketing opportunity by telecare suppliers. This in turn may limit the scope of the ATT assessment to only those devices available from a particular telecare supplier. Value network mapping yielded two groupings of ATTILA sites, based on the sector of telecare provider offering value to ATTILA participants. The

scope of assessors' roles (e.g. their remit to assess for a narrower or broader range of ATT) appeared to differ between network types. It seemed that the organisational resources available to assessors to deliver value to users of ATT differed between sectors. The contractual arrangements existing between CASSRs and other network actors and the extent to which they had contracted out ATT services may also have influenced the resources available to assessors, and thereby the scope of their roles. These must remain tentative findings, given the limitations in information available from a relatively small number of key informants.

The number of ATT needs identified were similar to the number of ATT installations (4.4 and 4.2 per participant, respectively), which on the surface might suggest that assessment recommendations generally translated into installations. Yet there were mismatches: devices were installed for needs that had not been identified in the assessment process, while half of the devices recommended as a result of assessment were not installed. Other ATT research suggests some reasons for these results: that difficulties identifying the resources to progress installation may have delayed or deterred installation [21, 22], that individuals may decline in cognitive capacity while awaiting installation so that the ATT is no longer appropriate [23], and that technology may be heavily promoted by suppliers, leading to over-prescribing of ATT [24].

Local authorities have identified the most important goals of ATT as being to delay the need for care and support, and to enhance quality of life for people with care and support needs [20]. This study found that ATT was most often recommended and installed for safety reasons. Indeed, it has been argued that the use of ATT for safety reasons could delay care by managing identified risks. ATT designed to increase quality of life through meaningful use of leisure time [9, 25-28] was rarely recommended or installed during the trial.

## Summary

The literature has argued for the importance of assessment in enabling assistive technology to maximise good outcomes, however, this trial identified assessment recommendations were routinely disregarded at the point of installation. It is, therefore, unlikely that assistive technology is being deployed in a way that will result in benefits for recipients. There is a subsequent need, therefore, to understand how to more effectively translate ATT needs into ATT recommendations and effective ATT installations. Two different delivery systems for deploying ATT were also identified which seemed to have different organisational resources available to assessors and clarity should be sought as to how this impacts on outcomes. Finally, ATT wasn't deployed to support meaningful use of leisure time which is inconsistent with local authority goals of enhancing quality of life and a review of barriers to deployment is needed.

Assistive technology and telecare; Assessment and practice for people with dementia. (Forsyth)

### **Acknowledgements**

Funding for the trial came from the NIHR Health Technology Assessment programme (HTA 10/50/02). Neither the funders nor the sponsor have been involved in the preparation or publication of this article. PB received financial support from TauRx Therapeutics. JOB received financial support from TauRx, GE Healthcare, Avid/Lilly and Eisai.

### **Supplementary Data**

# Assistive technology and telecare; Assessment and practice for people with dementia. (Forsyth)

## References

- [1] Prince M, Wimo A, Guerchet M, Ali G, Wu Y, Prina M. World Alzheimer Report 2015: The Global Impact of Dementia: An Analysis of Prevalence, Incidence, Costs and Trends. London: Alzheimer's Disease International; 2015 August 2015.
- [2] Luppia M, Luck T, Brahler E, Konig H, Riedel-Heller S. Prediction of Institutionalisation in Dementia: A Systematic Review. *Dement and Geriatr Cognitive Disorder*. 2008;26:65-78.
- [3] Marshall M, editor. *Astrid: a social and technological response to meeting the needs of individuals with dementia and their carers; a guide to using technology within dementia care*. UK; London: Hawker; 2000.
- [4] Department of Health. *Living well with dementia: A national dementia strategy*. Leeds, UK: Department of Health, UK Government; 2009 3 February 2009.
- [5] Department of Health, Older People and Disability Division. *Building Telecare in England*. UK; London: Department of Health; 2005 19 July 2005.
- [6] Bjerneby S, Topo P, Holthe T. *Technology, ethics and dementia: a guidebook on how to apply technology in dementia care*. Norwegian Centre for Dementia Care; 1999.
- [7] Sugarhood P, Wherton J, Procter R, Hinder S, Greenhalgh T. Technology as system innovation: A key informant interview study of the application of the diffusion of innovation model to telecare. *Disability and Rehabilitation: Assistive Technology*. 2014;9(1):79-87.
- [8] Greenhalgh T, Procter R, Wherton J, Sugarhood P, Hinder S, Rouncefield M. What is quality in assisted living technology? The ARCHIE framework for effective telehealth and telecare services. *BMC Med*. 2015;13:91.
- [9] Leroi I, Woolham J, Gathercole R, Howard R, Dunk B, Fox C, et al. Does Telecare Prolong Community Living in Dementia? A Study Protocol for a Pragmatic Randomised Controlled Trial. *Trials*. 2013;14:349.
- [10] Craig P, Dieppe P, Macintyre S, Michie S, Nazareth I, Petticrew M, on behalf of the Medical Research Council. Developing and evaluating complex interventions: Following considerable development in the field since 2006, MRS and NIHR have jointly commissioned an update of this guidance to be published in 2019. UK: Medical Research Council; 2008.
- [11] Hoffman T, Glasziou P, Boutron I, Milne R, Perera R, Moher D, et al. Better reporting of interventions: template for intervention description and replication (TIDieR) checklist and guide. *BMJ*. 2014 7 March 2014;348:1-12.
- [12] Ritchie J, Lewis J, editors. *Qualitative Research Practice: A Guide for Social Science Students and Researchers*. London: Sage Publications; 2003.
- [13] Ehrenhard M, Kijl B, Nieuwenhuis L. Market adoption barriers of multi-stakeholder technology: smart homes for the aging population. *Technological Forecasting and Soc Change*. 2014;89:306-15.
- [14] Kijl B, Nieuwenhuis L, Veld H, Vollenbroek-Hutten M, Hermens H. Deployment of e-health services - a business model engineering strategy. *J Telemed and Telecare*. 2010;16(6):344-53.
- [15] Molloy D, Standish T. A guide to the standardized Mini-Mental State Examination. *Int Psychogeriatr*. 1997;9(suppl 1):143-50.
- [16] Parkinson S, Forsyth K, Kielhofner G. *User's manual for the Model of Human Occupation Screening Tool (MOHOST)*. University of Illinois, Chicago: University of Illinois; 2002.
- [17] Siegel S. *Nonparametric statistics for the behavioral sciences*, New York, 1956. Google Scholar. 1956.
- [18] Pernecky R, Wagenpfeil S, Komossa K, Grimmer T, Diehl J, Kurz A. Regular Research Articles: Mapping Scores Onto Stages: Mini-Mental State Examination and Clinical Dementia Rating. *The American J of Geriatr Psychiatry*. 2006;14:139-44.
- [19] Schober P, Boer C, Schwart LA. Correlation coefficients: Appropriate use and interpretation. *Anesthesia & Analgesia*. 2018;126(5):1763-8.

## Assistive technology and telecare; Assessment and practice for people with dementia. (Forsyth)

[20] Woolham JG, Steils N, Fisk M, Porteus J, Forsyth K. The Utopia Project. Using Telecare for older people in adult social care: The findings of a 2016-2017 national survey of local authority telecare provision for older people in England. King's College, London, UK.: Social Care Workforce Research Unit, Kings College, London; 2018.

[21] Gibson G, Dickinson C, Brittain K, Robinson L. The everyday use of assistive technology by people with dementia and their family carers: a qualitative study. *BMC Geriatr*. 2015 07/24;15:89-.

[22] Gibson G, Newton L, Pritchard G, Finch T, Brittain K, Robinson L. The provision of assistive technology products and services for people with dementia in the United Kingdom. *Dementia (London)*. 2016 07;15(4):681-701.

[23] Canevelli M, Kelaiditi E, del Campo N, Bruno G, Vellas B, Cesari M. Predicting the rate of cognitive decline in Alzheimer's Disease: Data from the ICTUS study. *Alz Dis and Assoc Disorders*. 2016;30(3):237-42.

[24] Milligan C, Roberts C, Mort M. Telecare and older people: Who cares where? *Soc Sci Med*. 2011;72:347-54.

[25] Lund M, Christensen H, Caltenco H, Lontis E, Bentsen B, Struijk L. Inductive tongue-control of powered wheelchairs. 2010 Annual International Conference of the IEEE Engineering in Medicine and Biology; 31 Aug - 4 Sept 2010; Buenos Aires, Argentina. <https://doi.org/10.1109/IEMBS.2010.5627923>: IEEE; 2010.

[26] Windle K, Francis J, Coomber C. Preventing loneliness and social isolation: interventions and outcomes. Research Briefing. UK; London: Social Care Institute for Excellence; 2011 October 2011.

[27] Steptoe A, Shankar A, Demakakos P, Wardle J. Social isolation, loneliness, and all-cause mortality in older men and women. *PNAS [Internet]*. 2013 9 April 2013 [cited 10/10/2018];110(15):10/10/2018. Available from: <https://doi.org/10.1073/pnas.1219686110>.

[28] Pols J. The heart of the matter: About good nursing and telecare. *Health Care Analysis*. 2010;18(4):374-88.

-

Forsyth

Table 1 Participants' baseline characteristics

Characteristic		Participants N (%)
Age	<65 years	15/495 (3%)
	65-80 years	182/495 (37%)
	>80 years	298/495 (60%)
Gender	Male	205/495 (41%)
	Female	290/495 (59%)
Stage of Dementia (SMMSE)	Severe	59/495 (12%)
	Moderate	179/495 (36%)
	Mild	126/495 (25%)
	Questionable	51/495 (10%)
	No	0/495 (0%)
	Incomplete responses	80/495 (16%)
Marital Status of the Participant	Single (never married)	25/495 (5%)
	Married	191/495 (39%)
	Co-habiting	8/495 (2%)
	Civil Partnership	0/495 (0%)
	Separated	5/495 (1%)
	Divorced	22/495 (4%)
	Widowed	196/495 (40%)
	Blank	17/495 (3%)
Ethnic Background of the Participant	White- British	357/495 (72%)
	White- Irish	10/495 (2%)
	Other White Background	13/495 (3%)
	Black or Black British – Caribbean	45/495 (9%)
	Black or Black British – African	10/495 (2%)
	Other Black Background	0/495 (0%)
	Asian or Asian British – Indian	9/495 (2%)
	Asian or Asian British – Pakistani	0/495 (0%)
	Asian or Asian British – Bangladeshi	0/495 (0%)
	Chinese	1/495 (0%)
	Other Asian Background	4/495 (1%)
	Mixed – White and Black Caribbean	0/495 (0%)
	Mixed – White and Black African	0/495 (0%)
	Mixed – White and Asian	1/495 (0%)
	Other – Mixed Background	4/495 (1%)
	Other Ethnic Background	6/495 (1%)
	Blank	4/495 (1%)
Living Situation	Living with other	267/495 (54%)
	Living alone	228/495 (46%)
Risk of Wandering	Low	358/495 (72%)
	Moderate	100/495 (20%)
	High	37/495 (7%)
Safety Risk within the Home	Low	249/495 (50%)
	Moderate	205/495 (41%)
	High	41/495 (8%)
Caregiver age	<65 years	223/495 (45%)
	65-80 years	113/495 (23%)
	80+ years	48/495 (10%)
	Blank	111/495 (22%)
Caregiver Details	Live-in Caregiver	240/495 (48%)
	Caregiver visits at least once/day	122/495 (25%)
	Caregiver visits less than once/day	133/495 (27%)
Marital Status of the Caregiver	Single (never married)	59/495 (12%)
	Married	309/495 (62%)
	Co-habiting	25/495 (5%)

Forsyth

	Civil Partnership	0/495 (0%)
	Separated	4/495 (1%)
	Divorced	35/495 (7%)
	Widowed	10/495 (2%)
	Blank	22/495 (4%)
<b>Relationship of Caregiver to Participant</b>	Wife/husband	160/495 (32%)
	Partner	10/495 (2%)
	Daughter	153/495 (31%)
	Son	84/495 (17%)
	Daughter in law	10/495 (2%)
	Son in law	1/495 (0%)
	Sister	2/495 (0%)
	Brother	2/495 (0%)
	Other relative	26/495 (5%)
	Neighbour	5/495 (1%)
	Friend	7/495 (1%)
	Other	4/495 (1%)

## Forsyth

Table 2: Fidelity with ATT assessment standard and identified ATT needs

Sites ATT assessment areas/standard	Fidelity with ATT assessments standard			ATT needs (i.e., responses rated as some risk or significant multiple risk)		
1. Insight	241/451	53%		151/241	63%	
2. Values	245/451	54%		100/245	41%	
3. Wandering/Disorientation	284/451	63%		219/284	77%	
4. Daily activity	321/451	71%		298/321	93%	
5. Conversation	226/451	50%		100/226	44%	
6. Express Needs	175/451	39%		24/175	14%	
7. Memory	320/451	71%		284/320	89%	
8. Problem Solve	218/451	48%		181/218	83%	
9. Mobility	335/451	74%		224/335	67%	
10. Grip/Dexterity	147/451	33%		18/147	12%	
11. Space	140/451	31%		47/140	34%	
12. Resources	128/451	28%		26/128	20%	
13. Social Support	325/451	72%		183/325	56%	
14. The way the activity completed	162/451	36%		118/162	73%	
Total responses	3,267/6,314	52%		1973/3,267	60%	
	Fidelity with ATT assessments standard			Number ATT needs		
<b>Participant Characteristics</b>						
Gender	Median	Mean	%	Median	Mean	%
Female	8	7.67	62%	4	4.46	60%
Male	5	6.65	38%	3	4.25	40%
	P=.027			P.337		
Risk of wandering	Median	Mean	%	Median	Mean	%
Low	7	6.93	70%	4	4.10	68%
Medium	9	8.37	23%	5	5.04	23%
High	6	7.24	7%	4	5.24	9%
	P=.038			P=.005		
SMMSE score (18)	Median	Mean	%	Median	Mean	%
Questionable dementia (26-29)	7	7.22	13%	3	3.38	10%
Mild dementia (21-25)	7	6.6	27%	4	3.9	27%
Moderate dementia (11-20)	7	7.38	45%	4	4.27	44%
Severe dementia (0-10)	8.5	7.96	15%	5.5	5.79	19%
	P=.309			P<.000		
<b>Assessment Characteristics</b>						
Assessors	Median	Mean	%	Median	Mean	%
Health & Social Care Professionals	8	7.85	68%	4	4.66	67%
ATT assessor	5.5	6.51	29%	3	3.86	29%
	P=.051			P=.028		
Assessment method	Median	Mean	%	Median	Mean	%
In person at home	10	9.14	85%	5	5.06	82%
In person not at home	5	6.43	8%	3	3.38	8%
Telephone	2	3.42	6%	2	2.71	9%
Case notes	3	3.33	1%	2.5	3	1%
	P<.000			P<.000		
<b>Service structure</b>						
	Median	Mean	%	Median	Mean	%
Public telecare provider	7	7.59	73%	4	4.41	70%
Not-for-profit telecare provider	6	6.41	25%	4	4.31	28%
	P=.026			P=1.00		
	Mean fidelity with ATT assessment standard per participant			Mean number of responses per participant rated as an ATT need		
	7.2 assessment areas addressed (0-13)			4.4 ATT needs (0-12)		

Forsyth

Forsyth

Table 3: Recommended ATT devices matched to ATT devices installed at 24 weeks (Intervention arm only)

ATT Technology Checklist	ATT Recommended, N (%)	ATT Recommended + Installed at 24 wks. N (%)	ATT Recommended + Not Installed, N (%)*	ATT Installed at 24 wks. N (%)	ATT Not Recommended + Installed N (%)
<b>Control Group Technology</b>					
Pendant alarm	44/572 (8%)	22/44 (50%)	22/44 (50%)	89/704 (13%)	67/89 (75%)
Non-monitored smoke detector	0/572 (0%)	0(0%)	0(0%)	68/704 (10%)	68/68 (100%)
Non-monitored carbon monoxide	1/572 (0%)	0/1 (0%)	1 (100%)	36/704 (5%)	36/36 (100%)
Key safe	18/572 (3%)	9/18 (50%)	9/18 (50%)	89/704 (13%)	80/89 (90%)
Activity monitors assessment only	8/572 (1%)	4/8 (50%)	4/8 (50%)	5/704 (1%)	1/5 (20%)
Other devices	1/572 (0%)	0/1 (0%)	1/1 (100%)	6/704 (1%)	6/6 (100%)
<b>Intervention Group Technology</b>					
<b>Reminder or prompting devices</b>					
Date and time reminders	31/572 (5%)	13/31 (42%)	18/31 (58%)	46/704 (7%)	33/46 (72%)
Item locator devices	9/572 (2%)	8/9 (89%)	1/9 (11%)	11/704 (2%)	3/11 (27%)
Medication reminders/dispensers	56/572 (10%)	25/56 (45%)	31/56 (55%)	33/704 (5%)	8/33 (24%)
Voice recorders and memo minders	46/572 (8%)	27/46 (59%)	19/46 (41%)	38/704 (5%)	11/38 (29%)
Other reminder/prompting devices	1/572 (0%)	0/1 (0%)	1/1 (100%)	6/704 (1%)	6/6 (100%)
<b>Devices to promote safety</b>					
Activity monitors - on-going monitoring	5/572 (1%)	1/5 (20%)	4/5 (80%)	6/704 (1%)	5/6 (83%)
Fall detectors	75/572 (13%)	31/75 (41%)	44/75 (59%)	53/704 (8%)	22/53 (42%)
Continence management devices	1/572 (0%)	1/1 (100%)	0/1 (0%)	1/704 (0%)	0/1 (0%)
Alarm and pager units	5/572 (1%)	2/5 (40%)	3/5 (60%)	5/704 (1%)	3/5(60%)
Flood detectors and water temperature monitor	14/572 (2%)	9/14 (64%)	5/14 (36%)	11/704 (2%)	2/11 (18%)
Gas detectors	21/572 (4%)	8/21 (38%)	13/21 (62%)	19/704 (3%)	11/19 (58%)
Monitored carbon monoxide detectors	25/572 (4%)	8/25 (32%)	17/25 (68%)	22/704 (3%)	14/22 (64%)
Monitored smoke detectors	59/572 (10%)	39/59 (66%)	20/59 (34%)	47/704 (7%)	8/47 (17%)
Monitored extreme temperature sensors	26/572 (5%)	18/26 (42%)	15/26 (58%)	19/704 (3%)	8/19 (42%)
Lighting devices	2/572 (0%)	1/2 (50%)	1/2 (50%)	8/704 (1%)	7/8 (88%)
Other safety and security devices	15/572 (3%)	2/15 (13%)	13/15 (87%)	9/704 (1%)	7/9(78%)
<b>Safer walking technologies</b>					
To locate the user	43/572 (8%)	20/43 (47%)	23/43 (53%)	28/704 (4%)	8/28 (29%)
To alert the responder to movement	59/572 (10%)	25/59 (42%)	34/59 (58%)	37/704 (5%)	12/37 (32%)
<b>Communication devices</b>					
Intercoms	2/572 (0%)	0/2 (0%)	2/2 (100%)	1/704 (0%)	1/1 (100%)
Telephones	3/572 (1%)	0/3 (0%)	3/3 (100%)	7/704 (1%)	7/7 (100%)
Communication aids	0/572 (0%)	0/0 (0%)	0/0 (0%)	1/704 (0%)	1/1 (100%)
Other communication devices	1/572 (0%)	0/1 (0%)	1/1 (100%)	0/704 (0%)	0/0 (0%)
<b>Devices that support meaningful use of leisure time</b>					
Computer aids	0/572 (0%)	0/0 (0%)	0/0 (0%)	0 (0%)	0 (0%)
Dementia friendly TV/radio/music players	0/572 (0%)	0/0 (0%)	0/0 (0%)	0 (0%)	0 (0%)
Electronic photo albums/electronic reminiscence aids	0/572 (0%)	0/0 (0%)	0/0 (0%)	0 (0%)	0 (0%)
Electronic games	0/5572 (0%)	0/0 (0%)	0/0 (0%)	1/551 (0%)	1/1 (100%)
Other devices -support meaningful use of leisure time	1/572 (0%)	0/1 (0%)	1/1 (100%)	2/551 (0%)	2/2 (100%)
<b>Total</b>	572	266/572 (47%)*	306/572 (53%)	704	438/704 (62%)

\* ATT installed at any time point up to 24 weeks; if all ATT recommended was installed percentage would be 100%

Forsyth

Table 4: ATT installations 12 wks-104wks (for intervention arm only)

	12wks	24wks	52wks	104wks	Total (12wks-104wks)
<b>Control Group Technology installed</b>					
Basic ATT	235/580 (41%)	58/124 (47%)	45/87 (52%)	36/97 (37%)	374/888 (42%)
<b>Intervention Technology installed</b>					
Reminder/Prompting	116/580 (20%)	18/124 (15%)	9/87 (10%)	17/97 (18%)	160/888 (18%)
Safety	220/580 (38%)	45/124 (36%)	30/87 (35%)	43/97 (44%)	338/888 (38%)
Communication	8/580 (1%)	1/124 (0%)	2/87 (2%)	1/97 (1%)	12/888 (2%)
Support Leisure time	1/580 (0%)	2/124 (2%)	1/87 (1%)	0/97 (0%)	4/888 (0%)
Any other devices	0/580 (0%)	0/124 (0%)	0/87(0%)	0/97 (0%)	0/888 (0%)
Total installed	580	124	87	97	888
<b>Assessor</b>					
	12wks	24wks	52wks	104wks	Total (12wks-104wks)
Health/Social Care Professionals	126/580 (22%)	20/124(16%)	13/87 (15%)	17/97 (18%)	176/888 (20%)
ATT Assessor	152/580 (26%)	58/124 (47%)	23/87 (26%)	45/97 (46%)	278/888 (32%)
Other	68/580 (12%)	0/124 (0%)	3/87 (4%)	4/97 (4%)	75/888 (8%)
Unknown	234/580 (40%)	46/124 (37%)	48/87 (55%)	31/97 (32%)	359/888 (40%)
Total installed	580	124	87	97	888
<b>Assessment Method</b>					
	12wks	24wks	52wks	104wks	Total (12wks – 104wks)
In person at home	216/580 (37%)	70 (57%)	30 (34%)	55 (57%)	371/888 (41%)
In person not at home	7/580 (1%)	3 (2%)	1 (1%)	4 (4%)	15/888 (2%)
Telephone	50/580 (9%)	4 (3%)	1 (1%)	4 (4%)	59/888 (7%)
Using case notes	7/580 (1%)	0 (0%)	0 (0%)	2 (2%)	9/888(1%)
Other	56/580 (10%)	1 (1%)	2 (2%)	0 (0%)	59/888 (7%)
Unknown	244/580 (42%)	46 (37%)	53 (61%)	32 (33%)	375/888 (42%)
Total installed	580	124	87	97	888
<b>Monitoring</b>					
	12wks	24wks	52wks	104wks	Total (12wks – 104wks)
Yes	292/580 (51%)	56/124(45%)	32/87 (37%)	42/97 (43%)	422/888 (47%)
No	147/580 (25%)	45/124 (36%)	25/87 (29%)	40/97 (41%)	257/888 (29%)
Unknown	141/580 (24%)	23/124 (19%)	30/87 (34%)	15/97 (16%)	209/888 (24%)
Total installed	580	124	87	97	888
<b>Response</b>					
	12wks	24wks	52wks	104wks	Total (12wks – 104wks)
Formal Services	104/292 (36%)	29/56 (52%)	15/32 (47%)	14/42 (33%)	162/422 (38%)
Informal services	79/292 (27%)	11/56 (20%)	8/32 (25%)	16/42 (38%)	114/422 (27%)
Mixed services	106/292 36%)	14/56 (25%)	8/32 (25%)	12/42 (29%)	140/422 (33%)
Unknown	3/292 (1%)	2/56 (3%)	1/32 (3%)	0/42 (0%)	6/422 (2%)
Total installed	292	56	32	42	422


## Forsyth

Table 5: Current ATT practice with people with dementia using TIDieR format

TIDieR format [11]		Current ATT practice for people with dementia	
<b>When?*</b>	When did assessments, recommendation and installations happen?	Baseline (week 0), assessment and recommendations Week 12, 24, 52, 104, assessment & installation	
<b>What?</b>	What areas of assessment, within local ATT assessments, had higher fidelity to the ATT assessment standard?	Daily Activity, Memory, Mobility, & Social Support	
	What areas of assessment more frequently triggered the need for ATT?	Daily Activities, Memory, Problem Solving	
	What ATT was recommended more frequently within local ATT assessments?	Devices for safety issues & to remind/prompt with monitoring/formal response	
	What ATT was installed more frequently?	Devices for safety issues & to remind/prompt with monitoring/formal response and control arm devices (e.g., non-monitored smoke detectors)	
<b>How much?</b>	How much of the ATT assessment was completed?	52% of ATT assessment areas were completed 7.2 ATT assessment areas addressed on average (range 0-13)	
	How many ATT needs were present?	4.4 ATT needs on average (range 0-12 ATT risks)	
	How many ATT recommendations were identified?	3 ATT devices on average (range 1-14 ATT devices) 57% of participants had 1 or 2 ATT device recommended	
	How many installations happened?	4.2 ATT devices on average (range 1-15 ATT devices) (incl. control arm devices) 79% installed by week 24, with reduction of installation over time	
	How much monitoring and response happened?	47% of installed ATT required monitoring, of which 38% required formal response	
<b>Who?</b>	Who were the participants?	>80 years old, female, widowed, white British, not living alone, and had moderate dementia.	
	Who were the assessors of installed devices?	Baseline 57% Health & Social Care Professionals 33% ATT assessors 10% of assessors not known	Week 12-104 32% ATT assessors 20% Health & Social Care Professionals 40% of assessors not known 8% other
<b>Where?</b>	Where did the ATT assessment take place?	41% of installed devices were assessed by in-person home visits	
	Where did the installations take place?	Participant's home	
<b>Tailoring</b>	Was the devices tailored to the participants?	There was an expectation that ATT installations would be tailored to participants by the baseline ATT assessment, however, there was weak to moderate tailoring between.  a) Baseline ATT needs and ATT recommendations ( $\tau = 0.242$ ; $P < .000$ ) and  b) ATT recommendations and ATT installed ( $\tau = -0.470$ ; $P < .000$ ); 62% devices were installed for ATT needs that had not been identified in the assessment process, while 53% of the devices recommended as a result of assessment were not installed by week 24.	

\*pre-defined by Trial protocol

Figure 1 Cases available for analysis of ATT installations



Assistive technology and telecare; Assessment and practice for people with dementia (Forsyth)

### **Research in Context**

**Systematic Review:** The authors reviewed the literature using traditional (e.g. PubMed) sources and meeting abstracts and presentations. While assistive technology with people with dementia has not been described, there have been several studies, which advocate for assessment to drive installation. These relevant citations are appropriately cited.

**Interpretation:** Our findings are the first to describe assistive technology for people with dementia. It provides unique evidence that a) assistive technology practice is not consistent with best practice as promoted within the literature, b) assistive technology is deployed through at least two service delivery systems, namely, public and not-for-profit telecare provider.

**Future Directions:** The manuscript proposes a framework for the generation of new hypotheses and the conduct of additional studies. Examples include a) understanding how to translate recommendations into effective installations and b) understanding how different delivery systems for deploying assistive technology impacts on outcomes for people with dementia.

146 words