

**Supplementary Table 2. Results from the specific clinical & research Round 3 survey of experts (N = 19)**

	<b>HEALTHCARE PROFESSIONALS</b>	<b>FAMILY CAREGIVERS and PAID CAREGIVERS</b>
	Mode, Median (IQR)	Mode, Median (IQR)
Have techniques specific for ALS cognitive impairment (ALSci)	Very Important, 3 (1)	Important, 2 (1)
Have techniques specific for ALS behavioural impairment (ALSbi)	Very Important, 3 (1)	Important, 2 (1)
Have techniques specific for ALS cognitive and behavioural impairment (ALSbci)	Very Important, 3 (1)	Important, 2 (1)
Have techniques SPECIFIC to use in daily life when the diagnosis is ALS with FTD	Very Important, 3 (1)	Very Important, 2 (1)
Have general techniques which apply for all of the above altogether, i.e. do not distinguish between subtypes (“lumping as ALS plus”)	Important, 2 (1)	Important, 2 (1)
<b>Specific techniques:</b>		
Simplify decision making	Important, 2 (1.5)	Very Important, 3 (1)
Support problem-solving	Important, 2 (1)	Important, 2 (1)
Clarify Complex Information	Very Important, 3 (1)	Important, 2 (1)
Encourage & prompt engagement	Very Important, 3 (1)	Very Important, 3 (1)
Motivate everyone involved	Important, 2 (1)	Very Important, 3 (1)
Facilitate shared decision making	Very Important, 3 (0.5)	Very Important, 3 (1)
Support the family/caregiver	Very Important, 3 (0)	Very Important, 3 (1)

IQR = interquartile range, ALS = amyotrophic lateral sclerosis, FTD = frontotemporal dementia.