

1 ***Streptomyces* endophytes promote host health and enhance growth across**  
2 **plant species**

3  
4 Sarah F. Worsley<sup>1</sup>, Jake Newitt<sup>1,2</sup>, Johannes Rassbach<sup>2</sup>, Sibyl F. D. Batey<sup>2</sup>, Neil A. Holmes<sup>1,2</sup>,  
5 J. Colin Murrell<sup>3</sup>, Barrie Wilkinson<sup>2\*</sup> and Matthew I. Hutchings<sup>1,2\*</sup>  
6  
7

8 <sup>1</sup>School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich,  
9 Norfolk, United Kingdom. NR4 7TJ.

10 <sup>2</sup>Department of Molecular Microbiology, John Innes Centre, Norwich Research Park,  
11 Norwich, Norfolk, United Kingdom. NR4 7UH.

12 <sup>3</sup>School of Environmental Sciences, University of East Anglia, Norwich Research Park,  
13 Norwich, Norfolk, United Kingdom. NR4 7TJ.

14  
15 Email: [barrie.wilkinson@jic.ac.uk](mailto:barrie.wilkinson@jic.ac.uk); [matt.hutchings@jic.ac.uk](mailto:matt.hutchings@jic.ac.uk)  
16  
17  
18  
19

20 **Abstract.** *Streptomyces* bacteria are ubiquitous in soils and are well-known for producing  
21 secondary metabolites, including antimicrobials. Increasingly, they are being isolated from  
22 plant roots and several studies have shown they are specifically recruited to the rhizosphere  
23 and the endosphere of the model plant *Arabidopsis thaliana*. Here we test the hypothesis that  
24 *Streptomyces* bacteria have a beneficial effect on *A. thaliana* growth and could potentially be  
25 used as plant probiotics. To do this, we selectively isolated streptomycetes from surface washed  
26 *A. thaliana* roots and generated high quality genome sequences for five strains which we named  
27 L2, M2, M3, N1 and N2. Re-infection of *A. thaliana* plants with L2, M2 and M3 significantly  
28 increased plant biomass individually and in combination whereas N1 and N2 had a negative  
29 effect on plant growth, likely due to their production of polyene natural products which can  
30 bind to phytosterols and reduce plant growth. N2 exhibits broad spectrum antimicrobial activity  
31 and makes filipin-like polyenes, including 14-hydroxyisochainin which inhibits the Take-all  
32 fungus, *Gaeumannomyces graminis* var. *tritici*. N2 antifungal activity as a whole was  
33 upregulated ~2-fold in response to indole-3-acetic acid (IAA) suggesting a possible role during  
34 competition in the rhizosphere. Furthermore, coating wheat seeds with N2 spores protected  
35 wheat seedlings against Take-all disease. We conclude that at least some soil dwelling  
36 streptomycetes confer growth promoting benefits on *A. thaliana* while others might be  
37 exploited to protect crops against disease.

38  
39 **Importance.** We must reduce reliance on agrochemicals and there is increasing interest in  
40 using bacterial strains to promote plant growth and protect against disease. Our study follows  
41 up reports that *Arabidopsis thaliana* specifically recruits *Streptomyces* bacteria to its roots. We  
42 test the hypothesis that they offer benefits to their *A. thaliana* hosts and that strains isolated  
43 from these plants might be used as probiotics. We isolated *Streptomyces* strains from *A.*  
44 *thaliana* roots and genome sequenced five phylogenetically distinct strains. Genome mining  
45 and bioassays indicated that all five have plant growth promoting properties, including  
46 production of IAA, siderophores and ACC deaminase. Three strains significantly increased *A.*  
47 *thaliana* growth *in vitro* and in combination in soil. Another produces potent filipin-like  
48 antifungals and protected germinating wheat seeds against the fungal pathogen  
49 *Gaeumannomyces graminis* var. *tritici* (wheat Take-all fungus). We conclude that introducing  
50 *Streptomyces* strains into the root microbiome provides significant benefits to plants.

51

52 **Introduction.** The bacterial genus *Streptomyces* comprises more than 600 known species and  
53 they produce a diverse array of specialised metabolites that account for ~55% of the antibiotics  
54 currently used in human medicine (1). They are filamentous, spore-forming bacteria that are  
55 ubiquitous in soils where they play an important role in breaking down complex organic  
56 material (2, 3). Intriguingly, they only produce around 10% of their encoded secondary  
57 metabolites *in vitro* (2-4). Thus, understanding the role and regulation of their specialised  
58 metabolites in natural habitats is essential if we are to unlock the other 90% and discover new  
59 molecules (3). Increasingly, *Streptomyces* species are being recognised as important defensive  
60 symbionts of a wide range of invertebrate species including bees, beetles, digger wasps and  
61 ants (5-9). In addition to this, streptomycetes have also been shown to interact extensively with  
62 plant roots, inhabiting both the soil surrounding the plant root, called the rhizosphere, as well  
63 as the niche within and between root cells, referred to as the endophytic compartment (10, 11).  
64 It has even been suggested that their filamentous, hyphal growth and complex specialised  
65 metabolism may have evolved to facilitate interactions with plant roots, presumably to allow  
66 entry into root tissue and enable them to compete for food in the form of root exudates or more  
67 complex polymers that make up the plant cell wall (2).

68 Several recent studies have reported that streptomycetes are present, and sometimes enriched,  
69 in the endophytic compartment of the model plant *Arabidopsis thaliana* relative to the bulk soil  
70 (12-15) where they are attracted by plant metabolites in the root exudates such as salicylate  
71 and jasmonate (16, 17). They have also been isolated from the endosphere of many other plant  
72 species, including wheat, a crop of huge social and economic value (18-21). Due to their  
73 capacity to produce a large number of antimicrobial compounds and their ability to abundantly  
74 colonise plant roots, streptomycetes are gaining increasing interest from a biocontrol point of  
75 view (10, 11, 19, 22). A recent study demonstrated that certain strains can act as defensive  
76 mutualists of strawberry plants whereby they protect their host plant and pollinating bees  
77 against fungal infections (23). Many other isolates have been shown to protect important crops  
78 against infection and two strains have been developed into commercial biocontrol agents called  
79 Actinovate® and Mycostop® (10, 19, 22, 24). The ubiquity of streptomycetes in soil and their  
80 diverse specialised metabolism, combined with their ability to colonise plant roots, makes  
81 streptomycetes attractive for this purpose. Their spore-forming capabilities also makes them  
82 tolerant of many environmental pressures, allowing them to be applied as dried seed coatings  
83 which remain viable under various agricultural conditions.

84           The aim of this study was to test the hypothesis that plant-associated streptomycetes  
85 provide benefits to their host *A. thaliana* plant and that strains isolated from *A. thaliana* might  
86 confer benefits to important crop plants, such as wheat. We hypothesised that these strains may  
87 play defensive or plant growth promoting roles since this genus is consistently recruited to the  
88 plant root microbiome. To this end, we generated high quality genome sequences for five  
89 *Streptomyces* species which we isolated from the root microbiome of *A. thaliana* plants and  
90 which could recolonise *A. thaliana* roots when applied as seed coatings. All five strains, named  
91 L2, M2, M3, N1 and N2, encode a large number of secondary metabolite biosynthetic gene  
92 clusters (BGCs) and all strains inhibited at least one bacterial or fungal pathogen. Strain N2  
93 demonstrated broad spectrum antifungal and antibacterial activity and was able to inhibit  
94 growth of the Take all fungus, *Gaeumannomyces graminis* var. *tritici*, an economically  
95 important pathogen of wheat, both *in vitro* and on germinating wheat seeds. The antifungal  
96 activity of N2 was increased 2-fold *in vitro* in response to indole-3-acetic acid (IAA) and  
97 purification of the antifungal molecules identified a number of filipin-like compounds.  
98 Curiously, N2 reduced *A. thaliana* growth *in vitro* (but not in soil) which may have been caused  
99 by filipins targeting sterols in the plant cell membranes at high concentrations. Strains L2, M2  
100 and M3 all promoted *A. thaliana* growth *in vitro* and when applied in combination to seeds  
101 planted in soil, and they all have well characterised plant growth promoting (PGP) traits  
102 including the production of plant growth hormones, siderophores and ACC deaminase. We  
103 conclude that *A. thaliana* can acquire significant benefits from recruiting streptomycetes to  
104 their root microbiome which they likely attract through the deposition of root exudates and  
105 dead root material into the bulk soil. Additionally, mining plant-streptomycete interactions may  
106 yield novel biocontrol and plant-growth promoting agents that could be developed for future  
107 applications in agriculture.

108

## 109 **Results**

110 **Isolation and genome analysis of *A. thaliana* root-associated *Streptomyces* strains.** To  
111 culture bacteria from *A. thaliana* roots, the roots of 4 week old plants growing Levington's  
112 seed and modular compost were washed and sonicated in sterile buffer (as described in 12) to  
113 remove soil particles. This process removed all bacteria apart from those there were tightly  
114 bound to the plant root surface (the rhizoplane) or were living within the plant root tissue as  
115 endophytes (12). The roots were then crushed in sterile 10% glycerol and serial dilutions were  
116 plated onto soya flour mannitol, starch casein and minimal salts agar. Colonies resembling

117 streptomycetes were purified by restreaking and were then identified by colony PCR and 16S  
118 rRNA gene amplicon sequencing using the universal primers PRM341F and MPRK806R  
119 (Tables 1 and 2). Based on 16S rRNA sequencing, five phylogenetically distinct strains (L2,  
120 M2, M3, N1 and N2) were then selected for genome sequencing. We used the PacBio RSII  
121 platform (as described in 25) to generate high-quality genome sequences for the five isolates,  
122 in addition to three strains of *Streptomyces lydicus*, which are known plant endophytes (26,  
123 27). One of these strains was isolated from the horticultural product Actinovate® and the other  
124 two came from the ATCC culture collection (Table 2). All eight linear genomes are within the  
125 size range typical for this genus and do not show any significant reductions compared to the  
126 genomes of other sequenced *Streptomyces* species (Table 1). The genomes of the *A. thaliana*  
127 associated strains L2, M2, M3, N1 and N2 were uploaded to the automated multi-locus species  
128 tree (28) for phylogenetic classification. The highest average nucleotide identity (ANI) values  
129 of strains L2, M2 and M3 to strains in the database are 88.3%, 94.7% and 91.1%, respectively;  
130 these are below the 95% threshold that is generally used to assign strains to a known species  
131 (29, 30) so they could be new *Streptomyces* species. Strain N1 has a 98.7% ANI to  
132 *Streptomyces albidoflavus* and strain N2 has a 97.6% ANI to *Streptomyces griseofuscus*,  
133 suggesting they belong to these clades (Table 1).

134

135 ***Streptomyces* bacteria colonize *A. thaliana* roots.** To investigate whether the sequenced  
136 *Streptomyces* strains can promote plant growth and fitness, we established root infection assays  
137 in which seeds were coated with a suspension of pre-germinated *Streptomyces* spores. Tagging  
138 the strains with eGFP and the apramycin resistance (*aac*) gene allowed visual confirmation of  
139 root infection using confocal microscopy (Fig. 1) and selective re-isolation of the strains on  
140 agar plates containing apramycin, confirming that they were able to re-colonise plant roots.

141

142 ***Streptomyces* strains M2, M3 and L2 have growth-promoting effects in *A. thaliana*.** Next  
143 we wanted to determine if any of the *Streptomyces* strains isolated from *A. thaliana* roots can  
144 enhance plant growth, so we established root infection assays on agar plates, whereby  
145 *Streptomyces* spores were inoculated directly onto the roots of young *A. thaliana* seedlings.  
146 We tested all the genome-sequenced strains from this study and found that the inoculation of  
147 different strains had a significant effect on the dry weights of plants grown on agar, compared  
148 to a sterile control (Fig. 2,  $F_{(8,135)} = 27.63$ ,  $P < 0.001$  in an ANOVA test). Strains L2, M2 and  
149 M3 significantly increased *A. thaliana* dry biomass under these conditions when compared to  
150 sterile control plants (Fig. 2,  $P < 0.05$  in Tukey's Honestly Significant Difference (HSD) tests).

151 In comparison, the three strains of *S. lydicus* had no effect on *A. thaliana* plant biomass *in vitro*  
152 despite this species being previously noted to have plant-growth-promoting effects (26).  
153 *Streptomyces* strains N1 and N2 significantly reduced the growth of *A. thaliana in vitro* relative  
154 to control plants (Fig. 2;  $P < 0.05$  in Tukey's HSD).

155

156

157

158 To test whether each of the new isolates could promote plant growth in soil, spores of each  
159 strain were applied individually to *A. thaliana* seeds planted in Levington's F2 seed and  
160 modular compost. Since, it is known that some strains can work synergistically to promote  
161 plant growth (33-35), a mixture of spores of L2, M2 and M3 (the strains that promoted plant  
162 growth *in vitro*) was also added to seeds. Strain inoculation had a significant influence on the  
163 dry weight of plants grown in compost (ANOVA test on log-transformed dry weight:  $F_{(7,58)} =$   
164  $3.9358$ ,  $P = 0.001$ ). However, interestingly none of the strains had an effect on plant growth  
165 when applied individually (Fig. 3,  $P > 0.05$  in Tukey HSD test), but the application of a spore  
166 mixture of L2, M2 and M3 significantly increased plant dry weight from an average of  $12.69$   
167  $\text{mg} \pm 1.94$  (mean  $\pm$  SE) for control plants, to  $39.29 \text{ mg} \pm 4.39$  (Fig. 3).

168

169 **Characterisation of plant growth promoting traits.** KEGG pathway analysis revealed that  
170 the genomes of all our sequenced *Streptomyces* strains possess genes encoding proteins  
171 involved in the biosynthesis of IAA, which can contribute to shoot and root growth (Table S2).  
172 For example, all strains have genes encoding key proteins involved in the indole-3-acetamide  
173 (IAM) pathway, whereby tryptophan is converted to IAM via a tryptophan 2-monooxygenase  
174 enzyme (KEGG reaction R00679). IAM is then further converted to IAA through the action of  
175 an amidase enzyme (KEGG reaction R03096). Several strains also possessed genes encoding  
176 enzymes involved in the tryptamine (TAM) pathway. *In vitro* colorimetric assays using  
177 Salkowski reagent (as described in 36) qualitatively confirmed the ability of all strains to make  
178 IAA (Fig. S1). In addition to IAA, the genomes of all the *Streptomyces* isolates possess up to  
179 two copies of genes encoding the enzyme aminocyclopropane-1-carboxylate (ACC)  
180 deaminase. This cleaves ACC, which is the direct precursor for the plant phytohormone  
181 ethylene, into ammonia and 2-oxobutanoate, Kegg reaction: R00997 (Table S1). Bacteria can  
182 use the ammonium released in this reaction as a nitrogen source, and all the isolates are capable  
183 of utilising ACC as a sole nitrogen source in minimal medium (Fig. S2). There is evidence that  
184 the activity of the ACC deaminase enzyme can reduce plant damage and early-onset

185 senescence caused by excessive ethylene production under prolonged periods of plant stress,  
186 by removing the substrate for ethylene biosynthesis (37, 38).

187

188 **Bioactivity of root-associated *Streptomyces* isolates.** *Streptomyces* bacteria are well-known  
189 for their ability to produce a wide range of specialised metabolites which can have bioactivity  
190 against bacteria, fungi, viruses, nematodes, insects and plants (1, 3, 39). We reasoned that our  
191 strains likely make antimicrobial natural products and that strains N1 and N2 may encode  
192 herbicidal compounds given that they have an adverse effect on *A. thaliana* growth *in vitro*  
193 (Fig. 2). All eight sequenced genomes were submitted to the bacterial antiSMASH 5.0 portal  
194 (31) which can predict BGCs for major types of specialised metabolite. This identified between  
195 21 and 34 putative specialised metabolite BGCs in the sequenced genomes, which is within the  
196 typical range for this genus (Table S2). This includes BGCs predicted to encode polyketide  
197 synthases (PKS), non-ribosomal peptide synthases (NRPS), ribosomally encoded post  
198 translationally modified peptides (RiPPs) and terpenes (Table S2). All five strains encode  
199 multiple siderophore BGCs which are molecules that chelate metal ions, such as iron,  
200 generating soluble complexes that can be taken up by plant roots and contribute to plant growth  
201 (40). Siderophores also help microorganisms to compete in the soil, rhizosphere and  
202 endosphere, with the added benefit that this may act to exclude plant pathogens that are also  
203 competing for iron (41).

204 To test whether antimicrobial compounds could be produced *in vitro*, sequenced strains  
205 were tested for their ability to inhibit a range of bacterial and fungal pathogens, including the  
206 bacterial plant pathogen *Pseudomonas syringae* and the fungus *G. graminis* var. *tritici* which  
207 is the causative agent of wheat Take-all, one of the most economically damaging diseases of  
208 wheat worldwide (19). All eight strains exhibited antibacterial activity and N2 inhibited all of  
209 the bacterial strains that were tested: *Bacillus subtilis*, methicillin resistant *Staphylococcus*  
210 *aureus* (MRSA), *Escherichia coli* and *Pseudomonas syringae* (Fig. S3; Table S3). The N2  
211 genome encodes a rich repertoire of BGCs including several putative antibacterial BGCs  
212 predicted to encode the proteins responsible for the biosynthesis of albusnodin, albaflavenone  
213 and diisonitrile antibiotic SF2768-like antibiotics, the antiproliferative actinomycin D which is  
214 used clinically as an anti-cancer therapeutic, in addition to a possible analogue of the  
215 proteasome inhibitor cinnabaramide (Table S2). The actinomycins exhibit broad-spectrum  
216 bioactivity by binding to DNA and inhibiting transcription; thus it could result in the inhibition  
217 of all the pathogenic indicator strains (42). M2 is interesting because it inhibits *Pseudomonas*  
218 *syringae* but none of the other bacteria that were tested (Table S3). It encodes a relatively

219 modest number of BGCs with one for a putative aquamycin-type antibiotic (Table S2).  
220 However, this effect may be due to a siderophore because an antibacterial compound that  
221 inhibits *Pseudomonas* would also be expected to inhibit *B. subtilis* and *E. coli* (Table S3). L2  
222 encodes clusters for the biosynthesis of albaflavenone and thioviridamide-like molecules  
223 (Table S2) and inhibits *B. subtilis in vitro* (Table S4). M3 inhibits *B. subtilis* and *P. syringae*  
224 (Table S3) and encodes a type 3 PKS for the biosynthesis of alkylresorcinol-type phenolic  
225 lipids (Table S3) which may have antibacterial activity. Finally, N1 inhibits *B. subtilis* (Table  
226 S3) and encodes several BGCs encoding molecules with potential antibacterial activity  
227 including a putative polycyclic tetramate macrolactam (PTM) antibiotic which are widely  
228 distributed but often cryptic (43), and a desotamide like antibiotic (Table S2). It should be  
229 noted that up to 90% of specialised metabolite BGCs are cryptic (i.e. not expressed under  
230 laboratory conditions) in *Streptomyces* species, largely because we do not understand the  
231 environmental or host-derived signals that activate their expression (3). In addition, there are  
232 many known specialised metabolites for which the BGCs have not yet been identified so it is  
233 likely that at least some of the known BGCs in these genomes are not expressed under the  
234 growth conditions used in this study and/or that these strains encode antimicrobials whose  
235 BGCs cannot be predicted by antiSMASH 5.0.

236 The strains L2, M2 and M3 do not exhibit antifungal activity against *C. albicans*, *L.*  
237 *prolificans* or *G. graminis* var. *tritici in vitro* (Table S3), although it is possible that these strains  
238 may encode cryptic antifungal compounds that are not expressed under the conditions used in  
239 experiments. For example, M3 encodes a clavam-like cluster which has a similar structure to  
240 those that encode the antifungal compound clavamycin (Table S2). In comparison, strains N1  
241 and N2 both have type 1 (T1) PKS gene clusters which are predicted to encode the biosynthesis  
242 of known polyene antifungal compounds (Table S2). The N1 T1PKS gene cluster is predicted  
243 to encode candicidin and the N2 T1PKS gene cluster is predicted to encode filipin-type  
244 antifungals (Table S3). Both of these metabolites inhibit fungal growth by binding to sterols in  
245 their cell walls and it is known that filipins can also bind to phytosterols in plant cell membranes  
246 and reduce the growth of plant roots at high concentrations (44-46). Thus, it is possible that N1  
247 and N2 are making polyenes *in planta* and this is what caused the reduction in growth observed  
248 during *in vitro* inoculation experiments (Fig. 2). These strains did not influence plant growth  
249 in compost (Fig. 3) suggesting that the effect was diluted in a non-sterile system, or that the  
250 polyenes were not expressed under these conditions. Despite the presence of antifungal-like  
251 clusters in the other *A. thaliana* isolates, only N2 exhibited antifungal activity *in vitro* (Table  
252 S3). In fact, N2 demonstrated potent and broad-spectrum antifungal activity as it inhibited all



253 three test strains; these were the human pathogenic fungal species *Candida albicans* and multi-  
254 drug resistant *Lomentospora prolificans* as well as the plant pathogenic Take-all fungus  
255 *Gaeumannomyces graminis* var *tritici* (Table S3).

256 Production of antifungal compounds likely gives N2 an advantage in the rhizosphere  
257 and, consistent with this, N2 antifungal activity against *C. albicans* was shown to increase 2-  
258 fold *in vitro* in the presence of indole-3-acetic acid (IAA), as judged by the size of the inhibition  
259 zone (Fig. 4). IAA is the precursor to the plant phytohormone auxin which regulates processes  
260 involved in growth and development (47). It is also made by many microbial species in the  
261 rhizosphere, including the bacterial strains that were isolated in this study (Fig. S1 & Table 2),  
262 and has been noted previously for its role in both intra- and inter-kingdom signalling (47-49).  
263 Thus, N2 may be driven to produce antifungals in close proximity to plant roots when it is in  
264 competition with other microbes.

265

266 **Purification and identification of antifungal compounds from *Streptomyces* strain N2.** To  
267 extract the molecules with antifungal activity, strain N2 was plated on SFM agar (4 L, 120  
268 plates) and grown for eight days at 30 °C before extraction with ethyl acetate. The crude extract  
269 had an intense orange colour and HPLC analysis identified components with UV characteristics  
270 typical of polyene metabolites. Comparison to an authentic commercial sample of the filipin  
271 complex of *Streptomyces filipinensis* (Sigma Aldrich) confirmed the presence of several filipin  
272 related molecules, although the two major components eluted earlier than these molecules (Fig.  
273 S4). The two major components eluted very closely together and had *m/z* values consistent  
274 with the previously reported molecules pentamycin (also known as fungichromin) and 14-  
275 hydroxyisochainin (50-52). These filipin-like compounds (Fig. 5) can all be assigned to the  
276 type 1 PKS BGC in region 1 (Table S2, Fig. S17) (51, 53, 54). Despite a challenging elution  
277 profile sufficient separation was accomplished by applying multiple purification steps, and the  
278 isolated material enabled structural confirmation by 2D NMR (Figs S5-S16, Tables S4-S6)  
279 accompanied by comparison to published data (50-52) and bioinformatics analysis of the  
280 associated PKS encoding genes in strain N2 (53-55). The extract also contained a mixture of  
281 components with an absorption maximum at 444 nm and this mixture was red after separation  
282 from the polyene fraction. LCMS analysis indicated that these components were consistent  
283 with the known actinomycin congeners D, X<sub>2</sub> and X<sub>0β</sub> respectively (Fig. S4). These are most  
284 likely encoded by BGC5 in the N2 genome (Table S2) (56).

285 Pentamycin is active against *Candida albicans* and *Trichomonas vaginalis* and is used  
286 for the treatment of vaginal candidiasis, trichomoniasis, and some mixed infections (57-59). It

287 is identical in structure to filipin III apart from the presence of an additional secondary hydroxyl  
288 group at C14 (Fig. 5) which is added by a cytochrome P450 monooxygenase encoded by a  
289 gene located directly upstream of the first PKS gene (A1) in the *pentamycin* BGC as reported  
290 recently (51); the same BGC architecture is observed in *Streptomyces* strain N2 (Fig. S17). 14-  
291 hydroxyisochainin shares the same polyene core structure as pentamycin but carries an altered  
292 side chain lacking two carbon atoms and is indicative of different length extender units being  
293 utilized by the final module of the type I PKS during biosynthesis (Fig. 5). This observation is  
294 unexpected as the co-production of pentamycin and 14-hydroxyisochainin has only been  
295 observed as a result of precursor-directed biosynthesis (52). Thus, this example of co-  
296 production by *Streptomyces* strain N2 appears to be novel. Disc diffusion bioassays confirmed  
297 that pentamycin and 14-hydroxyisochainin inhibit the growth of *C. albicans* (Fig. S18), but  
298 only 14-hydroxyisochainin was able to inhibit *G. graminis* var. *tritici* (Fig. S19), suggesting  
299 that 14-hydroxyisochainin is responsible for the antifungal activity of N2 against the Take-all  
300 fungus, potentially in combination with lower quantities of other products of the filipin-like  
301 BGC. The complex consisting of actinomycin D, X<sub>2</sub> and X<sub>0β</sub>, which were co-purified from N2  
302 extracts (Fig. S4), did not have antifungal activity suggesting they are not responsible for N2  
303 antifungal bioactivity *in vitro* (Fig. S18). Neither pentamycin, 14-hydroxyisochainin or the  
304 actinomycin complex inhibit the growth of *E. coli* K12 (Fig. S18) suggesting that a previously  
305 undescribed compound is responsible for the observed inhibition of *E. coli* and *P. syringae* by  
306 N2 *in vitro* (Fig. S3, Table S3).

307

308 ***Streptomyces* strain N2 protects wheat against Take-all.** In order to test whether strain N2  
309 has the potential to protect wheat plants against *G. graminis* var. *tritici*, we inoculated surface  
310 sterilised wheat seeds (*Triticum aestivum* var. Paragon) with N2 spores or left them sterile as a  
311 control. These seeds were germinated next to a central plug of Take-all fungus on potato  
312 glucose agar (PGA) plates. On the control plates, *G. graminis* grew outwards across the agar  
313 plate and over the sterile wheat seedlings, whereas the seeds that had been inoculated with N2  
314 spores were resistant to *G. graminis* var. *tritici*, as indicated by a zone of inhibition around the  
315 germinating wheat seeds (Fig. 6). The most parsimonious explanation is that the N2 spores have  
316 germinated and are producing 14-hydroxyisochainin and possibly other filipins, that inhibit the  
317 Take-all fungus.

318 To further test the potential of the *Streptomyces* strain N2 to act as a biocontrol strain  
319 against Take-all *in vivo*, wheat seeds were soaked in N2 spores, allowed to dry, and then grown  
320 in sterile vermiculite containing *G. graminis* var. *tritici* mycelia. After 3 weeks of growth at

321 25°C, Take-all infection severity was scored on a scale of 0-8, using an infection scoring  
322 system as follows: : 0=no infection; 1=maximum of one lesion per root; 2=more than one lesion  
323 per root; 3=many small and at least one large lesion per root; 4=many large lesions per root;  
324 5=roots completely brown; 6=roots completely brown plus infection in stem; 7=roots  
325 completely brown, infection in stem and wilted yellow leaves; 8=entire plant brown and wilted  
326 (Fig. S20). The wheat plants that had germinated and grown in the absence of Take-all (with  
327 or without N2 spores) were healthy and infection free (Fig. 7). However, those seeds that had  
328 grown from un-inoculated seeds in the presence of the Take-all fungus showed extensive and  
329 severe levels of Take-all disease, with an average infection score of  $7.24 \pm 0.26$  SE (Fig. 7).  
330 Most of the plants in this treatment group exhibited infected roots, stems and leaves, which all  
331 appeared senescent and brown (Fig. 7). However, there was a significant effect of plant  
332 treatment on infection score (Kruskal-Wallis test  $H_{DF=3} = 83.41$ ,  $P = <0.001$ ). Plants that had  
333 grown from seeds coated in N2 spores (N = 25), demonstrated a small, but significant, decrease  
334 in average infection severity to  $5.47 \pm 0.59$ , compared to plants grown from uninoculated seeds  
335 in the presence of Take-all (Fig. 7, Dunn's test between inoculated and sterile wheat grown  
336 with *G. graminis*:  $P = 0.023$ ). As plants were grown in a sterile system in this experiment, it is  
337 likely that the streptomycete experienced low levels of nutrient availability compared to in a  
338 natural soil environment. We hypothesise that greater levels of competition and nutrients may  
339 fuel greater levels of antibiotic production by N2, and thus the strain could offer greater levels  
340 of protection against host infection in a more natural soil-plant system.

341

342 **Discussion.** *Streptomyces* species have traditionally been described as free-living soil bacteria  
343 but given that most bare soils are rapidly colonised by vegetation it is perhaps not surprising  
344 that they are also effective at colonising the rhizosphere and endosphere of plants (10). It has  
345 even been suggested that their filamentous growth may have evolved to facilitate plant root  
346 colonisation since this trait evolved 50 million years after plants first colonised land, around  
347 450mya (2, 10). Certainly, their ability to sporulate provides a useful mechanism for vertical  
348 transmission across plant generations, via the soil. In this study we followed up reports that  
349 *Streptomyces* bacteria are enriched in the *A. thaliana* rhizosphere and specifically recruited by  
350 plant-produced compounds such as the plant hormones salicylate and jasmonate (12-17). We  
351 aimed to isolate and characterise plant-associated streptomycetes from *A. thaliana* and test  
352 whether they can be beneficial to their plant host. We generated high quality genome sequences  
353 for five *A. thaliana* root-associated strains and three strains of the known endophyte *S. lydicus*.  
354 We found that three of the five root-associated strains significantly increased the biomass of *A.*

355 *thaliana* plants when they were applied to seeds or roots, both *in vitro* and when applied in  
356 combination in soil. Two others, N1 and N2, significantly decreased the biomass of *A. thaliana*  
357 *in vitro*, most likely because they make polyenes that bind to sterols, including the phytosterols  
358 found in plant cell walls (44-46), which probably had a negative effect on the plant in a sterile  
359 system. However, this effect was removed in a compost system with neither strain influencing  
360 plant growth. Our work suggests therefore, that while *Streptomyces* are consistently associated  
361 with *A. thaliana* roots and enriched in the root microbiome compared to the surrounding bulk  
362 soil, not all strains that are competitive in the rhizosphere and endosphere necessarily have a  
363 beneficial effect on host fitness. This is important from an ecological and applied perspective,  
364 the former because it helps us to better understand the microbial and host factors influencing  
365 plant microbiome assembly and the latter because tipping the balance in the plant's favour, for  
366 example by applying beneficial species as seed coatings or soil additives that can competitively  
367 colonise roots, could improve crop yields (19, 60, 61). Such strains could be used as biological  
368 growth promoters to replace the use of harmful pesticides and fertilisers which have negative  
369 effects on the wider ecosystem and also contribute to climate change (10, 11, 19, 60, 61). As a  
370 proof of this concept we took strain N2, which makes the polyene antifungals including  
371 pentamycin, 14-hydroxyisochainin and filipin III, and coated seeds of spring bread wheat with  
372 its spores. N2 was able to protect germinating wheat seedlings against the Take-all fungus *in*  
373 *vitro* and significantly reduce Take-all disease progression in wheat plants grown in sterile  
374 vermiculite. Although we do not have an assay for Take-all disease in soil the protective effect  
375 may be even greater because nutrients in the form of organic matter could provide a more  
376 beneficial growing environment for the *Streptomyces* strain than sterile vermiculite (62) and  
377 the presence of a greater diversity of microbes may fuel antimicrobial production by this strain  
378 as a result of interference competition (63). It is also intriguing that IAA increased the  
379 antifungal activity of strain N2 since it provides compelling evidence that environmental  
380 signals can alter the expression of secondary metabolites, and probably as an extension also  
381 influence interspecies competition (47, 49). Although there is much future work to do to  
382 understand this phenomenon our results provide a system to begin to understand how  
383 secondary metabolites are regulated and used by microbes in nature and may also provide new  
384 tools for activating the 90% of BGCs that are silent in these bacteria. Plant-associated  
385 *Streptomyces* strains may yet provide us with a new generation of antimicrobials for the clinic  
386 and might also be harnessed to improve our food security. Understanding the ecology of these  
387 bacteria and their associated natural products will be crucial if we are to achieve these goals.

388

## 389 **Materials and Methods**

390

391 **Isolation of root-associated *Streptomyces* strains.** Buffers and media recipes are listed in  
392 Table S7. Strains, primers and plasmids are listed in Table 2. Wild-type *A. thaliana* Col-0 seeds  
393 were sterilised by washing in 70% (v/v) ethanol for 2 minutes, 20% (v/v) sodium hypochlorite  
394 for 2 minutes, then five times in sterile water. Individual seeds were sown into pots of sieved  
395 Levington's F2 seed and modular compost, placed at 4°C for 24 hours, then grown for 4 weeks  
396 under a 12-hour photoperiod (12 hours light/12 hours dark) at 22°C. Plants were taken  
397 aseptically from pots and their roots tapped firmly to remove as much adhering soil as possible.  
398 Root material was placed into sterile PBS-S buffer for 30 minutes on a shaking platform and  
399 before being transferred into fresh PBS-S and washed for 30 minutes. Any remaining soil  
400 particles were removed with sterile tweezers. Cleaned roots were then transferred to 25ml of  
401 fresh PBS-S and placed in a sonicating water bath for 20 minutes to remove any residual  
402 material still attached to the root surface; this was to ensure that any remaining bacteria were  
403 either present in the endophytic compartment or were very firmly attached to the root surface  
404 ("the rhizoplane") (12). The roots were crushed in sterile 10% (v/v) glycerol and serial dilutions  
405 were spread onto either soya flour mannitol (SFM) agar, starch casein agar, or minimal salts  
406 medium agar containing sodium citrate. Plates were incubated at 30°C for up to 14 days.  
407 Colonies resembling streptomycetes were re-streaked onto SFM agar and identified by 16S  
408 rRNA gene PCR amplification and sequencing with the universal primers PRK341F and  
409 MPRK806R. Sequencing was carried out by Eurofins Genomics, Germany. Three strains of  
410 *Streptomyces lydicus*, which are known to associate with plant roots, were also used in  
411 experiments; *S. lydicus* WYEC108 was isolated from the commercial biocontrol product  
412 Actinovate® and two more *S. lydicus* strains (ATCC25470 and ATCC31975) were obtained  
413 from the American Type Culture Collection. *Streptomyces* strains were maintained on SFM  
414 agar (N1, N2, M2, M3), Maltose/Yeast extract/Malt extract (MYM) agar with trace elements  
415 (L2) or ISP2 agar (*S. lydicus* strains). Strains were spore stocked as described previously (64).

416

417

418 **Genome sequencing and analysis** High quality genome sequences were obtained for  
419 *Streptomyces* strains N1, N2, M2, M3, and L2, as well as the three known strains of  
420 *Streptomyces lydicus* using PacBio RSII sequencing technology at the Earlham Institute,  
421 Norwich, UK, as described previously (25). The automated Multi-Locus Species Tree  
422 (autoMLST) server (28) was used to phylogenetically classify the *Streptomyces* strains L2, M2,  
423 M3, N1 and N2. BGCs were predicted using antiSMASH 5.0 (31) and genomes were annotated  
424 using RAST (65). Amino acid sequences were uploaded to the KEGG Automatic Annotation  
425 Server (KAAS) for functional annotation of genes and metabolic pathway mapping (66).

426

427 **Generating eGFP-labelled *Streptomyces* strains.** Plasmid pIJ8660 containing a codon  
428 optimised eGFP gene under the control of the constitutive *ermE\** promoter and the *aac*  
429 apramycin resistance marker (67) was conjugated into *Streptomyces* strains (64). Exconjugants  
430 were selected and maintained on SFM agar plates (Table S7) containing 50  $\mu\text{g ml}^{-1}$  apramycin.  
431 For confocal microscopy, *A. thaliana* Col-0 seeds were germinated on MSk (Table S7) and  
432 grown vertically at 22 °C for 9 days under a 12-hour photoperiod. Seedlings were then  
433 transferred to MSk (1.5% agar, 0% sucrose, Table S7) and allowed to equilibrate for 24 hours  
434 before being inoculated with 1  $\mu\text{l}$  of spore suspension ( $10^6$  spores  $\text{ml}^{-1}$ ) of the eGFP-tagged M3  
435 or *Streptomyces coelicolor* M145 strains. As a known coloniser of plant roots, *S. coelicolor*  
436 was used as a control (32). Inoculated seedlings were then left to grow for 3 days before being  
437 washed in a 20% (v/v) solution of glycerol containing 1  $\mu\text{g ml}^{-1}$  276 SynaptoRed™ for 10  
438 minutes. A 20 mm section of root (taken from the base of the petiole) was then mounted onto  
439 a slide with 100  $\mu\text{l}$  of the SynaptoRed™/glycerol solution. Samples were imaged using a Zeiss  
440 LSM510 META laser-scanning confocal microscope with a PlanApochromat 63x (1.4 NA)  
441 objective. Green fluorescent protein was excited at 488 nm and emission collected through a  
442  $527.5 \pm 22.5$  nm bp filter, and FM4-64 excited at 543 nm and emission collected through a  
443  $587.5 \pm 27.5$  nm bp filter.

444

445 **Plant growth promotion assays.** *A. thaliana* Col-0 seeds were sterilised and plated onto MSk  
446 medium (1% (w/v) sucrose and 0.8% (w/v) agar, Table S7). These were then left at 4 °C in the  
447 dark for 24 hours before being placed, vertically, under long day growth conditions (12 h light/  
448 12 h dark) at 22°C for 10 days. Seedlings were then transferred to square agar plates containing  
449 MSk agar (as above, Table S7) and allowed to equilibrate, vertically, overnight at 22 °C. 1  $\mu\text{l}$

450 of *Streptomyces* spores ( $10^6$  ml<sup>-1</sup>) from each of the sequenced isolates was added to the top of  
451 the root system of each seedling and allowed to dry. 16 replicate seedlings were inoculated per  
452 sequenced *Streptomyces* strain. 10% (v/v) glycerol was added to control seedlings. Plates were  
453 grown vertically for 16 days, 12 h light/ 12 h dark at 22°C before measuring plant biomass (dry  
454 weight). The biomass of plants with different inocula were compared via ANOVA and Tukey's  
455 Honestly Significant Difference (HSD) tests using R 3.2.3 (68); biomass was log-transformed  
456 during analyses to ensure normality of residuals. Strains were also tested for their ability to  
457 promote *A. thaliana* growth in compost. Sterile *A. thaliana* Col-0 seeds were placed into a  
458 solution of 2xYT (Table S7) containing  $10^6$  pregerminated spores ml<sup>-1</sup> of each strain or no  
459 spores as a control. Seeds were incubated in the spore solution for 2 hours, before being  
460 transferred to pots containing sieved Levington's F2 seed and modular compost. An additional  
461 3 ml of pre-germinated spores (or sterile 2xYT) was pipetted into the soil surrounding each  
462 seed. The strains L2, M2 and M3 were also tested for their ability to promote plant growth in  
463 combination;  $10^3$  spores ml<sup>-1</sup> of each strain were mixed together and pregerminated in 2xYT  
464 before being used as above. Pots were then placed at 4°C for 48 hours before being grown for  
465 6 weeks under a photoperiod of 12 hours light/12 hours dark. There were 8 replicate pots per  
466 treatment. After 6 weeks, the plants were removed from pots and cleaned by washing in PBS-  
467 S (Table S7) and using tweezers to remove adhering soil particles. Plants were then dried in an  
468 oven at 50°C enabling plant dry weight to be calculated. An ANOVA test and Tukey's HSD  
469 tests were used (as above) to test for an effect of strain inoculation on plant dry weight. Dry  
470 weights were log-transformed prior to analysis.

471

472 **Indole 3 Acetic Acid (IAA) production assays.** *Streptomyces* isolates were grown on  
473 cellophane membranes covering YMD media supplemented with 5 mM tryptophan (Table S7).  
474 After 7 days, cellophane membranes with bacterial biomass were removed and plates were  
475 flooded with Salkowski reagent (as described in 36). A red colour indicates that IAA has  
476 leached into the medium.

477

478 **1-Aminocyclopropane-1-carboxylic acid (ACC) degradation assays.** To test for the use of  
479 ACC as a sole nitrogen source, *Streptomyces* strains were streaked onto Dworkin and Foster  
480 medium (69) in which 0.2% (w/v) NH<sub>4</sub>SO<sub>4</sub> or 0.051% (w/v) ACC was added as a sole nitrogen  
481 source, or no nitrogen source as a control. Plates were incubated for 10 days at 30°C before  
482 imaging.

483

484 **Antibiotic bioassays.** Spores ( $4 \mu\text{l}$  of  $10^6 \text{ ml}^{-1}$  solution) of individual *Streptomyces* isolates  
485 were pipetted onto the centre of agar plates and incubated at  $30^\circ\text{C}$  for 7 days before adding the  
486 pathogenic indicator strains (see Table 2). A clinical isolate of *Candida albicans* (gift from  
487 Prof Neil Gow), *Bacillus subtilis* 168 (from Prof Nicola Stanley Wall), a clinical isolate of  
488 Methicillin Resistant *Staphylococcus aureus* isolated from a patient at the Norfolk and Norwich  
489 University Hospital UK (70), an *Escherichia coli* K12 lab strain and the plant pathogen  
490 *Pseudomonas syringae* DC3000 (gift from Dr Jacob Malone) were grown overnight in 10 ml  
491 Lysogeny Broth (LB) at  $30^\circ\text{C}$ , 250 rpm. These were sub-cultured 1 in 20 (v/v) for a further 4  
492 hours at  $30^\circ\text{C}$  before being used to inoculate 100 ml of molten LB (0.5% agar), 3 ml of which  
493 was used to overlay each agar plate containing a *Streptomyces* colony. Plates were incubated  
494 for 48 hours at  $30^\circ\text{C}$ . Bioactivity was indicated by a clear halo around the *Streptomyces* colony.  
495 For bioassays using the fungal strains *Lomentospora prolificans* or *Gaeumannomyces graminis*  
496 var. *tritici* (Table 2), a plug of the fungus (grown for 14 days on potato glucose agar) was  
497 placed at the edge of the agar plate, 2 cm away from the growing streptomycete colony. Plates  
498 were incubated at  $25^\circ\text{C}$  for up to 14 days to assess inhibition of fungal growth. Bioassays were  
499 carried out on a range of different media, including minimal medium supplemented with  
500 Indole-3-Acetic Acid (IAA) (See Table S7 for media recipes).

501

502 **Purification and elucidation of filipin-like compounds from strain N2.** Spores of strain N2  
503 were spread onto 120 plates (4 L) of SFM agar and grown for eight days at  $30^\circ\text{C}$ . The resulting  
504 agar was then sliced into small pieces and extracted with ethyl acetate (3 x 1.5 L). An analytical  
505 sample was taken for analysis whereby the extract was filtered through gauze and the solvent  
506 removed under reduced pressure yielding 9.2 g of crude material. This was split into two halves  
507 and each treated identically: after dissolving in acetone (50 ml), loose normal phase silica gel  
508 (~30 g, Sigma Aldrich) was added and the solvent removed under reduced pressure. The  
509 impregnated silica gel was dry loaded onto a Biotage® SNAP Ultra cartridge (50 g, HP-Sphere  
510 normal phase silica). The resulting sample was chromatographed using a Biotage flash  
511 chromatography system to separate the polyene fraction (338 nm) and the actinomycin(s)  
512 containing fraction (444 nm; hereafter referred to as the 'actinomycin complex') using the  
513 following gradient with a flow rate of  $100 \text{ ml min}^{-1}$ : hold at 0% B for 1 CV; linear gradient 0-  
514 50% B over 10 CV; 50-100% B over 0.5 CV; and hold at 100 % B for 3 CV; (mobile phase A,  
515 chloroform; mobile phase B, methanol).



516 The polyene containing fractions were combined, the solvent removed under reduced  
517 pressure and the residues split into two fractions. Each fraction (in 800  $\mu$ L DMSO) was loaded  
518 onto a Biotage® SNAP Ultra cartridge (12 g, C<sub>18</sub>) and chromatography was achieved using the  
519 following gradient at a flow rate of 12 ml min<sup>-1</sup>: hold at 0% B for 5 CV; linear gradient 0-  
520 55% B over 1 CV, 55-85% B over 10 CV; 85-100% B over 2 CV; and hold at 100% B for  
521 1 CV; (mobile phase A, water; mobile phase B, methanol). The resulting fractions were  
522 analysed using LCMS and combined to yield three polyene samples of 53 mg, 34 mg and 9 mg  
523 after solvent was removed. LCMS spectra of these fractions and the original crude extract were  
524 then uploaded onto the GNPS (Global Natural Products Social Molecular Networking  
525 platform). The largest network containing the spectra of the filipin related compounds was  
526 manually adapted and processed in Cytoscape 3.6.1. The second fraction contained the most  
527 interesting compounds so only this sample was further purified by chromatography over a  
528 Synergi Fusion 4 micron C<sub>18</sub> 250 x 10 mm column (Phenomenex) using an Agilent 1100 series  
529 HPLC system fitted with a fraction collector and eluting at a flow rate of 3.5 ml min<sup>-1</sup> with the  
530 following gradient. 0-2 min, 45% B; 2-5 min, 45-50% B; 5-10 min, 50% B; 10.0-10.1 min, 50-  
531 45% B; 10.1-12.0 min, 45% B; (mobile phase A, 0.1% formic acid in water; mobile phase B,  
532 acetonitrile). This yielded pentamycin (4.2 mg) and 14-hydroxyisochainin (2.3 mg). Both  
533 structures were assigned using 2D NMR recorded on a Bruker Avance Neo 600 MHz  
534 spectrometer equipped with a helium-cooled cryoprobe and dissolved in DMSO-*d*<sub>6</sub>. Absolute  
535 structures were assigned based on the identical chemical shifts displayed in comparison to  
536 published data, and additional 1D-experiments were carried out in CD<sub>3</sub>OD for 14-  
537 hydroxyisochainin in order to compare directly with the published data for this compound (Figs  
538 S10-16 and Tables S5-8) (52).

539 Disc-diffusion bioassays were used to test whether purified pentamycin, 14-  
540 hydroxyisochainin and the actinomycin complex were active against the pathogenic strains *C.*  
541 *albicans* and *E. coli* (Table 2). Both indicator strains were grown in 10 ml of LB broth (Table  
542 S7) at 30°C and 200 rpm overnight. Cultures were then diluted 1 in 20 (v/v) into 10 ml of LB  
543 and grown for a further 4 hours. The 10 ml sub-culture was added to 50 ml of soft LB (0.5%  
544 agar) which was then poured into 10 cm square plates and allowed to set. Meanwhile, 6 mm  
545 sterile filter paper discs (Whatman) were inoculated with 40  $\mu$ l of each individual compound  
546 (three technical replicates of each compound were tested). 40  $\mu$ l of methanol was added to one  
547 disc per plate as a solvent control and 40  $\mu$ l of nystatin (5 mg ml<sup>-1</sup>) or hygromycin (50 mg ml<sup>-1</sup>)  
548 were used as positive controls on *C. albicans* and *E. coli* plates, respectively. Once dry, discs  
549 were placed onto plates 3 cm apart. These were then incubated at 30°C overnight. Inhibition

550 of the indicator strain was evidenced by a zone of clearing around the disc. Purified compounds  
551 were also tested for their ability to inhibit the wheat take-all fungus *G. graminis* (Table 2). For  
552 this, discs were placed onto PGA agar plates (Table S7) 2 cm away from an actively growing  
553 plug of *G. graminis*. Plates were left to grow at room temperature for 5 days before imaging.  
554 Three technical replicates were run for each purified compound (pentamycin, 14-  
555 hydroxyisochainin and actinomycin).

556

557 **Wheat seedling bioassays with *Streptomyces* strain N2.** Seeds of *Triticum aestivum* (var.  
558 Paragon, Table 2) were sterilized by placing them in 70% (v/v) ethanol for 2 minutes followed  
559 by a wash in 3% (v/v) NaOCl for 10 minutes. Seeds were then rinsed five times in sterile dH<sub>2</sub>O,  
560 before placing them into a solution of pregerminated spores (10<sup>7</sup> spores ml<sup>-1</sup>) of *Streptomyces*  
561 strain N2 (Table 2). Spores were pregerminated in 2xYT (Table S7) at 50°C for 10 minutes.  
562 Seeds were incubated in either the spore preparation or sterile 2xYT as a control, for 2 hours,  
563 before being allowed to dry in a Petri dish under sterile conditions. Two wheat seeds were then  
564 placed onto a 10 cm square plate of Msk agar (1.5% (w/v) agar, 0% (w/v) sucrose, Table S7),  
565 on either side of a plug of the *G. graminis* fungus, which was placed in the centre of the agar  
566 plate. Plugs were taken from an actively growing plate of *G. graminis* on PGA agar. Three  
567 replicate plates each of N2-coated seeds and sterile control seeds were used in each experiment.  
568 Plates were incubated for 5 days at room temperature after which inhibition of *G. graminis* was  
569 indicated by a zone of clearing around the wheat seeds.

570 A sterile vermiculite system was used to investigate the ability of *Streptomyces* strain  
571 N2 to protect older wheat seedlings against Take-all infection. 25 ml of sterile vermiculite was  
572 placed into the bottom of a 50 ml Falcon tube. Five plugs of *G. graminis* actively growing on  
573 PGA (Table S7), or uninoculated PGA plugs as a control, were placed on top of this layer,  
574 before being covered with a further 10 ml of vermiculite. Five seeds of *T. aestivum* (soaked in  
575 either N2 spores or uninoculated 2xYT, as described above) were then placed on top of this  
576 vermiculite layer, before the addition of a further 10 ml of vermiculite. The Falcon tubes were  
577 then sealed with parafilm and incubated at 25°C for 3 weeks, under a 12 hour light/ 12 hour  
578 dark photoperiod. There were five replicates tubes, each containing five replicate seeds, of each  
579 of the following combinations: PGA plugs with N2-coated seeds (wheat/*Streptomyces* control);  
580 *G. graminis* plugs with N2-coated seeds (wheat/*Streptomyces*/fungus treatment); PGA plugs  
581 with uninoculated seeds (wheat control); *G. graminis* plugs with uninoculated seeds  
582 (wheat/fungus treatment). After three weeks of incubation, plants were taken from the Falcon  
583 tubes and adhering vermiculite was removed from the roots. Take-all infection severity was

584 scored on a scale of 0-8, using an infection scoring system as follows: : 0=no infection;  
585 1=maximum of one lesion per root; 2=more than one lesion per root; 3=many small and at least  
586 one large lesion per root; 4=many large lesions per root; 5=roots completely brown; 6=roots  
587 completely brown plus infection in stem; 7=roots completely brown, infection in stem and  
588 wilted yellow leaves; 8=entire plant brown and wilted. Differences in infection score between  
589 treatments were analysed in R 3.2.3 (68) using a Kruskal-Wallis test, coupled with a post-hoc  
590 Dunn's multiple comparison test.

591

592 **Data Availability.** Genome accession numbers for the strains sequenced in this study are  
593 listed in Table 2.

594

595 **Acknowledgments.** SFW was funded by a Natural Environment Research Council (NERC)  
596 PhD studentship (NERC Doctoral Training Programme grant NE/L002582/1), JN was  
597 supported by a Biotechnology and Biological Sciences Research Council (BBSRC) PhD  
598 studentship (BBSRC Doctoral Training Programme grant BB/M011216/1), JR was funded by  
599 Leonardo Office Thuringia within the scope of Erasmus+ and the Deutschland stipendium,  
600 NAH was funded by NERC responsive mode grant NE/M015033/1 awarded to MIH, BW and  
601 JCM, and SFDB was funded by BBSRC responsive mode grant BB/P021506/1 to BW.  
602 Confocal microscopy was carried out at the Henry Wellcome Laboratory for Cell Imaging at  
603 UEA and we thank Dr Paul Thomas for training JN. This work was also supported by the  
604 Norwich Research Park (NRP) through a Science Links Seed Corn grant to MIH and JCM and  
605 by the NRP Earth and Life Systems Alliance (ELSA). We thank Anne Osbourn and Rachel  
606 Melton at the John Innes Centre (JIC) for assistance with the Take-all infection assays, Jacob  
607 Malone (JIC) for gifting us the *Pseudomonas* strain and Simon Orford (JIC) for the Paragon  
608 wheat seeds. Media recipes and other *Streptomyces* resources are available at ActinoBase.org.

## 609 **References**

- 610 1. Hutchings MI, Truman AW, Wilkinson B. 2019. Antibiotics: past, present and future.  
611 *Curr Opin Microbiol* 51:72-80.
- 612 2. Chater KF. 2016. Recent advances in understanding *Streptomyces*. *F1000Res* 5:2795.
- 613 3. van der Meij A, Worsley SF, Hutchings MI, van Wezel GP. 2017. Chemical ecology  
614 of antibiotic production by actinomycetes. *FEMS Microbiol Rev* 41:392-416.
- 615 4. Bentley SD, Chater KF, Cerdeno-Tarraga AM, Challis GL, Thomson NR, James KD,  
616 Harris DE, Quail MA, Kieser H, Harper D, Bateman A, Brown S, Chandra G, Chen  
617 CW, Collins M, Cronin A, Fraser A, Goble A, Hidalgo J, Hornsby T, Howarth S,  
618 Huang CH, Kieser T, Larke L, Murphy L, Oliver K, O'Neil S, Rabbinowitsch E,  
619 Rajandream MA, Rutherford K, Rutter S, Seeger K, Saunders D, Sharp S, Squares R,

620 Squares S, Taylor K, Warren T, Wietzorrek A, Woodward J, Barrell BG, Parkhill J,  
621 Hopwood DA. 2002. Complete genome sequence of the model actinomycete  
622 *Streptomyces coelicolor* A3(2). Nature 417:141-7.

623 5. Seipke RF, Kaltenpoth M, Hutchings MI. 2012. *Streptomyces* as symbionts: an  
624 emerging and widespread theme? FEMS Microbiol Rev 36:862-876.

625 6. Barke J, Seipke RF, Gruschow S, Heavens D, Drou N, Bibb MJ, Goss RJM, Yu DW,  
626 Hutchings MI. 2010. A mixed community of actinomycetes produce multiple  
627 antibiotics for the fungus farming ant *Acromyrmex octospinosus*. BMC Biol 8.

628 7. Kaltenpoth M. 2009. Actinobacteria as mutualists: general healthcare for insects?  
629 Trends Microbiol 17:529-535.

630 8. Kaltenpoth M, Goettler W, Dale C, Stubblefield JW, Herzner G, Roeser-Mueller K,  
631 Strohm E. 2006. ‘*Candidatus Streptomyces philanthi*’, an endosymbiotic  
632 streptomycete in the antennae of *Philanthus* digger wasps. Int J Syst Evol Microbiol  
633 56:1403-1411.

634 9. Hulcr J, Adams AS, Raffa K, Hofstetter RW, Klepzig KD, Currie CR. 2011. Presence  
635 and diversity of *Streptomyces* in *Dendroctonus* and sympatric bark beetle galleries  
636 across North America. Microb Ecol 61:759-68.

637 10. Viaene T, Langendries S, Beirinckx S, Maes M, Goormachtig S. 2016. *Streptomyces*  
638 as a plant's best friend? FEMS Microbiology Ecology:fiw119.

639 11. Schrey SD, Tarkka MT. 2008. Friends and foes: streptomycetes as modulators of  
640 plant disease and symbiosis. Antonie Van Leeuwenhoek 94:11-19.

641 12. Bulgarelli D, Rott M, Schlaeppi K, van Themaat EVL, Ahmadinejad N, Assenza F,  
642 Rauf P, Huettel B, Reinhardt R, Schmelzer E. 2012. Revealing structure and assembly  
643 cues for *Arabidopsis* root-inhabiting bacterial microbiota. Nature 488:91-95.

644 13. Lundberg DS, Lebeis SL, Paredes SH, Yourstone S, Gehring J, Malfatti S, Tremblay  
645 J, Engelbrektson A, Kunin V, Del Rio TG. 2012. Defining the core *Arabidopsis*  
646 *thaliana* root microbiome. Nature 488:86-90.

647 14. Bodenhausen N, Horton MW, Bergelson J. 2013. Bacterial communities associated  
648 with the leaves and the roots of *Arabidopsis thaliana*. PloS one 8:e56329.

649 15. Schlaeppi K, Dombrowski N, Oter RG, Ver Loren van Themaat E, Schulze-Lefert P.  
650 2014. Quantitative divergence of the bacterial root microbiota in *Arabidopsis thaliana*  
651 relatives. Proc Natl Acad Sci U S A 111:585-92.

652 16. Carvalhais LC, Dennis PG, Badri DV, Kidd BN, Vivanco JM, Schenk PM. 2015.  
653 Linking jasmonic acid signaling, root exudates, and rhizosphere microbiomes. Mol  
654 Plant Microbe Interact 28:1049-58.

655 17. Lebeis SL, Paredes SH, Lundberg DS, Breakfield N, Gehring J, McDonald M,  
656 Malfatti S, Del Rio TG, Jones CD, Tringe SG. 2015. Salicylic acid modulates  
657 colonization of the root microbiome by specific bacterial taxa. Science 349:860-864.

658 18. Coombs JT, Franco CM. 2003. Isolation and identification of actinobacteria from  
659 surface-sterilized wheat roots. Appl Environ Microbiol 69:5603-8.

660 19. Newitt JT, Prudence SMM, Hutchings MI, Worsley SF. 2019. Biocontrol of cereal  
661 crop diseases using streptomycetes. Pathogens 8.

662 20. Rey T, Dumas B. 2017. Plenty Is No Plague: *Streptomyces* Symbiosis with Crops.  
663 Trends Plant Sci 22:30-37.

664 21. Edwards J, Johnson C, Santos-Medellin C, Lurie E, Podishetty NK, Bhatnagar S,  
665 Eisen JA, Sundaresan V. 2015. Structure, variation, and assembly of the root-  
666 associated microbiomes of rice. Proc Natl Acad Sci U S A 112:E911-20.

667 22. Law JW, Ser HL, Khan TM, Chuah LH, Pusparajah P, Chan KG, Goh BH, Lee LH.  
668 2017. The potential of *Streptomyces* as biocontrol Agents against the Rice Blast  
669 Fungus, *Magnaporthe oryzae* (*Pyricularia oryzae*). Front Microbiol 8:3.

- 670 23. Kim DR, Cho G, Jeon CW, Weller DM, Thomashow LS, Paulitz TC, Kwak YS.  
671 2019. A mutualistic interaction between *Streptomyces* bacteria, strawberry plants and  
672 pollinating bees. *Nat Commun* 10:4802.
- 673 24. Lahdenperä ML, Simon E, Uoti J. 1991. Mycostop - a novel biofungicide based on  
674 *Streptomyces* bacteria, p 258-263. *In* Beemster ABR, Bollen GJ, Gerlagh M, Ruissen  
675 MA, Schippers B, Tempel A (ed), *Developments in Agricultural and Managed Forest*  
676 *Ecology*, vol 23. Elsevier.
- 677 25. Holmes NA, Innocent TM, Heine D, Bassam MA, Worsley SF, Trottmann F, Patrick  
678 EH, Yu DW, Murrell JC, Schiott M, Wilkinson B, Boomsma JJ, Hutchings MI. 2016.  
679 Genome analysis of two *Pseudonocardia* phylotypes associated with *Acromyrmex*  
680 leafcutter ants reveals their biosynthetic potential. *Front Microbiol* 7:2073.
- 681 26. Tokala RK, Strap JL, Jung CM, Crawford DL, Salove MH, Deobald LA, Bailey JF,  
682 Morra M. 2002. Novel plant-microbe rhizosphere interaction involving *Streptomyces*  
683 *lydicus* WYEC108 and the pea plant (*Pisum sativum*). *Appl Environ Microbiol*  
684 68:2161-2171.
- 685 27. Yuan WM, Crawford DL. 1995. Characterization of *Streptomyces lydicus* WYEC108  
686 as a potential biocontrol agent against fungal root and seed rots. *Appl Environ*  
687 *Microbiol* 61:3119-3128.
- 688 28. Alanjary M, Steinke K, Ziemert N. 2019. AutoMLST: an automated web server for  
689 generating multi-locus species trees highlighting natural product potential. *Nucleic*  
690 *Acids Res* doi:10.1093/nar/gkz282.
- 691 29. Jain C, Rodriguez RL, Phillippy AM, Konstantinidis KT, Aluru S. 2018. High  
692 throughput ANI analysis of 90K prokaryotic genomes reveals clear species  
693 boundaries. *Nat Commun* 9:5114.
- 694 30. Konstantinidis KT, Tiedje JM. 2005. Genomic insights that advance the species  
695 definition for prokaryotes. *Proc Nat Acad Sci USA* 102:2567.
- 696 31. Blin K, Shaw S, Steinke K, Villebro R, Ziemert N, Lee SY, Medema MH, Weber T.  
697 2019. antiSMASH 5.0: updates to the secondary metabolite genome mining pipeline.  
698 *Nucleic Acids Res* doi:10.1093/nar/gkz310.
- 699 32. Kumar J, Sharma VK, Singh DK, Mishra A, Gond SK, Verma SK, Kumar A,  
700 Kharwar RN. 2016. Epigenetic Activation of Antibacterial Property of an Endophytic  
701 *Streptomyces coelicolor* Strain AZRA 37 and Identification of the Induced Protein  
702 Using MALDI TOF MS/MS. *PLoS One* 11:e0147876.
- 703 33. Bradáčová K, Kandeler E, Berger N, Ludewig U, Neumann G. 2020. Microbial  
704 consortia inoculants stimulate early growth of maize depending on nitrogen and  
705 phosphorus supply. *Plant, Soil and Environment* 66:105-112.
- 706 34. Egamberdieva D, Jabborova D, Berg G. 2016. Synergistic interactions between  
707 *Bradyrhizobium japonicum* and the endophyte *Stenotrophomonas rhizophila* and their  
708 effects on growth, and nodulation of soybean under salt stress. *Plant and Soil* 405:35-  
709 45.
- 710 35. Jha CK, Saraf M. 2012. Evaluation of Multispecies Plant-Growth-Promoting  
711 Consortia for the Growth Promotion of *Jatropha curcas* L. *Journal of Plant Growth*  
712 *Regulation* 31:588-598.
- 713 36. Bric JM, Bostock RM, Silverstone SE. 1991. Rapid *in situ* assay for indoleacetic acid  
714 production by bacteria immobilized on a nitrocellulose membrane. *Appl Environ*  
715 *Microbiol* 57:535-538.
- 716 37. Glick BR. 2014. Bacteria with ACC deaminase can promote plant growth and help to  
717 feed the world. *Microbiol Res* 169:30-9.

- 718 38. Palaniyandi SA, Damodharan K, Yang SH, Suh JW. 2014. *Streptomyces* sp. strain  
719 PGPA39 alleviates salt stress and promotes growth of 'Micro Tom' tomato plants. *J*  
720 *Appl Microbiol* 117:766-73.
- 721 39. Hopwood DA. 2007. *Streptomyces* in nature and medicine: the antibiotic makers.  
722 Oxford University Press, New York.
- 723 40. Scavino AF, Pedraza RO. 2013. The role of siderophores in plant growth-promoting  
724 bacteria, p 265-285, *Bacteria in agrobiolgy: crop productivity*. Springer.
- 725 41. Kloepper JW, Leong J, Teintze M, Schroth MN. 1980. Enhanced plant growth by  
726 siderophores produced by plant growth-promoting rhizobacteria. *Nature* 286:885-886.
- 727 42. Katz E. 1967. Actinomycin, p 276-341. *In* Gottlieb D, Shaw PD (ed), *Biosynthesis*  
728 doi:10.1007/978-3-662-38441-1\_25. Springer Berlin Heidelberg, Berlin, Heidelberg.
- 729 43. Luo Y, Huang H, Liang J, Wang M, Lu L, Shao Z, Cobb RE, Zhao H. 2013.  
730 Activation and characterization of a cryptic polycyclic tetramate macrolactam  
731 biosynthetic gene cluster. *Nat Commun* 4:2894.
- 732 44. Bonneau L, Gerbeau-Pissot P, Thomas D, Der C, Lherminier J, Bourque S, Roche Y,  
733 Simon-Plas F. 2010. Plasma membrane sterol complexation, generated by filipin,  
734 triggers signaling responses in tobacco cells. *Biochim Biophys Acta* 1798:2150-9.
- 735 45. Milhaud J, Bolard J, Benveniste P, Hartmann MA. 1988. Interaction of the polyene  
736 antibiotic filipin with model and natural membranes containing plant sterols. *Biochim*  
737 *Biophys Acta* 943:315-25.
- 738 46. Moeller CH, Mudd JB. 1982. Localization of filipin-sterol complexes in the  
739 membranes of *Beta vulgaris* roots and *Spinacia oleracea* chloroplasts. *Plant Physiol*  
740 70:1554-61.
- 741 47. Duca D, Lorv J, Patten CL, Rose D, Glick BR. 2014. Indole-3-acetic acid in plant-  
742 microbe interactions. *Antonie Van Leeuwenhoek* 106:85-125.
- 743 48. Patten CL, Blakney AJ, Coulson TJ. 2013. Activity, distribution and function of  
744 indole-3-acetic acid biosynthetic pathways in bacteria. *Crit Rev Microbiol* 39:395-  
745 415.
- 746 49. Daddaoua A, Matilla MA, Krell T, Chini A, Morel B. 2018. An auxin controls  
747 bacterial antibiotics production. *Nucleic Acids Res* 46:11229-11238.
- 748 50. Cope AC, Johnson HE. 1958. Fungichromin. Determination of the structure of the  
749 pentaene chromophore. *J Am Chem Soc* 80:1504-1506.
- 750 51. Zhou S, Song L, Masschelein J, Sumang FAM, Papa IA, Zulaybar TO, Custodio AB,  
751 Zabala D, Alcantara EP, de Los Santos ELC, Challis GL. 2019. Pentamycin  
752 biosynthesis in Philippine *Streptomyces* sp. S816: cytochrome P450-catalyzed  
753 installation of the C-14 hydroxyl group. *ACS Chem Biol* 14:1305-1309.
- 754 52. Li Z, Rawlings BJ, Harrison PH, Vederas JC. 1989. Production of new polyene  
755 antibiotics by *Streptomyces cellulosae* after addition of ethyl (Z)-16-phenylhexadec-  
756 9-enoate. *J Antibiot (Tokyo)* 42:577-84.
- 757 53. Gao H, Gruschow S, Barke J, Seipke RF, Hill LM, Orivel J, Yu DW, Hutchings M,  
758 Goss RJM. 2014. Filipins: the first antifungal “weed killers” identified from bacteria  
759 isolated from the trap-ant. *RSC Advances* 4:57267-57270.
- 760 54. Payero TD, Vicente CM, Rumbero A, Barreales EG, Santos-Aberturas J, de Pedro A,  
761 Aparicio JF. 2015. Functional analysis of filipin tailoring genes from *Streptomyces*  
762 *filipinensis* reveals alternative routes in filipin III biosynthesis and yields bioactive  
763 derivatives. *Microb Cell Fact* 14:114.
- 764 55. Kwan DH, Schulz F. 2011. The stereochemistry of complex polyketide biosynthesis  
765 by modular polyketide synthases. *Molecules* 16:6092-115.

- 766 56. Keller U, Lang M, Crnovcic I, Pfennig F, Schauwecker F. 2010. The actinomycin  
767 biosynthetic gene cluster of *Streptomyces chrysomallus*: a genetic hall of mirrors for  
768 synthesis of a molecule with mirror symmetry. *J Bacteriol* 192:2583-95.
- 769 57. Frey Tirri B, Bitzer J, Geudelin B, Drewe J. 2010. Safety, tolerability and  
770 pharmacokinetics of intravaginal pentamycin. *Chemotherapy* 56:190-6.
- 771 58. Kranzler M, Syrowatka M, Leitsch D, Winnips C, Walochnik J. 2015. Pentamycin  
772 shows high efficacy against *Trichomonas vaginalis*. *Int J Antimicrob Agents* 45:434-  
773 7.
- 774 59. Xiong ZQ, Zhang ZP, Li JH, Wei SJ, Tu GQ. 2012. Characterization of *Streptomyces*  
775 *padanus* JAU4234, a producer of actinomycin X(2), fungichromin, and a new polyene  
776 macrolide antibiotic. *Appl Environ Microbiol* 78:589-92.
- 777 60. Dessaux Y, Grandclement C, Faure D. 2016. Engineering the rhizosphere. *Trends*  
778 *Plant Sci* 21:266-278.
- 779 61. Ryan PR, Dessaux Y, Thomashow LS, Weller DM. 2009. Rhizosphere engineering  
780 and management for sustainable agriculture. *Plant and Soil* 321:363-383.
- 781 62. Hoitink HAJ, Boehm MJ. 1999. Biocontrol within the context of soil microbial  
782 communities: a substrate-dependent phenomenon. *Annu Rev Phytopathol* 37:427-446.
- 783 63. Scheuring I, Yu DW. 2012. How to assemble a beneficial microbiome in three easy  
784 steps. *Ecol Lett* 15:1300-1307.
- 785 64. Kieser T, Bibb MJ, Buttner MJ, Chater KF, Hopwood DA. 2000. Practical  
786 *Streptomyces* Genetics. John Innes Foundation, Norwich.
- 787 65. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes  
788 S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA,  
789 McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R,  
790 Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST Server: rapid  
791 annotations using subsystems technology. *BMC Genomics* 9:75.
- 792 66. Moriya Y, Itoh M, Okuda S, Yoshizawa AC, Kanehisa M. 2007. KAAS: an automatic  
793 genome annotation and pathway reconstruction server. *Nucleic Acids Res* 35:W182-  
794 5.
- 795 67. Sun J, Kelemen GH, Fernandez-Abalos JM, Bibb MJ. 1999. Green fluorescent protein  
796 as a reporter for spatial and temporal gene expression in *Streptomyces coelicolor*  
797 A3(2). *Microbiology* 145 ( Pt 9):2221-7.
- 798 68. R Core Team. 2017. R: A language and environment for statistical computing.  
799 Accessed
- 800 69. Dworkin M, Foster JW. 1958. Experiments with some microorganisms which utilize  
801 ethane and hydrogen. *J Bacteriol* 75:592-603.
- 802 70. Qin Z, Munnoch JT, Devine R, Holmes NA, Seipke RF, Wilkinson KA, Wilkinson B,  
803 Hutchings MI. 2017. Formicamycins, antibacterial polyketides produced by  
804 *Streptomyces formicae* isolated from African *Tetraponera* plant-ants. *Chemical*  
805 *Science*.

807

808 **Table 1.** Genome features of root-associated *Streptomyces* strains sequenced for this study,  
 809 including their GenBank Accession numbers, their genome size in base pairs (bp) and the total  
 810 number of open reading frames (ORFs) per genome. Genomes were sequenced using the  
 811 PacBio RSII platform. Biosynthetic Gene Clusters (BGCs) were predicted using AntiSMASH  
 812 5.0 (31) and % ANI and closest relatives were determined using AutoMLST (28).

Strain	Accession no.	Size (bp)	ORFs	BGCs	Closest relative / ANI
L2	QBDT00000000	8,073,926	7079	29	<i>Streptomyces bungoensis</i> (88.3%)
M2	CP028834	8,718,751	8026	22	<i>Streptomyces</i> sp. HBG00200 (94.7%)
M3	QANR00000000	8,304,843	7561	25	<i>Streptomyces pratensis</i> (91.1%)
N1	QBDS00000000	7,207,104	6239	21	<i>Streptomyces albidoflavus</i> (98.7%)
N2	CP028719	8,428,700	7401	34	<i>Streptomyces griseofuscus</i> (97.6%)
Actinovate	RDTC00000000	9,139,876	7989	33	<i>S. lydicus</i>
ATCC25470	RDTD00000000	7,935,716	7084	25	<i>S. lydicus</i>
ATCC31975	RDTE00000000	9,244,118	8128	31	<i>S. lydicus</i>

813

814 Table 2. Strains and primers used in experiments.

Species/strain name	Description	Origin	Genome accession number
<i>Streptomyces</i> L2	Wild-type	<i>A. thaliana</i> root microbiome, this study	QBDT00000000




<i>Streptomyces</i> M2	Wild-type	<i>A.thaliana</i> root microbiome, this study	CP028834
<i>Streptomyces</i> M3	Wild-type	<i>A.thaliana</i> root microbiome, this study	QANR00000000
<i>Streptomyces</i> N1	Wild-type	<i>A.thaliana</i> root microbiome, this study	QBDS00000000
<i>Streptomyces</i> N2	Wild-type	<i>A.thaliana</i> root microbiome, this study	CP028719
<i>Streptomyces lydicus</i> ATCC25470	Wild-type	American Type Culture Collection	RDTD00000000
<i>Streptomyces lydicus</i> ATCC31975	Wild-type	American Type Culture Collection	RDTE00000000
<i>Streptomyces lydicus</i> Actinovate	Wild-type	Isolated from Actinovate™ by Elaine Patrick, UEA	RDTC00000000
<i>Bacillus subtilis</i>	Wild-type, strain 168	Gift from Nicola Stanley-Wall, University of Dundee	NA
Methicillin resistant <i>Staphylococcus aureus</i>	Clinical isolate	Norfolk and Norwich University Hospital (UK)	NA
<i>Escherichia coli</i> K12	Wild-type	Lab stock, UEA	NA
<i>Pseudomonas syringae</i> DC3000	Wild-type	John Innes Centre, Norwich, UK	NA
<i>Candida albicans</i>	Clinical isolate	Gift from Neil Gow, University of Exeter	NA
<i>Lomentospora prolificans</i>	Environmental isolate	American Type Culture Collection	NA
<i>Gaeumannomyces graminis</i> var. <i>tritici</i>	Environmental isolate	John Innes Centre, Norwich, UK	NA

<i>Arabidopsis thaliana</i> Col-0	Wild-type, ecotype Col-0	Lab stock, UEA	NA
<i>Triticum aestivum</i> var. Paragon	Wild-type, var. Paragon	John Innes Centre, Norwich, UK	NA
<b>Primer name</b>	<b>Sequence</b>		<b>Reference</b>
PRK341F	5'-CCTACGGGAGGCAGCAG-3'		Yu et al 2005
MPRK806R	5'-GGACTACHVGGGTWTCTAAT-3'		
<b>Plasmid name</b>	<b>Description</b>		<b>Reference</b>
pIJ8660	ermEp* driving constitutive production of codon optimised eGFP		Sun et al 1999

815


816



818

819 **Figure 1.** Confocal Laser Scanning Microscopy images of *A. thaliana* rhizoplane colonisation  
 820 by eGFP-tagged *Streptomyces* strains three days after inoculation. A and B show *A. thaliana*  
 821 roots (red) colonised by eGFP-tagged *Streptomyces coelicolor* M145 (green) which is a known  
 822 root endophyte and was used as a control (32). C and D show *A. thaliana* roots (red) colonised  
 823 by eGFP-tagged *Streptomyces* strain M3 which was isolated in this study (green).

824



825

826

827

828

829

830


831

832

833


834

**Figure 2.** Violin plots showing the biomass of *Arabidopsis thaliana* plants grown on agar plates, following inoculation with sequenced *Streptomyces* isolates. Biomass (dry weight in grams) was measured 16 days after inoculation. Sterile plants were grown as a control. N=16 plants per treatment. Box plots show the location of the median and quartiles, with whiskers reaching to 1.5 times the interquartile range. Blue diamonds represent mean values. The width of the outer shaded area illustrates the proportion of the data located there (the kernel probability density). Groups labelled with different letters have a significantly different plant biomass ( $P < 0.05$  in Tukey's HSD tests).



835

836 **Figure 3.** Violin plots demonstrating the total dry weight of *A. thaliana* plants grown in  
 837 Levington's F2 compost from seeds inoculated with spores of Actinovate, L2, M2, M3, N1  
 838 and N2 or a mixture of L2, M2 and M3 *Streptomyces* spores. Dry weight is shown in  
 839 milligrams. Sterile seeds were grown as a control. N=8 replicate plants per treatment. Box  
 840 plots show the location of the median and quartiles, with whiskers reaching to 1.5 times the  
 841 interquartile range. Blue diamonds represent mean values. The width of the outer shaded area  
 842 illustrates the proportion of data located there (the kernel probability density). Groups  
 843 labelled with different letters differ significantly in plant biomass ( $P < 0.05$  in Tukey's HSD  
 844 tests).



845

846

847

848

849




850

851

**Figure 4:** (A) Three biological replicates of strain N2 (centre) growing on minimal medium agar that has been overlaid with soft LB agar inoculated with *Candida albicans*. (B) is the same but the minimal medium agar contains 0.1 mg ml<sup>-1</sup> of indole-3-acetic acid. This experiment was repeated 4 times (each time with 3 replicates), with consistent results. Scale bars represent 2 cm.

852

853




854

855

856 **Figure 5.** Structures of filipin III, pentamycin and 14-hydroxyisochainin.

857





858

859 **Figure 6. Inhibition of *G. graminis* var. *tritici* in wheat seedlings.** Germinating wheat seeds  
860 (marked as “W”) are either A) sterile or B) inoculated with a spore preparation of *Streptomyces*  
861 isolate N2, growing next to a plug of *G. graminis* var. *tritici*, the Take-all fungus (marked as  
862 “P”). *G. graminis* is prevented from growing towards inoculated seeds, as demonstrated by the  
863 zone of inhibition (marked with arrowheads). Scale bars represent 5 mm.

864

865





866

867

868 **Fig. 7. Left:** The effect of *Streptomyces* strain N2 on wheat plant infection severity by *G.*  
 869 *graminis* var. *tritici*. Infections were scored after three weeks of growth. N2 control= seeds  
 870 coated in N2 spores/no *G. graminis*; Sterile control= sterile seeds/no *G. graminis*;  
 871 N2/fungus= seeds coated in N2 spores grown in the presence of *G. graminis*; Sterile  
 872 seeds/fungus= sterile seeds grown in the presence of *G. graminis*. N= 25 plants per treatment

873 group, error bars represent standard errors. Right: Wheat plants were grown A) from sterile  
874 seeds in the presence of *G. graminis* var. *tritici*, B) from seeds inoculated with N2 spores, in  
875 the presence of *G. graminis* var. *tritici* C) from sterile seeds, no *G. graminis* var. *tritici* D)  
876 from seeds coated with N2 spores, no *G. graminis* var. *tritici*. Scale bars represent 2 cm.

877