

Journal Pre-proof

The collagen synthesis response to an acute bout of resistance exercise is greater when ingesting 30 g versus 15 g and 0 g hydrolyzed collagen in resistance-trained young men

Joonsung Lee, Jonathan C.Y. Tang, John Dutton, Rachel Dunn, William D. Fraser, Kevin Enright, David R. Clark, Claire E. Stewart, Robert M. Erskine

PII: S0022-3166(23)72738-7

DOI: <https://doi.org/10.1016/j.tjnut.2023.10.030>

Reference: TJNUT 417

To appear in: *The Journal of Nutrition*

Received Date: 19 July 2023

Revised Date: 27 September 2023

Accepted Date: 26 October 2023

Please cite this article as: J. Lee, J.C.Y. Tang, J. Dutton, R. Dunn, W.D. Fraser, K. Enright, D.R. Clark, C.E. Stewart, R.M. Erskine, The collagen synthesis response to an acute bout of resistance exercise is greater when ingesting 30 g versus 15 g and 0 g hydrolyzed collagen in resistance-trained young men, *The Journal of Nutrition*, <https://doi.org/10.1016/j.tjnut.2023.10.030>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2023 The Author(s). Published by Elsevier Inc. on behalf of American Society for Nutrition.



The collagen synthesis response to an acute bout of resistance exercise is greater when ingesting 30 g versus 15 g and 0 g hydrolyzed collagen in resistance-trained young men

Joonsung Lee¹, Jonathan C. Y. Tang^{2,3}, John Dutton², Rachel Dunn^{2,3}, William D. Fraser^{2,3,4}, Kevin Enright¹, David R. Clark⁵, Claire E. Stewart¹, Robert M. Erskine*^{1,6}

¹School of Sport and Exercise Sciences, Liverpool John Moores University, Liverpool, UK;

²Bioanalytical Facility, Norwich Medical School, University of East Anglia, Norwich, UK

³Clinical Biochemistry, Departments of Laboratory Medicine, Norfolk and Norwich University Hospital NHS Foundation Trust, Norwich, UK.

⁴Departments of Diabetes and Endocrinology, Norfolk and Norwich University Hospital NHS Foundation Trust, Norwich, UK

⁵School of Health Sciences, Robert Gordon University, Aberdeen, UK

⁶Institute of Sport, Exercise and Health, University College London, London, UK.

Running title: Resistance exercise and the collagen dose-response

Data availability

Data described in the article, code book, and analytic code will be made available upon request from the corresponding author pending application and approval.

***Correspondence:**

Dr Rob Erskine, School of Sport and Exercise Sciences, Liverpool John Moores University, Liverpool, L3 3AF, UK.

Email: R.M.Erskine@ljmu.ac.uk;

J. Lee ORCID: [0000-0001-8248-3066](https://orcid.org/0000-0001-8248-3066)

J. C. Y. Tang ORCID: [0000-0001-6305-6333](https://orcid.org/0000-0001-6305-6333)

W. D. Fraser ORCID: [0000-0003-0556-3358](https://orcid.org/0000-0003-0556-3358)

K. Enright ORCID: [0000-0003-1775-6392](https://orcid.org/0000-0003-1775-6392)

C. E. Stewart ORCID: [0000-0002-8104-4819](https://orcid.org/0000-0002-8104-4819)

R. M. Erskine ORCID: [0000-0002-5705-0207](https://orcid.org/0000-0002-5705-0207)

Keywords: strength training, vitamin C, connective tissue, glycine, proline, hydroxyproline

Abbreviations

AUC, area under the curve

β-CTX, β-isomerized C-terminal telopeptide of type I collagen

CI, confidence interval

CV, coefficient of variation

ELISA, enzyme-linked immunosorbent assay

HFBA, heptafluorobutyric acid

η_p^2 , partial eta squared

FSR, fractional synthetic rate

HC, hydrolysed collagen

IGF-I, insulin-like growth factor-1

LC-MS-MS, liquid chromatography with tandem mass spectrometry

mTORC1, mammalian target of rapamycin complex 1

MTU, muscle-tendon unit

PICP, procollagen type I C-terminal propeptide

PINP, procollagen type I N-terminal propeptide

QC, quality controls

RE, resistance exercise

TGF-β, transforming growth factor beta

10-RM, 10 repetition maximum

Abstract

1 *Background:* Resistance exercise (RE) stimulates collagen synthesis in skeletal muscle and
2 tendon but there is limited and equivocal evidence regarding an effect of collagen
3 supplementation and exercise on collagen synthesis. Furthermore, it is not known if a dose-
4 response exists regarding the effect of hydrolyzed collagen (HC) ingestion and RE on collagen
5 synthesis.

6 *Objective:* We aimed to determine the HC dose-response effect on collagen synthesis following
7 high-intensity RE in resistance-trained young men.

8 *Methods:* Using a double-blind, randomized cross-over design, 10 resistance-trained men (age:
9 26 ± 3 years; height: 1.77 ± 0.04 m; mass: 79.7 ± 7.0 kg) ingested 0g, 15g or 30g HC with 50mg
10 vitamin C 1h prior to performing four sets' barbell back-squat RE at 10-repetition maximum
11 load, after which they rested for six hours. Blood samples were collected throughout each of
12 the three interventions to analyse procollagen type I N-terminal propeptide (PINP) and β -
13 isomerized C-terminal telopeptide of type I collagen (β -CTX) concentration, and the
14 concentration of 18 collagen amino acids.

15 *Results:* The serum PINP concentration \times time area-under-the-curve (AUC) was greater for 30g
16 (267 ± 79 $\mu\text{g}\cdot\text{L}^{-1}\cdot\text{h}$) than 15g (235 ± 70 $\mu\text{g}\cdot\text{L}^{-1}\cdot\text{h}$, $P=0.039$) and 0g HC (219 ± 88 $\mu\text{g}\cdot\text{L}^{-1}\cdot\text{h}$, $P=0.005$)
17 but there was no difference between 0g and 15g HC ($P=0.675$). The AUCs of glycine and
18 proline were greater for 30g than for 15g and 0g HC ($P<0.05$). Plasma β -CTX concentration
19 decreased from -1h to +6h ($P<0.05$), with no differences between interventions.

20 *Conclusion:* The greater PINP AUC suggests 30g HC ingested prior to high-intensity RE
21 augments whole body collagen synthesis more than 15g and 0g HC in resistance-trained young
22 men.

23 **Introduction**

24 Musculoskeletal tissues, such as skeletal muscle and tendon are crucial for generating
25 and transmitting force to the bone, enabling movement. The structure and function of these
26 tissues are therefore essential for musculoskeletal health and physical performance.
27 Unfortunately, however, injuries to these tissues are common in athletes, with soft-tissue
28 injuries making up the majority of all injuries in male athletes (1). One of the biggest risk
29 factors for soft-tissue injury is muscle weakness (2, 3) and one of the most common methods
30 used by athletes to mitigate this risk factor is to perform chronic resistance exercise (RE) (4).
31 Chronically overloading the muscle-tendon unit in this way causes the muscle to adapt by
32 hypertrophy and getting stronger (5), while the tendon also adapts by hypertrophy and
33 increasing its stiffness and elastic modulus (6, 7).

34 A stiffer tendon has a higher loading capacity, as there is a linear relationship between
35 Young's modulus and ultimate stress (8). Given that collagen (mainly type I) makes up 60–85%
36 tendon dry weight (9), it is considered a crucial component in the tendon's adaptation to RE,
37 particularly as gains in tendon stiffness are thought to be influenced by both tendon
38 hypertrophy and an increase in collagen fibril density (10, 11). An increase in type I collagen
39 fibril content over time is likely the product of an overload-induced increase in collagen
40 synthesis after each bout of RE.

41 Collagen synthesis can be assessed either directly from the overloaded tissue, e.g. by
42 measuring skeletal muscle or tendon collagen fractional synthetic rate (FSR), or indirectly from
43 serum concentration of procollagen type I C-terminal propeptide (PICP) or procollagen type I
44 N-terminal propeptide (PINP), which are both cleaved off during the maturation of procollagen
45 to collagen. Indeed, an acute bout of RE in young men has been shown to increase patellar

46 tendon collagen FSR (12) and serum PINP concentration (13). This response is likely due to
47 RE initiating mechanotransduction (i.e. mechanical stress initiating fibroblast intracellular
48 signalling) (14), and the secretion of growth factors [e.g. transforming growth factor beta
49 (TGF- β) and insulin-like growth factor-1 (IGF-I)], with these growth factors being crucial for
50 procollagen formation in tendon (15, 16). Furthermore, these newly synthesized procollagen
51 molecules undergo post-translational modifications, for which the presence of vitamin C is an
52 essential co-factor during collagen synthesis (17), transport and assembly into tendon (18).

53 Thus, RE appears crucial for inducing increases in serum PICP/PINP concentration and
54 muscle-tendon collagen FSR, which may lead to changes in connective tissue properties in the
55 longer term. Indeed, concomitant increases in serum PICP concentration and human Achilles
56 tendon collagen content after two months' chronic RE, followed by an increase in Achilles
57 tendon stiffness with a further month's RE training (19), suggest that augmented tendon
58 collagen synthesis and content are necessary to cause an increase tendon stiffness. Not only is
59 tendon stiffness important for mitigating soft-tissue injury risk but it can also influence
60 performance during 'explosive' actions, as a stiffer tendon can transmit muscle force more
61 effectively to the bone, thus increasing the rate of force development (20).

62 In addition to the role of exercise on collagen synthesis, ingestion of vitamin C-enriched
63 collagen appears to further augment an exercise-induced increase in collagen synthesis in a
64 dose-response manner, i.e. 15 g gelatin increased serum PINP concentration by more than two-
65 fold compared to 5 g and 0 g gelatin (21). This is currently the only study to investigate a dose-
66 response effect of collagen ingestion on changes in collagen synthesis following exercise, albeit
67 in jump-rope exercise not RE. Although no study has examined a collagen dose-response
68 relationship without exercise, chronic collagen supplementation alone has been shown to

69 induce improvements in bone mineral density (22) and cartilage health (23), suggesting
70 collagen ingestion might stimulate human connective tissue collagen synthesis independently
71 of exercise. Thus, just as ingestion of 40 g whey protein has been shown to augment the muscle
72 protein synthesis response to RE more than 20 g (24), it is possible that collagen ingestion may
73 further augment the RE-induced rise in collagen synthesis (12, 13) in a dose-response manner.

74 The aim of this study was therefore to investigate the effect of 30 g vs. 15 g vs. 0 g HC
75 ingested prior to high-intensity back squat RE on whole body collagen synthesis. We
76 hypothesized that 30 g HC would elicit a greater serum PINP response than 15 g HC, which
77 would induce a greater response than 0 g HC. We also hypothesized that 30 g HC would lead
78 to a greater blood availability of the amino acids necessary for collagen synthesis to occur, e.g.
79 glycine and proline.

80

81 **Methods**

82 **Participants**

83 Thirteen healthy young men volunteered to take part in the study. However, three were
84 excluded prior to participation due to not meeting the inclusion criteria ($n = 1$) and declining
85 to proceed with participation ($n = 2$) (**Figure 1**). Therefore, 10 resistance-trained, healthy
86 young men (mean \pm SD; age: 26 ± 3 years, height: 1.77 ± 0.04 m, body mass: 79.7 ± 7.0 kg, 4
87 ± 3 years' RE experience), who performed RE 4 ± 1 times per week, provided written informed
88 consent before completing this study. The study was registered at <https://clinicaltrials.gov/>
89 (identifier: NCT05932771), was approved by Liverpool John Moores University Ethics
90 Committee (approval number: 18/SPS/059) and complied with the Declaration of Helsinki.
91 Participants were recruited from a university student population and recruitment began in

92 January 2019 and data collection was completed in August 2019. To be eligible to participate,
93 volunteers had to be male, have at least 12 months' resistance training experience (including
94 barbell back squat exercise performed at least once a week) and to be free from musculoskeletal
95 injury. Volunteers were excluded if they had a history of patellar tendon pathology, were vegan
96 (due to the bovine source of HC), consumed nutritional supplements or medication purported
97 to have beneficial effects on muscle-tendon properties (e.g. antioxidants, protein, etc.), had
98 sustained a lower limb injury in the previous six months, smoked or were <18 or >30 years old.

99 *Insert Figure 1 near here*

100 **Experimental design**

101 This study was a double-blind, randomized cross-over design. All participants attended the
102 laboratory on four occasions, each separated by a week, and no strenuous physical activity was
103 performed 48 h prior to each visit. Visit 1 was used to assess barbell back squat 10-RM; while
104 visits 2-4 began with participants consuming a drink containing 0 g, 15 g or 30 g HC (each
105 containing 50 mg vitamin C), followed by four sets' 10-RM barbell back squat RE (with 2 min
106 rest in between sets), which typically took 20 min to complete. The three interventions (with a
107 seven-day wash-out period interspersed between each intervention) were performed at the same
108 time of day (08:00 – 15:00), following a 10 h overnight fast. After consuming the supplement
109 and completing the RE, participants rested for 6 h and 10 × 5-mL blood samples were collected
110 at different times points over a 7 h period (**Figure 2**). In addition to the supplement, only water
111 was allowed to be consumed (ad libitum) during each intervention. Participants were instructed
112 to record their dietary intake on the day before their first intervention and to replicate that
113 dietary behaviour on the day preceding each of the subsequent interventions.

114 *Insert Figure 2 near here*

115 **The barbell back squat 10-repetition maximum (RM) assessment and 10-RM bout**
116 **during each intervention**

117 The squat depth during the barbell back squat was standardized for all participants to induce
118 the same mechanical loading on the quadriceps femoris muscle-tendon unit during all three
119 experimental interventions. Participants were instructed to place a 20 kg Olympic barbell on
120 their shoulders (the high bar position), place their feet shoulder-width apart (foot location was
121 marked on the floor for subsequent sets) and descend until their knee joint angle reached 90°,
122 measured using a goniometer. While participants held the position at 90° knee flexion, the
123 vertical distance from the floor to the ischial tuberosity was measured. The 10-RM assessment
124 was performed in a squat rack and a resistance band was stretched across both sides of the squat
125 rack to indicate the participant's 90° depth (**Figure 3**). A warm-up comprised two dynamic
126 exercises (low lunge and squat to stand) prior to the actual 10-RM assessment, which
127 comprised the following sets of barbell back squat: 10 repetitions with the 20 kg barbell, 8
128 repetitions at 50% of the estimated 10-RM, 4 repetitions at 70% and 1 repetition at 90% of the
129 estimated 10-RM). After a 5-min rest period, participants performed 10-RM attempts separated
130 by 5-min rest periods until 10-RM load was obtained. Two researchers observed each test
131 procedure to provide a cue when the participant's proximal hamstrings/gluteus maximus
132 touched the elastic band and to spot the participant. The 10-RM bout during each experimental
133 intervention was preceded by a similar warm-up, i.e. two dynamic exercises followed by 10
134 repetitions' barbell back squat with the 20 kg barbell, 8 repetitions at 50% of the measured 10-
135 RM, 4 repetitions at 70% 10-RM and 1 repetition at 90% 10-RM. The barbell back squat 10-
136 RM load was 118 ± 21 kg during all three interventions.

137 *Insert Figure 3 near here*

138 Nutritional supplementation

139 Before commencing each intervention, a laboratory technician (independent to the study) made
140 up the supplement and randomly assigned the order of HC dose (Excel 2016, Microsoft,
141 Washington, USA) for each participant. For each intervention, the technician recorded the date,
142 randomly allocated intervention number (1, 2 or 3) and corresponding HC dose. The study
143 investigators and participants were blinded to HC dose until after all analyses were completed,
144 after which time the technician provided the lead researcher with the participants' intervention
145 numbers and corresponding HC doses. Three doses of HC (0 g, 15 g and 30 g, Myprotein,
146 Cheshire, UK) with 50 mg vitamin C powder (Holland and Barrett Retail Limited,
147 Warwickshire, UK) were dissolved in 300 mL water in an opaque drinks bottle. To match the
148 calories of 30 g HC in the other two interventions, 34.1 g and 15.4 g maltodextrin (Myprotein,
149 Cheshire, UK) was used in the 0 g and 15 g HC interventions, respectively. Although the
150 supplements were described by the manufacturers as "flavourless", 4 g non-caloric sweetener
151 (Truvia®, SilverSpoon, London, UK) was added in all drinks to mask any potential taste
152 difference between interventions. The amino acid profile of the HC supplement is shown in
153 **Table 1.**

154 *Insert Table 1 near here.*

155 Blood sampling

156 The BD Nexibia™ closed IV catheter system (22 G, Becton, Dickinson and Company, Franklin
157 Lakes, USA) was inserted into a peripheral vein in the right antecubital fossa by a trained
158 phlebotomist. A dressing band (3M™ Tegaderm™ I.V. Advanced Securement Dressing, 3M
159 Health Care, Loughborough, UK) then covered the catheter in order to secure the catheter site
160 and to keep it clean. Eight 5 mL venous blood samples were collected in specialized serum

161 collection tubes (BD Vacutainer™ Serum Separation Tube (SST™) II Advance, Dickinson and
162 Company, Franklin Lakes, USA) at the following time points: at rest immediately prior to HC
163 ingestion, 0.5 h post HC ingestion, 1 h post HC ingestion, 0.5 h post RE, 1 h post RE, 2 h post
164 RE, 4 h post RE and 6 h post RE for serum preparation (Figure 2). The samples were used to
165 analyse serum PINP and amino acid concentration. Two × 5 mL venous blood samples were
166 collected in EDTA plasma collection tubes (BD Vacutainer™ Hemogard Closure Plastic K2-
167 Ethylenediaminetetraacetic acid (EDTA) Tubes, Dickinson and Company, Franklin Lakes,
168 USA) at rest immediately prior to HC ingestion and 6 h post RE for plasma preparation. These
169 samples were used to analyse plasma β -CTX. The catheter was flushed by 3 mL sterile pre-
170 filled flush syringes containing sodium chloride 0.9% (BD PosiFlush™ Pre Filled Saline
171 Syringe, Dickinson and Company, Franklin Lakes, USA) every 30 min to clean and prevent
172 blood from clotting and blocking the catheter. The SSTs were stored in a tube rack for 30 min
173 for clotting at room temperature and the EDTA tubes were immediately placed on ice before
174 being centrifuged at 1000 g at 4°C for 10 min. The serum and plasma samples were then
175 aliquoted into 5 mL round-bottom polystyrene tubes (Falcon™, Thermo Fisher Scientific,
176 Whitby Canada) and stored at -80 °C until subsequent analysis.

177 **Blood analyses**

178 Markers of collagen synthesis and breakdown were analysed by measuring the circulating
179 concentration of PINP and β -CTX, respectively. Further, circulating collagen amino acid
180 concentrations were measured throughout the entirety of each intervention. PINP analyses were
181 performed at Liverpool John Moores University, while β -CTX and amino acid profile analyses
182 were performed at the Bioanalytical Facility, University of East Anglia.

183 *PINP*

184 Six serum samples (at rest prior to HC ingestion, 0.5 h-post RE, 1-h post RE, 2 h-post RE, 4 h-
185 post RE and 6 h-post RE) were used to measure serum PINP concentrations using an enzyme-
186 linked immunosorbent assay (ELISA) according to the manufacturer's instructions (USCN
187 Life Sciences, Wuhan, China). The intra-assay coefficient of variation (CV) was <10% and the
188 inter-assay CV was <12%, with a detection range of 2.47-200 $\mu\text{g}\cdot\text{L}^{-1}$, and sensitivity of <0.91
189 $\mu\text{g}\cdot\text{L}^{-1}$. The ELISA absorbance readings were performed at 450 nm, using a Clariostar
190 microplate reader (BMG Labtech, Ortenberg, Germany). The concentration \times time total area
191 under the curve (AUC) for PINP and amino acids (see below) were calculated using Prism
192 software (version 9.4.1, GraphPad Inc., San Diego, San Diego, California USA).

193 *β -CTX*

194 EDTA plasma concentrations of β -CTX were measured using electrochemiluminescence
195 immunoassay on a Cobas e601 analyser (Roche Diagnostics, Germany). The inter-assay CV
196 for β -CTX was $\leq 3\%$ between 0.2 and 1.5 $\mu\text{g}\cdot\text{L}^{-1}$ with the sensitivity of 0.01 $\mu\text{g}\cdot\text{L}^{-1}$.

197 *Amino acid profile*

198 Eight serum samples (at rest immediately prior to HC ingestion, 0.5 h post HC ingestion, 1 h
199 post HC ingestion, 0.5 h post RE, 1 h post RE, 2 h post RE, 4 h post RE and 6 h post RE) were
200 used to assess the concentration of 18 amino acids associated with collagen composition
201 (glycine, proline, hydroxyproline, glutamic acid, alanine, arginine, aspartic acid, lysine, serine,
202 leucine, valine, phenylalanine, threonine, isoleucine, histidine, tyrosine, methionine, and
203 glutamine, but not hydroxylysine). All 18 amino acid concentrations were measured
204 simultaneously using anionic ion-pair reverse phase liquid chromatography tandem mass
205 spectrometry (LC-MS/MS) system following derivatisation of the amino acid with *n*-butanol
206 hydrogen chloride. The LC-MS/MS system used a Micromass® Quattro Ultima™ Pt

207 (Manchester, UK) coupled to an Agilent 1100 series (Cheadle, UK) high performance liquid
208 chromatography binary pump. Electrospray ionisation source operating in positive ion mode,
209 mass detection for each amino acid butyl ester was achieved in multiple reaction monitoring
210 mode. Certified amino acid standards were purchased from Wacko Chemicals GmbH (Neuss,
211 Germany) and Sigma-Aldrich (Dorset, UK). Internal standards used were glucosaminic acid
212 and S-(2-Aminoethyl)-L-cysteine hydrochloride (Sigma-Aldrich, Dorset, UK) and L-
213 Citrulline-2,3,3,4,4,5,5-d7 (Isoscience, King of Prussia, PA, USA). Three internal quality
214 controls (QC) at low, medium and high concentrations were made from pooled human serum.
215 For each batch of analysis, 10 μL of standards, QC and test samples were added to a
216 microcentrifuge tubes, to which 440 μL of internal standards made up in 0.1M hydrochloride
217 in methanol was added. The mixture was vortexed twice, each time allowed to stand for 10
218 min, then centrifuged at $10,800 \times g$ for 5 mins. The supernatant was then transferred into a
219 borosilicate tube and dried to completeness under nitrogen gas at a temperature of 60°C . 100
220 μL of 3N *n*-butanol hydrogen chloride was added to the dried residue, vortex mixed, capped
221 and incubated at 60°C for 7 min. Following butylation, the mixture was dried completely under
222 nitrogen gas, and then reconstituted with 250 μL of 12% acetonitrile:water containing 0.025%
223 heptafluorobutyric acid (HFBA). After a final vortex mix, the samples were transferred to a
224 polypropylene autosampler vial for injection into the LC-MS/MS.

225 Chromatographic separation was achieved using a Modus AAC 100 x 2.1mm $3\mu\text{m}$ column
226 (Chromatography Direct Ltd, Runcorn, UK) maintained at 40°C . Anionic ion-pair reagent
227 HFBA was added to the mobile phases to improve analyte interaction with the stationary phase.
228 A gradient elution profile at a flow rate of $350 \mu\text{L}\cdot\text{min}^{-1}$ was used throughout. Initial conditions
229 were 88% mobile phase A (0.025% HFBA in water) and 12% mobile phase B (0.025% HFBA
230 in acetonitrile). This was held for 30 s. Mobile phase B was increased linearly to 20% at 10

231 min, with a further linear increase in mobile phase B to 60% at 15 min. This was held constant
232 until 16.9 min and returned to the initial conditions at 17 min. Injection volume was 10 μL with
233 an injection cycle time of 20 min. The assay range was 0 – 2000 $\mu\text{mol}\cdot\text{L}^{-1}$ for all 18 amino
234 acids studied. Inter-assay precision coefficient of variation (CV) for all amino acids were
235 between 3.3% to 10.3%.

236 **Statistical analyses**

237 Data were analysed using the statistical software package SPSS (Version 26, IBM Inc.,
238 Armonk, NY, USA). Sample size was estimated prior to conducting the study with G*Power
239 software (version 3.1.9.6, Heinrich-Heine-Universität Düsseldorf, Düsseldorf, Germany). The
240 estimation was performed using a large effect size ($\eta_p^2 = 0.22$), based on the results from Shaw
241 et al. (21), which demonstrated a two-fold increase in the serum PINP concentration \times time
242 area under the curve (AUC) following exercise with 15 g vs. 5 g gelatin ingestion. The results
243 from our a priori power calculation deemed a minimum of eight participants was necessary to
244 detect an effect of HC dose (one-way repeated measures analysis of variance (ANOVA); α :
245 0.05; power: 0.80). We recruited 10 participants to account for an expected 10-20% drop out.
246 Using the Shapiro-Wilk test, all data were deemed to be normally distributed except for the
247 amino acid data. The latter data were therefore log transformed prior to undergoing subsequent
248 statistical analyses. One-way within-subject ANOVA models were performed to compare
249 baseline (-1 h) concentrations of PINP and β -CTX in all three trails. Two-way within-subject
250 ANOVAs (dose \times time) were performed to detect changes in serum PINP and amino acid
251 concentrations over time. To detect changes in plasma β -CTX concentration, a two-way within-
252 subject ANOVA (dose \times time) was performed. One-way repeated measures ANOVA models
253 were performed to detect dose-dependent differences in concentration \times time AUCs for PINP

254 and each of the 18 amino acids analysed. Where Mauchly's test of sphericity had been violated,
255 Greenhouse-Geisser ($\epsilon < 0.75$) or Huynh-Feldt ($\epsilon > 0.75$) corrections were applied. Where a
256 main effect of HC dose existed, Fisher's LSD post-hoc pairwise comparisons were performed
257 to reveal which doses differed. Partial eta squared effect sizes (η_p^2) were reported for each
258 statistical model, and the thresholds for η_p^2 are defined as small ($\eta_p^2 = 0.01$), medium ($\eta_p^2 =$
259 0.06) and large ($\eta_p^2 = 0.14$) (25). All data analyses matched the research design, as there were
260 no missing data for any of the dependent variables. The level of statistical significance was set
261 at $P < 0.05$ and all data are presented as mean \pm standard deviations with 95% confidence
262 intervals (CI, where applicable), unless stated otherwise.

263

264 **Results**

265 *Serum PINP concentration and AUC*

266 Baseline serum PINP concentrations for the 0 g, 15 g and 30 g HC interventions did not differ
267 ($P = 0.990$, $\eta_p^2 = 0.001$; 0 g HC 32.7 ± 28.0 (95% CI: 12.7 – 52.7) $\mu\text{g}\cdot\text{L}^{-1}$; 15 g HC 32.7 ± 24.2
268 (95% CI: 15.4 – 49.9) $\mu\text{g}\cdot\text{L}^{-1}$; 30 g HC 32.3 ± 21.6 (95% CI: 16.9 – 47.7) $\mu\text{g}\cdot\text{L}^{-1}$. Regarding
269 serum PINP concentration, there was a main effect of HC dose ($P = 0.004$, $\eta_p^2 = 0.462$) and
270 time ($P = 0.013$, $\eta_p^2 = 0.458$) but no dose \times time interaction effect ($P = 0.732$, $\eta_p^2 = 0.071$,
271 **Figure 4A**). These results suggest the dose effect was not time specific. Post-hoc pairwise
272 comparisons revealed that 30 g HC had a greater PINP response than 0 g HC ($P = 0.002$) and
273 15 g HC ($P = 0.020$), while 15 g HC did not differ from 0 g HC ($P = 0.245$). Regarding the
274 serum PINP concentration \times time AUC, there was a main effect of HC dose ($P = 0.001$, $\eta_p^2 =$
275 0.517), and post-hoc pairwise comparisons revealed that 30 g HC had a greater AUC ($267 \pm$
276 79 [95% CI: 211 – 323] $\mu\text{g}\cdot\text{L}^{-1}\cdot\text{h}$ than 15 g HC (235 ± 70 [95% CI: 184 – 284] $\mu\text{g}\cdot\text{L}^{-1}\cdot\text{h}$, $P =$

277 0.013) and 0 g HC (219 ± 88 [95% CI: 155 – 281] $\mu\text{g}\cdot\text{L}^{-1}\cdot\text{h}$, $P = 0.002$), but 15 g HC AUC did
278 not differ from 0 g HC AUC ($P = 0.225$, **Figure 4B**).

279 *Insert Figure 4 near here.*

280 *Plasma β -CTX*

281 Baseline plasma β -CTX concentrations for the 0 g, 15 g and 30 g HC interventions did not
282 differ ($P = 0.311$, $\eta_p^2 = 0.122$; 0 g HC 0.8 ± 0.3 (95% CI: 0.6 – 1.0) $\mu\text{g}\cdot\text{L}^{-1}$; 15 g HC 0.8 ± 0.3
283 (95% CI: 0.6 – 0.9) $\mu\text{g}\cdot\text{L}^{-1}$; 30 g HC 0.8 ± 0.2 (95% CI: 0.6 – 0.9) $\mu\text{g}\cdot\text{L}^{-1}$. There was a main
284 effect of time ($P = 0.007$, $\eta_p^2 = 0.577$) but no main effect of HC dose ($P = 0.286$, $\eta_p^2 = 0.127$)
285 and no dose \times time interaction ($P = 0.748$, $\eta_p^2 = 0.031$), i.e. plasma β -CTX concentration
286 decreased from -1 h (prior to HC ingestion and RE) to 6 h post RE for all three interventions,
287 with no difference between intervention (**Figure 5**).

288 *Insert Figure 5 near here.*

289 *Serum amino acid concentrations*

290 Serum concentrations over the 7-h period of each intervention of the 18 amino acids that
291 constitute type I collagen are shown in **Figure 6**. The main effects of time, dose, and dose \times
292 time interaction effects for each amino acid are denoted in **Figure 6**. There were main effects
293 of dose for 14 amino acids (glycine, proline, hydroxyproline, glutamic acid, alanine, arginine,
294 aspartic acid, lysine, serine, leucine, valine, isoleucine, methionine, and glutamine), with 30 g
295 HC demonstrating higher serum concentrations than 0 g. Of those amino acids, glycine, proline,
296 hydroxyproline, arginine, lysine, serine, leucine, valine, isoleucine, and methionine in 30 g HC
297 showed higher serum concentrations than 15 g HC. All amino acids except for histidine ($P >$
298 0.05) showed a dose \times time interaction effect, and there was no main effect of dose for histidine

299 ($P = 0.451$).

300 *Insert Figure 6 near here.*

301 **Discussion**

302 This study is the first to investigate the effect of *high-intensity* resistance exercise (RE)
303 and 0, 15 and 30 g hydrolyzed collagen (HC) supplementation on whole body collagen turnover
304 in an homogenous group of resistance-trained, healthy, young men. We found that the serum
305 PINP concentration \times time area-under-the-curve (AUC) for the 30 g HC intervention was
306 greater than for the 15 g and 0 g HC interventions. Further, these results were consistent with
307 greater increases in the appearance of key amino acid constituents of collagen (e.g. glycine and
308 proline) within the blood following ingestion of 30 g HC versus 15 g and 0 g. Therefore, at
309 least 30 g HC is required to provide greater exogenous collagen amino acid availability, which
310 appears to be a key factor for optimising collagen synthesis following high-intensity RE in
311 resistance-trained, young men.

312 To address the aims of our study, we measured serum PINP following RE with different
313 doses of HC. We chose high-intensity back squat RE to target the quadriceps muscle-tendon
314 unit (MTU), because the human patellar tendon appears to hypertrophy only following
315 prolonged periods of high-intensity (6, 7) and not moderate-intensity (26) resistance training.
316 In the current study, the 7-h experimental design was based on a significant increase in muscle
317 and tendon collagen fractional synthetic rate (FSR) at 6 h post-exercise following 1-h RE in
318 healthy young men (12). Although the significantly elevated muscle collagen FSR was similar
319 at 6 h post- and 24 h post-RE (12), tendon collagen FSR appeared to be further augmented at
320 24 h post- compared to 6 h post-RE, although it is not stipulated in the article whether tendon
321 collagen FSR measured at these two time points differed significantly. In a separate study,

322 serum PICP concentration was significantly higher 48 h post-exercise following 50 maximal
323 concentric knee extensions in healthy young men (27). Thus, it is possible that, had each of our
324 three interventions lasted 24 – 48 h post-RE, we may have observed further increases in serum
325 PINP concentration, and possibly a larger effect of HC ingestion on these increases.

326 The 30 min high-intensity RE model we employed in our study was associated with
327 peak serum PINP concentrations of $\sim 60 \mu\text{g}\cdot\text{L}^{-1}$ (regardless of dose), which occurred 30-60 min
328 after the onset of RE (and 90-120 min after supplement ingestion). We chose to measure serum
329 PINP concentration because it is a reliable biomarker of collagen synthesis, being a procollagen
330 peptide that is cleaved off during maturation from procollagen to collagen (28). The similar
331 PINP concentration at +0.5 and +1 h post-RE for all three doses (including 0 g HC) suggests
332 the increase in PINP concentration observed within the first hour after starting the RE occurred
333 as a consequence of the RE, rather than HC ingestion. However, it is possible that our data at
334 +0.5 h post-RE may have been influenced by an increase in blood flow. A distinction should
335 be made between flux, i.e. the total amount of PINP passing through the blood registered at
336 any given time, and concentration, i.e. the ratio of PINP to the volume of serum. Blood flow
337 rises 20-fold and 7-fold in the calf muscle and peritendinous area of the Achilles tendon,
338 respectively, during repeated plantar flexion contractions in healthy individuals (29). However,
339 an increase in blood flow to the peritendinous tendon has been shown to return to resting levels
340 within a few minutes of finishing the same type of exercise (30). Therefore, the fact that serum
341 PINP concentration was still at its peak at +1 h in the current study, i.e. 30 min after RE ended,
342 when cardiac output would be expected to have returned to resting rates following lower-limb
343 RE (31), it is likely that this increase in PINP concentration was due to an increase in RE-
344 induced collagen synthesis, rather than blood flow. Serum PINP concentration decreased to
345 baseline values in the 0 g and 15 g interventions for the remainder of those interventions, while

346 it decreased but remained more elevated in the 30 g intervention, which resulted in the higher
347 AUC in the 30 g intervention compared to the 0 g and 15 g interventions.

348 In contrast to our results, Aussieker et al. (32) recently found that 30 g HC ingestion
349 with six sets of 8–15 repetitions at 60% estimated 1-RM barbell back squat did not augment
350 *vastus lateralis* muscle connective tissue protein FSR or circulating PINP concentration more
351 than RE with 30 g whey protein ingestion or RE alone in different groups of young men and
352 women. A number of differences in study design may help explain this discrepancy between
353 studies. Firstly, the between-group design used by Aussieker et al. (32) may have introduced
354 more within and between intervention variability (thus potentially confounding an effect of HC)
355 compared to a within-group cross-over design, as used in the current study. Secondly, estrogen
356 is known to affect skeletal muscle and tendon collagen synthesis in women (33, 34), and the
357 use of a mixed-sex cohort by Aussieker et al. (32), rather than a 100% male cohort as used in
358 the current study, may have increased within- and between-group variability in connective
359 tissue protein FSR. Thirdly, vitamin C was not consumed during the interventions by Aussieker
360 et al. (32), which began after an overnight fast. As vitamin C is required for the biosynthesis
361 of collagen (17) and humans are unable to store it in the body or synthesize it endogenously
362 (35), this may have limited muscle connective tissue protein FSR. Finally, it should be noted
363 that Aussieker et al. (32) measured connective tissue protein FSR in skeletal muscle and not
364 tendon or ligament. The latter tissues have a 70-85% type I collagen content (9) compared to
365 just ~5% in skeletal muscle (36).

366 Contrary to Aussieker et al. (32) and in accordance with our findings, Shaw et al. (21)
367 found that serum PINP concentration was greater following jump-rope exercise with gelatin
368 supplementation in a dose-response manner. Due to different time points used to measure
369 serum PINP concentration, a direct comparison of peak concentration between studies is not

370 possible. Nevertheless, the serum PINP AUC was greater in our 30 g HC intervention
371 compared to our 15 g and 0 g interventions (with no difference between our 0 g and 15 g
372 interventions, **Figure 4**), while Shaw et al. (21) found a greater effect of 15 g versus 5 g and 0
373 g gelatin. This suggests that the different exercise models used may require different doses of
374 exogenous collagen to optimise the collagen synthetic response for that particular exercise.

375 Regarding the amino acids that constitute collagen (e.g. glycine, proline,
376 hydroxyproline, etc.), these peaked in circulation around 1 – 1.5 h after ingestion of 30 g HC
377 in our study (**Figure 6**). This was in line with previous studies, which involved the ingestion
378 of 15 g gelatin or 20 g or 30 g collagen peptides in healthy young populations (21, 32, 37). This
379 similarity between studies indicates the maximal rate of amino acid absorption occurs
380 approximately an hour after ingestion of collagen in healthy young men, regardless of dose and
381 type of collagen supplementation. We also observed that the average concentrations of glycine,
382 proline, hydroxyproline, arginine, lysine, serine, leucine, valine, isoleucine, and methionine
383 were greater after ingestion of 30 g HC compared to 15 g and 0 g HC. The high availability of
384 collagen amino acids in the 30 g HC intervention might have promoted a greater collagen
385 synthetic response in two ways. Firstly, the greater abundance of key amino acids may have
386 simply provided more of the essential components to increase collagenous tissue content
387 following an overload-induced stimulation of collagen synthesis in the lower-limb MTUs.
388 Secondly, they may have stimulated mammalian target of rapamycin complex 1 (mTORC1)
389 phosphorylation independently of muscle contraction/stretch-activated mechanisms, in a
390 similar manner to amino acid stimulation of skeletal muscle myofibrillar protein synthesis via
391 mTORC1 activation (38). For example, after treating chondrocytes from bovine cartilage with
392 1.5 mM glycine, proline, or lysine for 15 days, type II collagen synthesis was 1.6 times greater
393 following glycine treatment compared to proline treatment and 2 times greater compared to

394 lysine treatment (39). Further, human skin fibroblasts treated with 5 mM proline for 48 h
395 demonstrated an increase in collagen type I $\alpha 1$ expression (40), while proline and
396 hydroxyproline also increased TGF- β expression in human fibroblasts (41), which would be
397 expected to cause phosphorylation of protein kinase B (Akt) and mTORC1 (thus explaining
398 the increase in collagen synthesis and gene expression). Considering the above mechanisms
399 and that collagen synthesis is stimulated via phosphorylation of Akt and mTORC1 in response
400 to mechanical loading in cultured human tendon-derived stromal cells (42), and that
401 mechanical loading increases in TGF- β expression and type I collagen expression in rat
402 Achilles tendon (43), independent RE- and amino acid-associated signalling pathways likely
403 explain our findings that high-intensity RE with 30 g HC supplementation stimulates collagen
404 synthesis more than RE with 15 g HC ingestion or RE alone.

405 Concerning collagen breakdown, β -CTX is released into circulation from mature type
406 I collagen during degradation, and therefore serves as a reliable biomarker of collagen
407 breakdown (44). In the current study, regardless of HC dose, plasma β -CTX concentration
408 decreased by ~30% from -1 to 6 h post RE (**Figure 5**). This may have been caused by the high-
409 intensity RE stimulating collagen synthesis (**Figure 4**), subsequently inhibiting collagen
410 breakdown. This hypothesis is supported by currently unpublished data from our laboratory in
411 middle-aged men, suggesting the decrease in circulating β -CTX concentration occurs
412 immediately after RE and remains lower for the subsequent six hours of rest, regardless of HC
413 dose. Alternatively, the reduction may have been associated with circadian rhythm, as Qvist et
414 al. (45) reported that resting serum β -CTX concentrations in men and pre- and postmenopausal
415 women (aged 24-73 years) peaked at 08:00, then sharply decreased between 11:00 and 14:00.
416 Nevertheless, as tissue turnover is determined by the rates of both synthesis and breakdown,
417 the role of collagen degradation on the regulation of collagen turnover is crucial. For example,

418 avian skeletal muscle exposed to mechanical loading led to an increase in collagen synthesis,
419 which was accompanied by a decrease in the degradation of newly synthesized collagen, as
420 well as an increase in the degradation of mature collagen (46). Thus, further research is
421 necessary to elucidate the effects of RE and HC supplementation on degradation of newly
422 synthesized and mature collagen for the regulation of collagen turnover.

423 A limitation of this study is that our assessment of collagen synthesis was indirect, i.e.
424 we measured serum PINP concentration rather than harvesting tendon biopsies, with which we
425 could have assessed tendon PINP concentration or collagen FSR directly. However, human
426 tendon (47) and serum PINP concentration (13) increases after an acute bout of exercise, and
427 serum PINP can remain elevated for up to four days following resistance-type exercise in
428 healthy young men (27). Nevertheless, future studies should investigate the effect of HC
429 ingestion with RE on connective tissue collagen synthesis by measuring circulating and tendon
430 PINP concentration, and tendon collagen FSR simultaneously.

431 In conclusion, we have demonstrated for the first time that a single bout of high-
432 intensity, lower-limb RE with 30 g HC ingestion increased whole body collagen synthesis more
433 than RE with 15 g or 0 g HC in resistance-trained young men. This higher response was likely
434 related to the greater availability of key amino acids following the ingestion of 30 g HC
435 compared to 15 g and 0 g. This may have implications for augmenting tendon adaptation to
436 high-intensity resistance training when 30 g HC is ingested in combination with resistance
437 exercise over a prolonged period of time. Future studies should also investigate if a dose-
438 response exists regarding high-intensity RE with HC supplementation in resistance-trained
439 young women.

440

441 **Conflict of interest**

442 The authors declare that the research was conducted in the absence of any commercial or
443 financial relationships that could be construed as a potential conflict of interest.

444

445 **Acknowledgements**

446 Authors contributed to this manuscript as follows: study conception and design (RME); data
447 collection (JL); data curation (JL); data analysis (JL, RME, JCYT, RD, JD); supervision (RME,
448 DRC, CES); writing – original draft (JL); writing – review, editing and approval of final draft
449 (RME, JCYT, RD, JD, WDF, KE, DRC, CES).

References

1. Ekstrand J, Hägglund M, Waldén M. Injury incidence and injury patterns in professional football: the UEFA injury study. *Br J Sports Med.* 2011;45(7):553-8.
2. Keller CS, Noyes FR, Buncher CR. The medical aspects of soccer injury epidemiology. *Am J Sports Med.* 1987;15(3):230-7.
3. Mendiguchia J, Alentorn-Geli E, Brughelli M. Hamstring strain injuries: are we heading in the right direction? *British Journal of Sports Medicine.* 2012;46(2):81.
4. Lauersen JB, Andersen TE, Andersen LB. Strength training as superior, dose-dependent and safe prevention of acute and overuse sports injuries: a systematic review, qualitative analysis and meta-analysis. *British Journal of Sports Medicine.* 2018;52(24):1557.
5. Erskine RM, Jones DA, Williams AG, Stewart CE, Degens H. Resistance training increases in vivo quadriceps femoris muscle specific tension in young men. *Acta physiologica (Oxford, England).* 2010;199(1):83-9.
6. Kongsgaard M, Reitelseder S, Pedersen TG, Holm L, Aagaard P, Kjaer M, et al. Region specific patellar tendon hypertrophy in humans following resistance training. *Acta Physiol (Oxf).* 2007;191(2):111-21.
7. Seynnes OR, Erskine RM, Maganaris CN, Longo S, Simoneau EM, Grosset JF, et al. Training-induced changes in structural and mechanical properties of the patellar tendon are related to muscle hypertrophy but not to strength gains. *J Appl Physiol (1985).* 2009;107(2):523-30.
8. LaCroix AS, Duenwald-Kuehl SE, Lakes RS, Vanderby R, Jr. Relationship between tendon stiffness and failure: a metaanalysis. *J Appl Physiol (1985).* 2013;115(1):43-51.
9. Kjaer M. Role of Extracellular Matrix in Adaptation of Tendon and Skeletal Muscle to Mechanical Loading. *Physiological Reviews.* 2004;84(2):649-98.

10. Heinemeier KM, Kjaer M. In vivo investigation of tendon responses to mechanical loading. *J Musculoskelet Neuronal Interact.* 2011;11(2):115-23.
11. Couppe C, Svensson RB, Skovlund SV, Jensen JK, Eriksen CS, Malmgaard-Clausen NM, et al. Habitual side-specific loading leads to structural, mechanical, and compositional changes in the patellar tendon of young and senior lifelong male athletes. *J Appl Physiol* (1985). 2021;131(4):1187-99.
12. Miller BF, Olesen JL, Hansen M, Dossing S, Cramer RM, Welling RJ, et al. Coordinated collagen and muscle protein synthesis in human patella tendon and quadriceps muscle after exercise. *J Physiol.* 2005;567(Pt 3):1021-33.
13. Huang T-H, Nosaka K, Chen TC. Changes in blood bone markers after the first and second bouts of whole-body eccentric exercises. *Scandinavian Journal of Medicine & Science in Sports.* 2022;32(3):521-32.
14. Chiquet M, Gelman L, Lutz R, Maier S. From mechanotransduction to extracellular matrix gene expression in fibroblasts. *Biochim Biophys Acta.* 2009;1793(5):911-20.
15. Heinemeier K, Langberg H, Olesen JL, Kjaer M. Role of TGF- β 1 in relation to exercise-induced type I collagen synthesis in human tendinous tissue. *Journal of Applied Physiology.* 2003;95(6):2390-7.
16. Olesen JL, Heinemeier KM, Haddad F, Langberg H, Flyvbjerg A, Kjaer M, et al. Expression of insulin-like growth factor I, insulin-like growth factor binding proteins, and collagen mRNA in mechanically loaded plantaris tendon. *J Appl Physiol* (1985). 2006;101(1):183-8.
17. Murad S, Grove D, Lindberg KA, Reynolds G, Sivarajah A, Pinnell SR. Regulation of collagen synthesis by ascorbic acid. *Proc Natl Acad Sci U S A.* 1981;78(5):2879-82.
18. Canty EG, Kadler KE. Procollagen trafficking, processing and fibrillogenesis. *Journal*

of Cell Science. 2005;118(7):1341.

19. Kubo K, Ikebukuro T, Maki A, Yata H, Tsunoda N. Time course of changes in the human Achilles tendon properties and metabolism during training and detraining in vivo. *Eur J Appl Physiol*. 2012;112(7):2679-91.

20. Bojsen-Møller J, Magnusson SP, Rasmussen LR, Kjaer M, Aagaard P. Muscle performance during maximal isometric and dynamic contractions is influenced by the stiffness of the tendinous structures. *J Appl Physiol* (1985). 2005;99(3):986-94.

21. Shaw G, Lee-Barthel A, Ross ML, Wang B, Baar K. Vitamin C-enriched gelatin supplementation before intermittent activity augments collagen synthesis. *Am J Clin Nutr*. 2017;105(1):136-43.

22. König D, Oesser S, Scharla S, Zdzieblik D, Gollhofer A. Specific Collagen Peptides Improve Bone Mineral Density and Bone Markers in Postmenopausal Women—A Randomized Controlled Study. *Nutrients*. 2018;10(1):97.

23. McAlindon TE, Nuite M, Krishnan N, Ruthazer R, Price LL, Burstein D, et al. Change in knee osteoarthritis cartilage detected by delayed gadolinium enhanced magnetic resonance imaging following treatment with collagen hydrolysate: a pilot randomized controlled trial. *Osteoarthritis Cartilage*. 2011;19(4):399-405.

24. Macnaughton LS, Wardle SL, Witard OC, McGlory C, Hamilton DL, Jeromson S, et al. The response of muscle protein synthesis following whole-body resistance exercise is greater following 40 g than 20 g of ingested whey protein. *Physiological Reports*. 2016;4(15):e12893.

25. Cohen J. *Statistical power analysis for the behavioural sciences* (Second edition). Second edition ed. Hillsdale, NJ: Lawrence Erlbaum Associates; 1988.

26. Lee J, Bridge JE, Clark DR, Stewart CE, Erskine RM. Collagen supplementation

- augments changes in patellar tendon properties in female soccer players. *Frontiers in Physiology*. 2023;14.
27. Virtanen P, Viitasalo JT, Vuori J, Vaananen K, Takala TE. Effect of concentric exercise on serum muscle and collagen markers. *J Appl Physiol* (1985). 1993;75(3):1272-7.
28. Heinemeier KM, Kjaer M, Magnusson SP. Methods of Assessing Human Tendon Metabolism and Tissue Properties in Response to Changes in Mechanical Loading. *Adv Exp Med Biol*. 2016;920:97-106.
29. Boushel R, Langberg H, Green S, Skovgaard D, Bulow J, Kjaer M. Blood flow and oxygenation in peritendinous tissue and calf muscle during dynamic exercise in humans. *J Physiol*. 2000;524 Pt 1(Pt 1):305-13.
30. Langberg H, Rosendal L, Kjaer M. Training-induced changes in peritendinous type I collagen turnover determined by microdialysis in humans. *J Physiol*. 2001;534(Pt 1):297-302.
31. Poton R, Polito MD. Hemodynamic response to resistance exercise with and without blood flow restriction in healthy subjects. *Clin Physiol Funct Imaging*. 2016;36(3):231-6.
32. Aussieker T, Hilken L, Holwerda AM, Fuchs CJ, Houben LHP, Senden JM, et al. Collagen Protein Ingestion during Recovery from Exercise Does Not Increase Muscle Connective Protein Synthesis Rates. *Med Sci Sports Exerc*. 2023.
33. Hansen M, Kongsgaard M, Holm L, Skovgaard D, Magnusson SP, Qvortrup K, et al. Effect of estrogen on tendon collagen synthesis, tendon structural characteristics, and biomechanical properties in postmenopausal women. *J Appl Physiol* (1985). 2009;106(4):1385-93.
34. Hansen M, Miller BF, Holm L, Doessing S, Petersen SG, Skovgaard D, et al. Effect of administration of oral contraceptives in vivo on collagen synthesis in tendon and muscle connective tissue in young women. *Journal of Applied Physiology*. 2009;106(4):1435-43.

35. Li Y, Schellhorn HE. New Developments and Novel Therapeutic Perspectives for Vitamin C1,2. *The Journal of Nutrition*. 2007;137(10):2171-84.
36. Babraj JA, Cuthbertson DJR, Smith K, Langberg H, Miller B, Krogsgaard MR, et al. Collagen synthesis in human musculoskeletal tissues and skin. *American Journal of Physiology-Endocrinology and Metabolism*. 2005;289(5):E864-E9.
37. Alcock RD, Shaw GC, Tee N, Burke LM. Plasma Amino Acid Concentrations After the Ingestion of Dairy and Collagen Proteins, in Healthy Active Males. *Frontiers in Nutrition*. 2019;6(163).
38. Drummond MJ, Dreyer HC, Fry CS, Glynn EL, Rasmussen BB. Nutritional and contractile regulation of human skeletal muscle protein synthesis and mTORC1 signaling. *Journal of Applied Physiology*. 2009;106(4):1374-84.
39. de Paz-Lugo P, Lupiáñez JA, Meléndez-Hevia E. High glycine concentration increases collagen synthesis by articular chondrocytes in vitro: acute glycine deficiency could be an important cause of osteoarthritis. *Amino Acids*. 2018;50(10):1357-65.
40. Szoka L, Karna E, Hlebowicz-Sarat K, Karaszewski J, Palka JA. Exogenous proline stimulates type I collagen and HIF-1 α expression and the process is attenuated by glutamine in human skin fibroblasts. *Molecular and Cellular Biochemistry*. 2017;435(1):197-206.
41. Surazynski A, Miltyk W, Prokop I, Palka J. Prolidase-dependent regulation of TGF β (corrected) and TGF β receptor expressions in human skin fibroblasts. *Eur J Pharmacol*. 2010;649(1-3):115-9.
42. Mousavizadeh R, Hojabrpour P, Eltit F, McDonald PC, Dedhar S, McCormack RG, et al. β 1 integrin, ILK and mTOR regulate collagen synthesis in mechanically loaded tendon cells. *Scientific Reports*. 2020;10(1):12644.
43. Heinemeier KM, Olesen JL, Haddad F, Langberg H, Kjaer M, Baldwin KM, et al.

Expression of collagen and related growth factors in rat tendon and skeletal muscle in response to specific contraction types. *J Physiol.* 2007;582(Pt 3):1303-16.

44. Gineyts E, Cloos PA, Borel O, Grimaud L, Delmas PD, Garnero P. Racemization and isomerization of type I collagen C-telopeptides in human bone and soft tissues: assessment of tissue turnover. *Biochem J.* 2000;345 Pt 3(Pt 3):481-5.

45. Qvist P, Christgau S, Pedersen BJ, Schlemmer A, Christiansen C. Circadian variation in the serum concentration of C-terminal telopeptide of type I collagen (serum CTx): effects of gender, age, menopausal status, posture, daylight, serum cortisol, and fasting. *Bone.* 2002;31(1):57-61.

46. Laurent GJ, McAnulty RJ, Gibson J. Changes in collagen synthesis and degradation during skeletal muscle growth. *Am J Physiol.* 1985;249(3 Pt 1):C352-5.

47. Langberg H, Skovgaard D, Petersen LJ, Bulow J, Kjaer M. Type I collagen synthesis and degradation in peritendinous tissue after exercise determined by microdialysis in humans. *J Physiol.* 1999;521 Pt 1:299-306.

Figure legends and Table

Figure 1. CONSORT flow diagram. HC, hydrolyzed collagen; BS, blood samples collected for serum procollagen type I N-terminal propeptide (PINP) concentration, plasma β -isomerized C-terminal telopeptide of type I collagen (β -CTX) concentration, and serum amino acid concentration.

Figure 2. Schematic diagram of the experimental protocol. HC, hydrolyzed collagen; RE, resistance exercise; -1, rest prior to HC intake; -0.5, 0.5 h-HC ingestion; 0, 1 h-HC ingestion; +0.5, 0.5 h-post RE; 1, 1 h-post RE; 2, 2 h-post RE; 4, 4 h-post RE; 6, 6 h-post RE.

Figure 3. 90° barbell back squat. A resistance band was attached to the squat rack to indicate when the participant had reached 90° knee flexion during each repetition of the back squat 10-RM.

Figure 4. Collagen synthesis following hydrolyzed collagen (HC) ingestion (-1 h) and performing resistance exercise. (A) serum PINP concentrations (B) serum PINP concentration \times time area under the curve following 0 g HC (green circles), 15 g HC (blue squares), and 30 g HC (red triangles) ingestion. *Greater than 0 g HC ($P = 0.005$); †Greater than 15 g HC ($P = 0.039$). Values represent mean \pm SEM.

Figure 5. Collagen breakdown following hydrolyzed collagen (HC) intake and performing resistance exercise. Plasma CTX-1 concentrations following 0 g HC (green circles), 15 g HC (blue squares), and 30 g HC (red triangles) ingestion. *Lower than -1 h ($P < 0.05$).

Figure 6. Concentrations of eighteen serum amino acid at before ingesting hydrolyzed collagen (-1 h), 1 h after ingesting 0 g (green circles), 15 g (blue squares), or 30 g (red triangles) (HC) (+1 h) and then performing barbell back squat exercise at time point 0 h. Values represent mean

± SEM.

Table 1. Amino acid composition of the hydrolyzed collagen supplement.

Amino acids	Weight (%)
Glycine	21.0
Proline	12.8
Hydroxyproline	12.2
Glutamic acid	10.3
Alanine	8.9
Arginine	7.3
Aspartic acid	6.0
Lysine	3.5
Serine	3.1
Leucine	2.7
Valine	2.4
Phenylalanine	2.1
Threonine	1.9
Hydroxylysine	1.5
Isoleucine	1.5
Histidine	1.1
Tyrosine	1.0
Methionine	0.9

Assessed for eligibility $n = 13$ **Excluded** $n = 3$

- Not meeting inclusion criteria ($n = 1$)
- Declined to participate ($n = 2$)
- Other reasons ($n = 0$)


Randomly assigned $n = 10$ **0 g HC** $n = 10$ **15 g HC** $n = 10$ **30 g HC** $n = 10$ **Completed** $n = 10$ **Completed** $n = 10$ **Completed** $n = 10$ **Analysed BS** $n = 10$ **Analysed BS** $n = 10$ **Analysed BS** $n = 10$

Enrollment

Allocation

Follow-Up

Analysis



HC



Exercise





Serum





Plasma











Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Journal Pre-proof