

Increasing global human exposure to wildland fires despite declining burned area

Seyd Teymoor Seydi¹, John T. Abatzoglou², Matthew W. Jones³, Crystal A. Kolden², Gabriel Filipelli⁴, Matthew D. Hurteau⁵, Amir AghaKouchak^{6,7}, Charles H. Luce⁸, Chiyuan Miao⁹, Mojtaba Sadegh^{1,10,*}

¹Department of Civil Engineering, Boise State University, Boise, ID, USA

²Management of Complex Systems Department, University of California, Merced, Merced, CA, USA

³Tyndall Centre for Climate Change Research, School of Environmental Sciences, University of East Anglia (UEA), Norwich, UK

⁴Department of Earth Science, Indiana University, Indianapolis, IN 46202, USA

⁵Biology Department, University of New Mexico, Albuquerque, NM, United States

⁶Department of Civil and Environmental Engineering, University of California, Irvine, CA, USA

⁷Department of Earth System Science, University of California, Irvine, CA, USA

⁸USDA Forest Service Rocky Mountain Research Station, Boise, Idaho, USA

⁹State Key Laboratory of Earth Surface Processes and Disaster Risk Reduction, Faculty of Geographical Science, Beijing Normal University, Beijing 100875, China

¹⁰United Nations University Institute for Water, Environment and Health, Richmond Hills, ON, Canada

*Correspondence to: mojtabasadegh@boisestate.edu

Abstract

Although half of Earth's population reside in the wildland-urban interface, human exposure to wildland fires remains unquantified. We show that population directly exposed to wildland fires increased 40% globally from 2002-2021, despite a 26% decline in burned area. Increased exposure was mainly driven by enhanced co-location of wildland fires and human settlements, doubling the exposure per unit burned area. We show that population dynamics accounted for one-quarter of the 440 million human exposures to wildland fires. While wildfire disasters in North America, Europe, and Oceania have garnered significant attention, 85% of global exposures occurred in Africa. The top 0.01% of fires by intensity accounted for 0.6% and 5% of global exposures and burned area, respectively, warranting enhanced efforts to increase fire resilience in disaster-prone regions.

Wildland fires – here defined as fires in vegetated land excluding commercial agriculture areas – have increasingly impacted social and environmental systems in various regions across the globe in recent years (1, 2), directly accounting for at least 2,500 human deaths and 10,500 injuries from 1990-2021 (3). Indirect impacts of wildland fires are several-fold larger and extend for years after the burn (4). For example, an annual 1.53 million deaths around the globe are attributable to landscape-fire-induced air pollution (5). Fire activity is directly linked with climate and weather (6), and climate change has increased the number of days conducive to extreme fire behavior over many fire-prone regions, culminating in a >54% increase in extreme fire weather from 1979-2022 globally (3), longer fire-weather seasons (7), and increased nighttime flammability (8).

Human activities can compound or confound the climate change impacts on fire activity (3, 9, 10). Human ignitions, both intentional and accidental, account for 84% of all wildfires in the contiguous United States (11) and >90% in Mediterranean Europe (12). Although lightning ignitions can dominate in more remote regions (13, 14), people strongly modify the timing and locations of fires (11, 15). People also modify vegetation type, structure, density and continuity, which, in turn, change regional fire regimes (16). For example, the introduction of invasive species in deserts of North America resulted in more frequent and larger wildfires (17), whereas agriculturally-induced land fragmentation in African savannas reduced burned areas (18). Outside of the tropics, humans commonly seek to limit burned area by suppressing fire, but as settlement expands into previously wildland areas, causing the Wildland-Urban Interface (WUI) to grow (19), human exposure to fire also grows (20). While the WUI covers only 4.7% of the global land surface, nearly half of Earth's population reside in WUI areas (21) and regional and global WUIs have been expanding (22, 23).

The confluence of changing social and environmental controls of fires resulted in 26% decline in global burned area in the past two decades, driven by trends in African savannas (24), despite increased fire extent in temperate and boreal forests (25) and increased occurrence of intense fire events (26). Here we examine the global and regional patterns and trends of human exposure to wildland fires. We further probe the characteristics of wildland fires that impact humans, and investigate the role of population dynamics in the overall exposure patterns and trends. Additionally, we investigate patterns, trends, and drivers of population exposure to intense fire events, as exposure to such fires has been tied to fire disasters (2). We use 18.6 million individual fire records from 2002-2021 from the MODIS-based Global Fire Atlas (18, 27) and gridded population data from WorldPop (28). We also use MODIS-based land cover and use (29), active fire records (30), and vegetation indices (31) to exclude non-wildland fires in the Global Fire Atlas (27). Our approach does not exclude deforestation

and other vegetation-clearing fires, hence does not differentiate between intentional fire use and wildfires. We define human exposure to wildland fire as the number of people who resided within the burned perimeters, noting that (a) burned perimeters include unburned patches, (b) consequential human impacts from fires extend well beyond burn perimeters, and (c) human and societal impacts of exposures to less frequent but more intense fires (e.g., in Western North America) are different from those in the case of frequent-low intensity fires (e.g., in African savannas). Exposure to wildland fires can be harmful and devastating, resulting in loss of lives and damage to infrastructure, noting that not all instances of exposure lead to negative outcomes.

Patterns of global exposure to wildland fire

A total of 440.2 million people were directly exposed to wildland fires globally from 2002-2021 (Fig. 1A), of which 85.6% occurred in Africa. While major wildfire disasters occurred in Oceania, Europe, and North America in recent years, these three continents cumulatively accounted for <2.5% of global exposures (Fig. 1B). Cumulative burned area in these wildland fires from 2002-2021 was 49.2 million km² (Fig. 1C), of which Africa accounted for 64.3% (Fig. 1D). Europe had the highest human exposure per unit burned area – hereafter, exposure density (17.7 persons/km²), followed by Africa (11.9 persons/km²). The global exposure density was 8.9 persons/km² (Fig. S1B).

Spatial patterns of cumulative human exposure to wildland fire, cumulative burned area, and per capita human exposure illustrate the variability in human relationships to fire across latitudes and continents (Figs. 1A,C,E). The highest levels of per capita exposure to wildland fire occurred in the tropical savannas of Africa (Fig. 1E; Figs. S2-S3), characterized by frequent human- and lightning-started low intensity fires where populations regularly use fire for land management and agricultural purposes (16). From 2002-2021, 1.8% of Africa's population was exposed to wildland fire, more than an order of magnitude larger than any other continent. Similarly, cumulative exposure to fire was highest in regions such as tropical Africa (16) with five tropical African countries accounting for half of the global exposures (Fig. 1A). High cumulative burned area in tropical Australia and Brazil (Fig. 1C) reflect both naturally-started fires and widespread intentional use of fire for regenerative and deforestation practices in low-to-unpopulated areas, while moderate cumulative burned area in sparsely populated regions of northern Russia and North America reflect increasing wildfire area burned associated with climate change.

Top-quintile events in terms of size, perimeter, duration, speed, and spread rate accounted for a majority of wildland fire exposures (see Section S1 and Figs. S4-S9).

Trends of global exposure to wildland fire

Global annual exposure to wildland fire increased by 7.7 million people (+39.6%) from 2002-2021 (382,700 persons/year; $p < 10^{-5}$; Fig. S1A). Africa accounted for nearly all of the increase (373,000 persons/year; $p < 10^{-7}$; +46.3%). South America (+33.8%), North America (+15.7%) and Asia (+2.8%) experienced non-significant increases in exposure to wildland fire from 2002-2021, whereas Oceania (-47.3%) and Europe (-17.3%) observed non-significant declines (Fig. S1A). Interannual variability of exposure to wildland fire and age distribution of the exposed populations are discussed in Sections S2 and S3 (Fig. S10), respectively.

Exposure density also increased globally, exposing an additional 6.2 (+93.4%) persons per km² of burned area in two decades ($p < 10^{-6}$; Fig. S1B). All continents experienced rising exposure density, with Africa and Europe showing the largest increases, exposing an additional 7.9 (+90.4%) and 6.8 (+42.3%) persons per km² of burned area, respectively, in two decades ($p < 10^{-7}$ and $p = 0.2$, respectively; Fig. S1B).

Role of population dynamics in exposure to wildland fire

By applying fixed 2002 population patterns, we developed a counterfactual model to account for the role of population dynamics versus fire dynamics in driving the observed exposure trends. Population dynamics – i.e., population growth and migration during 2002-2021 – accounted for 111.3 million exposures to wildland fire in the past two decades globally (25.3% of all global exposures; Fig. 2A). Without the effects of population dynamics, cumulative exposure to wildland fire from 2002-2021 would have been between 10.0% to 27.9% lower in Africa, Asia, North America, and South America, but 5.0% and 2.3% higher in Europe and Oceania, respectively (Fig. 2). Counterfactual trends in Europe and Oceania are driven by the marginal shrinking of the fire and human settlement overlap from 2002-2021 due a variety of factors including migration from rural to urban areas.

In the absence of population dynamics, the global exposure to wildland fire would have decreased at a rate of 278,000 people/year ($p < 10^{-4}$; 27.7% decline in two decades) commensurate with the decline in burned area (Fig. 2A) – with counterfactual trends primarily driven by declines in Africa (-250,000 people/year, $p < 10^{-4}$; -29.8%; Fig. 2E).

Without the population changes since 2002, exposure per unit burned area would be rather stable globally (1.8% decline in two decades; -0.006 persons/km²/year; $p = 0.83$), indicating that human influences, including population increase in and migration to fire-

prone areas, were the main driver of increasing exposures to wildland fires in the past two decades.

Population exposure to intense wildland fires

Following Cunningham et al. (26), we define “intense” fires as events with cumulative fire radiative power exceeding the 99.99th percentile of all wildland fires globally (Fig. 3A). Intense fires had a disproportionately large impact, representing just 0.01% of all wildland fires yet accounting for 0.6% of total population exposures (2.7 million people) and 5.0% of the cumulative burned area (2.4 million km²) globally. Similar to all wildland fires, 84.7% of global exposures to intense wildland fires occurred in Africa, while a significantly higher proportion was observed in North America (6.0% for intense fires compared to 1.2% for all wildland fires). Regional impacts of intense fires were pronounced in North America, Oceania and Europe, accounting for 3.1% (14.1%), 2.7% (15.2%), and 2.0% (4.1%) of all population exposures to wildland fire (burned area) in each continent (Fig. S11). See Fig. S12 for the contribution of population dynamics to exposure to intense fires.

Population exposure to intense fires remained stable globally from 2002-2021 (Fig. 3B), with mixed non-significant trends in different continents (Fig. S11). Global burned area declined both for intense fires (-47.2%; $p=0.12$) and other fires (-24.9%, $p < 0.01$) (Figs. 3B,C) in the past two decades. Regional trends, however, showed nuanced differences; for example, burned area by intense fires increased significantly in North America and South America from 2002-2021 (Fig. S11).

In wildfire disaster-prone regions (2) – Western North America, the Mediterranean, and Southern Australia, where loss of life and property to wildfires has occurred several times – 253,400 people were exposed to intense fires from 2002-2021, accounting for 32.0%, 4.1%, and 2.8%, of all exposures to wildland fire in each region, respectively (Fig. S13), far in excess of the 0.6% of global exposures to intense fires. Population exposure to intense fires showed non-significant increases in Western North America and Southern Australia from 2002-2021 (Fig. S13).

Discussion

Fires are complex social-ecological phenomena, and using burned area alone – often a common indicator of fire activity – fails to capture their full impact (20). The contrasting trends of burned area and exposure to wildland fire indicate multidimensional relationships between fire, humans, and landscapes (9, 10, 25, 32, 33). Specifically,

land fragmentation (24) induced by increasing human presence and expanding commercial agriculture in Africa reduced regional burned area, but co-located more humans with wildland fires – including fire use – and increased their exposure (3). Increasing human exposure to wildland fire in North America is due to both expansion of the WUI (19) and increasing burned area (14), and declining exposures in Europe is due to depopulation of rural areas (34), although climatic trends promoted more extreme potential fire behavior in both regions (35–37). In South America and Asia, population dynamics increased human exposure to wildland fire, despite a decline in burned areas from 2002-2021.

We found an increasing co-location of wildland fire and human settlements globally and in every continent. Global exposure density roughly doubled in the past two decades, mostly driven by trends in Africa. These trends are in accord with the expansion of the global WUI (22, 23). WUI expansion not only places humans at a higher risk to the negative effects of fire (19), but also increases the number of anthropogenic ignitions of fire (38).

Effects of intense fires were particularly significant in North America, Oceania, and Europe coinciding with global hotspots in fire disasters (2). Various wildfire mitigation strategies can be implemented in disaster-prone regions to reduce human impacts from fire exposure (39), as well as to limit human-caused fires (40). In wildfire-prone WUI areas, home hardening—incorporating structural and landscaping modifications to enhance fire resistance—is essential. There is also a need for increased intentional fire use as a vegetation management tool to mitigate wildfire disasters (41–43).

Our wildland fire exposure estimates focus solely on direct human exposure and do not account for indirect impacts such as exposure to smoke and postfire debris flow and flood (44). As a result, these figures are conservative, and the total number of people affected by wildland fires—whether directly or indirectly—is likely much higher. We note that extreme fire weather and fire behavior are expected to intensify in a warming climate (45), and fires are projected to increasingly impact humans especially given the population increases and WUI expansions (23).

Finally, our analysis indicates that the geography of wildfire disasters is distinctly different from the geography of major human exposures to wildland fires globally (2). Most wildfire disasters with the highest fatalities and societal losses has occurred in areas with Mediterranean climates in Australia, Southern Europe, South Africa, Western North America, and the west coast of South America (2, 26, 46). These wildfire disaster regions are not global hotspots in human exposure to wildland fire. By contrast, our analysis found the highest cumulative area burned and the highest human exposure to

wildland fire (both cumulative exposure and exposure density) in regions that have had relatively few wildfire disasters, such as in tropical Africa (2).

References

1. C. E. Synolakis, G. M. Karagiannis, Wildfire risk management in the era of climate change. *PNAS Nexus* **3**, pgae151 (2024).
2. D. M. Bowman *et al.*, Human exposure and sensitivity to globally extreme wildfire events. *Nat. Ecol. Evol.* **1**, 0058 (2017).
3. M. W. Jones *et al.*, Global and Regional Trends and Drivers of Fire Under Climate Change. *Rev. Geophys.* **60**, e2020RG000726 (2022).
4. C. A. Kolden, J. T. Abatzoglou, M. W. Jones, P. Jain, Wildfires in 2023. *Nat. Rev. Earth Environ.* **5**, 238–240 (2024).
5. R. Xu *et al.*, Global, regional, and national mortality burden attributable to air pollution from landscape fires: a health impact assessment study. *The Lancet* **404**, 2447–2459 (2024).
6. J. T. Abatzoglou, A. P. Williams, L. Boschetti, M. Zubkova, C. A. Kolden, Global patterns of interannual climate–fire relationships. *Glob. Change Biol.* **24**, 5164–5175 (2018).
7. W. M. Jolly *et al.*, Climate-induced variations in global wildfire danger from 1979 to 2013. *Nat. Commun.* **6**, 7537 (2015).
8. J. K. Balch *et al.*, Warming weakens the night-time barrier to global fire. *Nature* **602**, 442–448 (2022).
9. D. I. Kelley, I. Bistinas, R. Whitley, C. Burton, T. R. Marthews, N. Dong, How contemporary bioclimatic and human controls change global fire regimes. *Nat. Clim. Change* **9**, 690–696 (2019).
10. M. Forkel *et al.*, Emergent relationships with respect to burned area in global satellite observations and fire-enabled vegetation models. *Biogeosciences* **16**, 57–76 (2019).
11. J. K. Balch, B. A. Bradley, J. T. Abatzoglou, R. C. Nagy, E. J. Fusco, A. L. Mahood, Human-started wildfires expand the fire niche across the United States. *Proc. Natl. Acad. Sci.* **114**, 2946–2951 (2017).
12. A. Ganteaume, A. Camia, M. Jappiot, J. San-Miguel-Ayanz, M. Long-Fournel, C. Lampin, A Review of the Main Driving Factors of Forest Fire Ignition Over Europe. *Environ. Manage.* **51**, 651–662 (2013).
13. T. A. J. Janssen, M. W. Jones, D. Finney, G. R. van der Werf, D. van Wees, W. Xu, S. Veraverbeke, Extratropical forests increasingly at risk due to lightning fires. *Nat. Geosci.* **16**, 1136–1144 (2023).

- 269 14. J. T. Abatzoglou, C. A. Kolden, J. K. Balch, B. A. Bradley, Controls on interannual
270 variability in lightning-caused fire activity in the western US. *Environ. Res. Lett.* **11**, 045005
271 (2016).
- 272 15. A. D. Syphard, K. C. Clarke, J. Franklin, Simulating fire frequency and urban growth in
273 southern California coastal shrublands, USA. *Landsc. Ecol.* **22**, 431–445 (2007).
- 274 16. D. M. J. S. Bowman *et al.*, The human dimension of fire regimes on Earth. *J. Biogeogr.* **38**,
275 2223–2236 (2011).
- 276 17. C. J. McDonald, G. R. McPherson, Creating Hotter Fires in the Sonoran Desert:
277 Buffelgrass Produces Copious Fuels and High Fire Temperatures. *Fire Ecol.* **9**, 26–39
278 (2013).
- 279 18. N. Andela, G. R. Van Der Werf, Recent trends in African fires driven by cropland
280 expansion and El Niño to La Niña transition. *Nat. Clim. Change* **4**, 791–795 (2014).
- 281 19. V. C. Radeloff *et al.*, Rapid growth of the US wildland-urban interface raises wildfire risk.
282 *Proc. Natl. Acad. Sci. U. S. A.* **115**, 3314–3319 (2018).
- 283 20. A. Modaresi Rad *et al.*, Human and infrastructure exposure to large wildfires in the United
284 States. *Nat. Sustain.* **6**, 1343–1351 (2023).
- 285 21. F. Schug *et al.*, The global wildland–urban interface. *Nature* **621**, 94–99 (2023).
- 286 22. B. Chen *et al.*, Wildfire risk for global wildland–urban interface areas. *Nat. Sustain.* **7**, 474–
287 484 (2024).
- 288 23. W. Tang, C. He, L. Emmons, J. Zhang, Global expansion of wildland-urban interface (WUI)
289 and WUI fires: insights from a multiyear worldwide unified database (WUWUI). *Environ.*
290 *Res. Lett.* **19**, 044028 (2024).
- 291 24. N. Andela *et al.*, A human-driven decline in global burned area. *Science* **356**, 1356–1362
292 (2017).
- 293 25. M. W. Jones *et al.*, Global rise in forest fire emissions linked to climate change in the
294 extratropics. *Science* **386**, ead15889 (2024).
- 295 26. C. X. Cunningham, G. J. Williamson, D. M. Bowman, Increasing frequency and intensity of
296 the most extreme wildfires on Earth. *Nat. Ecol. Evol.* **8**, 1420–1425 (2024).
- 297 27. N. Andela *et al.*, The Global Fire Atlas of individual fire size, duration, speed and direction.
298 *Earth Syst. Sci. Data* **11**, 529–552 (2019).
- 299 28. A. J. Tatem, WorldPop, open data for spatial demography. *Sci. Data* **4**, 170004 (2017).
- 300 29. M. Friedl, D. Sulla-Menashe, MCD12Q1 MODIS/Terra+ aqua land cover type yearly L3
301 global 500m SIN grid V006. *No Title* (2019).
- 302 30. L. Giglio, L. Boschetti, D. P. Roy, M. L. Humber, C. O. Justice, The Collection 6 MODIS
303 burned area mapping algorithm and product. *Remote Sens. Environ.* **217**, 72–85 (2018).

- 304 31. E. Vermote, R. Wolfe, MODIS/Aqua Surface Reflectance Daily L2G Global 1km and 500m
305 SIN Grid V061, NASA Land Processes Distributed Active Archive Center (2021);
306 <https://doi.org/10.5067/MODIS/MYD09GA.061>.
- 307 32. W. Knorr, A. Arneth, L. Jiang, Demographic controls of future global fire risk. *Nat. Clim.*
308 *Change* **6**, 781–785 (2016).
- 309 33. C. Wu, S. Venevsky, S. Sitch, L. M. Mercado, C. Huntingford, A. C. Staver, Historical and
310 future global burned area with changing climate and human demography. *One Earth* **4**,
311 517–530 (2021).
- 312 34. F. Seijo, R. Gray, Pre-Industrial Anthropogenic Fire Regimes in Transition: The Case of
313 Spain and its Implications for Fire Governance in Mediterranean Type Biomes. *Hum. Ecol.*
314 *Rev.* **19**, 58–69 (2012).
- 315 35. P. Jain, Q. E. Barber *et al.*, Drivers and Impacts of the Record-Breaking 2023 Wildfire
316 Season in Canada. *Nat. Commun.* **15**, 6764 (2024).
- 317 36. M. Turco, S. Jerez, S. Augusto, P. Tarín-Carrasco, N. Ratola, P. Jiménez-Guerrero, R. M.
318 Trigo, Climate drivers of the 2017 devastating fires in Portugal. *Sci. Rep.* **9**, 13886 (2019).
- 319 37. C. S. Juang, A. P. Williams, J. T. Abatzoglou, J. K. Balch, M. D. Hurteau, M. A. Moritz,
320 Rapid Growth of Large Forest Fires Drives the Exponential Response of Annual Forest-
321 Fire Area to Aridity in the Western United States. *Geophys. Res. Lett.* **49**, e2021GL097131
322 (2022).
- 323 38. A. D. Syphard, V. C. Radeloff, T. J. Hawbaker, S. I. Stewart, Conservation Threats Due to
324 Human-Caused Increases in Fire Frequency in Mediterranean-Climate Ecosystems.
325 *Conserv. Biol.* **23**, 758–769 (2009).
- 326 39. T. Schoennagel *et al.*, Adapt to more wildfire in western North American forests as climate
327 changes. *Proc. Natl. Acad. Sci.* **114**, 4582–4590 (2017).
- 328 40. M. Kumar, A. AghaKouchak, J. T. Abatzoglou, M. Sadegh, Compounding effects of climate
329 change and WUI expansion quadruple the likelihood of extreme-impact wildfires in
330 California. *Npj Nat. Hazards* **2**, 17 (2025).
- 331 41. P. M. Fernandes, H. S. Botelho, A review of prescribed burning effectiveness in fire hazard
332 reduction. *Int. J. Wildland Fire* **12**, 117–128 (2003).
- 333 42. P. M. Fernandes *et al.*, Prescribed burning in southern Europe: developing fire
334 management in a dynamic landscape. *Front. Ecol. Environ.* **11** (2013).
- 335 43. C. A. Kolden, We're not doing enough prescribed fire in the Western United States to
336 mitigate wildfire risk. *Fire* **2**, 30 (2019).
- 337 44. R. Xu *et al.*, Global population exposure to landscape fire air pollution from 2000 to 2019.
338 *Nature* **621**, 521–529 (2023).
- 339 45. M. Turco *et al.*, I. Cvijanovic, Anthropogenic climate change impacts exacerbate summer
340 forest fires in California. *Proc. Natl. Acad. Sci.* **120**, e2213815120 (2023).

- 341 46. D. M. J. S. Bowman *et al.*, Human–environmental drivers and impacts of the globally
342 extreme 2017 Chilean fires. *Ambio* **48**, 350–362 (2019).
- 343 47. S. T. Seydi, J. Abatzoglou, M. Sadegh, M. Jones, Global human exposure to wildland fires
344 dataset: 2002-2021, Zenodo (2025); <https://doi.org/10.5281/zenodo.15549088>.
- 345 48. L. Giglio, J. Descloitres, C. O. Justice, Y. J. Kaufman, An enhanced contextual fire
346 detection algorithm for MODIS. *Remote Sens. Environ.* **87**, 273–282 (2003).
- 347 49. J. Eidenshink, B. Schwind, K. Brewer, Z.-L. Zhu, B. Quayle, S. Howard, A Project for
348 Monitoring Trends in Burn Severity. *Fire Ecol.* **3**, 3–21 (2007).
- 349 50. F. R. Stevens, A. E. Gaughan, C. Linard, A. J. Tatem, Disaggregating census data for
350 population mapping using random forests with remotely-sensed and ancillary data. *PloS*
351 *One* **10**, e0107042 (2015).
- 352 51. A. J. Tatem *et al.*, Millennium development health metrics: where do Africa’s children and
353 women of childbearing age live? *Popul. Health Metr.* **11**, 11 (2013).
- 354 52. A. Sorichetta, G. M. Hornby, F. R. Stevens, A. E. Gaughan, C. Linard, A. J. Tatem, High-
355 resolution gridded population datasets for Latin America and the Caribbean in 2010, 2015,
356 and 2020. *Sci. Data* **2**, 1–12 (2015).
- 357 53. L. Giglio, W. Schroeder, J. V. Hall, C. O. Justice, MODIS collection 6 active fire product
358 user’s guide revision A. *Dep. Geogr. Sci. Univ. Md.* **9** (2015).
- 359 54. L. Giglio, W. Schroeder, J. V. Hall, C. O. Justice, MODIS Collection 6 and Collection 6.1
360 Active Fire Product User’s Guide. *Natl. Aeronaut. Space Adm. Wash. DC USA*, 64 (2021).
- 361 55. T. Liu *et al.*, Missing emissions from post-monsoon agricultural fires in northwestern India:
362 regional limitations of MODIS burned area and active fire products. *Environ. Res.*
363 *Commun.* **1**, 011007 (2019).
- 364 56. L. Boschetti, D. P. Roy, L. Giglio, H. Huang, M. Zubkova, M. L. Humber, Global validation
365 of the collection 6 MODIS burned area product. *Remote Sens. Environ.* **235**, 111490
366 (2019).
- 367 57. L. Giglio, J. T. Randerson, G. R. Van Der Werf, Analysis of daily, monthly, and annual
368 burned area using the fourth-generation global fire emissions database (GFED4). *J.*
369 *Geophys. Res. Biogeosciences* **118**, 317–328 (2013).
- 370 58. A. Modaresi Rad *et al.*, Social vulnerability of the people exposed to wildfires in U.S. West
371 Coast states. *Sci. Adv.* **9**, eadh4615 (2023).
- 372 59. R. Jing, S. Heft-Neal *et al.*, Global population profile of tropical cyclone exposure from
373 2002 to 2019. *Nature* **626**, 549–554 (2024).
- 374 60. M. K. Adu, B. Agyapong, V. I. O. Agyapong, Children’s Psychological Reactions to
375 Wildfires: a Review of Recent Literature. *Curr. Psychiatry Rep.* **25**, 603–616 (2023).
- 376 61. Google Maps (2024). <https://www.google.com/maps>.

Acknowledgments

Funding: Joint Fire Science Program grant number L21AC10247 (MS, JTA), the U.S. National Science Foundation award OAI-2019762 (JTA) and 2429021 (MS), Boise State University Healthy Idaho Initiative (MS), Natural Environment Research Council (NERC) grant NE/V01417X/1 (MWJ). Any use of trade, firm or product names is for descriptive purposes only and does not imply endorsement by the US Government.

Author contributions:

Conceptualization: MS, JTA

Methodology: MS, JTA, STS, MWJ

Investigation: STS, JTA, MS, MWJ, CK, GF, MDH, AA, CHL, CM

Visualization: STS

Funding acquisition: MS, JTA

Project administration: MS

Supervision: MS, JTA

Writing – original draft: STS, JTA, MS, MWJ, CK, GF, MDH, AA, CHL, CM

Writing – review & editing: STS, JTA, MS, MWJ, CK, GF, MDH, AA, CHL, CM

Competing interests: The authors declare no competing interests.

Data and materials availability:

All data and materials used in this study are available at Seydi et al. (47).

Supplementary Materials

Materials and Methods

Supplementary Text, Supplementary sections S1-S4

Figs. S1 to S20

Table S1

Figure 1. Human exposure to wildland fires from 2002-2021. Cumulative human exposure to wildland fires (A) and cumulative burned area (C) from 2002-2021 at the country level. Share of continents in wildland fire exposure (B) and burned areas (D). Percentage of population exposed to wildland fire from 2002-2021 in each country (E) and continent (F). K stands for one thousand and M stands for one million.

Figure 2. Population dynamics' contribution to wildland fire exposure. Observed (black) and counterfactual constant-population scenario (purple) exposures to wildland fire from 2002-2021 across the globe (A) and in each continent (B-G), respectively. In observed exposures, both population and burned area data are dynamic; while in counterfactual exposures, population is fixed at year 2002 but burned area is dynamic as observed.

Figure 3. Population exposure to intense wildland fires. Location, occurrence year and exposure to individual intense fires that exposed more than 100 people from 2002-2021 (A), and share of continents in global population exposure to intense fires (B). Trends in annual exposure to and burned area by intense fires globally (C-D).