

## **REFERENCES**

- Amthor, H., Christ, B., Weil, M. and Patel, K. (1998). The importance of timing differentiation during limb muscle development. *Curr Biol* 8: 642-52.
- Basyuk, E., Suavet, F., Doglio, A., Bordonne, R., and Bertrand, E. (2003). Human let-7 stem-loop precursors harbor features of RNase III cleavage products. *Nucleic Acids Res.* 31, 6593–6597.
- Beltran, A.C., Dawson P.E. and Gottesfeld, J.M. Role of DNA sequence in the binding specificity of synthetic basic-helix–loop–helix domains, *Chem Bio Chem* 6(2005), pp. 104–113
- Bendall, A.J., Ding, J., Hu, G., Shen, M.M. and Abate-Shen, C. (1999). Msx1 antagonizes the myogenic activity of Pax3 in migrating limb muscle precursors. *Development* 126: 4965-76.
- Blais, A., Tsikitis, M., Acosta-Alvear, D., Sharan, R., Kluger, Y. & Dynlacht, B. D. (2005) *Genes Dev.* 19, 553–569
- Brand-Saberi, B., Muller, T.S., Wilting, J., Christ, B. and Birchmeier, C. (1996). Scatter factor/hepatocyte growth factor (SF/HGF) induces emigration of myogenic cells at interlimb level in vivo. *Dev Biol* 179: 303-8.
- Bryson-Richardson, R.J. and Currie, P.D. (2008) The genetics of vertebrate myogenesis. *Nat. Rev. Genet.* 9:632–646.
- Buckingham, M. (2001). Skeletal muscle formation in vertebrates. *Curr. Opin. Genet. Dev.* 11,440 -448.
- Buckingham, M., Bajard, L., Chang, T., Daubas, P., Hadchouel, J., Meilhac, S., Montarras, D., Rocancourt, D., Relaix, F., 2003. The formation of skeletal muscle: from somite to limb. *J. Anat.* 202, 59–68.
- Caudy, A.A., Myers, M., Hannon, G.J., and Hammond, S.M. (2002) Fragile X-related protein and VIG associate with the RNA interference machinery. *Genes Dev.* 16, 2491–2496

- Cerutti, L., Mian, N., and Bateman, A. (2000). Domains in gene silencing and cell differentiation proteins: the novel PAZ domain and redefinition of the Piwi domain. *Trends Biochem. Sci.* 25, 481–482.
- Chen, J.F., Mandel, E.M., Thomson, J.M., Wu, Q., Callis, T.E., Hammond, S.M., Conlon, F.L., Wang, D.Z., 2006. The role of microRNA-1 and microRNA-133 in skeletal muscle proliferation and differentiation. *Nat. Genet.* 38, 228–233.
- Christ, B., and Ordahl, C.P. (1995). Early stages of chick somite development. *Anat. Embryol.* 191, 381–396.
- Christ, B., Jacob, H.J. and Jacob, M. (1977). Experimental analysis of the origin of the wing musculature in avian embryos. *Anat Embryol (Berl)* 150: 171-86.
- Bartel D.P., (2004) MicroRNAs: Genomics, biogenesis, mechanism, and function, *Cell* 116, pp281-297
- Davin Townley-Tilson, W., H., Callis, T. E., Wang, D. (2009) MicroRNAs 1, 133 and 206: Critical Factor of Skeletal and Cardiac Muscle Development, Function and Disease. *The international Journal of Biochemistry & Cell Biology* 235:2185-2191.
- Delfini, M.C., and Duprez, D. (2000). Paraxis is expressed in myoblasts during their migration and proliferation in the chick limb bud. *Mech Dev* 96: 247-51.
- Duprez, D. (2002). Signals regulating muscle formation in the limb during embryonic development, *Int. J. Dev. Biol.* 46: pp. 915–925.
- Fallon, J.F., and Kelley, R.O. (1977). Ultrastructural analysis of the apical ectodermal ridge during vertebrate limb morphogenesis. II. Gap junctions as distinctive ridge structures common to birds and mammals. *J. Embryol. Exp. Morphol.* 41, 223–232.
- Gilbert, S. F., (2000) Paraxial and Intermediate Mesoderm. In: Sinauer A.D. (eds) *Developmental Biology* pp447-470, Sinauer Associates Inc., Publishers, Sunderland, Massachusetts.
- Hamburger, V., Hamilton, H.L. (1951). A series of normal stages in the development of the chick embryo. *J. Morphol.* 88, 49–92.

- Hammond, S.M., Boettcher, S., Caudy, A.C., Kobayashi, R., and Hannon, G.J. (2001). Argonaute2, a link between genetic and bio-chemical analyses of RNAi. *Science* 293, 1146–1150.
- Hutvagner, G., and Zamore, P.D. (2002). A microRNA in a multiple turnover RNAi enzyme complex. *Science* 297, 2056–2060.
- Ivarie R. (1993). The Myogenic Regulatory Factors and the Early Steps of Muscle Cell Differentiation and Growth *J. Anim. Sci.* 71(Suppl. 2): 23-32
- Johnson, R.L., and Tabin, C.J. (1997). Molecular models for vertebrate limb development. *Cell* 90, 979–990
- Kardon, G., (1998). Muscle and tendon morphogenesis in the avian hind limb. *Development* **125**: pp. 4019–4032.
- Kavlock R. J., Setzer R.W. (1996) The road to embryologically based dose-response models. *Environ Health Perspect* 104 (Suppl 1):107–121.
- Kawakami, Y., Capdevila, J., Buscher, D., Itoh, T., Esteban, C.R., and Belmonte, J.C. (2001). Wnt signals control FGF-dependent limb initiation and AER induction in the chick embryo. *Cell* 104, 891–900.
- Keynes, R., and C. Stern. (1988). Mechanisms of vertebrate segmentation. *Development* 103:413.
- Khvorova, A., Reynolds, A., and Jayasena, S.D. (2003). Functional siRNAs and miRNAs exhibit strand bias. *Cell* 115, 209–216.
- Kim, H.K., Lee, Y.S., Sivaprasad, U., Malhotra, A., Dutta, A. (2006). Muscle-specific microRNA miR-206 promotes muscle differentiation. *J. Cell. Biol.* 174, 677–687.
- Kuersten, S., and Goodwin, E.B. (2003). The power of the 3' UTR: translational control and development. *Nat. Rev. Genet.* 4: 626–637.
- Lagos-Quintana, M., Rauhut, R., Lendeckel, W., and Tuschl, T. (2001). Identification of novel genes coding for small expressed RNAs. *Science* 294, 853–858.
- Lau, N.C., Lim, L.P., Weinstein, E.G., and Bartel, D.P. (2001). An abundant class of tiny RNAs with probable regulatory roles in *Caenorhabditis*

- Laufer, E., Nelson, C.E., Johnson, R.L., Morgan, B.A., and Tabin, C. (1994). Sonic hedgehog and Fgf-4 act through a signaling cascade and feedback loop to integrate growth and patterning of the developing limb bud. *Cell* 79, 993–1003
- Lee, Y., Ahn, C., Han, J., Choi, H., Kim, J., Yim, J., Lee, J., Provost, P., Radmark, O., Kim, S., and Kim, V.N. (2003). The nuclear RNase III Drosha initiates microRNA processing. *Nature* 425, 415–419.
- Lee, Y., Jeon, K., Lee, J.T., Kim, S., and Kim, V.N. (2002). MicroRNA maturation: stepwise processing and subcellular localization. *EMBO J.* 21, 4663–4670.
- Lewis, B.P., Shih, I., Jones-Rhoades, M.W., Bartel, D.P., and Burge, C.B. (2003). Prediction of mammalian microRNA targets. *Cell* 115, 787–798.
- Li, X., C.S. Blagden, H. Bildsoe, M.A. Bonnin, D. Duprez, and S.M. Hughes. 2004. Hedgehog can drive terminal differentiation of amniote slow skeletal muscle. *BMC Dev. Biol.* 4:9.
- Lingel, A., Simon, B., Izaurralde, E., and Sattler, M. (2003). Structure and nucleic-acid binding of the *Drosophila* Argonaute 2 PAZ domain. *Nature* 426, 465–469.
- Ludolph DC and Konieczny SF. Transcription factor families: muscling in on the myogenic program. *FASEB J* 9: 1595–1604, 1995
- Nelson, C.E., Morgan, B.A., Burke, A.C., Laufer, E., DiMambro, E., Murtaugh, L.C., Gonzales, E., Tessarollo, L., Parada, L.F., and Tabin, C. (1996). Analysis of Hox gene expression in the chick limb bud. *Development* 122, 1449–1466.
- Ordahl, C. P., and Le Douarin, N. M. (1992). Two myogenic lineages within the developing somite. *Development* 114, 339–353.
- Ott, M., Bober, E., Lyons, G., Arnold, H. and Buckingham, M. (1991). Early expression of the myogenic regulatory gene, myf-5, in precursor cells of skeletal muscle in the mouse embryo. *Development* 111, 1097-1107
- Parr, B.A. and McMahon, A.P. (1995). Dorsalizing signal Wnt-7a required for normal polarity of D-V and A-P axes of mouse limb. *Nature* 374: 350-3.

- Pautou, M.P., Hedayat, I. and Kieny, M. (1982). The pattern of muscle development in the chick leg. *Arch Anat Microsc Morphol Exp* 71: 193-206.
- Rao, P.K., Kumar, R.M., Farkhondeh, M., Baskerville, S., Lodish, H.F. (2006). Myogenic factors that regulate expression of muscle-specific microRNAs. *Proc. Natl. Acad. Sci. U S A* 103: 8721–8726.
- Robson, L.G., Kara, T., Crawley, A. and Tickle, C. (1994). Tissue and cellular patterning of the musculature in chick wings. *Development* 120: 1265-76.
- Rong, P., TeiUet, M. and Ziller, C. (1990). Differential influence of the neural primordium on muscle development in myotome and limb. In *The Avian Model in Developmental Biology: From Organism to Genes* (ed. N. Le Douarin, F. Dieterlen-Lievre and J. Smith) pp. 153-164 CNRS, Paris.
- Scaal, M., Christ, B., (2008). Wnt Signalling in Somite Development. *Ann Anat* 190(3): 208-222
- Schramm, C. and Solourh, M. (1990). The formation of premuscle masses during chick wing bud development. *Anat Embryol (Berl)* 182: 235-47.
- Shijie Li, Michael P. Czubryt, John McAnally, Rhonda Bassel-Duby, James A. Richardson, Franziska F. Wiebel, Alfred Nordheim, and Eric N. Olson. (2005) Requirement for serum response factor for skeletal muscle growth and maturation revealed by tissue-specific gene deletion in mice. *PNAS*, 102(4): 1082–1087
- Simionato E, V Ledent, G Richards, M Thomas-Chollier, P Kerner, D Coornaert, BM Degnan & M Vervoort (2007), Origin and diversification of the basic helix-loop-helix gene family in metazoans: Insights from comparative genomics. *BMC Evol. Bio.* 7: 33. Glossary, Homoscleromorpha, Porifera.
- Summerbell, D., Lewis, J.H., and Wolpert, L. (1973). Positional information in chick limb morphogenesis. *Nature* 244, 492–496.
- Sweetman D, Goljanek K, Rathjen T, Oustanina S, Braun T, Dalmay T, Münsterberg A (2008). Specific requirements of MRFs for the expression of muscle specific microRNAs, miR-1, miR-206 and miR-133. *Dev Biol*, 321(2):491-499

TeiUet, M. and Le Douarin, N. (1983). Consequences of neural tube and notochord excision on the development of the peripheral nervous system in the chick embryo. *Devi. Biol.* 98, 192-211.

Yi, R., Qin, Y., Macara, I.G., and Cullen, B.R. (2003). Exportin-5 mediates the nuclear export of pre-microRNAs and short hairpin RNAs. *Genes Dev.* 17, 3011–3016.

Zhi, Q., Huang, R., Christ, B. and Brand-Saberi, B. (1996). Participation of individual brachial somites in skeletal muscles of the avian distal wing. *Anat Embryol (Berl)* 194: 327-39.

<http://www.the-scientist.com/blog/display/55439/>