

# Buildings of classical groups and centralizers of Lie algebra elements

P. BROUSSOUS AND S. STEVENS

*Février 2004*

## Abstract

Let  $F_o$  be a non-archimedean locally compact field of residual characteristic not 2. Let  $G$  be a classical group over  $F_o$  (with no quaternionic algebra involved) which is not of type  $A_n$  for  $n > 1$ . Let  $\beta$  be an element of the Lie algebra  $\mathfrak{g}$  of  $G$  that we assume semisimple for simplicity. Let  $H$  be the centralizer of  $\beta$  in  $G$  and  $\mathfrak{h}$  its Lie algebra. Let  $I$  and  $I_\beta^1$  denote the (enlarged) Bruhat-Tits buildings of  $G$  and  $H$  respectively. We prove that there is a natural set of maps  $j_\beta : I_\beta^1 \rightarrow I$  which enjoy the following properties: they are affine,  $H$ -equivariant, map any apartment of  $I_\beta^1$  into an apartment of  $I$  and are compatible with the Lie algebra filtrations of  $\mathfrak{g}$  and  $\mathfrak{h}$ . In a particular case, where this set is reduced to one element, we prove that  $j_\beta$  is characterized by the last property in the list. We also prove a similar characterization result for the general linear group.

## Introduction

In this paper we establish new functoriality properties between affine Bruhat-Tits buildings of classical reductive groups over local fields. More precisely let  $F_o$  be a non-archimedean local field of residual characteristic not 2 and  $G$  be the group of  $F_o$ -rational points of a classical group defined over  $F_o$ . We assume that  $G$  is the isometry group of an  $\varepsilon$ -hermitian form over an  $F$ -vector space, where  $F$  is a (commutative) extension of  $F_o$  of degree less than 2. We denote by  $\mathfrak{g}$  the Lie algebra of  $G$  and by  $I$  its affine building. Let  $\beta$  be an element of  $\mathfrak{g}$  that we assume to be semisimple for simplicity. Let  $H$  be the centralizer of  $\beta$  in  $G$ . Then  $H$  is the group of  $F_o$ -rational points of a product of groups of the form  $\text{Res}_{E_o/F_o} \mathbf{H}_i$ , where  $E_o/F_o$  is a field extension and where  $\text{Res}$  denotes the functor of restriction of scalars. Here  $i$  runs over a finite set  $\tilde{J}$ . Each  $\mathbf{H}_i$  is either a classical group as above or a general linear group. We denote by  $J_+ \subset \tilde{J}$  the (possibly empty) subset of indices corresponding to linear groups. We denote by  $\mathfrak{h}$  the Lie algebra of  $H$  and by  $I_\beta^1$  its (enlarged) affine building. Then there is a natural set of maps  $j_\beta : I_\beta^1 \rightarrow I$  which depend on identifications of the enlarged buildings of  $\mathbf{H}_i$ ,  $i \in J_+$ , with certain sets of lattice functions (see §4 below). In particular, when  $J_+ = \emptyset$ , there is a natural choice of  $j_\beta$ . The maps  $j_\beta$  enjoys the following properties:

- a) They are affine.
- b) They are  $H$ -equivariant.
- c) They map any apartment of  $I_\beta^1$  into an apartment of  $I$ .
- d) They are compatible with the Lie algebra filtrations of  $\mathfrak{g}$  and  $\mathfrak{h}$  (cf. §9).

In [BL] it was proved that when  $G$  is the general linear group and  $\beta$  is an elliptic element then, replacing the buildings by the non-enlarged buildings, there is such a natural map  $j_\beta$  satisfying the conditions above. It is actually characterized by properties

a) and b). However in the case of a classical group (and assuming that  $J_+ = \emptyset$ ) it is no longer true that properties a), b) and c) characterize  $j_\beta$ . The simplest counter-example is the following. Consider the case of  $G = \mathrm{Sp}_2(F_o) = \mathrm{SL}(2, F_o)$ . One may choose  $\beta$  is such a way that  $H$  is  $E^1$ , the group of norm 1 elements of a ramified quadratic extension  $E/F_o$ . Then  $I_\beta^1$  is reduced to a point and fixing a map  $j_\beta$  satisfying a) b) and c) amounts to choosing a point in  $I$  fixed by the torus  $E^1$ . But  $E^1$  is contained in an Iwahori subgroup of  $G$  and therefore fixes a chamber of  $I$ .

We prove that, in the case of a general linear group and of an elliptic element  $\beta$ , the map  $j_\beta$  of [BL] is actually characterized by property d). In the case of a classical group, we also prove that if  $J_+ = \emptyset$  and if a technical condition on  $\beta$  is satisfied then  $j_\beta$  is characterized by condition d). We conjecture that when  $J_+ = \emptyset$  then  $j_\beta$  is indeed characterized by property d).

In this work, we do not actually assume  $\beta$  to be semisimple but only to satisfy a weaker assumption (see hypothesis (H1) of §5). Such elements naturally appear in the generalization of the theory of strata due to Bushnell and Kutzko [BK] to the case of classical groups (see the work of the second author [S1], [S2]. Even though the work of the second author does not use the theory of affine buildings in a straightforward way (it uses the equivalent language of hereditary orders), the existence and properties of the maps  $j_\beta$  are applied to the representation theory of  $G$ , particularly in [S2].

The paper is organized as follows. In §2 we recall the structure of the maximal split tori of  $G$ . In §3,4, using ideas of Bruhat and Tits, we give a model of the affine building of  $G$  in terms of “self-dual lattice functions”. In §5 we study the centralizers in  $\mathfrak{g}$  and  $G$  of the Lie algebra element  $\beta$ . The construction of the maps  $j_\beta$  is done in §6 and their properties are established in §7,8 and 9. In §10 We prove the uniqueness result for the general linear group and finally §11 is devoted to the uniqueness result in the classical group case.

We thank G. Henniart, B. Lemaire, G. Prasad and J.-K. Yu for stimulating discussions.

## 1. Notation

Here  $F_o$  is the ground field; it is assumed to be non-archimedean, locally compact and equipped with a discrete valuation  $v$  normalized in such a way that  $v(F_o^\times)$  is the additive group of integers. We assume that the residual characteristic of  $F_o$  is not 2. We fix a Galois extension  $F/F_o$  such that  $[F : F_o] \leq 2$  and set  $\sigma_F = \mathrm{id}_F$  if  $F = F_o$  and take  $\sigma_F$  to be the generator of  $\mathrm{Gal}(F/F_o)$  in the other case. We still denote by  $v$  the unique extension of  $v$  to  $F$ . We fix  $\varepsilon \in \{\pm 1\}$  and a finite dimensional left  $F$ -vector space  $V$ . Recall that a  $\sigma_F$ -skew form  $h$  on  $V$  is a  $\mathbb{Z}$ -bilinear map  $V \times V \rightarrow F$  such that

$$h(\lambda x, \mu y) = \lambda^{\sigma_F} \mu h(x, y) , \quad \lambda, \mu \in F, \quad x, y \in V .$$

Such a form is called  $\varepsilon$ -hermitian if  $h(y, x) = \varepsilon h(x, y)^{\sigma_F}$  for all  $x, y \in V$ . From now on we fix such an  $\varepsilon$ -hermitian form on  $V$  and we assume it is non-degenerate (the orthogonal of  $V$  is  $\{0\}$ ).

For  $a \in \mathrm{End}_F(V)$ , we denote by  $a^{\sigma_h} = a^\sigma$  the adjoint of  $a$  with respect to  $h$ , i.e. the unique  $F$ -endomorphism of  $V$  satisfying  $h(ax, y) = h(x, a^\sigma y)$  for all  $x, y \in V$ .

We denote by  $\mathbf{G}$  the simple algebraic  $F_o$ -group whose set of  $F_o$ -rational points  $G$  is formed of the  $g \in \mathrm{GL}_F(V)$  satisfying  $g.h = h$  (it is not necessarily connected). Here  $g.h$  is the form given by  $g.h(x, y) = h(gx, gy)$ ,  $x, y \in V$ .

We know ([Sch](6.6), page 260) that in the case  $\sigma_F \neq \mathrm{id}_F$ , we may reduce to the case  $\varepsilon = 1$ . So we have three possibilities:

- $\sigma_F = \mathrm{id}_F$  and  $\varepsilon = 1$ , the orthogonal case;
- $\sigma_F = \mathrm{id}_F$  and  $\varepsilon = -1$ , the symplectic case;
- $\sigma_F \neq \mathrm{id}_F$  and  $\varepsilon = 1$ , the unitary case.

We abbreviate  $\tilde{G} = \mathrm{GL}_F(V)$  and  $\tilde{\mathfrak{g}} = \mathrm{End}_F(V)$ .

## 2. The maximal split tori of $\mathbf{G}$

Recall that a subspace  $W \subset V$  is totally isotropic if  $h(W, W) = 0$  and that maximal such subspaces have the same dimension  $r$ , the Witt index of  $h$ . Set  $I = \{\pm 1, \pm 2, \dots, \pm r\}$  and  $I_o = \{(0, k) ; k = 1, \dots, n - 2r\}$ . We fix a *Witt decomposition* of  $V$ , that is

two maximal totally isotropic subspaces  $V_+$  and  $V_-$ ,  
bases  $(e_i)_{i=1, \dots, r}$ ,  $(e_{-i})_{i=1, \dots, r}$ ,  $(e_i)_{i \in I_o}$  of  $V_+$ ,  $V_-$  and  $V_o := (V_+ + V_-)^\perp$ ,  
such that

$$\begin{aligned} h(e_i, e_i) &= 0, \quad i \in I, \\ h(e_i, e_j) &= 0, \quad \text{for } i, j \in I \text{ with } j \neq -i \text{ or } i \in I, j \in I_o, \\ h(e_i, e_{-i}) &= 1, \quad \text{for } i \in I \text{ with } i > 0, \\ h(x, x) &\neq 0, \quad \text{for } x \in V_o \text{ and } x \neq 0. \end{aligned}$$

The Witt decomposition gives rise to a maximal  $F_o$ -split torus  $\mathbf{S}$  whose group of  $F_o$ -rational points is

$$S = \{s \in G ; se_i \in F_o e_i, \quad i \in I \text{ and } (s - \mathrm{Id})V_o = 0\} .$$

It has dimension  $r$ , the  $F_o$ -rank of  $\mathbf{G}$ . Conversely any maximal  $F_o$ -split torus of  $\mathbf{G}$  is obtained from a Witt decomposition as above. The centralizer  $\mathbf{Z}$  of  $\mathbf{S}$  in  $\mathbf{G}$  has for  $F_o$ -rational points

$$Z = \{z \in G ; ze_i \in F e_i, \quad i \in I \text{ and } zV_o = V_o\} .$$

For each  $i \in I$ , we have a morphism of algebraic groups  $a_i : \mathbf{Z} \rightarrow \mathrm{Res}_{F/F_o}(\mathbf{G}_m)$  given by  $ze_i = a_i(z)e_i$ . Note that  $a_{-i}(z) = a_i(z)^{-\sigma}$ . We also denote by  $a_i : \mathbf{S} \rightarrow \mathbf{G}_m / F_o$  the character obtained by restriction. We have  $a_i = -a_{-i}$  in  $X^*(\mathbf{S})$ , the  $\mathbb{Z}$ -module of rational characters of  $\mathbf{S}$ . The  $a_i$ ,  $i \in I$ ,  $i > 0$ , form a basis of  $X^*(\mathbf{S})$ .

The normalizer  $\mathbf{N}$  of  $\mathbf{Z}$  in  $\mathbf{G}$  is the sub-algebraic group whose  $F_o$ -rational points are the elements of  $G$  which stabilize  $X_o$  and permute the lines  $V_i = F e_i$ ,  $i \in I$ . The group  $N = \mathbf{N}(F_o)$  is the semidirect product of  $Z$  by the subgroup  $N'$  formed of the elements which permute the  $\pm e_i$ ,  $i \in I$ .

## 3. MM-norms and self-dual lattice-functions

We keep the notation as in the previous sections.

Recall that a *norm* on  $V$  is a map  $\alpha : V \rightarrow \mathbb{R} \cup \{\infty\}$  satisfying:

- i)  $\alpha(x + y) \geq \mathrm{Inf}(\alpha(x), \alpha(y))$ ,  $x, y \in V$ ,

- ii)  $\alpha(\lambda x) = v(\lambda) + \alpha(x)$ ,  $\lambda \in F$ ,  $x \in V$ ,
- iii)  $\alpha(x) = \infty$  if and only if  $x = 0$ .

We denote by  $\text{Norm}^1(V)$  the set of norms on  $V$ .

**Definition 3.1.** (cf. [BT](2.1)) *Let  $\alpha \in \text{Norm}^1(V)$ . We say that  $\alpha$  is dominated by  $h$  if*

$$\alpha(x) + \alpha(y) \leq v(h(x, y)) \text{ for all } x, y \in V .$$

*We say that  $\alpha$  is an MM-norm for  $h$  (maximinorante in french), if  $\alpha$  is a maximal element of the set of norms dominated by  $h$ .*

In [BT](2.5) an involution  $\bar{\phantom{x}}$  is defined on  $\text{Norm}^1(V)$  in the following way. If  $\alpha \in \text{Norm}^1(V)$ , then

$$\bar{\alpha}(x) = \inf_{y \in V} [v(h(x, y)) - \alpha(y)] , \quad x \in V .$$

We then have

**Proposition 3.2.** (cf. [BT](Prop. 2.5)) *An element  $\alpha$  of  $\text{Norm}^1(V)$  is an MM-norm if and only if  $\bar{\bar{\alpha}} = \alpha$ .*

We are going to describe the set  $\text{Norm}_h^1(V)$  of MM-norms in terms of self-dual lattice-functions. Recall [BL] that a lattice-function in  $V$  is a function  $\Lambda$  which maps a real number to an  $\mathfrak{o}_F$ -lattice in  $V$  and satisfies:

- i)  $\Lambda(r) \subset \Lambda(s)$  for  $r \geq s$ ,  $r, s \in \mathbb{R}$ ,
- ii)  $\Lambda(r + v(\pi_F)) = \mathfrak{p}_F \Lambda(r)$ ,  $r \in \mathbb{R}$ ,
- iii)  $\Lambda$  is left-continuous.

Here  $\mathfrak{o}_F$  denotes the ring of integers of  $F$ ,  $\mathfrak{p}_F$  the maximal ideal of  $\mathfrak{o}_F$  and  $\pi_F$  a uniformizer of  $F$ . As in [BL], we denote by  $\text{Latt}_{\mathfrak{o}_F}^1(V)$  (or by  $\text{Latt}^1(V)$  when no confusion may occur) the set of  $\mathfrak{o}_F$ -lattice-functions in  $V$ . Recall [BL] that  $\text{Norm}^1(V)$  and  $\text{Latt}^1(V)$  may be canonically identified in the following way. To  $\alpha \in \text{Norm}^1(V)$ , we attach the function  $\Lambda = \Lambda_\alpha$  given by

$$\Lambda(r) = \{x \in V ; \alpha(x) \geq r\} , \quad r \in \mathbb{R} .$$

Conversely a lattice-function  $\Lambda$  corresponds to the norm  $\alpha$  given by

$$\alpha(x) = \sup\{r ; x \in \Lambda(r)\} , \quad x \in V .$$

For a  $\Lambda \in \text{Latt}^1(V)$  and  $r \in \mathbb{R}$ , set

$$\Lambda(r+) = \bigcup_{s > r} \Lambda(s) .$$

For an  $\mathfrak{o}_F$ -lattice  $L$  in  $V$ , we define its dual  $L^\sharp = L^{\sharp h}$  by

$$L^\sharp = \{x \in V ; h(x, L) \subset \mathfrak{p}_F\} .$$

Finally, we define the dual  $\Lambda^\sharp = \Lambda^{\sharp h}$  of a lattice-function  $\Lambda$  by

$$\Lambda^\sharp(r) = [\Lambda((-r) +)]^\sharp , \quad r \in \mathbb{R} .$$

We say that a lattice function  $\Lambda$  is self dual if  $\Lambda^\sharp = \Lambda$  and we denote by  $\text{Latt}_h^1(V)$  the corresponding set.

**Proposition 3.3.** *Given a norm  $\alpha \in \text{Norm}^1(V)$ , we have  $\Lambda_{\bar{\alpha}} = \Lambda_{\alpha}^{\sharp}$ .*

**Corollary 3.4.** *Let  $\alpha$  be a norm on  $V$ . Then  $\alpha$  is an MM-norm if and only if the attached lattice-function  $\Lambda$  is self-dual.*

*Proof of Proposition.* Let  $x \in V$  and  $r \in \mathbb{R}$ . Then the fact that  $x \in \Lambda_{\bar{\alpha}}(r) \setminus \Lambda_{\bar{\alpha}}(r+)$  is equivalent to the following points:

- i)  $\bar{\alpha}(x) = r$ ;
- ii) there exists  $y \in V$  such that  $v(h(x, y)) - \alpha(y) = r$ , and for all  $y \in V$ , we have  $v(h(x, y)) - \alpha(y) \geq r$ ;
- iii) there exists  $y \in V$  such that  $v(h(x, y)) = 0$  and  $\alpha(y) = -r$ , and for all  $y \in V$  such that  $\alpha(y) > -r$ , we have  $v(h(x, y)) > 0$  (scale by a suitable power of a uniformizer  $\pi_F$ );
- iv) there exists  $y \in \Lambda_{\alpha}(-r) \setminus \Lambda_{\alpha}(-r+)$  such that  $h(x, y) \in \mathfrak{o}_F \setminus \mathfrak{p}_F$ , and for all  $y \in \Lambda_{\alpha}(-r+)$  we have  $h(x, y) \in \mathfrak{p}_F$ ;
- v)  $x \in \Lambda_{\alpha}^{\sharp}(r) \setminus \Lambda_{\alpha}^{\sharp}(r+)$ .

This proves that the two lattice-functions  $\Lambda_{\bar{\alpha}}$  and  $\Lambda_{\alpha}^{\sharp}$  share the same discontinuity points and that at those points they take the same values; so there are equal.

Let  $\text{Norm}^2 \tilde{\mathfrak{g}}$  (resp.  $\text{Latt}^2 \tilde{\mathfrak{g}}$ ) denote the  $\tilde{G}$ -set of square norms in  $\tilde{\mathfrak{g}}$  (resp. of square lattice-functions in  $\tilde{\mathfrak{g}}$ ; see [BT1] and [BL]). Recall that a lattice-function  $\Lambda^2$  in the  $F$ -vector space  $\tilde{\mathfrak{g}}$  is square if there exists  $\Lambda \in \text{Latt}^1(V)$  such that  $\Lambda^2 = \text{End}(\Lambda)$ , where

$$\text{End}(\Lambda)(r) = \{a \in \tilde{\mathfrak{g}} ; a\Lambda(s) \subset \Lambda(s+r), s \in \mathbb{R}\}, r \in \mathbb{R} .$$

An additive norm on  $\tilde{\mathfrak{g}}$  is square if the corresponding lattice function is square. Recall [BT1] that  $\text{Norm}^1(V)$  and  $\text{Norm}^2 \tilde{\mathfrak{g}}$  (and therefore  $\text{Latt}^1(V)$  and  $\text{Latt}^2 \tilde{\mathfrak{g}}$  by transfer of structure) are endowed with affine structures : the barycenter of two points with positive weights is defined.

The involution  $\sigma$  acts on  $\text{Norm}^2 \tilde{\mathfrak{g}}$  via

$$\alpha^{\sigma}(a) = \alpha(a^{\sigma}), a \in \tilde{\mathfrak{g}}, \alpha \in \text{Norm}^2 \tilde{\mathfrak{g}} .$$

By transfer of structure,  $\sigma$  acts on  $\text{Latt}^2 \tilde{\mathfrak{g}}$  via

$$\Lambda^{\sigma}(r) = [\Lambda(r)]^{\sigma}, \Lambda \in \text{Latt}^2 \tilde{\mathfrak{g}}, r \in \mathbb{R} .$$

A square norm  $\alpha$  (resp. a square lattice function  $\Lambda$ ) is said to be self-dual if  $\alpha = \alpha^{\sigma}$  (resp.  $\Lambda = \Lambda^{\sigma}$ ). We denote by  $\text{Norm}_{\sigma}^2 \tilde{\mathfrak{g}}$  and  $\text{Latt}_{\sigma}^2 \tilde{\mathfrak{g}}$  the corresponding sets.

Now, in terms of lattice functions, Corollary 2 of [BT2], page 163, writes:

**Lemma 3.5** *The map  $\Lambda \mapsto \text{End}(\Lambda)$  induces a bijection from the set of self-dual lattice functions in  $V$  to the set of self-dual square lattice functions in  $\tilde{\mathfrak{g}}$ .*

In other words, for any  $\Lambda \in \text{Latt}_{\sigma}^2 \tilde{\mathfrak{g}}$ , there exists a unique  $\Lambda^2 = \Lambda_h^2 \in \text{Latt}_h^1(V)$  such that  $\text{End}(\Lambda) = \Lambda^2$ .

Note that the sets  $\text{Latt}_h^1(V)$ ,  $\text{Norm}_h^1(V)$ ,  $\text{Latt}_{\sigma}^2 \tilde{\mathfrak{g}}$  and  $\text{Norm}_{\sigma}^2 \tilde{\mathfrak{g}}$  are  $G$ -sets and that the various identifications among them are  $G$ -equivariant.

Let  $u \in F^{\times}$  and assume that  $uh$  is still an  $\varepsilon$ -hermitian form with respect to  $\sigma_F$ . Then the involution  $\sigma$  of  $\tilde{\mathfrak{g}}$  corresponding to  $uh$  remains the same and defines the same

unitary group  $G \subset \tilde{G}$ . For  $\Lambda \in \text{Latt}^1(V)$  and  $s \in \mathbb{R}$ , we denote by  $\Lambda + s$  the lattice function given by  $(\Lambda + s)(r) = \Lambda(s + r)$ ,  $r \in \mathbb{R}$ .

**Lemma 3.6.** *Let  $\Lambda^2 \in \text{Latt}_{\sigma}^2 \tilde{\mathfrak{g}}$  and  $\Lambda_h^2$  (resp.  $\Lambda_{uh}^2$ ) be the unique element of  $\text{Latt}_h^1(V)$  (resp. of  $\text{Latt}_{uh}^1(V)$ ) satisfying  $\text{End}(\Lambda_h^2) = \Lambda^2$  (resp.  $\text{End}(\Lambda_{uh}^2) = \Lambda^2$ ). Then  $\Lambda_{uh}^2 = \Lambda_h^2 - v(u)/2$ , that is  $\Lambda_{uh}^2(r) = \Lambda_h^2(r - v(u)/2)$ ,  $r \in \mathbb{R}$ .*

*Proof.* We easily check that for  $\Lambda \in \text{Latt}^1(V)$  and  $s \in \mathbb{R}$ , we have

$$\Lambda^{\sharp_{uh}} = u^{-\sigma} \Lambda^{\sharp_h} \text{ and } (\Lambda + s)^{\sharp_h} = \Lambda - s .$$

We certainly have  $\text{End}(\Lambda_h^2 - v(u)/2) = \text{End}(\Lambda_h^2) = \Lambda^2$ . So by a unicity argument, we must prove that  $\Lambda_h^2 - v(u)/2 \in \text{Latt}_{uh}^1(V)$ . But

$$\begin{aligned} (\Lambda_h^2 - v(u)/2)^{\sharp_{uh}} &= u^{-\sigma} (\Lambda_h^2 - v(u)/2)^{\sharp_h} \\ &= u^{-\sigma} (\Lambda_h^2 + v(u)/2) = \Lambda_h^2 + v(u)/2 - v(u^\sigma) = \Lambda_h^2 - v(u)/2 , \end{aligned}$$

as required.

#### 4. The building as a set of self-dual lattice-functions

Let  $I$  denote the building of the standard valuated root datum of  $G$  introduced in [BT2] and  $A$  denote the apartment of  $I$  attached to  $\mathcal{S}$ . Write  $V^* = X^*(\mathcal{S} \otimes \mathbb{R})$ ; this is an  $\mathbb{R}$ -vector space with basis  $(a_i)_{i=1, \dots, r}$ . Let  $V$  denote the linear dual of  $V^*$ . We identify  $A$  with  $V$ .

To a point  $p \in A \simeq V$ , we attach the norm  $\alpha_p$  on  $V$  defined by

$$\alpha_p \left( \sum_{i \in I} \lambda_i e_i + x_o \right) = \inf[\omega(x_o), \inf_{i \in I} (v(\lambda_i) - a_i(p))], \quad x_o \in V_o, \quad \lambda_i \in F \text{ for } i \in I .$$

Here  $\omega(x_o) = \frac{1}{2}v(h(x_o, x_o))$ ,  $x_o \in V_o$ .

Here are two important facts from [BT2].

**Proposition 4.1.** ([BT2](Prop. 2.9, 2.11(i))) *The map  $p \mapsto \alpha_p$  is a bijection from  $A$  to the set of MM-norms on  $V$  which split in the decomposition  $V = \bigoplus_{i \in I} F e_i \oplus V_o$ . It is  $N$ -equivariant.*

For the notion of splitting for norms, see [BT1](1.4).

**Proposition 4.2.** ([BT2](2.12)) *i) The map  $p \mapsto \alpha_p$  extends in a unique way to a  $G$ -equivariant and affine bijection  $j_h : I \rightarrow \text{Norm}_h^1(V)$  (in particular  $\text{Norm}_h^1(V)$  is a convex subset of  $\text{Norm}^1(V)$ ).*

*ii) The map  $j_h$  is the unique affine and  $G$ -equivariant map  $I \rightarrow \text{Norm}_h^1(V)$ .*

From §3, we get a unique affine and  $G$ -equivariant map  $I \rightarrow \text{Latt}_h^1(V)$  that we still denote by  $j_h$ .

For  $r \in \mathbb{R}$ , let  $\mathcal{V}_o^r$  be the lattice of  $V_o$  given by  $\{x_o \in V_o ; \omega(x_o) \geq r\}$ . For  $x \in \mathbb{R}$ , let  $[x]$  denote the least integer greater than or equal to  $x$ . Then the map  $j_h : I \rightarrow \text{Latt}_h^1(V)$  is given on  $A$  by  $j_h(p) = \Lambda_p$ , where

$$\Lambda_p(r) = \mathcal{V}_o^r \oplus \bigoplus_{i \in I} \mathfrak{p}_F^{\lceil r + a_i(p) \rceil} e_i , \quad r \in \mathbb{R} .$$

Let  $u$  be an element of  $F^\times$  such that  $uh$  remains  $\varepsilon$ -hermitian with respect to  $\sigma_F$ . It follows from the proof of Lemma (3.6) that if  $\Lambda \in \text{Latt}^1(V)$ , we have  $\Lambda \in \text{Latt}_h^1(V)$  if, and only if,  $\Lambda - v(u)/2 \in \text{Latt}_{uh}^1(V)$ . Since  $\text{End}(\Lambda + s) = \text{End}(\Lambda)$ , for  $\Lambda \in \text{Latt}^1(V)$  and  $s \in \mathbb{R}$ , the bijective map  $j_\sigma : I \rightarrow \text{Latt}_\sigma^2(V)$ , given by  $j_\sigma = \text{End} \circ j_h$ , does not depend on the choice of the form  $h$ , the involution  $\sigma$  being fixed. By construction it is affine and  $G$ -equivariant. It is uniquely determined by these two properties. Indeed if  $j'_\sigma : I \rightarrow \text{Latt}_\sigma^2(V)$  is affine and  $G$ -equivariant, so is  $(j'_\sigma)^{-1} \circ j_\sigma : I \rightarrow I$ . But such a map must be the identity map.

We also recall here the description of the enlarged building  $I^1$  of  $\tilde{G} = \text{GL}_F(V)$  in terms of lattice functions.

**Proposition 4.3.** ([BT1](2.11)) *i) There is a  $\tilde{G}$ -equivariant and affine bijection  $j : I^1 \rightarrow \text{Norm}^1(V)$ .*

*ii) If we have another affine and  $\tilde{G}$ -equivariant map  $j' : I^1 \rightarrow \text{Norm}^1(V)$  then there exists  $r \in \mathbb{R}$  such that, for all  $\alpha \in \text{Norm}^1(V)$ ,  $j'(\alpha) = j(\alpha) + r$ .*

From [BL] Proposition 2.4, for each  $j$  as in Proposition 4.3, we get an affine and  $\tilde{G}$ -equivariant map  $I^1 \rightarrow \text{Latt}^1(V)$  that we also denote by  $j$ .

## 5. Centralizers of Lie algebra elements

We denote by  $\mathfrak{g}$  the Lie algebra of  $G$ :

$$\mathfrak{g} = \{a \in \tilde{\mathfrak{g}} ; a + a^\sigma = 0\} .$$

We consider an element  $\beta$  of  $\mathfrak{g}$  satisfying

(H1) The  $F$ -algebra  $E := F[\beta] \subset \tilde{\mathfrak{g}}$  is a direct sum of fields.

We write  $\tilde{\mathfrak{h}}$  (resp.  $\mathfrak{h}$ ) for the centralizer of  $\beta$  in  $\tilde{\mathfrak{g}}$  (resp. in  $\mathfrak{g}$ ) and  $\tilde{H}$  (resp.  $H$ ) for the fixator of  $\beta$  in  $\tilde{G}$  (resp. in  $G$ ) for the adjoint action.

Since  $\sigma(\beta) = -\beta$ , we have easily that  $E \subset \tilde{\mathfrak{g}}$  is  $\sigma$ -stable. We write

$$E = \bigoplus_{i=1, \dots, t} (E_i \oplus E_{-i}) \oplus \bigoplus_{k=1, \dots, s} E_{(0,k)},$$

where, for each  $i$  in  $J = \{\pm 1, \dots, \pm t\}$  or  $J_o = \{(0, k) : k = 1, \dots, s\}$ ,  $E_i$  is a field extension of  $F$ , and we have labeled the components such that, for each  $i \in J_o \cup J$ ,

(H2)  $\sigma(E_i) = E_{-i}$ ,

with the understanding that  $i = -i$ , for  $i \in J_o$ . We remark that the torus  $E \cap G$  in  $G$  is anisotropic (modulo the centre) if and only if  $J = \emptyset$  and that every maximal anisotropic torus in  $G$  takes this form (see [Mor] Proposition 1.3).

For each  $i \in J_o$ , we set  $E_i^o = \{a \in E_i ; a = a^\sigma\}$ , so that  $E_i/E_i^o$  is a Galois extension of degree  $\leq 2$  and a generator of  $\text{Gal}(E_i/E_i^o)$  is  $\sigma_{E_i} := \sigma|_{E_i}$ . For  $i \in J_o \cup J$ , let  $\mathbf{1}_i$  be the idempotent of  $E$  attached to  $E_i$ ; from (H2), we have  $\sigma(\mathbf{1}_i) = \mathbf{1}_{-i}$ . We have the decomposition

$$V = \bigoplus_{i \in J_o \cup J} V_i, V_i = \mathbf{1}_i V .$$

Note that, if  $i \neq -k$ ,  $v \in V_i$  and  $w \in V_k$ , we have  $h(v, w) = h(\mathbf{1}_i v, w) = h(v, \mathbf{1}_i w) = 0$  so, for  $i \in J_o \cup J$ ,

$$V_i^\perp = \bigoplus_{k \neq -i} V_k.$$

For  $i \in J_o \cup J$ ,  $V_i$  is naturally an  $E_i$ -vector space and we have obvious isomorphisms of algebras and groups respectively:

$$\tilde{\mathfrak{h}} \simeq \prod_{i \in J_o \cup J} \text{End}_{E_i} V_i,$$

$$\tilde{H} \simeq \prod_{i \in J_o \cup J} \text{Aut}_{E_i} V_i.$$

The involution  $\sigma$  stabilizes  $\tilde{\mathfrak{h}} \subset \tilde{\mathfrak{g}}$  and, for each  $i$ ,  $\sigma(\text{End}_{E_i} V_i) = \text{End}_{E_{-i}} V_{-i}$ . For  $i \in J_o$ , we write  $\sigma_i = \sigma|_{\text{End}_{E_i} V_i}$ . Let us fix  $i \in J_o$ . The map  $\sigma_i$  is an involution of the central simple  $E_i$ -algebra  $\text{End}_{E_i} V_i$ . By a classical theorem ([Inv] Theorem 4.2), there exists  $\varepsilon_i \in \{\pm 1\}$  and a non-degenerate  $\varepsilon_i$ -hermitian form  $h_i$  on  $V_i$  relative to  $\sigma_{E_i}$  such that  $\sigma_i$  is the involution attached to  $h_i$ . Of course  $h_i$  is only defined up to a scalar in  $E_i^\times$ . Let

$$H_i = \{g \in \text{Aut}_{E_i} V_i ; gg^{\sigma_i} = 1\}$$

be the unitary group attached to  $h_i$ . On the other hand, for  $i \in J$ , we put

$$H_i = \text{Aut}_{E_i} V_i,$$

so that  $\sigma(H_i) = H_{-i}$  and  $H_i$  is isomorphic to  $\{g \in H_i \times H_{-i} : gg^\sigma = 1\}$  by  $h \mapsto (h, h^{-\sigma})$ . Then, putting  $J_+ = \{1, \dots, t\}$ , we have a natural group isomorphism

$$H \simeq \prod_{i \in J_o \cup J_+} H_i.$$

We may actually require a compatibility relation between the forms  $h_i$ ,  $i \in J_o$  and the form  $h$ . Let us fix  $i \in J_o$ . Let  $\lambda_i : E_i \rightarrow F$  be any  $\sigma$ -equivariant non-zero  $F$ -linear form. Such forms exist. Indeed choose a non-zero linear form  $\lambda_i^o : E_i^o \rightarrow F_o$ . If  $F = F_o$  then we put  $\lambda = \lambda_i^o \circ \text{Tr}_{E/E_i^o}$ . Otherwise  $E_i = FE_i^o$  and we can extend  $\lambda_i^o$  by linearity to get the required map  $\lambda_i$ . In all cases we have:

$$(5.1) \quad \lambda_i^o \circ \text{Tr}_{E_i/E_i^o} = \text{Tr}_{F/F_o} \circ \lambda.$$

We still write  $h$  for the restriction of  $h$  to  $V_i$ .

**Lemma 5.2.** *Let  $i \in J_o$ . There exists a unique  $\varepsilon$ -hermitian form  $h_i : V_i \times V_i \rightarrow E_i$  relative to  $\sigma_{E_i}$  such that*

$$(5.3) \quad h(v, w) = \lambda_i(h_i(v, w)), \quad \text{for all } v, w \in V_i.$$

*It is non-degenerate.*

*Proof.* Since we have the orthogonal decomposition

$$V = V_i \perp \bigoplus_{k \neq i} V_k,$$



the restriction  $h|_{V_i}$  is non-degenerate.

The  $F$ -linear map  $\text{Hom}_{E_i}(V_i, E_i) \rightarrow \text{Hom}_F(V_i, F)$ ,  $\varphi \mapsto \lambda_i \circ \varphi$  is an isomorphism of  $F$ -vector space. Indeed if  $\varphi$  lies in the kernel, we have  $\text{Im}(\varphi) \subset \text{Ker}(\lambda_i)$ , a strict subspace of  $E_i$ , and  $\varphi$  must be trivial. Moreover the two dual spaces have the same  $F$ -dimension. For  $v \in V_i$  let  $h_v$  be the element of  $\text{Hom}_F(V_i, F)$  given by  $h_v(w) = h(v, w)$ . There exists a unique  $\varphi_w \in \text{Hom}_{E_i}(V_i, E_i)$  such that  $h_v = \lambda_i \circ \varphi_w$ . It is now routine to check that  $h_i(v, w) := \varphi_v(w)$ ,  $v, w \in V_i$ , has the required properties.

We easily check that if  $h_i$  is as in the lemma, then the corresponding involution on  $\text{End}_{E_i} V_i$  is  $\sigma_i$ . In the following we assume that the forms  $h_i$ ,  $i \in J_o$ , satisfy (5.3).

For technical reasons, we need one more assumption on the  $\lambda_i$ ,  $i \in J_o$ . We fix  $i$  again. Let

$$\mathcal{J} = \{e \in E_i^o ; \lambda_i^o(e\mathfrak{o}_{E_i^o}) \subset \mathfrak{p}_{F_o}\} .$$

This is an  $\mathfrak{o}_{E_i^o}$ -lattice in  $E_i^o$  and must have the form  $t\mathfrak{p}_{E_i^o}$ , for some  $t \in (E_i^o)^\times$ . So replacing  $\lambda_i$  by  $e \mapsto \lambda_i(tx)$ , we may assume that  $\mathcal{J} = \mathfrak{p}_{E_i^o}$ . In the following we assume that the linear forms  $\lambda_i$ ,  $i \in J_o$ , have this property.

**Lemma 5.4.** *Fix  $i \in J_o$ . Let  $\lambda_i^1, \lambda_i^2 : E_i \rightarrow F$  be two linear forms as above and let  $h_i^1, h_i^2$  be the corresponding  $\varepsilon$ -hermitian forms on  $V_i$  (i.e.  $h_i^1$  and  $h_i^2$  satisfy (5.3)). Then there exists  $u \in \mathfrak{o}_{E_i^o}^\times$  such that  $h_i^2 = uh_i^1$ .*

*Proof.* Since  $h_i^1$  and  $h_i^2$  induce the same involution on  $\text{End}_{E_i} V_i$ , there exists  $u \in E_i^\times$  such that  $h_i^2 = uh_i^1$ . The fact that  $h_i^1$  and  $h_i^2$  are both  $\varepsilon$ -hermitian with respect to  $\sigma_{E_i}$  implies that  $u$  lies in  $E_i^o$ . Condition (5.3) writes

$$h(v, w) = \lambda_i^1(h_i^1(v, w)) = \lambda_i^2(uh_i^1(v, w)) , \quad v, w \in V_i .$$

So  $\lambda_i^1(e) = \lambda_i^2(ue)$ ,  $e \in E_i$ . By applying  $\text{Tr}_{F/F_o}$  to this equality, we get  $\lambda_i^{o,1}(e) = \lambda_i^{o,2}(ue)$ ,  $e \in E_i^o$ . Hence

$$\begin{aligned} \mathfrak{p}_{E_i^o} &= \{e \in E_i^o ; \lambda_i^{o,1}(e\mathfrak{o}_{E_i^o}) \subset \mathfrak{p}_{F_o}\} \\ &= \{e \in E_i^o ; \lambda_i^{o,2}(ue\mathfrak{o}_{E_i^o}) \subset \mathfrak{p}_{F_o}\} = u^{-1}\mathfrak{p}_{E_i^o} . \end{aligned}$$

So  $u \in \mathfrak{o}_{E_i^o}^\times$  as required.

Let us fix  $i$ . Let  $L$  be an  $\mathfrak{o}_{E_i^o}$ -lattice in  $V_i$ . Then  $L$  has a dual  $L^\sharp$  relative to the form  $h|_{V_i}$  and a dual  $L^{\sharp i}$  relative to the form  $h_i$ .

**Lemma 5.5.** *The lattices  $L^\sharp$  and  $L^{\sharp i}$  coincide.*

*Proof.* We have

$$\begin{aligned} L^\sharp &= \{v \in V_i ; h(v, L) \subset \mathfrak{p}_F\} \\ &= \{v \in V_i ; \text{Tr}_{F/F_o} h(v, L) \subset \mathfrak{p}_{F_o}\} \\ &= \{v \in V_i ; \lambda_o \circ \text{Tr}_{E_i/E_i^o} h_i(v, L) \subset \mathfrak{p}_{F_o}\} \\ &= \{v \in V_i ; \text{Tr}_{E_i/E_i^o} h_i(v, L) \subset \mathfrak{p}_{E_i^o}\} \\ &= \{v \in V_i ; f(v, L) \subset \mathfrak{p}_{E_i}\} \\ &= L^{\sharp i}, \end{aligned}$$

where the second and fifth equalities hold because  $F/F_o$  and  $E_i/E_i^o$  are at worst tamely ramified.

## 6. Embedding the building of the centralizer

We keep the notation as in the previous section. Assume for a moment that the extensions  $E_i/F$ ,  $i \in J_o \cup J$ , are separable. Then the group  $H$  is naturally the group of rational points of a reductive  $F$ -group  $\mathbf{H}$ . Indeed each  $H_i$ ,  $i \in J_o \cup J$ , is naturally the group of rational points of a classical  $E_i$ -group  $\mathbf{H}_i$  (we do not need  $E_i/F$ -separable here) and

$$\mathbf{H} \simeq \prod_{i \in J_o \cup J} \text{Res}_{E_i/F} \mathbf{H}_i .$$

The (enlarged) affine building of  $\mathbf{H}$ ,  $I_\beta^1 := I^1(\mathbf{H}, F)$ , is the cartesian product of the (enlarged) affine buildings  $I^1(\text{Res}_{E_i/F} \mathbf{H}_i, F)$ ,  $i \in J_o \cup J$ . For all  $i$ , the (enlarged) buildings  $I^1(\text{Res}_{E_i/F} \mathbf{H}_i, F)$  and  $I^1(\mathbf{H}_i, E_i)$  identify canonically. Note also that, for  $i \in J_o$ , the centre of  $\mathbf{H}_i$  is compact so the enlarged building is also the non-enlarged building; in particular, if  $J = \emptyset$  then all the buildings involved are non-enlarged.

Since we do not want any restriction on the extensions  $E_i/F$ , we shall take as a definition of the (enlarged) building  $I_\beta^1$  attached to the group  $H$ :

$$(6.1) \quad I_\beta^1 := \prod_{i \in J_o \cup J} I^1(\mathbf{H}_i, E_i)$$

We abbreviate  $I_i^1 = I^1(\mathbf{H}_i, E_i)$ ,  $i \in J_o \cup J$ .

We are going to construct a map  $j_\beta : I_\beta^1 \rightarrow I$ . We normalize the lattice-functions in  $\text{Latt}_{\mathfrak{o}_{E_i}}^1(V_i)$  by  $\Lambda_i(r + v_i(\pi_i)) = \mathfrak{p}_{E_i} \Lambda_i(r)$ ,  $r \in \mathbb{R}$ , where, for each  $i$ ,  $\pi_i$  denotes a uniformizer of  $E_i$  and  $v_i$  the unique extension of  $v$  to a valuation of  $E_i$ . It is straightforward that we have a well defined map

$$\begin{aligned} \tilde{j}_\beta : \prod_{i \in J_o \cup J} \text{Latt}_{\mathfrak{o}_{E_i}}^1(V_i) &\longrightarrow \text{Latt}^1(V) \\ (\Lambda_i)_{i \in J_o \cup J} &\mapsto \bigoplus_{i \in J_o \cup J} \Lambda_i \end{aligned}$$

where  $(\bigoplus_{i \in J_o \cup J} \Lambda_i)(r) = \bigoplus_{i \in J_o \cup J} \Lambda_i(r)$ , for  $r \in \mathbb{R}$ . This map is clearly injective and equivariant for the action of group  $\prod_{i \in J_o \cup J} \text{Aut}_{E_i} V_i \subset \text{Aut}_F V$ .

For  $i \in J_o$ , we denote by  $\sharp_i$  the involution on  $\text{Latt}_{\mathfrak{o}_{E_i}}^1(V_i)$  attached to  $h_i$ , and by  $\text{Latt}_{\mathfrak{o}_{E_i}, h_i}^1(V_i) \subset \text{Latt}_{\mathfrak{o}_{E_i}}^1(V_i)$  the set of fixed points. For  $i \in J$ , we denote by  $\sharp_i$  the map  $\text{Latt}_{\mathfrak{o}_{E_i}}^1(V_i) \rightarrow \text{Latt}_{\mathfrak{o}_{E_{-i}}}^1(V_{-i})$  given by

$$\Lambda_i^{\sharp_i}(r) = \{v \in V_{-i} ; h(v, \Lambda_i(-r+)) \subset \mathfrak{p}_F\} .$$

for  $\Lambda_i \in \text{Latt}_{\mathfrak{o}_{E_i}}^1(V_i)$ .

We define an involution  $b$  on  $\prod_{i \in J_o \cup J} \text{Latt}_{\mathfrak{o}_{E_i}}^1(V_i)$  by

$$(\Lambda_i)_{i \in J_o \cup J}^b = \left( \Lambda_{-i}^{\sharp_{-i}} \right)_{i \in J_o \cup J} ,$$

Then we have a bijection

$$\iota_h : \prod_{i \in J_o} \text{Latt}_{\mathfrak{o}_{E_i}, h_i}^1(V_i) \times \prod_{i \in J_+} \text{Latt}_{\mathfrak{o}_{E_i}}^1(V_i) \rightarrow \left( \prod_{i \in J_o \cup J} \text{Latt}_{\mathfrak{o}_{E_i}}^1(V_i) \right)^b,$$

given by  $(\Lambda_i)_{i \in J_o \cup J_+} \mapsto (\Lambda_i)_{i \in J_o \cup J}$ , with  $\Lambda_{-i} = \Lambda_i^{\sharp}$ , for  $i \in J_+$ .

**Lemma 6.2.** *For  $x \in \prod_{i \in J_o \cup J} \text{Latt}_{\mathfrak{o}_{E_i}}^1(V_i)$ , we have  $\tilde{j}_\beta(x^b) = \tilde{j}_\beta(x)^{\sharp h}$ . In particular  $\tilde{j}_\beta \circ \iota_h$  maps  $\prod_{i \in J_o} \text{Latt}_{\mathfrak{o}_{E_i}, h_i}^1(V_i) \times \prod_{i \in J_+} \text{Latt}_{\mathfrak{o}_{E_i}}^1(V_i)$  into  $\text{Latt}_h^1(V)$ .*

*Proof.* Fix  $(\Lambda_i)_{i \in J_o \cup J} \in \prod_{i \in J_o \cup J} \text{Latt}_{\mathfrak{o}_{E_i}}^1 V_i$  and set  $\Lambda = \tilde{j}_\beta \left( (\Lambda_i)_{i \in J_o \cup J_+} \right)$ . We have

$$\Lambda^{\sharp h}(r) = \Lambda(-r+)^{\sharp h} = \{v \in V ; h(v, \Lambda(-r+)) \subset \mathfrak{p}_F\}, \quad r \in \mathbb{R}.$$

Fix  $r \in \mathbb{R}$ . We have

$$\Lambda(-r+) = \bigoplus_{i \in J_o \cup J} \Lambda_i(-r+).$$

Let  $v = \sum_{i \in J_o \cup J} v_i$ , with  $v_i \in V_i$ , be an element of  $V$ . Since  $V_i^\perp = \bigoplus_{k \neq -i} V_k$ , we have  $v \in \Lambda^{\sharp h}(r)$  if and only if  $h(v_{-i}, \Lambda_i(-r+)) \subset \mathfrak{p}_F$ , for all  $i$ , that is if  $v_{-i} \in \Lambda_i^{\sharp}(r)$ , for all  $i$  (by lemma (5.5) for  $i \in J_o$  or by definition for  $i \in J$ ); the lemma follows.

With the notation of §4, for each set  $\{j_i\}_{i \in J_+}$  of maps  $j_i : I_i^1 \rightarrow \text{Latt}_{\mathfrak{o}_{E_i}}^1(V_i)$  given by Proposition 4.3, we define a map  $j_\beta : \prod_{i \in J_o \cup J_+} I_i^1 \rightarrow I$  by

$$j_\beta = j_h^{-1} \circ \tilde{j}_\beta \circ \iota_h \circ \left( \prod_{i \in J_o} j_{h_i} \times \prod_{i \in J_+} j_i \right).$$

These maps depend *a priori* on the forms  $h$ , and  $h_j$ ,  $j \in J_o$ .

**Theorem 6.3.** *Each map  $j_\beta$  is injective,  $H$ -equivariant. The set of such maps (as  $\{j_i\}_{i \in J_+}$  varies) depends only on the involution  $\sigma$ .*

In particular, if  $J = \emptyset$  then there is a unique map  $j_\beta$ , depending only on the involution  $\sigma$ .

*Proofs.* The first two properties are straightforward. Assume that  $h' = uh$ ,  $u \in F^\times$ , is another  $\varepsilon$ -hermitian form on  $V$ , with respect to  $\sigma_F$ , defining the same involution  $\sigma$  on  $\tilde{\mathfrak{g}}$ . Then  $u \in F_o$ . For  $i \in J_o$ , let  $h'_i$  be an  $\varepsilon$ -hermitian form on  $V_i$  satisfying

$$uh(v, w) = \lambda'_i(h'_i(v, w)) \quad v, w \in V_i,$$

where the  $\lambda'_i : E_i \rightarrow F$  are linear forms as above. Then by lemma (5.4), for all  $i \in J_o$ , there exists  $u'_i \in \mathfrak{o}_{E_i}^\times$  such that  $u^{-1}h'_i = u'_i h_i$ , that is  $h'_i = uu'_i h_i$ .

Let  $\{j_i\}_{i \in J_+}$  be as above; we show that, for a suitable choice of  $\{j'_i\}_{i \in J_+}$ , we have

$$j_h^{-1} \circ \tilde{j}_\beta \circ \iota_h \circ j_1 = j_{h'} \circ \tilde{j}_\beta \circ \iota_{h'} \circ j'_1,$$

and the result follows.

By Lemma (3.6), for  $i \in J_+$ , for all  $x_i \in I_i^1$ , we have  $j_{h'_i}(x_i) = j_{h_i}(x_i) - v(uu'_i)/2 = j_{h_i}(x_i) - v(u)/2$ . For  $i \in J_+$ , we choose  $j'_i$  such that  $j'_i(x) = j_i(x) - v(u)/2$  for  $x \in I_i^1$ , that is  $j'_i \circ j_i^{-1}(\Lambda_i) = \Lambda_i - v(u)/2$  for  $\Lambda_i \in \text{Latt}_{\mathfrak{o}_{E_i}}^1(V_i)$ . We abbreviate

$$j = \prod_{i \in J_o} j_{h_i} \times \prod_{i \in J_+} j_i, \quad j' = \prod_{i \in J_o} j_{h'_i} \times \prod_{i \in J_+} j'_i;$$

then, for  $(\Lambda_i)_{i \in J_o \cup J_+} \in \prod_{i \in J_o} \text{Latt}_{\mathfrak{o}_{E_i}, h_i}^1(V_i) \times \prod_{i \in J_+} \text{Latt}_{\mathfrak{o}_{E_i}}^1(V_i)$ , we have

$$j' \circ j^{-1} \left( (\Lambda_i)_{i \in J_o \cup J_+} \right) = (\Lambda_i - v(u)/2)_{i \in J_o \cup J_+}.$$

It is also straightforward to check that

$$\iota_{h'} \left( (\Lambda_i - v(u)/2)_{i \in J_o \cup J_+} \right) = \iota_h \left( (\Lambda_i)_{i \in J_o \cup J_+} \right) - v(u)/2,$$

for  $(\Lambda_i)_{i \in J_o \cup J_+} \in \prod_{i \in J_o} \text{Latt}_{\mathfrak{o}_{E_i}, h_i}^1(V_i) \times \prod_{i \in J_+} \text{Latt}_{\mathfrak{o}_{E_i}}^1(V_i)$ . Then we have

$$\begin{aligned} \tilde{j}_\beta \circ \iota_{h'} \circ j' \circ j^{-1} \left( (\Lambda_i)_{i \in J_o \cup J_+} \right) &= \tilde{j}_\beta \circ \iota_{h'} \left( (\Lambda_i - v(u)/2)_{i \in J_o \cup J_+} \right) \\ &= \tilde{j}_\beta \left( \iota_h \left( (\Lambda_i)_{i \in J_o \cup J_+} \right) - v(u)/2 \right) \\ &= \tilde{j}_\beta \circ \iota_h \left( (\Lambda_i)_{i \in J_o \cup J_+} \right) - v(u)/2. \end{aligned}$$

By Lemma (3.6) again, we have  $j_{h'}(x) = j_h(x) - v(u)/2$ ,  $x \in I$ , that is  $\Lambda - v(u)/2 = j_{h'} \circ j_h^{-1}(\Lambda)$ ,  $\Lambda \in \text{Latt}_h^1(V)$ . So

$$j_{h'} \circ j_h^{-1} \circ \tilde{j}_\beta \circ \iota_h = \tilde{j}_\beta \circ \iota_{h'} \circ j' \circ j^{-1},$$

and the lemma follows.

## 7. Affine structures

We keep the notation as in the previous sections. For  $x = (x_i)_{i \in J_o \cup J_+}$ ,  $y = (y_i)_{i \in J_o \cup J_+}$  in  $I_\beta^1 = \prod_{i \in J_o \cup J_+} I_i^1$  and  $t \in [0, 1]$ , we define the barycenter  $tx + (1-t)y$  to be

$$(tx_i + (1-t)y_i)_{i \in J_o \cup J_+}.$$

We define the barycenter of two points in  $\prod_{i \in J_o \cup J_+} \text{Latt}_{\mathfrak{o}_{E_i}}^1(V_i)$  in a similar way. Since, for  $i \in J_o$ ,  $\text{Latt}_{\mathfrak{o}_{E_i}, h_i}^1(V_i)$  is convex in  $\text{Latt}_{\mathfrak{o}_{E_i}}^1(V_i)$ , the subset  $\prod_{i \in J_o} \text{Latt}_{\mathfrak{o}_{E_i}, h_i}^1(V_i) \times \prod_{i \in J_+} \text{Latt}_{\mathfrak{o}_{E_i}}^1(V_i)$  of  $\prod_{i \in J_o \cup J_+} \text{Latt}_{\mathfrak{o}_{E_i}}^1(V_i)$  is convex also.

**Proposition 7.1.** *Let  $\beta$  be as in §5. Then each map  $j_\beta$  is affine: for all  $x, y \in I_\beta^1$ ,  $t \in [0, 1]$ , we have*

$$j_\beta(tx + (1-t)y) = tj_\beta(x) + (1-t)j_\beta(y) .$$

*Proof.* By construction it suffices to prove that the maps  $\tilde{j}_\beta$  and  $\iota_h$  are affine. We begin with  $\tilde{j}_\beta$ . Let  $(\Lambda_i)_{i \in J_o \cup J}$ ,  $(M_i)_{i \in J_o \cup J}$  be elements of  $\prod_{i \in J_o \cup J} \text{Latt}_{\mathfrak{o}_{E_i}}^1(V_i)$ . We must prove that

$$\bigoplus_{i \in J_o \cup J} (t\Lambda_i + (1-t)M_i) = t \left( \bigoplus_{i \in J_o \cup J} \Lambda_i \right) + (1-t) \left( \bigoplus_{i \in J_o \cup J} M_i \right) .$$

Let us recall the construction of the barycenter of two lattice functions (we do it for  $\text{Latt}^1(V)$ ). Let  $\Lambda, M \in \text{Latt}^1(V)$ . There exists an  $F$ -basis  $(e_1, \dots, e_n)$  of  $V$  which splits both  $\Lambda$  and  $M$  : there exist constants  $\lambda_1, \dots, \lambda_n, \mu_1, \dots, \mu_n$  in  $\mathbb{R}$  such that

$$\Lambda(r) = \bigoplus_{k=1, \dots, n} \mathfrak{p}_F^{\lceil r + \lambda_k \rceil} e_k , \quad M(r) = \bigoplus_{k=1, \dots, n} \mathfrak{p}_F^{\lceil r + \mu_k \rceil} e_k , \quad r \in \mathbb{R} .$$

Then for  $t \in [0, 1]$ ,  $t\Lambda + (1-t)M$  is given by

$$(t\Lambda + (1-t)M)(r) = \bigoplus_{k=1, \dots, n} \mathfrak{p}_F^{\lceil r + t\lambda_k + (1-t)\mu_k \rceil} e_k , \quad r \in \mathbb{R} .$$

The proof that  $\tilde{j}_\beta$  is affine is then to construct a common splitting basis for  $\bigoplus_{i \in J_o \cup J} \Lambda_i$  and  $\bigoplus_{i \in J_o \cup J} M_i$  from bases  $\mathcal{B}_i$  of  $V_i$ ,  $i \in J_o \cup J$ , where  $\mathcal{B}_i$  splits  $\Lambda_i$  and  $M_i$ . We leave this easy exercise to the reader.

Now we turn to  $\iota_h$ . Suppose  $i \in J_+$  and  $\Lambda_i \in \text{Latt}_{\mathfrak{o}_{E_i}}^1(V_i)$ , and let  $(e_1, \dots, e_n)$  be an  $E_i$ -basis of  $V_i$  which splits  $\Lambda_i$ . Let  $(e_{-1}, \dots, e_{-n})$  be the dual  $E_{-i}$ -basis of  $V_{-i}$ , such that  $h(e_{-k}, e_l) = \delta_{kl}$ , for  $1 \leq k, l \leq n$ . It is straightforward to check that this basis splits  $\Lambda_i^{\sharp_i}$  and that,

$$(7.2) \quad \text{if } \Lambda_i(r) = \bigoplus_{k=1, \dots, n} \mathfrak{p}_{E_i}^{\lceil r + \lambda_k \rceil} e_k \quad \text{then} \quad \Lambda_i^{\sharp_i}(r) = \bigoplus_{k=1, \dots, n} \mathfrak{p}_{E_{-i}}^{\lceil r - \lambda_k \rceil} e_{-k} .$$

To show that  $\iota_h$  is affine, we just need to check that, for  $i \in J_+$ ,  $\Lambda_i, M_i \in \text{Latt}_{\mathfrak{o}_{E_i}}^1(V_i)$  and  $t \in [0, 1]$ , we have

$$(t\Lambda_i + (1-t)M_i)^{\sharp_i} = t\Lambda_i^{\sharp_i} + (1-t)M_i^{\sharp_i} .$$

The details of the proof – which is to choose an  $E_i$ -basis of  $V_i$  which splits both  $\Lambda_i$  and  $M_i$ , take its dual basis and then use (7.2) – are again left to the reader.

## 8. The image of an apartment

We keep the notation of the previous sections. We will show that the image of an apartment of  $I_\beta^1$  under each map  $j_\beta$  is contained in an apartment of  $I$ .

Given a Witt decomposition  $V = V_+ \oplus V_o \oplus V_-$ , with basis  $(e_l)_{l=1,\dots,r}$  of  $V_+$  and the dual basis  $(e_{-l})_{l=1,\dots,r}$  of  $V_-$  (as in §2), we get a (self-dual) decomposition

$$V = \bigoplus_{l=1}^r V^l \oplus V_o \oplus \bigoplus_{l=1}^r V^{-l},$$

where  $V^l = Fe_l = \left( \bigoplus_{k \neq -l} V^k \oplus V_o \right)^\perp$ . Such a decomposition (which we will also call a Witt decomposition) corresponds to the choice of an apartment  $\mathcal{A}$  in  $I$ : in terms of lattice functions,  $j_h(\mathcal{A})$  is the set of self-dual lattice functions  $\Lambda$  such that

$$\Lambda(s) = \bigoplus_{l=1}^r (V^l \cap \Lambda(s)) \oplus (V_o \cap \Lambda(s)) \oplus \bigoplus_{l=1}^r (V^{-l} \cap \Lambda(s)), \quad \text{for all } s \in \mathbb{R},$$

that is,  $\Lambda$  is *split* by the decomposition (cf. Proposition 4.1).

Similarly, the choice of an (enlarged) apartment  $\mathcal{A}^1$  in  $I_\beta^1 = \prod_{i \in J_o \cup J_+} I_i^1$  is given by similar  $E_i$ -decompositions of  $V_i$  for  $i \in J_o$  and (without the self-duality restriction)  $i \in J_+$ .

**Proposition 8.1.** *Let  $\mathcal{A}^1$  be an (enlarged) apartment of  $I_\beta^1$ . Then there is an apartment  $\mathcal{A}$  of  $I$  such that  $j_\beta(\mathcal{A}^1) \subset \mathcal{A}$ .*

*Proof.* We write  $\mathcal{A}^1 = \prod_{i \in J_o \cup J_+} \mathcal{A}_i^1$ , with  $\mathcal{A}_i^1$  an (enlarged) apartment in  $I_i^1$ .

As above, for each  $i \in J_o$ , the apartment  $\mathcal{A}_i^1$  corresponds to a Witt  $E_i$ -decomposition of  $V^i$

$$V_i = \bigoplus_{l=1}^{r_i} V_i^l \oplus V_{i,o} \oplus \bigoplus_{l=1}^{r_i} V_i^{-l},$$

with  $V_i^l = \left( \bigoplus_{k \neq -l} V_i^k \oplus V_{i,o} \right)^\perp$ ,  $\dim_{E_i} V_i^l = 1$  and  $r_i$  the ( $E_i$ -)Witt index of  $V_i$ . We write  $\text{Latt}_{\mathfrak{o}_{E_i}}^{\mathcal{A}_i^1}(V_i)$  for the set of lattice functions split by this decomposition, and  $\text{Latt}_{\mathfrak{o}_{E_i}, h_i}^{\mathcal{A}_i^1}(V_i)$  for the subset of self-dual lattice functions, so that  $j_{h_i}(\mathcal{A}_i^1) = \text{Latt}_{\mathfrak{o}_{E_i}, h_i}^{\mathcal{A}_i^1}(V_i)$ .

Also, for each  $i \in J_+$ , the apartment  $\mathcal{A}_i^1$  corresponds to a decomposition of  $V_i$  as a sum of 1-dimensional  $E_i$ -subspaces,

$$V_i = \bigoplus_{l=1}^{r_i} V_i^l,$$

with  $r_i = \dim_{E_i} V_i$ . As above,  $j_i(\mathcal{A}_i^1) = \text{Latt}_{\mathfrak{o}_{E_i}}^{\mathcal{A}_i^1}(V_i)$ , the set of lattice functions split by this decomposition.

We also take the dual splitting of  $V_{-i}$  as a sum of 1-dimensional  $E_{-i}$ -subspaces,

$$V_{-i}^l = \left( \bigoplus_{k \neq l} V_i^k \right)^\perp.$$

We remark that, if  $\Lambda \in \text{Latt}_{\mathfrak{o}_{E_i}}^{\mathcal{A}_i^1}(V_i)$  then  $\Lambda_i^{\#i}$  is split by this decomposition.

Now, for  $i \in J_o \cup J_+$  and  $1 \leq l \leq r_i$ , we decompose  $V_i^l$  as a sum of 1-dimensional  $F$ -subspaces as follows: fix  $v \in V_i^l$ ,  $v \neq 0$ , and let  $\mathcal{B}_i$  be an  $F$ -basis for  $E_i$  which splits the  $\mathfrak{o}_F$ -lattice sequence  $s \mapsto \mathfrak{p}_{E_i}^{\lceil s/e(E_i/F) \rceil}$ ; then we take the decomposition

$$V_i^l = \bigoplus_{b \in \mathcal{B}_i} Fbv.$$

Note that any  $\mathfrak{o}_{E_i}$ -lattice sequence in  $V_i^l$  is split by this decomposition. For  $i \in J_o$ , we also take the dual decomposition of  $V_i^{-l}$  and, for  $i \in J_+$ , the dual decomposition of  $V_{-i}^l$ .

Now we need to decompose the anisotropic parts  $W := \bigoplus_{i \in J_o} V_{i,o}$  suitably, for which we cheat. Let  $\mathbf{G}_o$  denote the classical group associated to the restriction of the form  $h$  to  $W$  and, for  $i \in J_o$ , let  $\mathbf{H}_{i,o}$  denote the group associated to the restriction of the form  $h_i$  to  $V_{i,o}$ . Note that the groups  $H_{i,o}$  are compact so the building  $I_{\beta,o}^1 := I^1(\mathbf{H}_{i,o}, E_i)$  is reduced to a point.

Now, our constructions in §6 give an embedding of  $I_{\beta,o}^1$  in the building  $I_o^1 := I^1(\mathbf{G}_o, F)$  and the image is certainly contained in some apartment. Hence there is a Witt  $F$ -decomposition of  $W$  which splits the (unique) self-dual lattice sequence in  $W$  corresponding to  $I_{\beta,o}^1$ , and this is the decomposition we take.

Altogether, we have described a Witt  $F$ -decomposition of  $V$ , which corresponds to an apartment  $\mathcal{A}$  of  $I$ . We denote by  $\text{Latt}_{\mathfrak{o}_F, h}^{\mathcal{A}}(V)$  the set of self-dual lattice functions in  $V$  which are split by this splitting, so that  $j_h(\mathcal{A}) = \text{Latt}_{\mathfrak{o}_F, h}^{\mathcal{A}}(V)$ .

Finally, by construction it is clear that  $\tilde{j}_\beta \circ \iota_h$  maps  $\prod_{i \in J_o} \text{Latt}_{\mathfrak{o}_{E_i}, h_i}^{\mathcal{A}^1}(V_i) \times \prod_{i \in J_+} \text{Latt}_{\mathfrak{o}_{E_i}}^{\mathcal{A}^1}(V_i)$  into  $\text{Latt}_{\mathfrak{o}_F, h}^{\mathcal{A}}(V)$  so  $j_\beta(\mathcal{A}^1) \subset \mathcal{A}$ , as required.

## 9. Compatibility with Lie algebra filtrations

In this section, we fix  $H_k$ -equivariant identifications  $j_k : I^1(H_k, E_k) \rightarrow \text{Latt}_{\mathfrak{o}_{E_k}}^1(V_k)$ ,  $k \in J^+$ . They give rise to the map  $j_\beta : I_\beta^1 \rightarrow I(G, H)$  defined in §6.

Let  $x \in I(G, F) = I^1(G, F)$ , that we see as a self-dual lattice function  $\Lambda$  in  $\text{Latt}_h^1(V)$ . To  $x$  we can associate a filtration  $(\mathfrak{g}_{x,r})_{r \in \mathbb{R}}$  of the Lie algebra  $\mathfrak{g}$  as follows. First  $x$  defines a filtration  $(\tilde{\mathfrak{g}}_{x,r})_{r \in \mathbb{R}}$  of  $\tilde{\mathfrak{g}}$  by

$$\tilde{\mathfrak{g}}_{x,r} = \{a \in \tilde{\mathfrak{g}} ; a\Lambda(s) \subset \Lambda(s+r), s \in \mathbb{R}\}, r \in \mathbb{R}.$$

We then define

$$\mathfrak{g}_{x,r} := \tilde{\mathfrak{g}}_{x,r} \cap \mathfrak{g} = \{a \in \mathfrak{g} ; a\Lambda(s) \subset \Lambda(s+r), s \in \mathbb{R}\}, r \in \mathbb{R}. \quad (1)$$

Similarly a point  $x$  of  $I_\beta^1$  defines a filtration  $(\mathfrak{h}_{x,r})_{r \in \mathbb{R}}$  of  $\mathfrak{h}$ . Write  $x = (x_k)_{k \in J \cup J_o}$ ,  $x_k \in I^1(H_k, E_k)$ ; each  $x_k$  corresponding to a lattice function  $\Lambda_k$  of  $\text{Latt}_{\mathfrak{o}_{E_k}}(V_k)$  (with  $\Lambda_k^{\sharp k} = \Lambda_{-k}$ ,  $k \in J \cup J_o$ ). We then define

$$(2) \quad \mathfrak{h}_{x,r} := \bigoplus_{k \in J^+ \cup J_o} \mathfrak{h}_{x_k, r}^k, r \in \mathbb{R},$$

where

$$\mathfrak{h}_{x_k,r}^k = \{a \in \text{Lie}(H_k) ; a\Lambda_k(s) \subset \Lambda_k(s+r), s \in \mathbb{R}\}, r \in \mathbb{R}, k \in J^+ \cup J_o .$$

The filtration  $(\mathfrak{h}_{x,r})_{r \in \mathbb{R}}$  only depends on the image  $\bar{x}$  of  $x$  in the non-enlarged building  $I_\beta$ . One can prove that for  $x \in I(G, F)$ ,  $(\mathfrak{g}_{x,r})_{r \in \mathbb{R}}$  is the filtration of  $\mathfrak{g}$  attached to  $x$  defined by Moy and Prasad [MP]. Similarly, when  $\beta$  is semisimple and  $x \in I^1(H, F)$ ,  $(\mathfrak{h}_{x,r})_{r \in \mathbb{R}}$  is the filtration of  $\mathfrak{h}$  attached to  $\bar{x}$  defined in loc. cit. The proof of this fact is announced by B. Lemaire and J.-K. Yu [BY].

**Lemma 9.1.** *Let us see  $\mathfrak{h}$  as being canonically embedded in  $\tilde{\mathfrak{h}} = \text{End}_E V = \bigoplus_{k \in J \cup J_o} \text{End}_{E_k} V_k$*

via

$$(a_k)_{k \in J^+ \cup J_o} \mapsto (b_k)_{k \in J \cup J_o} ,$$

where  $b_k = a_k$ ,  $k \in J_o$ , and  $b_{-k} = -a_k^\sigma$ ,  $k \in J^+$ . Fix  $x \in I_\beta^1$  as before and consider the  $\mathfrak{o}_F$ -lattice function in  $V$  given by

$$\Lambda = \bigoplus_{k \in J \cup J_o} \Lambda_k \text{ (notation of §6) .}$$

For  $r \in \mathbb{R}$ , let

$$\tilde{\mathfrak{h}}_{x,r} = \{a \in \tilde{\mathfrak{h}} ; a\Lambda(s) \subset \Lambda(s+r), s \in \mathbb{R}\}, r \in \mathbb{R} .$$

Then we have  $\mathfrak{h}_{x,r} = \tilde{\mathfrak{h}}_{x,r} \cap \mathfrak{h}$ ,  $r \in \mathbb{R}$ .

*Proof.* Indeed, for all  $a = (a_k)_{k \in J \cup J_o} \in \text{End}_E V$ , we have  $a \in \tilde{\mathfrak{h}}_{x,r} \cap \mathfrak{h}$  if and only if  $a + a^\sigma = 0$  and  $a\Lambda(s) \subset \Lambda(s+r)$ ,  $s \in \mathbb{R}$ , i.e.

$$a_k \Lambda_k(s) \subset \Lambda_k(s+r), s \in \mathbb{R}, k \in J \cup J_o .$$

For  $k \in J_o$ , these conditions can be rewritten  $a_k \in \text{Lie}(H_k)$  and  $a_k \Lambda_k(s) \subset \Lambda_k(s+r)$ ,  $s \in \mathbb{R}$ , that is  $a_k \in \mathfrak{h}_{x,r}^k$ , as required. For  $k \in J$ , these conditions can be rewritten  $a_{-k} = -a_k^\sigma$  and

$$a_k \Lambda_k(s) \subset \Lambda_k(s+r), s \in \mathbb{R} \tag{a}$$

$$-a_k^\sigma \Lambda_k^\sharp(s) \subset \Lambda_k^\sharp(s+r), s \in \mathbb{R} . \tag{b}$$

So we must prove that conditions (a) and (b) are equivalent. By symmetry we only prove one implication. Applying the duality  $\sharp_k$  on lattices of  $V_k$  to inclusion (b), we obtain

$$\Lambda_k((-s-r)+) \subset [a_k^\sigma \Lambda_k^\sharp(s)]^\sharp, s \in \mathbb{R},$$

with

$$[a_k^\sigma \Lambda_k^\sharp(s)]^\sharp = \{v \in V_k ; a_k v \in \Lambda_k((-s)+)\}, s \in \mathbb{R} .$$

So we have

$$a_k \Lambda_k((-s-r)+) \subset \Lambda_k((-s)+) \subset \Lambda_k(-s), s \in \mathbb{R} ,$$

that is

$$a_k \Lambda(s+) \subset \Lambda_k(s+r), s \in \mathbb{R} .$$



On each open interval  $(u, v)$  where  $\Lambda_k$  is constant, we have

$$a_k \Lambda_k(s+) = a_k \Lambda_k(s) \subset \Lambda_k(s+r) ,$$

and (a) is true for  $s \in (u, v)$ . Finally if  $s_o$  is a jump of  $\Lambda_k$  with  $\Lambda_k$  constant on  $(t, s_o]$ , we have

$$a_k \Lambda_k(s_o) = a_k \Lambda_k(s_o+) \subset \Lambda_k(s_o+r) , \quad s \in (t, s_o) .$$

So

$$a_k \Lambda_k(s_o) \subset \bigcap_{s \in (t, s_o)} \Lambda_k(s+r) = \Lambda_k(s_o+r) ,$$

$\Lambda_k$  being left continuous, and (a) is then true for all  $s \in \mathbb{R}$ .

**Proposition 9.2.** *Let  $x \in I_\beta^1$ . Then we have*

$$\mathfrak{g}_{j_\beta(x), r} \cap \mathfrak{h} = \mathfrak{h}_{x, r} , \quad r \in \mathbb{R} .$$

*Proof.* Indeed, with the notation of (9.1) and by definition of  $j_\beta$ , we easily see that

$$\tilde{\mathfrak{g}}_{j_\beta(x), r} \cap \tilde{\mathfrak{h}} = \tilde{\mathfrak{h}}_{x, r} .$$

So our result is now a corollary of (9.1) since  $\mathfrak{h} = \mathfrak{g} \cap \tilde{\mathfrak{h}}$ .

## 10. A unicity result for the general linear group

As in [BL]§I.2, we define an equivalence relation  $\sim$  on  $\text{Latt}^1(V)$  by  $\Lambda_1 \sim \Lambda_2$  if there exists  $s \in \mathbb{R}$  such that  $\Lambda_1(s) = \Lambda_2(r+s)$ ,  $s \in \mathbb{R}$ . Then  $\sim$  is compatible with the  $\tilde{G}$ -action and the quotient  $\text{Latt}_{\mathfrak{o}_F}(V) := \text{Latt}^1(V) / \sim$  is naturally a  $\tilde{G}$ -set. We shall denote by  $\bar{\Lambda}$  an element of  $\text{Latt}_{\mathfrak{o}_F}(V)$ , where  $\Lambda$  is a representative in  $\text{Latt}^1(V)$ . As a consequence of [BL]§I.2 and [BT1], there is a unique affine and  $\tilde{G}$ -equivariant map  $j : \tilde{I} \rightarrow \text{Latt}_{\mathfrak{o}_F}(V)$ , where  $\tilde{I}$  denotes the non-enlarged building of  $\tilde{G}$ .

We fix an element  $\beta$  of  $\tilde{\mathfrak{g}}$  satisfying

$$(H) \quad E : F[\beta] \text{ is a field .}$$

As in §5 we denote by  $\tilde{\mathfrak{h}} = \text{End}_E V$  the centralizer of  $\beta$  in  $\tilde{\mathfrak{g}}$  and by  $\tilde{H} = \text{Aut}_E V$  its centralizer in  $\tilde{G}$ . There is a canonical identification of the non-enlarged affine building  $\tilde{I}_\beta$  of  $\tilde{H}$  with the  $\tilde{H}$ -set  $\text{Latt}_{\mathfrak{o}_E}(V)$ . Here we normalize the lattice functions of  $\text{Latt}_{\mathfrak{o}_E}^1(V)$  by the condition  $\Lambda(s + v(\pi_E)) = \pi_E \Lambda(s)$ ,  $s \in \mathbb{R}$ , where  $\pi_E$  is a uniformizer of  $E$ .

Any  $\bar{\Lambda} \in \text{Latt}_{\mathfrak{o}_F}(V)$  defines a filtration  $(\tilde{\mathfrak{g}}_{\bar{\Lambda}, r})_{r \in \mathbb{R}}$  by

$$\tilde{\mathfrak{g}}_{\bar{\Lambda}, r} = \{a \in \text{End}_F V ; a \Lambda(s) \subset \Lambda(r+s), \quad s \in \mathbb{R}\} .$$

Then the map  $\text{End}(\bar{\Lambda}) : r \mapsto \tilde{\mathfrak{g}}_{\bar{\Lambda}, r}$  is an element of  $\text{Latt}^1 \tilde{\mathfrak{g}}$ . The map  $\bar{\Lambda} \mapsto \text{End}(\bar{\Lambda})$ ,  $\text{Latt}_{\mathfrak{o}_F} V \rightarrow \text{Latt}^1 \tilde{\mathfrak{g}}$  is a  $\tilde{G}$ -equivariant injection (cf. [BL]§4) for the action of  $G$  on  $\text{Latt}^1 \tilde{\mathfrak{g}}$  by conjugation. Its image is  $\text{Latt}^2 \tilde{\mathfrak{g}}$ . From now on we shall canonically identify  $\tilde{I}$  (resp.  $\tilde{I}_\beta$  with  $\text{Latt}^2 \tilde{\mathfrak{h}}$ ).

Let us recall the main result of [BL].

**Theorem 10.1.** *There exists a unique affine and  $\tilde{H}$ -equivariant map  $\tilde{j}_\beta : \tilde{I}_\beta \rightarrow \tilde{I}$ . It is injective, maps any apartment into an apartment and is compatible with the Lie algebra filtrations in the following sense:*

$$(10.2) \quad \tilde{\mathfrak{g}}_{\tilde{j}_\beta, r} \cap \tilde{\mathfrak{h}} = \tilde{\mathfrak{h}}_{x, r}, \quad x \in \tilde{I}_\beta, \quad r \in \mathbb{R}.$$

Let us recall how  $\tilde{j}_\beta$  is constructed. If  $x \in \tilde{I}_\beta$  corresponds to  $\text{End}(\bar{\Lambda}) \in \text{Latt}^2 \tilde{\mathfrak{h}}$ , then  $\tilde{j}(x)$  simply corresponds to  $\text{End}(\bar{\Lambda})$ , where  $\Lambda$ , an  $\mathfrak{o}_E$ -lattice function in  $V$ , is now considered as an  $\mathfrak{o}_F$ -lattice function.

**Theorem 10.3.** *Let  $x \in \tilde{I}_\beta$  and  $y \in \tilde{I}$  satisfying*

$$\tilde{\mathfrak{g}}_{y, r} \cap \tilde{\mathfrak{h}} \supset \tilde{\mathfrak{h}}_{x, r}, \quad r \in \mathbb{R}.$$

*Then  $y = \tilde{j}_\beta(x)$ . As a consequence the map  $\tilde{j}_\beta$  is characterized by property (10.2).*

*Proof.* Assume that  $x$  and  $y$  correspond to elements  $\bar{\Lambda}_x$  and  $\bar{\Lambda}_y$  of  $\text{Latt}_{\mathfrak{o}_E}(V)$  and  $\text{Latt}_{\mathfrak{o}_F}(V)$  respectively.

**Lemma 10.4.** *Under the assumption of (10.2),  $\Lambda_y$  is an  $\mathfrak{o}_E$ -lattice function.*

*Proof.* To prove that  $\Lambda_y$  is an  $\mathfrak{o}_E$ -lattice function we must prove that it is normalized by  $E^\times = \langle \pi_E \rangle \mathfrak{o}_E^\times$ , or equivalently:

$$(10.5) \quad x \tilde{\mathfrak{g}}_{y, r} x^{-1} = \tilde{\mathfrak{g}}_{y, r}, \quad x \in E^\times, \quad r \in \mathbb{R}.$$

We first notice that  $\mathfrak{o}_E \subset \tilde{\mathfrak{h}}_{x, 0} \subset \tilde{\mathfrak{g}}_{y, 0}$ , so that  $\mathfrak{o}_E^\times \subset \tilde{\mathfrak{g}}_{y, 0}^\times$  and (10.5) is true for  $x \in \mathfrak{o}_E^\times$ . We are reduced to proving (10.5) when  $x = \pi_E$ .

We have  $\pi_E \in \tilde{\mathfrak{h}}_{x, 1/e} \subset \tilde{\mathfrak{g}}_{y, 1/e}$  and  $\pi_E^{-1} \in \tilde{\mathfrak{h}}_{x, -1/e} \subset \tilde{\mathfrak{g}}_{y, -1/e}$ , where  $e = e(E/F)$ . It follows that

$$(10.6) \quad \pi_E \tilde{\mathfrak{g}}_{y, r} \pi_E^{-1} \subset \tilde{\mathfrak{g}}_{y, 1/e} \tilde{\mathfrak{g}}_{y, r} \tilde{\mathfrak{g}}_{y, -1/e} \subset \tilde{\mathfrak{g}}_{y, r}, \quad r \in \mathbb{R}.$$

Consider the duality “\*” on subsets of  $\tilde{\mathfrak{g}}$  given by

$$S^* = \{a \in \tilde{\mathfrak{g}} ; \text{Tr}(aS) \subset \mathfrak{p}_F\}, \quad S \subset \tilde{\mathfrak{g}},$$

where  $\text{Tr}$  is the trace map. Recall ([BL](6.3)) that  $(\tilde{\mathfrak{g}}_{y, r})^* = \tilde{\mathfrak{g}}_{y, (-r)_+}$ ,  $r \in \mathbb{R}$ . Using a well known property of the trace map, we observe that

$$(\pi_E \tilde{\mathfrak{g}}_{y, r} \pi_E^{-1})^* = \pi_E (\tilde{\mathfrak{g}}_{y, r})^* \pi_E^{-1}, \quad r \in \mathbb{R}.$$

So applying the duality to (10.6), we obtain

$$\tilde{\mathfrak{g}}_{y, (-r)_+} \subset \pi_E \tilde{\mathfrak{g}}_{y, (-r)_+} \pi_E^{-1}, \quad r \in \mathbb{R}.$$

We have proved that on each open interval  $(r_1, r_2)$  where the lattice function  $(\tilde{\mathfrak{g}}_{y, r})_{r \in \mathbb{R}}$  is constant, we have both containments

$$\pi_E \tilde{\mathfrak{g}}_{y, r} \pi_E^{-1} \subset \tilde{\mathfrak{g}}_{y, r} \quad \text{and} \quad \pi_E \tilde{\mathfrak{g}}_{y, r} \pi_E^{-1} \subset \tilde{\mathfrak{g}}_{y, r}, \quad r \in \mathbb{R}.$$

So by continuity we have  $\pi_E \tilde{\mathfrak{g}}_{y,r} \pi_E^{-1} = \tilde{\mathfrak{g}}_{y,r}$ , for all  $r$ , as required.

Let us return to the proof of (10.3). Since  $\Lambda_y$  is an  $\mathfrak{o}_E$ -lattice function, we have

$$\tilde{\mathfrak{g}}_{y,r} \cap \tilde{\mathfrak{h}} = \tilde{\mathfrak{h}}_{x',r}, \quad r \in \mathbb{R},$$

where  $x' \in \tilde{I}_\beta$  is attached to  $\bar{\Lambda}_y$ ,  $\Lambda_y$  being seen as an  $\mathfrak{o}_E$ -lattice function. So by injectivity of the map  $\text{Latt}_{\mathfrak{o}_E}^1(V) \rightarrow \text{Latt}_{\tilde{\mathfrak{h}}}^2(V)$ , we have  $\bar{\Lambda}_x = \bar{\Lambda}_y$  and  $y = \tilde{j}_\beta(x)$  by definition.

## 11. A unicity result in the 1-block case and a conjecture

With the notation of §5, we consider an element  $\beta \in \mathfrak{g}$  satisfying:

$$(11.1) \quad E := F[\beta] \subset \tilde{\mathfrak{g}} \text{ is a field and } \beta \neq 0.$$

We fix an  $\varepsilon$ -hermitian form  $h_E$  on the  $E$ -vector space  $V$  relative to  $\sigma_E$  and we assume that it satisfies (5.3) as well as the condition  $\mathcal{J} = \mathfrak{p}_{E^\circ}$  of §5. This allows us to identify  $I_\beta^1$  with  $\text{Latt}_{h_E}^1(V)$ . Identifying  $I$  with  $\text{Latt}_h(V)$ , the map  $j_\beta$  of §6 is simply given by

$$j_\beta(\Lambda) = \Lambda, \quad \Lambda \in \text{Latt}_{h_E}^1(V),$$

where on the right hand side  $\Lambda$  is considered as an  $\mathfrak{o}_F$ -lattice function.

**Theorem 11.2.** *Under the assumption (11.1), let  $x \in I_\beta^1$  and  $y \in I$  satisfying*

$$(11.3) \quad \mathfrak{g}_{y,r} \cap \mathfrak{h} = \mathfrak{h}_{x,r}, \quad r \in \mathbb{R}.$$

*Then  $y = j_\beta(x)$ . In particular the map  $j_\beta$  is characterized by compatibility with the Lie algebra filtrations.*

*Proof.* The point  $x$  (resp.  $y$ ) corresponds to a self-dual lattice function  $\Lambda_x \in \text{Latt}_{h_E}^1(V)$  (resp.  $\Lambda_y \in \text{Latt}_h^1(V)$ ). We may see  $x$  and  $y$  as points of  $\text{Latt}_{\mathfrak{o}_E}^1(V)$  and  $\text{Latt}_{\mathfrak{o}_F}^1(V)$  respectively and they give rise to filtrations of  $\tilde{\mathfrak{h}}$  and  $\tilde{\mathfrak{g}}$  as in §9:  $(\tilde{\mathfrak{h}}_{x,r})_{r \in \mathbb{R}}$  and  $(\tilde{\mathfrak{g}}_{y,r})_{r \in \mathbb{R}}$ . Write

$$\mathfrak{g}_{y,r}^+ = \{a \in \tilde{\mathfrak{g}}_{y,r} ; a = a^\sigma\}, \quad r \in \mathbb{R}$$

and

$$\mathfrak{h}_{x,r}^+ = \{a \in \tilde{\mathfrak{h}}_{x,r} ; a = a^\sigma\}, \quad r \in \mathbb{R}$$

Since 2 is invertible in  $\mathfrak{o}_F$ , we have:

$$\tilde{\mathfrak{g}}_{y,r} = \mathfrak{g}_{y,r} \oplus \mathfrak{g}_{y,r}^+ \text{ and } \tilde{\mathfrak{h}}_{y,r} = \mathfrak{h}_{x,r} \oplus \mathfrak{h}_{x,r}^+, \quad r \in \mathbb{R}.$$

Write

$$r_o = v_{\Lambda_x}(\beta) := \text{Sup}\{r \in \mathbb{R} ; \beta \in \tilde{\mathfrak{h}}_{x,r}\}.$$

Since  $\beta \in E^\times$ , it normalizes  $\Lambda_x$  so that  $\beta \tilde{\mathfrak{h}}_{x,r} = \tilde{\mathfrak{h}}_{x,r+r_o}$ ,  $r \in \mathbb{R}$ . Moreover since  $\beta$  is central in  $\tilde{\mathfrak{h}}$ , we easily have that  $\mathfrak{h}_{x,r}^+ = \beta \mathfrak{h}_{x,r-r_o}^+$ ,  $r \in \mathbb{R}$ . Hence, for  $r \in \mathbb{R}$ , we have

$$\mathfrak{h}_{x,r}^+ = \beta(\mathfrak{g}_{y,r-r_o} \cap \mathfrak{h}) = \beta(\mathfrak{g}_{y,r-r_o} \cap \tilde{\mathfrak{h}}) \subset \mathfrak{g}_{y,r} \cap \tilde{\mathfrak{h}}.$$

It follows that, for  $x \in \mathbb{R}$ , we have:

$$\tilde{\mathfrak{h}}_{x,r} = \mathfrak{h}_{x,r} \oplus \mathfrak{h}_{x,r}^+ \subset \mathfrak{g}_{y,r} \cap \tilde{\mathfrak{h}} \oplus \mathfrak{g}_{y,r}^+ \cap \tilde{\mathfrak{h}} \subset \tilde{\mathfrak{g}}_{y,r} \cap \tilde{\mathfrak{h}}.$$

By applying (10.3), we obtain  $\bar{\Lambda}_y = \tilde{j}_\beta(\bar{\Lambda}_x)$ , that is  $\bar{\Lambda}_y = \bar{\Lambda}_x$ . In particular we have  $\text{End}(\Lambda_x) = \text{End}(\Lambda_y) \in \text{Latt}_\sigma^2 \tilde{\mathfrak{h}}$ . But by (3.5) we have  $\Lambda_x = \Lambda_y$ , as required.

Let us give an example. Assume that  $G = \text{Sp}_2(F) = \text{SL}(2, F)$  (here  $F = F_o$ ) and take  $\beta \in \mathfrak{g}$  such that  $E/F$  is quadratic and ramified. Then  $H$  is the group  $E^1$  of norm 1 elements in  $E$ . The building of  $H$  is reduced to a point  $\{x\}$ . The group  $E^\times$  fixes a unique chamber  $C$  of  $I$  and  $H \subset E^\times$  fixes  $C$  pointwise. There are infinitely many maps  $j : I_\beta^1 \rightarrow I$  which are affine and  $G$ -equivariant; indeed  $j(x)$  can be any point of  $C$ . On the other hand there is a unique map  $j : I_\beta^1 \rightarrow I$  which is compatible with the Lie algebra filtrations: it maps  $x$  to the isobarycenter of  $C$ .

We conjecture that when  $J = \emptyset$  (notation of §5) then the map  $j_\beta$  of §6 is characterized by condition (11.3). We may address the more general (but more informal) question. Being given two  $F$ -reductive groups  $\mathbf{H}$  and  $\mathbf{G}$ , as well as a morphism of algebraic groups  $\varphi : \mathbf{H} \rightarrow \mathbf{G}$ , is there an affine and  $\mathbf{H}(F)$ -equivariant map  $I(\mathbf{H}, F) \rightarrow I(\mathbf{G}, F)$  which is compatible with the Lie algebra filtrations defined by Moy and Prasad. When is it characterized by this last property?

## References

- [BK] C.J. Bushnell and P.C. Kutzko, *The admissible dual of  $\text{GL}(N)$  via compact open subgroups*, Ann. of Math. Studies 129, Princeton Univ. Press.
- [BL] P. Broussous and B. Lemaire, *Building of  $\text{GL}(m, D)$  and centralizers*, Transformations Groups, Vol. 7, No. 1, 2002, pp. 15–50.
- [BT1] F. Bruhat and J. Tits, *Schémas en groupes et immeubles des groupes classiques sur un corps local, 1ère partie: le groupe linéaire général*, Bull. Soc. Math. Fr., 112, 1984, pp. 259–301.
- [BT2] F. Bruhat and J. Tits, *Schémas en groupes et immeubles des groupes classiques sur un corps local, 2ème partie: groupes unitaires*, Bull. Soc. Math. Fr., 115, 1987, pp. 141–195.
- [BY] B. Lemaire and J.-K. Yu, Private communication.
- [Inv] M.-A. Knus, A. Merkurjev, M. Rost and J.-P. Tignol, *The book of involutions*, AMS Colloquium publications, Vol. 44, 1998.
- [Mor] L. Morris, *Some tamely ramified supercuspidal representations of symplectic groups*, Proc. London Math. Soc. (3), Vol. 63, 1991, pp. 519–551.
- [MP] A. Moy and G. Prasad, *Unrefined minimal  $K$ -types for  $p$ -adic groups*, Invent. Math., Vol. 116, 1994, pp. 393–408.
- [Sch] W. Scharlau, *Quadratic and Hermitian forms*, Grundlehren der Mathematischen Wissenschaften 270, Springer-Verlag, Berlin, 1985.
- [S1] S. Stevens, *Semisimple strata for  $p$ -adic classical groups*, Ann. Sci. École Norm. Sup. (4), Vol. 35, No. 3, 2002, pp. 423–435.

[S2] S. Stevens, *Semisimple characters for  $p$ -adic classical groups*, Preprint, September 2003.

Université de Poitiers  
Département de Mathématiques  
UMR 6086 du CNRS  
Bd Marie et Pierre Curie  
Téléport 2 – BP 30179  
86962 Futuroscope Chasseneuil Cedex  
France

School of Mathematics  
University of East Anglia  
Norwich NR4 7TJ  
United Kingdom