Influence of Climate Variables upon selected Infectious Diseases in Asir Region, Saudi Arabia

Turki Khalufa Faraj

A thesis submitted in fulfilment of the requirement for the Degree of Doctor of Philosophy to the University of East Anglia

> School of Environmental Sciences Norwich, England

> > September 2011

"This copy of the thesis has been supplied on condition that anyone who consults it is understood to recognise that its copyright rests with the author and that use of any information derived there from must be in accordance with current UK Copyright Law. In addition, any quotation or extract must include full attribution."

Abstract

Although the incidence of some infectious diseases (Malaria, Cutaneous Leishmaniasis (CL), and Schistosomiasis) in the Asir Region, Saudi Arabia, has decreased over the last decade, they still cause significant health problems. This study tests the hypotheses that there are relationships between climate variables (maximum, mean and minimum temperature, rainfall and relative humidity) and these diseases in Asir Region. Monthly malaria, CL and schistosomiasis cases and climate variable data were obtained for the study area between 1995 and 2006, 1996 and 2007 and 1998 and 2009 respectively.

Multiple regressions on the number of cases of each disease were performed against the climate variables. The climate variables were lagged to account for time gaps between weather events and disease. Different relationships with climate may occur at different times of the year and so separate models were created for each season.

Within Asir, there are two separate malaria endemic areas in the lowlands: Tehama of Asir (TA) and Tehama of Qahtan (TQ). The results revealed that in these areas, rainfall in the current and previous month was positively associated (P<0.05 and P<0.001 respectively) with an increase in malaria cases during the summer. This may be due to increased breeding and survival of vectors after the rainfall. Temperature was not an important factor.

Temperature was the most important factor associated with CL cases in all Asir and in the highlands area during the summer and beginning of autumn. If the maximum temperature in the previous 2 to 4 months was elevated there were more cases (P<0.001). Rainfall also plays a role in CL transmission in the lowlands. There were more cases of CL if rainfall was elevated in the previous 1 to 3 months (P<0.05).

Temperature was an important factor associated with schistosomiasis throughout Asir during the summer and beginning of autumn. If the maximum temperature in the previous 3 to 4 months was elevated, there were more cases of schistosomiasis (P<0.05). Rainfall also plays an important role in schistosomiasis transmission in the highlands. There were more cases if rainfall in the previous 6 to 7 months (P<0.005) was elevated. This occurs during spring and summer seasons

Based on these findings and an understanding of how these diseases are affected by climate it should be possible to design an early warning system using the weather to predict each disease's incidence. Therefore, the results of this study have important implications for policies to improve public health in this region.

Content	Page No.
Abstract	1
Table of Contents	2
List of Tables	9
List of Figures	14
List of Abbreviations	18
Acknowledgements	20
Chapter 1 Introduction	21
Background	22
1.1 What are Infectious Diseases?	22
1.2 The Global Importance of Infectious Diseases to Health	23
1.3 Climate Variables and Climate Change	24
1.4 The Influence of Climate Variables upon Infectious Diseases	25
1.5 Climate Change and its Potential Impact Health	26
1.6 Influence of Climate Variables upon Selected Infectious Diseases in the Asir Region, Saudi Arabia.	30
1.6.1 Overview	30
1.6.2 Importance of the Study	30
1.7 Structure of Thesis	31
Chapter 2 Study Area	32
2. Introduction	33
2.1 Saudi Arabia	33
2.1.1 Geography and Demography	33
2.1.2 Climate	37

Table of Contents

2.1.3 Education	37
2.1.4 Drinking Water Services	38
2.1.5 Health Service	38
2.1.6 Zoogeographical Areas of Saudi Arabia	41
2.1.7 Distribution of Malaria, Leishmaniasis and Schistosomiasis in Saudi Arabia	41
2.2 Asir Region	45
2.2.1 Geography and Demography	45
2.2.2 Climate in Asir	47
2.2.3 Drinking Water Services	47
2.2.4 Environmental Problems	48
2.2.5 Health Services	48
2.2.6 Distribution and Endemicity of the Malaria, Leishmaniasis and Schistosomiasis in the Asir Region	51
0	
Chapter 3 Malaria	52
Chapter 3 Malaria 3.1 Introduction	52
Chapter 3 Malaria 3.1 Introduction 3.2 Malaria Transmission	52 53 53
Chapter 3 Malaria 3.1 Introduction 3.2 Malaria Transmission 3.3 Life Cycle of Malaria	52 53 53 54
Chapter 3 Malaria 3.1 Introduction 3.2 Malaria Transmission 3.3 Life Cycle of Malaria 3.4 Symptoms of Malaria	52 53 53 54 55
Chapter 3 Malaria 3.1 Introduction 3.2 Malaria Transmission 3.3 Life Cycle of Malaria 3.4 Symptoms of Malaria 3.5 Preventing and Treating Malaria	52 53 53 54 55 56
Chapter 3 Malaria 3.1 Introduction 3.2 Malaria Transmission 3.3 Life Cycle of Malaria 3.4 Symptoms of Malaria 3.5 Preventing and Treating Malaria 3.6 Relationship Between Malaria and Climate	52 53 53 54 55 56 57
Chapter 3 Malaria 3.1 Introduction 3.2 Malaria Transmission 3.3 Life Cycle of Malaria 3.4 Symptoms of Malaria 3.5 Preventing and Treating Malaria 3.6 Relationship Between Malaria and Climate 3.7 Previous Studies into Malaria (Transmission, Vector and Incidents) and Climate Variables	52 53 53 54 55 56 57 58
Chapter 3 Malaria 3.1 Introduction 3.2 Malaria Transmission 3.3 Life Cycle of Malaria 3.4 Symptoms of Malaria 3.5 Preventing and Treating Malaria 3.6 Relationship Between Malaria and Climate 3.7 Previous Studies into Malaria (Transmission, Vector and Incidents) and Climate Variables 3.8 Malaria in Saudi Arabia	52 53 53 54 55 56 57 58 65
Chapter 3 Malaria 3.1 Introduction 3.2 Malaria Transmission 3.3 Life Cycle of Malaria 3.4 Symptoms of Malaria 3.5 Preventing and Treating Malaria 3.6 Relationship Between Malaria and Climate 3.7 Previous Studies into Malaria (Transmission, Vector and Incidents) and Climate Variables 3.8 Malaria in Saudi Arabia Topographical Classification of Asir Region and Endemic Areas	52 53 53 53 54 55 56 57 58 65 67
Chapter 3 Malaria 3.1 Introduction 3.2 Malaria Transmission 3.3 Life Cycle of Malaria 3.4 Symptoms of Malaria 3.5 Preventing and Treating Malaria 3.6 Relationship Between Malaria and Climate 3.7 Previous Studies into Malaria (Transmission, Vector and Incidents) and Climate Variables 3.8 Malaria in Saudi Arabia Topographical Classification of Asir Region and Endemic Areas 3.9 Controlling of Malaria Transmission in Asir Region	52 53 53 53 54 55 56 57 58 65 67 70

3.11 Data and Method	73
3.11.1 Sources of Data	73
3.11.2 Malaria Data Description	75
3.11.3 Temperature Data Description	76
3.11.4 Rainfall Data Description	78
3.11.5 Relative Humidity Data Description	79
3.11.6 Statistical Analysis Data Preparation	80
3.11.7 Statistical Analysis	82
3.12 Results	84
3.12.1 Malaria with Maximum Temperature in Asir	84
3.12.2 Malaria with Maximum Temperature in TA	85
3.12.3 Malaria with Maximum Temperature in TQ	86
3.12.4 Malaria with Mean Temperature in Asir	87
3.12.5 Malaria with Mean Temperature in TQ	88
3.12.6 Malaria with Minimum Temperature in Asir	89
3.12.7 Malaria with Minimum Temperature in TA	89
3.12.8 Malaria with Minimum Temperature in TQ	90
3.12.9 Malaria with Rainfall in Asir	91
3.12.10 Malaria with Rainfall in TA	91
3.12.11 Malaria with Rainfall in TQ	92
3.12.12 Malaria with Relative Humidity in Asir	93
3.12.13 Malaria with Relative Humidity in TQ	94
3.12.14 Multiple Regressions	95
A- Multiple Regressions of Malaria Cases in Over All Asir	95

B- Multiple Regressions of Malaria Cases in TA Sector	100
C- Multiple Regressions of Malaria Cases in TQ Sector	103
3.13 Discussion and Conclusion	108
Chapter 4 Leishmaniasis	112
4.1 Introduction	113
4.2 Geographic distribution	115
4.3 Leishmania Transmission and its Life Cycle	117
4.4 Previous Studies and Relationship Between Sandflies or Leishmaniasis with Climate and Weather Conditions	119
4.4.1 Sandflies with Climate and Weather	119
4.4.2 Leishmaniasis with Climate and Weather	123
4.4.3 Sandflies of Leishmaniasis with Climate and Weather in Saudi Arabia	125
4.4.4 Literature Review Summary	127
4.5 Sandfly and Leishmaniasis in Saudi Arabia	128
4.6 Leishmaniasis in Asir Region (Study Area)	131
4.7 Topographical Classification of Asir Region and Endemic Areas	132
4.8 Control of Leishmaniasis Transmission in Asir	133
4.9 Chapter Rationale and Hypotheses	135
4.10 Data and Method	135
4.10.1 Sources of Data	135
4.10.2 CL Geography	136
4.10.3 CL Data Description	139
4.10.4 Temperature Data Description	142
4.10.5 Rainfall Data Description	142
4.10.6 Relative Humidity Data Description	143

4.10.7 Statistical Analysis Data Preparation	143
4.10.8 Statistical Analysis	147
4.11 Results	148
4.11.1 CL with Maximum Temperature	148
4.11.2 CL with Mean Temperature	150
4.11.3 CL with Minimum Temperature	151
4.11.4 CL with Rainfall	153
4.11.5 CL with Relative Humidity	154
4.11.6 Multiple Regressions	155
A- Multiple Regression of CL in All Asir	155
B- Multiple Regression of CL in Highlands Group	158
C- Multiple Regression of CL in Lowlands Group	161
4.12 Discussion and Conclusion	163
4.12 Discussion and Conclusion Chapter 5 Schistosomiasis	163 168
4.12 Discussion and Conclusion Chapter 5 Schistosomiasis 5.1 Introduction	163 168 169
4.12 Discussion and Conclusion Chapter 5 Schistosomiasis 5.1 Introduction 5.2 Species of Schistosomes and Geographic Distribution	163168169170
4.12 Discussion and Conclusion Chapter 5 Schistosomiasis 5.1 Introduction 5.2 Species of Schistosomes and Geographic Distribution 5.3 Infection and Life Cycle of Schistosomiasis	163 168 169 170 171
4.12 Discussion and Conclusion Chapter 5 Schistosomiasis 5.1 Introduction 5.2 Species of Schistosomes and Geographic Distribution 5.3 Infection and Life Cycle of Schistosomiasis 5.4 Symptoms and Diagnosis of Schistosomiasis	163 168 169 170 171 172
4.12 Discussion and Conclusion Chapter 5 Schistosomiasis 5.1 Introduction 5.2 Species of Schistosomes and Geographic Distribution 5.3 Infection and Life Cycle of Schistosomiasis 5.4 Symptoms and Diagnosis of Schistosomiasis 5.5 Control of Schistosomiasis	163 168 169 170 171 172 173
4.12 Discussion and Conclusion Chapter 5 Schistosomiasis 5.1 Introduction 5.2 Species of Schistosomes and Geographic Distribution 5.3 Infection and Life Cycle of Schistosomiasis 5.4 Symptoms and Diagnosis of Schistosomiasis 5.5 Control of Schistosomiasis 5.5.1 Parasite Control (Treatment)	163 168 169 170 171 172 173
 4.12 Discussion and Conclusion Chapter 5 Schistosomiasis 5.1 Introduction 5.2 Species of Schistosomes and Geographic Distribution 5.3 Infection and Life Cycle of Schistosomiasis 5.4 Symptoms and Diagnosis of Schistosomiasis 5.5 Control of Schistosomiasis 5.5.1 Parasite Control (Treatment) 5.5.2 Control of Transmission 	163 168 169 170 171 172 173 174
4.12 Discussion and Conclusion Chapter 5 Schistosomiasis 5.1 Introduction 5.2 Species of Schistosomes and Geographic Distribution 5.3 Infection and Life Cycle of Schistosomiasis 5.4 Symptoms and Diagnosis of Schistosomiasis 5.5 Control of Schistosomiasis 5.5.1 Parasite Control (Treatment) 5.5.2 Control of Transmission 5.5.3 Health Education	163 168 169 170 171 172 173 174 175
4.12 Discussion and Conclusion Chapter 5 Schistosomiasis 5.1 Introduction 5.2 Species of Schistosomes and Geographic Distribution 5.3 Infection and Life Cycle of Schistosomiasis 5.4 Symptoms and Diagnosis of Schistosomiasis 5.5 Control of Schistosomiasis 5.5.1 Parasite Control (Treatment) 5.5.2 Control of Transmission 5.5.3 Health Education 5.6 Seasonal Influences on the Transmission of Schistosomiasis	163 168 169 170 171 172 173 173 174 175 176

5.7.1 The Influence of Weather on the Snails Hosts	177
5.7.2 The Influence of Weather on the Parasite of Schistosoma	179
5.7.3 The influence of Weather upon Schistosomiasis Illness	179
5.7.4 Studies in Saudi Arabia	180
5.7.5 Literature Review Summary	181
5.8 Schistosomiasis in Saudi Arabia	183
5.9 Schistosomiasis in Asir Region (Study Area)	185
5.10 Topographical Classification of Asir Region and Endemic Areas	186
5.11 Rationale and Hypotheses	187
5.12 Data and Method	187
5.12.1 Sources of Data	187
5.12.2 Classification of Data	188
5.12.3 Schistosomiasis Data Description	192
5.12.4 Snail Data Description	196
5.12.5 Temperature Data Description	197
5.12.6 Rainfall Data Description	198
5.12.7 Relative Humidity Data Description	199
5.12.8 Statistical Analysis Data Preparation	199
5.12.9 Statistical Analysis	205
5.13 Result	207
5.13.1 Schistosomiasis with Temperature Data	207
5.13.2 Schistosomiasis with Rainfall Data	211
5.13.3 Schistosomiasis with Relative Humidity Data	213
5.13.4 Snail Numbers with Weather Data	214

5.15.5 Schistosomiasis Cases with Snail Numbers	216
5.13.6 Multiple Regressions	217
A- Multiple Regressions of Schistosomiasis Cases in All Asir	217
B- Multiple Regressions of Schistosomiasis Cases in the Highlands	219
5.14 Discussion and Conclusion	221
Chapter 6 Conclusion	225
6.1 Introduction	226
6.2 Development of the Study	226
6.3 Summaries and Discussions	227
6.3.1 Infectious Diseases	227
6.3.2 Study Area	228
6.3.3 Malaria	228
6.3.4 CL Leishmaniasis	228
6.3.5 Schistosomiasis	229
6.4 Conclusion of the Results of the Three Illnesses	230
6.5 The Importance of Climate Change in this Study	232
6.6 Study Limitations and Future Work	233
References	235
Appendix	253
Appendix A. Malaria. Appendix A.1	254
Appendix A.2	281
Appendix A.3	307
Appendix B. CL	310
Appendix C. Schistosomiasis	319

List of Tables

Content	Page No.
Table 2.1: Comparison of demographic indicators among selectedneighbouring countries of Saudi Arabia during 2004	35
Table 2.2: Demographic and health economic indicators of Saudi Arabia in between 2007 and 2008	36
Table 2.3: Key indicators of Saudi Arabia	36
Table 2.4 Literacy rate of each region of Saudi Arabia with unemployed rate and traditional houses during 2007.	37
Table 2.5 The Health resources rate for Saudi Arabia during 2008	39
Table 2.6 presents the reported cases of notifiable infectious diseases in overall the country during 2008	41
Table 2.7: The population distribution of the emirate quarter and governorates of Asir region	46
Table 2.8: Reported cases of notifiable infectious diseases in Asir during 2008	49
Table 2.9: Comparison of some health infrastructures and selected infectious diseases among some sectors of Asir Region 2004	50
Table 3.1: Minimum, optimum and maximum temperature for anopheles mosquito	58
Table 3.2: Summary of the main studies relating to malaria	64
Table 3.3; Seasonal spray of houses in Asir Region	72
Table 3.4: Descriptive statistics for Ln malaria of overall Asir	81
Table 3.5: Descriptive statistics of Ln rainfall in Khamise	81
Table 3.6: DT Ln malaria and maximum temperature in Asir (Whole Year)	84
Table 3.7: DT Ln malaria and maximum temperature in Asir (April to Aug)	85
Table 3.8: DT Ln malaria and maximum temperature in TA (Whole Year)	85
Table 3.9: DT Ln malaria and maximum temperature in TA (April to Aug)	86
Table 3.10: DT Ln malaria and maximum temperature in TQ (whole year)	86

Table 3.11: DT Ln malaria and maximum temperature in TQ (April to September)	87
Table 3.12: DT Ln malaria and mean temperature in Asir (April to August)	87
Table 3.13: DT Ln malaria and mean temperature in TQ (April to September)	88
Table 3.14: DT Ln malaria and minimum temperature in Asir (April to August)	89
Table 3.15: DT Ln malaria and minimum temperature in TA (April to August)	89
Table 3.16: DT Ln malaria and minimum temperature in TQ (April to September)	90
Table 3.17: DT Ln malaria and Ln rainfall in Asir (April to August)	91
Table 3.18: DT Ln malaria and with Ln rainfall in TA (Whole year)	91
Table 3.19: DT Ln malaria and with Ln rainfall in TA (April to August)	92
Table 3.20: DT Ln malaria and Ln Rainfall in TQ (whole year)	92
Table 3.21: DT Ln malaria and Rainfall in TQ (April to September)	93
Table 3.22: DT Ln malaria and relative humidity in Asir (Whole Year)	93
Table 3.23: DT Ln malaria and relative humidity in Asir (April to August)	94
Table 3.24: DT Ln malaria and relative humidity in TQ (whole year)	94
Table 3.25: DT Ln malaria and relative humidity in TQ (April to September)	95
Table 3.26: Regressions of DT Ln malaria cases of Asir for maximum temperature, rainfall and relative humidity for whole year	96
Table 3.27: Regressions of DT Ln malaria cases of Asir with multiple weather (maximum temperature with rainfall and maximum temperature with relative humidity and for all of them together) for whole year	97
Table 3.28: Regressions of DT Ln malaria cases of Asir for maximum temperature, rainfall and relative humidity for April to August	98
Table 3.29: Regressions of DT Ln malaria cases of Asir with multiple weather (maximum temperature with rainfall and maximum temperature with relative humidity and for all of them together) for April to August	99

Table 3.30: Regressions of DT Ln Malaria cases of ta for maximum temperature and rainfall for whole year	101
Table 3.31: Regressions of DT Ln malaria cases of TA with multiple weather (maximum temperature and rainfall) for whole year	101
Table 3.32: Regressions of DT Ln malaria cases of TA for maximum temperature and rainfall for April to August	102
Table 3.33: Regressions of DT Ln malaria cases of TA with multiple weather (maximum temperature and rainfall) for April to August	102
Table 3.34: Regressions of DT Ln malaria cases of TQ for maximum temperature, rainfall and relative humidity for whole year	104
Table 3.35: Regressions of DT Ln malaria cases of TQ with multiple weather (maximum temperature with rainfall and maximum temperature with relative humidity and for all of them together) for whole year	105
Table 3.36: Regressions of DT Ln malaria Cases of TQ for maximum temperature, rainfall and relative humidity for April to September	106
Table 3.37: Regressions of DT Ln malaria cases of TQ with multiple weather (maximum temperature with rainfall and maximum temperature with relative humidity and for all of them together) for April to September	107
Table 4.1: Summary of the main studies relating to Leishmaniasis	127
Table 4.2 Number of visceral leishmaniasis cases in all Asir Region (1996-2007)	131
Table 4.3: Descriptive Statistics for Ln. CL Cases in Whole Air Region	144
Table 4.4: Descriptive Statistics for Ln. CL Cases in Highlands Group	144
Table 4.5: Descriptive Statistics for Ln. CL Cases in Lowlands Group	145
Table 4.6: Descriptive Statistics of Ln Rainfall in Khamise	145
Table 4.7: DT Ln CL and maximum temperature in Asir during June to	149
Table 4.8: DT Ln CL and maximum temperature in the Highlands during	149
June to September	
Table 4.9: DT Ln CL and Maximum temperature in the Lowlands during March to November	150
Table 4.10: DT Ln CL and mean temperature in Asir during June to September	150
Table 4.11: DT Ln CL and mean temperature in the highlands during June	151
to September	
June to September	151

Table 4.13: DT Ln CL and Minimum Temperature in the Highlands	152
Table 4.14: DT Ln CL and Minimum Temperature in the Lowlands during March to November	152
Table 4.15: DT Ln CL and Rainfall in All Asir during June to September	153
Table 4.16: DT Ln CL and Rainfall in the Highlands during June to September	153
Table 4.17: DT Ln CL and Rainfall in the Lowlands during March to November	154
Table 4.18: DT Ln CL and Relative Humidity in All Asir during June to September	154
Table 4.19: DT Ln CL and Relative Humidity in the highlands during June to September	155
Table 4.20: Regressions of DT Ln CL cases of Asir for maximum temperature, rainfall and relative humidity for June to September	156
Table 4.21: Regressions of DT Ln CL cases of Asir with multiple weather (maximum temperature with rainfall and maximum temperature with relative humidity and for all of them together) during June to September for average of 2 to 4 previous months	157
Table 4.22: Regressions of DT Ln CL cases of the highlands for maximum temperature, rainfall and relative humidity during June to September, for average of 2 to 4 previous months	159
Table 4.23 : Regressions of DT Ln CL cases of the highlands with multiple weather (maximum temperature with rainfall and maximum temperature with relative humidity and for all of them together) during June to September for average of 2 to 4 previous months	160
Table 4.24: Regressions of DT Ln CL cases of the lowlands for maximum temperature and rainfall during March to November for average of 1 to 3 previous months	162
Table 4.25: Regressions of DT Ln CL cases of the lowlands with multiple weather (maximum temperature and rainfall) during March to November for average of 1 to 3 previous months	162
Table 5.1: Summary of the main studies relating to schistosomiasis	182
Table 5.2: Descriptive Statistics for Ln. Schist Cases in Asir Region	201
Table 5.3: Descriptive Statistics for Ln. Schist Cases in Highlands	201
Table 5.4: Descriptive Statistics for Ln. Schist Cases in Lowlands	202
Table 5.5: Descriptive Statistics for Ln. Schist Cases in Sabt-Alalaya	202
Table 5.6: Descriptive Statistics for Ln. Snail in Asir	203
Table 5.7: DT Ln Schist and Maximum Temperature in all Asir (June to Sep)	208

Table 5.8: DT Ln Schist and mean temperature in all Asir (June to Sep)	209
Table 5.9: DT Ln Schist and minimum temperature in all Asir (June to Sep)	210
Table 5.10: DT Ln Schist and Ln Rainfall in all Asir (Feb to May)	211
Table 5.11: DT Ln Schist and Ln Rainfall in all Asir (June to Sep)	212
Table 5.12: DT Ln Schist and Ln Rainfall in Highland (Feb to July)	213
Table 5.13: DT Ln Snails and Maximum Temperature in all Asir Region (Dec to April)	214
Table 5.14: DT Ln Snails and Maximum Temperature in all Asir Region(May to Nov)	215
Table 5.15: DT Ln snails and Ln rainfall in all Asir Region (Dec to April)	216
Table 5.16: DT Ln Schist and Ln Snails in all Asir Region	217
Table 5.17: Regressions of the coefficient of DT Ln schistosomiasis cases of whole Asir for averages of 3 and 4 previous months of maximum temperature and rainfall during June to September	218
Table 5.18: Regressions of the coefficient of DT Ln schistosomiasis cases of whole Asir with multiple weather averages of 3 and 4 previous months of maximum temperature and rainfall during June to September	218
Table 5.19: Regressions of the coefficient of DT Ln schistosomiasis cases of highlands group for averages of 6 and 7 previous months of maximum temperature and rainfall during February to July	220
Table 5.20: Regressions of the coefficient of DT Ln schistosomiasis cases of the highlands group with multiple weather averages of 6 and 7 previous months of maximum temperature and rainfall during February to July	220
Table 6.1: Summary of the significant relationships between the climate variables and selected infectious diseases in Asir (Malaria, CL and Schistosomiasis)	231

List of Figures and Maps

Content				
Figure 1.1: Four main transmission cycle for infectious diseases.	23			
Figure 1.2: The links between climate change and vector & water-borne diseases incidence.	27			
Figure 1.3: Impact of climate change with some other factors upon health	28			
Figure 2.1: Map of Saudi Arabia with its administrative regions.	34			
Figure 2.2: Population distribution by regions by demographic survey 2007.	35			
Figure 2.3 Projects locations of water desalinated & distribution locations in SA.	38			
Figure 2.4: The different sources of households of water supply in SA	38			
Figure 2.5: Reporting System for infectious diseases in MOH, Saudi Arabia.	40			
Figure 2.6: Malaria distribution in Saudi Arabia and prevalence of the vector	42			
Figure 2.7: distribution of endemicity of schistosomiasis in Saudi Arabia during 2009.	44			
Figure 2.8: The schistosomiasis endemic regions in Saudi Arabia during 2009.	44			
Figure 2.9: Map of Asir Region with its emirate quarter and governorates.	45			
Figure 2.10: Topography of Asir Region.	46			
Figure: 3.1: Schema of the life cycle of malaria	55			
Figure 3.2: Number of malaria cases in Saudi Arabia (1980-2009)	67			
Figure 3.3: Endemic Sectors in Asir Region. Adapted from	68			
Figure 3.4: Topographic map for Asir Region including the weather stations	70			
Figure 3.5: Spraying the houses (Indoor) with insecticides that have residual effect in endemic area of Asir Region.	71			
Figure 3.6: Removal of breeding habitats and unnecessary water body beside the valleys	71			
Figure 3.7: Monthly averages of malaria cases in Asir Region (1995 - 2006)	75			

Figure 3.8: Monthly averages of malaria cases in TA and TQ (1995 - 2006)	76			
Figure 3.9: Maximum, mean and minimum averages temperature in Khamise City 1995 -2006				
Figure 3.10: Maximum and minimum averages temperature in Majaredah 1995 -2006				
Figure 3.11: Monthly averages rainfall in Khamise City (1995 -2006)	78			
Figure 3.12: Monthly averages rainfall in Rejal-Alma and Sarat-Abida 1995 -2006				
Figure 3.13: Monthly averages relative humidity in Khamise 1995 – 2006	79			
Figure 3.14: Frequency of natural logarithm of malaria (Ln.M) in overall Asir Region (1995 – 2006)	80			
Figure 3.15: Frequency of logarithm rainfall (Ln. R) in Khamise (1995 – 2006)	81			
Figure 3.16: Yearly Ln malaria in Asir Region (1995 - 2006)	82			
Figure 3.17: Yearly Ln malaria in TA and TQ sectors (1995 - 2006)	82			
Figure 4.1: Phlebotomine Sandfly	114			
Figure 4.2: Clinical Presentations of Leishmaniasis	115			
Figure 4.3: Distribution of CL, MCL and VL in the World	116			
Figure 4.4: Life Cycle of Leishmaniasis	118			
Figure 4.5 Monthly Averages of CL Cases in Saudi Arabia (2002-2006)	129			
Figure 4.6: The Number of (CL) Cases in Saudi Arabia (1983-2009)	130			
Figure 4.7: The Relationship between Altitude and Density of Some Species of Sandfly in Saudi Arabia	131			
Figure 4.8: The correlation between density of sandfly and the number of CL cases in Al-Dereva City, Saudi Arabia				
Figure 4.9: Topographic map of Asir Region including weather stations	133			
Figure 4.10: Annual rate and Monthly Average of CL Cases in Sectors of Asir Region (1996-2007)	138			
Figure 4.11: Monthly Averages of CL Cases in All Asir Region (1996 - 2007)	140			
Figure 4.12: Monthly averages of CL cases in the first group (Highlands) of Asir Region (1996 - 2007)				
Figure 4.13: Monthly averages of CL cases in the second group (Lowlands) of Asir Region (1996 - 2007)				
Figure 4.14: Frequency of Natural Logarithm of CL (Ln. CL) in All Asir Region (1996 – 2007)				

Figure 4.15: Frequency of Natural Logarithm of CL (Ln. CL) in Highlands Group (1996 – 2007)				
Figure 4.16: Frequency of Natural Logarithm of CL (Ln. CL) in Lowlands Group (1996 – 2007)				
Figure 4.17: Frequency of Natural Logarithm Rainfall (Ln R) in Khamise (1996 – 2007)				
Figure 4.18: Monthly CL Cases in all Asir Region (1996 – 2007)				
Figure 4.19: Monthly CL cases in highlands group in Asir Region (1996 – 2007)				
Figure 4.20: Monthly CL cases in lowlands group in Asir Region (1996 – 2007)	147			
Figure 5.1: Different single parasites of Schistosoma	170			
Figure 5.2: Single egg of Schistosoma	170			
Figure 5.3: Schistosoma snail host	170			
Figure 5.4: Distribution of Schistosomiasis	171			
Figure 5.5: Life Cycle of Schistosomiasis	172			
Figure 5.6: The enlargement of the liver and the spleen	173			
Figure 5.7: Spraying of molluscicides in locations where snails breed in Asir, Saudi Arabia				
Figure 5.8: Removing vegetation, filling rain pools and removing snails in Asir	175			
Figure 5.9: The Number of (schist) Cases in Saudi Arabia (1980- 2009)	184			
Figure 5.10: Comparison of Schistosomiasis among regions of Saudi Arabia 2005	185			
Figure 5.11: Monthly Averages of schistosomiasis Cases in sectors of Asir Region (1998 - 2009)	191			
Figure 5.12: Topographic map for Asir Region including the weather stations	192			
Figure 5.13: Monthly Averages of schistosomiasis Cases in All Asir Region (1998 - 2009)	193			
Figure 5.14: Monthly Averages of schistosomiasis Cases of Highlands Group (1998 - 2009)	194			
Figure 5.15: Monthly Averages of Schistosomiasis Cases in Lowlands Group (1998 - 2009)	195			
Figure 5.16: Monthly Averages of Schistosomiasis Cases in Sabt-Alalaya, Asir Region (1998 - 2009)				
Figure 5.17: Monthly averages of snail numbers in all Asir Region (1998 - 2009)				
Figure 5.18: Maximum and Minimum Temperature in Sabt-Alalaya (1998 -2009)				
Figure 5.19: Monthly Averages rainfall in Sabt-Alalaya Sector (1998 - 2009)				

Figure 5.20: Frequency of Natural Logarithm of Schist (Ln. Schist) in Asir				
Region (1998 – 2009)				
Figure 5.21: Frequency of Natural Logarithm of Schist (Ln. Schist) in	201			
Highlands (1998 – 2009)	-01			
Figure 5.22: Frequency of Natural Logarithm of Schist (Ln. Schist) in	202			
Lowlands (1998 – 2009)	202			
Figure 5.23: Frequency of Natural Logarithm of Schist (Ln. Schist) in	202			
Sabt-Alalaya (1998 – 2009)	202			
Figure 5.24: Frequency of Natural Logarithm of Snail (Ln. Snail) in Asir	203			
(1998 – 2009)	205			
Figure 5.25: Monthly Schistosomiasis Cases in all Asir Region (1998 -	203			
2009)	205			
Figure 5.26: Monthly Schistosomiasis Cases in Highlands Group (1998 -	204			
2009)				
Figure 5.27: Monthly Schistosomiasis Cases in Lowlands Group (1998 -	204			
2009)	201			
Figure 5.28: Monthly Schistosomiasis Cases in Sabt-Alalaya Sector (1998	205			
- 2009)	205			
Figure 5.29: Monthly Snail numbers in the whole of Asir Region (1998 –	205			
2009)	205			

List of Abbreviations

CBD	The Convention of Biological Diversity
CDC	Centres for Diseases Control and Prevention
CDSI	Central Department of Statistic & Information Saudi Arabia,
CFSPH	The Centre for Food Security and Public Health
CL	Cutaneous Leishmaniasis
CPRD	Costal Plain Red Sea
ECDC	European Centre for Diseases Prevention and Control
El Nino	If the change of temperature over the Pacific Ocean is positive
ENSO	El Nino southern oscillation
HIV/AIDS	Human Immunodeficiency virus/ Acquired Immune Deficiency Syndrome
IPCC	Intergovernmental Panel on Climate Change
IUCN	International Union for Conservation of Nature
IWGCCH	Interagency Working Group on Climate Change and Health
La Nina	If the temperature change over the Pacific Ocean is negative
MCL	Mucocutaneous Leishmaniasis
MEPS	Meteorological and Environmental Protection Section
МОН	Ministry of Health
MOWE	Ministry of Water and Electricity

NIAID	National Institute of Allergy and Infection Diseases
NIEHS	The National Institute of Environmental Health Sciences
NWS	National Weather Service
Schist	The Schistosomiasis diseases
SPSS	Statistical Package for the Social Sciences
ТА	Tehama of Asir, Saudi Arabia
TQ	Tehama of Qahtan, Saudi Arabia
UNFCCC	United Nations Framework Convention on Climate Change
VCA	The Vector Control Administration, in Health Affairs Directorate in Asir Region.
VL	Visceral Leishmaniasis
WHO	World Health Organisation
WSH	Water Sanitation and Health

Acknowledgements

This PhD research would not have been possible without the help, guidance and support of almighty Allah, so I praise and sincerely thank Him for this achievement.

Firstly, I would like to express my deepest appreciation and sincere gratitude to my supervisor Dr Iain Lake for his copious knowledge, continuous support, patience, kindness, insightful comments and his encouragement throughout my studies. My sincere thanks go to the rest of my thesis committee, Prof. Graham Bentham and Dr Steve Dorling for their advice, useful comments and thought provoking discussions.

Acknowledgements to Health Sciences College in Riyadh, King Saud University, for giving me this opportunity to complete my postgraduate study, and many thanks to all of my colleagues in this college for their help and co-operation. I would also like to thank the many individuals in Saudi Arabia, who during my fieldwork provided me with the opportunity to learn from their rich experience, particularly Dr Abdulmehsen Abdoon, who spent a lot of time with me and gave me much useful information regarding Malaria, Leishmaniasis and Schistosomiasis diseases in Asir, and to Mr. Ali Alshahrani, Mr. Ahmed Thabet and Mr. Mohamed Ali Faraj, from the Vector Control Administration in Asir for their help and co-operation. These thanks extend to Dr Mohamed Zahrani and Dr Adnan Aldaham from the Infectious Diseases Department in MOH, and to all of those who supported me in any respect during the completion of this project.

I can hardly find the words to express my gratitude and thanks to my beloved Father and Mother who I consider as the highest valued greatest humans in the world. I would like to extend this gratitude to my brothers and sister, particularly my brother Ahmed. I will never forget all his essential assistance. These thanks also go to all of my relatives and friends in Saudi Arabia, who have encouraged and supported me from time to time.

It is a great pleasure to thank everyone who has helped me and my family in Norwich, specifically Dr Mansour Alghamdi, who has given me considerable assistance from my first day in this country, and this thanks goes as well to Dr Mansour Almazroui, Dr Turki Habeeballah, Dr Majed Alshamari, Dr Rady Othman, Ahmed Abu-Amr Alghamdi, Mubarak Aldosary and Hamad Aldosary and others for their friendship, help and advice.

Lastly, I offer my sincere regards and love to my small family, my dear wife for her continued support, encouragement and patience, and to my lovely children, Faisal, Ahmed and Abdulaziz for their love, patience and help.

Chapter 1 <u>Introduction and Overview</u>

Background

1.1 What are Infectious Diseases?

Infectious diseases, also known as communicable diseases, are diseases which can be passed from one person to another. It is also possible for such diseases to spread indirectly from animals to people (zoonotic diseases) or through unhygienic conditions (Patz et al., 2002; Ryan et al., 2010). These diseases may cause damage or injury to the host due to the presence of pathogenic microbial agents such as viruses, bacteria, fungi, protozoa and parasites (Ryan et al., 2010). Examples of these diseases include herpes, malaria, mumps, HIV/AIDS, influenza, chicken pox, leishmaniasis and schistosomiasis (CDC, 2010). Vector-borne diseases are type of infectious diseases which are transmitted from one person to another by an insect or other arthropod vectors that carry the pathogens. The vector is an agent such as a micro organism, animal or person which carries and transmits an infectious agent. Common vectors include domestic animals, mammals or arthropods which assist in transmitting the parasitic organisms to humans (Schmidt and Roberts, 1996). An example of a vector-borne disease is malaria where the mosquito acts as a vector. Water-borne diseases are those diseases where the micro-organisms spend part of their lifecycle in water. An example is schistosomiasis which is transmitted by worms through contact with contaminated water (WHO, 2010).

There are different classifications of infectious diseases. The transmission mode of these diseases can either be anthroponoses or zoonoses according to the pathogen natural reservoir. Anthroponoses are where the reservoir is human, whereas zoonoses occur when the reservoir is animal. Transmission can also be direct or indirect. Direct transmission is where transmission occurs directly from person to person or from animal to animal. In the latter case the animal then gives the illness to a person. Indirect transmission is where human to human or animal to animal transmission is dependent upon a vector (Patz et al., 2002; Ryan et al., 2010). Figure 1.1 shows these different types of transmission cycle for infectious diseases.

Infectious diseases are transmitted by a variety of different mechanisms including (Schmidt and Roberts, 1996; Ryan et al., 2010; ECDC, 2010):

- 1. Ingesting contaminated food or water. These are known as gastrointestinal diseases (e.g. salmonellosis, cholera).
- 2. Contact with bodily fluids, often through sexual activity blood transfusions and sharing needles. (e.g. HIV/ADIS, hepatitis B).



Figure 1.1: Four main transmission cycles for infectious diseases. Adapted from Patz et al., (2002)

- 3. Aerosolized droplets spread by sneezing, coughing, talking and kissing (e.g. flu, meningitis).
- 4. Contact with a contaminated inanimate object such as clothes which can pass disease from one person to another (e.g. scabies).
- 5. Transmission of infectious diseases by mechanical vector. An example would be houseflies that pick up an infectious agent on the outside of its body and transmits it to an individual in a passive method (e.g. Typhoid, anthrax).
- 6. Transmission of infectious diseases by biological vectors such as mosquitoes and sandflies that deliver pathogens from one host to another (e.g. dengue fever and leishmaniasis).

1.2 The Global Importance of Infectious Diseases to Health

Throughout history, many types of these infectious diseases have killed millions of people. For example, an estimated 75 to 200 million people were killed by

the Black Death during the 14th century and this disease continues to kill about 3000 people each year (Philipkoski, 2008). Another example occurred in Mexico during the 15th and 16th centuries, when the introduction of smallpox, measles, and typhus led to disease pandemics which killed over 80% of the population (Dobson et al., 1996). Moreover, the first European influenza epidemic occurred during the 16th century which is estimated to have killed 20% of the population. Tuberculosis killed an estimated quarter of the adult population of Europe during the 19th century. In 1918 the influenza pandemic killed up to 50 million people, and today this disease is responsible for about 0.5 million deaths worldwide each year (The Navy Department Library, 2005; WHO, 2010).

Infectious diseases still represent a major health problem today and the World Health Organization (2002) has estimated that currently there are more than 14.8 million deaths per year globally due to infectious diseases. This represents 25.9% of all deaths over the world (The World Health Report, 2004). Malaria is the most important vector-borne disease, and WHO estimates that 2.4 billion people live in malarious regions of the world (WHO/SDE/PHE, 1999). The exact number of deaths caused by malaria is unknown but it has been estimated at 1.5 - 3 million annually (De Savigny et al., 2004).

1.3 Climate Variables and Climate Change

Weather has been defined as "the short term changes in temperature, rainfall, wind speed, and some other atmospheric conditions" (WHO, 2003). Weather events usually occur on a daily time scale and these variables are associated with many health impacts (Harold et al., 2000). Climate is the total experience of weather over a longer time scale (conventionally 30 years) (IPCC, 2007). Atmospheric variables such as temperature, rainfall, wind, relative humidity and cloudiness are described as climate variables (McMichael et al., 2003). Climate variability occurs on many time scales, it usually occurs on a daily time scale and these variables are associated with many impacts on health. Climate variability occurs at other scales such as seasonal variability and also over longer periods. One important example of climate variability over longer time scales is the El Nino southern oscillation (ENSO). It is a global climate phenomenon, which affects temperature (a warming or cooling of at least 0.5 °C) over the tropical Pacific Ocean. Time periods are known as "El Nino" if the

change of temperature over the Pacific Ocean is positive. If the temperature change is negative it is known as "La Nina". These events occur every 3 to 5 years on average and lead to changing of the weather patterns across much of the planet (Lipp et al., 2002: NWS, 2005). El Nino creates an increase in rainfall across the east-central and eastern Pacific Ocean. It also causes extreme weather conditions such as floods and droughts in many regions of the world.

Human activities that burn fossil fuels such as oil increase the amount of greenhouse gases in the atmosphere. Although these gases also occur naturally, their recent increase may contribute to climate change (IPCC, 2007). During the 20th century the global surface temperature increased 0.74 ± 0.18 °C (IPCC, 2007). Scientists predict that climate change will cause a mass extinction of several species of plants and animals, lead to the flooding of low-lying areas, and increase the frequency of extreme weather events (IUCN, 2009). The United Nations Framework Convention on Climate Change declared that climate change "contributes to the global burden of disease and premature death" (UNFCCC, 2007). There are four climatic zones in Asia; boreal, arid and semi-arid, tropical and temperate (UNFCCC, 2007). Martens et al., (1999) state that the principal impacts of climate change on health in Asia will be upon epidemics of vector-borne diseases.

1.4 The Influence of Climate Variables upon Infectious Diseases

The vectors of vector-borne diseases (a part of infectious diseases) such as mosquitoes and sandflies are sensitive to climate factors specifically temperature, rainfall and relative humidity (Hana et al., 2001; Cross et al., 1996). Therefore the weather will have an influence upon the transmission dynamics of these diseases. This will affect their spatial and temporal patterns. For example elevated temperatures in an area could increase the occurrence of vector-borne diseases such as malaria. Also Pampana, 1969 noted that the absolute temperature and its duration can affect the life cycle of malaria, but that this varies by parasite species.

There are many other examples. Air-borne infectious diseases can be affected by climate variables, although little is known about these types of diseases (Lipp et al., 2002). An example is influenza where Langford et al., (1995) found that the peak number of cases of flu and cold occur during winter in the Northern Hemisphere indicating that the transmission of the viruses is more active during winter season, thus increasing the cases. Food-borne diseases are also affected by climate variables. Lake et al., (2009) examined the correlation between temperature and food-borne infectious diseases. There were positive correlations (P<0.01) with each temperature lagged at the current week to 5 weeks. These were attributed to increased bacterial reproduction during warm weather as well as to changes in human behaviour (e.g. the popularity of barbecues during warm weather). Climate (e.g. heavy rainfall, flooding and increased temperature) is also recognized to be an important factor influencing the epidemiology of water borne disease (Hunter, 2003). Therefore water-borne diseases may increase due to changes in environmental conditions. Outbreaks in water-borne disease have been positively associated to extreme rainfall in the USA between 1948 and 1994 (Curriero, 2001).

There have been many previous studies which have indicated that climate variables have an impact on infectious diseases, particularly vector, air, food and water-borne diseases (Tulu, 1996; Woube, 1997; Ye et al., 2007; Alam et al., 2008; Lake et al., 2009; Oluleye and Akinbobola, 2010). Some have shown that the weather influences the abundance of vectors (Cross and Hyams, 1996; Amerasinghe et al., 1999; Abdoon and Alshahrani, 2003; Boussaa et al., 2005; Kasap et al., 2009) while others have found that vector abundance had an effect on the incidence of these type of diseases (Al-Amru, 2002; Al-Tawfiq et al., 2004; Al-Ibrahim et al., 2005).

1.5 Climate Change and its Potential Impact Health

Climate changes can affect human health directly as well as indirectly by changing ecological and biological processes (WHO et al., 2003; Cowie, 2007; Habib, 2011). These changes can cause death, injuries, illness, suffering and increases in the risk of infection diseases. In addition there may be adverse effects on the quality and quantity of food and water (Confalonieri et al., 2007). Postigo, (2007) has estimated that there are 150,000 deaths per year resulting from climate change and this number may increase to 300,000 by 2030. Recently, heavy rainfall events occurred in Jeddah, Saudi Arabia, in November 2009 and also in January 2011 which caused loss of life for hundreds of people and much damage to the infrastructure (Almazroui, 2001). These may have been associated with climate change.

The previous section has show that weather can affect the incidence of infectious disease. Therefore any change in climate may impact upon infectious diseases. These changes can influence the transmission patterns of many infectious diseases (Lindgren et al., 2010). In addition to the importance of climate change, other factors may influence the incidence of infectious diseases in the future. Examples include environmental influences such as agricultural development, deforestation and urbanization (Grasso et al., 2010). There are also socio demographic factors such as transportation, nutrition and human migration which may all have an impact (WHO, 2006).

A report in the United States, published by Environmental Health Perspectives and the National Institute of Environmental Health Sciences (NIEHS), has linked climate change to nine broad human health categories including infectious diseases such as vector-borne diseases and water-borne diseases (IWGCCH, 2010). The link between climate change and vector or water-borne disease incidence is presented in Figure 1.2. This figure shows that many factors, both direct and indirect, can influence water-borne diseases and vector-borne diseases, starting with climate change but noting other effects such as human behaviour, activities, host pathogen transmission, vector and vector habitat.



Figure 1.2: The links between climate change and vector & water-borne diseases incidence. Adapted from ECDC, 2010

Figure 1.3 shows another example of the impact of climate change with other factors upon health and presence of some infectious diseases.



Figure1.3: Impact of climate change with some other factors upon health. Adapted from (MOH, Climate change, 2006)

WHO, (2011) has listed a number of facts about climate change including some significant observations relating to health such as the changing patterns of infectious diseases in extreme high temperature. Sea levels can rise as a consequence of global warming and resultant flooding will increase the risks of infection from water and vector borne diseases. Shortages of fresh water or water of poor quality can occur in areas where rainfall amounts vary and this affects hygiene and health. This occurs especially where water has to be carried long distances as this increases the likelihood of contamination. Life threatening diseases transmitted through water via vectors such as mosquitoes are sensitive to climatic conditions as increased temperatures provide optimum conditions for mosquito breeding. As global temperatures rise, new geographical areas may provide these conditions. In the highlands of Kenya, for example, many people died of malaria in the summer of 1997. This was blamed on climate change as the population there had not previously been exposed to this disease. There are many potential consequences of climate change and some examples and their impacts will be presented. Drought may affect the transmission of some mosquito-borne diseases. During droughts in an endemic area, the activity of mosquito is usually reduced. This may reduce case incidence but also the immunity of the population. When this drought breaks, increases in mosquito abundance may occur. This leads to a large increase in cases because of more vectors and weakened immunity in the population (Woodruff et al., 2002). Another example is heat waves which are related to increases in mortality in different areas across the world. In India several heat waves were reported during the last decade of the last century. These caused thousands of deaths (Mohanty et al., 2003). Furthermore, flood is a frequent disaster. This event is a result of interaction of rainfall, wind, sea level, surface runoff and natural topography (Confalonieri et al., 2007). In Venezuela more than 30,000 people lost their lives due to storms and floods during 1999 (Guha Sapir et al, 2004). More than 130 million people in China were affected by floods during 2003.

There is already evidence that climate change is affecting disease. Increases in malaria and leishmaniasis transmission and infectious diseases in general have been reported in Bolivia during the end of the last century. There was no responding from the related governmental agencies to adapt these increases (Confalonieri et al., 2007). However in Canada when increases in heat wave related to the number of deaths, due to the increases in air pollution, vector-borne diseases and allergic disorders, the relevant private and governmental agencies have tried to adapt. A number of different steps have been taken including monitoring for the emergence of the infectious diseases, measuring to reduce the heat island effect, upgrading water and wastewater treatment facilities, preparing plans for emergency management and early warning systems (Confalonieri et al., 2007; Riedel, 2004).

The effects of climate change on human health as well as the effects on the environment have been widely discussed. There is an almost universal scientific consensus on the many direct effects of climate change that will occur in the coming decades. There has been much research in these areas but the impact of climate changes on health and specifically on infectious diseases is an area where more research is needed.

1.6 Influence of Climate Variables upon Selected Infectious Diseases in the Asir Region, Saudi Arabia.

1.6.1 Overview

This study investigates the influence of climate variables on three related vector and water -borne diseases in the Asir region, Saudi Arabia. The diseases that have been chosen for the study are malaria, cutaneous leishmaniasis and schistosomiasis. These diseases are the most wide spread vector and water-borne diseases which have the highest impact on public health in Asir (Dr A. Abdoon 2007 pers. Comm.).

Each disease has been considered separately and has a complete chapter devoted to it. Asir is an interesting study area because it has a variety of different geographical regions from the low plains of the sea coast to mountainous areas over 3000m. Because of this, it has been possible to study three diseases in different climatic conditions within a single area in Saudi Arabia. The extent of each disease in a particular sector of the Asir Region is discussed in the individual chapters on each disease.

The hypothesis for this study is that climate variables, specifically temperature, rainfall and humidity, have an influence on these diseases within Asir Region.

1.6.2 Importance of the study

This study is important for three main reasons. First, it is innovative research as it is the first time that there has been a specific and rigorous investigation of the influence of climate variables upon diseases in Saudi Arabia. Other Saudi Arabian studies have mentioned the influence of climate briefly but have not dealt with it in detail or analyzed its influence rigorously. Secondly, it is a subject that fully justifies research because these diseases have a large impact on the lives of people in Saudi Arabia who become infected, with high costs to public health. Prevention of a disease is better than cure and it is hoped that using the results of this study it should be possible to design an early warning system from weather patterns to predict the incidence of each selected disease. Finally as the climate changes in Saudi Arabia associated with climate change, this research could help predict the impact of any changes upon the health of the population.

1.7 Structure of Thesis

The first chapter (Introduction) will give an overview of infectious diseases and their importance to health. Climate and climate change will then be described followed by their influence upon infectious diseases. The chapter will then consider this study and its importance in detail.

The study area (second chapter) will be presented. This introduces the country of study, Saudi Arabia, including information on topography, climate, demography, education and the health of the population. The second half of this chapter will examine Asir Region in more detail. This is the region in the South West of the Country which forms the study area for this thesis.

There are three main chapters; 3, 4 and 5 which will examine the influence of weather upon malaria, leishmaniasis and schistosomiasis respectively within the Asir Region. Due to the similar methods used for studying each disease these three chapters have a similar structure. Each one will start by providing an overview of the disease globally. Its transmission will then be described in detail followed by information on the symptoms and causes of the disease. Each chapter will then show how this illness may be prevented and treated. The influence of weather upon disease is the focus of this thesis and so we will then present a literature review of the illness and previous studies that have looked at its relationships with weather. This will conclude with a brief summary of the literature and how this information will be developed into the design for each study. The study area will then be introduced focusing specifically on the prevalence of the disease. This will be followed by a description of the data sources to be used in each chapter and a description of the methods to be employed. The results of the analysis will then be presented, specifically evidence of any relationships with weather. Each of these three chapters will conclude with a section discussing the results obtained.

The last chapter (6) will present the summary and conclusions of this study. This will include a discussion of its strengths and limitation. It will also discuss how research developed over the 3 chapters. Finally it will make recommendations for future research. Chapter 2 <u>Study Area</u>

.

2. Introduction

2.1 Saudi Arabia

2.1.1 Geography and Demography:

Saudi Arabia is located in the south west of the Asian continent and is the largest country in the Middle East. It occupies four-fifths of the Arabian Peninsula covering an area of 2.25 million km². According to the system of regions issued during 1992, this country has been divided into 13 administrative regions (Figure 2.1) (Central Department of Statistic & Information (CDSI), 2010). The population in 2010 was 27.14 million (18.7 million Saudis, 8.4 million non-Saudi) and the annual population growth rate between 2004 and 2010 was 3.2% (CDSI, 2010). Figure 2.2 shows the distribution of population by regions according to a demographic survey undertaken during 2007. This figure shows that Riyadh and Makkah regions have the highest population number followed by Eastern Region then Asir Region which has the fourth highest population in Saudi Arabia. The lowest population number is in Northern Border followed by Jouf region. The survey also shows that the population of males is higher than females in all regions with the greatest difference in the regions of Riyadh, Makkah and Eastern Province.

Saudi Arabia is bordered by 7 countries, Jordan on the northwest, Iraq on the north and northeast, Kuwait, Qatar, Bahrain, and the Arab Emirates on the east, Oman on the southeast and Yemen on the south. It is also bordered by two seas the Arabian Gulf to the North East and the Red Sea on the west (Garout, 2000). The west coastal plain which lies along the Red Sea is 1,100 km long. The most significant mountains chain are the Sarawat Mountains (rising to 3000 meters in the south and gradually falling to 1000 meters in the north) located to the east of the west coastal plain. From these mountains there are many valleys which slope westward to the sea and eastward into the hills and the desert (CDSI, 2010).

As the country has a large area it can be divided into several different geographical areas including (Garout, 2000; CDCI, 2010):

• Najd: located in the middle of Saudi Arabia including some plateaus, hills and oases, e.g. Riyadh City (Capital City) and Beraidah City.

- Hejaz: spreads along the western coast, and includes many cities such as Makkah, Madinah and Jeddah.
- Asir Region: Located in the south west of Saudi Arabia, with three altitudes; Highlands (2000 – 3000m) which from part of the Sarawat Mounties range (e.g. Abha City), Tehama which lies on the coastal plain with foothills (e.g. Muhail City) and the east plateaus (e.g. Bishah City).
- Eastern Region which includes Al Hasa oases (e.g. Al Hasa City), coastal plains on the Arabian Gulf (e.g. Dammam City) and a large part of the Empty Quarter desert (Alrub Alkhali).
- Northern Region: contains some oases and part of the Nufoud desert.



Figure 2.1: Map of Saudi Arabia with its administrative regions (CDSI, 2008)



Figure 2.2: Population distribution by regions by demographic survey 2007 (CDSI, 2008)

Table 2.1 illustrates a comparison of demographic indicators among selected neighbouring countries of Saudi Arabia during 2006. This figure shows that Saudi Arabia has the largest area compared to the neighbouring countries, and has the second highest population. It also has the second lowest population density. More than 85% of people in Saudi Arabia live in urban areas.

	KSA	UAE	Qatar	Bahrain	Kuwait	Iraq	Yemen	Oman
Area (000) km ²	2250	83.6	11.58	0.72	17.82	435.1	555	309.5
Pop. (Million)	23.7	4.1	0.84	0.74	3.10	28.00	20.90	2.60
Density person/km2	10.5	49.1	72.4	1031.9	171.2	64.3	37.6	8.3
Pop. <15 (%)	32.9	19.5	22.5	27.3	21.8	43.3	45	32.4
Population +65 (%)	2.8	2.9	1.2	2.5	1.6	2.8	2.9	2.2
Urban Pop. %	85	82	100	100	100	67	26	71
Crude birth rate ‰	24.9	15.7	16.8	20.9	17.8	38	24.2	39.2
Crude death rate ‰	4	1.5	2.1	3.1	1.7	10	2.5	11.4

Table 2.1: Comparison of demographic indicators among selected neighbouring countries of Saudi Arabia during 2004 (WHO, 2007; CDSI, 2006)

Pop.: population
Tables 2.2 and 2.3 demonstrate some selected demographic and health economic indicators for Saudi Arabia between 2007 and 2010. They indicated that advances in health services, industry, education, commerce, agriculture and construction have turned the country into one of the fastest developing countries in the world (CDCI, 2010).

2007 and 2008. Adapted from (MOH, 2008; CDSI, 2008					
Estimated population	24,807,273				
Crude birth rate /1000 pop	24.1				
Pop. Growth Rate	2.23%				
Population Under 5 years	11.56%				
Population under 15 years	32.26%				
Population 15 - 64 years	64.95%				
Population from 65 & above	2.79%				
Total Fertility rate	3.04				
Life expectancy at birth	73.4				
Male	72.4				
Female	74.5				
GDP per capita	\$ 19206				
MOH Budget (of Governmental Budget)	5.6%				
MOH Expenditure per capita	\$ 271				

Table 2.2: Demographic and health economic indicators of Saudi Arabia between2007 and 2008. Adapted from (MOH, 2008; CDSI, 2008)

 Table 2.3: Key indicators of Saudi Arabia (CDSI, 2010)

Population 2010 (people)	27,136,977
Population density (person / sq km) 2010	14
GDP growth at (CDSI), 2010 constant prices 2009	0.6
The contribution of the private sector in GDP at constant prices 2009	48%
Per capita GDP at current prices in 2009 (SAR)	52,853
Export growth in 2009	-37.44%
Import growth in 2009	-9.08%
The contribution of exports to GDP at current prices in 2009	54%
Unemployment Rate (2009)	5.4%
General index for the cost of living 2009	122.4
Change in the index of cost of living (inflation) for the year 2009%	5.06
Growth rate of GDP per employed person (2008)	19.3
The proportion of the working population to population (2009 m)	32.1
Gross enrolment rate in primary education (2009 m)%	99
Rate and infant mortality (per thousand live births) 2009	14

2.1.2 Climate

The climate of Saudi Arabia is arid and semiarid with extremes of high temperature during summer. It is cool during the winter in Najd and some parts of the Eastern Region. However temperatures are more moderate in South Western and Northern regions during the summer and cooler in the winter. Frost and snow may occur in the Northern region. In contrast, the temperature during the summer season is high in most of the country (>400C). The humidity is high along the coastal areas (MEPS, 2007; CDSI, 2010). Average annual rainfall for Riyadh, Jeddah and Jazan is 100mm, 54mm and 500mm respectively (MOH, Workshop, 2006).

2.1.3 Education

Literacy rate %

Unemployed %

Traditional

houses %

79.5

4.7

51.2

82.2

3.0

19.8

The government of Saudi Arabia gives a high priority to education which is almost free and the literacy rate is growing steadily. In 2007, 86.3% of the total population was literate compared to only 40% during 1950. Table 2.4 shows the literacy rate, unemployed rate and the percentage of the population living in traditional houses for each administrative region of Saudi Arabia. Literacy rate, unemployed rate and percentage of traditional houses are considered indicators of the living standards. The region of Riyadh has the best living standards with high literacy, low unemployment and a low percentage of traditional homes. This is followed by the Eastern Region. The lowest living standards were in Jazan Region, although the highest unemployment rate was in Albahah, followed by Hail. Asir Region comes in the middle of living standards in Saudi Arabia.

	Al- Riyadh	Makkah	Al- Madinah	Al- Qaseem	Eastern Region	Asir	Tabouk
Literacy rate %	90.1	86.3	86.1	85.0	89.9	82.5	87.1
Unemployed %	1.5	2.5	3.2	3.1	1.8	4.5	2.2
Traditional houses %	12.2	31.8	31.9	28.0	17.0	40.7	30.4
Continue table 2.4							
	Hail	Northern Borders	Jazan	Najran	Al-Baha	Al- Jouf	Total

76.5

4.9

73.4

80.9

3.2

40.3

80.2

5.6

41.0

86.0

2.3

25.5

Table 2.4 Literacy rate of each region of Saudi Arabia with unemployed rate and
traditional houses during 2007(CDSI, 2010)

86.3

2.7

27.8

2.1.4 Drinking Water Services

Saudi Arabia draws its water from four main sources (Ministry of Water and Electricity (MOWE), 2005):

- Surface water, such as dams, springs and wells (<30meters depth) which is to be found mostly in the west and south-west of the country (7%).
- Ground water which is held in aquifers. Some are naturally replenished while others are non-renewable (23%).
- Desalinated seawater, a source of water production in which this country is now a world leader. There are 29 desalination plants providing drinking water to major urban and industrial centres through a network of water pipes (Figure 2.3). During 2004 desalination met around 70% of the Kingdom's drinking water requirements.
- Reclaimed wastewater (<1%).

In terms of how this water is distributed to households, Figure 2.4 shows the different sources of water supply in Saudi Arabia during 2004. The figure shows that most of the households (>70%) use public piped water followed by those who use catchment tanks (~24%). Less than 5% use wells and less than 1% use other sources (MOWE, 2005).



Figure 2.3 Projects locations of water desalinated& distribution locations in SA (MOWE, 2005)

Figure 2.4: The different sources of households of water supply in SA (MOWE, 2005)

2.1.5 Health service

All parts and sectors in Saudi Arabia provide health care services under the supervision of the Ministry of Health with some other governmental health sectors

such as the National Guard health services, health services of the Ministry of Defense & Aviation and Security Force health services. These sectors have their own hospitals that provide health care for their employees and their dependents (Garout, 2000). In addition, there is a private sector, and this service is increasingly being used by business companies to provide medical care for their employees.

The Ministry of Health has three main assistant agencies; curative preventive, prevention medicine and an assistant agency for planning and research with many other programs and services. The Primary Health Care Centers (PHCC) are the first health care point for the community. The Ministry of Health has made considerable efforts to develop these centers. In 2008 there were more than 1986 centers covering each part of the country (MOH, 2009; CDSI, 2010). PHCCs provide basic health care services including maternity and child health and immunization programs. More than 60% of these centers have laboratory facilities, 25% have an X-ray service and 45% have dental services. The second level of health care is hospitals, which are widely distributed throughout each main part of the country. In 2007 there were 393 hospitals with 58.8% (231) under the supervision of the MOH with 31720 beds (MOH, 2008; CDSI, 2010). Table 2.5 shows the health resources indicators for Saudi Arabia during 2008.

	-
Physicians	21.5
Dentists	2.8
Pharmacists	6.4
Nurses	40.8
Allied health Personnel	20.8
Hospitals beds,	21.72
Primary health care centers	0.8
Governmental hospital beds	17.14
Private hospital beds	4.58

Table 2.5 The Health resources rate for Saudi Arabia during 2008(Rate per 10,000 Pop) (CDSI, 2010)

Infectious Diseases

In order to eliminate infectious diseases the MOH is continuously upgrading its prevention and control programs. As a result of these efforts during the last decades, they have eliminated several diseases such as diphtheria and poliomyelitis. The number of cases of the remaining diseases such as tetanus and measles have been reduced. This is due to the control and surveillance of the epidemiological control units spread over the whole of the country (MOH, 2008). There has been a reduction in the incidence of some vaccination target diseases such as whooping cough and mumps, due to the high immunization coverage achieved over the last decade.

Despite advances in the health services, the control of infectious diseases remains a challenge. For example, a massive spray against Rift Valley Fever lasting for more than six months started in Asir and Jazan during 2000. This action was carried out by many related governmental agencies using airplanes, cars and manpower (Dr A. Abdoon 2007 pers. Comm.).

Figure 2.5 shows the reporting system for infectious diseases in MOH, Saudi Arabia. The first level for the weekly notification in each region of this country is the primary health care centres, clinics, hospitals and any related health center. Monthly report is sent by the related department into the health directorate, primary health care department & preventive medicine division for each administrative region. Each health directorate of a region sends these reports to the infectious diseases department in the Ministry of Health.



Figure 2.5: Reporting System for infectious diseases in MOH, Saudi Arabia (Epidemiology Field, MOH, 2007)

Table 2.6 presents the reported cases of notifiable infectious diseases in the country during 2008 (MOH, 2009).

Cholera	7	Malaria	1497
Diphtheria	0	Leishmaniasis	2321
Whooping cough	30	Schistosomiasis	699
Tetanus neonatorum	13	Rift valley fever	0
Tetanus, other forms	4	Plague	0
Poliomyelitis	0	Yellow fever	0
Measles	158	Guillian - Barre syndrome	121
Mumps	31	Dengue fever	913
Rubella	15	Rabies	0
Chickenpox	60007	Echinococcus hydatid disease	6
Brucellosis	3447	Salmonellosis	1292
Meningococcal	7	Shigellosis	188
Pneumococcal	25	Amoebic dysentery	3311
Haemophilus influenza.	4	Typhoid & paratyphoid	269
Meningococcal, other forms	270	Unspecified hepatitis	255
Tuberculosis (TB)	3900	Hepatitis A	1678
Pulmonary TB	2635	Hepatitis B	5066
Extra – pulmonary TB	1265	Hepatitis C	2733

 Table 2.6: Reported Cases of notifiable infectious diseases by Region, Saudi

 Arabia during 2008 (MOH, 2009)

2.1.6 Zoogeographical areas of Saudi Arabia

Saudi Arabia lies at the crossroads of three of the world's major zoogeographical realms, the Palaearctic zone, the Afro-tropical and the Indo-Malayan and is itself at the centre of the Eremian zone (or Saharo - Sindian) desert region which is the vast desert belt extending from Morocco to western China (CBD, 2007). Asir and Jazan Regions are located in the Afro-tropical zone as some countries in the east of Africa. This could provide suitable climate conditions for some vectors of vector-borne diseases such as malaria and leishmaniasis (Abdoon and Alshahrani, 2003).

2.1.7 Distribution of Malaria, Leishmaniasis and Schistosomiasis in Saudi Arabia

The focus of the current study will be on malaria, leishmaniasis and schistosomiasis. The distribution of these illnesses varies from one region to another. It even varies within regions and from one year to another.

Malaria transmission is confined to the southwest of Saudi Arabia (Jazan, Asir, and A1-Baha Regions) but also occurs in some isolated rural areas in Makah and Madinah Regions. Environmental conditions play an important role in the distribution of malaria, as they affect the mosquito species to be found in an area, its life cycle, the

life span of the plasmodium and the density of mosquitoes. These conditions may vary from year to year. Malaria transmission is also affected by the control measures taken against it. Figure 2.6 shows the distribution of malaria across Saudi Arabia, the pathogenic mosquitoes in each area and the endemicity of malaria in each area (MOH, 2006; Al-Saghayer S. and Alattas M, 2006). The distribution in the map shows that the regions of Asir and Jizan are higher endemic areas followed by Makkah and Albahah Regions. The low endemic areas are in Najran, Madinah, and Tabuk Regions. The other Saudi Arabian regions do not report indigenous cases.



Figure 2.6: Malaria distribution in Saudi Arabia and prevelace of the vector (The main vector in the high- endemic foothills area is arabiensis, and sergenti in the coastal plain) Adapted from (Al-Saghayer S. and Alattas M, 2006)

There are two types of leishmaniasis in Saudi Arabia; cutaneous leishmaniasis (CL) and visceral leishmaniasis (VL). CL and VL have differing distributions across Saudi Arabia. According to MOH, (2008) CL occurs in most Saudi Arabian regions except the Northern region, AlJouf, Qurayat, Hafr-Albaten and Qunfedah Regions. The peak of CL occurs during winter, and the lowest incidence occurs during the end of spring and the beginning of summer (MOH, 2008). VL has a different distribution and during 2008 90.6% of cases occurred in Jazan, 6.3% in Asir and most other cases (3.1%) were reported in Qaseem (MOH, 2008).

The geographical distribution of schistosomiasis in Saudi Arabia varies depending on the type of the disease. During 2008, urinary schistosomiasis was recorded only in Jazan and Asir. Intestinal schistosomiasis was recorded in Taif, Albahah, and Asir. However, combined schistosomiasis was recorded only in Asir, as environmental predisposing factors are present in this region (MOH, 2008). The geographical distribution of the intermediate host of schistosomiasis is different from the distribution of cases of the disease. The host (snail) of urinary schistosomiasis is endemic in Makkah, Madinah, Jazan, Asir and Lith regions. The host of Intestinal schistosomiasis is endemic in Taif, Albahah, Asir, Najran, Makkah, Madinah, Hail, Jeddah and Jazan regions. Both types of host are present in Jazan, Makkah, Asir and Madinah (MOH, 2008). Figures 2.7 and 2.8 demonstrate the distribution of endemicity of schistosomiasis in the regions of Saudi Arabia in 2009. The distribution in the map shows that the regions of Asir Jizan and Albahah have low incidence of schistosomiasis followed by Makkah Region which has irregular incidence. There were no cases recorded in the regions of Najran, Riyadh, Madinah Tabuk, Hail and Al-Jouf. The rest of Saudi Arabia regions are free incidence. The diagram in Figure 2.8 shows that Asir and Taif regions have the highest schistosomiasis cases followed by Jazan, with lowest cases in Alqunfidah regions. It is worth mentioning that all of these regions are located in southern Saudi Arabia, with the exception of Taif region which is close to these regions.



Figure 2.7: distribution of endemicity of schistosomiasis in Saudi Arabia during 2009 (Dr M. Alzahrani 2010 pers. Comm.)



Figure 2.8: The schistosomiasis endemic regions in Saudi Arabia during 2009

2.2 Asir Region

2.2.1 Geography and Demography:

The Asir region is located in the south west of Saudi Arabia. It is a large administrative region, with its capital at Abha. It is divided into the emirate quarter and eleven governorates (Central Department and Information 2004; 2010) as illustrated in Figures 2.9 and 2.10. It has a varied terrain ranging from highlands, high mountains (2000 to ~3000m above sea level) and the lowlands (Tehama) parallel to the Red Sea (Gaafar et al., 1997). It also includes a large area of the desert to the north and east as far as the cities of Bishah and Tathlith. The mountain range has the highest rainfall in Saudi Arabia (MEPS, 2007; Subyani, 2000). The highest place in Saudi Arabia is in this region (Alsoudah, >3000m).



Central Department of Statistics & Information

Figure 2.9: Map of Asir Region with its emirate quarter and governorates (CDSI, 2005)



Figure 2.10: Topography of Asir Region.

There were more than 20 cities, 915 community centers and 7760 villages in this region in 2008. The main cities are Abha, Khamise, Bishah and Muhail. The population of Asir Region during 2010 was more than 1.9 million. Table 2.7 shows the distribution of population of the emirate quarters and governorates of this region (CDSI, 2010). The population of the sectors of Abha, Khamise and Bishah are more than 56% of the total population. Khamise has the highest population followed by Abha. The lowest population is in the Animas sector.

Abha	366551
Khamise	512599
Bishah	205346
Animas	54119
Muhail	228979
Sarat-Abidah	67120
Tathlith	59188
Rejal-Alma	65406
Ahad-Rifadah	113043
Dhahran-Aljanoub	63119
Balgarn	74391
Majaredah	103531
Total	1913392

Table 2.7: The population distribution of the emirate quarter and governorates of Asir region (CDSI 2010)

2.2.2 Climate in Asir

The topography of Asir has a large influence upon its climate. Therefore Asir can be divided climatically into three areas:

- Asir Sarawat Mountains (Highlands): A series of high mountains extending from the north of Asir Region to the south (2000 -~3000m above sea level). The annual average temperature for this area is 17.7^oC, average rainfall is 500mm (the highest rainfall in Saudi Arabia), and the annual average relative humidity is 53.1%.
- Asir Plateau (a part of Highlands Group): In the NE of Asir with an elevation range of between 1000 and 2000m, decreasing to the east. The annual average temperature for this sector is 19.5^oC, with 250mm annual rainfall and an annual average relative humidity of 44.4%.
- 3. Tehama (Lowlands): This area can be divided into two sections
 Foothills (300 900m) Red Sea Costal Plain (0 200m).

This area receives an annual rainfall of 350mm, and temperature ranges between 25° C and 49° C during summer. The relative humidity varies from 55% in summer to 70% in winter.

2.2.3 Drinking Water Services

Sources of water in this region are wells, dams and springs in addition to desalinated water. Desalinated water arrived in Abha for the first time in 1988 and the amount of water produced in 2007 was 47.68 million tons. Water is distributed throughout the region by a network of pipes which were initially installed in Abha during 1995. This network distribution of drinking water in Asir now covers four cities, Abha, Khamise, Ahad-Rifadah and Muhail. Within each city the proportion of homes on network supply is 70%, 30%, 50% and 30% respectively. This region suffers from a shortage of drinking water and most people who live in the countryside depend on water from wells or dams which may have health consequences. In addition, in this region the shortage of drinking water increases during summer with the beginning of the tourism season (General Directorate of Water and Electricity in Asir, 2008).

2.2.4 Environmental Problems

The region of Asir covers a large area and it has a border with Yemen where there are a lot of infectious diseases. These include vector-borne diseases such as malaria and leishmaniasis. It also has a border with Jazan region which suffers from the same problems. Therefore, it is influenced by conditions in neighbouring regions. Asir has many man-made dams (56) and lakes which form after rainfall. Both these may encourage the presence of mosquitoes or the parasites of schistosomiasis or other harmful insects. Some parts of Asir, such as the foothills are hot and humid during summer, and this weather may encourage some types of insects to multiply. Furthermore, the rural areas and most of the countryside areas in the region use cesspools which may contribute to the contamination of underground water and the environment (General Directorate of Water and Electricity in Asir, 2008; Dr A. Abdoon 2007 pers. Comm.; MOH, 2008).

2.2.5 Health Services

The Health Affairs Directorate in Asir represents the MOH in this region. This directorate provides the health care services in the region including curative medicine and preventive medicine. During 2007, there were 290 primary health care centers, 22 hospitals (18 public & 4 private) and one main referral hospital (Asir Central Hospital) in Abha as the capital city of this region. In addition, there are many assistant preventive medicine departments related to public health and environmental health.

Table 2.8 demonstrates the number of reported cases of some infectious diseases in Asir during 2008. This table shows that the highest number of cases of the infectious diseases in this region is from chickenpox followed by brucellosis, hepatitis B and amoebic dysentery. Other diseases exist in Asir including climate-sensitive diseases such as leishmaniasis and schistosomiasis. Sixteen infectious diseases had no cases in Asir during 2008.

	(1)1 C	, 2007)	
Cholera	0	Malaria	31
Diphtheria	0	CL Leishmaniasis	130
Whooping cough	1	Schistosomiasis	115
Tetanus neonatorum	0	Rift valley fever	0
Tetanus, other forms	0	Plague	0
Poliomyelitis	0	Yellow fever	0
Measles	1	Guillian - Barre syndrome	11
Mumps	0	Dengue fever	0
Rubella	1	Rabies	0
Chickenpox	4300	Echinococcus hydatid disease	0
Brucellosis	563	Salmonellosis	9
Meningococcal	0	Shigellosis	0
Pneumococcal	0	Amoebic dysentery	135
Haemophilus infi.	0	Typhoid & paratyphoid	30
Meningococcal, other forms	9	Unspecified hepatitis	13
Tuberculosis (TB)	125	Hepatitis A	93
Pulmonary TB	84	Hepatitis B	296
Extra – pulmonary TB	41	Hepatitis C	105

Table 2.8: Reported cases of notifiable infectious diseases in Asir during 2008 (MOH. 2009)

Table 2.9 shows comparisons among the main sectors of Asir Region 2004 for the rate of climate-sensitive borne diseases such as malaria, leishmaniasis and schistosomiasis. This table also illustrates the health infrastructure for each sector including the number of hospitals and primary health care centers.

Sector	Altitude L	Schist R	Malaria R	Parasite R	CL. R	Hospitals No.	PHCCs No.	Pop. that PHCCs cover	Pop. per every PHCC	No. of patients in month	% patients per pop. in month
Abha & its sectors	2500	0.0	0.0	14.7	0.2	3	44	152971	3477	43782	28.62
Khamise	1800	0.1	0.0	23.3	1.4	1	15	177700	11847	45601	25.66
Ahad-Rifadah	1700	0.0	0.0	60.6	0.0	1	9	64669	7185	16182	25.02
Al Madhah	1600	0.0	0.0	47.4	0.0	0	17	33950	1997	14898	43.88
Tathlith	975	0.0	0.0	17.5	0.4	1	18	50435	2802	26282	52.11
Sarat-Abeeda	2400	1.0	0.0	42.5	3.1	1	15	47716	3181	17106	35.85
Dhahran Al Janoub	2020	1.8	1.5	72.4	2.4	1	21	45308	2158	19658	43.39
Tanoma	2600	2.3	0.0	28.9	1.7	2	15	35182	2345	16588	47.15
Sabt-Alalaya	2200	6.6	0.0	175.4	0.7	2	18	59038	3280	20743	35.13
Muhail	450	3.2	0.7	6.39	0.0	1	22	120474	5476	38625	32.06
Al Majaredah	450	0.4	0.9	18.71	0.3	1	19	66814	3517	24601	36.82
Rejal	700	0.0	5.6	21.7	2.0	1	19	39180	2062	10995	28.06
Alberek	20	0.0	0.0	0	1.1	1	7	18200	2600	4291	23.58
Al Gahma	20	0.0	0.7	14.6	0.0	1	4	13619	3405	5352	39.30
Marabah	600	-	1 N	-	-	-	-	-	-	-	-
Al Farshah	450	-	51 N	-	-	-	-	-	-	-	-
Ballesmer	2250	-	-	-	-	1	9	10665	1185	2287	21.00
Central Laboratory	2200	11 N	32 N	279 N	68 N	-	-	-	-	-	-
Bishah	1020	-	6 N	-	-	3	35	-	-	-	-
Total	-	-	-	-	-	21	287	935921	-	306991	-

 Table 2.9: Comparison of some health infrastructures and selected infectious diseases among some sectors of Asir Region 2004 (Department of preventive medicine, Health Affairs Directorate in Asir, MOH, 2006)

R: No. of cases /10000 population. N: No. of cases. L: above sea level. Schist: Schistosomiasis. PHCC: Primary health care. Pop: population. -: No information available

2.2.6 Distribution and Endemicity of Malaria, Leishmaniasis and Schistosomiasis in the Asir Region

According to the Vector Control Administration in Asir, environmental factors such as climate variables and topography play an important role in determining the distribution of many infectious diseases, in particular, vector and water-borne diseases. However the methods of control are different depending on the type of disease. Control methods may also vary between seasons and from one place to another. This fact is appreciated by the health decision makers in the Ministry of Health and in this sector.

This study will focus on and discuss the impact of climate variables upon three diseases (malaria, leishmaniasis and schistosomiasis). Each disease will be examined separately.

Chapter 3

<u>Influence of Climate Variables upon</u> <u>Malaria Cases in Asir Region, Saudi Arabia</u>

3.1 Introduction

Malaria is one of the major health problems facing tropical and subtropical areas of the world. It is the most important vector-borne disease, and WHO estimates that 2400 million people live in malarious regions of the world (WHO/SDE/PHE, 1999). It is responsible for an estimated 300–500 million clinical attacks globally (Keiser et al., 2004). The exact number of deaths caused by malaria is unknown but has been estimated at 1.5 - 3 million annually (De Savigny et al., 2004). Malaria transmission is sensitive to weather and climate conditions, so a changing climate is likely to change transmission dynamics in many regions (e.g. increased temperature or abnormal rainfall). Indeed, it is perceived as the vector-borne disease most likely to be affected by global climate change (WHO/WMO/UNEP, 1996; Mc Michael et al., 1996; Hana et al., 2001).

This chapter will start by providing an overview of malaria transmission and its life cycle involving two hosts; a human and a mosquito, and goes on to describe the symptoms and causes of malaria and how this illness may be prevented and treated. We will then present the literature review of malaria and previous studies that have looked at relationships with climate variables. The study area will then be introduced followed by a description of the data sources used and the method employed. This data will then be analyzed statistically to look at any relationships with climate variables. Finally we will present the results followed by discussion and conclusion.

3.2 Malaria Transmission

Malaria is a vector borne disease caused by protozoan parasites of the genus Plasmodium. It is caused by any of four species: *P. falciparum, P. vivax, P. ovale,* and *P. malaria.* These parasites are transmitted from person to person by the bite of an infective female anopheles mosquito (WHO, 2005). P. falciparum is responsible for 90% of deaths but only about 15% of infections. However, P. vivax is the most common cause of infection, and responsible for about 80% of all malaria cases (Mendis et al., 2001).

About 30 (out of 400) species of anopheles mosquitoes in the world are malaria vectors of major importance. However, throughout the world different species are responsible for malaria and these species vary in their ability to transmit the parasite. For example, Funestus, Gambia and Arabians, are highly efficient vectors which are spread throughout sub-Saharan Africa, whereas Stephensi and Minimus predominate in Asian countries are less efficient parasite transmitters (WHO, 2005).

3.3 Life Cycle of Malaria

The US Centre for Disease Control and Prevention (CDC, 2010) describes the life cycle of the malaria parasite and this is presented in Figure 3.1. The life cycle of malaria parasite consists of two host stages involving human and mosquitoes. In the initial stage, female mosquito (anopheles) injects and inoculates sporozoites into the human host during a blood meal. The sporozoites travel through the blood into the liver cells (hepatocytes) where they mature into schizonts which further ruptures into merozoites. In plasmodium vivax and ovale, the maturation into schizonts may not be achieved spontenously and can remain in the liver as dormant stage (hypnozoites) for a long while and may cause a relapse by re-infecting the blood. The schizoites containing mature merozoites in the liver cells undergo asexual multiplication in the red blood cells (erythrocytes). While in the erythrocytes, the merozoites grow into a ring-shaped form called trophozoite. Some merozoites then differentiate into male or female sexual forms called gametocytes (The only forms found in the blood). The gametocytes are then ingested by anopheles mosquito during blood meal where they fertilize each other, forming motile zygotes called ookinetes in the mosquitoe's midgut. The ookinetes develop further into oocytes which mature and rapture into sporozoites. The sporozoites then migrate into mosquito's salivary gland where they stay until they are inoculated into a human host for further development of the life cycle of the parasite (CDC, 2010; NIAID, 2010).



Figure: 3.1: Schema of the life cycle of malaria (CDC, 2010)

3.4 Symptoms of Malaria

The common first symptoms of malaria usually appear ten to fifteen days after the infection but this period can be longer due to the dormant stage which was discussed earlier. Severe malaria is almost exclusively caused by P. falciparum infection and usually arises 6-14 days after infection. It can cause severe often fatal malaria if not treated promptly with effective medicines (WHO, 2005). Malaria can have many symptoms (WHO/HTM/MAL, 2005; Hana et al., 2001) such as headache, nausea, vomiting, muscle pain and malaise, rigors consisting of severe shakes or muscle spasms and chills. The severity of the infection depends on which species of the malaria parasite is involved, and it can cause (WHO/HTM/MAL, 2005; Hana et al., 2001):

- 1. Enlargement of the spleen.
- 2. Liver tenderness.
- 3. Brain problems including seizures, and coma.
- 4. Problems of the immune system.
- 5. Affect on the structure of the blood corpuscles, anemia.

- 6. Kidney failure.
- 7. Respiratory failure.

Some groups of people have a higher risk of contracting malaria (WHO, 2010). These include young children, as they have not yet developed immunity and are therefore at risk of the severe forms of this illness. Also pregnant women who have no immunity to the disease, or are only semi-immune, are also at risk from an increased rate of miscarriage and maternal death, and their babies may have a low birth weight. In addition the risk of malaria for pregnant women infected with HIV is significantly increased, and they also risk passing the HIV infection to their babies. This heightened risk of malaria applies to all people infected with the HIV/AIDS virus. People who live in non endemic areas lack immunity to endemic areas.

3.5 Preventing and Treating Malaria

Malaria control is one of the main requisites in development of health care, and this disease requires early diagnosis and speedy treatment (Ridley, 2001). Once malaria is diagnosed there are currently a range of drugs available to treat malaria which can reduce the duration of the disease and prevent the majority of deaths (Ridley, 2001). In the future vaccines to prevent this disease hold out promise for future disease control (World Resources Institute (WRI) et al., 1998) but are not available at present.

The aim of malaria control is to reduce the numbers of parasite-infected mosquitoes and therefore hinder transmission of the vector of disease. Two main interventions are available for controlling the malaria vector; spraying insecticide indoors and use of insecticidal nets. All of that can be complemented with other locally-appropriate methods, such as larval control or environmental management (WHO, 2010). Some communities have controlled breeding of the vector by removing algae from breeding places, and use it as manure. Also coordinated efforts among multinational agencies, and private companies can help to reduce the disease incidence (WRI et al., 1998). Based on the insecticide regulation from the World Health Organization (WHO, 2010) there are only 12 types of these insecticides can be use for indoor residual spraying.

The most common drug treatments for malaria include; chloroquine, qualaquin hydroxychloroquine, mefloquine and a combination of atovaquone and proguanil (CDC, 2010). In some parts of the world there may be drug resistance for various reasons (WHO, 2010). For example, some patients do not complete their treatment and stop taking the malaria drugs in the early stages of recovery when the parasite is still in the blood. In these cases such as these the parasite will survive and be drug resistant if it is transmitted to another person. Drug resistance may develop and spread to a wider community, presenting a considerable public health problem due to lack of alternative treatments (WHO, 2010).

3.6 Relationship between Malaria and Climate

Data of atmospheric variables such as: temperature, rainfall, wind, relative humidity and cloudiness are described as climate variables (McMichael et al., 2003). Climate variability occurs on many time scales, however weather events usually occur on a daily time scale and these variables are associated with many health impacts. Climate variability at other scales also affects health (Harold et al., 2000).

The vector of malaria (Anopheline mosquito) and its parasite are sensitive to three main climate factors; temperature, rainfall and relative humidity (Hana et al., 2001). Climate predicts, to a large degree, the natural distribution of malaria (Bouma et al., 1996). Pampana, (1969) noted that the absolute temperature and its duration can affect the life cycle of malaria, but that this varies by parasite species. The development time of the Anopheline egg decreases as the temperature increases from 21°C to 27°C (Patz et al., 1998; Patz et al., 2000). Also McMichael et al., (1996) and Hana et al., (2001) indicate that 25 - 27°C is the optimum breeding temperature for both the mosquito and parasite, with 40°C being the maximum temperature. The life cycle of some malaria species below 20°C is limited, but colder than this the species can survive as the vector often lives in houses where the temperature is warmer. 8-10°C has been identified as the minimum temperature required for mosquito development (Hana et al., 2001) (Table 3.1).

	Mosquito	Parasite
Minimum	8–10°C	14–19°C
Optimum	25–27°C	25–27°C
Maximum	$40^{\circ}C$	40°C

 Table 3.1: Minimum, optimum and maximum temperature for anopheles mosquito (from Hana et al., 2001; McMichael et al., 1996)

Rainfall is another factor affecting survival of anopheline mosquitoes, by providing breeding sites, as this vector breeds in water habitats (Oaks et al., 1991). However, the amount of rainfall is important as breeding larvae can be flushed away by excessive rainfall or rainfall associated with stormy conditions (Pampana, 1969). Also time of year whether in the wet or dry season, and taking into account the amount and intensity of rainfall can have an impact upon malaria survival rates.

Rainfall also increases relative humidity and modifies temperature, thus it can affects malaria transmission through these mechanisms (Pampana, 1969). Although plasmodium parasites are unaffected by relative humidity, the activity and survival of anopheline mosquitoes can be limited when the average monthly relative humidity is below 60% (Pampana, 1969).

Regarding climate change, increase of temperature by time can increase the survival of mosquitoes, and this could be the reason that malaria can now survive at higher altitudes (Patz and Lindsay, 1999). Also it would be able to transmit some species of malaria that were not present in free malaria area depending on the change of the environmental conditions for the parasite or mosquito (Pampana, 1969).

3.7 Previous Studies into Malaria (Transmission, Vector and Incidents) and Climate Variables

This section on previous studies into malaria and weather will focus on semi arid areas across the world. One study which has looked at the relationship between malaria and environmental factors was undertaken in eastern Sudan (Hamad et al., 2002). It was carried out in two villages close to Gedaref city. Malaria case incidence was compared with the number of mosquitoes detected and with rainfall, relative humidity, maximum, minimum and mean temperatures, wind speed and direction. Climate data was also compared to data on the numbers of both adult anopheline mosquitoes and their immature stages. These were sampled bi-monthly in each village for two dry seasons and two transmission seasons from April 1994 to October 1995. The study showed that the abundance of mosquitoes was positively related to monthly rainfall, maximum temperature and relative humidity. From the end of the short rainy season in October until February there was a gradual drop in the number of mosquitoes. Vectors re-appeared in June as the humidity rose with the onset of rain. Despite the hot dry season between February and May, sporadic asymptomatic malaria infections were detected in the two villages.

In SW Uganda (Kabarole District), researchers analyzed the incidence of reported clinical malaria cases and its correlation with rainfall between 1995 and 1998. The data showed a positive relationship exists between rainfall and malaria incidence. In October - December 1997 rainfall was above the upper 95% confidence limit of previous years, and this was associated with a big increase in plasmodium falciparum infection (Kilian et al., 1999).

Malaria rates in Zimbabwe during 1913 - 1993 (Freeman et al., 1996) were compared to weather data and the results showed that, rainfall has little effect on the number of malaria cases. However, temperature appears to have an impact on malaria. Greater than average mean temperatures in September were correlated with an increase in the severity of malaria in the following year.

In Pakistan, malaria transmission was associated with the seasonality of rainfall over September and October. A relationship was also observed with temperature during November and December, and with humidity in December (Bouma et al., 1996).

Another study in Sri Lanka (Briet et al., 2008) (a) found that the relationship between rainfall and malaria cases in most districts varied during study period (1972 – 2005). Four analytical methods were explored: (I) Cross-correlation (monthly malaria cases and monthly total rainfall). (II) Cross-correlation with pre-whitening: (Crosscorrelation with the seasonality and autocorrelation removed by simple pre-whitening allows for detection of the time lags of rainfall preceding malaria). (III) Inter-annual, and seasonal inter-annual regression (the series of differenced transformed annual malaria cases was studied to determine the correlation to total annual rainfall). The cross-correlation results showed that malaria cases in most districts were positively associated with rainfall with a 0-4 month lag. However, they were also negatively associated with rainfall with a 4-9 month lag. Also low seasonal rainfall was associated with low seasonal malaria, and high seasonal rainfall was associated with high seasonal malaria. Using the cross-correlation with pre-whitening method (including an autocorrelation term to control for the previous months malaria) malaria was only significantly associated with rainfall in five out of 25 districts at a lag of 0-1 month. Only 6 from the 25 districts showed a negative relationship with rainfall at a 2-5 month lag. Although malaria decreases with annual rainfall by inter-annual analysis method for eight districts in the centre-west of the country, the seasonal this method showed that the effect of rainfall on malaria varied according to the season and geography. Districts in the east showed positive correlations with rainfall, however the relationship in the centre-west and in the north shows significant negative correlations for rainfall during February to June. A spatial malaria study was undertaken using the same data as the previous paper (Briet et al., 2008) (b). This confirmed that after controlling for spatial autocorrelation season and rainfall were positively associated with malaria.

One study in the highlands of Kenya showed that climate did not play an important role in malaria transmission because average temperature and rainfall did not change during the time that malaria rates changed (Malakooti et al., 1998). Another study in the highland Tea Estates of Kericho, Western Kenya (Dennis et al., 2002) confirmed that while the total hospital admissions of malaria over 30 years (1966–1995) increased significantly, all meteorological variables showed no significant relationships with disease. This was observed when these variables were combined into a monthly suitability index for malaria transmission. Therefore the researchers concluded that recent climate changes have not caused the highland malaria resurgence in this area. However, epidemics of malaria in different areas of the highlands of Kenya in third study found that increased rainfall was related to the epidemic (Kigotho, 1997).

A study in Central India investigated whether climate and malaria were related in a highly malarious village during the periods 1967-2000. The result showed no relationship between rainfall and malaria incidence (Singh et al., 2002). Another study examined the relationship between climate change and malaria in India (Bhattacharya et al., 2006). It noted a positive correlation between malaria transmission and relative humidity when the range 55 to 80 %. It also found that maximum number of reported cases of malarial occurred when the average temperature between 15 to 30°C.

In the Ethiopian highlands (Debre), there was a positive correlation between malaria incidence and with increasing of temperature from 1968 to 1993 (Tulu, 1996). Another study in Ethiopia (Woube, 1997) found mixed associations between malaria and rainfall. In 1993 there was high rainfall and no malaria incidence, but in 1984 -85 there was very little rainfall but a high malaria incidence. The study noted that epidemics were usually attributed to higher temperatures, rainfall and relative humidity. However, another Ethiopian study (Teklehaimanot et al., 2004) compared malaria incidence between two areas depending on the temperature. In the hot area rainfall was followed quickly by an increase in malaria cases. Minimum temperature was non-significant. In the cold area rainfall was also correlated with increasing malaria but after a longer delay. The same delay was associated with minimum temperature.

A study in Nouna and the villages of Cisse and Goni in Burkina Faso (Ye et al., 2007) studied the effect of meteorological conditions upon clinical malaria risk among children. It confirmed that weather has a positive significant influence on malaria rates among children under five years old. Mean temperature was the best predictor of clinical malaria incidence, and the risk of clinical malaria increased with mean temperature up to 27 °C. For rainfall only the effect was observed when it was over 100mm. In highland regions the limiting factor for malaria appeared to be the low temperature because of the high altitude.

In Like Bufundi, Kilibwoni, East Africa, located at an altitude above 2000m, the associations between climate and malaria morbidity were studied (Kristan et al., 2008). A peak in rainfall during April-May was followed by a peak in malaria occurrence in June. The second peak of rainfall in August was preceded by a low temperature in July. This was followed by a peak in malaria incidence in October.

Alam et al., (2008) studied the relationship between weather and the incidence of malaria in the Chittagong Hill Tracts in Bangladesh (1997 – 2005). The analysis

was conducted by dividing the seasons into two. The first season from April to October is warm and wet and the other November to March is cool and dry. They noted that during first season there is a considerable increase in malaria cases. In the second, mosquitoes are less active with a decline in cases of malaria. They mentioned that optimal relative humidity for mosquito survival is 60 %, and a high relative humidity lengthens the life of the mosquito.

Another study investigated the impact of temperature and rainfall upon malaria in Lagos Nigeria between 1990 and 2003 (Oluleye and Akinbobola, 2010). Temperature and malaria were highly positively correlated between February and December but negative associations occurred in November. Rainfall and malaria were highly positively correlated during January, September, November, and December. Negative associations were observed in the months of April, June and October.

Another study examined the correlation between monthly climatic variables and malaria in Shuchen County, China (Bi P et al., 2003). The climatic variables were maximum, mean and minimum temperature, relative humidity and rainfall. The dependant variable was the malaria incidence. The study examined the correlation between malaria incidence and each climatic variable while controlling for autocorrelation in the data. There were highly positive correlations (P<0.0001) with each climate variable lagged at one month. The study noted that minimum temperature seemed to play a more important role than maximum temperature in affecting malaria incidence. The study notes that other factors can play role in the transmission of malaria such as ecological and environmental issues, also social and economic determinants. However, for forecasting malaria climatic variables are best suited.

The previous studies have all examined the Influence of climate variables upon malaria cases. The review will now focus upon studies which have examined the influence of climate variables upon vector abundance. An entomological study was conducted into the spatial and temporal variations in malaria transmission in a low endemicity area in Tanzania (Oesterholt et al., 2006). Part of this study investigated the relationship between vector abundance (Anopheles arabiensis) and precipitation. The study noted that the peak mosquito numbers followed the peak of rainfall. Amerasinghe et al., (1999) studied malaria vectors (Anopheline mosquitoes) in a traditional dry zone village in Sri Lanka. Monthly malaria incidence, during the study period, was not significantly correlated with monthly rainfall. However, it was correlated indirectly with rainfall. Abundance of some species of Anopheline mosquito were correlated with the previous months rainfall. Positive significant correlations were found for *An. annularis, An. subpictus, An. Vagus* and *An. varuna*. Malaria incidence in the study was also correlated with the abundance of Anopheline species. A positive significant correlation was seen between an outbreak of malaria and the numbers of some species Anopheline one month earlier.

The focus of this thesis is Saudi Arabia and in this country there are several studies related to malaria and weather. In Gellwa, Al-Baha Region, located in SW Saudi Arabia a malaria outbreak occurred (A1-Abdullatif et al., 1996). Between January 1 to March 31, 1996, 476 confirmed malaria cases were reported from in and around the Eliab Valley. This was an increase from the usual numbers of cases reported each year from 1990 to 1995 (between 32 and 176). In 1996, following heavy rainfall in Tehama, many valleys were flooded and the malaria increase was attributed to this. The study also reports that seasonal peak in malaria incidence in Saudi Arabia occurs between January and March.

Another study Gaafar et al., (1998) examined the clinical aspects of malaria in the Asir Region, Saudi Arabia. The researchers reported 334 confirmed malaria cases in Asir Central Hospital, Abha, during the period 1991-1995. This study has emphasized that malaria is still a health problem in the Asir region. Cases were seen throughout the year, with two peaks, in February and June. These peaks are most likely to be related to the rainy and hot seasons, respectively. In Riyadh Alkhalife, (2003) studied malaria infections between 1996 and 2001, and found that an annual peak of malaria incidence occurred between February and April.

Another important study was conducted in Jazan SW Saudi (Al-Jaser, 2006). They tried to determine the yearly correlation between weather data and malaria between 1997 and 2002. The data showed a significant positive correlation between relative humidity and malaria cases, but a significant negative correlation between temperature and malaria. The researcher noted that the number of malaria cases has declined significantly after the year 2000, when an outbreak of the Rift Valley fever happened in Jazan and Asir.

Table 3.2 summarises the main studies (statistically) of the correlation between malaria or its vector and climate variables.

No	The author	Place	Dependent Variable	Independent Variable	The relationship & Lag time (month)
1	Hamad et al., 2002	Easter Sudan	Abundance of mosquitoes	Rainfall Max Temperature Humidity	Positive 1-3
2	Kilian, et al., 1999	SW Uganda	No. of malaria cases	Rainfall	Positive
3	Freeman et al., 1996	Zimbabwe	No. of malaria cases	Temperature	Positive
4	Bouma et al., 1996	Pakistan	malaria incidence	Rainfall Temperature humidity	Positive 0-1 Positive 1-2 Positive 2
5	Briet (a) et al., 2008	Sri Lanka	No. of malaria cases	Rainfall	Positive 0-4 Negative 4-9
6	Malakooti et al., 1998	highlands of Kenya	Malaria transmission	Different Climate variables	No relationship
7	Kigotho, 1997	Different highlands of Kenya	No. of malaria cases	Rainfall	positive
8	Bhattacharya et al., 2006	India	malaria incidence	Temperature humidity	positive
9	Tulu, 1996	Ethiopian highlands	No. of malaria cases	Rainfall Temperature humidity	positive
10	Ye et al., 2007	Burkina Faso	No. of malaria cases for the children	Mean temperature Rainfall	Positive
11	Kristan et al., 2008	Kilibwoni, East Africa	malaria incidence	Rainfall	Positive 1-2
12	Oluleye and Akinbobola, 2010	Lagos Nigeria	malaria incidence	Temperature Rainfall	Mixed relationship depending upon the time of year
13	Bi P et al., 2003	Shuchen County, China	malaria incidence	max, mean and min temperature Humidity Rainfall	Positive 1
14	Amerasinghe et al., 1999	dry zone in Sri Lanka	Abundance of mosquitoes	Rainfall	Positive 1
15	A1-Abdullatif et al., 1996	SW Saudi Arabia	No. of malaria cases	Rainfall	Positive 1-3
16	Gaafar et al., 1998	Asir Region, Saudi Arabia	No. of malaria cases	Temperature Rainfall	Positive
17	Al-Jaser, 2006	Jazan Saudi	No. of malaria cases	Temperature Humidity	Negative Positive

Table 3.2: Summary of the main studies relating to malaria

The literature review highlights a number of important points:

- 1- A wide range of study designs have been used ranging from purely observational to those conducting statistical analyses. The complexity of statistical analyses varies as well.
- 2- Numerous different climate variables have been used including temperature, rainfall and humidity
- 3- Most but not all studies showed that climate variables have an impact upon malaria incidence. However, the impact of specific weather variables can be either positive or negative depending upon the setting and also the season. Lagging the climate variables is an important feature of some studies.
- 4- Other studies support these findings by showing that weather influences the abundance of the malaria vector (anopheles mosquito), and others found that the vector abundance has an effect upon the incidence of malaria.
- 5- Weather conditions are not the only factor which can affect malaria, and that there are many other factors such as ecological and environmental conditions, social and economical determinants. Some studies noted that climatic variables are best suited to predict future trends (Bi P et al., 2003).
- 6- There have been few studies in Saudi Arabia focusing upon the influence of climate variables upon the disease.

Based on the previous studies, it is clear that there is an urgent need to study and research the influence of climate variables upon the presence of malaria in Saudi Arabia, and in endemic areas such as Asir Region specifically. Building upon the literature review such, a study should examine a wide range of weather variables and also consider that any relationships with weather may vary seasonally. A statistical analysis seems the most appropriate study design and this may need to consider factors such as autocorrelation in these data as some previous studies have done.

3.8 Malaria in Saudi Arabia

Malaria is endemic in parts of the country with high rainfall. Transmission is confined to the SW (Jazan, Asir and A1-Baha Regions) but also occurs in some isolated rural areas in Makah and Madinah Regions. Figure 2.1 (Chapter 2) shows the distribution of malaria in across of Saudi Arabia and some species of pathogenic mosquitoes in each endemic or low endemic area (MOH, 2006; Al-Saghayer and Alattas, 2006).

Figure 3.2 shows the number of malaria cases in Saudi Arabia between 1980 and 2009. These data show a sharp decrease in malaria cases in the last ten years except an outbreak which happened at the end of 1997 and into 1998. The transmission of malaria is seasonal with peak incidence from October to April (Al-Saghayer et al., 1996). More than 50% of the total cases per year occur in Jazan, Asir, and Makah Regions (MOH, 2006). The population living in the most endemic areas is 1.7 million (~5% of the country's population) (Al-Harthy and Jamjoom, M, 2007). Many of the current cases of malaria reported in Saudi Arabia are imported from other regions or from overseas. Some cases are imported by pilgrims during Hajj to the Holy Cities (Makkah & Madinah). As an example, in 2008 the Ministry of Health in Saudi Arabia reported 2864 positive cases of malaria. From this 249 (8.7%) were indigenous cases (MOH Statistical Books, 2009). Of these cases 109 cases (43.8%) were from the Asir Region, 35.7% from Jazan and 12.9% from Qunfedah Aria. There were no indigenous cases in Riyadh and Al-Ahsa, Eastern Region, Taife, Qassim, Tabuk, Hail, the northern border, Najran, Al Baha and Qurayat. A malaria expert working in the Ministry of Health in Saudi Arabia (Dr S. Algaryah 2009 pers. Comm.; Al-Saghayer, and Alattas, 2006) noted that weather variables especially rainfall, play an important role in the incidence of indigenise malaria in the country. He gave the example of an outbreak of malaria cases during 1998 after a large amount of rainfall in some parts of Saudi Arabia during the end of 1997 and beginning of 1998 (see Figure 3.2).



Figure 3.2: Number of malaria cases in Saudi Arabia (1980-2009) (MOH, the Statistical Books 2000-2009; Al-Saghayer S, and Alattas M, 2006)

- Topographical Classification of Asir Region and Endemic Areas:

Asir Region is located in the SW Saudi Arabia and is divided into the emirate quarter and eleven governorates (CDSI, 2005) as illustrated in Figure2.9. This region is in the tropical zone as are some countries in the east of Africa (Abdoon and Alshahrani, 2003). It has a varied terrain ranging from highlands, high mountains (2000 to ~3000m above sea level) and the lowlands (Tehama) parallel to the Red Sea (Gaafar et al., 1998). It also includes a large area of the desert and hills to the north and east as far as Bishah and Tathlith. The mountain range has the highest rainfall in Saudi Arabia (MEPS, 2007).

Malaria is endemic in the lowlands (Tehama) in Asir Region (Figure 3.3) (Haddad, 1990). In the last few years the number of malaria cases has spread to adjoining areas of higher elevation, such as Abha and Khamise city (MOH, 2006).



Figure 3.3: Endemic sectors in Asir Region. Adapted from (The Vector Control Administration in Asir Region, 2007)

Depending on the endemic areas of malaria and topography, this region can divide into four main malarial sectors (Dr A. Abdoon 2007 pers. Comm.; MEPS, 2007) (Figure 3.4):

- 1. Tehama of Qahtan (TQ) is located among high mountains such as Farshah Al-Jauah, Al-Ergain and Wadi-Alhaiah. This sector is in the lowlands with an altitude between 800 and 1300m, and most of its areas called the foothills. Farshah Area has the main health center in TQ, and high malaria incidence. More than 90% of the pathogenic mosquitoes in these areas are Arabians.
- 2. Tehama of Asir (TA), area can divide into two sections:
 - a) Foothills such as Al-Majaredah, Muhail and Rejal-Alma, which have the highest malaria incidence in Asir. It is among the mountains and the valleys with many appropriate places for the vector (mosquito). The altitude varies between 300 and 900m. The main vector in this area is Arabians.
 - b) Red Sea Costal Plain (RSCP) such as Al-Gahma, Al-Berk, Al-Huraidah. These areas have low malaria incidence and the main vector is Sergenti. All

of the vectors in this area dependent on open water tanks and swampland. The altitude in this area is between 0 and 200m.

- North-East of Asir such as Bishah, Tathlith. The altitude for this sector is 1000 2000m. This sector has had no local transmission of malaria for over ten years. It had a few local cases before 1995, but all were imported from endemic areas.
- 4. Highlands (The Sarawat Mountains) which have altitudes 2000 ~3000m such as Abha, Khamise, Al-Numas. These areas have the highest rainfall in Saudi Arabia (542mm) (MEPS, 2007) but no local transmission of malaria. However, there are many reported cases in some health centers and the main region laboratory in this sector. These cases are likely to have come from endemic areas in the lowlands.

The endemic areas in Asir (TA and TQ) are drained by valleys, which are usually wet except in some dry seasons, and so provide suitable habitats for mosquito breeding (Alzahrani, 2007). These valleys maintain a continuous steady flow of water into the foothills area especially in TQ and some parts of TA.

Omar et al., (1999) described that in the malaria endemic areas of Asir there are two categories of the population at risk. The first are permanent residents living in the foothills (Tehama) or in the coastal plain. The second group is residents from the non-malaria's areas in the highlands, who travel for recreational activities to the endemic lowland areas during the transmission season.



Figure 3.4: Topographic map for Asir Region including the weather stations

3.9 Controlling of Malaria Transmission in Asir Region

There is a strategy for anti-malaria control measures in each endemic area of Saudi Arabia and this strategy depends upon the following guidelines (MOH, The Statistical Book, 2006; Dr A. Abdoon 2007 pers. Comm.)

- a. Spraying of the breeding foci of mosquito to control the larva. This method is wide spread all over the Saudi Arabia except the mosquito free areas.
- b. Spraying the houses (indoor) with insecticides that have residual effect. It is used in the areas of high density of malaria (Figure 3.5)
- c. Spraying the narrow spaces with a spray of minute particulates to reach to deep spaces. This method is used in areas with high cases of malaria.
- d. Mechanical control method.
- e. Wide spread of nets impregnated with insecticides.
- f. Removal of breeding habitats and water body unnecessary beside the valleys (Figure 3.6).
- g. Health education program.
- h. Collaboration with the governmental and private sector in control measures.



Figure 3.5: Spraying the houses (Indoor) with insecticides that have residual effect in endemic areas of Asir Region. (The Vector Control Administration in Asir Region, 2007)



Figure 3.6: *Removal of breeding habitats and unnecessary water body beside the valleys (The Vector Control Administration in Asir Region, 2007)*

The Vector Control Administration in Asir Region has made important progress regarding the control of malaria. In addition to the general strategy of ministry, their improved plan for Asir depends on four main issues (Dr A. Abdoon 2007 pers. Comm.):

 Control of places which have water born malaria parasite or transmission occurrences by spraying particular types of pesticides depending on the vector type. This usually happens in open water areas such as valleys, water tanks and dams. This type of spraying occurs all year round by a huge number of technical workers and a massive amount of equipment, cars, tents and sprays etc.
- 2. Seasonal spraying of houses, using pesticide which has a residual effect up to three months. This step happens in three stages (Table 3.3) (Figure 3.5):
 - A. First stage in TQ between August and October.
 - B. Second stage in TQ and TA between November and January.
 - C. Third stage in TQ and TA between February and April.
- 3. Space Spraying (Fogging), vehicle-mounted machines and hand-held sprayers used for outdoor (ULV and fogging) and indoor (ULV) coverage to control any unusual increase in cases or vectors. These occur in the affected and nearby areas.
- Intersectoral Collaboration, A joint committee meetings with related department in the region (e.g. Water & Electricity Department, Agriculture, Education, and Municipality addition to the Ministry of Culture and Information).

N _a A		Time	First stage	Second stage	Third stage
NO. A	Ireas	Time	Aug - Oct	Nov – Jan	Feb - Apr
1 7	TQ	9 months	Yes	Yes	Yes
2 7	TA	6 months	No	Yes	Yes

Table 3.3: Seasonal spray of houses in Asir Region

TQ: Tehama of Qahtan TA: Tehama of Asir

Malaria cases in Asir have decreased sharply over the last 15 years. The Vector Control Administration in Asir gave five main reasons for this decrease (Dr A. Abdoon 2007 pers. Comm.):

- 1. Climate change, as after 1998 rainfall has declined during these years, therefore the cases decreased.
- 2. Improved facilities and an increased number of professionals, technicians and workers.
- 3. Good immigration control to free this region from illegal refugees who are coming from endemic areas carrying this illness.
- Cooperation with The Vector Control Administration in the neighbouring Jazan Region in the border areas.
- A massive spray against rift valley fever for more than six months which started in Asir and Jazan 2000. This was carried out by many related governmental agencies using airplanes, cars and manpower.

3.10 Chapter Rationale and Hypotheses

The influence of climate variables upon malaria is important to understand for a number of reasons:

- Based on the findings and understanding of how malaria is affected by weather it should be possible to design an early warning system from the weather patterns to predict incidence (Bouma & Van Derkaay, 1996). This would be useful for current health service planning, for example identifying periods when enhanced insecticide spraying is required.
- 2. If weather changes as a result of climate change then this data could help predict the impact of this change on malaria in Asir Region.
- 3. It can help clarify the important modes of transmission in the region.

This study tests the hypotheses that there are relationships between weather (maximum, mean and minimum temperature, rainfall and relative humidity) and malaria cases in Asir Region.

3.11 Data and Method

3.11.1 Sources of Data

Official letters were issued, instructing authorized personnel to collect health and weather data. A standard form was provided to the personnel, so that the data could be received in a standard format. The total number of malaria cases in the whole of Asir Region each month from the first of Jan 1995 to the end of Dec 2006 was reported. In addition, the total number of cases in each epidemic sector in Asir for the same period was reported separately. All the malaria data were obtained from the Vector Control Administration, Health Affairs Directorate in Asir. This department represents The Ministry of Health in this region. The data has been collected from all hospitals and primary health care centers (public or private) and also the Main Parasitology Laboratory in Abha (The Vector Control Administration 1995 -2006). All the reported malaria cases were confirmed by a positive blood film (mainly asexual forms). Dr A. Abdoon 2007 pers. Comm. states that more than 90% of reported cases are indigenous cases, however, there are some other indigenous cases from Asir were reported in other regions of Saudi Arabia or abroad. These malaria data needed to be associated with weather data for this region. Unfortunately it was only possible to obtain weather data for a few locations and this had a large influence on the design of the analysis. Three models were constructed. The first was an analysis of malaria in the whole of Asir, and the next two focused upon malaria in the two endemic sectors TA and TQ. Further details are provided below:

- Malaria cases in overall Asir Region against Khamise weather data (Maximum, mean and minimum temperature, rainfall and relative humidity, Station 1 Figure 3.4). Khamise is the second main city in the region, and covers the largest occupied sector in the middle of this region (MOH, 2006). Weather data were available for all months.
- 2. Malaria cases in TA (Muhail, Rejal-Alma, Marabah, and Majaredah). These sectors were grouped as they all had similar malaria seasonality. They were related to the maximum and minimum temperature of Majaredah (See station 3 Figure 3.4, data available for 132 out of 144 months) and rainfall of Rejal-Alma (See station 4 Figure 3.4), data available for 132 out of 144 months). No humidity data was available for this area
- 3. Malaria cases in TQ against maximum, mean and minimum temperature, relative humidity of Khamise (Station 1 Figure 3.4) and rainfall of Sarat-Abeda (Station 2 Figure 3.4, data available for 132 out of 144 months). These stations are outside the sector but were used because there were no weather stations within, so there was no alternative but to use nearby stations.

Weather data for Khamise were obtained from The Center of Information and Documentation in the Meteorological and Environmental Protection Section (MEPS) of the Ministry of Defence and Aviation in Saudi Arabia. Weather data for Sarat-Abeda, Rejal-Alma and Majaredah were obtained from the Ministry of Water and Electricity. Monthly averages were obtained by dividing the monthly totals by the number of days in each month.

3.11.2 Malaria Data Description

Monthly averages of malaria cases in the Asir Region between 1995 and 2006 (Figure 3.7) show transmission of the parasite throughout the year. However, there is a clear seasonality with peak incidence between December and April, with the highest average cases in March (298.5) followed by February (276.3 cases). The averages of cases during the summer are lower, with the lowest number of cases in July (37 cases) and there is a slight peak in September. The standard deviation bars on the graph indicate that there is a large variability from year to year in monthly cases of malaria.



Figure 3.7: Monthly averages of malaria cases in Asir Region (1995 - 2006)

Figure 3.8 shows the data for the two main endemic sectors in Asir (TA and TQ). There is a clear seasonality for these sectors at different time of the year. In TA there is a peak of incidence between December and April, with the highest average cases in March (262.1 cases) followed by February (226.6 cases). The average number of cases during the summer is lower, with the lowest number in August (9.9 cases). There is some seasonality in TQ with a slight peak in incidence between August and October. The highest average number of cases was in September (93.6 case) followed by October (44 cases). The average number of cases during the rest of the year is lower, with the lowest number in February (7.9 cases). The standard

deviation bars on the graph again indicate much variability from year to year in monthly malaria cases for both sectors.



Figure 3.8: Monthly averages of malaria cases in TA and TQ (1995 - 2006)

3.11.3 Temperature Data Description

The average maximum, mean and minimum temperature in Khamise city is shown in Figure 3.9. The highest average maximum and mean temperatures occur in June (32.3, 24.6^oC respectively). The highest average minimum temperature occurs during July (18.1^oC). The lowest average temperature occurs in January for maximum, mean and minimum temperature (21.7, 14.6 and 8.5^oC respectively).



Figure 3.9: Maximum, mean and minimum averages temperature in Khamise 1995 -2006

The averages maximum and minimum temperature for Marjaredah is shown in Figure 3.10. The highest averages maximum temperature occur in June (41.98^oC) and the highest minimum in July (29.55^oC). The lowest averages maximum and minimum temperature occur in January (31.85 and 20.34^oC respectively).



Figure 3.10: Maximum and minimum averages temperature in Majaredah 1995 -2006

3.11.4 Rainfall Data Description

The monthly average rainfall in Khamise City (Figure 3.11) shows peak rainfall between March and August with the highest occurring in April (40.5mm) followed by August (30.9mm). During the rest of the year rainfall is low with a minimum occurring in February (1.6mm). The standard deviation bars on the graph indicates extreme variability from year to year.



Figure 3.11: Monthly averages rainfall in Khamise City (1995 -2006)

The monthly average of rainfall in Rejal-Alma (Figure 3.12) shows seasonality with peak rainfall between December and April. The highest averages were in March (80.1mm) followed by April (64.4mm). The average rainfall during the summer and autumn are lower with the lowest averages occurring in October (13.5mm). A small peak occurs in August. The monthly averages rainfall in Sarat-abeda (Figure 3.12) shows seasonality with two peaks occurring during spring and autumn. The highest average rainfall was in August (34.4mm) followed by October (30.0mm). The averages during the summer and winter are lower with the lowest rainfall occurring in June (3.2mm). The standard deviation bars on the graph indicate that there is a large variability from year to year in monthly rainfall for both sectors.



Figure 3.12: Monthly averages rainfall in Rejal-Alma and Sarat-Abida 1995 -2006

3.11.5 Relative Humidity Data Description

The monthly average relative humidity in Khamise City (Figure 3.13) shows a peak during the months of November to March with the highest percentage occurring in Jan (66.26%). The lowest percentage of humidity was in September (37.13%).



Figure 3.13: Monthly averages relative humidity in Khamise 1995 – 2006

3.11.6 Statistical Analysis Data Preparation

The analysis to examine the relationship between monthly cases of malaria and the selected weather data proceeded in a number of stages.

A histogram of the distribution of all malaria cases in Asir Region was constructed and this showed a positively skewed distribution, with a majority of lower values and a few somewhat higher ones. This was confirmed through descriptive statistics. This may cause problems when we come to perform regression analysis. We therefore took the natural logarithm (adding 1 to zero values) for malaria cases to produce Ln malaria. The distribution of this is shown in Figure 3.14 and descriptive statistics are presented in Table 3.4. These demonstrate that taking the natural logarithm of malaria largely corrected this problem. For similar reasons the natural logarithm was taken of malaria cases in TA and TQ.

In terms of the independent variables, rainfall of Khamise City also demonstrated a skewed distribution. Therefore, we took the natural logarithm of rainfall (adding 1 to zero values) to get (Ln rainfall). Figure 3.15 shows a histogram of rainfall after transformation. Table 3.5 presents the descriptive statistic of this distribution. The same action was been taken for rainfall in TA and TQ.



Figure 3.14: Frequency of natural logarithm of malaria (Ln.M) in overall Asir Region (1995 – 2006)

In	Ν	Minimum	Maximum	Mean	Std. Deviation	Skev	vness
Malaria 1	144	00	774	2.4	1 75	Statistic	Std. Error
	144	.00	1.14	3.4	1.75	.223	.202

Table 3.4: Descriptive statistics for Ln malaria of overall Asir



Figure 3.15: Frequency of logarithm rainfall (Ln. R) in Khamise (1995 – 2006)

Ln	N	Minimum	Maximum	Mean	Std. Deviation	Ske	ewness
Rainfall	144	0	5 2	1 70	1 55	Statistic	Std. Error
	144	0	5.5	1.72	1.55	0.27	0.20

Table 3.5: Descriptive statistics of Ln rainfall in Khamise

The data were adjusted for effects that could bias the results. The long term trend in Ln malaria in Asir region is shown in Figure 3.16, and the long term in Ln malaria in TA and TQ are shown in Figure 3.17. Over the study period the total number of malaria cases indicates a sharp decline from the end of the nineties with a peak of cases in 1998. In order to control for this effect the natural logarithm of malaria cases were put into a regression against time (each month numbered from 1 to 144). The unstandardised residual from this model represent the natural logarithm of malaria cases with the long term trend in incidence removed. This was performed on the malaria data for the whole of Asir region, TA and TQ.



Figure 3.16: Yearly Ln malaria in Asir Region (1995 - 2006)



Figure 3.17: Yearly Ln malaria in TA and TQ sectors (1995 - 2006)

3.11.7 Statistical Analysis

The literature review has demonstrated that the relationship between malaria and weather may vary according to the time of the year. There may also be a lag in the relationship between weather and malaria. Therefore, the analysis proceeded in stages.

In the first stage scatter plots were created of the detrended natural logarithm of malaria (DT Ln malaria) for every month against each of the weather variables. The purpose of this analysis was to determine which months should be grouped together based upon relationships with the weather variables (i.e. group consecutive months with positive or negative relationships with weather together). Unfortunately, this method proved difficult to implement because it was often unclear which months should be grouped. Also the relationship between weather and malaria may be lagged and so theoretically the analysis should be repeated for every lagged weather variable. Therefore this method was not taken further (for results see Appendix A.1), and replaced with one where the annual malaria time series was examined and groups of months determined based upon the trends in this graph (i.e. months of high incidence during the winter). For overall Asir this procedure led to three separate time periods which were April to August, September to November and December to March (See Figure 3.7). TA was separated into two time periods April to August and September to March (See Figure 3.8). TQ was separated into April to September and October to March (See Figure 3.8).

For each of time periods linear regressions of DT Ln Malaria against each of the lagged weather variables were created (lagged up to 6 months). We then examined these relationships and combined the weather from the two months with the strongest association. In most cases the strongest relationships were between DT Ln Malaria and the weather in the current and previous month. Therefore, for consistency these two lags were chosen in all cases. Because these models contained data from multiple months it became important to control for seasonality. Therefore, all models were analyzed incorporating dummy variables for each of the individual months. Finally in these analyses it was not possible to assume that the malaria rate in one month is unrelated to the malaria rate in the next month, and this is known as autocorrelation. Therefore the malaria in the previous month was added into these models as an independent variable. Due to the low numbers of cases in some months and the problems that this might cause in a regression an ordinal logistic regression of Ln malaria cases was additionally applied. This was achieved by recoding Ln malaria into four categories of increasing malaria incidence. These were then used in an ordinal logistic regression to produce an analysis that is more robust to the distribution of the data.

Multiple regression analysis was examined in the final stage of the statistical analysis. This method describes the effect of the two or three weather data together against detrended logarithm of malaria cases. The purpose of this test is to identify which weather variable has more influence upon malaria cases than others.

All of the data were analyzed using SPSS v.16 / v.18 for Windows and Microsoft Excel 2003 / 2007.

3.12 Results

3.12.1 Malaria with Maximum Temperature in Asir

Linear regressions and ordinal logistic regressions of DT Ln malaria against maximum temperature are presented for the whole year in Table 3.6 and April to August model in Table 3.7. Models for the rest of year (September to November and December to March) for this variable and all similar variables which do not have any significant relationship are not presented in this chapter but appear in AppendixA.2. In these tables 0P represents the models where the independent variables are not lagged, 1P represents the independent variables lagged by 1 month and so on. The results for the combined lags are presented on the final line of each table. In the analysis we also investigated weather instead of using the average maximum temperature each month it would be better to use the maximum temperature from any one day in the month. Such an analysis was performed (See AppendixA.3) but as the results were no different from using the average maximum temperature each month this was not taken any further.

		Li	inear re	egressio	on			С	rdinal	logist	ic	
Model	Ma tempe on	ax. rature ly	M tempe w seaso	Max. Emperature with easonality Max. temperature with auto correlation& seasonality		ax. rature auto ation& nality	Max. temperature only		re Max. temperature with seasonality		Max. temperature with auto correlation& seasonality	
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	104	.001	108	0.166	080	.195	140	.002	720	.001	750	.034
1 P	109	.001	132	.096	.080	.193	155	.001	738	.001	.515	.091
2 P	092	.001	176	.025	.095	.125	122	.008	678	.001	201	.497
3 P	044	.041	158	.044	037	.558	091	.041	891	.001	907	.011
4 P	.013	.548	163	.039	064	.315	026	55	788	.001	291	.363
5 P	.061	.005	176	.036	065	.303	.022	.620	915	.001	690	.040
6 P	.095	.001	165	.040	063	.323	.056	.213	109	.001	845	.021
0-1 P	118	.001	167	.075	114	.120	158	.001	-1.13	.001	929	.028

 Table 3.6: DT Ln malaria and maximum temperature in Asir (Whole Year)

0P: Current month. P: previous month. Cof.: Coefficients. P: P value (= or<). 0-1 P: Average of 0P+1 P

		Li	near re	gressi	on			(Ordinal	logist	ic	
Model	Ma temper on	ıx. rature ly	Ma temper wi seasor	ax. rature th nality	Ma temper with correla seasor	ux. rature auto ution& nality	Ma temper on	ıx. rature ly	Ma tempe wi season	ux. rature th nality	M tempe with correl seaso	ax. erature auto ation& nality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
<mark>0 P</mark>	<mark>200</mark>	<mark>.00</mark>	<mark>200</mark>	<mark>.038</mark>	<mark>151</mark>	<mark>.032</mark>	<mark>28</mark>	<mark>.041</mark>	<mark>70</mark>	<mark>.035</mark>	<mark>-1.02</mark>	<mark>.069</mark>
<mark>1 P</mark>	<mark>138</mark>	<mark>.001</mark>	<mark>212</mark>	<mark>.013</mark>	<mark>120</mark>	<mark>.050</mark>	<mark>196</mark>	<mark>.028</mark>	<mark>-1.06</mark>	<mark>.002</mark>	<mark>886</mark>	<mark>.049</mark>
2 P	099	.001	201	.015	069	.274	134	.090	-1.09	.001	419	.315
3 P	098	.001	117	.021	062	.279	175	.045	-1.12	.001	789	.087
4 P	092	.039	109	.146	031	.578	251	.056	851	.002	310	.406
5 P	.014	.831	166	.040	086	.149	410	.025	-1.37	.001	-2.05	.035
6 P	.138	.004	133	.140	.006	.932	129	.326	-1.59	.001	-2.05	.036
<mark>0-1 P</mark>	<mark>175</mark>	<mark>.001</mark>	<mark>292</mark>	<mark>.006</mark>	<mark>191</mark>	<mark>.014</mark>	<mark>247</mark>	<mark>.027</mark>	<mark>-1.38</mark>	<mark>.002</mark>	<mark>-1.49</mark>	<mark>.034</mark>

Table 3.7: DT Ln malaria and maximum temperature in Asir (April to Aug)

The results indicated that in Asir over the whole year malaria cases appear to be negatively associated with maximum temperature in the current and previous month. However, in the linear regression, not all these associations were significant. When only the April to August period is examined, maximum temperature in the current and previous month is significantly associated with malaria incidence in all variants of the model.

3.12.2 Malaria with Maximum Temperature in Tehama of Asir (TA)

Linear regressions and ordinal logistic of DT Ln malaria against maximum temperature in TA are presented for the whole year and April to August in Tables 3.8 and 3.9.

		Li	near re	gressi	on			C	Ordinal	logisti	ic	
Model	Ma temper on	ax. rature ly	Ma temper wi seasor	ix. rature th nality	Max. temperature with auto correlation& seasonality		Max. temperature only		Max. temperature with seasonality		Max. temperature with auto correlation& seasonality	
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	115	.001	.121	.128	038	.506	079	.092	064	.718	.046	.908
1 P	164	.001	.037	.639	048	.399	098	.037	39	.823	.205	.628
2 P	170	.001	061	.443	087	.125	121	.011	223	.215	666	.083
3 P	126	.001	103	.193	061	.287	082	.080	295	.107	500	.202
4 P	046	.079	108	.173	036	.527	027	.556	375	.050	308	.363
5 P	.051	.052	056	.486	.021	.720	.012	.800	501	.013	456	.062
6 P	.131	.001	.056	.487	.095	.097	.050	.277	.335	.076	.238	.474
0-1 P	156	.001	.124	.215	08	.399	101	.044	091	.684	161	.746

Table 3.8: DT Ln malaria and maximum temperature in TA (Whole Year)*

* Maximum temperature data: available for 132out of 144 months

		Li	near re	gressi	on			C	Ordinal	logisti	ic	
Model	Ma temper on	ıx. rature ly	Ma temper wi seasor	ux. rature th nality	Ma temper with correla seasor	ax. rature auto ation& nality	Ma temper on	ıx. rature ly	Ma temper wi seasor	ux. rature th nality	Ma tempe with correla season	ax. rature auto ation& nality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	199	.021	128	.292	006	.944	080	.665	.029	.925	052	.912
1 P	186	.001	.001	.997	087	.347	077	.435	001	.996	.019	.973
2 P	149	.001	117	.250	115	.119	098	.218	375	.187	736	.103
3 P	147	.001	258	.005	150	.035	107	.173	614	.046	101	.035
4 P	159	.001	194	.036	075	.290	113	.270	506	.068	.271	.475
5 P	070	.381	207	.039	063	.415	367	.047	-1.04	.008	-6.55	.034
6 P	.173	.001	050	.629	.143	.077	033	.757	529	.082	.428	.396
0-1 P	240	.001	.106	.500	075	.523	099	.477	.022	.954	035	.956

Table 3.9: DT Ln malaria and maximum temperature in TA (April to Aug)

In the TA sector malaria cases are not consistently associated with maximum temperature in the current and previous month. This result applies to the whole year model and to the April to August model.

3.12.3 Malaria with maximum temperature in Tehama of Qahtan (TQ)

Linear regressions and ordinal logistic regressions of DT Ln malaria against maximum temperature in TQ are presented for the whole year in Table 3.10 and April to September in Table 3.11.

		Li	near re	gressi	on		Î	0	Ordinal	logisti	ic	
Model	Ma temper on	ux. rature ly	Ma temper wi seasor	ux. rature th nality	Ma temper with correla season	ux. rature auto ation& nality	Ma temper on	ax. rature ly	Ma tempe wi seasoi	ux. rature th nality	Ma tempe: with correla seasor	ax. rature auto ation& nality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	021	.422	114	.260	032	.693	092	.039	611	.002	487	.040
1 P	.038	.151	083	.420	021	.791	042	.341	478	.013	207	.379
<mark>2 P</mark>	<mark>065</mark>	<mark>.014</mark>	<mark>243</mark>	<mark>.018</mark>	<mark>192</mark>	<mark>.016</mark>	<mark>012</mark>	<mark>.784</mark>	<mark>696</mark>	<mark>.001</mark>	<mark>679</mark>	<mark>.008</mark>
3 P	.076	.002	309	.002	162	.037	.012	.786	792	.001	526	.031
4 P	073	.003	266	.006	090	.263	.021	.636	843	.001	608	.020
5 P	.044	.079	294	.002	129	.092	.030	.500	794	.001	487	.064
6 P	.007	.798	299	.002	119	.126	.016	.722	908	.001	721	.010
0-1 P	.005	.844	148	.223	044	.644	078	.096	817	.001	533	.064

 Table 3.10: DT Ln malaria and maximum temperature in TQ (whole year)

		Li	near re	gressi	on			C	Ordinal	logisti	c	
Model	Ma tempe on	ax. rature ly	Ma tempe wi season	ax. rature th nality	Ma tempe with correla season	ax. rature auto ation& nality	Ma tempe on	ax. rature ly	Ma tempe wi season	ax. rature th nality	Ma tempe with correla season	ax. rature auto ation& nality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
<mark>0 P</mark>	<mark>082</mark>	<mark>.031</mark>	<mark>375</mark>	<mark>.025</mark>	<mark>306</mark>	<mark>.009</mark>	<mark>085</mark>	<mark>.192</mark>	<mark>714</mark>	<mark>.025</mark>	<mark>-1.50</mark>	<mark>.013</mark>
<mark>1 P</mark>	<mark>089</mark>	<mark>.045</mark>	<mark>317</mark>	<mark>.034</mark>	<mark>241</mark>	<mark>.022</mark>	<mark>032</mark>	<mark>.079</mark>	<mark>712</mark>	<mark>.013</mark>	<mark>598</mark>	<mark>.043</mark>
<mark>2 P</mark>	<mark>.074</mark>	<mark>.077</mark>	<mark>337</mark>	<mark>.012</mark>	<mark>249</mark>	<mark>.008</mark>	<mark>023</mark>	<mark>.730</mark>	<mark>839</mark>	<mark>.002</mark>	<mark>986</mark>	<mark>.036</mark>
3 P	.082	.127	363	.005	170	.073	022	.711	-1.08	.001	843	.038
4 P	.094	.133	291	.012	101	.276	058	.440	932	.001	622	.105
5 P	.058	.468	362	.002	205	.004	.198	.104	-1.10	.001	-1.08	.063
6 P	083	.259	363	.005	109	.239	271	.039	-1.32	.001	-1.08	.079
0-1 P	<mark>056</mark>	<mark>.038</mark>	<mark>484</mark>	<mark>.009</mark>	<mark>381</mark>	<mark>.003</mark>	<mark>073</mark>	<mark>.144</mark>	<mark>-1.05</mark>	<mark>.006</mark>	<mark>-1.38</mark>	<mark>.020</mark>

Table 3.11: DT Ln malaria and maximum temperature in TQ (April to September)

The results of TQ indicated that malaria cases are not consistently associated with temperature over the whole year. When the April to September period is focused upon maximum temperature in the current and previous month is significantly negatively associated with malaria in all but one variant of the model.

3.12.4 Malaria with Mean Temperature in Asir

Linear regressions and ordinal logistic regressions of DT Ln malaria against mean temperature in Asir are presented for April to August model in Table 3.12.

		Li	near re	gressi	on			0	Ordinal	logisti	ic	
Model	Me tempe on	an. rature ly	Me tempe wi seasor	an. rature th nality	Me tempe with correla season	an. rature auto ation& nality	Me tempe on	an. rature ly	Me tempe wi seasor	an. rature th nality	Me tempe with correla season	an. rature auto ation& nality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
<mark>0 P</mark>	<mark>243</mark>	<mark>.001</mark>	<mark>324</mark>	<mark>.016</mark>	<mark>235</mark>	<mark>.015</mark>	<mark>366</mark>	<mark>.021</mark>	<mark>-1.52</mark>	<mark>.003</mark>	<mark>-2.53</mark>	<mark>.017</mark>
<mark>1 P</mark>	<mark>157</mark>	<mark>.001</mark>	<mark>317</mark>	<mark>.008</mark>	<mark>183</mark>	<mark>.038</mark>	<mark>236</mark>	<mark>.023</mark>	<mark>-2.52</mark>	<mark>.001</mark>	<mark>-2.7</mark>	<mark>.011</mark>
2 P	111	.001	286	.013	088	.323	149	.095	-1.55	.001	718	.255
3 P	102	.002	134	.244	.012	.887	151	.113	-1.28	.003	356	.563
4 P	110	.024	088	.518	037	.687	165	.223	-1.06	.015	023	.974
5 P	054	.470	053	.693	015	.875	232	.244	-1.88	.001	-1.26	.060
6 P	.198	.001	025	.874	.012	.916	.068	.622	-1.27	.022	484	.535
<mark>0-1 P</mark>	<mark>205</mark>	<mark>.001</mark>	<mark>463</mark>	<mark>.002</mark>	<mark>303</mark>	<mark>.006</mark>	<mark>309</mark>	<mark>.018</mark>	<mark>-3.42</mark>	<mark>.001</mark>	<mark>-6.39</mark>	<mark>.009</mark>

Tables 3.12: DT Ln malaria and mean temperature in Asir (April to August)

The results indicated that in Asir over April to August malaria cases appear to be significantly negatively associated with mean temperature in the current and previous month for the linear regression and the ordinal logistic models. This correlation is highly significant in all variant of the model.

3.12.5 Malaria with Mean Temperature in TQ

Linear regressions and ordinal logistic regressions of DT Ln malaria against mean temperature in TQ are presented for April to September model in Table 3.13.

		Li	near re	gressi	on			(Ordinal	logisti	ic	
Model	Me temper on	an. rature ly	Me tempe wi seasor	an. rature th nality	Mean. temperature with auto correlation& seasonality		Me tempe on	an. rature ly	Mean. temperature with seasonality		Mean. temperature with auto correlation& seasonality	
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
<mark>0 P</mark>	<mark>.013</mark>	<mark>.885</mark>	<mark>545</mark>	<mark>.018</mark>	<mark>516</mark>	<mark>.001</mark>	<mark>172</mark>	<mark>.238</mark>	<mark>-1.95</mark>	<mark>.001</mark>	<mark>-4.17</mark>	<mark>.002</mark>
<mark>1 P</mark>	<mark>.080</mark>	<mark>.166</mark>	<mark>472</mark>	<mark>.022</mark>	<mark>379</mark>	<mark>.008</mark>	<mark>041</mark>	<mark>.657</mark>	<mark>-1.54</mark>	<mark>.001</mark>	<mark>-1.48</mark>	<mark>.032</mark>
<mark>2 P</mark>	.092	.057	<mark>616</mark>	<mark>.001</mark>	<mark>379</mark>	<mark>.006</mark>	015	.942	<mark>-1.92</mark>	<mark>.001</mark>	<mark>-1.75</mark>	<mark>.016</mark>
3 P	.113	.006	351	.067	065	.643	.007	.918	-1.24	.002	355	.508
4 P	.129	.008	302	.100	174	.204	03	.970	973	.008	582	264
5 P	.190	.011	212	.290	186	.152	08	.948	871	.025	262	.685
6 P	.023	.710	319	.140	260	.064	101	.442	-1.39	.003	-2.07	.026
<mark>0-1 P</mark>	<mark>.069</mark>	<mark>.348</mark>	<mark>754</mark>	<mark>.004</mark>	<mark>650</mark>	<mark>.001</mark>	<mark>090</mark>	<mark>.443</mark>	<mark>-2.96</mark>	<mark>.001</mark>	<mark>-5.11</mark>	<mark>.003</mark>

 Table 3.13: DT Ln malaria and mean temperature in TQ (April to September)

The results indicated that in TQ during April to September, malaria cases appear to be highly negatively associated with mean temperature in the current and previous month for all models except those including the temperature variables only.

3.12.6 Malaria with Minimum Temperature in Asir

Linear regressions and ordinal logistic regressions of DT Ln malaria against minimum temperature in Asir are presented for April to August model in Table 3.14.

		Li	near re	gressi	on			C	Ordinal	logisti	ic	
Model	Mi temper on	n. rature ly	Mi tempe wi seasor	in. rature th nality	Mi temper with correla season	n. rature auto ation& nality	Mi tempe on	in. rature ly	Mi tempe wi seasoi	n. rature th nality	Mi temper with correla season	n. rature auto ation& nality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
<mark>0 P</mark>	<mark>251</mark>	<mark>.001</mark>	<mark>288</mark>	288 .044		.082	314	.050	<mark>-1.67</mark>	<mark>.019</mark>	<mark>-4.27</mark>	<mark>.031</mark>
<mark>1 P</mark>	<mark>164</mark>	<mark>.001</mark>	<mark>341</mark>	<mark>.038</mark>	<mark>264</mark>	<mark>.024</mark>	<mark>208</mark>	<mark>.010</mark>	<mark>-2.29</mark>	<mark>.001</mark>	<mark>-7.22</mark>	<mark>.020</mark>
2 P	121	.001	221	.134	075	.488	113	.246	588	.200	.036	.958
3 P	098	.007	.127	.312	023	.789	069	.476	.197	.603	1.85	.020
4 P	095	.042	.103	.373	095	.042	012	.924	.365	.304	.357	.544
5 P	.064	.340	.162	.125	.075	.329	.075	.666	008	.981	140	.774
6 P	.214	.001	.093	.417	082	.337	.255	.097	.412	.254	.657	.256
<mark>0-1 P</mark>	<mark>208</mark>	<mark>.001</mark>	<mark>498</mark>	<mark>.026</mark>	<mark>392</mark>	<mark>.012</mark>	<mark>263</mark>	<mark>.050</mark>	<mark>-3.21</mark>	<mark>.001</mark>	<mark>-10.7</mark>	<mark>.025</mark>

 Table 3.14: DT Ln malaria and minimum temperature in Asir (April to August)

The results in Asir indicated that during April to August, malaria cases appear to be significantly negatively associated with minimum temperature in the current and previous month for linear regression and ordinal logistic models.

3.12.7 Malaria with Minimum Temperature in TA

Linear regressions and ordinal logistic regressions of DT Ln malaria areas against minimum temperature in TA are presented for April to August in Table 3.15.

		Li	near re	gressi	on			C	Ordinal	logisti	с	
Model	Mi tempe on	n. rature ly	Mi tempe wi seasor	n. rature th nality	Mi tempe with correla seasor	n. rature auto ation& nality	Mi tempe on	in. rature ly	Mi tempe wi seasor	n. rature th nality	Mi tempe with correla seasor	n. rature auto ation& nality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	216	.009	.116	.298	100	.255	.043	.811	.450	.152	338	.440
1 P	187	.001	.019	.810	090	.305	049	.639	.201	.499	915	.133
2 P	172	.001	073	.617	048	.657	040	.679	.485	.218	2.086	.047
3 P	.1.59	.001	073	.617	048	.657	040	.679	.485	.218	2.086	.047
4 P	122	.043	.037	.805	027	.806	.004	.975	.366	.357	.011	.987
5 P	.071	.552	192	.215	213	.056	057	.817	157	.689	-1.14	.162
6 P	.338	.001	.062	.767	.221	.149	.147	.426	.535	.324	1.71	.073
0-1 P	231	.001	.088	.494	119	.224	030	.826	.408	.234	694	.208

 Table 3.15: DT Ln malaria and minimum temperature in TA (April to August)

In the TA sector malaria cases are not consistently associated with minimum temperature in the current and previous month for the April to August period.

3.12.8 Malaria with Minimum Temperature in TQ

Linear regressions and ordinal logistic regressions of DT Ln malaria against minimum temperature in TQ are presented for April to September in Table 3.16.

		Li	near re	gressi	on			C	Ordinal	logisti	c	
Model	Mi tempe on	n. rature ly	Mi tempe wi seasor	in. rature th nality	Mi tempe with correla season	n. rature auto ation& nality	Mi tempe on	in. rature ly	Mi tempe wi seasor	in. rature th nality	Mi tempe with correla seasor	in. rature auto ation& nality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
<mark>0 P</mark>	<mark>.004</mark>	<mark>.971</mark>	<mark>757</mark>	<mark>.021</mark>	<mark>541</mark>	<mark>.020</mark>	<mark>112</mark>	<mark>.472</mark>	<mark>-2.48</mark>	<mark>.001</mark>	<mark>-3.52</mark>	<mark>.020</mark>
<mark>1 P</mark>	<mark>.115</mark>	<mark>.073</mark>	<mark>569</mark>	57 .021 69 .040		<mark>.032</mark>	<mark>.005</mark>	<mark>.962</mark>	<mark>-1.41</mark>	<mark>.011</mark>	<mark>787</mark>	<mark>.356</mark>
2 P	.109	.030	805	.001	378	.037	.012	.879	-1.55	.003	925	.236
3 P	.142	.001	.098	.642	.297	.065	.059	.426	.049	.890	1.15	.071
4 P	.151	.002	.056	.764	141	.312	.073	.388	.278	.402	037	.946
5 P	.250	.001	.262	.126	.087	.445	.166	.171	.449	.158	.838	.186
6 P	.125	.116	.122	.485	130	.267	.099	.455	.229	.463	425	.434
<mark>0-1 P</mark>	<mark>.094</mark>	<mark>.251</mark>	<mark>994</mark>	<mark>.007</mark>	<mark>724</mark>	<mark>.005</mark>	<mark>035</mark>	<mark>.787</mark>	<mark>-3.05</mark>	<mark>.001</mark>	<mark>-3.09</mark>	<mark>.032</mark>

Table 3.16: DT Ln malaria and minimum temperature in TQ (April to September)

The results indicated that in TQ only over the April to September model malaria cases appear to be significantly negatively associated with minimum temperature in the current and previous month for both linear regression and ordinal logistic method but not when only the temperature was included in the model.

3.12.9 Malaria with Rainfall in Asir

Linear regressions and ordinal logistic regressions of DT Ln malaria against rainfall in Asir are presented for April to August model in Table 3.17.

		Li	near re	gressi	on			C	Ordinal	logisti	ic	
Model	Rain on	ıfall ly	Rair wi seasor	nfall th nality	Rain with correla season	nfall auto ation& nality	Rair on	nfall ly	Rair wi seasor	nfall th nality	Rair with correla seasor	nfall auto ation& nality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
<mark>0 P</mark>	<mark>.227</mark>	<mark>.003</mark>	<mark>.217</mark>	<mark>.007</mark>	<mark>.115</mark>	<mark>.050</mark>	<mark>.414</mark>	<mark>.069</mark>	<mark>.447</mark>	<mark>.036</mark>	<mark>.717</mark>	<mark>.031</mark>
1 P	<mark>.169</mark>	<mark>.014</mark>	<mark>.170</mark>	<mark>.015</mark>	<mark>.107</mark>	<mark>.036</mark>	<mark>.500</mark>	<mark>.021</mark>	<mark>.652</mark>	<mark>.010</mark>	<mark>.568</mark>	<mark>.018</mark>
2 P	116	.053	.043	.552	.023	.668	.108	.499	.518	.039	.079	.806
3 P	100	.110	.055	.482	.029	.607	.083	.612	.376	.141	.470	.218
4 P	076	.241	019	.804	013	.823	.025	.880	.088	.708	375	.302
5 P	.045	.559	.023	.773	.022	.697	.202	.308	.221	.346	399	.319
6 P	028	.768	008	.927	016	.791	.081	.735	.130	.611	.323	.477
<mark>0-1 P</mark>	<mark>.440</mark>	.001	<mark>.400</mark>	<mark>.001</mark>	<mark>.244</mark>	<mark>.002</mark>	<mark>1.10</mark>	<mark>.003</mark>	<mark>1.27</mark>	<mark>.002</mark>	<mark>4.58</mark>	<mark>.025</mark>

Table 3.17: DT Ln malaria and Ln rainfall in Asir (April to August)

Only the April to August model is presented. Rainfall in the current and previous month is highly positively significantly associated with malaria incidence in both the ordinal logistic and the linear regression models.

3.12.10 Malaria with rainfall in TA

Linear regressions and ordinal logistic regressions of DT Ln malaria cases against rainfall in TA are presented for whole year in Table 3.18 and April to August model in Table 3.19.

		01201					<u> </u>					
		Li	near re	gressi	on			C	rdinal	logisti	c	
Model	Rain on	ıfall ly	Rain wi seasoi	ıfall th nality	Rain with correla season	nfall auto ation& nality	Rain on	nfall ly	Rain wi seasoi	ıfall th nality	Rain with a correla seasor	fall auto tion& nality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
<mark>0 P</mark>	<mark>.247</mark>	<mark>.001</mark>	<mark>.210</mark>	<mark>.001</mark>	<mark>.093</mark>	<mark>.015</mark>	<mark>.462</mark>	<mark>.001</mark>	<mark>.532</mark>	<mark>.001</mark>	<mark>.541</mark>	<mark>.037</mark>
<mark>1 P</mark>	<mark>.220</mark>	<mark>.001</mark>	<mark>.222</mark>	<mark>.001</mark>	<mark>.057</mark>	<mark>.045</mark>	<mark>.509</mark>	<mark>.001</mark>	<mark>.585</mark>	<mark>.001</mark>	<mark>.553</mark>	<mark>.023</mark>
2 P	.180	.003	.219	.001	.060.	.145	.468	.001	.562	.001	.301	.177
3 P	.177	.004	.224	.001	.058	.159	.555	.001	.737	.001	.694	.007
4 P	.095	.121	.150	.004	.13	.750	.482	.001	.643	.001	.273	.226
5 P	.030	.628	.101	.059	0.02	.962	.374	.001	.504	.001	.020	.925
6 P	.015	.810	.067	.215	.024	.541	.505	.001	.676	.001	.643	.013
<mark>0-1 P</mark>	<mark>.321</mark>	<mark>.001</mark>	<mark>.288</mark>	<mark>.001</mark>	<mark>.119</mark>	<mark>.013</mark>	<mark>.714</mark>	<mark>.001</mark>	<mark>.829</mark>	<mark>.001</mark>	<mark>.802</mark>	<mark>.014</mark>

Table 3.18: DT Ln malaria and with Ln rainfall in TA (Whole year)*

*Rainfall data: available for 132out of 144 months

		Li	inear re	gressio	on			C	Ordinal	logisti	с	
Model	Rainfa	ll only	Rainfa seaso	ll with nality	Rainfa au correla season	ll with to ation& nality	Rainfa	ll only	Rainfa seaso	ll with nality	Rain with correla seasor	ifall auto ition& nality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
<mark>0 P</mark>	<mark>.269</mark>	<mark>.001</mark>	<mark>.249</mark>	<mark>.001</mark>	<mark>.119</mark>	<mark>.028</mark>	<mark>.288</mark>	<mark>.005</mark>	<mark>.277</mark>	<mark>.035</mark>	<mark>.617</mark>	<mark>.036</mark>
<mark>1 P</mark>	<mark>.254</mark>	<mark>.001</mark>	<mark>.215</mark>	<mark>.001</mark>	.069	.205	<mark>.696</mark>	<mark>.003</mark>	<mark>.695</mark>	<mark>.004</mark>	1.06	.069
2 P	.181	.011	.222	.001	.048	.396	.970	.001	1.06	.001	.769	.068
3 P	.158	.021	.176	.003	.033	.505	1.44	.002	1.71	.002	1.0	.035
4 P	.045	.497	.073	.233	.081	.086	1.16	.001	1.35	.002	.602	.187
5 P	.041	.549	.056	.369	025	.586	.953	.001	1.18	.003	1.16	.121
6 P	.058	.419	.031	.631	.102	.229	.868	.001	1.12	.002	2.34	.104
<mark>0-1 P</mark>	<mark>.396</mark>	<mark>.001</mark>	<mark>.337</mark>	<mark>.001</mark>	<mark>.155</mark>	<mark>.025</mark>	<mark>.742</mark>	<mark>.006</mark>	<mark>.736</mark>	<mark>.007</mark>	<mark>1.11</mark>	<mark>.048</mark>

Table 3.19: DT Ln malaria and with Ln rainfall in TA (April to August)

The results indicated that in TA over the whole year malaria cases appear to be significantly positively associated with rainfall in the current and previous month for both linear regression and ordinal logistic models. For the April to August period, malaria incidence is significantly associated with rainfall in the current and previous month in all variants of the model.

3.12.11 Malaria with Rainfall in TQ

Linear regressions and ordinal logistic regressions of DT Ln malaria against rainfall in TQ are presented for the whole year in Table 3.20 and April to September in Table 3.21.

	14	idie J.2	20. DI	Ln mu	iuriu u		unjun	m r Q	whole	e yeur)		
		Li	inear re	gressio	on			C	rdinal	logistic	с	
Model	Rainfa	ll only	Rainfa seasor	ll with nality	Rainfa au correla season	ll with to ation& nality	Rainfa	ll only	Rainfa season	ll with nality	Rain with a correla seasor	fall auto tion& nality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
<mark>0 P</mark>	<mark>.377</mark>	<mark>.001</mark>	<mark>.388</mark>	<mark>.001</mark>	<mark>.238</mark>	<mark>.001</mark>	<mark>.654</mark>	<mark>.001</mark>	<mark>.808</mark>	<mark>.001</mark>	<mark>.771</mark>	<mark>.001</mark>
<mark>1 P</mark>	<mark>.329</mark>	<mark>.001</mark>	<mark>.346</mark>	<mark>.001</mark>	<mark>.0128</mark>	<mark>.041</mark>	<mark>.605</mark>	<mark>.001</mark>	<mark>.802</mark>	<mark>.001</mark>	<mark>.554</mark>	<mark>.006</mark>
2 P	.268	.001	.287	.001	.077	.202	.628	.001	.767	.001	.519	.006
3 P	.094	.165	.150	.028	.031	.571	.368	.002	.472	.001	.081	.658
4 P	015	.813	.021	.751	075	.161	.276	.019	.331	.010	.093	.571
5 P	.026	.696	.014	.836	.001	.994	.362	.003	.395	.003	.345	.048
6 P	038	.596	056	.403	063	.216	.206	.074	.278	.033	.073	.667
<mark>0-1 P</mark>	<mark>.494</mark>	<mark>.001</mark>	<mark>.496</mark>	<mark>.001</mark>	<mark>.271</mark>	.001	1.02	.001	<mark>1.3</mark>	.001	<mark>1.10</mark>	<mark>.001</mark>

Table 3.20: DT Ln malaria and Ln rainfall in TQ (whole year)*

*Rainfall data: available for 132out of 144 months

		Li	inear re	gressio	on			(Drdinal	logistic	2	
Model	Rainfa	ll only	Rainfa seasoi	ll with nality	Rainfa au correla seasor	ll with to ation& nality	Rainfa	ll only	Rainfa seaso	ll with nality	Rain with correl seaso	nfall auto ation& nality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof	Р
<mark>0 P</mark>	<mark>.439</mark>	<mark>.001</mark>	<mark>.481</mark>	<mark>.001</mark>	<mark>.319</mark>	<mark>.001</mark>	<mark>.733</mark>	<mark>.001</mark>	<mark>.929</mark>	<mark>.001</mark>	<mark>1.36</mark>	<mark>.015</mark>
<mark>1 P</mark>	<mark>.411</mark>	<mark>.001</mark>	<mark>.477</mark>	<mark>.001</mark>	<mark>.243</mark>	<mark>.003</mark>	<mark>.611</mark>	<mark>.001</mark>	<mark>.889</mark>	<mark>.001</mark>	<mark>.857</mark>	<mark>.046</mark>
2 P	.230	.028	.330	.002	.064	.449	.350	.042	.523	.013	.125	.691
3 P	.080	.444	.196	.071	.052	.519	.176	.279	.251	.176	.568	.123
4 P	045	.649	060	.549	139	.053	.047	.769	.036	.834	.466	.120
5 P	.001	.998	079	.440	010	.874	.141	.385	.097	.579	.08	.980
6 P	181	.064	235	.116	146	.920	.054	.730	.079	.643	.497	.175
<mark>0-1 P</mark>	<mark>.576</mark>	<mark>.001</mark>	<mark>.626</mark>	<mark>.001</mark>	<mark>.391</mark>	<mark>.001</mark>	<mark>1.02</mark>	<mark>.001</mark>	<mark>.141</mark>	<mark>.001</mark>	<mark>1.58</mark>	<mark>.012</mark>

 Table 3.21: DT Ln malaria and rainfall in TQ (April to September)

The results in TQ over the whole year and April to September model indicate that malaria cases appear to be significantly positively associated with rainfall in the current and previous month for both linear regression and ordinal logistic models.

3.12.12 Malaria with Relative Humidity in Asir

Linear regressions and ordinal logistic regressions of DT Ln malaria against relative humidity in Asir are presented for the whole year in Table 3.22 and April to August model in Table 3.23.

		Li	inear re	gressio	on			С	rdinal	logisti	с	
Model	Hum on	idity ly	Hum wi seaso	idity th nality	Hum with correla seaso	idity auto ation& nality	Hum on	idity ly	Humi wit seasor	idity th nality	Humi with a correlat season	dity auto cion & ality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
<mark>0 P</mark>	<mark>.031</mark>	<mark>.001</mark>	<mark>.029</mark>	<mark>.036</mark>	<mark>.027</mark>	<mark>.013</mark>	<mark>.036</mark>	<mark>.023</mark>	<mark>.032</mark>	<mark>.032</mark>	<mark>.132</mark>	<mark>.024</mark>
<mark>1 P</mark>	<mark>.039</mark>	<mark>.001</mark>	<mark>.051</mark>	<mark>.001</mark>	<mark>.036</mark>	<mark>.001</mark>	<mark>.047</mark>	<mark>.005</mark>	<mark>.059</mark>	<mark>.032</mark>	<mark>.104</mark>	<mark>.075</mark>
2 P	.029	.001	.039	.005	.010	.408	.032	.047	.030	.371	.050	.385
3 P	.013	.082	.045	.001	.023	.046	.021	.185	.083	.021	.186	.005
4 P	007	.369	.033	.020	.007	.536	003	.838	.077	.029	.042	.464
5 P	.021	.007	.042	.003	.024	.035	.010	.504	.091	.012	.074	.175
6 P	030	.001	.024	.101	004	.743	012	.453	.086	.016	.029	.605
<mark>0-1 P</mark>	<mark>.044</mark>	<mark>.001</mark>	<mark>.059</mark>	<mark>.001</mark>	<mark>.047</mark>	<mark>.001</mark>	<mark>.053</mark>	<mark>.004</mark>	<mark>.070</mark>	<mark>.034</mark>	<mark>.180</mark>	<mark>.013</mark>

Table 3.22: DT Ln malaria and relative humidity in Asir (Whole Year)

		Li	inear re	gressio	on			С	rdinal	logisti	с	
Model	Hum on	idity ly	Hum wi seasor	idity th nality	Hum with correla seaso	idity auto ation& nality	Hum on	idity ly	Humi wit seasor	idity th nality	Humi with a correlat season	dity auto cion & ality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
<mark>0 P</mark>	<mark>.041</mark>	<mark>.004</mark>	<mark>.021</mark>	<mark>.022</mark>	<mark>.013</mark>	<mark>.015</mark>	<mark>.082</mark>	<mark>.009</mark>	<mark>.090</mark>	<mark>.039</mark>	<mark>.230</mark>	<mark>.029</mark>
<mark>1 P</mark>	<mark>.050</mark>	<mark>.001</mark>	<mark>.036</mark>	<mark>.034</mark>	<mark>.023</mark>	<mark>.043</mark>	<mark>.100</mark>	<mark>.006</mark>	<mark>.204</mark>	<mark>.003</mark>	<mark>.212</mark>	<mark>.043</mark>
2 P	.030	.005	.024	.155	.004	.767	.065	.039	.180	.006	.078	.409
3 P	.039	.001	.036	.026	.016	.181	.062	.077	.103	.057	.168	.055
4 P	.036	.018	.026	.123	.001	.998	.049	.234	.062	.232	010	.903
5 P	.017	.271	.042	.007	.023	.043	.021	.605	.054	.269	.019	.794
6 P	023	.015	.030	.057	.006	.631	013	.582	.051	.304	.056	.453
<mark>0-1 P</mark>	<mark>.066</mark>	<mark>.001</mark>	<mark>.046</mark>	<mark>.034</mark>	<mark>.029</mark>	<mark>.048</mark>	<mark>.135</mark>	<mark>.005</mark>	<mark>.241</mark>	<mark>.004</mark>	<mark>.387</mark>	<mark>.017</mark>

Table 3.23: DT Ln malaria and relative humidity in Asir (April to August)

The results in Asir over the whole year indicate that malaria cases appear to be significantly positively associated with relative humidity in the current and previous month. In the April to August model all the associations remain significant

3.12.13 Malaria with Relative Humidity in TQ

Linear regressions and ordinal logistic regressions of DT Ln malaria against relative humidity in TQ are presented for the whole year in Table 3.24 and April to September in Table 3.25.

		Li	inear re	gressi	on			Ċ	Drdinal	logisti	ic	
Model	Hum on	idity ly	Hum wi seasor	idity th nality	Hum with correla season	idity auto ation& nality	Hum on	idity ly	Hum wi seasor	idity th nality	Humi with a correlat season	dity uto ion & ality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
<mark>0 P</mark>	<mark>.097</mark>	<mark>.040</mark>	<mark>.013</mark>	<mark>.035</mark>	<mark>.023</mark>	<mark>.033</mark>	<mark>.024</mark>	<mark>.015</mark>	<mark>.060</mark>	<mark>.047</mark>	<mark>.091</mark>	<mark>.037</mark>
<mark>1 P</mark>	<mark>.081</mark>	<mark>.010</mark>	<mark>.038</mark>	<mark>.035</mark>	<mark>.033</mark>	<mark>.021</mark>	<mark>.013</mark>	<mark>.012</mark>	<mark>.074</mark>	<mark>.031</mark>	<mark>.058</mark>	<mark>.042</mark>
2 P	.008	.371	.054	.003	.032	.027	.003	.869	.074	.030	.050	.233
3 P	.011	.247	.066	.001	.036	.012	.010	.752	.102	.004	.090	.042
4 P	.011	.239	.046	.008	.010	.561	.004	.794	.086	.015	.044	.307
5 P	005	.558	.043	.014	.015	.266	.001	.925	.084	.018	.056	.205
6 P	.002	.797	.029	.101	.001	.942	.003	.852	.067	.052	.026	.539
<mark>0-1 P</mark>	<mark>.091</mark>	<mark>.031</mark>	<mark>.039</mark>	<mark>.038</mark>	<mark>.041</mark>	.017	<mark>.023</mark>	.018	<mark>.108</mark>	<mark>.042</mark>	<mark>.113</mark>	<mark>.031</mark>

Table 3.24: DT Ln malaria and relative humidity in TQ (whole year)

		L	inear re	egressi	on			C	Ordinal	logisti	ic	
Model	Hum on	idity ly	Hum wi seasor	idity th nality	Hum with correla seaso	nidity auto ation& nality	Hum on	idity ly	Humi wi seasor	idity th nality	Humi with a correlat season	dity auto tion & ality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	.24	.211	.009	.762	.023	.268	.036	.249	.095	.067	.149	.087
1 P	.001	.968	.025	.383	.031	.113	.045	.142	.139	.101	.167	.042
2 P	016	.349	.058	.041	.035	.086	.021	.440	.147	.008	.142	.110
3 P	018	.211	.080	.002	.035	.076	.025	.293	.173	.002	.129	.075
4 P	018	.292	.051	.038	.004	.0831	.028	.338	.100	.033	.014	.854
5 P	.010	.638	.057	.016	.018	.278	.039	.274	.08	.081	.032	.667
6 P	.017	.229	.046	.059	.001	.965	.028	.253	.110	.026	.089	.274
0-1 P	.018	.441	.026	.462	.042	.091	.061	.110	.189	.007	.254	.020

Table 3.25: DT Ln malaria and relative humidity in TQ (April to September)

The results over the whole year indicated that malaria cases in TQ appear to be positively associated with relative humidity in the current and previous month for both linear regression and ordinal logistic models. This is significant in all but the most basic form of the model. This correlation mostly disappears in the April to September model.

3.12.14 Multiple Regressions

In this section we will present the results from the normal and multiple regressions of DT Ln malaria cases against averages (current and one previous month) of three main weather data (Maximum Temperature, Rainfall and Relative Humidity). For temperature, we will present maximum temperate only in the multiple regressions as it has a similar seasonality to minimum and mean temperature and appeared to have the strongest associations with malaria.

A- Multiple Regressions of Malaria Cases in All Asir

The normal regressions for malaria in all Asir against maximum temperature, rainfall and relative humidity, in the current and previous month, for whole period are illustrated in Tables 3.26. Table 3.27 presents the multiple regressions for these variables.

The normal regressions for malaria in all Asir against maximum temperature, rainfall and relative humidity, in the current and previous month, for the April to August period are illustrated in Table 3.28. Table 3.29 presents the multiple regressions for these variables.

Aver. of				Linea	ar regre	ssion							Ord	inal log	istic			
0 P&1 P For	V	∕ariable Only	es	se	with easonali	ty	V COI Se	Vith aut rrelatior easonali	to n & ity	V	/ariable Only	es.	Se	With easonali	ity	V cor Se	Vith aut relation easonali	o 1 & ty
All year Cases	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.
Max.Tem. Only	118	.001	.189	167	.075	.187	114	.120	.507	158	.001	-92.1	-1.13	.001	-83.2	929	.028	-40.7
Rainfall Only	.133	.056	.039	.057	.553	.177	.071	.338	.506	.076	.590	-98.9	.576	.014	-91.0	.401	.286	-43.1
Humidity Only	.044	.001	.187	.059	.001	.250	.047	.001	.550	.053	.004	94.8	.070	.034	92.7	.180	.013	40.2

Table 3.26: Regressions of DT Ln malaria cases of Asir for maximum temperature, rainfall and relative humidity for whole year

Aver. of				Line	ear regr	ession							Orc	linal log	gistic			
0 P&1 P Variables	,	Variables Only	5	S	With easonalit	y	C	With auto orrelation Seasonalit) & У	,	Variable Only	8	S	With easonalit	y	correlat	With auto ion & Sea	o asonality
In Multiple Method	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.
Max.Tem	119	.001	183	178	.090	181	103	.206	503	203	.001	-	-1.05	.001	-	908	.045	-40 66
Rainfall	.007	.916	.105	025	.818	.101	.24	.777	.505	.327	.046	90.07	.175	.518	82.95	.057	.896	10.00
Max.Tem	070	.044	200	024	.813	220	.004	.962	5.40	145	.074	-	-1.17	.001	-	621	.184	-
Humidity	.023	.085	.200	.057	.003	.238	.048	.001	.542	.006	.842	92.11	.016	.036	83.10	.133	.037	39.11
Rainfall	.044	.001	200	087	.382	240	039	.609	517	.104	.473	-	.497	.049	-	.029	.947	40.21
Humidity	113	.070	.200	.066	.001	.249	.050	.001	.347	.053	.044	.94.51	.036	.019	90.72	.178	.021	-40.21
Max.Tem	50	.232		059	.582		011	.893		.282	.008		-1.11	.001		673	.186	
Rainfall	061	.426	.198	111	.297	.238	.047	.571	.540	.407	.031	- 89.68	.215	.445	- 82.81	179	.707	-39.04
Humidity	028	.059		.063	.002		.050	.001		.031	.374	02.00	026	.597	02.01	.140	.080	

 Table 3.27: Regressions of DT Ln Malaria cases of Asir with multiple weather (maximum temperature with rainfall and maximum temperature with relative humidity and for all of them together) for whole year

Aver. of				Linea	ar regre	ssion							Ord	inal log	istic			
0 P&1 P	V	/ariable	S		with		V COI	Vith aut	0 1 &	V	ariable	S		With		V	Vith aut	0
Variables		Only		se	easonali	ty	Se	easonali	ty		Only		se	easonali	ty	Se	asonali	ty
In Multiple Method	Cof.	Р	Adj. R ²	Cof.	Cof. P Adj. R ²		Cof.	Р	Adj. R ²	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.
Max.Tem. Only	175	.001	.318	292	.006	.316	191	.014	.643	247	.027	-38.1	-1.38	.002	-33.4	-1.49	.034	-16.1
Rainfall Only	.440	.001	.258	.400	.001	.403	.244	.002	.665	1.10	.003	-35.1	1.27	.002	-33.9	4.58	.025	-16.0
Humidity Only	.066	.001	.300	.046	.034	.276	.029	.048	.624	.135	.005	-36.0	.241	.004	-34.5	.387	.017	-14.7

Table 3.28: Regressions of DT Ln malaria cases of Asir for maximum temperature, rainfall and relative humidity for April to August

Aver. of				Linea	ar regre	ssion	•	U	v		<i>, ,</i>		Ord	inal log	istic			
0 P&1 P Variables	١	ariable only	S	se	with easonali	ty	V coi Se	Vith aut relation easonali	o 1 & ty	١	/ariable Only	¢S	se	With asonali	ty	V cor Se	Vith aut relatior easonali	o 1 & ty
In Multiple Method	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.
Max.Tem	140	.001	441	152	.152	415	117	.140	672	177	.140	-	-1.13	.018	-	-1.45	.052	-
Rainfall	.321	.001	.++1	.336	.002	.415	.197	.019	.072	.997	.008	33.95	1.01	.026	30.60	1.59	.050	13.71
Max.Tem	108	.050	224	338	.063	210	160	.086	(29	046	.801	-	-1.04	.038	-	999	.233	-
Humidity	.033	.122	.334	.019	.457	.310	.011	.552	.038	.150	.050	36.00	.148	.114	32.04	.307	.076	13.89
Rainfall	.256	.020	252	.394	.002	202	231	.015	650	.809	.046	-	.890	.055	-	.944	.238	-
Humidity	.046	.003	.335	.002	.924	.392	.05	.790	.039	.084	.125	33.86	.155	.101	32.47	.313	.071	13.89
-																		
Max.Tem	115	.004		184	.124		135	.132		082	.676		999	.049		-1.06	.216	
Rainfall	.342	.002	.432	.366	.003	.408	.214	.023	.667	.865	.044	- 33.77	.865	.075	- 30.32	1.12	.223	-
Humidity	.01	.713		.016	.544		.010	.659		.054	.546	55.17	.075	.465	50.52	.206	.291	15.10

 Table 3.29: Regressions of DT Ln malaria cases of Asir with multiple weather (maximum temperature with rainfall and maximum temperature with relative humidity and for all of them together) for April to August

The results of multiple regressions for malaria in Asir through whole year demonstrated that in no models were more than one weather variable significant. We conclude therefore that the best model for the whole year is that of DT Ln malaria against humidity. Humidity was positively significant in all variants of the normal regression and in most cases had a better goodness of fit (adjusted R^2 and log likelihood ratio) and is a stronger significance than any of the other weather variables.

Multiple regressions for malaria in Asir during April to August demonstrated that there were no models where all the weather variables were consistently significant. We therefore conclude that the final model should be a normal regression with one weather variable. Both rainfall and humidity are positively significantly in all variants of the normal regression. However, we have chosen the model including rainfall as the final model as in most cases the significances and model fit is better than the model containing humidity

B- Multiple Regressions of Malaria Cases in TA Sector

The normal regressions for malaria in TA areas against maximum temperature and rainfall, in the current and previous month, for whole period are illustrated in Tables 3.30. Table 3.31 presents the multiple regressions for these variables.

The normal regressions for malaria in TA areas against maximum temperature and rainfall for April to August period are illustrated in Tables 3.32. Table 3.33 presents the multiple regressions for these variables.

Aver. of				Line	ar regre	ssion							Ord	inal log	istic			
0 P&1 P For	١	/ariable only	2S	se	with easonali	ty	V COI Se	Vith aut rrelatior easonali	o n & ty	V	/ariable only	S	Se	With easonali	ty	V coi Se	Vith aut relation easonali	o 1 & ty
All year Cases	Cof. P Adj. R ²		Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.
Max.Tem. Only	156	.001	.239	.124	.215	.322	08	.399	.649	101	.044	-88.5	091	.684	-86.8	161	.746	-25.2
Rainfall Only	.321	.001	.162	.288	.001	.450	.119	.013	.670	.714	.001	-86.1	.829	.001	-62.3	.802	.014	-18.4

Table 3.30: Regressions of DT Ln malaria cases of TA for maximum temperature and rainfall for whole year

 Table 3.31: Regressions of DT Ln malaria cases of TA with multiple weather (maximum temperature and rainfall) for whole year

Aver. of				Line	ar regre	ssion							Ord	inal log	istic			
0 P&1 P Variables	۲	/ariable only	2S	S€	with easonali	ty	V coi Se	Vith aut rrelatior easonali	to n & ity	V	/ariable only	S	Se	With easonali	ity	V cor Se	Vith aut relation easonali	o 1 & ty
In Multiple Method	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.
Max.Tem	163	.001	508	.112	.262	540	001	.987	686	112	.095	-	.222	.468	-	.797	.303	-
Rainfall	.365	.001	.500	.341	.001	.540	.168	.001	.000	.827	.001	56.78	.875	.001	54.08	1.05	.031	11.32

Aver. of				Line	ar regre	ssion							Ord	inal log	istic			
0 P&1 P For	V	√ariable only	es	se	with easonali	ty	V COI Se	Vith aut rrelatior easonali	o n & ty	V	/ariable Only	es.	S€	With easonali	ty	V coi Se	Vith aut relation easonali	o ı & ty
April To Aug	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.
Max.Tem. Only	240	.001	.239	.106	.500	.274	075	.523	.612	099	.477	-37.8	.022	.954	-33.5	035	.956	-12.6
Rainfall Only	.396	.001	.351	.337	.001	.491	.155	.025	.669	.742	.006	-28.9	.736	.007	-28.7	1.11	.048	-7.33

Table 3.32: Regressions of DT Ln malaria cases of TA for maximum temperature and rainfall for April to August

Table 3.33: Regressions of DT Ln malaria cases of TA with multiple weather (maximum temperature and rainfall) for April to August

Aver. of				Line	ar regre	ssion							Ord	inal log	istic			
0 P&1 P Variables	۲	/ariable only	es	se	with easonali	ty	V coi Se	Vith aut rrelatior easonali	to n & ity	V	√ariable Only	es	Se	With easonali	ty	V COI Se	Vith aut rrelation easonali	o 1 & ty
In Multiple Method	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.
Max.Tem	199	.001	514	078	.610	513	089	.502	665	029	.870	-	543	.230	-	.443	.600	-
Rainfall	.351	.001	.514	.342	.001	.515	.156	.038	.005	.759	.005	25.80	.781	.006	24.73	1.11	.046	7.085

The results of multiple regressions for malaria in TA through whole year and in the April to August period no two weather variables were significant. Therefore, the regression containing only one weather variable was chosen as the final model. In these, rainfall was consistently positively significant in contrast to temperature. Additionally the models containing rainfall had better of fit. Therefore, the best model to describe DT Ln malaria in TA over the whole year and in the April to August period was one containing rainfall only.

C-Multiple Regressions of Malaria Cases in TQ Sector

The normal regressions for malaria in TQ against maximum temperature, rainfall and relative humidity, in the current and previous month, for whole period are illustrated in Tables 3.34. Table 3.35 presents the multiple regressions for these variables.

The normal regressions for malaria in TQ against maximum temperature, rainfall and relative humidity, in the current and previous month, for the April to August period re illustrated in Table 3.36. Table 3.37 multiple regressions for these variables.

Aver. of				Line	ar regre	ssion							Ord	inal log	istic			
0 P&1 P For	۲	/ariable only	S	se	with easonali	ty	V COI Se	Vith aut relatior easonali	.0 1 & .ty	V	/ariable only	S	Se	With easonali	ty	V cor Se	Vith aut relatior easonali	o 1 & ty
All year Cases	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.
Max.Tem. Only	.005	.844	.01	148	.223	.089	044	.644	.434	078	.096	-96.0	817	.001	-88.5	533	.064	-67.7
Rainfall Only	.494	.001	.278	.496	.001	.375	.271	.001	.549	1.02	.001	-69.1	1.3	.001	-63.1	1.10	.001	-47.5
Humidity Only	.091	.031	.01	.039	.038	.102	.041	.017	.465	.023	.081	-98.0	.108	.042	93.1	.113	.031	-67.6

Table 3.34: Regressions of DT Ln malaria cases of TQ for maximum temperature, rainfall and relative humidity for whole year

Aver. of				Linea	ar regre	ssion	•	0	v				Ord	inal log	istic			
0 P&1 P Variables	V	/ariable only	2S	Se	With easonali	ity	V coi Se	Vith aut relatior easonali	o 1 & ty	١	/ariable only	es.	Se	With easonali	ty	V cor Se	Vith aut relatior easonali	o 1 & ty
In Multiple Method	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.
Max.Tem	.040	.128	272	.095	.397	250	.055	.567	526	023	.688	-	569	.048	-	304	.370	-
Rainfall	.497	.001	.272	.507	.001	.539	.278	.001	.330	.986	.001	68.69	1.22	.001	60.63	1.04	.001	46.97
Max.Tem	030	.526	01	069	.610	002	.076	.476	155	061	.440	-	698	.009	-	320	.317	-
Humidity	017	.354	01	.031	.210	.095	.047	.016	.455	.008	.785	96.00	.051	.283	87.89	.086	.138	66.56
	1				1	1		1			1						1	
Rainfall	.505	.001	302	.490	.001	371	.254	.001	551	1.01	.001	-	1.29	.001	-	1.06	.001	-
Humidity	023	.221	.302	.010	.626	.571	.022	.209	.551	.004	.848	69.11	.103	.084	61.45	.096	.157	46.44
r		_																
Max.Tem	17	.692		.148	.240		.132	.214		029	.759		441	.160		124	.738	
Rainfall	.496	.001	.282	.505	.001	.359	.271	.001	.543	.987	.001	68.68	1.23	.001	- 60.10	1.04	.001	- 46.28
Humidity	028	.098		.022	.345		.033	.098		003	.932	00.00	.065	.312	00.10	.086	.247	+0.20

 Table 3.35: Regressions of DT Ln malaria cases of TQ with multiple weather (maximum temperature with rainfall and maximum temperature with relative humidity and for all of them together) for whole year

Aver. of				Line	ar regre	ssion							Ord	inal log	istic			
0 P&1 P For	١	/ariable only	S	Se	With easonali	ty	V COI Se	Vith aut relation easonali	o 1 & ty	١	/ariable Only	S	se	With asonali	ty	V cor Se	Vith aut relation easonali	o 1 & ty
April To Sep	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.
Max.Tem. Only	056	.038	.01	484	.009	.175	381	.003	.606	073	.144	-49.6	-1.05	.006	-44.6	-1.38	.020	-21.2
Rainfall Only	.576	.001	.338	.626	.001	.470	.391	.001	.697	1.02	.001	-35.7	.141	.001	-32.1	1.58	.012	-14.9
Humidity Only	.018	.441	.01	.026	.462	.091	.042	.091	.569	.061	.110	-48.3	.189	.007	44.8	.254	.020	-21.2

 Table 3.36: Regressions of DT Ln malaria Cases of TQ for maximum temperature, rainfall and relative humidity for April to September

Aver. of				Linea	ar regre	ssion					· • •		Ord	inal log	istic			
0 P&1 P Variables	١	ariable/ only	S	Se	With easonali	ty	V coi Se	Vith aut relatior easonali	o 1 & ty	١	/ariable Only	es	se	With asonali	ty	V cor Se	Vith aut relatior easonali	o 1 & ty
Multiple Method	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.
Max.Tem	.193	.001	440	108	.524	165	191	.133	704	.363	.043	-	196	.627	-	506	.462	-
Rainfall	.682	.001	.442	.601	.001	.405	.342	.001	.704	1.34	.001	33.64	1.37	.001	31.48	1.41	.035	14.62
Max.Tem	.044	.677	017	556	.010	169	361	.017	600	.220	.194	-	756	.075	-	982	.131	-
Humidity	.005	.886	017	027	.502	.108	.007	.803	.000	.126	.048	47.45	.128	.098	43.13	.176	.130	19.96
Rainfall	.664	.001	124	.629	.001	161	.379	.001	605	1.06	.001	-	1.47	.001	-	1.76	.018	-
Humidity	068	.001	.434	007	.820	.401	.019	.443	.095	021	.677	35.57	.257	.014	28.46	.345	.085	12.81
Max.Tem	.123	.138		159	.407		185	.198		.629	.014		.024	.964		364	.676	
Rainfall	.686	.001	.443	.597	.001	.459	.342	.001	.699	1.29	.001	31.90	1.47	.001	- 28.46	1.91	.026	12.73
Humidity	.033	.302		.021	.561		.003	.916		.160	.075	21.75	.260	.025	_00	.390	.093	12.1.5

Table 3.37: Regressions of DT Ln malaria cases of TQ with multiple weather (maximum temperature with rainfall and maximum temperature with relative humidity and for all of them together) for April to September
The results of multiple regressions for malaria in TQ through whole year and the April to September period demonstrated that no two weather variables were consistently significant. Therefore, the final models chosen included one weather variable only. Over the whole year and in the April to September period rainfall was clearly the most significant positive variable with better model fits than any of the other weather variables.

3.13 Discussion and Conclusion

Malaria is endemic in lowlands of Asir Region (Haddad, 1990), which can be divided into two areas; Tehama of Asir which includes foothills with coastal plain, and Tehama of Qahtan. This disease is less endemic in the coastal plain, but more endemic along the valleys and foothills (Abdoon and Alshahrani, 2003). This study tests the hypotheses that there are relationships between weather and malaria cases in this region, and examines the influence of climate variables upon malaria cases.

During this study several difficulties were encountered. The main one was the limited weather data for some selected areas in Asir, and missing information from within this data. This meant that it was not possible to produce models for each malaria endemic area in Asir. For example, we have combined several sectors for malaria data in TA including Muhail, Rejal-Alma, Marabah and Majaredah, into one group. Although all these areas are in the lowland (foothills) and have the same seasonality of malaria, it would have been preferable to examine each separately but the weather data was not available for each sector. It also would have been good to examine the impact of humidity upon malaria in TA but data was not available.

The situation in TQ was even more difficult as no weather stations were available within this area. Instead we had to use the temperature and humidity from Khamise and rainfall of Sarat-Abeda which were outside the sector boundaries. These stations were used because there were no weather stations within. Although these stations were chosen because of their close similarity to the weather in TQ, this can cause problems in the analysis if the variations in temperature, rainfall and humidity in TQ are different to those in the two chosen weather stations. In the results, malaria incidence was compared to temperature across Asir. The results presented little evidence that temperature is an important factor affecting variations in malaria cases. Temperature was significant in some of the original regressions but no temperature variables were significant in the final models. In all cases, other weather variables demonstrated stronger associations with malaria cases.

This lack of association was surprising because the optimum temperature for the vector of malaria (mosquito) to live and breed is 25 to 27 0 C and maximum temperature for both vectors and parasites, is 40°C (McMichael et al., 1996; Hana et al., 2001). In TA the average maximum temperature (Majaredah) reaches over 40 0 C between May and August. Therefore this high temperature could kill the mosquito, which may provide an unsuitable condition for the transmission of malaria. As temperature is not significant in the models, this suggests that there is a little variability in these high temperature has been shown to be associated with malaria in a study in Jazan Region (Southwest of Saudi Arabia) (Al-Jaser, 2006) which has almost similar conditions to the lowlands of Asir Region. In the literature review many other studies were identified which highlighted the importance of temperature to malaria transmission. Examples are Burkina Faso, (Ye et al., 2007), China (Bi P et al., 2003) and Bangladesh (Alam et al., 2008).

In TA and TQ and in overall Asir, the final model included rainfall only. We therefore conclude that rainfall is the strongest predictor of malaria in the summer period and over the whole year. However, because rainfall was not significant in the models for other times of the year, we conclude that the whole year result is driven by the summer result. The results indicate that less rainfall leads to a lower number of malaria cases. Figure 3.12 shows that rainfall is low during the summer period which may explain why it is not important at other times of the year. At other times of the year there may be enough rainfall for mosquito breeding. It is also interesting to note the strategy of malaria control in Asir (see table 3.3). This shows that no control activities happen between May and July in TQ and between May and October in TA, although the pesticides which are used in the third stage of seasonal spraying (February to end of April), have a residual effect for up to three months. This lack of

summer control may be another reason why associations with weather are only observed during this period.

These results are consistent with previous studies that have found that rainfall is one of the weather phenomena which play a significant role in malaria. Biologically, this is plausible as Oaks et al., (1991) mentioned that rainfall is one of the main factors affecting survival of anopheline mosquitoes by helping to provide suitable breeding conditions. Hence, the survival of anopheline (vector of malaria) may lead to an increase malaria cases. Pampana, (1969) discusses that too much rainfall, or rainfall accompanied by storm conditions, can flush away breeding larvae and so reduce survival. There was no evidence of this in the present study. Flash flooding does not usually happen in Asir Region, and the region has a lot of dams which can hold the water for long time. These dams can provide suitable breeding conditions for the vector (Dr A. Abdoon 2007 pers. Comm.). Not only the amount and intensity of rainfall, but also the time of year, whether in the wet or dry season, affects malaria survival. This is seen in this study with rainfall only being important in the dry summer period. Also low amounts of rainfall may not be appropriate for the mosquito as it needs breeding sites (Briet (b) et al., 2008).

The importance of rainfall highlighted in this study is consistent also with other studies the same zone of the world (The Afro-tropical Zone) and in Saudi Arabia. In Sudan, Hamad et al., (2002) showed that, the abundance of mosquitoes in this area was positively related to monthly rainfall. In Ethiopia, Teklehaimanot et al., (2004) noted that rainfall was followed within a short time by an increase in malaria cases. In SW Saudi Arabia an unusual increase of malaria happened at the beginning of 1996 coinciding with heavy rainfall when many valleys were flooded (A1-Abdullatif et al., 1996). In Asir region in 2008 the peak in malaria cases coincided with high unexpected rainfall (MOH, The Statistical Book, 2009). During last three months in 2008 most of Asir Region received high unexpected rainfall which may have led to that increase of malaria cases (Dr A. Abdoon 2007 pers. Comm.).

The time lag used in this study (rainfall in the current and previous month) is also consistent with previous research. Bi P et al., (2003) mentioned that there was a highly positive correlation between monthly rainfall and malaria in Shuchen County, China lagged at one month. In Sri Lanka Briet (a) et al., (2008) found a correlation between malaria and rainfall lagged 0-4 months lag. In Like Bufundi, Kilibwoni, East Africa, a peak in rainfall during April-May was followed by a peak in malaria occurrence one to 2 months later in June (Kristan et al., 2008). The current study in Asir Region is consistent with these studies for the lag time between rainfall and malaria incidence.

The results also indicated that relative humidity appears to be of importance for transmission of malaria as there is some evidence that when the humidity is high, there are more cases of malaria. However, humidity was only significant in the model for all Asir which combines malaria data from all over the region. Due to data limitations it was not possible to test its effect in TA. In TQ it was not significant. The possible importance of humidity is consistent with Al-Jaser, (2006) which found a significant positive correlation between relative humidity and malaria cases in Jazan. Pampana, (1969) noted that if the average monthly relative humidity is below 60%, it is believed that the life of the mosquito is so shortened that there is no malaria transmission. Alam et al., (2008) demonstrated that, the optimal relative humidity for mosquito survival is 60%, and a high relative humidity lengthens the life of the mosquito. In the current study, averages relative humidity were much lower than 60% over the summer which may explain humidity's importance. In India, Bhattacharya et al., (2006) noted that the average relative humidity range (55 to 80 per cent) remains correlated with malaria transmission.

In summary, this study showed that weather variations factors play an important role in affecting malaria cases in Asir Region during the summer months. At other times of the year, weather variations do not appear important. Rainfall was the most important factor associated with malaria in TA and TQ with greater rainfall in the current and previous month leading to more malaria cases. Relative humidity may also play a role in malaria transmission. Although this study shows that weather has an influence upon malaria in Asir, it is important to recognize that malaria is becoming less of a problem in general but still a significant problem in some endemic areas. Overall, the study shows that the total number of yearly malaria cases has declined sharply, more than 90% from the end of the nineties compared to the end of the study period. The reasons for this were provided in the literature review.

Chapter 4

Influence of Climate Variables upon Leishmaniasis Cases in Asir Region, Saudi <u>Arabia</u>

4.1 Introduction

Leishmaniasis is an important public health problem in several countries and is considered to be the vector borne disease with the second highest number of affected people in the world after malaria (Chaves et al., 2006). It has been estimated by World Health Organization that globally there are 14 million people who are infected, and each year about 2 million new cases occur (WHO, 2009; Faulde et al, 2009). Annually there are over 70,000 deaths (WHO, 2009). This disease is considered the 6th most important endemic diseases in the world by the World Health Organisation (WHO, 2007). In addition to immediate health consequences leishmaniasis is associated with about 2.4 million disabilities globally. One of the most common forms, cutaneous leishmaniasis, is rarely fatal but it can lead to tissue destruction, scarring and serious visible impairment. These can lead to social stigma due to deformation of the face with permanent scarring (Murray et al., 2005; Reithinger et al., 2007; CFSPH, 2009). Other forms can affect the mucous membranes which cause disfiguring nose lesions (mucocutaneous leishmaniasis), or damage internal organs (visceral leishmaniasis) (CFSPH, 2009).

Leishmaniasis is a group of zoonotic diseases transmitted to humans and animals by the bite of phlebotomine sandflies (Figure 4.1). The infective agents are several species of flagellated protozoan parasites (Reithinger et al., 2001). There are more than 1000 known species of sandfly in the world, of which only 70 species are known to transmit diseases such as Leishmaniasis (Murray et al., 2005). Only female phlebotomine sandflies suck blood leading to Leishmaniasis. Male sandflies rarely bites humans and do not transmit this disease. Sandflies transmit other viral and bacterial diseases, but they are particularly the vector of leishmaniasis (Reithinger et al., 2001). Sandflies feed on mammals, birds and both sexes takes sugar meals to maintain their daily activity (TDR, 2002). Sandflies do not make noise when they fly and are also small being around one-third the size of a mosquito. Their bite is also painless (CDC, 2010). This means that people may be unaware of sandflies and their biting.



Figure 4.1: Phlebotomine sandfly (Kamiura, A. and Reforma, L, 2007)

There are several different clinical forms of leishmaniasis, but the most common are; cutaneous leishmaniasis (CL), visceral leishmaniasis (VL) and mucocutaneous leishmaniasis (MCL) (Figure 4.2). In addition to these three major forms, there are others such as diffuse cutaneous leishmaniasis, recidivans leishmaniasis, and post-kala-azar dermal leishmaniasis (Croft et al., 2006). CL causes skin sores caused by leishmania major. It is common in rural areas with dogs and bush mammals being reservoir hosts. Another cause of CL L tropica is more common in urban areas and leads to CL in humans and dogs. L. Aethiopica also leads to CL with hyraxes being the reservoir hosts (Wittner et al., 2000). One of the most important vectors of CL is Phlebotomus papatasi, a common human-biting sandfly species found all over North Africa, through Eurasia, and into India (Hanafi et al., 2007). VL affects the internal organs of the body such as spleen, liver, and bone marrow and it can be fatal. It is caused by leishmania infantum species and the common vector is phlebotomus sergenti, a known vector of phlebotomus Tropica (TDR, 2002). MCL is a complication of simple CL, and it involves the mucosa, soft tissue and cartilage of the respiratory system (WHO, 2009).

Leishmania transmission is sensitive to weather and climate conditions, and Cross et al., (1996) state that: one of the most important factors in leishmaniasis distributions are environmental conditions such as weather.



Figure 4.2: Clinical presentations of leishmaniasis (Narciso et al., 2007; Kamiura, A. and Reforma, L, 2007)

This chapter will start by providing an overview of the geographic distribution of leishmaniasis. It will then present information on the transmission of leishmania and its life cycle. How this illness may be prevented and treated will then be discussed. We will then present a literature review of leishmaniasis and how it is related to weather.

The analysis section of the chapter will begin by presenting information on leishmaniasis in Saudi Arabia and how it is currently controlled. The aims and hypothesis for the study will then be discussed. The data sources to be used for the study will then be presented followed by an overview of the methods. The results will then be presented which will analyze statistically the relationships with weather. The chapter will end with a discussion and conclusion.

4.2 Geographic Distribution

Leishmaniasis is distributed in around 90 countries of the world most of which are in the developing countries (16 developed countries and 74 developing countries), and there are approximately 350 million people are at risk of leishmaniasis in these areas (Schriefer et al, 2009; Cardenas et al, 2006). Tropical and subtropical countries are the most affected by this illness (WHO, 2000). The settings in which leishmaniasis is found ranges from rain forests in Central and South America to deserts in West Asia (WHO, 1994). In terms of the types of leishmaniasis, more than 90% of CL occurs in parts of Afghanistan, Algeria, Iran, Iraq, Saudi Arabia, Syria, Brazil and Peru (CDC, 2008; TDR, 2002), but it is found also in Mexico, Central America, and South America, southern Europe and particularly East and North Africa (WHO, 2007) (Figure 4.3). Currently Kabul is thought to have the highest incidence of CL in the world (76500 - 200,000 cases/year with a population of 3.5 - 5 million) (Reithinger et al., 2005). More than 90% of VL cases are found in India, Bangladesh, Nepal, Sudan, some parts of the Middle East and Brazil (WHO, 2007). Ninety percent of MCL occurs in Central and South America, Bolivia, Peru and Brazil, and in East of Africa (Costa, 2011).

Al-Hammad, (2003) states that the distribution and spread of leishmaniasis across the world is affected by several factors including: topography, population migration, hygiene factors, micro-environmental factors, host factors, occupational factors, weather factors and global warming.



Figure 4.3: Distribution of CL, MCL and VL in the World (Kamiura, A. and Reforma, L, 2007)

4.3 Leishmania Transmission and its Life Cycle

Transmission of leishmania is most often zoonotic (transmitted to humans from animals) where an animal is the reservoir hosts (such as dogs and small rodents). In the case of leishmaniasis sandflies act as the vector. It also can be anthropontic where man is the reservoir host and source of infections (TDR, 2002; Rey, 2007). Leishmaniasis in humans and other mammals is caused by 21 of the 70 species of pathogenic sandfly (Al-Amru, 2002). However this is disputed by Reithinger et al., (2007) who state that there are about 40 species or subspecies which are vectors. Another 40 species may also be involved in leishmaniasis transmission.

Figure 4.4 shows the life cycle of leishmaniasis according to the US Centre for Disease Control and Prevention (CDC, 2010). The cycle consists two host stages involving humans and sandflies. It starts when an infected female sandfly (phlebotomine) injects promastigotes into the skin of an animal or person during its blood meal. In the human stages these promastigotes transform into amastigotes and multiply in infected cells inside macrophages. They then affect different tissues depending on the specie of leishmania. In the sandfly stages, an uninfected sandfly takes its blood meal from an infected person or animal and then becomes infected when it ingests the amastigotes. The amastigotes transform into promastigotes and multiply in the midgut of the fly. The promastigotes divide in the midgut and transfer to the mouthparts, where the insect is ready to make another infection and thus the cycle begins again (CDC, 2010; Rey, 2007).

Sandflies are usually most active in twilight, the evening, and night-time hours (from dusk to dawn) and are less active during the hottest time of the day. However, they will bite if they are disturbed, such as when a person brushes up against the trunk of a tree where sandflies are resting (Al-Amru, 2002).



Figure 4.4: Life cycle of leishmaniasis (CDC, 2010)

In terms of transmission the female sandfly lays its eggs in the bark of trees, old buildings, in cracks in the walls of houses, in animal houses and in household rubbish. This because it needs the shelter and humidity provided by these environments for egg development. In terms of range the female sandfly covers an area from a few hundred to several hundred metres around its habitat during its search for blood (WHO, 2010).

Gubler et al., (2001) and Hunter, (2003) discuss how changes in temperature may influence the transmission of vector borne diseases. Possible influences include increases or decreases in vector survival, changes to the rate of vector population growth and also changes in feeding behaviour. In addition changes to the susceptibility of vectors to pathogens, alterations to incubation periods and lastly changes in the seasonality of vector activity and pathogen transmission are all possible. Also they mention that rainfall changes can impact the risk of transmission of vector-borne disease. Increased rainfall can increase vegetation and allow expansion in the population of vertebrate hosts whilst flooding may eliminate habitat for both vectors and vertebrate hosts. Finally flooding may force vertebrate hosts into closer contact with humans. Therefore a changing climate is likely to change transmission dynamics in many regions, e.g. increased temperature, abnormal rainfall or higher relative humidity. Leishmaniasis is perceived as the vector-borne disease most likely to be affected by global climate change (Al-Amru, 2002).

4.4 Previous Studies and Relationship between Sandflies or Leishmaniasis with Climate and Weather Conditions

There are only a few studies that look at the direct influence of weather upon leishmaniasis cases. However, there is good evidence of a close linkage between the abundance of sandflies and leishmaniasis cases (Amru, 2002; Al-Tawfiq et al., 2004; Al- Ibrahim et al., 2005). Therefore this section will try to present studies which examine the relationship between the environment and weather conditions with the abundances of both sandflies and leishmaniasis cases. In this section we will try to discuss studies from around the world before focussing upon studies conducted in the Middle east and finally in Saudi Arabia and the study area Asir Region.

4.4.1 Sandflies with Climate and Weather

Cross and Hyams, (1996) through research in Southwest Asia state that one of the most important factors for the distribution of the sandfly species *P. papatasi* (the main vector for leishmaniasis) and hence leishmaniasis distributions are environmental conditions such as weather. Through a review of the literature they state that temperature and humidity play an important role in the lifecycle and development stages of the sandfly. Adults and larvae are sensitive to high temperatures and low humidity. In laboratory experiments, all adult sandflies died within two hours at temperatures above 40°C. However, below this temperature, increasing temperature can shorten the incubation period and life cycle of the fly which could increase disease transmission. Temperatures below 10°C are unfavourable for sandfly survival. However, cold does not necessarily lead to death and under cold conditions as sandfly larvae have the ability to become dormant until conditions improve. In Southeast Asia the colder winter period is absent of sandflies with an increasing population at the beginning of spring. The number of sandfly survivors increase with increased relative humidity (Cross and Hyams, 1996).

Temperature may also affect the daily activity rates of sandflies. In terms of the daily activity, a study of P. papatasi in Iraq demonstrated that biting starts after sunset and increases gradually till the maximum activity in the middle of the night when the temperature is lower and the humidity higher. Biting activity in Central Iraq showed a decrease to 13% at 03:00 hr and stopped after sunrise (Cross and Hyams, 1996).

The researchers predicted the seasonal and geographical distribution of papatasi in this region based on 115 weather stations via an appropriate analysis model. Interestingly this model also simulated the effect of global warming. Currently leishmaniasis was endemic at 62% of the weather stations. However, under climate change they predict that a further 12% of stations could become endemic if the temperature increase by 1°C. An additional 15% of stations would become endemic if the increase was 3°C. If the increase is 5°C the additional percentage will be 10%. In addition to changes in endemic stations they also conclude that seasonal disease transmission could be extended throughout all months of year in 6% of locations with a 3°C rise in temperature. This figure will increase to 25% with a 5°C rise (Cross and Hyams, 1996).

Moving away from South East Asia, in Pondicherry, India Srinivasan et al., (1993) studied the population dynamics of P. papatasi. They selected three villages to monitor the density of sandflies for the period from March 1988 to February 1990 in different seasons; summer which is from March to June, rainy from July to October and winter from November to February. The findings showed a positive correlation between daily survival of sandflies and relative humidity and a negative correlation with temperature. The high temperature and low relative humidity during summer may have affected the survival of sandfly. The declining trend of sandflies population was observed in winter, when the temperature was low, and also during summer, when temperature was high. Rainfall was found to have a significant influence on the density of sandflies as the rainfall in June and July was followed by increase in the population of sandflies till October. However, a sudden decrease in sandfly numbers was observed in November and December which, the author's state may have been due to heavy rain destroying breeding sites.

In Rajasthan, India Singh, (1999) investigated the role of climatological factors in the seasonality of phlebotomine sandflies. They found a total of eight phlebotomine species in houses. The phlebotomines were found only at temperatures between 17 and 36°C. However the optimum temperature distribution varied with

species. For example the optimum range for *P. papatasi* was from 28 to 34°C, *P. sergenti* from 31 to 33°C. The species *Sergentomyia* are also vectors of leishmaniasis and the optimum range for *S. punjabensis* is from 27 to 30°C, *S. babu* from 28 to 33°C, *S. baghdadis* from 27 to 34°C and *S. christophersi* from 29 to 33°C. The maximum phlebotomine prevalence (39.4%) was recorded in September when the average temperature and RH were 32.1°C and 58.1%, respectively. The minimum phlebotomine prevalence was found in January (1.2%) having an average temperature of 20.8°C and 46.3% relative humidity. This result suggested that these species are well adapted to the conditions of high temperatures and low RH. The statistical analyses noted that temperature has a positive significant correlation with phlebotomine abundance.

In the Cukurova Plain, Turkey, Kasap et al., (2009) found that the maximum abundance of sandflies was recorded in the hottest season (26.9 to 31.2 °C) and also in the driest season when the relative humidity was comparatively low (46.6 to 56.8%) during the period from June to September. In terms of relative humidity the study contradicts with Cross et al., (1996) who state that the number of sandfly survivors increases with increased relative humidity. This contradiction could be due to different locations and also the fact that the increase in humidity occurs at the same time as high temperatures which are favourable to sandfly survival.

In Morocco, leishmaniasis is endemic and widespread throughout the country and is considered a serious public health problem (Guernaoui et al., 2006). Seasonal fluctuations of sandfly populations in the urban area of Marrakech, Morocco were studied by Boussaa et al., (2005). Sandflies in this area were collected, between October 2002 and September 2003. They found that the sandflies were active throughout the year, but that the major activity was in two periods: The first was in October and November with a peak on November and the second one, which had higher activity, was during the months of April–July with a peak in May. The maximum and minimum monthly temperatures in the two periods were between 23 and 36°C, and 11 and 19°C, respectively. The study suggested that male sandflies were more sensitive to temperature than females. They concluded that the seasonal patterns related to environmental factors such as topography and climate variables and specifically temperature in this study. The relationship between sandflies and altitude in the High-Atlas Mountains of Morocco was studied by Guernaoui et al, (2006). Sandflies were captured from 25 stations at various altitudes (400–1400m). It was noted that altitude can play a role in terms of the distribution of sandfly species and the density of each vector. The highest abundance for *P. papatasi* was at (400-599m) and the lowest at more than 1199m. However, the highest density for *P. perniciosus* was at (1000-1199m) with the lowest at (400-599m). It showed there were fewer of all species at altitude over 1199.

In a study carried out in Iran between May 1995 and May 1996 Yaghoobi et al., (1999) noted that the activity of P. papatasi starts from the last days of April or early May and extends to mid October. They found two peaks in abundance one in mid or late June and the second one in early or mid September. There was a decrease in sandfly density at the end of October which they attributed to the rains which can destroy or damage the breeding sites (Hunter, 2003). This decrease remained until the end of March which was attributed to the cold weather. There were no sandflies found in the area between November and March. Another study in the centre of Iran (Yaghoobi et al., 2001) demonstrated that sandflies in this area started to appear in late May and disappeared at the end of October. There are two peaks in the density curve of P. papatasi, one in late May or June and the second at the end of September as the active season in this time. A decrease of sandfly density was observed in the end of October and the researchers thought it is due to the rains. Also due to cold weather there was no sandflies during November to March.

The sandfly species pattern was studied in some parts in Argentina by Salomon et al., (2008). This study reports that the epidemic risk may be associated with climate. The activity season of the most abundant sandfly species begins from March to April (autumn), with its highest peak in November (spring). However during dry and cold weather in winter, only a few flies were found. The abundance of these species increased during the rainy season and decreased during the months of lower rainfall or higher temperatures.

Molina et al., (2008) found that in Colombia the seasonal density of the most abundant sandfly species were correlated with the average monthly rainfall, maximum temperature, and relative humidity. The mean monthly density of this species increased during the rainy months from April to October and decreased during the months without rainfall (January to March). A multiple linear regression between the most abundant species and rainfall, maximum temperature, and relative humidity showed a positive significant correlation for these species (P=0.001, 0.003 and 0.002 respectively).

4.4.2 Leishmaniasis with Climate and Weather

In a study of the seasonality transmission of leishmaniasis carried out in the state of Campeche, Mexico between February 1993 and March 1995. Narvaez et al., (2003) report that leishmaniasis transmission occurs from November to March. They concluded that the winter season with its high humidity and low temperature have ideal ecological conditions for transmission. If the incubation period (approximately 3 months) is taken into account the total number of leishmaniasis cases peak between March and July. Very few cases are reported between June and November.

In North-eastern Colombia, Cardenas et al., (2006) studied impact of climate variability in the occurrence of leishmaniasis. This study focuses on the impact of the El Nino Southern Oscillation between 1985 and 2002. During El Nino (dry seasons) in 1987, 1992–1994, 1997, and 2002, leishmaniasis cases increased, however during La Nina (wet season). In 1988–1989, 1995–1996, and 1998–2001 the number of cases decreased. The study concluded that there is evidence linking leishmaniasis to climate variables, and suggested that annual changes in weather conditions play an important role in varying the incidence of vector borne diseases including leishmaniasis. They also conclude that the occurrence of drought followed by rainfall is likely to increase leishmaniasis incidence. This is because drought can lead to waning immunity in humans and a consequent increase in cases when transmission conditions become favourable (Cardenas et al., 2006). This is supported by Gage et al., (2008) who state that the decrease in transmission in endemic areas after a reduction in rainfall may actually increase the occurrence of the disease due to a decline in the immunity of the populations in this area.

In the state of Bahia, Brazil, rainy seasons following the El-Nino led to increases in both the densities of sandfly and subsequent infection rates of leishmaniasis in high-risk populations (Franke et al., 2002). The authors state that the lag between leishmaniasis cases and the rainy season is around 2-6 months. In

addition the time from the beginning of symptoms to diagnosis approximately is 3 months. This study also reported that the number of positive cases increased for 2 years following droughts associated with El Nino. One result of these droughts was low vector densities which led to reductions in the immunity of the susceptible humans and lower sandfly populations in endemic areas.

Faulde et al., (2008) studied a CL outbreak in Mazar-e Sharif, northern Afghanistan. Case leishmaniasis data (Anthroponotic Cutaneous leishmaniasis ACL and Zoonatic ZCL) were collected in this area from March 2004 to March 2006. ACL is caused by Leishmania Tropica and mainly transmitted in this area by the sandfly Phlebotomus sergenti. ZCL is caused by Leishmania major. More than 95% of the CL cases were ZCL while less than 5% were ACL. The peak transmission of ZCL was in August and September and is correlated with dry and hot season. ACL transmission usually starts in November with the first case in December and it may be positive associated with the cold and wet winter. The result showed that ACL is endemic in this area with peak during February and March. There were no cases reported from September to November. The incubation period varies between leishmaniasis types and for example ACL the period was found to be 4 - 6 weeks. However it is 8 - 12 weeks for ZCL.

Cross et al., (1996) used weather data to predict the seasonal, geographic and distribution of *P. papatasi* in SW Asia. They hypothesised that the distribution of *P. papatasi* might be highly dependant on the temperature and relative humidity (environmental conditions). They designed a computer model with the occurrence of sandfly as dependent variables and temperature and relative humidity data as independent variables. They found that the greatest activity of sandfly may lead to a high risk of leishmaniasis infection during the months of spring and summer with positive association. As an extension to this study the researchers (Cross et al., 1996) studied the potential effect of global warming on the geographic and seasonal distribution of this species of sandfly. They found that higher temperatures could greatly increase both the geographic and seasonal distribution of sandfly. The geographic distribution may also extend to nearby areas where the sandfly populations are currently too small to maintain endemicity.

CL in some parts of Yemen was studied by Khatri et al., (2004). They suggest that altitude plays a big role in the distribution of CL as the majority of patients originated from the plateau and mountainous range (1800–2000m) of Hajjah city and a few patients from the adjacent regions.

4.4.3 Sandflies of Leishmaniasis with Climate and Weather in Saudi Arabia

In Saudi Arabia much research on sandflies and leishmaniasis has been carried out. Al-Amro., (2002) studied the geographical distribution of 24 sandfly species in Saudi Arabia and also the relationship between the seasonal insect density and the number of leishmaniasis cases. It demonstrates that Riyadh, Qasime (Central regions) and Alahsa Regions (Eastern Saudi Arabia) have the highest case numbers of CL. Jazan and Asir Regions have the highest case numbers of VL. The researchers report that the Asir Region, where this study will be focussed, has around 13 species of sandfly including *P. papatasi, P. bergeroti, P. chinesis P. arabicus, P. orientalis, P. saevus* and *P. sergenti*. The study also report that the highest daily activity period of the insects vary by species but that almost twilight is the best time especially if the weather condition are suitable (windless and cloudy).

The Ministry of Health in Saudi Arabia, 2000 reports a highly significant correlation between the density of sandfly and positive cases of CL. However, the lag between sandflies and illness is between two and five months, because the illness will occur after an incubation period following a sandfly bite (MOH, 2000). This lag depends upon the species of sandfly and on the environmental conditions. They also state that significant under-reporting of cases occurs as some patients do not go to any medical centre for treatment as they treat themselves. Other cases are only seen by medical centres when the disease has reached advanced stages.

El-badry et al., (2009) studied the distribution of sandfly in four surrounding areas of Al Madenah Region (Al-Nekheil), Saudi Arabia. The average temperature is high over summer and moderate during winter (August mean 38°C and January mean 25°C and). They collected 621 sandflies, from March 2006 to November 2007, and found six species of sandfly; *papatasi* represented more than 70% of the sandfly population. They noted that there is an increase of this insect from May to September due to the weather conditions.

An epidemiological study of CL over 46 years in eastern Saudi Arabia (Al-Tawfiq et al., 2004) presents evidence that the seasonal pattern of the disease correlates with the known activity of the lag time from a person being bitten to the development of skin lesions to the presentation of patients to physicians. There was a peak of sandfly activity in April and September followed by a peak of CL cases in September and January, and this lag explains the differences of these peaks.

One of the most important research studies on the vector of leishmaniasis in Asir Region was conducted by Ibrahim et al, (2005). A total of 1896 sandflies were collected from 3 topographically different zones during June 1999 to May 2001. They note that altitude has an effect on sandfly, as they found the density of the insect was higher in the lowland (foothills) compared to the highland (mountains). This may indicate that the activation of this insect and their development stages (eggs – larva - pupae – adults) prefer warm lowland conditions. Low temperature, fog and strong winds in the highlands may make these areas unsuitable for sandflies. This study indicated that the highest abundance of sandflies was recorded during spring and summer, and the lowest density was during November to February. They suggest that this is due to the low temperature, high relative humidity and strong winds during winter.

4.4.4 Literature Review Summary

Table 4.1 summarises the main studies (statistically) of the correlation between leishmaniasis or its vector and climate variables.

No	The author	Place	Dependent Variable	Independent Variable	The relationship & Lag time (month)	
1	Cross and Hyams, 1996	Southwest Asia	density of Sandfly	Temperature Humidity	Positive	
2	Srinivasan et al., 1993	Pondicherry, India	ondicherry, density of Humidity India sandflies Rainfall Rainfall		Negative Positive Positive 2-4 Negative 5-7	
3	Singh, 1999	Rajasthan, India	phlebotomines	temperature	Positive	
4	Kasap et al., 2009	Cukurova Plain, Turkey	abundance of sandflies	Temperature Humidity	Positive Negative	
5	Guernaoui et al., 2006	Morocco	abundance of sandflies	Temperature	Positive	
6	Yaghoobi et al., 1999	Iran	density of sandflies	Temperature	Negative	
7	Salomon et al., 2008	some parts in Argentina	abundance of sandflies	Rainfall Temperature	Positive	
8	Molina et al., 2008	Colombia	density of sandfly	Temperature Rainfall Humidity	Positive Positive 1-6 Positive	
9`	Cardenas et al., 2006	North-eastern Colombia	leishmaniasis cases	Rainfall	Negative	
10	Narvaez et al., 2003	Campeche, Mexico	L. incidence	Temperature Humidity	Negative 3 Positive 3	
11	Franke et al., 2002	Bahia, Brazil	Number of leishmaniasis cases	Rainfall	Positive 2-6	
12	Faulde et al., 2008	Mazar-eSharif, Afghanistan	Number of leishmaniasis cases	Temperature Rainfall	Negative Positive	
13	Al-Amro., 2002	Saudi Arabia	Number of CL cases	density of sandfly	Positive 2-5	
14	Al-Tawfiq et al., 2004	eastern Saudi Arabia	Number of CL cases	density of sandfly	Positive 1-5	
15	Ibrahim et al, 2005	Asir Region, Saudi Arabia	Density of sandfly	Temperature	Positive	

Table 4.1: Summary of the main studies relating to leishmaniasis

The literature review highlights a number of important points:

- A wide range of study designs have been used to investigate the relationships between leishmaniasis and weather. These range from purely observational (e.g. more leishmaniasis in summer), but very few conduct statistical analyses of leishmaniasis cases and associations with weather.
- 2. Numerous different climate variables have been used in previous studies including temperature, rainfall, wind and relative humidity.

- Most, but not all studies, showed that climate variables have an impact upon CL incidence. However, the impact of specific climate variables can be either positive or negative depending upon the geographical setting and season.
- Lagging the weather variables is an important feature of these studies. Studies in Saudi Arabia suggest a 2-5 month lag between weather and cases. This lag also depends on the type of leishmaniasis.
- 5. Many studies support these findings by showing that weather influences the abundance of the leishmaniasis vector (sandfly). Others found that the vector abundance has an effect upon the incidence of leishmaniasis.
- 6. Weather conditions are not the only factor which can affect Leishmaniasis. There are many other factors such as ecological and environmental conditions, social and economical determinants.
- 7. There have been few studies in Saudi Arabia focusing upon the influence of weather upon this disease.

Based on the previous studies, it is clear that there is an urgent need to study the influence of weather variables upon the presence of CL in Saudi Arabia, and in endemic areas such as Asir Region. The previous studies appear very location specific and so the results cannot be easily translated to Asir. Within Asir this also means that different models may have to be produced for different geographical areas. Building upon the literature review such a study should examine a wide range of weather variables and also consider that any relationships with weather may vary seasonally. Variables will need to be lagged to account for incubation periods and reporting lags. A statistical analysis seems the most appropriate study design and this may need to consider factors such as autocorrelation in these data. Few previous studies have conducted a rigorously statistical analysis of CL.

4.5 Sandfly and Leishmaniasis in Saudi Arabia

There are two main species of leishmaniasis in the country; CL and VL (Al-Amru, 2002). The Ministry of Health (MOH, 2006) showed that there were 3980 cases of leishmaniasis (CL and VL) in Saudi Arabia in 2005 and 4165 cases in 2004. Alkhawajah, (1998) states, that the most common manifestation of leishmaniasis in this country is CL. It is found all over the country but there is a great variation in the prevalence rate between different regions (MOH, 2008). In terms of geographical

distribution of CL during 2007 the region of Qaseem had the highest percentage of cases (25.9%) followed by Al-Ahsa (24.9%) and Madinah (18.8%). No cases have been reported in Northern region, Al-Jouf, Qurayyat and Qunfedah Regions (MOH, 2008). VL occurs in some parts of the south and south east of Saudi such as Jazan, Asir and Albahah Regions with a few cases in the rest of Saudi (Al-Amru, 2002).

The seasonal variation in CL in Saudi Arabia is presented in Figure 4.5. This figure demonstrates a peak of CL during the winter (December to March) and these four months account for 37 % of the total cases occurring each year. Another small peak occurs between July and October. The least number of cases was reported in May representing less than 4% of the annual total. During the period 2002-2006 the annual cases of CL were 4454, 3842, 4123, 3883 and 3602 respectively. In 2007 the incidence of CL was 13.9/100,000 population while it was 15.2/100,000 population in 2006 (MOH, 2008).



Figure 4.5 Monthly averages of CL cases in Saudi Arabia (2002-2006)

In terms of trends over time Figure 4.6 presents the number of CL cases in Saudi Arabia from 1983 to 2009. This figure shows a dramatic decrease of cases through out the last two decades, and cases are now less than a quarter of their levels in the 1980's.



Figure 4.6: The number of (CL) cases in Saudi Arabia (1983- 2009) (MOH, The Statistical Books 2000-2009; MOH, 2006)

There are only a few studies on the vector sandfly abundances in Saudi Arabia and one study has investigated the relationship between altitude and density of sandfly in this country. This is presented in Figure 4.7. This figure shows that the highest density of sandfly was for *P. papatasi* specie at 500 - 750m altitude followed by *P. sergenti* at the same altitude, then *P. orientals* at 1750 - 2000m. In the highest altitudes (>1750m) *P. orientals* and *P. arabicus* are the most common sandfly species.



Figure 4.7: The relationship between altitude and density of some species of sandfly in Saudi Arabia. Adapted from (Buttiker et al., 1983; Al-Amru, 2002)

In terms of the relationships between sandfly numbers and leishmaniasis Figure 4.8 plots the result of both the density of sandfly and positive cases of CL in Al-Dereya city (20 km from Riyadh) during October 1997 to February 1999 (Al-Amru, 2002), however, it shows no association between sandfly numbers and CL.



Figure 4.8: The correlation between density of sandfly and the number of CL cases in Al-Dereya City, Saudi Arabia. Adapted from (Buttiker et al., 1983; Al-Amru, 2002)

4.6 Leishmaniasis in Asir Region (Study Area)

This study will occur in the Asir Region which is located in SW Saudi Arabia. During the period of this study (1996 – 2007) there were few VL cases in all Asir Region (Table 4.2). Therefore this study will focus upon CL. This is more common in this region (see previous section) and occurs on the foothills and high plateau of Asir Region (Ibrahim et al., 2005). The distribution of this illness is influenced by little known geographical and climate factors (Al-Zahrani et al., 1989).

Table 4.2 Number of visceral leishmaniasis cases in all Asir Region (1996-2007)(VCA, MOH)

Year	96	97	98	99	00	01	02	03	04	05	06	07
No. of	1/	5	4	1	8	13	10	3	3	2	3	3
cases	14	5	4	4	0	15	10	5	5	4	5	5

4.7 Topographical Classification of Asir Region and Endemic Areas

Topographically, climatically and endemically this region can be divided into three main sectors for CL (Dr A. Abdoon 2007 pers. Comm.; MEPS, 2007; Gaafar et al., 1997) (Figure 4.9 and 4.10):

- Asir Sarawat Mountains (became a part of Highlands Group): A series of high mountains extending from the north of Asir Region to the south along the coastal plains of the Red Sea (2000 -~3000m above sea level). This area covers districts such as Abha, Khamise, Tanoma and Dhahran. The annual average temperature for this sector is 17.7^oC, average rainfall is 500mm (the highest rainfall in Saudi Arabia), and annual average relative humidity of 53.1%. These areas have the highest occurrence of leishmaniasis in Asir Region, and *P. arabicus* specie is the main vector.
- 2. Asir Plateau (became a part of Highlands Group): In the NE Asir with an elevation range of between 1000 and 2000m, decreasing to the east. This area covers districts such as Tathlith, Bishah and Tareeb. The annual average temperature for this sector is 19.5°C, with 250mm annual rainfall and an annual average relative humidity of 44.4%. This sector has many wadies which all flow eastward. It has_little leishmaniasis incidence and *P. arabicus* specie is the main vector.
- 3. Tehama (Lowlands Group became a part Tehama): Receives an annual rainfall of more than 350mm, and temperature ranges between 25°C and 49°C, and the relative humidity varies from 55% in summer to 70% in winter. This area can be divided into two sections:
 - a. Foothills such as Al-Majaredah, Muhail and Rejal-Alma, where the altitude is between 300 and 900m. This area has the highest occurrence of leishmaniasis after the highlands in Asir Region and *P. papatasi* specie is the main vector.
 - b. Red Sea Costal Plain (RSCP) such as Al-Gahma, Al-Berk, Al-Huraidah. The altitude in this area is between 0 and 200m. The lowest leishmaniasis occurs in the region, and *P. papatasi* specie is the main vector.



Figure 4.9: Topographic map of Asir Region including weather stations

4.8 Control of Leishmaniasis Transmission in Asir

The control methods against leishmaniasis can act to control the vector, the host (reservoir) or human behaviours which bring people into contract with the vector. The World Health Organisation has set up regional programmes to improve health systems to allow access to innovation (WHO, 2009). The control of leishmaniasis can be reached by breaking the transmission cycle of an agent (Al-Hammed, 2003). WHO, (2007) recommends two main interventions to control the vector; spraying insecticide indoors and use of insecticidal nets (Bed nets). Control of sandfly may occur inadvertently through other interventions. For example in 1950s and 1960s DDT was used in some part of the world for malaria control and was thought to have markedly reduced the sandfly population as well (Al-Hammed, 2003). A variety of insecticides can be used to control sandfly including DDT, Malathion, Propoxus, Fenitrothion, and synthetic pyrethroids. Insecticide control can be effective for adults of sandfly, pupae of sandfly and larva (Al-Hammed, 2003). The activity of sandflies

play an important role in determining the period of maximum risk of leishmaniasis transmission and hence when control programmes are necessary (El-badry et al., 2009). In addition to the control of the vector, there are other control methods for the disease, such as control of reservoirs, protection of people and health education (Al-Hammad, 2003).

Malaria control programmes, based on indoor residual insecticide spraying have had a side benefit for leishmaniasis incidence (Farrell, 2002). Saudi Arabia has adopted an appropriate programme for the control of both diseases; malaria and leishmaniasis (Ref. Section 3.9). Therefore the malaria programme has had an impact upon the incidence of leishmaniasis particularly in Asir and Jazan Regions (Dr A. Abdoon 2007 pers. Comm.). In Asir region, the MOH has a strategy for the control of disease vectors in each endemic area. This strategy depends upon several guidelines (Al-Hammad, 2003; MOH, year statistical book, 2006; Dr A. Abdoon 2007 pers. Comm.):

- a. Spraying of the breeding foci of vector to control the larva. This method is widespread all over the endemic areas of Saudi Arabia and starts at the onset of sandfly activity.
- b. Spraying the houses (deep cracks in walls, dark corners and humid areas) caves, rodent burrows, garbage areas and abattoirs with insecticides that additionally have a residual effect.
- c. Control of carrier animals and rodents.
- d. Mechanical control method, which can involve the use of various nets to protect against the vector (sandfly). Examples include bed nets, fine-mesh netting for windows and doors, and use of insecticide treated nets.
- e. Widespread distribution of nets impregnated with insecticides.
- f. Health education programmes in schools and gathering places to educate people how to protect their selves from this disease.
- g. Collaboration through cooperative agreements with local and central health departments, academic institutions and other governmental or private-sector organizations related to public health.

4.9 Chapter Rationale and Hypotheses

The influence of weather upon CL cases is important to understand for a number of reasons:

- Based on the findings of this study and an understanding of how CL is affected by weather it should be possible to design an early warning system to predict incidence. This would be useful for health service planning, for example identifying periods when enhanced insecticide spraying is required.
- 2. If weather alters under climate change then this data could help predict the impact of this change on CL in Asir Region.
- 3. It can help clarify the important modes of transmission in the region.
- 4. It improves our understanding disease transmission dynamics (Boussaa et al., 2005).

This study tests the hypotheses that there are relationships between weather (maximum, mean and minimum temperature, rainfall and relative humidity) and CL cases in Asir Region.

4.10 Data and Method

4.10.1 Sources of Data

Official letters were issued, instructing authorized personal to collect health and weather data. The numbers of CL cases in each month from the first of Jan 1996 to the end of Dec 2007 for the whole of Asir Region were obtained from the Vector Control Administration, Health Affairs Directorate in Asir. This department represents The Ministry of Health in this region. The data represents all reported cases of CL in Asir region and has been collected from every hospital and primary health care centres (public or private) and the Main Parasitology Laboratory in Abha (Annual reports of the Vector Control Administration, Health Affairs Directorate in Asir, 1996- 2007, VCA, MOH). A case of cutaneous leishmaniasis is a person showing clinical signs (skin or mucosal lesions) with parasitological confirmation of the diagnosis by positive smear or culture. For each case the sector where it was reported was known, as well as the month that the sample was reported and analysed. This however, presents a number of issues. Firstly, because we only have the reporting sector and not the address where the patient lived, it is possible that the person acquired CL outside that sector. This is likely to be the case with the Main Parasitology Laboratory in Abha which collects data from across the whole region.

Once the CL data was collected the aim was to obtain corresponding weather data for each district within Asir. However, this proved impossible due to the limited availability of data for many districts within Asir. Weather data (maximum, mean and minimum temperature, rainfall and humidity) were collected from Khamise city. This city is the second main city in the region, and covers the largest occupied sector in the middle of the region (MOH, 2006). These data were obtained from The Centre of Information and Documentation in the Meteorological and Environmental Protection section of the Ministry of Defence and Aviation in Saudi Arabia. Other weather data for the lowlands were obtained from Majaredah (maximum and minimum temperature for 132 out of 144 months) and Rejal-Alma (rainfall for 132 out of 144 months) from the Ministry of Water and Electricity. All locations of weather stations in Asir Region are illustrated in Figure 4.9.

4.10.2 CL Geography:

As indicated in Section 4.7 there are wide differences in topology and weather within Asir. Consequently there are likely to be differences in leishmaniasis incidence and seasonality between sectors. To investigate this further the overall incidence rates and histograms of CL were produced for each of the sectors within Asir. Data exists for nine sectors: Abha, Khamise, Sarat-Abedah, Muhail, Majaredah, Rejal-Alma, Dhahran-Aljanoub, Tanoma and Sabt-Alalaya. One sector was not included in the analysis (Ahad-Rifadah) as it only had a few cases during the period of this study (8 cases). In addition a few cases (11 cases) existed for unclassified sectors during the study period (Dr A. Abdoon 2007 pers. Comm.).

The overall leishmaniasis incidence rates for each sector and histograms of monthly incidence are presented in Figure 4.10. In most of the highlands areas such as Abha, Khamise, Sarat-Abidah, Tanoma and Sabt-Alalaya there are clear seasonalities with peak incidence between October and March. The average numbers of cases during the summer are lower. The rates (representing the number of positive cases in the year/100,000 population) of these sectors are 36.6, 8.8, 35.5, 67.7 and 27.0 respectively. In lowlands such as Majaredah and Rejal-Alma there is much less seasonality that in the highlands, with peak averages of cases occurring between

February and June, and the averages of cases in the rest of the year are little lower. The rates for these sectors are 1.31 and 4.37 respectively. The rest of the infected sectors in Asir are Muhail and Dhahran-Aljanoub which do not give us clear seasonality with rates of 1.05 and 1.02 respectively.

Differences in incidence between sectors may be due to differences in reporting. However, varying seasonalities are more likely to be due to transmission differences between sectors. The limited weather data available across Asir imply that it is not possible to examine the associations between CL cases and weather for all sectors in Asir individually.



For the analysis four models were constructed. The first was an analysis of CL in all Asir against Khamise weather data (Maximum, mean and minimum temperature, rainfall and relative humidity, Station 1 Figure 4.9). The following sectors were grouped together based upon sectors with similar seasonal patterns described in the previous paragraph and presented in Figure 4.10. The most appropriate weather data for each group was also selected:

- First Group (Highlands Group): CL cases in 7 sectors: Abha, Khamise, Sarat-Abedah, Tanoma, and Sabt-Alalaya against Khamise weather data (Maximum, mean and minimum temperature, rainfall and relative humidity, Station 1 Figure 4.9).
- Second Group (Lowlands Group): CL case in Majaredah and Rejal-Alma against Majaredah temperature (maximum and minimum, Station 2 Figure 4.9) and Rejal-Alma rainfall (Station 3 Figure 4.9). No relative humidity data was available for this area.
- Third Group (Different Lands Group): Muhail and Dhahran-Aljanoub. The data for this group has not been analysed for two important reasons: As shown in Figure 4.10 a low number of cases were reported during the study period. The second reason is that no representative weather data exists for this group of sectors.

4.10.3 CL Data Description

The monthly averages of CL cases in the whole of Asir Region during the study period are presented in Figure 4.11 and show a transmission of the parasite throughout the year. However, there is a clear seasonality with peak incidence between October and March, with the highest average number of cases in January (37.5) followed by February (30.5). The average number of cases during the summer is lower, with the lowest number occurring in June (16.8). The standard deviation bars on the graph indicate that there is a large variability from year to year in monthly cases of CL.



Figure 4.11: Monthly averages of CL cases in All Asir Region (1996 - 2007)

Most of the CL cases in Asir are in the first group (highlands) and so the monthly patterns of cases are similar between the two. Figure 4.12 shows the monthly averages of CL cases in the highlands group during study period. There is a clear seasonality with peak averages of cases between October and March, with the highest average cases in Jan (28.2) followed by November (27.8). The average numbers of cases during the summer are lower, with the lowest number of cases in June (11.08). The standard deviation bars on the graph indicate that there is a large variability from year to year in monthly cases of CL.



Figure 4.12: Monthly averages of CL cases in the first group (Highlands) of Asir Region (1996 - 2007)

Figure 4.13 shows monthly averages of CL cases in the lowlands group (Group 2) in Asir Region during study period. There is much less seasonality than in the highlands group with the peak cases occurring between February and June, with the highest average cases in March (5.5) followed by June (4.1). The averages of cases in the rest of the year are little lower, with the lowest averages of cases in September (2.1). The standard deviation bars on the graph indicates that there is a large variability from year to year in monthly cases of CL.



Figure 4.13: Monthly averages of CL cases in the second group (Lowlands) of Asir Region (1996 - 2007)

4.10.4 Temperature Data Description:

The temperature data for Khamise will be used in the analysis of cases across the whole of Asir Region as well as the first group (highlands). Descriptions of these data were presented in Section 3.11.3. Although this analysis is for a different year range (1996-2007 vs. 1995-2006) the seasonal trends in these data are similar to those presented previously.

The temperature in Majaredah will be used as representative of conditions in the lowlands group and these data were again presented previously for a slightly different range of years in Section 3.11.3.

4.10.5 Rainfall Data Description

Khamise rainfall will be assumed to be representative of all Asir and the highlands group. The monthly average rainfall in this city during the study period was presented in Section 3.11.4 for a slightly different yearly range.

Rejal-Alma is assumed to be representative of rainfall of the lowlands group, and data for this weather station was presented also in Section 3.11.4.

4.10.6 Relative Humidity Data Description

The relative humidity in Khamise will be assumed to be representative of all Asir and the highlands group. These data were presented previously in section 3.115.

4.10.7 Statistical Analysis Data Preparation

The analysis to examine the relationship between monthly cases of CL and the selected weather data proceeded in a number of stages;

A histogram of the number of CL cases each month in the whole of Asir shows a positively skewed distribution, with the majority of values between 7 and 35 cases. There are a few months with more than 40 cases and a few with less than 5 cases. This was confirmed through descriptive statistics. This may cause problems when performing regression analysis and so the natural logarithm of CL cases (adding 1 to zero values) was performed for the cases to produce Ln CL and this is illustrated in Figure 4.14. Descriptive statistics of this distribution are presented in Table 4.3 and demonstrated that taking the natural logarithm of CL produces a variable that is nearly normally distributed. For similar reasons the natural logarithm was taken of CL cases in highlands (Figures 4.15) (Table 4.4) and lowlands groups (Figures 4.16) (Table 4.5).

In terms of the independent variables, the rainfall data from Khamise also demonstrated a positively skewed distribution. Therefore, we took the natural logarithm of rainfall (adding 1 to zero values) to produce Ln rainfall. Figure 4.17 shows histograms of the rainfall after transformation, and Table 4.6 presents the descriptive statistics of this distribution.

The data were then adjusted for effects that could bias the results. Over time the number of CL cases in Asir has declined and this long term trend is shown in Figure 4.18. Since the 1990's cases have declined from the peak of cases in 1998. The long term of trend in CL in each of the two groups is shown in Figures 4.19 and 4.20. All these groups indicate a decrease in cases since the late 1990's.

In order to control for this effect, for the whole of Asir, the highlands and lowlands data, the natural logarithm of CL cases were put into a regression against time (each month numbered from 1 to 144). The unstandardised residual from this model represent the natural logarithm of CL cases with the long term trend in incidence removed.


Figure 4.14: Frequency of natural logarithm of CL (Ln. CL) in All Asir Region (1996 – 2007)

Table 4.3: Descriptive Statistics for Ln. CL Cases in Whole Air Region



Figure 4.15: Frequency of natural logarithm of CL (Ln. CL) in the highlands group (1996 – 2007)

Table 4.4: Descriptive Statistics for Ln. CL Cases in Highlands Group

	N	Minimum	Movimum	Maan	Std Doviation	Skev	vness
Ln. CL	N	wiiiiiiiiiiiiiiiiiiiiiiiiiiiiiiiiiiiiii	Maximum	Iviean	Stu. Deviation	Statistic	Std. Error
	144	0.69	4.25	2.81	0.686	- 0.380	.202



Figure 4.16: Frequency of natural logarithm of CL (Ln. CL) in the lowlands group (1996 – 2007)

Table 4.5: Descriptive statistics for Ln. CL cases in the lowlands group



Figure 4.17: Frequency of natural logarithm rainfall (Ln.R) in Khamise (1996 – 2007)

Ln.Rainfall.Khamise

Table 4.6: Descriptive Statistics of Ln Rainfall in Khamise

Ln	N	Minimum	Maximum	Mean	Std. Deviation	Ske	ewness
Rainfall	144	0	5.27	1 0 2 6	1 557	Statistic	Std. Error
		0	5.27	1.820	1.557	0.145	0.202



Figure 4.18: Monthly CL cases in all Asir Region (1996 – 2007)



Figure 4.19: Monthly CL cases in the highlands group in Asir Region (1996 – 2007)



Figure 4.20: Monthly CL cases in the lowlands group in Asir Region (1996 – 2007)

4.10.8 Statistical Analysis

The literature review has demonstrated that the relationship between CL cases and weather may vary according to the time of the year. Therefore, the analysis proceeded in stages

- Within each group (all Asir, highland, lowlands) months were grouped together. The models for all Asir and the highlands region consider three time periods; February to May, June to September and October to January (See Figure 4.11 for all Asir and Figure 4.12 for the highlands). The results for the lowlands region are presented for 2 time periods; March to November and December to February (See Figure 4.13). The grouping for all Asir, highlands and lowlands for each period was depended on the pattern of cases throughout the year.
- To produce an analysis that is more robust to the distribution of the data an ordinal logistic of Ln CL cases for every month against each of the weather variables was produced as well. This was achieved by dividing Ln CL into five categories with equal numbers of months.
- 3. In the regular regression analysis it was not possible to assume that the CL rate in one month is unrelated to the CL rate in the next month (i.e. the data is autocorrelated). Therefore all models were analyzed in three ways. The first did

not control for seasonality or autocorrelation. The second controlled for seasonality by incorporating dummy variables for each of the individual months. The final controlled for autocorrelation by adding the CL cases in the previous months into these models.

- 4. The same actions were repeated for the ordinal logistic analysis.
- 5. The regressions were performed against the current month's weather data as well as the previous month's weather data up 6 previous months. The literature review indicated that lagged relationships may exist up to this period. We then examined these relationships and combined the weather from the three months with the strongest associations with CL.
- 6. In most cases the strongest relationships were between DT Ln CL and temperature in 2-4 previous months and between DT Ln CL and rainfall in 1-3 previous months. Therefore, for consistency these lags were used in all models.

All of the data were analyzed using SPSS v.16 / v.18 for Windows and Microsoft Excel 2003 / 2007.

4.11 Results

The analysis is presented by first looking at relationships between CL and temperature (maximum, mean and minimum) followed by rainfall and relative humidity. This is performed for all of the Asir region, the highlands and the lowlands separately.

4.11.1 CL with Maximum Temperature

Linear regressions and ordinal logistic of DT Ln CL cases against maximum temperature are presented for all Asir during June to September in Table 4.7, the highlands June to September in Table 4.8 and the lowlands during March to November in Table 4.9. Models for the rest of year for this variable and all similar variables which have a few significant relationships are not presented in this chapter but appear in Appendix B1. In these tables 0P represents the models where the independent variables are not lagged, 1P represents the independent variables lagged by 1 month and so on.

		Li	near re	gressi	on			C	rdinal	logisti	ic	
Model	Ma temper on	ıx. rature ly	Ma tempe wi seasor	ax. rature th nality	Ma temper with correla seasor	ax. rature auto ation& nality	Ma temper on	ıx. rature ly	Ma temper wi seasor	ux. rature th nality	Ma temper with correla seasor	ature auto auto alition&
	Cof.	Р	Cof.	Р	P Cof. P		Cof.	Р	Cof.	Р	Cof.	Р
0 P	.030	.624	.130	.106	.130	.109	.270	.374	.940	.044	.890	.113
1 P	.070	.128	.070	.259	.080	.218	.360	.177	.430	.218	.640	.139
<mark>2 P</mark>	<mark>.065</mark>	<mark>.003</mark>	<mark>.161</mark>	<mark>.002</mark>	<mark>.164</mark>	<mark>.001</mark>	<mark>.207</mark>	<mark>.007</mark>	<mark>.730</mark>	<mark>.026</mark>	<mark>.909</mark>	<mark>.020</mark>
<mark>3 P</mark>	<mark>.041</mark>	<mark>.009</mark>	<mark>.119</mark>	<mark>.017</mark>	<mark>.146</mark>	<mark>.006</mark>	<mark>.090</mark>	<mark>.017</mark>	<mark>.060</mark>	<mark>.030</mark>	<mark>.360</mark>	<mark>.040</mark>
<mark>4 P</mark>	<mark>.050</mark>	<mark>.006</mark>	<mark>.130</mark>	<mark>.004</mark>	<mark>.130</mark>	<mark>.003</mark>	<mark>.080</mark>	<mark>.035</mark>	<mark>.040</mark>	<mark>.035</mark>	<mark>.350</mark>	<mark>.041</mark>
5 P	.040	.033	.070	.127	.070	.101	.050	.637	190	.455	060	.833
6 P	.080	.005	.090	.023	.100	.021	.080	.651	.010	.960	.630	.056
<mark>2-4 P</mark>	<mark>.055</mark>	<mark>.003</mark>	<mark>.221</mark>	<mark>.001</mark>	<mark>.247</mark>	<mark>.001</mark>	<mark>.323</mark>	<mark>.030</mark>	<mark>.432</mark>	<mark>.032</mark>	<mark>.851</mark>	<mark>.034</mark>

Table 4.7: DT Ln CL and maximum temperature in Asir during June to September

The results indicated that in Asir over June to September CL cases appear to be a positively significantly associated with average maximum temperature in the 2-4 previous months for linear regression and ordinal logistic regression.

		Li	near re	gressi	on			С	rdinal	logisti	c	
Model	Ma temper on	ix. rature ly	Ma temper wi seasor	ux. rature th nality	Ma temper with correla seasor	ax. rature auto ation& nality	Ma temper on	ıx. rature ly	Ma temper wi seasor	ax. rature th nality	Ma temper with correla seasor	ux. rature auto ution& nality
	Cof.	Р	Cof.	Cof. P .190 .077		Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	.020	.776	.190	.077	.200	.069	.210	.472	.940	.049	.670	.192
1 P	.080	.246	.030	.769	.040	.709	.410	.177	.330	.382	.260	.550
<mark>2 P</mark>	<mark>.082</mark>	<mark>.008</mark>	<mark>.170</mark>	<mark>.021</mark>	<mark>.169</mark>	<mark>.024</mark>	<mark>.290</mark>	<mark>.049</mark>	<mark>.783</mark>	<mark>.023</mark>	<mark>1.08</mark>	<mark>.007</mark>
<mark>3 P</mark>	<mark>.050</mark>	<mark>.012</mark>	<mark>.145</mark>	<mark>.040</mark>	<mark>.179</mark>	<mark>.018</mark>	<mark>.111</mark>	<mark>.025</mark>	<mark>.176</mark>	<mark>.036</mark>	<mark>.220</mark>	<mark>.044</mark>
<mark>4 P</mark>	<mark>.062</mark>	<mark>.007</mark>	<mark>.160</mark>	<mark>.009</mark>	<mark>.165</mark>	<mark>.008</mark>	<mark>.103</mark>	<mark>.038</mark>	<mark>.315</mark>	<mark>.031</mark>	<mark>.342</mark>	<mark>.041</mark>
5 P	.070	.010	.120	.046	.140	.029	.140	.251	.070	.773	.230	.392
6 P	.090	.028	.100	.084	.100	.079	.001	.985	010	.693	.040	.894
<mark>2-4 P</mark>	<mark>.073</mark>	<mark>.005</mark>	<mark>.262</mark>	<mark>.002</mark>	.100 .073 .02 .278 .001		<mark>.164</mark>	<mark>.047</mark>	<mark>.512</mark>	<mark>.033</mark>	<mark>.718</mark>	<mark>.036</mark>

 Table 4.8: DT Ln CL and maximum temperature in the highlands during June to

 September

P: previous month. Cof.: Coefficients. P: **P** value (= or<).

The results indicated that in highlands over June to September CL cases appear to be a positively significantly associated with average maximum temperature in the 2-4 previous months for linear regression and ordinal logistic regression.

		Li	near re	gressi	on			C	Ordinal	logisti	ic	
Model	Ma temper on	ıx. rature ly	Ma temper wi seasor	ux. rature th nality	Ma temper with correla season	ix. rature auto ation& nality	Ma temper on	ıx. rature ly	Ma temper wi seasor	ux. rature th nality	Ma temper with correla seasor	ax. rature auto ation& nality
	Cof.	Р	Cof.	Р	Cof.	of. P Co		Р	Cof.	Р	Cof.	Р
0 P	010	.763	010	.853	020	.720	040	.590	010	.956	.030	.912
1 P	031	.065	.027	.540	.020	.642	110	.082	.069	.683	.081	.644
2 P	040	.006	.030	.557	.020	.714	120	.015	050	.749	040	.825
3 P	040	.001	015	.742	024	.591	120	.010	138	.426	123	.491
4 P	030	.005	.010	.805	.020	.709	100	.032	260	.200	150	.473
5 P	020	.069	.001	.955	.001	.933	060	.238	160	.339	080	.701
6 P	.001	.824	.030	.955 .001 .464 .030		.429	.010	.822	.001	.981	.090	.621

 Table 4.9: DT Ln CL and Maximum temperature in the lowlands during March to

 November

Temp. available for 132 out of 144 months for the whole of year model

In the lowlands sector CL cases over March to November are not consistently associated with maximum temperature for any of the previous months.

4.11.2 CL with Mean Temperature

Linear regressions and ordinal logistic of DT Ln CL cases against mean temperature are presented for all Asir during June to September in Table 4.10 and the highlands in Table 4.11. Mean temperature data was not available for the lowlands.

		Li	near re	gressi	on			С	rdinal	logisti	c	
Model	Me temper on	an rature ly	Me temper wi seasor	an rature th nality	Me tempe with correla seasor	ean rature auto ation& nality	Me temper on	an rature ly	Me temper wi seasor	an rature th nality	Me temper with correla seasor	an rature auto ition& nality
	Cof.	Р	Cof.	Р	Cof. P		Cof.	Р	Cof.	Р	Cof.	Р
0 P	.010	.883	.190	.073	.190	.082	620	.158	660	.262	1.10	.165
1 P	.070	.222	.070	.190 .073 .070 .358		.358	130	.652	500	.246	.220	.702
2 P	.070	.005	.200	.003	.200	.004	.120	.384	.170	.043	1.10	.037
3 P	.040	.018	.120	.070	.150	.036	.070	.510	.250	.496	.180	.670
<mark>4 P</mark>	<mark>.050</mark>	<mark>.012</mark>	<mark>.160</mark>	<mark>.009</mark>	<mark>1.70</mark>	<mark>.007</mark>	.070	.494	.200	.573	.340	.409
5 P	.050	.043	.080	.277	.080	.251	.030	.783	800	.054	570	.212
6 P	.080	.017	.110	.117	.120	.108	.001	.998	750	.088	.360	.489

Table 4.10: DT Ln CL and mean temperature in Asir during June to September

The results indicated that in Asir over June to September CL cases appear to be a positively significantly associated with mean temperature 2- 4 months previously but only in the linear regression. This effect mostly disappears in the ordinal logistic regression.

		Li	near re	gressi	on			С	Ordinal	logisti	ic	
Model	Me temper on	Mean temperature only		an rature th nality	Me tempe with correla seasor	ean rature auto ation& nality	Me tempe on	an rature ly	Me temper wi seasor	an rature th nality	Me temper with correla seasor	an rature auto ation& nality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	of. P Cof. P		Р	Cof.	Р
0 P	.010	.917	.280	.059	.290	.057	740	.109	800	.215	030	.966
1 P	.090	.245	.050	.660	.060	.622	050	.877	560	.225	120	.826
<mark>2 P</mark>	<mark>.090</mark>	<mark>.011</mark>	<mark>.230</mark>	<mark>.024</mark>	<mark>.230</mark>	<mark>.027</mark>	.200	.090	.300	.056	<mark>1.30</mark>	<mark>.021</mark>
3 P	.060	.025	.180	.206	.150	.139	.080	.428	290	.422	450	.267
4 P	.060	.022	.140	.111	.150	.103	.100	.371	120	.742	.260	.501
5 P	.070	.023	.130	.169	.150	.143	.090	.478	440	.272	310	.445
6 P	.090	.054	.090	.374	.090	.358	020	.905	800	.055	400	.378

Table 4.11: DT Ln CL and mean temperature in the highlands during June toSeptember

The results indicated that in the highlands between June and September CL cases appear to be positively and mostly significantly associated with mean temperature only in the 2 previous months for the linear regression and ordinal logistic regression.

4.11.3 CL with Minimum Temperature

Linear regressions and ordinal logistic of DT Ln CL cases against minimum temperature are presented for all Asir during June to September in Table 4.12, the highlands in Table 4.13 and the lowlands during March to November in Table 4.14.

		Li	near re	oressi	on			C	rdinal	logisti	C	
Model	Mi temper on	n. rature ly	Mi temper wi seasor	n. rature th nality	Mi temper with correla seasor	n. rature auto ation& nality	Mi temper on	n. rature ly	Mi temper wi seasor	n. rature th nality	Mi temper with correla seasor	n. rature auto ution& nality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	030	.647	.100	.372	.100	.384	400	.216	760	.212	.440	.542
1 P	.060	.259	020	.845	020	.856	200	.499	-1.20	.029	480	.445
<mark>2 P</mark>	<mark>.070</mark>	<mark>.014</mark>	<mark>.200</mark>	<mark>.032</mark>	<mark>.190</mark>	<mark>.037</mark>	<mark>.120</mark>	<mark>.428</mark>	<mark>.189</mark>	<mark>.039</mark>	<mark>.910</mark>	<mark>.035</mark>
3 P	.040	.041	.060	.483	.080	.395	.080	.493	330	.486	130	.810
4 P	.040	.049	.070	.358	.070	.369	.090	.434	130	.768	.070	.882
5 P	.040	.092	010	.869	010	.883	.080	.497	170	.640	350	.381
6 P	.040	.195	030	.574	030	.596	.030	.835	450	.170	410	.253

Table 4.12: DT Ln CL and minimum temperature in All Asir during June toSeptember

The results indicated that in all Asir over June to September CL cases appear to be a positively significantly associated with minimum temperature only in the 2 previous months for both linear and ordinal logistic regression.

		Li	near re	gressi	on			С	rdinal	logisti	ic	
Model	Mi temper on	n. rature ly	Mi temper wi seasor	n. rature th nality	Mi temper with correla seasor	n. rature auto ation& nality	Mi temper on	n. rature ly	Mi temper wi seasor	n. rature th nality	Mi temper with correla seasor	n. rature auto ation& nality
	Cof.	Р	Cof.	Р	Cof. P		Cof.	Р	Cof.	Р	Cof.	Р
0 P	090	.001	150	.120	150	.120	280	.017	700	.099	420	.349
1 P	100	.001	130	.120	100	.222	270	.260	300	.392	170	.661
<mark>2 P</mark>	<mark>.040</mark>	<mark>.182</mark>	<mark>.120</mark>	<mark>.018</mark>	<mark>.150</mark>	<mark>.034</mark>	.210	.194	.100	.769	.660	.058
3 P	.080	.152	.100	.176	.080	.254	.001	.996	120	.686	030	.924
4 P	.090	.004	.060	.412	.020	.789	.190	.137	180	.574	290	.399
5 P	.060	.008	.070	.428	.060	.493	.130	.155	480	.221	030	.941
6 P	.060	.014	.800	.491	.090	.414	.140	.150	500	.329	.090	.394

Table 4.13: DT Ln CL and minimum temperature in the highlands during June toSeptember

The results indicated that in the highlands over June to September CL cases appear to be a positively significantly associated with mean temperature only in the 2 previous months. The effect is only seen in the linear regression.

 Table 4.14: DT Ln CL and minimum temperature in the lowlands during March to

 November

		Li	near re	gressi	on			С	rdinal	logisti	ic	
Model	Mi temper on	Min. temperature only		n. rature th nality	Mi temper with correla seasor	n. rature auto ation& nality	Mi temper on	n. rature ly	Mi temper wi seasor	n. rature th nality	Mi temper with correla seasor	n. rature auto ation& nality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	.020	.396	.040	.362	040	.412	.030	.628	.230	.257	.220	.244
1 P	040	.030	040	.315	060	.187	120	.079	.010	.946	160	.372
2 P	050	.002	050	.270	040	.341	120	.030	.070	.706	.040	.797
3 P	050	.001	.001	.980	.001	.966	110	.045	.030	.164	.230	.264
4 P	.040	.015	.080	.165	.090	.152	070	.200	470	.049	.420	.096
5 P	040	.029	.010	.842	.001	.968	.060	.311	.430	.151	.190	.543
6 P	010	.563	030	.614	.001	.936	.010	.854	.190	.402	.120	.568

Minimum temperature data: available for 132 out of 144 months for the whole of year model

In the lowlands sector CL cases over March to November are not consistently associated with minimum temperature for any of the previous months.

4.11.4 CL with Rainfall

Linear regressions and ordinal logistic of DT Ln CL cases against rainfall are presented for all Asir during June to September in Table 4.15, the highlands June to September in Table 4.16 and the lowlands during March to November in Table 4.17.

		Li	near re	gressi	on			С	rdinal	logisti	ic	
Model	Rain on	fall ly	Rain wi seasor	fall th nality	Rain with correla season	ifall auto ation& nality	Rain	ıfall ly	Rain wi seasor	ıfall th nality	Rain with correla season	fall auto tion& nality
	Cof.	Р	Cof. P		Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	030	.331	Cof. P 050 .189		050	.214	090	.587	010	.642	420	.120
1 P	.040	.342	020 .686		.140	.754	.140	.507	.130	.578	010	.973
2 P	084	.014	075	.059	074	.063	170	.393	197	.365	402	.122
3 P	053	.099	059	.093	061	.089	210	.243	220	.287	430	.067
4 P	.020	.417	030	.472	040	.423	020	.891	230	.336	.120	.670
5 P	.020	.536	030	.448	030	.458	010	.941	180	.430	100	.693
6 P	030	.498	030	.426	040	.397	.020	.909	060	.799	160	.564

Table 4.15: DT Ln CL and rainfall in All Asir during June to September

In the whole of Asir CL cases over June to September are not consistently associated with rainfall for any of the previous months.

		Li	near re	gressi	on			С	rdinal	logisti	ic	
Model	Rain	ıfall ly	Rain wi seasor	ıfall th nality	Rain with correla season	nfall auto ation& nality	Rain	ıfall ly	Rain wi seasor	fall th nality	Rain with correla season	fall auto tion& nality
	Cof.	Р	Cof. P		Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	080	.056	140	.015	140	.018	110	.538	190	.428	490	.083
1 P	.050	.297	140 .015 .050 .391		.050	.446	.120	.552	.170	.467	.060	.818
2 P	056	.252	041	.469	040	.498	092	.649	099	.659	347	.173
3 P	074	.094	081	.103	082	.103	140	.439	070	.408	180	.409
4 P	.043	.295	023	.702	027	.656	.110	.515	040	.868	.230	.351
5 P	.010	.901	070	.231	070	.232	110	.506	320	.159	420	.077
6 P	010	.873	040	.538	040	.493	.020	.915	120	.607	.130	.637

 Table 4.16: DT Ln CL and rainfall in the highlands during June to September

In the highlands sector CL cases over June to September are not consistently associated with rainfall for any of the previous months.

		Li	near re	gressi	on			С	Ordinal	logisti	ic	
Model	Rain on	ıfall ly	Rain w seaso	nfall ith nality	Rain with correla seasor	fall auto tion& nality	Rain on	ıfall ly	Rain wi seasoi	ıfall th nality	Rain with correla seasor	fall auto tion& nality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	.070	.020	.050 .082		.070	.110	.420	.001	.420	.001	.270	.036
<mark>1 P</mark>	<mark>.075</mark>	<mark>.008</mark>	.076 .005 .		<mark>.070</mark>	<mark>.009</mark>	<mark>.460</mark>	<mark>.001</mark>	<mark>.531</mark>	<mark>.001</mark>	<mark>.355</mark>	<mark>.010</mark>
2 P	.009	.001	.020 .548		.030	.314	.300	.005	.300	.007	.040	.744
<mark>3 P</mark>	<mark>.079</mark>	<mark>.003</mark>	<mark>.076</mark>	<mark>.004</mark>	<mark>.076</mark>	<mark>.003</mark>	<mark>.535</mark>	<mark>.001</mark>	<mark>.559</mark>	<mark>.001</mark>	<mark>.409</mark>	<mark>.002</mark>
4 P	.030	.219	.040	.168	.020	.395	.400	.001	.490	.001	.050	.726
5 P	.020	.539	.020	.384	.020	.568	.410	.001	.490	.001	.210	.115
6 P	.010	.742	.030	.244	.030	.259	.460	.001	.620	.001	.450	.001
<mark>1-3 P</mark>	.099 .004 .085 .012		<mark>.078</mark>	<mark>.022</mark>	<mark>.800</mark>	<mark>.001</mark>	<mark>.829</mark>	<mark>.001</mark>	<mark>.492</mark>	<mark>.008</mark>		
Rainfall d	ata: avai	ilable fo	r 132 o	t of 14	4 month	s for th	e whole	of vear	model			

Table 4.17: DT Ln CL and rainfall in the lowlands during March to November

le for 132 out of 144 months for the whole of year model

The results indicated that in the lowlands over March to November CL cases appear to be positively and significantly associated with average rainfall of 1-3 months previously for both the linear and ordinal logistic regression.

4.11.5 CL with Relative Humidity

Linear regressions and ordinal logistic of detrended logarithm CL cases against humidity are presented for all Asir June to September in Table 4.18 and the highlands June to September Table 4.19. No humidity data was available for the lowlands.

		Li	near re	gressi	on			С	rdinal	logisti	ic	
Model	Rela humi on	tive dity ly	Rela humi wi seasoi	tive dity th nality	Rela humi with correla seasor	tive dity auto ation& nality	Rela humi on	tive dity ly	Rela humi wi seasor	tive dity th nality	Rela humi with correla seasor	tive dity auto ttion& nality
	Cof.	Р	Cof. P		Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	.001	.987	.001 .803		.001	.832	020	.646	020	.690	040	.573
1 P	.010	.211	.010 .483		.010	.479	.030	.530	.020	.649	.001	.950
2 P	015	.021	013	.132	013	.151	027	.464	026	.587	093	.094
3 P	.013	.015	.013	.120	.015	.091	020	.422	.010	.867	.060	.248
4 P	.020	.001	.030	.073	.030	.082	.070	.058	.090	.066	.140	.021
5 P	010	.038	010	.198	010	.152	060	.125	070	.191	010	.885
6 P	.015	.021	.020	010 .198 .020 .132		.110	.090	.032	.100	.037	.140	.016

Table 4.18: DT Ln CL and relative humidity in All Asir during June to September

		Li	near re	gressi	on			С	rdinal	logisti	c	
Model	Rela humi on	tive dity ly	Rela humi wi seasor	tive dity th nality	Rela humi with correla seasor	tive dity auto ation& nality	Rela humi on	tive idity ly	Rela humi wi seasor	tive dity th nality	Rela humi with correla seasor	tive dity auto tion& nality
	Cof.	Р	Cof.	Cof. P		Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	.001	.080	010 .756		010	.760	.001	.958	.010	.936	.001	.961
1 P	.010	.262	.010 .429		.010	.431	.010	.790	.010	.784	030	.587
2 P	013	.145	008	.529	007	.588	036	.344	028	.548	077	.144
3 P	011	.159	.001	.907	.001	.941	010	.784	.040	.373	.040	.418
4 P	024	.007	024	024 .057		.063	.110	.060	.060	.071	.100	.054
5 P	020	.031	020 .237		020	.202	090	.049	090	.107	090	.146
6 P	020	.033	020 .237 020 .129		020	.131	060	.188	060	.229	070	.221

 Table 4.19: DT Ln CL and relative humidity in the highlands during June to

 September

In the all Asir and highlands sector CL cases over June to September are not consistently associated with relative humidity for any of the previous months.

4.11.6 Multiple Regressions

In this section we will present the results from the normal and multiple regressions for the coefficient of DT Ln CL cases against averages of three main variables of weather data (Maximum Temperature, Rainfall and Relative Humidity). For temperature, we will present maximum temperate only as it has a similar seasonality to minimum and mean temperature and appeared to have the strongest associations with CL. In the multiple regressions we have presented only models for times of the year when one of the independent variables was significant.

A- Multiple Regression of CL in All Asir

The strongest results from the univariable linear and ordinal logistic regression for CL in all Asir against maximum temperature, rainfall and relative humidity, for average of 2, 3 and 4 previous months during June to September period are illustrated in Table 4.20. Table 4.21 presents the multiple regressions for all combination of these variables.

Table 4.20: Regressions of DT Ln CL cases of Asir for maximum temperature, rainfall and relative humidity during June to September for
average of 2 to 4 previous months

A				Linea	ar regre	ssion							Ord	inal log	istic			
of 2-4 previous months –	١	/ariable Only	S	se	With asonali	ty	V coi Se	Vith aut relatior easonali	.o 1 & .ty	V	/ariable Only	S	se	With easonali	ty	V cor Se	Vith aut relation easonali	o & ty
monuis	Cof.	Р		Cof.	Р		Cof.	Cof. P Adj. R ²		Cof.	Р		Cof.	Р		Cof.	Р	
Max.Tem. Only	.055	.003	.155	.221	.001	.291	.247	.001	.319	.323	.030	-58.1	.432	.032	-56.4	.851	.034	-51.1
Rainfall Only	130	.043	.066	158	.056	.175	162	.055	.171	512	.166	-68.5	777	.068	-67.4	845	.061	-51.3
Humidity Only	024	.031	.201	030	.080	.121	030	.089	.123	056	.169	-68.6	069	.271	-68.5	168	168	-52.5

Table 4.21: Regressions of DT Ln CL cases of Asir with multiple weather (maximum temperature with relative humidity and for all of them together) during June to September for average of 2 to 4 previous months

Average				Line	ear regr	ession							Ord	inal log	gistic			
of 2-4 previous months		Variable: Only	S	S	with easonalit	у	Co	With auto orrelation Seasonalit	D L & Ly		Variable Only	S	S	With easonali	ty	co S	With auto rrelation easonalit) & у
Variables In Multiple Method	Cof.	Р	Adj. R ²	Cof.	Р	$\begin{array}{c} \text{Adj.} \\ \text{R}^2 \end{array}$	Cof.	Р	Adj. R ²	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.
Max.Tem	.059	.001	257	.183	.005	300	.207	.002	332	.153	.049	-	.169	.044	-	.600	.024	-
Rainfall	151	.010	.237	088	.214 .300		091	.192	.552	605	.107	61.48	638	.248	58.41	554	.283	50.53
_																		
Max.Tem	.008	.016	104	.183	.009	200	.217	.003	210	.019	.050	-	.303	.048	-	.411	.041	-
Humidity	021	.113	.184	012	.344	.289	009	.477	.312	053	.478	68.29	040	.602	68.29	127	.167	50.16
Rainfall	022	.002	220	114	.173	100	124	.144	102	457	.217	-	792	.138	-	357	.532	-
Humidity	103	.078	238	020	.138	.199	020	.156	.192	049	.224	67.80	.004	.963	67.39	133	.158	50.25
Max.Tem	.083	.045		.170	.018		.204	.006		.178	.027		.201	.039		.361	.031	
Rainfall	136	.038	.244	072	.365	.287	.086	.277	.315	633	.144	67.48	752	.640	67.29	291	.618	50.03
Humidity	007	.607		006	.667		002	.901		.011	.898	07.10	.020	.819	07.27	103	.323	50.05

The results of multiple regressions for CL in Asir during June to September demonstrated that in no models were more than one weather variable significant. We conclude therefore that the best model for period is that of DT Ln CL against maximum temperature in the previous 2-4 months. This temperature variable was positively significant in all variants of the normal and ordinal logistic regression. In all cases models including this variable had a better fit (adjusted R^2 and log likelihood ratio) and a stronger significance than any of the other weather variables. Higher maximum temperature was associated with elevated CL cases.

B- Multiple Regression of CL in Highlands Group

The strongest results from the univariable linear and ordinal logistic regression for CL in the highlands against maximum temperature, rainfall and relative humidity, for the average of the previous 2-4 months during June to September are illustrated in Table 4.22. Table 4.23 presents the multiple regressions for all combinations of these variables.

Table 4.22: Regressions of DT Ln CL cases of the highlands for maximum temperature, rainfall and relative humidity during June to September,
for average of 2 to 4 previous months

Average				Line	ar regre	ssion							Ord	inal log	istic			
of 2-4 previous months	١	/ariable Only	S	se	with easonali	ty	V COI Se	Vith aut rrelatior easonali	o 1 & ty	٧	/ariable Only	÷S	Se	With easonali	ty	V cor Se	Vith aut relation asonali	o 1 & ty
montins	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.
Max.Tem. Only	.073	.005	.140	.262	.002	.194	.278	.001	.198	.164	.047	-64.8	.512	.033	-64.1	.718	.036	-54.3
Rainfall Only	096	.289	.030	175	.080	.066	178	.079	.050	110	.765	-65.8	411	.339	-64.6	345	.427	-55.5
Humidity Only	022	.026	.083	018	.288	.022	017	.309	.002	048	.266	-65.2	029	.663	-64.9	084	.249	-55.1

Table 4.23: Regressions of DT Ln CL cases of the highlands with multiple weather (maximum temperature with rainfall and maximum temperature with relative humidity and for all of them together) during June to September for average of 2 to 4 previous months

Average				Line	ar regre	ssion							Orc	linal log	gistic			
of 2-4 previous months Variables	۲	/ariable only	S	se	with easonali	ty	co S	With au orrelatic easona	ito on & lity		Varia On	bles ly	s	With easonal	ity	co: Se	With au rrelation easonal	to n & ity
In Multiple Method	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Adj R ²	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.
Max.Tem	.077	.003	162	.245	.012	178	.263	.008	180	.177	.02	4 _	.440	.039	-	.740	.042	-
Rainfall	123	.143	.102	040	.706	.170	034	.747	.100	222	.55	4 64.57	.176	.714	63.09	.046	.927	53.16
							_		_				_		_	-		_
Max.Tem	.085	.036	102	.314	.003	100	.355	.001	207	.206	.03	- C	.632	.035	-	.676	.040	-
Humidity	.006	.764	.125	.016	.381	.190	.023	.235	.207	.019	.814	4 63.20	.037	.652	64.03	012	.891	54.24
							<u>.</u>					·						
Rainfall	069	.430	076	175	.168	044	183	.156	027	083	.82	3 -	461	.385	-	080	.881	-
Humidity	021	.037	.076	.001	.994	.044	.001	.944	.027	047	.174	4 65.17	.014	.871	64.61	076	.393	55.10
Max.Tem	.129	.019		.295	.006		.336	.003		.297	.020		.579	.019		.697	.040	
Rainfall	170	.073	.167	104	.384	.186	116	.237	.206	330	.433	-64.41	344	.524	-63.83	.091	.869	-54.23
Humidity	.023	.271		.025	.242		.032	.135		.052	.572		.064	.409		.019	.847	

The results of the multiple regressions for CL in the highlands during June to September demonstrated that in no models with more than one weather variable was positively significant. We conclude therefore that the best model for this period is that of DT Ln CL against maximum temperature in the previous 2-4 months. This weather variable was positively significant in all variants of the normal linear and ordinal logistic regressions and in all cases had a better fit (adjusted R^2 and log likelihood ratio) and a stronger significance than any of the other weather variables. Higher maximum temperature was associated with elevated CL cases.

C- Multiple Regression of CL in Lowlands Group

The strongest results from the univariable linear and ordinal logistic regression for CL in the lowlands against maximum temperature, and rainfall in the previous 1-3 months are illustrated in Table 4.24. Table 4.25 presents the multiple regressions for all combinations of these variables.

Table 4.24: Regressions of DT Ln CL cases of the lowlands for maximum temperature and rainfall during March to November for average of 1to 3 previous months

A				Linea	ar regre	ssion							Ord	inal log	istic			
of 1-3 previous months	V	/ariable Only	S	se	with asonali	ty	V coi Se	Vith aut relatior easonali	.0 1 & .ty	V	/ariable Only	es.	Se	With easonali	ty	V coi Se	Vith aut relation easonali	o & ty
	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.
Max. Only	038	.010	.058	.048	.426	.091	.032	.587	.122	137	.019	- 127.1	045	.843	- 124.2	018	.939	- 103.1
Rainfall. Only	.099	.004	.080	.085	.012	.142	.078	.022	.152	.800	.001	- 105.2	.829	.001	- 101.6	.492	.008	85.01

 Table 4.25: Regressions of DT Ln CL cases of the lowlands with multiple weather (maximum temperature and rainfall) during March to November for average of 1 to 3 previous months

Average				Line	ar regre	ssion							Ord	inal logi	istic			
of 1-3 previous months	Variables only			se	with easonali	ty	V COI Se	Vith aut relatior easonali	to n & ty	V	/ariable Only	es	S€	With easonali	ty	V cor Se	Vith aut relation easonali	o 1 & ty
 Multiple Method	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.
Max.Tem	028	.070	130	.035	.567	189	.023	.710	195	067	.322	-	.213	.443	-	.120	.665	-
Rainfall	.097	.006	.150	.086	.012	.107	.080	.022	.175	.853	.001	90.09	.893	.001	86.79	.565	.005	75.58

The results of multiple regressions for CL in the lowlands group during March to November demonstrated that in no models was more than one weather variable significant. We conclude therefore that the best model for this period is that of DT Ln CL against rainfall. Average rainfall in the previous 1-3 months between March and November was positively significant in all variants of the normal linear and ordinal logistic regression and in all cases had a better fit (adjusted R^2 and log likelihood ratio) and a stronger significance than maximum temperature. Higher rainfall was associated with elevated CL.

4.12 Discussion and Conclusion

There are two main species of leishmaniasis cases in Saudi Arabia; VL and CL (Al-Amru, 2002). This study focuses on CL in Asir Region as this form of illness is endemic and spread over the foothills and highlands of this region (Ibrahim et al., 2005). Al-Zahrani et al., (1989) state that the distribution of this disease in Asir region is influenced by some little known climate factors and geographical situation.

There were some difficulties during this study, as we only had limited data for weather in the selected areas in Asir. For example, the weather in Khamise City is not representative of the whole of Asir as the elevation in this region varies from 0 to 3000m. Weather and hence sandfly distributions and leishmaniasis cases are almost certain to vary across Asir. Also it was not possible to analyse the data for some sectors due to lack of weather data and the low number of cases. For example no weather data was available for the Different Lands Group and the numbers of cases reported for this sector was low. Finally, some of the Leishmaniasis cases were unclassified and so could not be assigned to a sector.

Overall the CL cases show that the total number of yearly cases is declining from a peak (outbreak) in 1998. This outbreak may have been due to the heavy rainfall throughout this region during some months of year 1997 and 1998 (Dr A. Abdoon 2007 pers. Comm.). However, our results indicated that rainfall is not associated with CL cases in the highlands area where most cases occur. Vector control is a likely explanation for this reduction in cases. The statistical analysis presented in this chapter was based on the hypothesis that there is a relationship between weather and CL. This was assumed because the distribution of leishmaniasis is tightly linked to geography and may be sensitive to weather and climate influences (Reithinger et al., 2007). There is also evidence that climate affects sandfly numbers which has an impact upon leishmaniasis cases. Cross et al., (1996) noted that the vector is affected by environmental conditions especially temperature and rainfall. In Argentina Salomon et al., (2008) showed that the activity season of one sandfly species sharply increased during the rainy season and decreased during the months of lower rainfall or higher temperatures. Hunter, (2003) mentioned that the increased rainfall can lead to an increase in vegetation, allowing expansion in the population of vertebrate host. An increase in sandfly populations can lead to an increase in the occurrence of leishmaniasis (Gage et al., 2008). There is therefore, much evidence that weather is important for leishmaniasis.

Monthly averages of CL cases in all Asir Region during the study period show transmission throughout the year. However, there is a clear seasonality with peak incidence between October and March, with the highest average number of cases in January. The average number of cases during the summer is lower, with the lowest number of cases in June. There was a high standard deviation in most months which indicates that there is a large variability from year to year in monthly CL cases. Most of the CL cases in Asir are in the first group (highlands) and so the monthly patterns of cases are similar between the two. There is much less seasonality in the lowlands than in the highlands group with the peak cases occurring between February and June, with the highest cases in March. Cases numbers in the rest of the year are little lower, with the lowest cases in September.

In the results, CL incidence was compared to temperature across Asir. The results presented evidence that temperature is an important factor affecting variation in CL cases but only during the summer (June to September). Elevated maximum temperature in the 2-4 previous months was highly significantly associated with leishmaniasis cases (P<0.001). This result was nearly identical in the highlands group, which is unsurprising as most cases in Asir occur in the highlands. This result is plausible as the lags indicate that temperature during the end of winter and early spring affects leishmaniasis cases in the summer. During the end of winter and early

spring temperatures are generally low (around 10 to 20° C) and so elevated temperature may have positive effects upon the vector. There was no effect in the lowlands indicating that temperatures are never a limiting factor for Leishmaniasis. In the lowlands winter temperatures are usually higher (20 to 25° C). The influence of temperature is supported in the literature and Cross et al, (1996) state that the vector is affected by environmental conditions, especially temperature. Hunter, (2003) and Gubler et al., (2001) state that there are some possible mechanism changes in the influence of temperature on the transmission of vector borne disease. Most of the CL cases in this study are from highland areas, and the highest average maximum temperature in the representative area occurs in June (32.26). In some parts of India one of the studies revealed that the sandfly were found only at temperatures between 17 and 36°C (Singh, 1999). Therefore the highlands might provide the optimum temperature for the vector during summer and then the activity of sandfly is likely to increase with this temperature.

The study shows that the increase of temperature during summer can lead to an increase of CL cases in the whole of the Asir Region and also in highlands group. However, there was no evidence for a relationship between maximum temperature and CL cases in the lowlands group. This is slightly unexpected as the highest average maximum temperature occurs in June which reaches over 40^oC. Adults and larvae are sensitive to high temperature and in laboratory experiments, all adult sandflies died within two hours at temperatures above 40°C (Cross and Hyams, 1996). A negative association with maximum temperature in the summer might therefore be expected. The lack of a result suggests that sandflies find mechanisms to avoid these high temperatures

Maximum, mean and minimum temperatures in Asir are highly correlated. However, maximum temperature in Asir was most strongly associated with leishmaniasis cases. This suggests that maximum temperature is most related to the temperatures experienced by the sandflies but the reasons for this are unclear.

Within the whole of Asir and in the highlands there was no evidence that rainfall impacted on CL cases. This may indicate that rainfall has no impact or influence on the species of sandflies in the highlands. In the lowlands group, there was evidence that increased rainfall can lead to increased cases of Leishmaniasis. There is a highly positive significant relationship between the average rainfall in the 1-3 previous months and CL cases between (March and November). This indicates that rainfall provides suitable conditions for the vector but in the lowlands only. Hunter, (2003) noted that increased rainfall can increase vegetation growth and allow expansion in the population of vertebrate hosts, therefore this can increase the infected sandfly numbers and thus the possibility of CL cases. MOH in Saudi Arabia, 2000 has mentioned that there is a highly significant correlation between the density of sandfly and positive cases. This is because after rainfall it takes time for increases in sandflies to occur.

In Pondicherry, India, summer rainfall is followed by an increase in the population of sandflies (Sirnivasan et al., 1993). Also some studies in Colombia stated that drought followed by rainfall increases the occurrence of CL (Cardenas et al., 2006; Gage et al, 2008). Cross et al., (1996) noted that the vector is affected by rainfall. Therefore, these conditions can lead to increase the vector, thus the occurrence of CL. Our results suggest that in the lowlands area of Asir there is no enough rainfall for vegetation. This means that when there is rainfall there is more sandflies and hence leishmaniasis. The weather in these areas is drier. It is therefore not surprising that more rainfall may lead to more vegetation and hence more sandflies and leishmaniasis.

Some studies note that humidity is one of the most important weather factors for sandfly survival, development and activity (Cross and Hyams, 1996; Sirnivasan et al., 1993; Kasap et al., 2009), but the results of this study do not show evidence that there was a significant relationship between relative humidity and CL cases for Asir or any of the endemic areas. This interpretation is slightly limited by the absence of humidity data in the lowlands. This contradiction could be due to the different location of this study. It also could be due to the increase in humidity which is associated with other weather factors, such as high temperature, which have shown themselves to be significant in the model.

In summary, this study showed that weather factors play an important role in affecting CL cases. Temperature was the most important factor associated with CL in all Asir and in the highlands area during the summer and beginning of autumn. If the maximum temperature in the previous 2 to 4 months was elevated there were more

cases. Rainfall plays an important role in CL transmission in the lowlands with greater cases if rainfall is elevated in the previous 1 to 3 months. This occurs in all months except during the winter season. Although this study shows that weather has an influence upon CL in Asir, it is important to recognize that leishmaniasis cases are becoming less of a problem in this Region. Overall, the study shows that the total number of yearly CL cases has declined to one third of cases in the nineties by the end of the study period. The reasons for this were provided in the literature review.

Chapter 5

<u>Influence of Climate Variables upon</u> <u>Schistosomiasis Cases in Asir Region, Saudi</u> <u>Arabia</u>

5.1 Introduction

Schistosomiasis, also known as Bilharzia, is a parasitic disease caused by parasitic worms called schistosomes that leads to chronic ill health in humans and is second only to malaria in terms of public health importance (WHO, 2005). Over one billion humans are at risk worldwide, and it is estimated that 200 million people worldwide are infected with the snail-transmitted, water-borne parasite leading to 1.53 million disability-adjusted life year (Gryseels et al., 2006). Also 20,000 deaths are associated with the severe consequences of infection, including bladder cancer or renal failure (Schistosoma *haematobium*) and liver fibrosis and portal hypertension (S. *mansoni*) (CDC, 2010). It is also easily transmitted from developing countries to developed countries (Hussein et al., 2008).

Schistosoma (Figure 5.1), are blood-flukes, including flatworms which are responsible for the parasitic infection which causes schistosomiasis (WHO, 2007). The worms responsible for the disease were discovered in 1851 by Theodor Bilharz, a German pathologist, from whom this disease took its name, Bilharziasis (Appleton et al., 1977).

Schistosoma needs fresh water to survive, so it can be found in and around water, i.e. rivers, streams, dams and lakes in endemic areas. Wherever people are in contact with water during their normal daily activities, such as recreation and irrigation, they are at risk. Children are nearly always the most heavily infected since they spend some of their time playing and swimming in water, and also because their natural immunity is less than in adults (Amin et al., 1995). Therefore, this disease has an especially detrimental effect on the growth and development of children (Hussein et al., 2008).

Adult worms can lay hundreds of eggs (Figure 5.2) each day while they are inside humans, and these eggs are excreted in human faeces or urine. This leads to the water becoming contaminated by Schistosoma. The parasite lifecycle is completed when these hatch into miracidia which then infect the intermediate snail host (Figure 5.3) (WHO, 2007). The snails release cercaria into the water which may infect other people who come into contact with this water.



Figure 5.1: Different single parasites of schistosoma (WHO, 2009)



Figure 5.2: Single egg of schistosoma (CDC, 2010)



Figure 5.3: Schistosoma snail host (CDC, 2010)

5.2 Species of Schistosomes and Geographical Distribution

Schistosomiasis is found in tropical countries in Africa, Caribbean, eastern South America, East Asia and in the Middle East (Figure.5.4) (WHO, 2004). There are five species of water-borne flatworm, or blood flukes, called schistosomes (WHO, 2007; CDC, 2010):

- Schistosoma *mansoni* which cause intestinal schistosomiasis, occurs in Africa, the Eastern Mediterranean, the Caribbean and South America,
- S. *Japonicum* group of parasites, which cause Oriental or Asiatic intestinal schistosomiasis. It is endemic in South-East Asia and in the Western Pacific region.
- S. *mekongi* in the Mekong river basin which cause Oriental or Asiatic intestinal schistosomiasis. It is endemic also in South-East Asia and in the Western Pacific region.
- S. *intercalatum* has been reported from central African countries. It causes another Intestinal schistosomiasis.

• S. *haematobium* is endemic in Africa and the Eastern Mediterranean. It causes Urinary schistosomiasis.



Figure 5.4: Distribution of schistosomiasis

5.3 Infection and Life Cycle of Schistosomiasis

Infection occurs when people come into contact with contaminated fresh water including infected snails (which carry schistosomes). The life cycle of the human schistosomes are generally similar, and involves many steps (CDC, 2010; Amin et al., 1995).

Figure 5.5 (CDC, 2010) shows the life cycle of schistosomiasis. The cycle begins when infected people urinate or defecate in fresh water. This water becomes contaminated by Schistosoma eggs. The eggs hatch releasing larva (miracidium), and this larva searches for appropriate snails in the water. The miracidium penetrate into snails, grow and develop inside the snails and transform into a mother sporocyst. About a week later this mother will give rise to many daughters which after 3-4 weeks will produce cercarae. These are sexually differentiated, into male and female. The

cercaria of schistosomes is usually released 25 to 30 days after the snail has been infected. It can penetrate the skin of people who come into contact with contaminated water. During penetration the cercarae lose their tails and become schistosomes. Within several weeks, the worms grow, multiply and develop inside the blood stream. They then pass to the right side of the heart and the lung and produce eggs when they reach the liver. Some of the eggs go to the bladder or intestines and are passed into urine or stools. This human cycle is usually completed in 30 to 45 days.



Figure 5.5: Life cycle of schistosomiasis (CDC, 2010)

5.4 Symptoms and Diagnosis of Schistosomiasis

Many infections are sub clinically symptomatic (CDC, 2010; WHO, 2005), with mild anaemia and malnutrition being common in endemic areas. Acute schistosomiasis may occur 7-14 weeks after the initial infection depending on the species of snail, parasite, climate variables, environment and source of water (Amin at al., 1995). There are two main stages of schistosomiasis; the early stage of schistosomiasis includes itchiness of the skin, bloody mucous stools, fever, cough, diarrhoea, abdominal pain and dysenteric attacks. The late stage of schistosomiasis

includes; Hepatosplenomegaly (the enlargement of both the liver and the spleen) (Figure 5.6), weight loss, severe liver disease, anaemia and jaundice. It can also cause bladder cancer, kidney problems or complications of the liver and spleen.

Microscopic identification of eggs in the stool or urine is the most practical method for diagnosis. The stool examination is the more common of the two (Amin et al., 1995).



Figure 5.6: The enlargement of the liver and the spleen (Medical Ecology, 2004)

5.5 Control of Schistosomiasis

WHO state that there are three main methods that may be used to control schistosomiasis (CDC, 2010; WHO, 2005; Amin at al., 1995): Parasite control (Treatment), control of transmission and health education.

5.5.1 Parasite Control (Treatment)

The main schistosomiasis treatment drug is praziquantel using a single oral dose annually. When outbreaks of schistosomiasis occur it is often not practical to test all individuals for schistosomiasis especially as some individuals may show few symptoms. Therefore WHO, 2005 has developed guidelines for community treatment of schistosomiasis based on the impact of the illness on children (children being more susceptible) in endemic villages:

1. If more than 50% of children have blood in their urine (a symptom of schistosomiasis) then everyone in the village should receive a treatment.

- If 20 to 50% of children have blood in their urine, only school-age children are treated
- 3. But if less 20% of children have these symptoms, mass treatment is not implemented.

5.5.2 Control of Transmission

Methods of control of transmission include snail control and prevention of human water contact.

Control of snail:

- Molluscicides: Snails can be controlled through the spraying of molluscicides in locations where snails breed. Examples include valleys, wells, dams, streams, vegetation, algae and any areas where individuals may be exposed to snails (Figure 5.7) (Amin at al., 1995). Hand or power spraying is the recommended method for stagnant or slow running water in ponds, marshes, lake shores and wells where the parasite and snails may be present. For flowing water in small rivers, streams, irrigation canals, the application of molluscicide through drip-feed methods in recommended. In these the chemical is applied continuously at one dispensing point upstream of any stretch of water to be treated and is carried downstream killing snails along the length of water course.
- Environmental management: This includes the elimination of snails and changing their habitats by removing vegetation, filling rain pools and removing snails (Figure 5.8) (Medical Ecology.org., 2004). Secondly, improving water supply and sanitation for communities, as the provision of safe water for drinking and domestic activities and adequate sanitation systems have been proved to be effective in reducing human water contact. This is the top intervention priority for the control of transmission (Medical Ecology.org., 2004). Safe drinking water can be provided through the use of deep and surface wells, and the filtration and treatment of stream water. Prevention of human water contact can also be effected by fencing off stretches of water, bridging streams and providing safe laundry facilities. Also improving sanitation facilities can also be effected (Amin at al., 1995).



Figure 5.7: Spraying of molluscicides in locations where snails breed in Asir, Saudi Arabia (The Vector Control Administration in Asir Region, 2007)



Figure 5.8: Removing vegetation, filling rain pools and removing snails in Asir (*The Vector Control Administration in Asir Region, 2007*)

5.5.3 Health Education

Co-operation among related agencies is required to communicate to people that this illness is dangerous to their health. People should be urged co-operate with the health team, avoid contact with water at high risk sites, and avoid urination or defecation in or beside water. The message can be spread through schools, gatherings e.g. religious gathering, or through face to face contacts in primary health care centres or any health point. Leaflets and posters at water contact points or schools are other methods through which such information can be spread. As children are most at risk education can be challenging because children love playing in water (Amin at al., 1995).

5.6 Seasonal Influences on the Transmission of Schistosomiasis.

Snails play a very important role in the transmission of schistosomiasis, and snail population sizes, rates of infection, and the production of cercaria are strongly influenced by climate. This therefore affects the rates of transmission of the disease. There are several environmental factors that can affect the density and viability of snail populations (Medical Ecology, 2004; Davis et al., 2002):

- Water levels, affected by rainfall, are critical as the optimal snail habitat occurs in a narrow zone of elevation above mean low water. In arid areas rainfall is required to provide suitable snail habitat. Annual floods in some environments can drown many, if not the majority of the adult snails. In areas where there is continuous flooding some species of snail live for about one year. Where there is no flooding, the species can live twice as long, and often longer.
- Temperature can also have an impact on the snail population as when temperatures in the water fall below 10°C or exceed 30°C both adult snails and their eggs may die.
- Elevation also plays an important role in the density of snail populations. The optimum habitats for the snails are flat, mid-level land with some dry-season ponds, and also thick grass.

5.7 Previous Studies

There are very few studies that look at the direct influence of weather upon schistosomiasis cases. Therefore this section will try to present the available studies which examine the relationship of weather conditions to snail populations followed by the parasite. The few studies on weather and schistosomiasis cases will then be presented. We will present studies from around the world before focussing upon those conducted in the Middle East and finally in Saudi Arabia and the study area Asir Region.

5.7.1 The Influence of Weather on the Snails Hosts

One study in Senegal examined seasonality in the transmission of schistosomiasis (Sturrock et al., 2001). Quantitative snail surveys were conducted over 3 years from 1992 at sites representing different habitats in and around an irrigation scheme. Water temperature affected snail survival and reproduction and this concurred with Appleton's research (1977). All snail populations grew rapidly and numbers peaked in late spring and early summer depending on the habitat. They then dropped sharply in the heat of the summer as water temperatures reached their peak. Numbers remained relatively low until the following spring, with most transmission between May and August each year.

Pimentel-Souza, et al (1988) studied the influence of experimental illumination and seasonal variation on the snail Biomphalaria Glabrata, a snail vector of schistosomiasis in Brazil. The data were counted throughout the year under two regimes (12h light and 12h dark from 7am to 10pm). The researchers state that, as found in some previous studies, water temperature has an influence on egg-laying, hatching and survival for the schistosomiasis vector snails. An increase of temperature during winter and autumn leads to more mating activity, but when the temperature starts to decrease the mating count decreased. Another study in Brazil (Pimentel-Souza et al., 1990) stated that a higher rate of egg production was observed at water temperatures between 20.0 and 27.5°C, and the number of eggs / snails / day reached a maximum within this temperature range. A statistically significant lower number of all these was observed at lower temperatures around 17.5°C.

Barbosa et al., (1987) studied the effect of temperature on the reproductive rate in snails as these are a vector of schistosoma manisoni. The snail data were collected between summer 1981 and spring 1982 and the experiments were in the middle of each season. Reproduction was measured by egg production, number of clutches and hatching, and the highest rates occurred during winter and low rates during the_summer. Although there was not a high correlation between the rates of reproduction and temperature, the highest mean number of hatching, clutches, and number of snails per day were recorded in winter. Following this lower numbers occurred during the autumn then spring with the lowest occurring during the summer. The variation in temperature for this study during the year was from 19.5 to 26.3°C.

This study confirmed that during winter the density of snails increases in tropical zones such as Egypt (El-Hassan, 1974), and Brazil (Barbosa, 1970). However it increases during summer in temperate zones such as South Africa. This is probably due to the optimum temperature for snails, as in temperate zones temperatures are most suitable in the summer (Appleton, 1977).

Michelson, (1960) studied the effects of temperature on the growth and reproduction of australorbis glabratus (a genus of planorbid freshwater snails) in the laboratory. Michelson mentioned that temperature may influence the ability of miracidia to infect snails and also the development of the schistosoma larvae within snails. It may also affect cercarial emergence and thus the infectivity of cercaria for the mammalian host. He states that the optimal temperature for rapid growth was at 30°C. However this range is different to that found in Brazil (Pimentel-Souza et al., 1990) which stated that a higher rate of egg production was observed at temperatures between 20.0 and 27.5°C.

A study of infections with larval stages in snails was carried out in Qena Upper Egypt (Hussein et al., 2008). The result of this study showed that the highest numbers of infected snails collected were during January followed by September, whereas the lowest number appeared in June as this is the hottest month in Egypt. The influence of climate was clearly shown and the number of snails peak in winter and autumn, and decrease in summer and spring.

In the Jequitinhonha Valley, Brazil, Kloos et al., (2004) studied the schistosoma in all aquatic snail habitats in a rural area. Snail and environmental surveys were carried out from July 2001 to November 2002. One of the major objectives of this study was to examine the seasonal distribution of schistosoma *mansoni* infections in this area. Larger snail numbers were found in standing waters from October to March during the rainy and hot season, but the densities were higher in streams and canals from April to September during the dry season. Also a significantly larger mean number of one of the snail species (*B. glabrata*) was found in the rainy season P < 0.001.

In a study in the Jos metropolis area of Nigeria, Agi (1995) noted that the population of *Bulinus sp.* snail increased during the summer with the peak occurring

in June and that the number of these snails decreased during March. The researcher mentioned that the seasonality of the snail population was attributed to changes in the rainfall pattern. The rainy season in this part of the country occurs during summer when more snails were found.

5.7.2 The Influence of Weather on the Parasite of Schistosoma

Adult infection-free schistosoma japonicum harboured in Oncomelania hupensis were collected from Jiangsu province in eastern China by Yang et al., (2007). In one of the experiments carried out under laboratory conditions, they investigated the effect of temperature on the development of *S. japonicum*, and have also tried to determine the lowest temperature at which the hibernation of *O. hupensis* occurs.

The findings state that a positive relationship exists between temperature and the development of *S. japonicum*. The effect of temperature on the prepatent (early stage) periods was highly statistically significant (P<0.001) as the average development rate / day of the snails increased with rising of temperature. At 21°C, 24°C, 27°C and 30°C the development rates were 0.008, 0.011, 0.014 and 0.016, respectively. The lowest thermal temperature derived from this regression is $15.3^{\circ}C$ (lowest temperature at which development occurs). These findings indicate that the transmission of schistosomiasis will be increased if climate change leads to a warmer China.

Coelho et al., (2006) evaluated the influence of temperature on schistosoma *mansoni* infection in Biomphalaria glabrata. Lower temperatures led to the lowest infection level. The influence of temperature upon infection at 15°C was significantly less than it was at 20°C (P < 0.01). At 20°C it was significantly less than it was at 30° C (P<0.01).

5.7.3 The influence of weather upon schistosomiasis illness

A recent epidemiological study for schistosomiasis disease was carried out in two local areas of Benue State in Nigeria (Houmsou et al., 2009). Seven hundred and fifty urine samples were obtained from different parts of the community between November 2008 and March 2009. The greatest risk and prevalence of infection was
during the dry season (P<0.01), but there was no significant difference in infection rates between one month and another. The highest incidence of schistosoma infection was recorded during February and March (52%) for each. The paper suggests that this may be due to the hot season in this area during these months.

In Morocco climate variability may have contributed to the prevalence of schistosomiasis during the 1980's. The severe drought in that period led to a lack of water, and a reduction in the number of intermediate snail hosts of the parasite of this illness. Also the lack of rainfall especially in the Ouarzazate area of Morocco, led to the drying out of the breeding sites of Bulinus truncatus, hence stopping the transmission cycle of the schistosomiasis (Laamrani et al., 2000).

5.7.4 Studies in Saudi Arabia

Several studies on schistosomiasis in Saudi Arabia have been carried out, but most of them do not examine the influence of weather in detail. Problems in the control of schistosomiasis in Asir Region have been studied by (Al-Madani, 1991). This study discussed some factors of this region such as geography, climate, hydrographical conditions, sources and spread of infection, and the role of primary health care centres. This information is used to establish appropriate plans for the control of this disease.

This study mentioned that rainfall and streams are the major sources of water in Asir region and that vector snail habitats are found in streams in the highlands. After heavy rainfall these streams carry the snails from the highlands down to the lowlands where the flood water accumulates and the snails exist in pools and dams. Sources of the infection of this illness are also mentioned, for example when expatriates from endemic countries come to work in this region and become an important source of infection. Also in the past there were not enough specialists and field technicians in the region to implement an appropriate program to control schistosomiasis. The paper makes many suggestions to improve the control program including; Activate the cooperation and coordination between different ministries and agencies involved in the control program and joint committee meetings with related department in this region (e.g. Water & Electricity Department, Agriculture and Municipality). Also provide safe drinking water supplies, an adequate sanitation system and provide health education to the residents. Furthermore specialized training programs should be established such as health education programmes and training for the professionals of the primary health care centres. Routine population check-ups for this disease should be carried out by these centres thus giving them a more active role in schistosomiasis prevention.

A recent study in Asir region, Saudi Arabia, has been carried out and considers the factors affecting the prevalence of schistosomiasis (Shati, 2009). He compared two sectors in Asir; Abha (highlands) and Tehama (lowlands), over eight years (2000-2007). Over the period a decline in the number of cases was noted and there were significantly more cases in the highlands in comparison to the lowlands. The infection rate in the highlands during summer was high, but there were no significant differences in infection rates between seasons in the lowlands. So the prevalence of the illness was significantly affected by season (summer) in the highlands only. The study mentioned that many biotic factors can play an important role in schistosomiasis prevalence such as rainfall which usually increases during summer in some parts of the highlands. This increase can be followed by a rise in schistosomiasis cases in the highlands area. As was found in other studies, the researcher noted that the increase of schistosomiasis prevalence in this area usually peaks during the wet season. In the highlands the temperature during winter can drop to less than ten degrees, and the literature review has demonstrated that such temperatures are unsuitable for schistosomiasis transmission. However in the lowlands temperatures during winter season are more suitable. In the summer temperatures are very high in the lowlands which may provide unsuitable conditions for schistosomiasis transmission. Thus temperature can affect schistosomiasis transmission. The study showed that there was a decrease in the prevalence during the study period, and this may be due to Ministry of Health control program.

5.7.5 Literature Review Summary

Table 5.1 summarises the main studies (statistically) of the correlation between schistosomiasis or snails and climate variables.

No	The author	Place	Dependent Variable	Independent Variable	The relationship & Lag time (month)
1	Appleton, 1977	Senegal	Number of snails	Water Temperature	Negative
2	Pimentel-Souza, et al 1988	Brazil	The mating count of snail	Water Temperature	Positive
3	Barbosa et al., 1987	Tropical. South Africa	density of snails	Water Temperature	Negative in tropical Positive in South Africa
4	Hussein et al., 2008	Qena Upper Egypt	numbers of infected snails	Water Temperature	Negative
5	Kloos et al., 2004	Jequitinhonha Valley, Brazil,	numbers of snails	Water Temperature Rainfall	Positive Positive
6	Agi, 1995	Nigeria	numbers of snails	Water Temperature Rainfall	Positive Positive
7	Yang et al., 2007	Eastern China	Snails (S. japonicum)	Temperature	Positive
8	Coelho et al., 2006	laboratory conditions	schistosoma mansoni	Temperature	Positive
9	Houmsou et al., 2009	Benue State, Nigeria	infection rates	Temperature Rainfall	Positive Negative
10	Laamrani et al., 2000	Morocco	Transmission of schistosomiasis	Rainfall	Positive
11	Al-Madani, 1991	Asir Region, Saudi Arabia	abundance of snails	Rainfall	Positive
12	Shati, 2009	Asir Region, Saudi Arabia	schistosomiasis infection	Temperature	Positive in highlands Not significant in Lowland

Table 5.1: Summary of the main studies relating to schistosomiasis

The literature review highlights a number of important points:

- 1. There are few studies which look at the influence of weather upon schistosomiasis cases.
- 2. The weather influences vary by location.
- 3. Time lags between weather and infection are likely to be between 3 and 9 months depending on the location and type of schistosomiasis, weather variables and other factors.
- 4. Different climate variables such as temperature and rainfall may influence schistosomiasis.
- 5. Some studies investigated the influence of weather upon the populations, survival and reproduction of the schistosomiasis host (snail). Others found that the host abundance has an effect upon the incidence of the illness.

6. Some studies showed that climate variables have an impact upon schistosomiasis incidence. However, the impact of specific weather variables can be either positive or negative depending upon the geographical setting and season.

Based on the findings of the previous studies, it is clear that there is an urgent need to research the influence of weather upon the host (snail) and parasite of schistosomiasis in the endemic areas of Saudi Arabia such as Asir Region. Building upon the literature review such a study should examine a wide range of weather variables and also consider that any relationships with the weather may vary seasonally. A statistical analysis for this disease seems the most appropriate study design and this may need to consider factors such as autocorrelation in the data.

5.8 Schistosomiasis in Saudi Arabia

There are three main species of schistosomiasis in this country; urinary, intestinal and combined schistosomiasis. The geographical distribution during 2007 for urinary schistosomiasis (MOH, 2008), shows that this illness is endemic in Makkah, Madinah, Jazan, Asir and Lithe Region. However intestinal schistosomiasis is endemic in Taif, Albahah, Asir, Bishah, Najran, Makkah, Madinah, Hail, Jeddah and Jazan Regions. The snail responsible for both types of schistosomiasis exists only in Jazan, Makkah, Asir and Madinah Region. Combined schistosomiasis has been recorded in Asir Region (MOH, 2008).

Schistosomiasis used to be one of the most important public health problems in Saudi Arabia, especially in southern regions. In terms of trends over time Figure 5.9 presents the number of schistosomiasis cases in Saudi Arabia from 1980 to 2009. This figure shows a sharp decrease of cases throughout this period, with the highest number of cases occurring in 1983 (25985 cases). There was a dramatic increase from 1980 (9880 cases) to 1983. The number of cases in 2009 is less than 3% of the level in 1980.



Figure 5.9: The number of (schist) cases in Saudi Arabia (1980-2009)

The prevalence rates of this illness in 2007, 2008 and 2009 were 1.52/100,000, 2.80/100,000 and 1.14/100,000 respectively. Urinary schistosomiasis in all of Saudi Arabia in 2007 was 24.6% of all cases while intestinal schistosomiasis was 75% and the combined illness represented just 0.4% (MOH, 2010).

The statistical year book (MOH, 2006) states that the highest percentage of all cases types in 2005 was found in Asir Region (39%), followed by Jazan, Bishah and Albahah Regions (27.6%, 16.3% and 9.9% respectively) (Figure 5.10). All these regions are located in the south of Saudi Arabia.

One study in Asir Region (AI-Madani, 1991), reports that both intestinal and urinary schistosomiasis have been reported. The topography of Asir region and different weather conditions between sectors create difficulties for schistosomiasis control in the Region. However, improvements to primary health care centres are the most important factor for better control of this disease (AI-Madani, 1991).



Figure 5.10: Comparison of schistosomiasis among regions of Saudi Arabia 2005 (MOH, 2006)

5.9 Schistosomiasis in Asir Region (Study Area)

This study will focus on schistosomiasis cases in Asir Region and its subsectors. The prevalence of this illness and its transmission is influenced by little known geographical and climate factors such as temperature, rainfall and humidity (AI-Madani, 1991; Amin et al., 1995; Shati, 2009).

The topography of Asir Region may be one of the main factors which contribute to the difficulty in controlling of schistosomiasis. This topography helps in disseminating the snails from the highlands to lower areas thus repopulating free habitats in the lowlands with the intermediate hosts (Al-Madani, 1991). Also the presence of many inaccessible water sources in the highlands which are infested with snails makes control difficult. Additionally some parts of this region are inhabited by bedouins who move from one area to another searching for water. These people use water in the valleys as the main source for their daily activities. It is therefore difficult to stop schistosomiasis transmission as valley water is part of their everyday live. Education may also be problematic. Within Bedouin tribes it is also difficult to locate the infected individuals and provide them proper treatment. Without treatment these individuals become part of the life cycle of schistosomiasis (Almadini, 1991; Shati, 2009).

5.10 Topographical Classification of Asir Region and Endemic Areas

Topographically, climatically and endemically this region can be divided into three main sectors in terms of schistosomiasis transmission (Dr A. Abdoon 2007 pers. Comm.) (Figures 5.11 and 5.12):

- Highlands Areas: a series of high lands which extend from the north of Asir Region to the south along the coastal plains of the Red Sea (2000 -~3000m above sea level). It is one of the coldest areas in Saudi Arabia and has the highist annual rainfall (500mm), raining almost every month, in spring and summer (MEPS, 2007; Subyani, 2000). This area covers districts such as Abha, Khamise, Tanoma, and Dhahran. The annual average temperature for this sector is 17.7^oC, and the annual average relative humidity 53.1%. These areas have the highest occurrence of schistosomiasis in and it can be divided into two sections depending on the endemicity:
 - a. Southern Asir Sarawat Mountains and their sectors
 - b. Sabt-Alalaya Sector (North)
- 2. Asir Plateau (a part of Highlands); is in the east of Asir region with an elevation range between 1000 and 2000m decreasing to the east. This area covers the districts such as Tathlith, Almadha and Bishah. The annual average temperature for this sector is 19.5°C, with 250mm annual rainfall and annual average relative humidity of 44.4%. This sector has many wadies all flowing eastward. According to local sources (Dr A. Abdoon 2007 pers. Comm.) there are few cases of schistosomiasis in this area as a result of a low amount of rainfall during the last two decades in this sector.
- 3. Tehama (Lowlands): This area receives an annual rainfall of more than 350mm, and temperature ranging between 25°C and 49°C, with relative humidity varying from 55% in summer to 70% in winter. Depending on the altitude this area can be divided into two areas: foothills between 300 and 900m such as Al-Majaredah, Muhail and Rejal-Alma. This area has a moderate occurrence of schistosomiasis. However the coastal area which is between 0 and 200m (Costal Plain Red Sea)

has the fewest cases of schistosomiasis in the region such as Al-Gahma and Al-Berk.

5.11 Rationale and Hypotheses

The influence of weather upon schistosomiasis cases is important to understand for a number of reasons:

- Based on the findings of this study and an understanding of how schistosomiasis is affected by weather it should be possible to design an early warning system to predict schistosomiasis cases. This would also be useful for health service planning and hence to develop good and effective control programmes, for example identifying periods when enhanced molluscicides spraying is required.
- 2. If weather alters under climate change then this data could help predict the impact of this change on schistosomiasis in Asir Region.
- 3. It can help clarify the important modes of transmission in the region.
- 4. It improves our understanding of disease transmission dynamics.

This study tests the hypotheses that there are relationships between weather (maximum, mean and minimum temperature, rainfall and relative humidity) and schistosomiasis cases in Asir Region. In addition this study was also able to make use of a dataset of snail abundance to examine how numbers of the host in Asir Region are affected by the same weather factors. It also enabled us to examine the relationships between schistosomiasis cases and snail numbers.

5.12 Data and Method

5.12.1 Sources of Data

Official letters were issued, instructing authorized personal to collect health and weather data. The numbers of schistosomiasis cases and snails in each month from the first of Jan 1998 to the end of Dec 2009 for the whole of Asir Region were obtained from the Vector Control Administration, Health Affairs Directorate in Asir. The data represents all reported cases of schistosomiasis in Asir region and have been collected from all hospitals and primary health care centres (public or private) and the Main Parasitology Laboratory in Abha (Annual reports, 1998- 2009). The reported schistosomiasis cases were confirmed in different ways depending on the type of this disease. Urinary schistosomiasis cases were confirmed by a person with visible haematuria or with a positive reagent strip for haematuria or with the eggs of S. haematobium in urine (microscope). Intestinal schistosomiasis were confirmed by a person with eggs of S. mansoni, or S. japonicum / mekongi in stools (microscope) whether chronic or recurrent intestinal cases. This study will not consider each type of this disease separately but will consider both types together for the statistical analysis. The snail data were collected by teams from the Vector Control Administration and Primary Health Care Centres in different endemic areas across Asir region. The control program for Schistosomiasis in Asir consists of collecting and removing snails from areas where they are likely to occur. These include valleys, wells, streams, water pools and dams. It is acknowledged that these methods of control have not been very effective (Al-Madani, 1991). As part of this process the removed snails are counted and typed. These counts are the input data used in this study. The professionals and experts of this disease with their field teams collect snails throughout the year particularly during the seasons when snails are anticipated (Dr A. Abdoon 2007 pers. Comm.; Annual reports, 1998-2009).

Weather data for Khamise, the lowlands and Sabt-Alalaya sector were obtained from sources mentioned in the last 2 chapters. All the locations of weather stations in Asir Region are illustrated in Figure 5.12. The aim of this collection was to obtain corresponding weather data for each sector within Asir. However, this proved impossible due to the limited availability of data for many sector within Asir.

5.12.2 Classification of Data:

As a consequence of wide differences in topology and weather within Asir, there are likely to be differences in schistosomiasis between sectors. There are fourteen sectors in Asir (Figure 5.11): Abha, Khamise, Ahad-Rifadah, Al-Madhah, Tathlith, Sarat-Abedah, Dhahran-Aljanoub, Tanoma, Rejal-Alma, Majaredah, Muhail, Al-Berk, Al-Gahma and Sabt-Alalaya.

The limited weather data available across Asir imply that it is not possible to examine the associations between schistosomiasis cases and weather for each endemic sector in Asir individually. It is therefore, necessary to group sectors together. Differences in incidence between sectors may be due to differences in reporting. However, varying seasonalities are more likely to be due to transmission differences between sectors. To group sectors together graphs of schistosomiasis seasonality in each sector were produced Figure 5.11. However, these did not reveal consistent seasonality between groups of sectors partly due to the low numbers of cases in subsectors. This made the grouping of sectors, using the method used in previous chapters impossible.

Therefore, sectors were grouped depending on the topography of this region, and also according to the classifications of the Vector Control Administration, Health Affairs Directorate in Asir. Shati, (2009) uses similar classification of Asir Region. Groupings were created which were highlands, lowlands, Sabt-Alalaya. Using these three groups four models were constructed. The first was an analysis of schistosomiasis in the whole of Asir. Then three models of schistosomiasis cases of the sectors; the highlands, the lowlands and in Sabt-Alalaya sector. Further details are provided below:

- Schistosomiasis cases in overall Asir Region against Khamise weather data (Maximum, mean and minimum temperature, rainfall and relative humidity, Station 1 Figure 5.12). Khamise is the second main city in the region, and covers the largest occupied sector in the middle of this region (MOH, 2006). Weather data was available for all months.
- Schistosomiasis cases in the highlands group (eight sectors): Abha, Khamise, Ahad-Rifadah, Al-Madhah, Tathlith, Sarat-Abedah, Dhahran-Aljanoub, and Tanoma. Khamise weather data will represent this group as well (See station 1 Figure 5.12).
- 3. Schistosomiasis cases in the lowlands group (five sectors): Rejal-Alma, Majaredah, Muhail, Al-Berk and Al-Gahma. Majaredah temperature (See station 2 Figure 5.12) and rainfall of Rejal-Alma (See station 3 Figure 5.12, data available for 132 out of 144 months) will represent this group. No humidity data was available for this area.
- Schistosomiasis cases in Sabt-Alalaya Sector, Sabt-Alalaya station weather data will represent this sector. (See station 4 Figure 5.12). No humidity data was available for this area.

Monthly snail numbers are available for the whole of Asir only, Therefore in the final model these data will be analysed against Khamise weather data (Maximum, mean and minimum temperature, rainfall and relative humidity, Station 1 Figure 5.12).



Figure 5.11: Monthly Average of Schistosomiasis Cases in Sectors of Asir Region (1998-2009) Adapted from (CDSI, 2005)



Figure 5.12: Topographic map for Asir Region including the weather stations

5.12.3 Schistosomiasis Data Description

The monthly averages of schistosomiasis cases in the whole of Asir Region during the study period are presented in Figure 5.13 and show transmission of the parasite throughout the year. However, there is some seasonality with peak incidence between March and June, with the highest average number of cases in May (24.83) followed by April (24.75). The average number of cases during the autumn and winter are lower, with the lowest number occurring in November (9.83). The standard deviation bars on the graph indicate that there is a large variability from year to year in monthly cases of schistosomiasis.



Figure 5.13: Monthly Averages of schistosomiasis Cases in All Asir Region (1998 - 2009)

Figure 5.14 shows the monthly averages of schistosomiasis cases in the highlands group during the study period. There is a clear seasonality with peak averages of cases between February and June, with the highest average cases in May (13.83) followed by June (13.58). The average numbers of cases during the late summer and autumn and early winter are lower, with the lowest number of cases in November (5.17). The standard deviation bars on the graph indicate that there is a large variability from year to year.



Figure 5.14: Monthly averages of schistosomiasis cases of the highlands group (1998 - 2009)

Figure 5.15 shows monthly averages of schistosomiasis cases in the lowlands group in Asir Region during study period. There is less seasonality that in the highlands group with the peak averages of cases occurring between February and June, with the highest average cases in May (4.92) followed by March (4.83). The average number cases in the rest of the year were a little lower with a small peak in October (but this may simply be random variation), and the lowest averages of cases was in September (1.5). The standard deviation bars on the graph indicates that there is a large variability from year to year.



Figure 5.15: Monthly averages of schistosomiasis cases in the lowlands group (1998 - 2009)

Figure 5.16 shows monthly average number of schistosomiasis cases in Sabt-Alalaya sector, which is in the north of the highlands in Asir Region, during the study period. There is a little seasonality in this sector, with the peak averages of cases occurring between March and August, with the highest average cases in April (7.33) followed by August (7.17). The lowest average of cases was in November (2.42). The standard deviation bars on the graph indicates that there is a large variability from year to year.



Figure 5.16: Monthly averages of schistosomiasis cases in Sabt-Alalaya, Asir Region (1998 - 2009)

5.12.4 Snail Data Description

The monthly average number of snails in all Asir Region during the study period is presented in Figure 5.17. There is a clear seasonality with peak incidence between January and June, with the highest average number in May (117.5) followed by April (114.8). The average number in the next half of the year is lower, with the lowest number occurring in December (68.9). The standard deviation bars indicate that there is variability from year to year in the monthly number of snails.



Figure 5.17: Monthly averages of snail numbers in all Asir Region (1998 - 2009)

5.12.5 Temperature Data Description

The temperature data for Khamise City will be used in the analysis of cases across the whole of Asir Region as well as the highlands group. It will also be used for the snail data. Descriptions of these data were presented in Section 3.11.3. Although this analysis is for a different year range (1998-2009 vs. 1995-2006) the seasonal trends in these data are similar to those presented previously.

The temperature in Majaredah will be used as representative of conditions in the lowlands group and these data were again presented previously for a slightly different range of years in Section 3.11.3.

The temperature in Sabt-Alalaya station will be used as representative of conditions in this area. These data have not been used in previous chapters. From this station only maximum and minimum temperature are available and the data are shown in Figure 5.18. The highest average maximum temperature occurs in June $(30.6^{\circ}C)$ and the highest minimum in July $(18.1^{\circ}C)$. The lowest average maximum and minimum temperature occurs in January (17.4 and 8.7°C respectively).



Figure 5.18: Maximum and minimum temperature in Sabt-Alalaya (1998 -2009)

5.12.6 Rainfall Data Description

Khamise rainfall will be assumed to be representative of all Asir and the highlands group. The monthly average rainfall in this city during the study period was presented in Section 3.11.4 for a slightly different yearly range. Rejal-Alma is assumed to be representative of rainfall of the lowlands group, and similar data for this weather station was presented also in Section 3.11.4.

The monthly average rainfall in Sabt-Alalaya sector over the study period in Figure 5.19 shows a slight seasonality with a peak incidence between November and January and also between March and May. The highest averages were in April (49.1mm) followed by May (37.3 mm). However the average rainfall between June and October also in February are lower. The lowest average occurs in July (2.4mm), with a slight peak in August (12.2mm). The standard deviation bars on the graph indicate that there is an extreme variability from year to year.



Figure 5.19: Monthly averages rainfall in Sabt-Alalaya sector (1998 - 2009)

5.12.7 Relative Humidity Data Description

The relative humidity in Khamise will be assumed to be representative of all Asir and the highlands group, and these data were presented previously in section 3.115.

5.12.8 Statistical Analysis Data Preparation

The analysis to examine the relationship between monthly cases of schistosomiasis and the selected weather data proceeded in a number of stages. Identical stages were used for analysis of the snail data.

A histogram of the number of schistosomiasis cases each month in the whole of Asir Region was produced. There are few months with more than 40 cases. The skewed nature of the distribution may cause problems when performing regression analysis and so the natural logarithm of schistosomiasis cases (adding 1 to zero values) was performed. This produced Figure 5.20, and descriptive statistics of this distribution are presented in Table 5.2 and demonstrate that taking the natural logarithm of schistosomiasis produces a variable that is nearly normally distributed. The same action has been carried out for the highlands group, the lowlands group, Sabt-Alalaya cases and snail numbers in all Asir. Figure 5.21 shows histograms of the highlands after transformation. Table 5.3 presents the descriptive statistic of this distribution. Figure 5.22 shows histograms of the lowlands after transformation. Table 5.4 presents the descriptive statistic of this distribution. Figure 5.23 shows histograms of the Sabt-Alalaya sector after transformation. Table 5.5 presents the descriptive statistic of this distribution. Figure 5.23 shows histograms of the Sabt-Alalaya sector after transformation. Table 5.5 presents the descriptive statistic of this distribution. Figure 5.24 shows histograms of the snail numbers of all Asir after transformation. Table 5.6 presents the descriptive statistic of this distribution. In terms of the independent variables, rainfall for each group, also demonstrated a skewed distribution. Therefore, we took the natural logarithm of rainfall (adding 1 to zero values) to get (Ln rainfall) for each one.

The schistosomiasis data were then adjusted for any effects that could bias the results. Over time the number of schistosomiasis cases in Asir has declined and this long term trend is shown in Figure 5.25. Since the middle of 2001 the number of cases has declined from the peak number of cases in June. The long term trend in schistosomiasis in the highlands, lowlands and Sabt-Alalaya are shown in Figures 5.26, 5.27.and 5.28 respectively. The long term trend in snail numbers in the whole of Asir Region is shown in Figure 5.29. Over time the number of snails in Asir is almost constant with only a small decline over time.

In order to control for this effect, for the whole of Asir, highlands, lowlands and Sabt-Alalaya, the natural logarithm of schistosomiasis cases were put into a regression against time (each month numbered from 1 to 144). The unstandardised residual from this model represent the natural logarithm of schistosomiasis cases with the long term trend in incidence removed. The same procedure was applied to the snail data to produce the natural logarithm of snail counts with the long term trend in incidence removed.

The final stage in the analysis was to examine whether the monthly numbers of snails were related to monthly cases of schistosomiasis and this was carried out using regression analysis.



Figure 5.20: Frequency of natural logarithm of schist (Ln. Schist) in Asir Region (1998 – 2009)

Table 5.2: Descriptive statistics for Ln. schist cases in Asir Region

	N	Minimum	Morimum	Moon	Std.	Skewness		
Ln. Schist.	IN	winninum	Maximum	Mean	Deviation	Statistic	Std. Error	
	144	0.0	4.25	2.62	0.906	- 1.011	0.202	



Ln.Highlands.Sch

Figure 5.21: Frequency of Natural Logarithm of Schist (Ln. Schist) in Highlands (1998 – 2009)

Ln. Highlands Schist	N	Minimum	Marimum	Maan	Std.	Skewness		
Ln. Highlands Schist	1	WIIIIIIIII	waxiiiuiii	Mean	Deviation	Statistic	Std. Error	
	144	0.0	4.04	1.94	0.957	371	.202	

Table 5.3: Descriptive Statistics for Ln. Schist Cases in Highlands



Figure 5.22: Frequency of Natural Logarithm of Schist (Ln. Schist) in Lowlands (1998 – 2009)

Table 5.4: Descriptive Statistics for Ln. Schist Cases in Lowlands

	NT	M::	Minimum Maximum		Std.	Skewness		
Ln. Lowlands Schist	1	WIIIIIIII	waxiiiuiii	Mean	Deviation	Statistic	Std. Error	
Semst	144	0.0	3.04	1.488	0.820	434	.202	



Figure 5.23: Frequency of Natural Logarithm of Schist (Ln. Schist) in Sabt-Alalaya (1998 – 2009)

Ln. Highlands Schist	N	Minimum Maximum		Maan	Std.	Skewness		
Ln. Highlands Schist	1	MIIIIIIIIIIIIIIIII	waxiiiuiii	Mean	Deviation	Statistic	Std. Error	
	144	0.0	3.04	1.102	0.838	.209	.202	

Table 5.5: Descriptive Statistics for Ln. Schist Cases in Sabt-Alalaya



Figure 5.24: Frequency of natural logarithm of snail (Ln. Snail) in Asir (1998 – 2009)

Table 5.6: Descriptive statistics for Ln. snail in Asir

Ln.Snail All Asir	N	Minimum	Marimum	Maan	Std.	Skewness		
Ln.Snail All Asir	n	WIIIIIIII	waxiiiuiii	Mean	Deviation	Statistic	Std. Error	
	144	2.94	5.37	4.40	.394	548	.202	



Figure 5.25: Monthly schistosomiasis cases in all Asir Region (1998 – 2009)



Figure 5.26: Monthly schistosomiasis cases in the highlands group (1998 – 2009)



Figure 5.27: Monthly schistosomiasis cases in the lowlands group (1998 – 2009)



Figure 5.28: Monthly schistosomiasis cases in Sabt-Alalaya sector (1998 – 2009)



Figure 5.29: Monthly snail numbers in the whole of Asir Region (1998 – 2009)

5.12.9 Statistical Analysis

According to the literature review the relationship between schistosomiasis and weather may vary depending on the time of the year. The same may be true for the snail data. Therefore, the analysis proceeded in a number of stages: Depending on the trend of schistosomiasis cases in Asir region and the other groups we separated the year into different time periods. In the Asir region the schistosomiasis was separated into three time periods: Feb to May, June to September and October to January (See Figure 5.13). The highlands group was separated into two time periods November to April and May to October (See Figure 5.14). The lowlands group was separated into two time periods; June to November and December to May (See Figure 5.15). Sabt-Alalaya sector was separated into two time periods May to October and November to April (See Figure 5.16). The snail numbers during the study period also was separated into two times, May to November and December to April (See Figure 5.17).

For each of time periods linear regressions of DT Ln schistosomiasis or DT Ln snails against each of the lagged weather variables were created (lagged up to 12 months). Longer time lags than used in the malaria and leishmaniasis chapter were produced as the incubation period for schistosomiasis and its lifecycle is more uncertain and may be longer. We then examined these relationships and combined the weather from the two or three months with the strongest association. In most cases the strongest relationships varied depending on the weather variables examined and the sector the model was produced for. Because these models contained data from multiple months it became important to control for seasonality. Therefore, all models were analyzed, incorporating dummy variables for each of the individual months. Finally in these analyses it was not possible to assume that the schistosomiasis rate in one month is unrelated to the schistosomiasis rate in the next month, and this is known as autocorrelation. Therefore the schistosomiasis in the previous month was added into these models as an independent variable. An identical procedure was used for the analysis of DT Ln snails.

Due to the low numbers of schistosomiasis cases and snails in some months and the problems that this might cause in a linear regression, an ordinal logistic regression of Ln schistosomiasis cases and Ln snails was additionally applied. This was achieved by recoding Ln schistosomiasis and Ln snails into four categories of increasing schistosomiasis incidence or snail numbers. These were then used in an ordinal logistic regression to produce an analysis that is more robust to the distribution of the data. Multiple regression analysis was examined in the final stage of the statistical analysis. This method describes the effect of the two or three weather variables together against DT Ln of schistosomiasis cases or DT Ln snail numbers. The purpose of this test is to identify which weather variable has more influence upon schistosomiasis cases and snail numbers than others.

All of the data were analyzed using SPSS v.16 / v.18 for Windows and Microsoft Excel 2003 / 2007.

5.13 Results

The analysis is presented by first looking at relationships between schistosomiasis and temperature (maximum, mean and minimum) followed by rainfall and relative humidity. This is performed for whole Asir, highlands, lowlands and Sabt-Alalaya sector separately, but only where meteorological data is present. An identical sequence of models is then produced for the snail data against weather predictors. Finally schistosomiasis incidence is compared to snail numbers using regression analysis.

The models which do not have any significant relationship are not presented in this chapter but appear in Appendix C.1. The results for the combined lags are presented on the final line of each table.

5.13.1 Schistosomiasis with Temperature Data

This section presents the significant results for the relationships between schistosomiasis cases and temperature in all Asir, highlands group, lowlands group and Sabt-Alalaya sector. Linear regressions and ordinal logistic of detrended logarithm schistosomiasis cases against temperature data will be illustrated. Maximum temperature is considered first, followed by mean and minimum temperature. Maximum temperature for overall Asir model between June and September is presented in Table 5.7

	Linear Regression Ordinal Log							Logist	gistic			
lel	Ma	ıx.	Ma	1X. rature	Ma tempe with	ax. rature	Ma	ax.	Ma	IX. rature	Ma tempe with	ax. rature
Mod	tempe on	rature ly	wi	th	correl	ation	tempe on	rature ly	wi	th	correl	lation
			seaso	nality	seasor	ia nality			seaso	nanty	seaso	na nality
	Cof.	Р	Cof.	Р	Cof.	P	Cof.	Р	Cof.	Р	Cof.	P
0 P	41	.818	52	.797	89	.641	.13	.721	.12	.818	.06	.910
1 P	.81	.311	-1.2	.57	61	.714	.08	.615	.02	.954	02	.955
2 P	1.5	.048	1.6	.284	1.3	.353	1.1	.013	.96	.012	.96	.013
3 P	1.4	.063	1.7	.136	1.3	.129	.83	.041	.57	.072	.38	.063
<mark>4 P</mark>	<mark>2.8</mark>	<mark>.011</mark>	<mark>2.6</mark>	<mark>.034</mark>	<mark>2.4</mark>	<mark>.030</mark>	<mark>1.2</mark>	<mark>.011</mark>	<mark>.72</mark>	<mark>.017</mark>	<mark>.686</mark>	<mark>.023</mark>
5 P	.63	.414	.45	.715	-1.5	.895	.33	.413	.23	.359	.17	.654
6 P	71	.663	65	.568	27	.793	.13	.971	.02	.936	.02	.944
7 P	.41	.654	37	.766	.21	.855	.28	.533	.14	.626	.19	.506
8 P	.65	.791	.46	.743	.84	.517	.09	.777	.05	.855	.12	.709
9 P	.2	.818	12	.939	.13	.930	03	.875	08	.801	01	.977
10 P	2.9	.081	3.1	.244	3.3	.718	.83	.117	.72	.206	1.1	.071
11 P	3.3	.036	4.1	.061	2.8	.190	1.3	.008	.96	.042	.77	.110
12 P	.21	.294	.16	.940	73	.713	.81	.376	.44	.351	.20	.685
<mark>3-4 P</mark>	<mark>3.6</mark>	<mark>.001</mark>	<mark>4.0</mark>	<mark>.021</mark>	<mark>3.5</mark>	<mark>.029</mark>	<mark>1.0</mark>	<mark>.008</mark>	<mark>1.2</mark>	<mark>.006</mark>	<mark>1.0</mark>	<mark>.008</mark>
<mark>2-4 P</mark>	<mark>4.0</mark>	<mark>.031</mark>	<mark>4.67</mark>	<mark>.021</mark>	<mark>4.0</mark>	<mark>.032</mark>	<mark>1.2</mark>	<mark>.010</mark>	<mark>1.9</mark>	<mark>.001</mark>	<mark>1.6</mark>	<mark>.004</mark>

 Table 5.7: DT Ln schist and maximum temperature in all Asir (June to Sep)

0: current month. P: Previous month. Cof: Coefficient

The results indicated that in Asir over June to September schistosomiasis cases appear to be positively associated with maximum temperature in the 3 - 4 previous months in all variants of the model for both linear and ordinal logistic regression. A positive association is also observed for the 2 - 4 previous months. The results for maximum temperature against schistosomiasis cases in the highlands, the lowlands and Sabt-Alalaya sector did not show any consistency in their significant associations. The results for mean temperature in the all Asir model between June and September are presented in Table 5.8.

		Li	near Re	egressi	on			0	rdinal	Logist	tic		
Model	Me tempe on	ean rature ly	Mean temperature with seasonality		Me tempe with correl ar seaso	Mean temperature with auto correlation and seasonality		Mean temperature only		ean rature th nality	Mean temperature with auto correlation and seasonality		
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	
0 P	1.5	.555	1.3	.652	1.95	.456	1.0	.545	.08	.899	.03	.598	
1 P	-2.7	.265	-2.5	.281	-2.5	.243	07	.225	06	.910	03	.953	
2 P	.90	.624	1.1	.584	.78	.683	.90	.614	.81	.098	.88	.070	
3 P	1.4	.038	1.9	.102	1.5	.094	.34	.068	.52	.036	.28	.064	
<mark>4 P</mark>	<mark>2.3</mark>	<mark>.079</mark>	<mark>2.2</mark>	<mark>.027</mark>	<mark>2.3</mark>	<mark>.047</mark>	<mark>.23</mark>	<mark>.048</mark>	<mark>.74</mark>	<mark>.041</mark>	.75	.070	
5 P	-2.5	.160	-2.7	.134	-2.4	.145	55	.268	37	.342	53	.193	
6 P	-2.1	.448	-2.1	.269	52	.773	-2.1	.444	38	.333	50	.903	
7 P	3.9	.018	4.4	.026	3.5	.069	1.9	.019	1.05	.029	.90	.060	
8 P	-1.7	.632	-1.5	.510	.79	.725	41	.662	496	.324	.24	.696	
9 P	-3.3	.204	-4.2	.119	-2.7	.303	-1.3	.224	-1.2	.061	85	.189	
10 P	2.7	.447	2.4	.441	2.2	.440	.17	.437	.06	.928	.66	.351	
11 P	2.1	.506	2.9	.438	1.1	.715	.21	.516	.38	.579	.231	.737	
12 P	-4.4	.098	-4.9	.111	-4.8	.084	44	.088	66	.324	63	.346	
<mark>3-4 P</mark>	<mark>1.8</mark>	<mark>.047</mark>	<mark>2.2</mark>	<mark>.049</mark>	1.9	.086	<mark>2.9</mark>	<mark>.053</mark>	<mark>.65</mark>	<mark>.038</mark>	.70	.061	

Table 5.8: DT Ln schist and mean temperature in all Asir (June to Sep)

The results indicated that in all Asir between June and September schistosomiasis cases appear to be positively associated with mean temperature in the 3 - 4 previous months in some, but not all variants of the model.

The results for mean temperature against schistosomiasis cases in the highlands, the lowlands and Sabt-Alalaya sector do not show any consistency in their significant associations.

Minimum temperature for the all Asir model between June and September is presented in Table 5.9.

		L11	near Re	egressi	on			0	rdinal	al Logistic		
					М	in					M	in
5	M	in	М	in	tempe	rature	м	in	Μ	in	temper	rature
ode	temne	rature	tempe	rature	with	auto	temne	ni rature	tempe	rature	with	auto
Ŭ	on	lv	wi	th	correl	lation	on	lv	wi	th	correl	ation
, ,	-	5	season	nality	an	nd		5	season	nality	an	d
	a c	D	<u> </u>	D	season	nality	a c	D	a c	D	season	nality
	Cof.	P	Cof.	P	Cof.	Р	Cof.	P	Cof.	Р	Cof.	Р
0 P	1.2	.432	42	.736	63	.802	2.1	.532	14	.829	22	.720
1 P	-1.5	.325	-2.3	.371	-1.2	.632	-2.1	.336	39	.473	18	.749
2 P	2.2	.496	1.6	.528	1.0	.658	1.9	.396	.48	.423	.61	.314
<mark>3 P</mark>	<mark>1.3</mark>	<mark>.015</mark>	<mark>3.8</mark>	<mark>.040</mark>	3.0	.057	1.6	.043	.67	.061	.58	.087
<mark>4 P</mark>	<mark>2.0</mark>	<mark>.037</mark>	<mark>.17</mark>	<mark>.044</mark>	.21	.063	2.6	.047	.01	.078	.09	.086
5 P	-2.2	.163	-3.1	.029	-2.2	.105	-2.3	.132	.69	.046	.69	.050
6 P	-2.5	.288	-1.4	.282	65	.600	-1.5	.278	44	.111	25	.396
7 P	4.8	.034	2.7	.039	2.6	.028	2.6	.040	.92	.004	.91	.004
8 P	89	.427	67	.656	.57	.691	64	.400	22	.484	.229	.445
9 P	-1.9	.122	-1.2	.522	86	.613	-2.0	.214	33	.370	20	.600
10 P	2.4	.466	1.5	.553	1.3	.566	2.6	.445	.08	.870	.25	.634
11 P	2.5	.449	.82	.782	.06	.983	2.0	.389	.05	.938	24	.700
12 P	-4.2	.164	-5.6	.045	-4.7	.070	-3.4	.204	91	.156	82	.194
<mark>3-4 P</mark>	<mark>5.9</mark>	<mark>.030</mark>	<mark>5.2</mark>	<mark>.043</mark>	2.3	.062	2.1	.047	1.1	.062	.50	.087

 Table 5.9: DT Ln schist and minimum temperature in all Asir (June to Sep)

The results indicated that in all Asir between June and September schistosomiasis cases appear to be positively associated with minimum temperature in the 3 - 4 previous months in some, but not all variants of the model.

The results for minimum temperature against schistosomiasis cases in the highlands, the lowlands and Sabt-Alalaya sector do not show any consistency in their significant associations.

5.13.2 Schistosomiasis with rainfall Data

The all Asir model with rainfall between Feb and May is presented in Table 5.10.

	Table	Table 5.10: DT Ln schist and Ln rainfall in all Asir (Feb to May)												
		Liı	near Re	egressi	ion			0	rdinal	Logist	ic			
Model	Rair on	nfall ly	Rainfall with seasonality		Rainfall with auto correlation and seasonality		Rair on	Rainfall only		Rainfall with seasonality		Rainfall with auto correlation and seasonality		
	Cof.	P	Cof.	P	Cof.	P	Cof.	P	Cof.	P	Cof.	P		
0 P	3.2	.047	1.6	.294	1.6	.297	.31	.395	.25	.288	.24	.301		
1 P	1.3	.230	3.3	.033	3.5	.025	.47	.034	.58	.028	.62	.022		
2 P	.24	.861	26	.861	01	.996	11	.627	07	.777	06	.820		
3 P	-1.4	.327	99	.523	95	.541	45	.324	30	.217	31	.205		
4 P	.670	.623	.432	.757	.48	.735	.12	.749	.05	.837	.05	.835		
<mark>5 P</mark>	<mark>2.8</mark>	<mark>.040</mark>	<mark>2.8</mark>	<mark>.043</mark>	<mark>2.8</mark>	<mark>.049</mark>	<mark>.56</mark>	<mark>.024</mark>	<mark>.45</mark>	<mark>.046</mark>	<mark>.47</mark>	<mark>.044</mark>		
6 P	.46	.695	1.9	.192	1.8	.220	.25	.268	.19	.393	.195	.397		
7 P	.23	.846	.83	.610	.86	.600	.44	.214	.41	.117	.41	.114		
8 P	1.8	.133	2.3	.180	3.2	.187	.67	.147	.56	.038	.56	.039		
9 P	-1.3	.340	-1.4	.421	-1.1	.549	08	.745	07	.878	03	.904		
10 P	86	.510	-1.5	.392	-1.5	.401	21	.507	14	.601	15	.596		
11 P	.64	.591	1.0	.517	1.02	.523	.42	.124	.33	.213	.33	.215		
12 P	57	.611	-3.0	.089	-3.2	.075	.74	.084	.56	.050	.57	.046		
<mark>5-6 P</mark>	<mark>2.4</mark>	<mark>.139</mark>	<mark>3.7</mark>	<mark>.037</mark>	<mark>3.7</mark>	<mark>.048</mark>	<mark>3.6</mark>	<mark>.081</mark>	<mark>4.6</mark>	<mark>.025</mark>	<mark>4.5</mark>	<mark>.039</mark>		

 Table 5.10: DT Ln schist and Ln rainfall in all Asir (Feb to May)

 Lipear Pagression

 Ordinal Logistic

The results indicated that in all Asir between Feb and May schistosomiasis cases appear to be positively associated with rainfall in the 5 - 6 previous months for all variants of the model except the rainfall only variants.

The all Asir model for rainfall between June and September is presented in Table 5.11

	Linear Regression Ordinal Logist								Logisti	ic		
Model	Rain on	ıfall ly	Rainfall with seasonality		Rainfall with auto correlation and seasonality		Rainfall only		Rainfall with seasonality		Rainfall with auto correlation and seasonality	
	Cof.	Р	Cof.	Р	Cof.	Р	Cof. P		Cof.	Р	Cof.	Р
0 P	83	.344	-1.7	.123	-1.2	.235	741	.290	31	.193	27	.236
1 P	38	.685	26	.803	.22	.819	315	.554	22	.341	08	.737
2 P	19	.836	-1.4	.199	-1.1	.252	.798	.026	.55	.025	.60	.021
3 P	43	.609	93	.350	76	.400	353	.214	30	.175	19	.404
4 P	-2.4	.001	-2.3	.031	-1.9	.056	.515	.016	76	.105	64	.116
5 P	1.1	.184	.02	.090	1.4	.074	.147	.621	.10	.112	.15	.104
6 P	.83	.430	1.5	.070	.77	.059	.318	.200	.27	.044	.29	.131
<mark>7 P</mark>	<mark>3.3</mark>	<mark>.002</mark>	<mark>2.8</mark>	<mark>.009</mark>	<mark>1.8</mark>	<mark>.046</mark>	<mark>.789</mark>	<mark>.008</mark>	<mark>.63</mark>	<mark>.015</mark>	<mark>.51</mark>	<mark>.022</mark>
8 P	27	.798	.61	.540	.35	.717	.212	.348	.32	.144	.13	.592
9 P	.55	.604	.67	.504	.43	.641	.354	.451	.22	.307	.14	.520
10 P	.75	.393	42	.689	-3.8	.694	117	.524	10	.648	22	.347
11 P	.17	.843	-1.5	.183	-1.1	.306	218	.589	16	.491	14	.552
12 P	.94	.290	1.5	.194	2.1	.950	.373	.099	.29	.255	.44	.105
<mark>5-7 P</mark>	<mark>1.8</mark>	<mark>.042</mark>	<mark>3.5</mark>	<mark>.038</mark>	<mark>2.8</mark>	<mark>.034</mark>	<mark>.754</mark>	<mark>.049</mark>	<mark>.62</mark>	<mark>.037</mark>	<mark>.67</mark>	<mark>.030</mark>

 Table 5.11: DT Ln schist and Ln rainfall in all Asir (June to Sep)

The results indicated that in Asir between June and September schistosomiasis cases appear to be positively associated with rainfall in the 5-7 previous months and in all variants of the model.

The highland group model for rainfall between Feb and July is presented in Table 5.12.

		Li	near Re	egressi	on			C	Logist	stic			
					Rair	nfall					Rair	nfall	
lel	Rair	fall	Rair	nfall	with	auto	Rair	nfall	Rainfall		with auto		
00	on	lv	wi	th	correl	ation		lv	wi	th	correl	ation	
Z	011	1y	seaso	nality	ar	ıd	01	ily.	seaso	nality	an	ıd	
					seaso	nality					seasor	nality	
	Cof.	P	Cof.	P	Cof.	P	Cof. P		Cof.	P	Cof.	P	
0 P	07	.266	15	.104	09	.273	07	.196	22	.164	082	.184	
1 P	07	.284	21	.020	153	.063	061	.191	28	.067	131	.050	
2 P	06	.350	194	.040	070	.439	051	.265	214	.081	094	.477	
3 P	115	.103	.128	.176	050	.560	124	.129	.166	.122	088	.480	
4 P	011	.931	.011	.960	.055	.488	012	.861	.014	.860	.076	.421	
5 P	.102	.249	.086	.365	.061	.467	.147	.278	.099	.311	.089	.397	
<mark>6 P</mark>	<mark>.119</mark>	<mark>.019</mark>	<mark>.249</mark>	<mark>.006</mark>	<mark>.200</mark>	<mark>.014</mark>	<mark>.249</mark>	<mark>.029</mark>	<mark>.527</mark>	<mark>.010</mark>	<mark>.559</mark>	<mark>.007</mark>	
<mark>7 P</mark>	<mark>.150</mark>	<mark>.041</mark>	<mark>.274</mark>	<mark>.004</mark>	<mark>.187</mark>	<mark>.034</mark>	<mark>.321</mark>	<mark>.031</mark>	<mark>.590</mark>	<mark>.006</mark>	<mark>.530</mark>	<mark>.016</mark>	
8 P	038	.603	05	.634	124	.145	062	.524	015	.584	214	.095	
9 P	.024	.746	.093	.360	.172	.055	.034	.702	.083	.311	.222	.085	
10 P	018	.815	091	.409	092	.340	054	.765	077	.329	082	.301	
11 P	033	.689	053	.620	.032	.733	067	.612	036	.602	.044	.513	
12 P	011	.881	06	.605	03	.794	009	.791	051	.545	023	.684	
<mark>5-7 P</mark>	<mark>.212</mark>	<mark>.023</mark>	<mark>.428</mark>	<mark>.001</mark>	<mark>.342</mark>	<mark>.007</mark>	<mark>.467</mark>	<mark>.020</mark>	<mark>1.02</mark>	<mark>.001</mark>	<mark>1.05</mark>	<mark>.001</mark>	
<mark>6-7 P</mark>	<mark>.290</mark>	<mark>.015</mark>	<mark>.454</mark>	<mark>.001</mark>	<mark>.327</mark>	<mark>.003</mark>	<mark>.583</mark>	<mark>.026</mark>	<mark>.943</mark>	<mark>.006</mark>	<mark>.995</mark>	<mark>.006</mark>	

Table 5.12: DT Ln schist and Ln rainfall in the highlands (Feb to July)

The results indicated that in the highlands group between February and July schistosomiasis cases appear to be positively associated with rainfall in the 5 - 7 previous months, and in the 6 - 7 previous months in all variants of the model.

The results for rainfall data against schistosomiasis cases in the lowlands and Sabt-Alalaya did not show any consistency in their significant associations.

5.13.3 Schistosomiasis with relative humidity Data

The results for relative humidity data against schistosomiasis cases in all Asir, the highlands, the lowlands and Sabt-Alalaya did not show any consistency in their significant associations.

5.13.4 Snail numbers with Weather Data

The next analysis examined the relationships between snail number and weather data. This is performed for the whole Asir Region as the snail data is only available for whole the region.

Linear regressions and ordinal logistic regressions of DT snail numbers against weather data in all Asir will be illustrated. Maximum temperature over December to April and May to November periods are presented in Tables 5.13 and 5.14 respectively.

Model	Linear Regression							Ordinal Logistic						
	Max. temperature only		Max. temperature with seasonality		Max. temperature with auto correlation and seasonality		Max. temperature only		Max. temperature with seasonality		Max. temperature with auto correlation and seasonality			
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р		
<mark>0 P</mark>	<mark>6.5</mark>	<mark>.001</mark>	<mark>8.01</mark>	<mark>.047</mark>	<mark>8.2</mark>	<mark>.042</mark>	<mark>.271</mark>	<mark>.008</mark>	<mark>.382</mark>	<mark>.041</mark>	<mark>.341</mark>	<mark>.045</mark>		
1 P	1.18	.516	3.4	.446	2.3	.607	2.34	.487	4.5	.322	2.8	.536		
2 P	121	.178	106	.398	.001	.994	344	.115	350	.244	127	.735		
3 P	093	.120	151	.342	143	.255	172	.175	445	.209	433	.377		
4 P	192	.098	128	.554	.029	.870	103	.379	233	.605	.304	.621		
5 P	141	.058	347	.186	193	.357	139	.389	965	.096	940	.289		
6 P	255	.169	172	.526	.102	.637	217	.575	565	.316	.164	.848		
7 P	137	.002	.117	.264	.063	.402	.091	.224	112	.611	.209	.732		
8 P	.167	.001	.053	.611	033	.656	109	.088	057	.789	.351	.550		
9 P	171	.001	015	.899	069	.414	129	.034	105	.658	418	.577		
10 P	.177	.001	.058	.642	.003	.976	.105	.108	041	.868	.048	.930		
11 P	244	.001	011	.928	016	.863	139	.133	235	.383	351	.632		
12 P	072	.405	.070	.556	.070	.408	133	.349	160	.520	092	.885		

 Table 5.13: DT Ln snails and maximum temperature in all Asir Region
 (Dec to April)

The results indicated that in all of Asir between December and April, the number of snail number appear to be positively associated with the maximum temperature in the current month only in all variants of the model.

	Linear Regression						Ordinal Logistic						
Model	Max. temperature only		Max. temperature with seasonality		Max. temperature with auto correlation and seasonality		Max. temperature only		Max. temperature with seasonality		Max. temperature with auto correlation and seasonality		
	Cof.	P	Cof.	P	Cof.	P	Cof.	P	Cof.	P	Cof.	P	
0 P	1.6	.112	164	.696	-2.79	.588	.218	.186	83	.393	-1.02	.342	
1 P	-1.93	.358	.183	.311	.071	.076	240	.264	2.01	.093	1.40	.091	
2 P	-4.79	.940	024	.918	14	.452	561	.046	-1.18	.033	644	.424	
3 P	2.8	.004	5.48	.123	5.4	.131	1.9	.007	6.01	.254	4.1	.098	
<mark>4 P</mark>	<mark>2.31</mark>	<mark>.004</mark>	<mark>7.11</mark>	<mark>.032</mark>	<mark>6.75</mark>	<mark>.048</mark>	<mark>2.91</mark>	<mark>.011</mark>	<mark>6.66</mark>	<mark>.025</mark>	<mark>7.21</mark>	<mark>.041</mark>	
5 P	1.9	.027	2.9	.358	2.2	.496	2.08	.030	3.4	.265	2.91	.254	
6 P	3.25	.264	1.48	.113	1.23	.381	2.75	.126	1.38	.047	1.59	.667	
7 P	2.50	.635	157	.215	-1.33	.759	2.81	.718	.124	.650	299	.488	
8 P	-2.30	.337	-2.9	.494	-2.27	.814	1.71	.699	911	.670	248	.567	
9 P	2.59	.649	2.52	.697	2.80	.464	.939	.885	.920	.941	928	.641	
10 P	1.83	.549	1.73	.603	1.38	.736	.430	.181	.408	.219	.572	.204	
11 P	-2.21	.937	-1.57	.689	-2.25	.830	041	.843	.236	.456	.464	.352	
12 P	-1.06	.277	-1.07	.965	-1.77	.524	152	.450	085	.780	345	391	
<mark>3-4 P</mark>	<mark>2.6</mark>	<mark>.003</mark>	<mark>10.7</mark>	<mark>.015</mark>	<mark>10.2</mark>	.022	<mark>3.2</mark>	<mark>.001</mark>	<mark>8.75</mark>	<mark>.011</mark>	<mark>9.24</mark>	<mark>.034</mark>	
<mark>2-4 P</mark>	<mark>3.3</mark>	<mark>.001</mark>	<mark>10.0</mark>	<mark>.047</mark>	<mark>10.92</mark>	.044	<mark>3.1</mark>	<mark>.007</mark>	<mark>9.23</mark>	<mark>.033</mark>	<mark>9.47</mark>	<mark>.039</mark>	

Table 5.14: DT Ln snails and maximum temperature in all Asir Region (May to Nov)

The results indicated that in all Asir between May and November, the number of snails appear to be positively associated with maximum temperature in the 3 - 4 previous months and in 2 - 4 previous months in all variants of the models.
The result of rainfall for all Asir model between December and April period is presented in Table 5.15

		Li	near Re	egressi	on		Ordinal Logistic							
Model	Rainfall only		Rainfall with seasonality		Rainfall with auto correlation and seasonality		Rainfall only		Rainfall with seasonality		Rainfall with auto correlation and seasonality			
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р		
<mark>0 P</mark>	<mark>.72</mark>	<mark>.815</mark>	<mark>-7.8</mark>	<mark>.048</mark>	<mark>-8.2</mark>	<mark>.036</mark>	<mark>.67</mark>	<mark>.511</mark>	<mark>-3.46</mark>	<mark>.033</mark>	<mark>-2.49</mark>	<mark>.028</mark>		
1 P	-3.2	.392	-4.3	.252	-3.5	.350	-1.18	.219	-3.11	.112	-3.7	.169		
2P	1.1	.781	3.4	.379	3.7	.321	.954	.621	2.0	.412	1.91	.249		
3 P	.120	.658	.041	.731	058	.533	.119	.119	151	.585	.177	.697		
4 P	122	.758	116	.697	066	.790	16	.668	-1.06	.099	519	.662		
5 P	615	.869	356	.462	375	.348	819	.913	-1.02	.298	469	.267		
6 P	.662	.085	234	.364	.123	.600	.718	.795	-1.27	.424	705	.251		
7 P	.556	.226	264	.305	37	.874	616	.857	-1.80	.305	-1.28	.261		
8 P	238	.626	186	.404	.156	.445	152	.335	-1.01	.047	562	.304		
9 P	198	.008	247	.219	086	.625	212	.179	-1.11	.018	915	.072		
10 P	.348	.011	.200	.721	.390	.311	.227	.946	-1.47	.210	782	.229		
11 P	012	.047	.94	.245	.324	.311	037	.054	.842	.362	.254	.318		
12 P	2.0	.540	-6.0	.161	-5.3	.207	1.3	.454	-4.12	.174	-4.84	.222		

 Table 5.15: DT Ln snails and Ln rainfall in all Asir Region (Dec to April)

The results indicated that in all Asir over December to April, the number of snails appears to be negatively associated with rainfall in the current month only in most variants of the model.

5.13.5 Schistosomiasis Cases with Snail Numbers

The final stage of the analysis investigated whether there were associations between schistosomiasis cases and snail numbers. Linear regressions and ordinal logistic regressions of DT Ln schistosomiasis cases against DT Ln the number of snails in all Asir are presented in Table 5.16.

		Li	near R	egressi	on			0	rdinal	Logist	ic	
					Sna	ails					Sna	uils
lel	Sna	ils	Sn	ails	with	with auto Snails		Snails		with auto		
100	only		W	ith	correl	ation	on	lv	wi	th	correlation	
2		5	seaso	nality	an	id	_	5	seasor	nality	an	d
	C.f. D		0.6	D	seasonality					D	season	
	Cof.	P	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	P	Cof.	P
0P	<mark>.082</mark>	.001	<mark>.048</mark>	.031	<mark>.06</mark>	<mark>.036</mark>	<mark>.017</mark>	<mark>.001</mark>	<mark>.012</mark>	<mark>.026</mark>	<mark>.014</mark>	<mark>.011</mark>
1 P	<mark>.101</mark>	<mark>.001</mark>	<mark>.066</mark>	<mark>.016</mark>	<mark>.06</mark>	<mark>.031</mark>	<mark>.020</mark>	<mark>.001</mark>	<mark>.016</mark>	<mark>.003</mark>	<mark>.015</mark>	<mark>.007</mark>
2 P	.047	.070	.017	.541	.010	.817	.037	.084	.042	.431	.012	.757
3 P	.028	.286	.027	.336	.024	.384	.029	.271	.034	.387	.024	.244
4 P	.021	.422	.042	.132	.038	.173	.031	.472	.032	.149	.079	.142
5 P	019	.475	.022	.448	.015	.603	029	.485	.041	.499	.027	.543
6 P	06	.026	018	.518	02	.434	05	.036	033	.555	10	.384
7 P	.073	.006	04	.140	040	.167	.078	.004	014	.187	042	.097
8 P	05	.064	023	.431	02	.580	04	.055	033	.466	026	.521
9 P	.038	.155	.057	.047	.06	.032	.033	.167	.086	.067	.046	.089
10 P	.036	.192	.026	.375	.02	.561	.031	.184	.019	.178	.022	.561
11 P	.032	.248	.010	.936	01	.948	.039	.227	.011	.857	012	.838
12 P	<mark>.110</mark>	<mark>.001</mark>	<mark>.082</mark>	<mark>.005</mark>	<mark>.08</mark>	<mark>.005</mark>	<mark>.017</mark>	<mark>.001</mark>	<mark>.013</mark>	<mark>.019</mark>	<mark>.014</mark>	<mark>.015</mark>
<mark>0-1 P</mark>	<mark>.140</mark>	<mark>.001</mark>	<mark>.094</mark>	<mark>.007</mark>	<mark>.094</mark>	<mark>.007</mark>	<mark>.029</mark>	<mark>.001</mark>	<mark>.024</mark>	<mark>.001</mark>	<mark>.023</mark>	<mark>.001</mark>

Table 5.16: DT Ln schist and DT Ln snails in all Asir Region

The results over the whole year indicated that in all Asir schistosomiasis cases appear to be positively associated with the number of snails collected, in average of the current month and one previous month and in all variants of the models.

5.13.6 Multiple Regressions

In this section we will present the results from the normal and multiple regressions of DT Ln schistosomiasis cases against the weather variables. For temperature, we will present maximum temperature only in the multiple regressions as it has a similar seasonality to minimum and mean temperature and appeared to have the strongest associations with schistosomiasis. No multiple regressions were produced for the snail data as in no period was more than one set of weather variables significant.

A. Multiple Regressions of Schistosomiasis Cases in All Asir

The normal regressions for schistosomiasis in all Asir against maximum temperature and rainfall, in the averages of 3 and 4 previous months, during June to September are illustrated in Tables 5.17. Table 5.18 presents the multiple regressions for these variables.

Table 5.17: Regressions of the coefficient of DT Ln schistosomiasis cases of whole Asir for averages of 3 and 4 previous months of maximumtemperature and rainfall during June to September

				Line	ar regre	ssion				Ordinal logistic								
Vormal gression	Variables only			se	with easonali	ty	V coi Se	Vith aut rrelation easonali	o 1 & ty	V	/ariable only	es	Se	With easonali	ty	V cor Se	Vith aut relatior easonali	o ı & ty
Re	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.
Max.Tem. Only	3.62	.001	.085	4.0	.021	.216	3.51	.029	.337	1.0	.008	-75.0	1.23	.006	-69.2	1.04	.008	-69.0
Rainfall Only	-2.62	.012	.109	-2.72	.045	.193	-2.24	.074	.312	44	.043	-75.0	481	.116	-71.6	480	.114	-71.5

 Table 5.18: Regressions of the coefficient of DT Ln schistosomiasis cases of whole Asir with multiple weather averages of 3 and 4 previous months of maximum temperature and rainfall during June to September

po				Line	ar regre	ssion							Ord	inal log	istic			
riables In ple Meth	١	/ariable only	s	se	with easonali	ty	COI Se	Vith aut rrelatior easonali	o 1 & ty	٧	/ariable only	S	Se	With easonali	ty	V coi Se	Vith auto relation & easonality	
Va Multi	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.
Multiple Max.Tem	.808	.039	155	2.95	.047	212	2.75	.043	220	.049	.019	-	1.06	.036	-	1.10	.033	-
Multiple Rainfall	-2.2	.033	.155	-1.41	.377	.215	-1.03	.485	.329	470	.038	72.83	09	.811	69.12	0.82	.817	69.01

The results of multiple regressions for schistosomiasis in all Asir during June to September demonstrate that in no models were the two weather variables significant. We conclude therefore that the best model for this period is that of DT Ln schistosomiasis against maximum temperature. Maximum temperature was positively significant in all variants of the normal and multiple regressions.

B. Multiple Regressions of Schistosomiasis Cases in the Highlands

The normal regressions for schistosomiasis cases in the highlands against maximum temperature and rainfall, in the averages of 6 and 7 previous months, during February to July are illustrated in Tables 5.19. Table 5.20 presents the multiple regressions for these variables.

Table 5.19: Regressions of the coefficient of DT Ln schistosomiasis cases of highlands group for averages of 6 and 7 previous months ofmaximum temperature and rainfall during February to July

				Line	ar regre	ssion							Ord	inal log	istic			
Vormal gression	Variables only			se	with easonali	ty	V coi Se	Vith aut rrelation easonali	o 1 & ty	V	∕ariable only	S	Se	With easonali	ty	V cor Se	Vith aut relatior easonali	o 1 & ity
Re	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.
Max.Tem. Only	025	.406	.01	34	.087	.001	179	.323	.202	.010	.993	- 102.8	580	.185	- 100.2	231	.593	-96.0
Rainfall Only	.290	.015	.064	.454	.001	.165	.327	.003	.307	.583	.026	-99.9	.943	.006	-94.3	.995	.006	-89.7

Table 5.20: Regressions of the coefficient of DT Ln schistosomiasis cases of the highlands group with multiple weather averages of 6 and 7 previous months of maximum temperature and rainfall during February to July

po				Line	ar regre	ssion							Ord	inal log	istic			
riables In ple Meth	Variables only			se	with easonali	ty	V coi Se	Vith aut rrelation easonali	o 1 & ty	٧	ariable/ only	S	Se	With easonali	ty	V coi Se	Vith aut relatior easonali	o 1 & ty
Vaı Multi	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Adj. R ²	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.	Cof.	Р	Log Likl.
Multiple Max.Tem	060	.053	104	084	.668	152	.003	.987	205	.070	.293	_	.024	.959	-	.472	.327	-
Multiple Rainfall	.287	.004	.104	.408	.001	.155	.327	.005	.295	.559	.011	99.40	1.03	.002	94.30	1.18	.001	89.24

The results of multiple regressions for schistosomiasis in the highlands during February to July demonstrate that in no models were the two weather variables significant. We conclude therefore that the best model for this period is that of DT Ln schistosomiasis against rainfall. Rainfall was positively significant in all variants of the normal and multiple regressions.

5.14 Discussion and Conclusion

Several studies of schistosomiasis disease in Saudi Arabia have been carried out, but most of them have not examined the importance of climate and weather upon the existence of the disease. This is an important research gap as schistosomiasis used to be one of the most serious public health problems in this country, especially in the southern regions (Statistical year book MOH, 2002). The highest rate of this disease in Saudi Arabia during 2005 was in Asir Region (16.3 cases/100,000) followed by Albahah and Jazan Regions, (13.0, 11.5 cases/100,000) respectively. This justifies the importance of examining schistosomiasis within the Asir region. Both intestinal and urinary schistosomiasis have been reported in Asir (Statistical year book MOH, 2006).

During this study some difficulties were encountered. The main one was the limited weather data for some selected endemic areas in Asir, and also information missing from within this data. This meant that it was not possible to produce models for each schistosomiasis endemic area in Asir. For example, we have combined several sectors for endemic schistosomiasis data in the highlands including Abha, Khamise Sarat-Abidah Ahad-Rifadah, Tathlith, Almadha, Dhahran-Aljanoub and Tanoma, into one group, and a similar combination of sectors has been carried out for the lowlands. Although all these areas are in the highlands and have the same seasonality of schistosomiasis, it would have been better to examine each separately but the weather data was not available for each sector. It was therefore necessary to group sectors together. However, there were even greater limitations with the snail data which were only available for the whole of Asir. Obtaining these data for sectors within Asir would be helpful in understanding variations in snail numbers throughout the region. Also these data appeared to correlate well with schistosomiasis cases in the same month. This indicates that they do not give an accurate picture of variations in

snail numbers in the region. In the future, it would be helpful to examine whether more useful information on snail numbers in the region could be extracted.

The average number of schistosomiasis cases in the whole of Asir in the last year (2009) of the study is less than one third of the average of cases in the first year (1998), This is a lower reduction in reported cases than for leishmaniasis and malaria and indicates that there has been less progress with this illness than with malaria and leishmaniasis.

In the results presented in this chapter, temperature was associated with schistosomiasis incidence in Asir, and there is a positive association between the maximum temperature and the number of cases for the whole of Asir during June to September. The lag for this relationship is from 3 to 4 months which fits with lags reported in the literature. Mean and minimum temperature showed similar trends. This relationship did not appear in the rest of year. However, a link with temperature was not apparent in any of the groups making up Asir, specifically the highlands sector, the lowlands or in Sabt-Alalaya sector. If such a relationship exists, it is difficult to understand why it would not be apparent in any of the 3 sectors making up Asir. However, further support for this relationship comes from the number of snails collected between May and November which also appear to be associated with maximum temperature averages of the previous 3 and 4 months. In the literature there is support for an association with temperature. Pimentel-Souza., et al (1988) state that in Brazil, an increase in temperature during winter and autumn leads to more mating activity amongst snails which may lead to more schistosomiasis. Michelson, 1960 mentioned that temperature may influence the ability of miracidia to infect snails and also the development of the schistosoma larvae within snails. In Biomphalaria glabrata Coelho et al., (2006) showed that lower temperatures led to the lowest infection level. These could all be explanations for the possible positive associations between temperature and schistosomiasis and snails observed in Asir. It is important to note that positive associations are not observed in all locations. In Senegal, Sturrock et al., (2001) noted that snail numbers drop sharply in the heat of the summer. It is likely that temperatures in that country rise to a level where snail survival is affected. In Asir, temperatures in the highlands (where most cases occur) are low enough that elevated temperatures are unlikely to have a negative effect upon snail numbers.

The strongest relationship emerging from this chapter is that rainfall appears to be positively associated with schistosomiasis cases for the highlands only. Elevated rainfall 6 and 7 months previously led to elevated schistosomiasis cases between February and July. In the literature rainfall / water is constantly cited as being important for schistosomiasis. The impact of rainfall upon schistosomiasis in this study was only in the highlands. This is unsurprising as most cases (more than 80%) are reported from the highlands. The positive association between rainfall and schistosomiasis occurs between February and July. It refers to rainfall 6 and 7 months previously which is the time of year when rainfall at its lowest in the highlands (e.g. September – December). During these dry periods elevated rainfall may provide opportunities for snails to breed. This then leads to schistosomiasis cases 6-7 months later.

There is widespread support for an association between schistosomiasis and rainfall. In Saudi Arabia Al-Madani (1991) mentioned that rainfall and streams are the major sources of water in Asir region and that vector snail habitats are found in streams in the highlands and associated with precipitation. Kloos et al., (2004) state that larger snail numbers were found in standing waters from October to March during the rainy and hot season in a rural area in Brazil. Agi (1995) in the Jos metropolis area of Nigeria noted that the rainy season in this part of the country occurs during summer when more snails were found. A lack of rainfall in the Ouarzazate part of Morocco, led to the drying out of the breeding sites of Bulinus truncatus, hence stopping of transmission cycle of the schistosomiasis (Laamrani et al., 2000).

One novel feature of this chapter was the snail data which might help to explain the relationships between schistosomiasis and weather. However, as already discussed, these data had several limitations as they were available only for the whole of Asir and not for each endemic area. The results for snails against weather data indicated that in the whole of the region over winter and spring seasons, the number of snails appeared to be negatively associated with rainfall in the current month only. This is different to the schistosomiasis data which indicated a positive association with rainfall 6-7 months earlier. One hypothesis for this relationship could be that rainfall washes snails from the bodies of water being investigated leading to lower numbers being collected. An alternative explanation could be that rainfall has a negative effect on the ability of collection teams to search for snails.

The third relationship examined was between schistosomiasis cases and the number of snails. The results over the whole year indicated that in the whole of this region schistosomiasis cases appear to be associated with the average number of snails, in average of the current month and one previous month. The reason for this is not clear as the literature review has indicated that the development of the parasite of schistosoma in the snails and the incubation of the disease take more than two months. Instead the relationship may indicate that the people who are responsible for the collection of snails may try to collect more snails from the environmental sources when they realise that there is increase of the schistosomiasis cases.

In summary, this study showed that weather variation factors play a role in affecting schistosomiasis cases. The results indicated that temperature may be an important factor associated with schistosomiasis in the whole of Asir during the summer and beginning of autumn but it was unclear why this impact was not seen in any of the sectors making up Asir. The results showed that if the maximum temperature in the previous 3 to 4 months was elevated, there were more cases of schistosomiasis. There was stronger support for the role of rainfall, and transmission in the highlands was elevated if rainfall was elevated in the previous 6 to 7 months. This occurred during the spring and summer seasons only. The results also indicated that weather was not an important factor associated with snails in Asir. This may be partly due to limitations of the snail data especially the close association with the number of reported cases. Although this study shows a decrease of schistosomiasis cases during the study period, it is important to recognize that this disease is a still a significant health problem particularly in some endemic sectors in this Region.

Chapter 6

Conclusion

6.1 Introduction

Habib, (2011) states that in the Arab countries, malaria, and leishmaniasis are the major vector-borne diseases of concern. The prevalence of these diseases has been linked to climate-related environmental changes. Increased temperatures, variable humidity and rainfall trends may affect the vector population density, the patterns of transmission, and infection rates (WHO et al., 2003; Habib, 2011). The consequences of these changes may cause vector-borne illnesses to emerge in some non endemic areas or to intensify in endemic areas particularly in Arab countries (Habib, 2011).Costello et al., (2009) state that no effective vaccines exist for many climatesensitive infectious diseases such as malaria, dengue fever, schistosomiasis, and leishmaniasis. Reliance upon finding a vaccine solution is not appropriate

The three illnesses, malaria, leishmaniasis and schistosomiasis, in this study have been chosen for some reasons. These illnesses still represent major public health problems in Saudi Arabia, and particularly in Asir region. Costello et al., (2009) state that these three diseases are sensitive to climate (climate-sensitive infectious diseases), so it was useful to study these diseases in Asir.

6.2 Development of the Study

As we have discussed, the methodology of this study was developed predominantly in the first main chapter, malaria. The other two chapters, CL and schistosomiasis, built upon the methods created in this chapter meaning that difficulties or obstacles encountered in the Malaria chapter did not need to be repeated. A good example of this was with the first stage of the malaria study in which much time was spent investigating how to account for the fact that the relationships with weather may vary over the year. Initially we produced scatter plots of malaria against the independent variables for every month. The purpose of this analysis was to determine which months should be grouped together (i.e. group consecutive months with positive or negative relationships with weather together). Unfortunately, this method proved difficult to implement because it was often unclear which months should be grouped. The situation was complicated further by the fact that some of the relationships between weather and malaria may be lagged. Therefore a new method was developed and grouped months together based upon those with similar trends in the malaria time series over the year (e.g. months of high incidence during the winter). This method was taken forward to the CL and schistosomiasis chapter and scatter plots were not used again. In the malaria chapter much effort also went into deciding that most appropriate statistical model to use. In this chapter we decided upon conducting both a multiple regression and an ordinal logistic regression. These developments were taken forward to the CL and schistosomiasis chapter.

In the first two data chapters weather was compared to disease incidence. A major development of this idea occurred in the schistosomiasis chapter where we were able to obtain data on the abundance of the environmental host of this disease (snails). It was hoped that this development might explain the relationships between schistosomiasis and weather observed. Although an exciting addition to the study it was concluded that these data had major limitations as they seemed to be a poor indicator of the numbers of snails in the environment and were more affected by sampling effort. In addition, this data was only available for Asir a whole and not for each endemic area which limited its use.

6.3 Summaries and discussions

The results of this study were divided into three main data in addition to two secondary chapters (Introduction and Study area). A brief summary of each will now be given.

6.3.1 Infectious Diseases (Chapter 1)

The study has presented an overview of infectious diseases and highlighted their global importance to health. We also demonstrated why many of these are sensitive to climate variability. One important reason to examine climate variability is because of the likely impact of climate change upon incidence. An overview of global climate change and its adverse impact upon health were presented in this chapter. The chapter ended by justifying the three illnesses (malaria, CL and schistosomiasis) which are the infectious diseases to be studied in this thesis.

6.3.2 Study Area (Chapter 2)

The next chapter presented Saudi Arabia and the Asir Region. The choice of this region was justified through its location in the Afro-tropical Zone which extends from tropical countries in Africa to the Arab Peninsula (Khalifa 2009). It was shown how this region had a number of infectious diseases and how its varied topography and weather made it an ideal study choice. Based on the literature review of the three illnesses, and all the above information, it was shown clearly that there was an urgent need to study and research the influence of climate variability upon the presence of these illnesses in Asir Region.

6.3.3 Malaria (Chapter 3)

The first main data chapter considered malaria, as this illness is the most important vector-borne disease globally (WHO/SDE/PHE, 1999) and in the study area (Asir Region) (Dr A. Abdoon 2007 pers. Comm.). Malaria is endemic in lowlands of Asir Region (Haddad, 1990). The study showed that weather variations play an important role in affecting the number of malaria cases in Asir Region. The results indicated that rainfall was the most important factor associated with malaria during the summer months in Tehama of Asir (TA) and Tehama of Qahtan (TQ) (which include valleys and foothills). Greater rainfall in the current and previous month led to more malaria cases. Although rainfall was also significant over the whole year, because rainfall was not significant in models for other times of the year we concluded that the whole year result was driven by the summer result. Because summer is the driest period any rainfall during this period may provide opportunities for mosquito breeding. At other times of the year there may always be enough breeding sites for mosquitoes and so additional rainfall is unimportant.

6.3.4 CL Leishmaniasis (Chapter 4)

The second climate-sensitive infectious disease examined in this study was Leishmaniasis. As there were only limited cases of visceral leishmaniasis in Asir over the study period this study focused upon cutaneous leishmaniasis (CL). Leishmaniasis is considered to be the vector-borne disease with the second highest number of affected people in the world after malaria (Chaves et al., 2006). This study showed that weather factors play an important role in affecting the incidence of CL cases. Temperature was the most important factor associated with CL in all Asir and in the highlands area. Between summer and the beginning of autumn if the maximum temperature in the previous 2 to 4 months was elevated there were more cases of leishmaniasis. The highlands in Asir might provide the optimum temperature for the vector during summer, and elevated temperatures may increase sandfly activity. However, there was no evidence for a relationship between temperature and CL cases in the lowlands areas. Because these areas are much hotter, a negative association with maximum temperature in the summer was anticipated. The lack of a result suggests that sandflies find mechanisms to avoid these high temperatures. Although temperature does not appear to be important rainfall plays an important role in CL transmission in the lowlands. A greater number of cases occurs if rainfall is elevated in the previous 1 to 3 months. This occurs in all months except during the winter season. Rainfall may provide more suitable conditions for sandflies.

6.3.5 Schistosomiasis (Chapter 5)

The influence of climate variability upon this disease is a significant research gap as schistosomiasis used to be one of the most important public health problems in Saudi Arabia, particularly in the southern regions (Statistical year book MOH, 2002). This study showed that weather variations play a role in affecting schistosomiasis cases. The results indicated that temperature may be an important factor associated (positive association) with schistosomiasis in all Asir during the summer and beginning of autumn. It was unclear why this impact was not seen in any of the individual sectors within Asir. The results showed that if the maximum temperature in the previous 3 to 4 months was elevated there were more recorded cases of schistosomiasis. The strongest relationship emerging from this study is that rainfall appears to be positively associated with schistosomiasis cases in the previous 6 to 7 months for the highlands only. This occurred during the spring and summer seasons. Rainfall may provide suitable breeding sites for snails. One novel feature of this study was the data on snail abundance which it was hoped might help to explain the relationships between schistosomiasis and weather. However, the associations between these data and the schistosomiasis incidence data suggest that the snail counts were not a true reflection of snail abundance within Asir. Instead they suggest that sampling effort is increased when schistosomiasis cases are more abundant.

6.4 Conclusion of the results of the three illnesses

This study shows associations between weather and malaria, schistosomiasis and leishmaniasis in Asir. Overall the total numbers of yearly cases for all three diseases have declined sharply since the 1990's. The Vector Control Administration, Health Affairs Directorate in Asir Region has made important progress regarding the control of the vector-borne diseases in Asir, in addition to the general strategy adopted by the Ministry of Health. This progress provides an explanation for the reduction in the number of cases in Asir. This is as a result of the improved facilities and an increased number of professionals, technicians and workers concerned with the problem. Different methods for the control of malaria and control of leishmaniasis have been introduced by the Vector Control Administration and some Primary Health Care centres. Two methods both involved spraying pesticides to combat malaria: (Seasonal spraying) in the houses which has a residual effect and space spraying (Fogging) which can make a significant impact upon the vector of leishmaniasis (sandfly) as well as the mosquito vectors. The greatest effort has been concentrated on the control of malaria as this has been the most prevalent disease. However, there was a massive spray against rift valley fever disease for more than six months which started in Asir and Jazan in 2000. This action had an important impact upon the vectors of malaria and leishmaniasis.

However, it is important to recognize that these diseases still represent significant health problems in Asir, especially schistosomiasis and Leishmaniasis with a concern of malaria incidence in some endemic areas if there is any control shortage in these areas (Dr A. Abdoon 2007 pers. Comm.).

To summarize, Table 6.1 shows the most important results for the relationships of between malaria, CL and schistosomiasis with the climate variables (Temperature, rainfall and relative humidity) in Asir Region during the study period.

	¥	Temp	erature	Rai	nfall	Humi	dity
Disease	Lag	Location	Time	Location	time	Location	time
Malaria	0-1 P	-	-	TA&TQ	Summer	All Asir	Whole year
	2-4 P	All Asir	Summer	-	-	-	-
CI	2-4 P	Highland	of Autumn	-	-	-	-
CL	1-3 P	-	-	Lowland	Spring + Summer + Autumn	-	-
Schistosom iasis	3-4 P	All Asir	Summer + beginning of Autumn	-	-	-	_
	6-7 P	-	-	Highland	Spring + Summer	-	-

 Table 6.1: Summary of the significant relationships between the climate variables and selected infectious diseases in Asir (Malaria, CL and Schistosomiasis)

P: Previous month. TA: Tehama of Asir. TQ: Tehama of Qahtan

The previous table illustrates the following points:

- Weather variations play an important role upon the three illnesses. However, the influence varies by region and different weather variable have different lags. The shortest lag times were found for malaria cases. The longest lag times were for schistosomiasis. These are biologically plausible.
- Rainfall is positively associated with all three illnesses indicating the importance of rainfall on disease transmission in Asir. In general rainfall during drier periods appears to be important.
- 3. Temperature has an influence upon only CL and Schistosomiasis. Summer and beginning of autumn temperatures appear to be important for both these diseases.
- 4. Humidity does not seem to be an important factor for many illnesses. Although it is significant for malaria in all Asir, it is not significant in any of the sectors. In this study the ability to detect a humidity effect was limited by the lack of humidity data in many parts of Asir.

5. Most associations between weather and disease appear for illnesses reported in the summer season.

6.5 The importance of climate change in this Study

The key measures of climate change in any particular region may depend on changes in temperature and rainfall (Islam, 2009). Cross and Hyams, 1996 carried out the influences on human health. This study was not designed to examine how the distribution of diseases in Saudi Arabia and Asir might change. Instead it provides evidence on how disease rates may change in areas where they are already present.

During a presentation in Amman, Jordan 2006 "Regional Workshop on Adaptation Strategies to Protect Health under Climate Change Variability and Change in Water Stressed Countries in EMR" it was mentioned that the annual temperature change in the Middle East between 1955 and 1994 was a 0.5°C increase. It is forecast that the increase by 2030 and 2050 will be 1 - 2°C respectively (MOH, workshop, 2006). Almazroui, 2011 states that forecasting the future climate of Saudi Arabia, projections using previously calibrated data reveals that rainfall will have a tendency to increase by 1.04% per decade. The tendency of maximum temperature will be to increase 0.66°C - 0.64°C per decade, compared to the base period. The projected rainfall will tend to increase particularly over the coastal areas along the central parts of the Red Sea as well as over the south-south-western areas of Saudi Arabia (Almazroui, 2011) i.e. Asir.

In this study there was a positive association between temperature 2-4 months previously and CL in summer and beginning of autumn. Should climate change lead to higher temperatures as is indicated this might increase the incidence of CL in highland areas in summer and the beginning of autumn. Rainfall 0-1 month previously was associated with elevated cases of malaria in TA and TQ. Again should climate change increase rainfall as indicated then malaria cases could increase in TA and TQ. Temperature 3-4 months previously was associated with increased cases of schistosomiasis. Should climate change lead to increase the incidence of this disease also in summer and beginning of autumn. Again with an increase in rainfall due to climate change, as rainfall 6-7 month previously

was associated with the increased cases of schistosomiasis in the highlands of Asir Region.

In addition to these changes in incidence is also possible that areas which do not currently have malaria, schistosomiasis and CL could become endemic or that transmission could be extended throughout most or all months of year.

6.6 Study limitations and future work

During this study several difficulties were encountered. The main one was the limited weather data for some selected areas in Asir, and missing information from within this data. This meant that it was not possible to produce models for each endemic area in Asir. The situation in some endemic areas was even more difficult as no weather stations were available within some of these areas. Instead we had to use the weather data from nearby stations which were sometimes outside the sector boundaries. Although these stations were chosen because of their close similarity to the weather in the endemic areas, this can cause problems in the analysis if the variations in temperature, rainfall and humidity in those areas are different to that in the chosen weather stations. The problem was greatest with humidity where data was only available for one weather station. This meant that in all three data chapters the influence of humidity in many sectors could not be assessed. This highlights the importance of weather data collected to epidemiological studies. Should improved weather data exist then it would be possible to explore the relationships in this study further.

There were also limitations with the health data as only monthly reports were available. For diseases such as malaria where time lags between weather and illness are short, an analysis of weekly malaria cases would be preferable and would be an important addition to this study. Also the degree of under-reporting in these data is unknown.

Although this study was based in Asir the study could easily be extended to other areas where malaria, CL or schistosomiasis are endemic. In Saudi Arabia such regions as Jizan or Albahah would be ideal. It might also be possible to combine health data (e.g. Asir (Tehama) with some areas in Jizan) as these regions are similar topographically and environmentally. This might help overcome problems of low numbers.

It is also possible that similar methods could be used to investigate the influence of the climate variables upon other diseases such as food-borne diseases or air-borne diseases. Chapter 1 indicated that amoebic dysentery and brucellosis have a relatively high incidence in Asir so could be investigated in future research.

Understanding how weather affects vector populations is important to understanding weather health relationships. In this study the relationship between schistosomiasis cases and the number of snails was examined as this data might explain the relationships between schistosomiasis and weather. However, as already discussed this data had several limitations as it was available only for Asir as a whole and not for each endemic area.

Although our study may have some limitations, we strongly believe that our findings are relevant from a public health perspective to better understand the epidemiology of these illnesses. This study adds to the growing evidence linking the three diseases to climate variability. It suggests that variations in the incidence of these diseases in Asir are associated to changes in weather conditions. Although during the study period there were activities to control these diseases in this region, the number of cases for each disease correlated statistically, with climate variables. The result of this study could contribute to an increased awareness of these diseases and help to diagnosis the problem for the decision-makers in related health agencies in this region particularly and in Saudi Arabia generally.

References

- Abdoon, A. and Alshahrani, A. (2003). Prevalence and distribution of anopheline mosquitoes in malaria endemic areas of Asir Region, Saudi Arabia. *Eastern Mediterranean Health Journal* 9 (3), 240-247.
- Agi, P.I. (1995). Distribution of freshwater snail vectors of schistosome parasites in Jos metropolis (Nigeria). *Acta Hydrobiologica* **37**, 59-67.
- AI-Madani, A. A. (1991). Problems in the control of schistosomiasis in Asir Province, Saudi Arabia. *Journal of Community Health* **16** (**3**), 143-149.
- Al-Abdullatif, Z., Abdelaziz, B., Al-Hamdan N. and Al-Sagheer S. (1996). Malaria outbreak in Gellwa, Al-Baha, Saudi Arabia 1996. Field Epidemiology Training Program, *Saudi Epidemiology Bulletin* 3 (4), 27-28.
- Alam, M., Shafiqur, R., Sabbir, H. and Shajedul, H. (2008). *Relationship between climate change and incidence of malaria in Chittagong Hill Tracts in Bangladesh*. Presentation made at the 2nd International Workshop on Community Based Adaptation (CBA) to Climate Change, Dhaka.
- Al-Amru, S. (2002). The importance of sandfly (Psychodidae: Diptera) in KSA. Entomology Department, Agriculture College, King Saud University. (Report, Arabic Version).
- Al-Hammed, N. (2003). *Cutaneous Leishmaniasis in Saudi Arabia*. A dissertation for the degree of Master of Science in Clinic Dermatology, University of London.
- Al-Harthy, S. and Jamjoom, M. (2007). Study of the prevalence of malaria parasites in Jizan region, Saudi Arabia using PCR technique. Final Report, Department of Parasitology, Faculty of Medicine, Umm Al-Qura, University, Makkah, KSA.
- Al-Jaser, M. H. (2006). Studies on the epidemiology of malaria and visceral leishmaniasis in Jizan Area, Saudi Arabia, Riyadh. *Journal of King Saud University* **19** (**1**), 9-19.
- Alkhalife, I. S. (2003). Imported malaria infections diagnosed at the malaria referral laboratory in Riyadh, Saudi Arabia. *Saudi Medical Journal* **24** (**10**), 1068-1072.

- Alkhawajah, A. (1998). Recent trends in the treatment of cutaneous leishmaniasis (Review Article). *Annals of Saudi Medicine* **18** (**5**), 412-416.
- Almazroui, M. (2011). Present and Future Climate of Saudi Arabia as Simulated by the PRECIS Regional Climate Model Center of Excellence for Climate Change Research. Department of Meteorology, King Abdulaziz University, Jeddah, Saudi Arabia.
- Al-Saghayer S. and Alattas M. (2006). *Roll Back Malaria in Saudi Arabia*, (Report). <u>http://www.emro.who.int/rbm/presentations/5.%20KSA%20-</u> <u>%20Country%20Presentation.ppt#364,7,Slide7</u> Last accessed: April, 2011.
- Al-Saghayer S. and Tamim, O. (1996). Guidelines for treatment of malaria. Department of Malaria, Preventive Medicine, Ministry of Health. Field Epidemiology Training Program. *Saudi Epidemiology Bulletin* 3 (4), 30.
- Al-Tawfiq, J. A. and Abu-Khams, A. (2004). Cutaneous leishmaniasis: a 46-year study of the epidemiology and clinical features in Saudi Arabia (1956-2002). *International Journal of Infectious Diseases* 8, 244-250.
- Al-Zahrani, M. A., Peters, W., Evans, D. A., Smith, V. and Ching Chin, I. (1989). Leishmania infecting man and wild animals in Saudi Arabia. 6. Cutaneous leishmaniasis of man in the south west. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 83, 621-628.
- Al-zahrani, M. H. (2007). *Impact of irrigation systems on malaria and RVF transmission in Jazan, Saudi Arabia.* PhD Thesis, Liverpool University, Liverpool.
- Amerasinghe PH., Amerasinghe FP., Konradsen F., Fonseka KT. and Wirtz RA. (1999). Malaria vectors in a traditional dry zone village in Sri Lanka. *American Journal of Tropical Medicine and Hygiene* 60, 421-429.
- Amin, M. A. and Al-Saghayer, S. M. (1995). *Manual on the control of schistosomiasis for primary health care workers*. Book. Ministry of Health, Saudi Arabia, No. 2484/15, Radmac list 2-03-603-9960.
- Appleton, C. C. (1977). The influence of temperature on the life-cycle and distribution of Biomphalaria pfeifferi, Krauss, 1948 in south-eastern Africa. *International Journal for Parasitology* 7, 335-345.

- Barbosa, F., Pimentel, S. and Sampaio, I. B. M. (1987). The effect of seasonal temperature and experimental illumination on reproductive rate in the snail biomphalaria glabrata. *Brazilian Journal of Medical and Biological Research* **6**, 31-59.
- Bhattacharya, S. C., Sharma, R., Dhiman, H. and Mitra, A. P. (2006). Climate change and malaria in India, Special Section: *Climate Change and India, Current Science* 90 (3), 369-376.
- Bi, P., Tong, S., Donald, K., Parton KA. and Ni, J. (2003). Climatic variables and transmission of malaria: a 12-year data analysis in Shuchen County, China. *Public Health Reports* 118, 65-71.
- Bouma, M.J., Dye, C. and Van, d. HJ. (1996). Falciparum malaria and climate change in the North West Frontier Province of Pakistan. *American Journal of Tropical Medicine and Hygiene* 55 (2), 131-137.
- Boussaa, S., Guernaoui, S., Pesson, B. and Boumezzough, A. (2005). Seasonal fluctuations of phlebotomine sand fly populations (Diptera: Psychodidae) in the urban area of Marrakech, Morocco. *Acta Tropica* **95**, 86-91.
- Briet, O. J.T., Vounatsou, P., Gunawardena, D. M., Gawrie, Galappaththy, NL. G. and Amerasinghe, H. P. (2008) (a). Temporal correlation between malaria and rainfall in Sri Lanka. *Malaria Journal* 7, 1-14.
- Briet, O.J.T., Vounatsou, P. and Amerasinghe, H. P. (2008) (b). Malaria seasonality and rainfall seasonality in Sri Lanka are correlated in space. *Geospatial Health* **2** (**2**), 183-190.
- Buttiker, W. and Lewis, DJ. (1983). Insects of Saudi Arabia: detection of faunal biotypes of medical importance in South-Western Saudi Arabia using landsat images. *Fauna Saudi Arabia* **3**, 10-27.
- Cardenas, R., Claudia, M., Alfonso, J. and Franco-Paredes, C. (2006). Studied impact of climate change variability in the occurrence of leishmaniasis in Northeastern Colombia. *American Journal of Tropical Medicine and Hygiene* **75** (2), 273–277.
- Centers for Disease Control and Prevention, (CDC). (2010). Leishmaniasis.
 http://www.cdc.gov/parasites/leishmaniasis/. Last updated: November, 2010, last accessed: April, 2012.

• Centres for Disease Control and Prevention, (CDC). (2010). Leishmaniasis, Schema of the Life Cycle of Leishmaniasis.

http://www.cdc.gov/parasites/leishmaniasis/biology.html . Last updated: November, 2010, last accessed: April, 2012.

- Centers for Disease Control and Prevention, (CDC). (2010). Malaria. http://www.cdc.gov/malaria/ . Last updated: April, 2012, last accessed: April, 2012.
- Centers for Disease Control and Prevention, (CDC). (2010). Schema of the Life Cycle of Malaria.
 http://www.cdc.gov/malaria/about/biology/. Last updated: February, 2010, last accessed: April, 2012.
- Centers for Disease Control and Prevention, (CDC). (2010). Schistosomiasis.
 <u>http://www.cdc.gov/parasites/schistosomiasis/index.html</u>. Last updated: November, 2010, last accessed: April, 2012.
- Centres for Disease Control and Prevention, (CDC). (2010). Prevention and Control, Leishmaniasis. <u>http://www.cdc.gov/parasites/leishmaniasis/prevent.html</u> last updated: November, 2010, last accessed: April 2012.
- Central Department of Statistic & Information, (CDSI). (2004-2010). Ministry of Economy and Planning, Riyadh, KSA.
 <u>http://www.cdsi.gov.sa/english/index.php</u> last accessed: April, 2012.
- Chaves, L.F. and Pascual, M. (2006). Climate Cycles and Forecasts of Cutaneous Leishmaniasis, a Nonstationary Vector-Borne Disease. *PLoS Medicine* **3** (**8**), 1320-1328.
- Coelho, J. R. and Fernando, SM. B. (2006). The effects of temperature change on the infection rate of biomphalaria glabrata with Schistosoma mansoni. *Memorias do Instituto Oswaldo Cruz* 101 (2), 223-224.
- Confalonieri, U., Menne, B., Akhtar, R., Ebi, K.L., Hauengue, M., Kovats, R.S., Revich, B. and Woodward, A. (2007). Human health. Climate Change: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, IPCC Climate Change. *Cambridge University Press* 8, 391-431.

 Costa, J. (2011). Mucocutaneous leishmaniasis, epidemiological and clinical modalities. Harvard School of Public Health Centro Moniz, Cruz, Brazil Studies Program, DRCLAS, Harvard University.

http://www.drclas.harvard.edu/files/Harvard%20Course%202011%20Leishmaniasis%20-%20Jackson%20Costa.pdf_last accessed: April, 2012.

- Costello, A., Abbas, M., Allen, A., Ball, S., Bellamy, R., Sharon, F., Groce, N., Johnson, A., Kett, M., Lee, M., Levy, C., Maslin, M., McCoy, D., McGuire, B., Montgomery, H., Napier, D., Pagel, C., Patel, J., Oliveira, J., Redclift, N., Rees, H., Rogger, D., Scott, J., Stephenson, J., Twigg, J., Wolff, J. and Patterson, C. (2009). Managing the health effects of climate change. *The Lancet Commissions* 373, 1693-1733.
- Croft, S. L., Sundar, S. and Alan H. F. (2006). Drug Resistance in Leishmaniasis. *Clinical Microbiology Reviews* 19 (1), 111-126.
- Cross, E. and Hyams, K. (1996) (a). The potential effect of global warming on the geographic and seasonal distribution of phlebotomus papatasi in Southwest Asia. *Environmental Health Perspectives* 104 (7), 724-727.
- Cross, E.R., Newcomb, WW. and Tucker, CJ. (1996) (b). Use of weather data and remote sensing to predict the geographic and seasonal distribution of phlebotomus papatasi in Southwest Asia. *American Journal of Tropical Medicine and Hygiene* 56, 530-536.
- Curriero, F.C. (2001). The association between extreme precipitation and waterborne disease outbreaks in the United States, 1948–1994. *American Journal of Public Health* 91 (8), 1194-1199.
- De Savigny, D., Mayombana, C., Mwageni, E., Masanja, H., Minhaf, A., Mkilindi, Y., Mbuya, C., Kasale, H. and Reid, G. (2004). Care-seeking patterns for fatal malaria in Tanzania. *Malaria Journal* 3, 1-15.
- Dennis, S. G., Simon H. I., David, S. I., Kimutai, B. and Robert, W. (2002). Meteorologic influences on plasmodium falciparum malaria in the Highland Tea Estates of Kericho, Western Kenya. *Emerging Infectious Diseases Journal* 8 (12), 1404-1408.
- Dobson, P. A. and Carter, E. R. (1996). Infectious diseases and human population history. *Bioscience* 46, 115-126.

- El-Badry, A., Al-Juhani, A., El-Kheir, I. and Al-Zubiany, S. (2009). Distribution of sandflies in El-Nekheil Province, in Al-Madinah Al-Munawwarah Region, Western of Saudi Arabia. *Research Journal of Parasitology* 4 (1), 1-11.
- El-Hassan, A. A. (1974). Laboratory studies on the direct effect of temperature on bulinis truncatus and biomphalaria alexandrina the snail intermediate host of schistosomes in Egypt. *Folia Parasitologica* **21**, 181-187.
- Farrell, J. P. (2002). The immunology of cutaneous leishmania: experimental infections and human disease. World Class Parasites Leishmania, Kluwer Academic Publishers, Boston, Dordrecht and London. *World Class Parasites* 4, 193.
- Faulde, M., Schrader, J., Heyl, G., Amirih, M. and Hoerauf, A. (2008). Zoonotic cutaneous leishmaniasis outbreak in Mazar-e Sharif, northern Afghanistan: an epidemiological evaluation. *International Journal of Medical Microbiology* 298, 543-550.
- Faulde, M., Schrader, J., Heyl, G. and Hoerauf, A., (2009). High efficacy of integrated preventive measures against zoonotic cutaneous leishmaniasis in northern Afghanistan, as revealed by Quantified Infection Rates. *Acta Tropica* **110**, 28-34.
- Franke, CR., Ziller, M., Staubach C. and Latif, M. (2002). Impact of the El Nino/southern oscillation on visceral leishmaniasis, Brazil. *Emerging Infectious Diseases* **8**, 914-917.
- Freeman, T. and Bradley, M. (1996). Temperature is predictive of severe malaria years in Zimbabwe. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 90, 232-238.
- Freeman, T. (1994). Manyuchi Dam malaria outbreak. Harare, Zimbabwe. Investigation. Proposed by P.M.D Midland Province GTZ. <u>http://malair.org/Docs/Manyuchi.1994.doc</u> last accessed: April, 2012.
- Gaafar, M., Malik, M., Osheik S., El-Taher, A. and Abdin, S. M. (1998). Clinical Aspects of Malaria in the Asir Region, Saudi Arabia. *Annals of Saudi Medicine* **18** (**1**), 15-17.
- Gage, L. K., Thomas, R. B., Rebecca J. E. and Edward, B. H. (2008). Climate and vectorborne diseases. *American Journal of Preventive Medicine* **35** (**5**), 436-450.

- Garout, M. A. (2000). *Malaria control in Saudi Arabia*. A thesis of Master degree, School of Epidemiology and Pubic health, Facility of Medicine, University of Manchester, UK.
- Global Alert and Response, (GAR). (2009). WHO. Report on Global Surveillance of Epidemic-prone Infectious Diseases - Leishmaniasis: <u>http://www.who.int/csr/resources/publications/CSR_ISR_2000_lleish/en/index.html</u> last accessed: April, 2012.
- Gramiccia, M. and Gradoni, L. (2005). The current status of zoonotic leishmaniasis and approaches to disease control. *International Journal for Parasitology* **35**, 1169-1180.
- Grasso, M., Matteo, M., Aline, C. and Markandya, A. (2010). *The health effects of climate change*. A survey of recent quantitative research working paper. The Center for Research on Energy and Environmental Economics and Policy at Bocconi University.
- Gryseels, B., Polman, K., Clerinx, J. and Kestens, L. (2006). Human Schistosomiasis. Seminar. *Lancet* 368, 1106-1118.
- Gubler, D. J., Reiter, P., Ebi, K. L., Yap, W., Nasci, R. and Patz, J. A. (2001). Climate variability and change in the United States: potential impacts on vector- and rodent-borne diseases. *Environmental Health Perspectives* **109** (**2**), 223-233.
- Guernaoui, S., Boumezzough, A. and Laamrani, A. (2006). Altitudinal structuring of sand flies (Diptera: Psychodidae) in the High-Atlas mountains (Morocco) and its relation to the risk of leishmaniasis transmission. *Acta Tropica* **97**, 346-351.
- Habib, R. (2011). *Improving Health in a Changing Climate*. Department of Environmental Health, Faculty of Health Sciences, American University of Beirut.
- Haddad, O. (1990). Malaria: clinical aspects and treatment. *Annals of Saudi Medicine* **10**, 346-348.
- Hamad, A. A., Abd-Elhamid, D. N., David, E. A., Haider, A. G., Abdel-Muhsin, A. A., Gwiria, M.H., Satti, T. G., Theander, M. C., Hamza, A. and Babiker, D. (2002). An Elnaiem A marked seasonality of malaria transmission in two rural sites in eastern Sudan. *Elsevier Science Acta Tropica* 83, 71-82.
- Hana, C. and Windmuller, E. (2001). *Malaria control and its effects on the environment*. Environmental studies, University of Arhus, Finlandsgade Denmark, Miljolaere.

- Hanafi, A. H., David, J. F., Govind, B. M., Moustafa, O. I. and Andrew, J. M. (2007). Bionomics of phlebotomine sandflies at a peacekeeping duty site in the north of Sinai, Egypt. *Acta Tropica* 101, 106-114.
- Harold, W., Anthony, J. and McMichael, A. (2000). *Climate variables and health*. Senior honors thesis in environmental science. Center for Environmental Studies, Brown University, USA.
- Houmsou, R., Kela, S.L., Suleiman, M. M. and Ogidi, J. A. (2009). Endemicity and intensity of vesical schistosomiasis: epidemiological profile of two local government areas of Benue State, Nigeria. *The Internet Journal of Health* 9 (2), 1251-1258.
- Hunter P.R. (2003). Climate change and waterborne and vector-borne disease. The Society for Applied Microbiology. *Journal of Applied Microbiology* **94**, 37S-46S.
- Hussein, A. and Bin-Dajem, S. M. (2008). Prevalence of urinary schistosomiasis and infections with trematode larval stages in bulinus truncatus snails from Qena, Upper Egypt. *Journal of Applied Sciences Research* 4 (11), 1610-1617.
- Ibrahim, A. A. and Abdoon, A. M. (2005). Distribution and population dynamics of phlebotomus sandflies in an endemic area of cutaneous leishmaniasis in Asir Region. *Journal of Entomology* 2 (1), 102-108.
- IPCC, Solomon, S. (2007). Summary for policy makers, climate change. The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, U.K. and New York, USA.
- Islam, M. N. (2009). Rainfall and temperature scenario for Bangladesh. *Open Atmospheric Science Journal* **3**, 93-103.
- IUCN, (2009). Species and Climate Change: More than Just the Polar Bear. IUCN/Species Survival Commission. Cambridge, UK. Available at: <u>http://www.iucn.org/what/tpas/biodiversity/resources/publications/?4562/Species-and-Climate-Change</u>. Last updated: January, 2010, last accessed: April, 2012.
- Interagency Working Group on Climate Change and Health, (IWGCCH). (2010). *A Human Health Perspective on Climate Change*. USA: Environmental Health Perspectives

and the National Institute of Environmental Health Sciences. Interagency Working Group on Climate Change and Health.

- Khalifa, M. A. (2009). *Education Directory of workers in the national program to eradicate malaria in Saudi Arabia*. MOH, Malaria Department. (Arabic Version).
- Kamiura, A. and Reforma, L. (2007). *Parasitic mechanisms of immune evasion*: leishmania. Human Biology, Stanford University: <u>http://www.stanford.edu/class/humbio153/ImmuneEvasion/index.html</u> last accessed: April, 2012.
- Kasap, O., Belen, A., Kaynas, S., Simsek, F., Biler, L., Ata, N. and Alten, B. (2009). Activity patterns of sand fly (diptera: psychodidae) species and comparative performance of different traps in an endemic cutaneous leishmaniasis focus in Cukurova Plain, Southern Anatolia, Turkey. *Acta Veterinaria Brno* 78, 327-335.
- Keiser, J., Utzinger, J., De Castro, M., Smith, T., Tanner, M. and Singer B. (2004). Urbanization in Sub-Saharan Africa and implication for malaria control. *The American Society of Tropical Medicine and Hygiene* **71** (12), 118-127.
- Khatri, M., Haider, N., Di Muccio, T. and Marina, G. (2006). Cutaneous leishmaniasis in Yemen; clinicoepidemiologic features and a preliminary report on species identification. *International Journal of Dermatology* 45, 40-45.
- Kigotho, AW. (1997). Services stretched as malaria reaches Kenyan highlands. *Lancet* 350, 422-425.
- Kilian A., Langi P., Talisuna, A. and Kabagambe, G. (1999). Rainfall pattern, El Nino and malaria in Uganda. Short report. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 93, 22-23.
- Kristan, M., Abeku, T., Beard, J., Okia, M., Rapuoda, B., Sang, J. and Cox, J. (2008). Variations in entomological indices in relation to weather patterns and malaria incidence in East African highlands: implications for epidemic prevention and control. *Malaria Journal* 7, 231.
- Laamrani, H., Mahjour, J., Madsen, H., Khallaayoune, K. and Gryseels, B. (2000). Schistosoma haematobium in Morocco, moving from control to elimination. *Parasitology Today* 16 (6), 257-260.

- Lake, I. R., Gillespie, I.A., Bentham, G., Nichols, G. L., Lane, C., Adak, G.K. and Threlfall, E. J. (2009). A re-evaluation of the impact of temperature and climate change on food-borne illness. *Epidemiology and Infection* 137 (11), 1538-1547.
- Langford, I.H., and Bentham, G. (1995). The potential effects of climate change on winter mortality in England and Wales. *International Journal of Biometeorology* **38**, 141-147.
- Lindgren, E., Johannesson, M. and Ebi, K. L. (2010). *Climate change and communicable diseases in the EU Member States*. Handbook for national vulnerability, impact and adaptation assessments. Stockholm, ISBN 978-92-9193-206-1, doi 10.2900/27967. European Centre for Disease Prevention and Control (ECDC).
- Lindsay, SW. and Martens WJM. (1998) Malaria in the African highlands: past, present and future. *Bull World Health Organ* **76**, 33-45.
- Lipp, K., Huq, A. and Colwell, R. (2002). Effects of global climate on infectious disease: the cholera model. Clinical microbiology reviews. *American Society for Microbiology* 15 (4), 757-770.
- Malakooti, M.A., Biomndo, K. and Shanks, G.D. (1998). Reemergence of epidemic highland malaria in the highlands of western Kenya. *Emerging Infectious Diseases* 4 (4), 671-676.
- Martens, P., Kovats, R.S., Nijhof, S., De Vries, P., Livermore, M.T.J., Bradley, D.J., Cox, J. and McMichael, A.J. (1999). Climate Change and Future Populations at Risk of Malaria. *Global Environmental Change* 9, S89-S107.
- McMichael, A.J., Haines, A., Slooff, R. and Kovats, S. (1996). *Climate change and human health*. Geneva, World Health Organization WHO/WMO/UNEP (WHO/EHG/96.7).
- McMichael, A.J., Campbell-Lendrum, D.H., Corvalan, C.F., Ebi, K.L., Githeko, A.K., Scheraga, J.D. A. (2003). *Woodward, Climate change and human health, risks and responses.* World Health Organization, Geneva.
- Medical Ecology Organisation, (MEO). (2004). Summary of basic science and clinical information for schistosomiasis.
 http://www.medicalecology.org/water/schistosomiasis/schistosomiasis.htm Last accessed: 2004.

- Mendis, K., Sina, B., Marchesini, P. and Carter, R. (2001). The neglected burden of plasmodium vivax malaria. *American Journal of Tropical Medicine and Hygiene* 64 (1), 97-105.
- Meteorological and Environmental Protection Section (MEPS). (2004). The Center of Information and Documentation in the Ministry of Defence and Aviation in Saudi Arabia, Meteorological Department. *Annual report*.
- Meteorological and Environmental Protection Section (MEPS). (2007). The Center of Information and Documentation in the Ministry of Defence and Aviation in Saudi Arabia, Meteorological Department. *Annual report*.
- Meteorological and Environmental Protection Section (MEPS). (2008). The Center of Information and Documentation in the Ministry of Defence and Aviation in Saudi Arabia, Meteorological Department. *Annual report*.
- Meteorological and Environmental Protection Section (MEPS). (2009). The Centre of Information and Documentation in the Ministry of Defence and Aviation in Saudi Arabia, Meteorological Department. *Annual report*.
- Michelson, H. E. (1960). The effects of temperature on growth and reproduction of Australorbis Glabratus in the laboratory. *American Journal of Epidemiology* **73**, 66-74.
- Ministry of Health, KSA, Statistic Department. The Statistical Books (2000 2009). A review of health situation, Leishmaniasis.
 <u>http://www.moh.gov.sa/Ministry/Statistics/Book/Pages/default.aspx</u>. Last updated: August, 2011, last accessed: April, 2012.
- Ministry of Health, (MOH), KSA, Statistic Department. The Statistical Books (2002 2009. A review of health situation, Malaria.
 <u>http://www.moh.gov.sa/Ministry/Statistics/Book/Pages/default.aspx</u>. Last updated: August, 2011, last accessed: April, 2012.
- Ministry of Health, (MOH), KSA, Statistic Department. The Statistical Books (2002 2009). A review of health situation, Schistosomiasis.
 <u>http://www.moh.gov.sa/Ministry/Statistics/Book/Pages/default.aspx</u>. Last updated: August, 2011, last accessed: April, 2012.

- Ministry of Health, (MOH). (2006). Presentation to the Regional Workshop on Adaptation Strategies to Protect Health under Climate Change Variability and Change in Water Stressed Countries in EMR. Fact file.
- Mohanty, P. and Panda, U. (2003). Heatwave in Orissa: A study based on heat indices and synoptic features – Heatwave conditions in Orissa. Regional Research Laboratory, Bhubaneswar. *Institute of Mathematics and Applications* 15, 251-254.
- Molina, J., Alberto, O., Ivan, M. and Guhl, F. (2008). Seasonality of Lutzomyia fairtigi (Diptera: Psychodidae: Phlebotominae). A species endemic to Eastern Colombia. *Rio de Janeiro* 103 (5), 477-482.
- Murray, H., Berman, J.D., Davies, C.R., Saravia, N.G., (2005). Advances in leishmaniasis. *Lancet* 366, 1561-1577.
- Narciso, H. and Ruelan, J. (2007). Infectious Public Health Challenges. Stanford University Parasites & Pestilence: <u>http://www.stanford.edu/class/humbio103/ParaSites2006/Leishmaniasis/Index.htm</u> last accessed: April, 2012.
- Narvaez, J.F., Lara, S.B., Wynsberghe, C., Nicole R., Rebollar-Tellez, E.A., Vargas-Gonzalez, A. and Albertos-Alpuche, N.E. (2003). Seasonal transmission of leishmania Mexicana in the state of Campeche, Yucatan Peninsula. *Memorias do Instituto Oswaldo Cruz* 98 (8), 995-998.
- National Institute of Allergy and Infection Diseases, (NIAID). (2010)
 <u>http://www.niaid.nih.gov/topics/malaria/pages/lifecycle.aspx</u>. Last updated: April, 2012, last accessed: April, 2012.
- National Weather Service, (NWS). (2005), Climate Predict Centre: <u>http://www.cpc.ncep.noaa.gov/products/analysis_monitoring/ensostuff/ensofaq.shtml#HOWOFTEN</u>. Last updated: April, 2012, last accessed: April, 2012.
- Oaks, S.C. J., Mitchell, V.S., Pearson, G.W., and Carpenter, C.C. J. (1991). *Malaria: obstacles and opportunities*. A report of the committee for the study on malaria prevention and control: status review and alternative strategies. Division of International Health, Institute of Medicine. Washington, DC: National Academy Press.
- Oesterholt, M.J.A.M., Bousema, JT., Mwerinde, O.K., Harris1, C., Lushino, P., Masokoto, A., Mwerinde, H., Mosha, F.W. and Drakeley, CJ. (2006). Spatial and

temporal variation in malaria transmission in a low endemicity area in northern Tanzania. *Malaria Journal* **5** (**9**), 2875-2898.

- Oluleye, A. and Akinbobola, A. (2010). Malaria and pneumonia occurrence in Lagos, Nigeria: role of temperature and rainfall. *African Journal of Environmental Science and Technology* 4 (8), 506-516.
- Omar, M.S., Gaafar, M.M., Al-Amari, O.M., Abdalla, S.E. and Moosa, R.A. (1999). The rapid manual paraSightTH-F test for diagnosing plasmodium falciparum malaria in Saudi Arabia. *Annals of Saudi Medicine* 19 (2), 159-162.
- Pampana, E. (1969). *A Textbook of malaria eradication* (Second edition). London, Oxford University Press.
- Patz, J.A., Strzepek, K., Lele, S., Hedden, M., Greene, S., Noden B., Hay, S.I., Kalkstein, L. and Beier, J.C. (1998). Predicting key malaria transmission factors, biting and entomological inoculation rates, using modelled soil moisture in Kenya. *Tropical Medicine and International Health* 3, 818-827.
- Patz, J.A. and Lindsay, SW. (1999). New challenges, new tools: the impact of climate change on infectious diseases. *Current Opinion in Microbiology* **2**, 445-451.
- Patz, J.A, Thaddeus, K., Graczyk, N. G. and Amy, Y. V. (2000). Effects of environmental change on emerging parasitic diseases. Elsevier Science Ltd. *International Journal for Parasitology* 30, 1395-1405.
- Patz, J.A., Githeko, A., McCarty, J., Hussein, S., Confalonieri, U. and Wet, N. (2002). Climate change and infectious diseases. *Climate Change and Human Health* **6**, 103-132.
- Philipkoski, K. (2008). Black Death's Gene Code Cracked (Article). <u>http://www.wired.com/medtech/health/news/2001/10/47288</u> last accessed: April, 2012.
- Pimentel-Souza, N.B. and Resende, D-F. (1990). Effect of Temperature on the Reproduction of the Snail Biomphalaria GlabrataBraz. *Brazilian Journal of Medical and Biological Research* 23 (5), 441-449.
- Pimentel-Souza, N.B., Schettino, M. and Lautner, R.J. (1988). Influence of experimental illumination and seasonal variation on crossbreeding mating in the snail Biomphalaria Glabrata. *Memorias do Instituto Oswaldo Cruz* 83 (1) 79-85.

- Reithinger, R., Teodoro, U. and Davies, C.R. (2001). Topical Insecticide Treatments to Protect Dogs from Sandfly Vectors of Leishmaniasis, London School of Hygiene & Tropical Medicine, London, United Kingdom. *Emerging Infectious Diseases* 7 (5), 872-876.
- Reithinger, R., Aadil, K., Kolaczinski, J., Mohsen, M. and Hami, S. (2005). Social impact of leishmaniasis, Afghanistan. *Emerging Infectious Diseases* **11** (**4**), 634-636.
- Reithinger, R., Dujardin, J., Louzir, H., Pirmez, C., Alexander, B. and Brooker, S. (2007). Cutaneous leishmaniasis. Review. *The Lancet Infectious Diseases* **7**, 581-596.
- Rey, J.R. (2007). *Leishmaniasis*. University of Florida, *ENY*-738. http://edis.ifas.ufl.edu/pdffiles/IN/IN70500.pdf_last accessed: April, 2012.
- Ridley, R. (2001). Malaria drug development & Malaria Africa Emerging Prevention and Control Strategies. Reported from a symposium held at the February 17, 2001 AAAS. Annual Meeting, San Francisco.
- Riedel, D. (2004). Human health and well-being. *Climate change: impacts and adaptation A Canadian Perspective*. D. Lemmen and F. Warren, Eds., Climate Change Impacts and Adaptation Directorate. Natural Resources Canada, Ottawa, 151-171.
- Ryan, K., Ray, C., George, N., Drew, A., Lawrence, W. and Plorde, J. (2010). *Sherris Medical Microbiology Pathology*. (Book). Microbiology College of Medicine University of Arizona Tucson, United State.
- Salomon, D.O., Rosa1, J.R., Marina, S., Maria, G.Q., Maria, S.F., Andres M., Visintin, G.R., Spinelli, M.M., Bogado, D. and Molinari, M.L. (2008). Phlebotominae (Diptera: Psycodidae) fauna in the Chaco Region and cutaneous leishmaniasis transmission patterns in Argentina. *Memorias do Instituto Oswaldo Cruz* 103 (6), 578-584.
- Schmidt, G. and Larry, R. (1996). *Foundations of Parasitology*. Wm. C. Brown, 1996 Science – 659 pages. The University of Wisconsin Madison, United States.
- Schriefer, A., Luiz, H., Guimaraes, P. R. L., Machado, M. L., Helio A., Lessa, E.L., Guilherme-Ritt, A., Goes-Neto, A. L.F., Schriefer, L., Riley, W. and Carvalho, E.M. (2009). Geographic Clustering of Leishmaniasis in Northeastern Brazil1. *Emerging Infectious Diseases* 15 (6), 871-876.

- Shati A.A. (2009). Factors affecting the prevalence of human schistosomiasis in Aseer Region, Saudi Arabia. *Journal of biological sciences* 9 (8), 815-819.
- Singh, K. (1999). Studies on the role of climatological factors in the distribution of phlebotomine sandflies (Diptera: Psychodidae) in semi-arid areas of Rajasthan, India. *Journal of Arid Environments* 42, 43-48.
- Singh, N. and Sharma, V. (2002). Patterns of rainfall and malaria in Madhya Pradesh, Central India. *Annals of Tropical Medicine and Parasitology* **96** (4), 349-359.
- Srinivasan, R., Panicker, K.N. and Dhanda, V. (1993). Population dynamics of Phlebotomus papatasi (Diptera: Phlebotomidae) in Pondicherry, India. *Acta Tropica* 54, 125-130.
- Sturrock, R. F., Diaw, I., Talla, M., Niang, P. and Capron, A. (2001). Seasonality in the transmission of schistosomiasis and in populations of its snail intermediate hosts in and around a sugar irrigation scheme at Richard Toll, Senegal. *Parasitology* **123**, S77-S89.
- Subyani, A.M. (2000). Topographic and Seasonal Influences on Precipitation Variability in Southwest Saudi Arabia. JKAU, *Earth Science* **11**, 89-102.
- TDR. (2002). *Strategic Direction for Research, Programme for Research and Training in Tropical Diseases*. Strategic Direction: Leishmaniasis, Avenue Appia 20, 1211 Geneva.
- Teklehaimanot, H.D., Lipsitch, M., Teklehaimanot, A. and Schwartz, J. (2004). Weatherbased prediction of plasmodium falciparum malaria in epidemic-prone regions of Ethiopia. Patterns of lagged weather effects reflect biological mechanisms. *Malaria Journal* 3, 1-11.
- The Center for Food Security & Public Health, (CFSPH). (2009). Leishmaniasis cutaneous and visceral. Factsheets. Institute for International Cooperation in Animal Biologics, Iowa State University, College of Veterinary Medicine.
 http://www.cfsph.iastate.edu/Factsheets/pdfs/leishmaniasis.pdf last accessed: April, 2012.
- The Convention of Biological Diversity, (CBD). Saudi Arabia. (2007) http://www.cbd.int/doc/world/sa/sa-nbsap-01-en.pdf last accessed: April, 2012.

- The Navy Department Library. (2005). *Influenza of 1918 (Spanish flu) and the US Navy*. The Navy Naval Historical Centre Washington navy yard Washington dc 20374-5060. <u>http://www.history.navy.mil/library/online/influenza main.htm</u> last accessed: April, 2012.
- The Vectors Control Administration, (VCA). Annual reports of the Vector Control Administration, (1995 – 2009) Health Affairs Directorate in Asir, Malaria, Leishmaniasis and Schistosomiasis, MOH, Saudi Arabia.
- The World Health Report. (2004). Statistical Annex 1, Deaths by cause, sex and mortality stratum in regions, estimates for 2002, computed by WHO, Annex1, Table2. http://www.who.int/whr/2004/annex/topic/en/annex_2_en.pdf last accessed: April, 2012.
- Trenberth, K. (1997). The Definition of El Nino. Bulletin of American Meteorological Society 78 (12), 2771-2777.
- Tulu, AN. (1996). Determinants of malaria transmission in the highlands of Ethiopia: the impact of global warming on morbidity and mortality ascribed to malaria, (Book). London School of Hygiene and Tropical Medicine.
- United Nations Framework Convention on Climate Change, (UNFCCC). (2007). *Climate change, Impacts, vulnerabilities and adaptation in developing countries,* http://unfccc.int/resource/docs/publications/impacts.pdf last accessed: April, 2012.
- Vergel, C., Palacios, R. and Cadena, H. (2006). Evidence for leishmania (Vienna) parasites in the skin and blood of patients before and after treatment. *The Journal Infectious Diseases* 194 (4), 503-511.
- Wittner, M. and Tanowitz, H.B. (2000). Leishmaniasis in infants and children. *Seminars in Pediatric Infectious Diseases* **11** (**3**), 196-201.
- Woodruff, R.E., Guest, C.S., Garner, M.G., Becker, N., Lindesay, J., Carvan, T. and Ebi, K. (2002): Predicting Ross River virus epidemics from regional weather data. *Epidemiology* 13 (4), 384-393.
- World Health Organization, (WHO). (1994). *The leishmaniasis*. Report of Expert Committee, World Health Organization Technical Report Series 701, WHO, Geneva 1994.

- World Health Organization, (WHO). (1999). Sustainable Development and Healthy Environments, El Nino and Health. Protection of the Human Environment, (WHO/SDE/PHE). Task Force on Climate and Health, Geneva.
- World Health Organization, (WHO). (2000). The leishmaniasis and Leishmania/HIV coinfections.

http://www.knowledgebank.irri.org/factsheetsPDFs/Health and Nutrition/whoFS hiv leishmaniasesCoInf ect.pdf last accessed: April, 2012.

- World Health Organization, WHO/CDS/CSR/ISR/. (2000). Department of Communicable Disease Surveillance and Response. <u>http://www.who.int/csr/resources/publications/surveillance/Leishmaniasis.pdf</u>. Last updated: February, 2004, last accessed: April, 2012.
- World Health Organization, (WHO). (2003). *Climate change and human health*. Risks and Responses, World Health Organization, WHO Press.
- World Health Organization, (WHO). (2005). WHO/HTM/MAL/. *Malaria control in complex emergencies*. An inter-agency field handbook, 1107.
- World Health Organization, (WHO). (2006). *Control of leishmaniasis*. Report of a WHO Expert Committee. Geneva, World Health Organization. Technical Report (**793**).
- World Health Organization, (WHO). (2006). *Preventing disease through healthy environments: Towards an Estimate of the Environmental Burden of Disease*. World Health Organization.
- World Health Organization, (WHO). (2006). WHO/HTM/MAL. Systems for the early *detection of malaria epidemics in Africa*. An analysis of current practices and future priorities.
- World Health Organization, (WHO). (2009). Leishmaniasis.
 <u>http://www.who.int/neglected_diseases/integrated_media_leishmaniasis/en/index.html</u>. Last updated: March, 2010, last accessed: April, 2012.
- World Health Organization, (WHO). (2010). The disease and its epidemiology, the vector. <u>http://www.who.int/leishmaniasis/disease epidemiology/en/index.html</u> last accessed: April, 2012.
- World Health Organization, (WHO). (2010). Malaria. Fact sheet.
 http://www.who.int/mediacentre/factsheets/fs094/en/. Last updated: April, 2012, last accessed: April, 2012.
- World Health Organization, (WHO). (2010). Schistosomiasis. Fact sheet. <u>http://www.who.int/mediacentre/factsheets/fs115/en/index.html</u>. Last updated: January, 2012, last accessed: April, 2012.
- World Health Organization, (WHO). (2011). 10 Facts on Climate Change and Health. Fact Sheet. <u>http://www.who.int/features/factfiles/climate_change/facts/en/index.html</u> last accessed: April, 2012.
- World Health Organization, (WHO). (2011). Water Sanitation and Health (WSH) Waterrelated disease.
 http://www.who.int/water_sanitation_health/diseases/en/last accessed: April, 2012.
- World Resources Institute, (WRI). (1998). Environmental change and human health, Controlling disease vectors in the local environment. The United Nations Environment Programme, the United Nations Development Programme, and the World Bank. World Resources 1998-1999.

http://www.wri.org/publication/content/8371 last accessed: April, 2012.

- Woube, M. (1997). Geographical distribution and dramatic increases in incidences of malaria: consequences of the resettlement scheme in Gambela, SW Ethiopia. *Indian Journal of Malariology* 34 (3), 140-63.
- Yaghoobi-Ershadi, A.A. (1999). Entomological survey of sandflies (Diptera: Psychodidae) in a new focus of zoonotic cutaneous leishmaniasis in Iran. Elsevier Science. *Acta Tropica* 73, 321-326.
- Yaghoobi-Ershadi, A.A., Hanafi-Bojd, A.A., Akhavan, A.R., Zahrai-Ramazani, C. and Mohebali, M. (2001). Epidemiological study in a new focus of cutaneous leishmaniasis due to Leishmania major in Ardestan town, central Iran. *Acta Tropica* 79, 115-121.
- Yang, G., Utzinger, J., Le-Ping, Q., Sun, V., Penelope, T.M. and Xiao-Nong, Z. (2007). Effect of temperature on the development of Schistosoma japonicum within Oncomelania hupensis, and hibernation of O. hupensis. *Journal of Parasitology Research* 100, 695-700.
- Ye, Y., Louis, V.R., Simboro, S. and Sauerborn, R. (2007). Effect of meteorological factors on clinical malaria risk among children. An assessment using village-based meteorological stations and community-based parasitological survey. *BioMed Central Public*

Appendix

Appendix A

<u>Malaria</u> <u>Appendix A.1</u>

Knannse		
Month	Unstandardized Coefficients	Significance
Jan	-0.10	0.70
Feb	-0.06	0.86
Mar	0.03	0.90
Apr	-0.20	0.31
May	-0.16	0.49
Jun	-0.43	0.09
Jul	-0.10	0.62
Aug	-0.36	0.48
Sep	0.02	0.99
Oct	0.37	0.44
Nov	0.04	0.88
Dec	-0.20	0.40

Table 1. Correlation of DT Ln malaria in all Asir against maximum temperature in Khamise



Figure 1. DT Ln malaria in Asir with maximum temperature in Khamise for each

Majaredan		
Month	Unstandardized Coefficients	Significance
Jan	-0.151	0.677
Feb	0.168	0.532
Mar	0.187	0.599
Apr	-0.29	0.920
May	0.302	0.292
Jun	0.493	0.262
Jul	-0.29	0.931
Aug	0.137	0.360
Sep	-0.052	0.722
Oct	0.260	0.443
Nov	-0.362	0.515
Dec	0.680	0.044

Table 2. Correlation of DT Ln malaria in TA against maximum temperature in Majaredah



Figure 2: DT Ln malaria in TA with maximum temperature in Majaredah for each

<u> </u>		
MONTH	Unstandardized Coefficients	Significance
Jan	-0.248	0.272
Feb	0.252	0.427
Mar	0.705	0.026
Apr	-0.082	0.838
May	-0.350	0.316
Jun	-0.886	0.034
Jul	-0.471	0.047
Aug	-1.406	0.161
Sep	-0.383	0.683
Oct	-0.448	0.470
Nov	-0.029	0.947
Dec	-0.255	0.188

Table 3. Correlation of DT Ln malaria in TQ against maximum temperature in Khamise



Figure 3. DT Ln malaria in TQ with maximum temperature of Khamise for each

Month	Unstandardized Coefficients	Significance
Jan	-0.05	0.92
Feb	-0.46	0.37
Mar	0.03	0.92
Apr	-0.52	0.10
May	-0.13	0.62
Jun	-0.32	0.35
Jul	-0.40	0.29
Aug	-0.58	0.14
Sep	0.25	0.82
Oct	0.96	0.18
Nov	-0.12	0.74
Dec	-0.43	0.46

Table 4. Correlation of DT Ln malaria in Asir against mean temperature in Khamise



MONTH	Unstandardized Coefficients	Significance
Jan	0.125	0746
Feb	-0.160	0.767
Mar	1.01	0.019
Apr	-0.455	0.491
May	-0.172	0.661
Jun	-0.719	0.210
Jul	-0.860	0.051
Aug	-1.457	0.063
Sep	-1.027	0.344
Oct	-0252	0.795
Nov	-0.684	0.223
Dec	-0473	0.330

Table 5 Correlation of DT Ln malaria in TQ against mean temperature in Khamise



Khamise		
MONTH	Unstandardized Coefficients	Significance
Jan	0.33	0.27
Feb	-0.18	0.70
Mar	0.22	0.49
Apr	-0.63	0.27
May	-0.37	0.42
Jun	0.01	0.97
Jul	-0.44	0.30
Aug	-0.01	0.98
Sep	-1.24	0.38
Oct	0.63	0.34
Nov	-0.17	0.51
Dec	0.25	0.46

Table 6 :Correlation of DT Ln malaria in all Asir against minimum temperature in Khamise



Figure 6 : DT Ln malaria in all Asir with minimum temperature for each month

MONTH	Unstandardized Coefficients	Significance
Jan	1.560	0.09
Feb	0.118	0.832
Mar	0.124	0.673
Apr	-0.109	0.839
May	-0.170	0.643
Jun	0.216	0.550
Jul	0.343	0.073
Aug	-0.013	0.912
Sep	-0.073	0.469
Oct	-0.673	0.283
Nov	-0.940	0.137
Dec	1.211	0.035

Table 7: Correlation of DT Ln malaria in TA against mean temperature in Majaredah



Figure 7: DT Ln malaria in TA with minimum temperature in Majaredah for each

Knamise		
MONTH	Unstandardized Coefficients	Significance
Jan	0.379	0.128
Feb	-0.272	0.578
Mar	0.798	0.131
Apr	-1.472	0.181
May	-0.480	0.499
Jun	-0.198	0.767
Jul	-0.980	0.052
Aug	-1.097	0.240
Sep	-0723	0.621
Oct	0.072	0.934
Nov	-0.522	0.175
Dec	0.307	0.285

Table 8: Correlation of DT Ln malaria in TQ against minimum temperature in Khamise



Figure 8: DT Ln malaria in TQ with minimum temperature in Khamise for each

	0	
MONTH	Unstandardized Coefficients	Significance
Jan	0.06	0.81
Feb	-0.38	0.38
Mar	-0.23	0.10
Apr	0.19	0.36
May	0.33	0.20
Jun	0.26	0.03
Jul	0.32	0.11
Aug	-0.16	0.44
Sep	0.17	0.73
Oct	-0.13	0.57
Nov	-0.15	0.42
Dec	-0.07	0.85

Table 9: Correlation of DT Ln malaria in
Asir against Ln. rainfall in Khamise



MONTH	Unstandardized Coefficients	Significance
Jan	0216	0.205
Feb	0.299	0.097
Mar	0.205	0.257
Apr	0.430	0.009
May	0.343	0.014
Jun	0.413	0.033
Jul	-0.026	0.874
Aug	0.204	0.010
Sep	0.106	0.453
Oct	-0.188	0.441
Nov	0.139	0.466
Dec	0.312	0.244

Table 10: Correlation of DT Ln malaria in TA against rainfall in Rejal



Table 11: Correlation of DT Ln malaria in

MONTH	Unstandardized Coefficients	Significance
Jan	0.325	0.069
Feb	0.204	0.435
Mar	0.082	0.794
Apr	0.631	0.039
May	0.400	0.145
Jun	0.461	0.148
Jul	0.307	0.112
Aug	0.404	0.078
Sep	0.564	0.007
Oct	0.373	0.016
Nov	0.464	0.133
Dec	0.404	0.022

TQ against rainfall in Sarat



MONTH	Unstandardized Coefficients	Significance
Jan	0.03	0.51
Feb	0.11	0.29
Mar	0.01	0.85
Apr	0.07	0.09
May	0.02	0.55
Jun	-0.01	0.78
Jul	-0.02	0.71
Aug	0.01	0.73
Sep	0.08	0.45
Oct	0.03	0.57
Nov	0.00	0.88
Dec	0.06	0.20

Table 12: Correlation of DT Ln malaria in Asir against humidity in Khamise



Figure 12: DT Ln malaria in all Asir with humidity in Khamise for each month

MONTH	Unstandardized Coefficients	Significance
Jan	0.044	0.268
Feb	0.140	0.163
Mar	-0.115	0.171
Apr	0.060	0.483
May	-0.011	0.843
Jun	-0.017	0.805
Jul	0.00	0.995
Aug	0.009	0.911
Sep	0.009	0.936
Oct	0.032	0.648
Nov	-0.024	0.611
Dec	0.053	0.160

Table 13: Correlation of DT Ln malaria in TQ against humidity in Khamise



Figure 13: DT Ln malaria in TQ with humidity in Khamise for each month

Appendix A.2

1. Maximum Temperature with Malaria in the whole of Asir

			L	R					Ĉ	DL		
Model	M tempe or	Max. temperature only Cof P Cof		ax. erature ith nality	Ma tempe with corre an seaso	ax. erature auto lation nd nality	Ma tempe or	ax. erature lly	Max. temperature with seasonality		Max. temperature with auto correlation and seasonality	
	0.10	P 000	124	P 625	192	P 417	052	P 607	720	P	201	715
0 P	-0.10	.888	.134	.025	.185	.417	055	.097	129	.192	.391	./15
1 P	.010	.906	192	.612	279	.366	.034	.830	-1.08	.165	435	.159
2 P	134	.477	397	.145	262	.253	096	.794	380	.485	499	.791
3 P	288	.213	215	.305	.113	.617	546	.259	527	.290	432	.652
4 P	-021	.889	135	.430	.003	.984	262	.373	563	.157	.324	.755
5 P	013	.861	195	.256	.085	.566	099	.501	627	.090	-1.08	.341
6 P	036	.571	259	.091	103	.451	182	.167	-1.28	.003	963	.261
0-1 P	053	.751	165	.751	.695	.302	.038	.801	925	.170	-235	.611

Table 14: DT Ln malaria and maximum temperature (Sept to Nov)

Table 15: December to March

			L	R					(DL		
Model	Ma temper on	x. rature ly	Max. temperature with seasonality		Ma temper with correl an seasor	ix. rature auto ation id nality	Ma tempe or	ax. erature lly	Max. temperature with seasonality		Max. temperature with auto correlation and seasonality	
	Cof.	Р	Cof	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	.035	.700	- .094	.447	086	.380	327	.107	724	.021	013	.976
1 P	033	.724	- .037	.785	007	.945	394	.064	442	.144	.035	.894
2 P	087	.164	- .070	.640	084	.476	165	.219	259	.429	.035	.939
3 P	071	.153	- .064	.718	.060	.671	138	.191	674	.110	1.24	.058
4 P	096	.058	- .469	.067	349	.090	113	.295	963	.124	629	.498
5 P	119	.100	- .112	.642	.059	.764	074	.627	.287	.560	.143	.841
6 P	168	.295	.050	.827	207	.254	058	.860	016	.971	.177	.777
0-1 P	.038	.129	.058	.611	054	.702	.350	.085	.538	.101	.032	.911

2. Maximum Temperature with Malaria in Tehama of Asir

			L	R		-			0	L		
Model	M tempe or	in. erature lly	M tempe wi seaso	in. rature th nality	M tempe with correl ar seaso	in. rature auto lation nd nality	M tempe on	in. rature lly	M tempe wi seaso	in. rature th nality	Min. temperature with auto correlation and seasonality	
	Cof.	Р	Cof. P		Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	226	.002	.049	.651	.049	.521	137	.279	.042	.855	133	.847
1 P	217	.000	071	.503	071	.344	134	.114	044	.839	301	.687
2 P	195	.000	01 .961		.036	.615	126	.071	056	.786	081	.885
3 P	206	.000	.041	.662	.045	.493	117	.113	066	.738	.068	.879
4 P	204	.000	.01	.931	013	.855	132	.136	.004	.984	.245	.648
5 P	105	.108	027	.725	022	.696	081	.448	.077	.704	.136	.820
6 P	.115	.033	.022	.808	.023	.726	.072	.416	.099	.604	.198	.564
0-1 P	.275	.000	025	.869	024	.826	174	.121	025	.935	471	.654

Table 16: September to March

3. Maximum Temperature with Malaria in Tehama of Qahtan

Table 17: October to March

			L	R					0	L	L	
Model	Ma tempe on	ax. rature ly	Ma tempe wi season	ax. rature th nality	Ma tempe with correl ar seaso	ax. rature auto lation nd nality	Ma tempe on	ax. rature ly	Max. temperature with seasonality		Max. temperature with auto correlation and seasonality	
	Cof.	Р	Cof. P		Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	.122	.064	.049	.698	.131	.233	107	.371	541	.031	371	.177
1 P	.127	.003	.148	.290	.196	.104	.028	.718	258	.327	020	.947
2 P	.082	.117	093	.561	087	.526	.024	.695	489	.114	495	.159
3 P	.054	.129	212	.170	122	.363	.021	.758	351	.275	205	.548
4 P	.055	.215	191	.296	011	.947	.007	.938	615	.124	486	.260
5 P	21	.745	124	.474	.093	.551	059	.655	214	.546	019	.961
6 P	181	.005	199	.175	084	.514	157	.245	439	.162	446	.204
0-1 P	.145	.008	.116	.459	.213	.119	023	.816	639	.044	355	.310

4. Mean Temperature with Malaria in the whole of Asir

			L	R					0	L		
Model	Me tempe on	Mean. temperature only Mean temperat with seasonal		an. rature th nality	Me tempe with correl ar seaso	an. rature auto lation nd nality	Me tempe on	an. rature ly	Mean. temperature with seasonality		Mean. temperature with auto correlation and seasonality	
	Cof.	Р	Cof. P 192 .106		Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	110	.000	192	.106	108	.249	156	.001	-1.89	.000	-1.81	.002
1 P	118	.000	268	.024	152	.105	173	.001	-2.00	.000	-1.17	.022
2 P	101	.000	332	.005	159	.093	149	.002	-1.92	.000	-1.01	.056
3 P	046	.048	170	.155	.065	.503	093	.050	-1.78	.000	799	.126
4 P	.014	.552	227	.059	123	.195	022	.637	-1.68	.000	879	.104
5 P	.072	.002	179	.138	043	.654	.036	.447	-1.82	.000	-1.22	.017
6 P	.110	.000	252	.037	140	.142	.074	.123	-1.76	.000	-1.22	.017

Table 18: Whole Year

Table 19: September to November

			L	R					О	L		
Model	Me tempe on	an. rature ly	Me tempe wi seaso	Mean. temperature with seasonality Cof. P		an. rature auto lation nd nality	Me tempe on	ean. erature lly	Mean. temperature with seasonality		Mean. temperature with auto correlation and seasonality	
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	01	.896	.198	.596	.199	.520	075	.571	-2.71	.007	995	.606
1 P	027	.781	837	.050	952	.007	087	.645	-3.06	.008	-2.08	.112
2 P	582	.045	-1.01	.008	564	.116	-1.75	.021	-3.63	.004	-2.91	.279
3 P	466	.120	588	.106	.020	.953	-2.18	.006	-3.36	.004	-1.19	.560
4 P	013	.935	245	.335	049	.822	519	.126	-1.80	.012	694	.769
5 P	014	.866	273	.241	110	.580	154	.359	-1.38	.014	-2.92	.246
6 P	040	.612	364	.096	149	.418	218	.186	-1.83	.003	-1.02	.342

			L	R					0	L		
Model	Me tempe on	an. rature ly	Me tempe wi seaso	Mean. to emperature with co seasonality		an. rature auto lation nd nality	Me tempe on	an. rature ly	Mean. temperature with seasonality		Mean. temperature with auto correlation and seasonality	
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	.093	.358	183	396	071	.677	.240	.264	-2.01	.003	-1.40	.091
1 P	.009	.940	024	.918	.140	.452	561	.046	-1.18	.033	.644	.424
2 P	087	.192	127	.631	177	.393	248	.082	-1.88	.008	-1.18	.192
3 P	066	.167	.027	.931	.253	.306	131	.191	-2.21	.008	-1.74	.102
4 P	102	.050	634	.050	530	.041	-1.47	.808	-3.99	.001	-4.07	.014
5 P	146	.085	372	.300	.011 .970		189	.314	-2.70	.006	981	.445
6 P	484	.030	471	.133	462	.061	-1.44	.011	-2.73	.006	-1.79	.167

Table 20: December to March

5. Mean Temperature with Malaria in TQ Table 21: Whole Year

			L	R					0	L		
Model	Me tempe on	an. rature ly	Me tempe wi seaso	an. rature th nality	Mean. temperature ure with auto correlation lity and seasonality		Me tempe on	an. rature ly	Me tempe wi seaso	ean. erature ith nality	Mean. temperature with auto correlation and seasonality	
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	019	.492	209	.178	089	.467	094	.048	-1.28	.000	797	.030
1 P	.038	.176	275	.077	148	.228	048	.309	-1.33	.000	921	.014
2 P	.071	.012	475	.002	304	.012	08	.858	-1.6	.000	-1.19	.014
3 P	.096	.000	317	.035	016	.896	.026	.573	-1.25	.000	554	.139
4 P	.088	.001	291	.047	109	.363	.039	.406	-1.21	.000	731	.048
5 P	.065	.015	202	.171	019 .869		.051	.281	921	.003	378	.313
6 P	.025	.365	284	.053	157	.169	.036	.449	-1.25	.000	-1.09	.007

		LINE	AR RE	GRES	SION			ORE	DINAL	LOGI	ISTIC		
Model	Me tempe on	an. rature ly	Me tempe wi seaso	an. rature th nality	Me tempe with correl ar seaso	an. rature auto lation nd nality	Me tempe or	ean. erature lly	Me tempe wi seaso	an. rature ith nality	Me tempe with corre ar seaso	an. rature auto lation id nality	
	Cof.	Р	Cof. P		Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	
0 P	.124	.081	.100	.630	.288	.115	054	.670	788	.055	230	.620	
1 P	.121	.006	.018	.940	.187	.366	.024	.760	-1.05	.029	690	178	
2 P	.087	.016	194	.459	123	.584	.030	.640	-1.17	.027	896	.102	
3 P	.062	.085	234	.364	.123	.600	.018	.795	-1.27	.024	705	.251	
4 P	.056	.226	264	.305	37	.874	016	.857	-1.80	.005	-1.28	.061	
5 P	038	.626	186	.404	.156	.445	152	.335	-1.01	.047	562	.304	
6 P	198	.008	247	.219	086	.625	212	.179	-1.11	.018	915	.072	
0+1 P	.148	.011	.100	.721	.390	.111	.007	.946	-1.47	.010	782	.229	

Table 22: October to March

	r		-		10 201 1		OL					
			L	К					0	L		
Model	M tempe on	in. rature lly	M tempe wi seaso	in. rature th nality	M tempe with correl ar seaso	in. rature auto lation nd nality	M tempe on	in. rature ly	M tempe wi seaso	in. erature ith nality	Min. temperature with auto correlation and seasonality	
	Cof.	Р	Cof. P		Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	104	.000	.027	.817	.068	.448	133	.008	762	.010	535	.219
1 P	113	.000	06	.956	.004	.962	155	.002	985	.002	882	.097
2 P	099	.000	075	.522	060	.513	138	.006	-1.0	.002	641	.190
3 P	045	.059	.149	.206	.215	.180	070	.152	313	.266	1.33	.073
4 P	.016	.516	.126	.286	.018	.844	.000	.990	272	.324	047	.919
5 P	.079	.001	.156	.189	.074	.428	.066	.179	316	.253	272	.532
6 P	.120	.000	.041	.731	058	.533	.119	.019	151	.585	.177	.697

6. Minimum Temperature with Malaria in the whole of Asir Table 23: The whole of year

Table 24: September to November

	LR						OL					
Model	Min. temperature only		Min. temperature with seasonality		Min. temperature with auto correlation and seasonality		Min. temperature only		Min. temperature with seasonality		Min. temperature with auto correlation and seasonality	
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	022	.758	116	.697	066	.790	06	.668	-1.06	.099	519	.662
1 P	015	.869	356	.462	375	.348	019	.913	-1.02	.298	-7.69	.267
2 P	105	.590	692	.137	236	.576	234	.541	-1.76	.083	1.78	.450
3 P	040	.913	178	.655	.300	.389	190	.030	-2.48	.014	-1.18	.587
4 P	.112	.506	.175	.629	.358	.232	329	.329	-1.98	.027	-1.31	.518
5 P	.039	.707	.160	.465	.132	.666	073	.719	-1.03	.176	-2.21	.386
6 P	.025	.791	.110	.723	.165	.518	072	.712	651	.298	928	.625
Model			L	R					0	L		
-------	---------------------------------------	------	--	------	---	---	------------------	----------------------	---------------------------	---------------------------------	---	---
Model	Min. temperature only Cof. P		Min. temperature with seasonality		M tempe with correl ar seaso	in. rature auto lation nd nality	M tempe on	in. rature lly	M tempe wi seaso	in. erature ith nality	M tempe with correl ar seaso	in. rature auto lation nd nality
	Cof.	Р	Cof.	Р	Cof. P		Cof.	Р	Cof.	Р	Cof.	Р
0 P	165	.056	.185	.275	.225	.092	001	.994	344	.352	069	.883
1 P	.137	.250	.194	.274	.177	.177	384	.142	462	.213	.407	.500
2 P	056	.431	.120	.506	034	034 .814		.048	-1.27	.016	-1.51	.067
3 P	055	.286	.288	.195	.328	.060	-1.11	.291	448	.366	.754	.338
4 P	074	.154	.159	.575	121	.602	116	.297	-1.26	.078	733	.444
5 P	114	.135	.104	.797	027	.934	129	.429	-1.53	.084	499	.678
6 P	320	.086	317	.388	344	.237	586	.159	-2.27	.013	-1.33	.266

Table 25: December to March

7. Minimum Temperature with Malaria in TA

Table 26: Whole of Year

Model			L	R					0	L		
Model	M tempe on	in. erature lly	Min. temperature with seasonality		M tempe with corre ar seaso	in. rature auto lation nd nality	M tempe or	in. rature ly	M tempe wi seaso	in. rature th nality	M tempe with corre ar seaso	in. erature auto lation nd nality
	Cof. P Cof. P Cof.		Р	Cof.	Р	Cof.	Р	Cof.	Р			
0 P	103	.000	.074	.347	011	.845	074	.180	.200	.286	274	.473
1 P	189	.000	041	.608	081	.149	100	.070	.042	.809	691	.139
2 P	149	.000	018	018 .816		.870	115	.040	055	.754	618	.216
3 P	142	.000	.018	.814	.027	.618	064	.242	.054	.757	.570	.151
4 P	051	.105	015	.854	017	.773	015	.778	.080	.648	.185	.681
5 P	.045	.149	049	.455	048	.311	.032	.561	.027	.882	451	.362
6 P	.134	.000	.027	.738	.046 .419		.098	.076	.149	.407	.436	.198
0+1 P	181	.000	.028	.783	.078	.291	100	.091	.176	.434	624	.195

Model			L	R			OL					
	M tempe on	in. rature ly	Min. temperature with seasonality		M tempe with correl ar seaso	in. rature auto lation Id nality	M tempe on	in. erature lly	M tempe wi seaso	in. rature th nality	M tempe with corre ar seaso	in. rature auto lation nd nality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	226	.002	.049	.651	.049	.521	137	.279	.042	.855	133	.847
1 P	217	.000	071	.503	071	.344	134	.114	044	.839	301	.687
2 P	195	.000	01	.961	.036	.615	126	.071	056	.786	081	.885
3 P	206	.000	.041	.662	.045	.493	117	.113	066	.738	.068	.879
4 P	204	.000	.01	.931	013	.855	132	.136	.004	.984	.245	.648
5 P	105	.108	027	.725	022	.696	081	.448	.077	.704	.136	.820
6 P	.115	.033	.022	.808	.023 .726		.072	.416	.099	.604	.198	.564
0+1 P	.275	.000	025	.869	024	.826	174	.121	025	.935	471	.654

Table 27: September to March

8. Minimum Temperature with Malaria in TQ Table 28: Whole Year

Model		LINE	AR RE	GRES	SION			ORE	DINAL	LOGI	STIC	
Model	M tempe on	in. erature lly	Min. temperature with seasonality		M tempe with corre ar seaso	in. erature auto lation nd nality	M tempe or	in. rature ly	M tempe wi seaso	in. rature th nality	M tempe with corre ar seaso	in. rature auto lation nd nality
	Cof.	Р	Cof.	Р	P Cof. P Cof.		Р	Cof.	Р	Cof.	Р	
0 P	.023	.421	063	.674	.035	.768	075	.125	314	.211	.141	.677
1 P	.036	.220	210	.169	153	.200	040	.406	709	.015	715	.035
2 P	.079	.006	253	253 .097		.341	.008	.876	675	.020	402	.236
3 P	.110	.000	.130	.382	303	.008	.055	.259	074	.783	.525	.140
4 P	.100	.000	.068	.640	016	.888	.069	.158	087	.747	107	.751
5 P	.085	.002	.228	.114	.184	.096	.089	.074	.221	.417	.488	.200
6 P	.044	.120	.099	.447	049	.662	.070	.159	059	.827	304	.368
0+1 P	.007	.822	227	.247	098	.526	060	.240	876	.020	527	.246

Model		LINE	AR RE	GRES	SION			ORE	DINAL	LOGI	STIC	
Model	M tempe on	in. rature lly	Min. temperature with seasonality		M tempe with corre ar seaso	in. ature auto lation nd nality	M tempe or	in. rature lly	M tempe wi seaso	in. erature ith nality	M tempe with corre ar seaso	in. rature auto lation nd nality
	Cof.	Р	Cof.	Cof. P Cof. P		Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	.110	.132	.131	.426	.426 .192 .176		.048	.710	.156	.605	.478	.176
1 P	.123	.010	043	.812	025	.871	.032	.709	392	.245	517	.158
2 P	.100	.007	.134	.489	.093	.572	.049	.470	161	.645	126	.738
3 P	.072	.038	.171	.412	.313	.082	.032	.635	241	.561	.119	.804
4 P	.064	.148	.093	.702	.159	.443	.004	.962	953	.076	610	.278
5 P	002	.180	.104	.725	.336	.190	057	.695	587	.321	046	.945
6 P	192	.019	.018	.952 .021 .93		.932	191	.252	-1.11	.070	-1.02	.123
0+1 P	.153	.017	.089	.694	.159	.415	.054	.638	144	.726	.001	.998

Table 29: October to March (TQ)

9. Rainfall with Malaria in the Whole of Asir

Table 30: Whole Year

			L	R			OL					
Model	Rainfa	ll only	Rainfall with seasonality		Rainfa au corre ar seaso	ll with to lation nd nality	Rainfa	ll only	Rainfa seaso	ll with nality	Rainfa au corre ar seaso	ll with to lation nd nality
	Cof.	Р	Cof.	P Cof. P		Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	057	.280	009	.890	.027	.609	.060	.575	.259	.106	.200	.466
1 P	096	.069	.050	.455	.044	.401	.018	.866	.295	.068	.193	.429
2 P	085	.110	.107	.109	.066 .206		016	.855	.263	.104	.024	.921
3 P	093	.084	.088	.193	.018	.730	.080	.463	.448	.010	.564	.041
4 P	059	.274	.041	.831	045	.398	.055	.612	.304	.065	085	.753
5 P	.021	.697	.000	.998	012	.826	.123	.262	.214	.177	089	.721
6 P	.102	.061	.091	.181	.099 .061		.229	.042	.287	.076	.381	.196
0+1 P	.133	.056	.057	.553	.071	.338	.076	.590	.576	.014	.401	.286

-			L	R			OL						
Model	Rainfa	ll only	Rainfall with seasonality		Rainfa au corre ar seaso	ll with to lation nd nality	Rainfa	ll only	Rainfa seaso	ll with nality	Rainfa au corre ar seaso	ll with to lation nd nality	
	Cof.	Р	Cof.	Р	Cof. P		Cof.	Р	Cof.	Р	Cof.	Р	
0 P	.082	.588	.079	.612	021	.872	.140	.636	.155	.608	.471	.492	
1 P	075	.504	054	.760	017	.907	089	.687	117	.736	.003	.997	
2 P	.196	.099	.440	.023	.338	.039	.069	.771	.052	.894	210	.836	
3 P	.145	.240	.163	.299	017	.400	.423	.118	.599	.073	1.08	.348	
4 P	047	.704	.024	.877	122	.356	.420	.116	.791	.031	-1.07	.354	
5 P	014	.907	042	.777	010	.935	.332	0.174	.471	.125	.001	.890	
6 P	.264	.041	.258	.061	.212	.064	.257	.346	.253	.370	.510	.631	

Table 31: September to November

Table 32: December to March

			L	R			OL						
Model	Rainfa	ll only	Rainfall with seasonality		Rainfa au correl ar seaso	ll with to lation nd nality	Rainfa	ll only	Rainfa seaso	ll with nality	Rainfa au corre ar seaso	ll with to lation nd nality	
	Cof.	Р	Cof.	Р	Cof. P		Cof.	Р	Cof.	Р	Cof.	Р	
0 P	.050	.635	157	.215	033	.759	.081	.718	.124	.650	299	.488	
1 P	130	.337	049	.494	027	.814	.000	.999	.011	.970	248	.567	
2 P	.059	.649	.052	.697	.080	.080 .444		.885	.020	.941	028	.641	
3 P	.083	.549	.073	.603	.038	.736	.430	.181	.408	.219	.572	.204	
4 P	008	.937	.057	.689	025	.830	041	.843	.236	.456	.464	.352	
5 P	106	.277	.007	.965	077 .524		152	.450	085	.780	345	.391	
6 P	027	.792	.041	.770	.139	.219	.378	.085	.548	.088	.689	.177	

10. Malaria with rainfall in TA

Table 33: September	to March

Model			L	R					0	L		
Model	Rainfa	ll only	Rainfall with seasonality		Rainfa au correl ar seaso	ll with to lation nd nality	Rainfa	ll only	Rainfa seaso	ll with nality	Rainfa au corre ar seaso	ll with to lation nd nality
	Cof.	Р	Cof.	Р	Cof. P		Cof.	Р	Cof.	Р	Cof.	Р
0 P	.239	.003	.188	.008	.085	.109	.612	.000	.784	.000	.396	.025
1 P	.236	.005	.226	.002	.083	.148	.486	.001	.524	.002	.378	.060
2 P	.220	.012	.217	.005	.081	.081 .163		.027	.346	.025	.020	.946
3 P	.252	.007	.271	.001	.090	.090 .159		.015	.448	.009	475	.117
4 P	.196	.042	.233	.006	.064	.317	.277	.070	.361	.032	.101	.707
5 P	.038	.693	.149	.082	.036	.564	.122	.402	.220	.173	.333	.252
6 P	.053	.572	.155	.061	.048 .425 .		.323	.031	.480	.007	.380	.165
0+1 P	.314	.001	.259	.002	.109	.097	.793	.000	.897	.000	.315	.041

11. Malaria with rainfall in TQ

Table 34: October to March

Model		LINE	AR RE	GRES	SION			ORE	DINAL	LOGI	STIC	
Model	Rainfa	ll only	Rainfall with seasonality		Rainfa au corre ar seaso	ll with to lation nd nality	Rainfa	ll only	Rainfa seaso	ll with nality	Rainfa au corre ar seaso	ll with to lation nd nality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	.311	.000	.529	.030	.173	.021	.581	.001	.710	.001	.620	.011
1 P	.263	.003	.244	.006	.050	.050 .149		.001	.733	.001	.518	.031
2 P	.286	.001	.253	.004	.100	.251	.969	.000	1.02	.000	.900	.001
3 P	.077	.370	.110	110 .193		.844	.567	.003	.725	.001	.533	.034
4 P	.010	.948	.101	.250	.040	.956	.539	.004	.729	.001	.593	.016
5 P	.062	.467	.102	.242	.046	.545	.705	.001	.809	.001	.740	.005
6 P	.125	.151	.140	.140 .120 .074		.351	.429	.021	.553	.010	.426	.073
0+1 P	.413	.000	.382	.000	.176	.045	1.03	.000	1.22	.000	1.02	.003

			T	R					0	T		
Model	Hum on	idity ly	Humidity with seasonality		Hum with correl ar seaso	idity auto lation nd nality	Hum on	idity ly	Hum wi seaso	idity ith nality	Hum with corre ar seaso	idity auto lation nd nality
	Cof.	Р	Cof.	of. P Cof. P		Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	.003	.856	.018	.549	.011	.669	012	.704	025	.673	.029	.798
1 P	.018	.463	.081	.029	.061	.050	.006	.904	.015	.843	.047	.749
2 P	.043	.082	.101	.019	.044	.282	.027	.590	.040	.645	.025	.889
3 P	.021	.455	.028	.484	.000	.990	.064	.277	.122	.150	256	.276
4 P	005	.862	.012	.726	.001	.966	.077	.193	.134	.070	257	.359
5 P	.017	.410	.042	.167	.031	.221	.077	.089	.173	.018	.424	.115
6 P	.025	.264	.048	.113	.023	.381	.075	.126	.138	.047	.059	.667

12. Malaria with Relative Humidity in Asir Table 35: September to November

Table 36: December to March

			L	R					0	L		
Model	Hum on	idity ly	Hum wi seaso	idity ith nality	Hum with correa ar seaso	iidity auto lation nd nality	Hum or	idity ly	Hum wi seaso	idity th nality	Hum with corre ar seaso	idity auto lation nd nality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof. P		Cof.	Р	Cof.	Р
0 P	.021	.371	.045	.038	.057	.006	.000	.996	.021	.707	.122	.152
1 P	.039	.067	.051	.035	.038	.045	.000	.993	.031	.551	.030	.717
2 P	.020	.090	.034	.152	.012	.541	.002	.935	.090	.096	.146	.080
3 P	.029	.011	.062	.014	.040	.057	.024	.340	.038	.499	.415	.017
4 P	.029	.051	.059	.036	.021	.395	.001	.982	.054	.393	.166	.172
5 P	.007	.744	.044	.238	.015	.633	.003	.939	.067	.410	.060	.615
6 P	024	.301	031	.410	041	.173	.063	.197	.100	.212	077	.526

				Iuo	10 57. 0	501000		uen				
		LINE	AR RE	GRES	SION			ORE	DINAL	LOGI	STIC	
Model	Hum on	idity lly	Hum wi seaso	idity th nality	Hum with corre ar seaso	idity auto lation nd nality	Hum or	idity lly	Hum wi seaso	idity ith nality	Hum with corre ar seaso	idity auto lation nd nality
	Cof.	Р	Cof.	Р	P Cof. P		Cof.	Р	Cof.	Р	Cof.	Р
0 P	019	.139	.015	.504	.022	.261	.003	.383	.036	.402	.059	.239
1 P	015	.161	.050	.036	.038	.067	006	.745	.023	.595	.008	.867
2 P	005	.624	.051	.030	.031	.127	008	.678	.022	.617	.010	.840
3 P	.005	.650	.050	.033	.034	.102	.001	.950	.035	.459	.042	.444
4 P	.015	.308	.039	.115	.011	.621	.037	.211	.066	.208	.049	.404
5 P	.028	.136	.022	.401	.005	.828	.059	.120	.091	.107	.075	.228
6 P	.015	.372	.005	.838	007	.765	003	.934	15	.775	017	.767
0+1 P	020	.116	.048	.094	.043	.075	.003	.906	.051	.344	.050	.405

13. Malaria with Relative Humidity in TQ Table 37: October to March

Multiple regressions

			111	u/MIIIu	in iom	perutu		Tunnu	11			
Whole			L	R					0	L		
year Cur.	Varia on	ables ly	wi seaso	th nality	With correla Seaso	auto tion & nality	Varia on	ables ly	wi seaso	th nality	With correla seaso	auto tion & nality
Month	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	090	.003	042	.631	014	.836	154	.026	798	.001	480	.217
Rainfall	.006	.538	.025	.100	.026	.035	.015	.819	.030	.028	.094	.042
1 prev.			L	R					0	L		
Month	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	064	.031	006	.941	.010	.876	.131	.058	775	.001	330	.341
Rainfall	.021	.040	.050	.001	.037	.002	.011	.647	.083	.039	.073	.034
Aver. Curr.			L	R					0	L		
& 1prev.	Cof. P Cof. P			Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	
Max.T	070	.044	024	.813	.004	.962	145	.074	-1.17	.000	621	.184
Rainfall	.023	.045	.057	.003	.048	.001	.006	.842	.016	.036	.133	.037

Table 38: Multiple regressions of Malaria in All Asir Maximum Temperature with rainfall

Table 39: Multiple regressions of Malaria in All Asir Maximum Temperature with rainfall

						±						
April			L	R					0	L		
To Aug. Cur.	Varia on	ables ly	wi seaso	th nality	With correla Seaso	auto tion & nality	Varia on	ables ly	wi seasor	th nality	With correla Seaso	auto tion & nality
Month	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	171	.000	131	.185	121	.098	220	.119	588	.087	875	.113
Rainfall	.129	.065	.182	.029	.082	.184	.302	.210	.314	.255	.614	.241
1 prev.			L	R					0	L		
Month	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	126	.000	148	.113	081	.238	170	.076	890	.012	774	.097
Rainfall	.101	.096	.114	.135	.078	.164	.435	.049	.396	.148	.448	.272
Aver. Curr.			L	R					0	L		
& 1prev.	Cof.PCof.PCof.				Р	Cof.	Р	Cof.	Р	Cof.	Р	
Max.T	140	.000	152	.152	117	.140	177	.140	-1.13	.018	-1.45	.052
Rainfall	.321	.000	.336	.002	.197	.019	.997	.008	1.01	.026	1.59	.050

			1110	4/111101	n temp	cruture	/ // Itil I	lamai	cy.			
Whole			L	R					0	L		
year Cur.	Varia on	ables ly	wi seaso	th nality	With correla Seaso	auto tion & nality	Varia on	ables ly	wi seaso	th nality	With correla Seaso	auto tion & nality
Month	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	090	.003	042	.631	014	.836154 .026789					480	.217
R.Hum.	.006	006 .538 .025 .100 .026 .0						.819	.030	.028	.094	.042
1 prev.			L	R					0	L		
Month	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	064	.031	.006	.941	.010	.876	131	.058	775	.001	330	.341
R.Hum.	.021	.040	.050	.001	.037	.002	.011	.647	.083	.039	.073	.064
Aver. Curr.			L	R					0	L		
& 1prev.	Cof.	Cof. P Cof. P				Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	070	.044	024	.813	.004	.962	145	.074	-1.17	.000	621	.184
R.Hum.	.023	.085	.057	.003	.048	.001	.006	.842	.016	.036	.133	.037

Table 40: Multiple regressions of Malaria in all Asir Maximum temperature with humidity

Table 41: Multiple regressions of Malaria in all Asir Maximum temperature with humidity

April			L	R					0	L		
To Aug. Cur.	Varia on	ables ly	wi seasoi	th nality	With correla Seaso	auto tion & nality	Varia on	ables ly	wi seaso	th nality	With correla seasor	auto tion & nality
Month	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	188	.001	192	.094	155	.060	181	.280	578	.120	583	.358
R.Hum.	.01	.717	.004	.842	.001	.932	.048	.349	.040	.538	.181	.117
1 prev.			L	R					0	L		
Month	Cof.	Cof. P Cof. P			Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	081	.098	160	.111	084	.25	076	.644	762	.041	510	.354
R.Hum.	.025	.169	018	.350	.014	.332	.126	.059	.131	.082	.142	.250
Aver. Curr.			L	R					0	L		
& 1prev.	Cof. P Cof. P Cof.						Cof.	Р	Cof.	Р	Cof.	Р
Max.T	108	.050	338	.063	160	.086	046	.801	-1.04	.038	999	.233
R.Hum.	.033	.122	.019	.457	.011	.552	.150	.050	.148	.114	.307	.076

Whole			L	R					0	L		
Year 	Varia on	ables ly	wi seasor	th nality	With correla Seaso	auto tion & nality	Varia on	ables ly	wi seasor	th nality	With correla seaso	auto tion & nality
Mon.	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	090	.010	064	.480	025	.726	240	.005	770	.002	550	.179
Rainfall	001	.985	069	.358	033	.570	.259	.059	.103	.584	207	.570
R.Hum.	.007	.581	.029	.068	.027	.030	031	.266	036	.360	.104	.118
1 Prev.			L	R					0	L		
Mon.	Cof.	Cof. P Cof. P			Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	.034	.318	027	.757	.000	.998	180	.029	752	.002	350	.330
Rainfall	096	.086	064	.384	.032	.580	.152	.261	.071	.706	063	.833
R.Hum.	.031	.009	.054	.001	.039	.002	.004	.889	.001	.980	.077	.256
Aver. Cur.			L	R					0	L		
& 1 prev	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	50	.232	059	.582	011	.893	.282	.008	-1.11	.000	673	.186
Rainfall	061	.426	111	.297	.047	.571	.407	.031	.215	.445	179	.707
R.Hum.	028	.059	.063	.002	.050	.001	.031	.374	026	.597	.140	.080

Table 42: Multiple regressions of Malaria in All Asir Maximum Temperature with rainfall and relative humidity

April			L	R					0	L		
To Aug.	Varia or	ables 11y	wi seaso	ith nality	With correla Seaso	auto ation & nality	Vari or	ables 11y	wi seaso	ith nality	With correla Seaso	auto aution & nality
Curr. Mon.	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	195	.000	168	.127	145	.074	195	.248	564	.135	560	.378
Rainfall	.170	.047	.211	.022	.102	.137	.259	.368	.295	.330	.270	.665
R.Hum.	.016	.396	.017	.431	.011	.481	.017	.785	.011	.878	.157	.221
1 Prev.			L	R					0	νL		
Mon.	Cof. P Cof. P Cof.					Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	93	.063	121	.244	.059	.436	.022	.900	900677 .076			.389
Rainfall	.080	.221	.102	.196	.069	.237	.328	.159	.296	.294	.320	.461
R.Hum.	.015	.422	.012	.557	.010	.514	.092	.196	.113	.146	.113	.388
Aver.			L	R					0	٥L		
& 1 prev	LR Cof. P Cof. P Cof.					Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	115	.044	184	.124	135	.132	082	.676	999	.049	-1.06	.216
Rainfall	.342	.002	.366	.003	.214	.023	.865	.044	.865	.075	1.12	.223
R.Hum.	.01	.713	.016	.544	.010	.659	.054	.546	.075	.465	.206	.291

Table 43: Multiple regressions of Malaria in All Asir Maximum Temperature with rainfall and relative humidity

Multiple for TA

			111	u//////		Joraiai	e wittii	Iumu				
Whole			L	R					0	L		
Year Cur.	Varia on	ibles ly	wi seaso	th nality	With correla Seaso	auto tion & nality	Varia on	ables ly	wi seaso	th nality	With correla seaso	auto tion & nality
Month	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	.014	.867	053	.427	.031	.625	074	.193	.086	.695	.212	.686
Rainfall	.257	.000	.101	.021	.128	.002	.518	.000	.553	.000	.497	.097
1 prev.			L	R					0	L		
Month	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	168	.000	.014	.867	053	.422	104	.081	.114	.606	1.07	.197
Rainfall	.255	.000	.257	.000	.101	.021	.595	.000	.645	.000	1.22	.033
Aver. Curr.			L	R					0	L		
& 1prev.	Cof.	Cof. P Cof. P			Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	163	.000	.112	.262	001	.987	112	.095	.222	.468	.797	.303
Rainfall	.365	.000	.341	.000	.168	.001	.827	.000	.875	.000	1.05	.031

Table 44: Multiple regressions of Malaria in TA Maximum temperature with rainfall

Table 45: Multiple regressions of Malaria in TA Maximum temperature with rainfall

April			L	R					0	L		
To Aug. Cur.	Varia on	ables ly	wi seasor	th nality	With correla Seaso	auto tion & nality	Varia on	ables ly	wi seaso	th nality	With correla seaso	auto tion & nality
Month	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	183	.037	096	.430	018	.855	.064	.769	.327	.361	045	.947
Rainfall	.297	.000	.266	.000	.134	.026	.365	.050	.356	.032	.787	.033
1 prev.			L	R					0	L		
Month	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	158	.000	.010	.972	103	.323	.028	.826	.428	.266	.699	.389
Rainfall	.229	.001	.233	.001	.068	.270	.708	.004	.733	.004	1.27	.083
Aver. Curr.			L	R					0	L		
& 1prev.	Cof. P Cof. P				Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	199	.001	078	.610	089	.502	029	.870	543	.230	.443	.600
Rainfall	.351	.000	.342	.000	.156	.038	.759	.005	.781	.006	1.11	.046

Multiple of TQ

			111	u/MIIIu	in iom	perutu		Tunnu	11			
Whole			L	R					0	L		
Year Cur.	Varia on	ables ly	wi seaso	th nality	With correla Seaso	auto tion & nality	Varia on	ables ly	wi seaso	th nality	With correla season	auto tion & nality
Month	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	.005	.828	.068	.478	.077	.325	047	.357	411	.067	321	.247
Rainfall	.375	.000	.397	.000	.251	.000	.636	.000	.759	.000	.735	.001
1 prev.			L	R					0	L		
Month	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	.063	.015	.050	.618	.002	.979	006	.910	341	.123	170	.518
Rainfall	.337	.000	.349	.000	.126	.050	.595	.000	.757	.000	.533	.008
Aver. Curr.			L	R					0	L		
& 1prev.	Cof. P Cof. P			Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	
Max.T	.040	.128	.095	.397	.055	.576	023	.688	569	.048	304	.370
Rainfall	.497	.000	.507	.000	.278	.000	.986	.000	1.22	.000	1.04	.000

Table 46: Multiple regressions of Malaria in TQ Maximum Temperature with rainfall

Table 47: Multiple regressions of Malaria in TQ Maximum Temperature with rainfall

			1,1	an in in a		perata		Ium	11			
April			L	R					0	L		
To Aug. Cur.	Varia on	ables ly	wi seasor	th nality	With correla Seaso	auto tion & nality	Varia on	ables ly	wi seaso	th nality	With correla Seaso	auto tion & nality
Month	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	.118	.113	096	.549	120	.288	.170	.268	366	.337	652	.390
Rainfall	.477	.000	.462	.000	.295	.000	.806	.000	.885	.001	1.16	.038
1 prev.	LR								0	L		
Month	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	.166	.000	125	.380	187	.082	.156	.119	444	.168	387	.421
Rainfall	.519	.000	.452	.000	.200	.018	.742	.001	.848	.001	.834	.048
Aver. Curr.			L	R					0	L		
& 1prev.	Cof. P Cof. P			Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	
Max.T	.193	.001	108	.524	191	.133	.363	.043	196	.627	506	.462
Rainfall	.682	.000	.601	.000	.342	.000	1.34	.000	1.37	.000	1.41	.035

			1110	minui	n temp	oruture	/ WILLIII.	lumui	<i>cy</i>			
Whole			L	R					0	L		
Year Cur.	Varia on	ables ly	wi seasoi	th nality	With correla Seaso	auto tion & nality	Varia on	ables ly	wi seasoi	th nality	With correla Seaso	auto tion & nality
Month	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	81	.042	105	.361	.035	.702	086	.198	570	.009	338	.193
R.Hum.	028	.048	.004	.857	.025	.118	.002	.916	.016	.667	.065	.172
1 prev.									0	L		
Month	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	.053	Cof. P .053 .187		.899	.074	.401	029	.659	371	.081	072	.785
R.Hum.	.007	.615	.039	.050	.038	.016	.006	.803	.045	.234	.052	.263
Aver. Curr.			L	R					0	L		
& 1prev.	Cof.	Cof. P Cof. P			Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	030	.526	069	.610	.076	.476	061	.440	698	.009	320	.317
R.Hum.	017	.354	.031	.210	.047	.016	.008	.785	.051	.283	.086	.138

Table 48: Multiple regressions of Malaria in TQ Maximum temperature with humidity

Table 49: Multiple regressions of Malaria in TQ Maximum temperature with humidity

April			L	R	•				0)L		
To Aug. Cur.	Varia on	ables ly	wi seaso	th nality	With correla Seaso	auto tion & nality	Varia on	ables ly	wi seaso	th nality	With correla seaso	auto tion & nality
Month	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	073	.417	445	.019	310	.019	020	.891	83	.093	-1.33	.038
R.Hum.	033	.139	025	.423	.002	.941	.034	.046	.055	.341	.069	.457
1 prev.			L	R					0	L		
Month	Cof. P Cof.			Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	.157	Cof. P Cof. P .157 .035 339 .054				.087	.161	.192	466	.150	210	.657
R.Hum.	.045	.114	008	.802	.011	.636	.094	.057	.098	.108	.146	.119
Aver. Curr.			L	R					0	L		
& 1prev.	Cof. P Cof. P Cof					Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	.044	.044 .677556 .010361 .0					.220	.194	756	.075	982	.131
Rainfall	.005	.886	027	.502	.007	.803	.126	.048	.128	.098	.176	.130

Whole			L	R					0	L		
Year 	Varia on	ables ly	wi seasor	th nality	With correla Seaso	auto tion & nality	Varia on	ables ly	wi seaso	th nality	With correla seaso	auto tion & nality
Mon.	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	076	.040	.073	.498	.114	.192	088	.256	400	.106	261	.394
Rainfall	.378	.000	.398	.000	.250	.000	.644	.000	.759	.000	.737	.001
R.Hum.	039	.004	.002	.918	.015	.339	021	.479	.005	.917	.026	.648
1 Prev.			L	R					0	L		
Mon.	Cof.	Cof. P Cof. P 064 101 120 210				Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	.064	.101	.139	.210	.093	.319	.022	.776	164	.502	.090	.759
Rainfall	.337	.000	.350	.000	.126	.049	.593	.000	.794	.000	.575	.007
R.Hum.	.000	.982	.038	.064	.039	.024	.014	.630	.088	.073	.118	.044
Aver. Cur.			L	R					0	L		
& 1 prev	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	17	.692	.148	.240	.132	.214	029	.759	441	.160	124	.738
Rainfall	.496	.000	.505	.000	.271	.000	.987	.000	1.23	.000	1.04	.000
R.Hum.	028	.098	.022	.345	.033	.098	003	.932	.065	.312	.086	.247

Table 50: Multiple regressions of Malaria in TQ Maximum Temperature with rainfall and relative humidity

April			L	R					0	L		
To Aug.	Varia on	ables ly	wi seaso	th nality	With correla Seaso	auto tion & nality	Varia on	ables lly	wi seaso	th nality	With correla seaso	auto tion & nality
Curr. Mon.	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	.012	.881	149	.405	121	.337	.193	.272	148	.728	346	.677
Rainfall	.506	.000	.458	.000	.295	.000	.796	.000	.921	.001	1.25	.040
R.Hum.	055	.008	020	.497	.000	.978	.012	.789	.095	.197	.114	.412
1 Prev.			L	R					0	L	•	
Mon.	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	.167	.015	175	.285	193	.119	.277	.073	152	.685	.174	.758
Rainfall	.518	.000	.453	.000	.200	.019	.703	.002	.937	.001	1.29	.044
R.Hum.	.001	.972	019	.526	.002	.927	.064	.288	.133	.077	.266	.077
Aver.			L	R					0	L		
& 1 prev	LR Cof. P Cof. P Cof.					Р	Cof.	Р	Cof.	Р	Cof.	Р
Max.T	.123	.138	159	.407	185	.198	.629	.014	.024	.964	364	.676
Rainfall	.686	.000	.597	.000	.342	.000	1.29	.000	1.47	.000	1.91	.026
R.Hum.	.033	.302	.021	.561	.003	.916	.160	.075	.260	.025	.390	.093

Table 51: Multiple regressions of Malaria in TQ Maximum Temperature with rainfall and relative humidity

Appendix A.3

<u>Malaria Data with The Hottest</u> <u>Temperature days in the Whole of Asir</u>

			L	R					0	L		
Model	Hot tempe on	test rature ly	Hot tempe wi seaso	test rature th nality	Hot tempe with correl ar seaso	test rature auto lation nd nality	Hot tempe on	test rature ly	Hot tempe wi seaso	test erature ith nality	Hot tempe with correl ar seaso	test rature auto lation nd nality
	Cof.	Р	Cof. P C		Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	136	.000	161	.027	117	.041	187	.001	737	.001	381	.236
1 P	142	.000	164	.026	077	.187	207	.001	787	.000	547	.092
2 P	124	.000	165	.026	053	.368	192	.001	831	.000	417	.154
3 P	075	.006	109	.007	092	.122	147	.011	907	.000	618	.055
4 P	007	.812	148	.047	026	.671	072	.202	791	.000	338	.262
5 P	.054	.052	180	.015	089	.135	012	.837	.999	.000	971	.011
6 P	.103	.000	167	.025	058	.331	.035	.533	-1.07	.000	765	.037
0+1P	-2.15	.000	190	.085	130	.100	356	020.	-1.28	.005	880	.158

All Asir Table52: Whole Year

Table 53: April to August

			Ι	R					0	L		
Model	Hot tempe or	ttest erature lly	Ho tempe w seaso	ttest erature ith onality	Ho tempe with corre as seaso	ttest erature auto elation nd pnality	Hot tempe on	test rature ly	Ho tempe w seaso	ttest erature ith nality	Ho tempe with corre au seaso	ttest rature auto lation nd mality
	Cof.	Р	Cof.	Р	Cof.	Cof. P		Р	Cof.	Р	Cof.	Р
0 P	248	.000	156	.045	094	.019	441	.024	914	.019	447	.041
1 P	136	.000	138	.122	103	. 06	258	.028	908	.007	849	.052
2 P	137	.000	153	.061	054	.374	247	.023	-1.18	.001	757	.088
3 P	128	.001	216	.002	104	.052	257	.025	-1.14	.001	641	.128
4 P	149	.003	149	.026	048	.351	477	.007	997	.001	649	.094
5 P	133	.062	148	.037	038	.489	720	.005	-1.01	.001	909	.043
6 P	.117	.088	075	.345	.002	.969	381	.051	-1.22	.001	963	.072
0+1 P	15	.000	22	.008	13	.044	218	.000	-1.20	.000	69	.046

			L	R					0	L		
Model	Hot tempe on	test trature lly	Hot tempe wi seaso	test rature ith nality	Hot tempe with correl ar seaso	test auto lation nd nality	Hot tempe on	test rature lly	Hot tempe wi seaso	test grature ith nality	Hot tempe with correl ar seaso	test rature auto lation id nality
	Cof.	Р	Cof.	Р	Cof. P		Cof.	Р	Cof.	Р	Cof.	Р
0 P nt	046	.551	186	.412	092	.627	088	.561	723	.151	793	.533
1 P	021	.806	279	.369	.007	.980	049	.768	890	.162	-2.35	.341
2 P	206	.297	626	.053	386	.168	348	.380	-1.18	.101	.441	.816
3 P	111	.702	232	.464	.206	.470	958	.118	-1.26	.067	-2.26	.247
4 P	121	.570	167	.496	.036	.864	582	.188	739	.143	1.43	.388
5 P	022	.829	194	.217	092	.539	261	.210	-1.01	.022	-1.96	.248
6 P	073	.327	312	.030	161	.217	245	.118	-1.11	.006	-1.52	.162

Table54: September to November

Table 55: December to March

			L	R					0	L		
Model	Hot tempe on	test rature ly	Hot tempe wi seaso	test rature ith nality	Hot tempe with corre ar seaso	test auto lation nd nality	Hot tempe on	test erature lly	Hot tempe wi seaso	test erature ith nality	Hot tempe with corre ar seaso	test rature auto lation id nality
	Cof.	Р	Cof.	Р	Cof. P		Cof.	Р	Cof.	Р	Cof.	Р
0 P nt	046	.621	157	.148	133	.120	312	.152	628	.034	.322	424
1 P	105	.324	168	.151	068	.477	551	.057	624	.056	202	.627
2 P	121	.178	106	.398	.001	.994	344	.115	350	.244	127	.735
3 P	093	.120	151	.342	143	255	172	.175	445	.209	433	377
4 P	192	.098	128	.554	.029	.870	103	.379	233	.605	.304	.621
5 P	141	.058	347	.186	393	.057	139	.389	965	.096	940	.289
6 P	255	.169	172	.526	102	.637	217	.575	565	.316	.164	.848

Appendix B <u>CL</u>

		Ι	inear re	gressio	n			X	Ordinal	logistic	;	
Model	Ma temper on	ux. rature ly	Ma temper wi seasor	ax. rature th nality	Ma tempe with correla season	ax. rature auto ation& nality	Ma temper on	ax. rature ly	Ma temper wi seasor	ax. rature th nality	Ma tempe with correla season	ax. rature auto ation& nality
	Cof.	Р	Cof. P		Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	060	.002	020	Cof. P 020 .632		.624	190	.041	340	.184	310	.261
1 P	060	.004	030	.546	020	.690	220	.044	400	.118	200	.442
2 P	070	.035	010	.775	010	.828	330	.047	310	.186	210	.380
3 P	030	.405	.010	.882	.020	.626	080	.607	220	.390	.090	.724
4 P	.060	.007	.000	.875	.000	.999	.100	.378	280	.350	330	.264
5 P	.040	.015	040	.536	050	.427	.100	.266	210	.525	320	.343
6 P	.050	.007	030	.715	030	.729	.120	.227	610	.206	420	.404

1. CL with Maximum Temperature in Asir Table 1: Maximum Temperature in Asir (February to May)

Table 2: October to January

		Ι	_inear re	gressio	n				Ordinal	logistic	:	
Model	Ma temper on	ıx. rature ly	Ma temper wi seasor	ax. rature th nality	Ma tempe with correla season	ax. rature auto ation& nality	Ma temper on	ıx. rature ly	Ma temper wi seasor	ux. rature th nality	Ma tempe with correla season	ax. rature auto ation& nality
	Cof.	Р	Cof.	Cof. P		Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	050	.023	100	Cof. P 0 100 .091 -		.220	140	.206	430	.122	300	.309
1 P	030	.141	.000	.957	.050	.500	050	.614	060	.849	.610	.081
2 P	040	.064	170	.106	180	.066	080	.394	920	.082	580	.323
3 P	040	.159	040	.718	100	.330	050	.741	.410	.347	680	.218
4 P	030	.643	.050	.617	.030	.719	.050	.863	.400	.365	290	.533
5 P	.050	.360	.020	.824	030	.672	.190	.522	.250	.520	040	.916
6 P	.030	.226	010	.892	020	.778	.030	.827	040	.897	660	.071

		Ι	Linear re	gressio	n		Ŭ	,	Ordinal	logistic	,	
Model	Ma tempe on	ax. rature ly	Ma tempe wi seasor	ax. rature th nality	Ma tempe with correla season	ax. rature auto ation& nality	Ma tempe on	ax. rature ly	Ma tempe wi seaso	ax. rature th nality	Ma tempe with correla season	ax. rature auto ation& nality
	Cof.	Р	Cof.	Cof. P		Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	060	.006	040	Cof. P 0 040 .514 -		.518	190	.031	260	.307	180	.484
1 P	080	.003	060	.225	050	.337	280	.012	440	.080	340	.174
2 P	050	.189	.000	.978	.010	.814	270	.090	230	.329	140	.550
3 P	.040	.300	.000	.958	.020	.776	.010	.926	240	.335	.010	.962
4 P	.070	.019	.020	.744	040	.485	.150	.178	270	.350	410	.163
5 P	.050	.030	070	.368	070	.335	.130	.198	270	.393	400	.232
6 P	.050	.022	.030	.783	030	.805	.130	.171	770	.115	570	.256

2. CL with Maximum Temperature in the highlands Table 3: Maximum Temperature in the highlands (February to May)

Table 4: October to January

		Ι	Linear re	gression	n				Ordinal	logistic		
Model	Ma tempe on	ax. rature ly	Ma tempe wi seasor	ax. rature th nality	Ma tempe with correla season	ax. rature auto ation& nality	Ma tempe on	ax. rature ly	Ma tempe wi season	ax. rature th nality	Ma tempe with correla season	ax. rature auto ation& nality
	Cof.	Р	Cof.	Cof. P		Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	060	.030	080	Cof. P 080 .246		.858	180	.109	440	.103	060	.835
1 P	030	.141	.050	.577	.010	.236	070	.473	.040	.903	.310	.124
2 P	040	.116	130	.270	170	.116	100	.308	-1.00	.047	-1.4	.024
3 P	030	.438	050	.687	060	.606	040	.769	.330	.517	150	.774
4 P	.010	.894	.080	.462	.080	.445	.050	.867	.410	.380	110	.826
5 P	.060	.184	.020	.866	030	.738	.270	.369	.130	.735	260	.500
6 P	.060	.058	.020	.816	.000	.979	.120	.377	.080	.790	300	.409

3. CL with Maximum Temperature in the lowlands

Table 5: December to February

		Ι	Linear re	gressio	n			-	Ordinal	logistic	2	
Model	Ma temper on	ux. rature ly	Ma tempe wi seasor	ax. rature th nality	Ma tempe with correla season	ax. rature auto ation& nality	Ma temper on	ax. rature ly	Ma temper wi seasor	ux. rature th nality	Ma tempe with correla season	ax. rature auto ation& nality
	Cof.	Р	Cof.	Р	seasonalityCof.		Cof.	Р	Cof.	Р	Cof.	Р
0 P	.060	.319	Cof. P .030 .674		.270	.174	.070	.711	080	.751	.090	.755
1 P	020	.807	.090	.423	.130	.260	170	.424	020	.958	.310	.436
2 P	020	.571	.040	.778	.040	.757	090	.422	.160	.698	.100	.814
3 P	040	.337	010	.952	.000	.988	140	.260	090	.766	.060	.849
4 P	070	.245	050	.564	050	.524	250	.277	140	.647	260	.370
5 P	060	.463	040	.622	030	.698	140	.570	040	.879	.020	.946
6 P	110	.259	080	.419	060	.601	450	.151	370	.329	270	.531

			-	Tab	le 6: F	ebruar	y to Ma	ay				
		Ι	Linear re	gressio	n				Ordinal	logistic	;	
Model	Me temper on	an rature ly	Me tempe wi seasor	an rature th nality	Me tempe with correla season	ean rature auto ation& nality	Me temper on	an rature ly	Me temper wi seasor	an rature th nality	Me tempe with correla season	ean rature auto ation& nality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	060	.002	020	.744	030	.646	220	.034	680	.059	510	.183
1 P	070	.002	040	.544	030	.632	290	.028	930	.018	510	.223
2 P	080	.029	.070	.317	.080	.272	410	.026	640	.081	.000	.999
3 P	.050	.285	.090	.253	.090	.224	260	.228	910	.034	320	.508
4 P	.070	.003	.090	.332	.060	.534	.090	.459	920	.053	860	.090
5 P	.050	.005	.050	.642	.020	.826	.080	.334	-1.50	.011	850	.170
6 P	.050	.004	.060	.585	.060	.588	.110	.286	-1.70	.011	750	.276

4. CL with Mean Temperature in Asir

Table 7: October to January

		Ι	linear re	gressio	n				Ordinal	logistic	;	
Model	Me temper on	an rature ly	Me temper wit seasor	an rature th nality	Me tempe with correla season	an rature auto ation& nality	Me temper on	an rature ly	Me temper wi seasor	an rature th nality	Me tempe with correla season	ean rature auto ation& nality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	050	.037	Cof. P 200 .058		170	.082	170	.158	-1.80	.001	-1.20	.032
1 P	030	.155	.090	.457	.160	.186	070	.438	-1.10	.058	.460	.480
2 P	040	.095	140	.304	230	.079	100	.294	-2.10	.003	-1.70	.024
3 P	050	.140	110	.465	190	.184	160	.332	920	.162	-1.90	.014
4 P	230	.013	250	.057	260	.036	-1.30	.005	-2.30	.001	-2.50	.002
5 P	.010	.628	020	.822	040	.638	360	.245	840	.074	560	.269
6 P	.040	.191	010	.924	020	.861	050	.733	840	.050	-1.00	.053

5. CL with Mean Temperature in the highlands Table 8: February to May

				1 40	10 0.1	coruar	y to 141	uy				
		Ι	Linear re	gressio	n				Ordinal	logistic	2	
Model	Me tempe on	an rature ly	Me temper wi seasor	an rature th nality	Me tempe with correla season	ean rature auto ation& nality	Me temper on	an rature ly	Me temper wi seasor	an rature th nality	Me tempe with correla seaso	ean rature auto ation& nality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	070	.004	060	Cof. P 060 .487		.431	250	.017	640	.078	350	.359
1 P	090	.002	100	.253	080	.344	330	.009	970	.017	690	.104
2 P	060	.149	.120	.185	.150	.091	350	.049	430	.253	.240	.579
3 P	.080	.129	.110	.259	.110	.273	050	.819	750	.088	180	.714
4 P	.080	.004	.090	.433	.020	.867	.180	.141	620	.183	700	.154
5 P	.050	.100	.060	.767	.040	.778	.130	.151	-1.10	.053	590	.321
6 P	.060	.020	.100	.502	.090	.526	.120	.217	-1.30	.042	520	.434

		Ι	linear re	gressio	n				Ordinal	logistic	2	
Model	Me temper on	an rature ly	Me temper wi seasor	an rature th nality	Me tempe with correla season	ean rature auto ation& nality	Me temper on	an rature ly	Me temper wi seasor	an rature th nality	Me tempe with correla season	ean rature auto ation& nality
	Cof.	Р	Cof.	Р	seasonality Cof. P		Cof.	Р	Cof.	Р	Cof.	Р
0 P	070	.027	150	.211	100	.379	220	.070	-1.80	.001	-1.30	.020
1 P	030	.134	.140	.329	.190	.146	090	.323	570	.172	.470	.450
2 P	030	.193	120	.434	250	.081	110	.283	-2.30	.002	-2.50	.001
3 P	030	.453	050	.752	110	.505	140	.383	930	.168	-1.10	.093
4 P	270	.014	330	.027	320	.017	-1.60	.002	-2.70	.000	-2.90	.000
5 P	.100	.178	010	.939	.000	.991	-1.20	.707	770	.112	400	.457
6 P	.070	.045	.050	.620	.030	.751	.040	.793	630	.144	810	.104

 Table 9: October to January

6. CL with Minimum Temperature in all Asir Table 10: February to May

				140			<i>j</i> to 10					
		Ι	linear re	gressio	n				Ordinal	logistic	;	
Model	Mi temper on	n. rature ly	Mi temper wi seasor	n. rature th nality	Mi tempe with correla season	in. rature auto ation& nality	Mi temper on	n. rature ly	Mi temper wi seasor	n. rature th nality	Mi tempe with correla season	in. rature auto ation& nality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	070	.001	Cof. P 090 .243		100	.190	230	.044	680	.096	610	.164
1 P	080	.001	070	.262	070	.294	210	.089	220	.514	.030	.944
2 P	060	.064	.090	.133	.010	.088	210	.196	.030	.934	.610	.082
3 P	.040	.346	.070	.187	.060	.276	160	.446	190	.496	080	.805
4 P	.070	.004	.050	.384	.030	.675	.100	.452	310	.334	350	.309
5 P	.050	.005	.040	.591	.030	.710	.080	.366	810	.051	390	.377
6 P	.050	.004	.070	.457	.080	.383	.120	.225	570	.252	.490	.395

Table 11: October to January

		Ι	Linear re	gressio	n				Ordinal	logistic	2	
Model	Mi tempe on	n. rature ly	Mi temper wi seasor	n. rature th nality	Mi tempe with correla season	in. rature auto ation& nality	Mi tempe on	n. rature ly	Mi temper wi seasor	n. rature th nality	Mi tempe with correla season	in. rature auto ation& nality
	Cof.	Р	Cof.	Р	seasonalityCof.P		Cof.	Р	Cof.	Р	Cof.	Р
0 P	040	.164	.010	.844	020	.729	130	.297	490	.151	350	.330
1 P	030	.223	.110	.614	.110	.187	070	.421	-5.50	.157	260	.547
2 P	030	.154	.060	.614	010	.912	080	.393	950	.084	630	.272
3 P	040	.161	010	.955	070	.598	110	.430	560	.389	-1.80	.014
4 P	120	.108	160	.233	150	.239	550	.094	-1.50	.028	-1.00	.129
5 P	.070	.310	.000	.987	010	.978	240	.428	-1.10	.063	280	.656
6 P	.050	.160	020	.897	.000	.987	020	.871	-1.10	.045	850	.151

		Ι	_inear re	gressio	n		<u> </u>	2	Ordinal	logistic		
Model	Mi tempe on	in. rature ly	Mi tempe wi seasor	in. rature th nality	Mi tempe with correla season	in. rature auto ation& nality	M tempe on	in. rature ly	M tempe wi seaso	in. rature th nality	Mi tempe with correla season	in. rature auto ation& nality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	090	.001	150	Cof. P 150 .120		.120	280	.017	700	.099	420	.349
1 P	100	.001	130	.120	100	.222	270	.260	300	.392	170	.661
2 P	050	.182	.120	.118	.150	.034	210	.194	.100	.769	.660	.058
3 P	.080	.152	.100	.176	.080	.254	.000	.996	120	.686	030	.924
4 P	.090	.004	.060	.412	.020	.789	.190	.137	180	.574	290	.399
5 P	.060	.008	.070	.428	.060	.493	.130	.155	480	.221	030	.941
6 P	.060	.014	.800	.491	.090	.414	.140	.150	500	.329	.090	.394

7. CL with Minimum Temperature in the highlands Table 12: February to May

Table13: October to January

		Ι	linear re	gressio	n				Ordinal	logistic		
Model	Mi tempe on	n. rature ly	Mi tempe wi seasor	n. rature th nality	Mi tempe with correla season	n. rature auto ation& nality	Mi tempe on	in. rature ly	Mi tempe wi seasor	in. rature th nality	Mi tempe with correla season	n. rature auto ation& nality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	060	.091	Cof. P .030 .713		010	.881	190	.165	460	.179	570	.120
1 P	030	.163	.090	.380	.060	.496	090	.345	320	.413	090	.830
2 P	030	.182	.010	.915	090	.490	090	.343	860	.112	-1.00	.079
3 P	040	.308	090	.583	160	.291	110	.468	630	.353	-1.20	.080
4 P	080	.359	210	.172	180	.231	500	.137	-1.60	.019	-1.40	.046
5 P	.150	.049	.070	.640	.080	.507	.020	.946	750	.205	030	.962
6 P	.700	.051	.080	.547	.040	.754	.060	.711	790	.145	830	.153

7. CL with Minimum Temperature in the lowlands

Table 14: December to February

		Ι	linear re	gressio	n			-	Ordinal	logistic	:	
Model	Mi temper on	n. rature ly	Mi temper wi seasor	n. rature th nality	Mi tempe with correla season	in. rature auto ation& nality	Mi temper on	n. rature ly	Mi temper wi seasor	n. rature th nality	Mi tempe with correla season	n. rature auto ation& nality
	Cof.	Р	Cof.	Cof. P 180 265		Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	.180	.122	Cof. P .180 .265		.180	.302	.500	.227	.780	.160	1.20	.057
1 P	070	.461	020	.937	020	.931	320	.300	.340	.648	100	.902
2 P	080	.258	290	.176	290	.185	250	.274	360	.619	530	.483
3 P	130	.030	140	.067	110	.155	360	.102	360	.198	240	.374
4 P	100	.832	.010	.899	.040	.564	060	.681	.050	.820	.230	.363
5 P	080	.034	120	.051	130	.058	250	.680	310	.162	490	.052
6 P	040	.576	020	.854	010	.953	010	.681	050	.871	.080	.820

			I uole I	5. Itu	iniun n	1 1 1011	(1 0010	ur y 10	maj			
		Ι	Linear re	gressio	n				Ordinal	logistic	2	
Model	Rain on	ıfall ly	Rain wi seasor	ıfall th nality	Rain with correla seaso	nfall auto ation& nality	Rain on	ıfall ly	Rain wi seasor	ıfall th nality	Rain with correla season	nfall auto ation& nality
	Cof.	Р	Cof.	Cof. P		Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	060	.069	Cof. P .000 .978		010	.915	170	.294	110	.619	100	.696
1 P	070	.027	020	.663	020	.662	280	.093	210	.344	200	.396
2 P	030	.502	010	.915	.000	.986	060	.778	.030	.891	.000	.997
3 P	.000	.971	030	.559	030	.556	.160	.497	.090	.749	020	.931
4 P	070	.107	060	.177	060	.141	.090	.679	.140	.546	120	.610
5 P	030	.549	010	.794	.010	.838	.240	.342	.320	.221	.410	.126
6 P	.040	.279	.000	.970	.020	.703	.150	.392	.190	.445	.220	.389

8. CL with Rainfall in all Asir

Table 15: Rainfall in Asir (February to May)

Table16: October to January

		L	inear reg	gression	1				Ordinal	logistic	2	
Model	Rai	nfall nly	Rain wi seasoi	ifall ith nality	Rair with correla seasor	nfall auto ation& nality	Rain on	ıfall ly	Rain wi seasoi	ıfall th nality	Rair with correla seasor	nfall auto ntion& nality
	Cof.	Р	Cof. P 030 .582		Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	020	.693	030	.582	040	.408	.020	.942	010	.951	140	.519
1 P	050	.278	060	030 .582 060 .238		.223	130	.546	120	.582	150	.501
2 P	020	.652	.010	.788	.020	.621	040	.792	030	.910	.080	.748
3 P	060	.116	040	.442	030	.541	140	.414	210	.412	.050	.857
4 P	.000	.996	.020	.738	.060	.268	.010	.934	.130	.580	.610	.032
5 P	010	.846	030	.602	030	.587	.130	.508	.140	.557	.080	.740
6 P	040	.406	030	.587	040	.437	.030	.892	.030	.909	150	.530

9. CL with Rainfall in all the highlands

Table 17: Rainfall in the highlands (February to May)

		Ι	Linear re	gressio	n			2				
Model	Rain	ıfall ly	Rain wi season	Rainfall with seasonality		Rainfall with auto correlation& seasonality		ıfall ly	Rain wi seasor	ıfall th nality	Rair with correla season	nfall auto ation& nality
	Cof.	Р	Cof. P 201090		Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	070	.062	.010 .902		010	.898	270	.106	120	.624	110	.635
1 P	050	.214	.010	.010 .848		.844	210	.209	080	.719	050	.812
2 P	040	.419	020	.734	020	.696	090	.664	050	.845	110	.657
3 P	+.02	.763	010	.846	010	.907	.180	.441	.060	.542	.060	.823
4 P	100	.072	040	.126	800	.128	020	.921	.030	.896	200	.406
5 P	040	.575	010	.846	.030	.639	.130	.611	.220	.411	.390	.160
6 P	.060	.154	.000	.948	.020	.709	.250	.158	.190	.462	.260	.315

		Ι	linear re	gressio	n				Ordinal	logistic	:	c	
Model	Rain	ıfall ly	Rainfall with seasonality		Rainfall with auto correlation& seasonality		Rain on	ıfall ly	Rain wi seasor	ıfall th nality	Rair with correla seasor	nfall auto ation& nality	
	Cof.	Р	Cof. P		Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	
0 P	010	.894	020 .757		020	.732	.010	.977	040	.852	080	.715	
1 P	070	.211	090	090 .117		.118	170	.448	190	.406	220	.327	
2 P	060	.175	010	.864	.010	.830	090	.591	.030	.909	.260	.311	
3 P	060	.141	050	.138	040	.476	110	.534	070	.516	070	.777	
4 P	.050	.285	.040	.534	.070	.213	.140	.423	.260	.290	.730	.010	
5 P	010	.901	.010	.917	.000	.939	.050	.781	.150	.537	.020	.941	
6 P	070	.181	040	.469	060	.254	070	.720	170	.756	310	.218	

Table 18: October to January

10. CL with Rainfall in all the lowlands

Table 19: December to February

		Ι	Linear re	gressio	n			į	Ordinal	logistic	;	
Model	Rain	ıfall ly	Rainfall with seasonality		Rainfall with auto correlation& seasonality		Rain on	Rainfall only		ıfall th nality	Rair with correla seasor	nfall auto ation& nality
	Cof.	Р	Cof. P		Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	030	.519	030 .614		030	.599	.520	.005	.520	.004	.230	.347
1 P	030	.656	040	.524	040	.483	.370	.052	.350	.065	090	.710
2 P	.020	.776	.020	.713	.010	.892	.540	.012	.570	.011	.230	.402
3 P	100	.112	180	.076	110	.087	.130	.538	.120	.578	220	.363
4 P	.060	.375	.060	.370	.080	.210	.260	.231	.310	.169	.340	.171
5 P	030	.629	020	.708	040	.501	.170	.409	.170	.397	120	.595
6 P	.060	.385	020	.708	.050	.483	.180	.369	.190	.354	.120	.575

11. CL with relative humidity in all Asir

Table 20: relative humidity in Asir (February to May)

		Ι	_inear re	gressio	n				Ordinal	logistic	ic		
Model	Rela humi on	tive idity ly	Relative humidity with seasonality		Relative humidity with auto correlation& seasonality		Relative humidity only		Rela humi wi seasor	tive idity ith nality	Rela hum with correla seaso	tive idity auto ation& nality	
	Cof.	Р	Cof. P		Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	
0 P	.020	.008	.010 .464		.010	.476	.030	.350	030	.629	.000	.963	
1 P	.020	.043	.000	.875	.000	.970	.020	.565	030	.556	.000	.981	
2 P	.020	.013	.010	.204	.010	.155	.030	.490	010	.890	.060	.231	
3 P	.000	.690	.010	.316	.000	.598	020	.596	030	.526	030	.537	
4 P	010	.071	.010	.431	.000	.699	030	.190	030	.495	040	.424	
5 P	010	.052	.010	.250	.010	.385	020	.115	030	.569	020	.676	
6 P	.000	.895	.010	.179	.010	.190	020	.509	.000	.960	.050	.363	

		Ι	linear re	gressio	n			,	Ordinal	logistic	:	
Model	Rela humi on	Relative humidity only		tive dity th nality	Relative humidity with auto correlation& seasonality		Relative humidity only		Rela humi wi seasor	tive dity th nality	Rela humi with correla season	tive idity auto ation& nality
	Cof.	Р	Cof. P		Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	.010	.065	.010 .340		.010	.609	.010	.760	020	.713	.010	.838
1 P	.010	.089	.010	.464	.010	.661	.000	.930	050	.242	060	.235
2 P	.010	.233	.020	.172	.010	.256	.000	.997	040	.471	.020	.727
3 P	.000	.790	.020	.069	.020	.078	020	.533	040	.508	.060	.405
4 P	.000	.603	.010	.310	.010	.232	010	.888	.010	.870	.090	.129
5 P	010	.492	010	.215	010	.292	.000	.953	010	.806	.010	.792
6 P	010	.089	010	.222	010	.294	.000	.950	.000	.983	.010	.881

Table 21: October to January

12. CL with relative humidity in the highlands

Table 22: relative humidity in the highlands (February to May)

		Ι	linear re	gressio	n				Ordinal	logistic	c		
Model	Rela humi on	tive dity ly	Relative humidity with seasonality		Relative humidity with auto correlation& seasonality		Relative humidity only		Rela humi wi seasor	tive dity th nality	Rela humi with correla season	tive idity auto ation& nality	
	Cof.	Р	Cof. P		Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	
0 P	.020	.022	.010 .561		.010	.645	.040	.241	020	.741	010	.940	
1 P	.020	.036	.000	.786	.000	.817	.030	.376	030	.496	.000	.985	
2 P	.020	.070	.010	.379	.010	.251	.030	.503	010	.777	.060	.244	
3 P	.000	.779	.010	.265	.010	.459	030	.368	020	.654	020	.641	
4 P	010	.082	.010	.307	.010	.516	040	.085	010	.807	010	.767	
5 P	010	.160	.020	.143	.010	.318	030	.143	.000	.995	.020	.635	
6 P	.000	.577	.020	.114	.010	.205	.000	.872	.020	.643	.070	.186	

Table 23: October to January

		Ι	linear re	gressio	n				Ordinal	logistic	2	
Model	Rela humi on	tive dity ly	Relative humidity with seasonality		Relative humidity with auto correlation& seasonality		Relative humidity only		Rela humi wi seasor	tive dity th nality	Rela humi with correla season	tive idity auto ation& nality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	.010	.027	.010 .383		.010	.641	.010	.497	020	.649	020	.655
1 P	.010	.168	.000	.743	.000	.851	.000	.975	050	.315	060	.232
2 P	.000	.650	.000	.783	.000	.956	020	.534	050	.347	030	.576
3 P	.000	.719	.010	.458	.010	.413	030	.494	050	.331	.030	.679
4 P	.020	.073	.020	.122	.020	.103	.020	.644	.030	.563	.090	.124
5 P	010	.220	020	.205	020	.138	030	.411	040	.394	070	.197
6 P	020	.018	020	.163	010	.252	020	.620	.000	.931	.040	.517

<u>Appendix C</u> <u>Schistosomiasis</u>

		I	inear Re	egressio	n	emper	Ordinal Logistic Max.					
Model	Ma tempe on	ax. rature ly	Ma tempe wi seaso	ax. rature th nality	Ma tempe with correl ar seaso	ax. rature auto lation nd nality	Ma tempe on	ax. rature ly	Ma tempe wi season	ax. rature th nality	Ma tempe with correl an season	ax. rature auto ation ad nality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	23	.697	-2.5	.152	24	.167	13	.697	38	.171	37	.189
1 P	1.1	.109	1.9	.256	2.2	.207	.94	.181	.45	.119	.48	.104
2 P	05	.96	47	77	43	.788	15	.81	09	.744	08	.772
3 P	44	.68	45	.796	35	.844	24	.61	.03	.903	.05	.865
4 P	95	.196	-2.1	.291	-2.2	.274	84	.211	27	.376	28	.364
5 P	84	.147	3.4	.074	-3.8	.010	71	.168	.81	.050	.83	.053
6 P	56	.356	-2.3	.540	-1.8	.639	65	.401	.29	.610	.32	.583
7P	51	.563	1.8	.565	1.5	.624	49	.511	.09	.835	.09	.848
8 P	-1.2	.557	.76	.801	.57	.853	-1.0	.598	.12	.815	.11	.822
9 P	1.5	.442	21	.937	39	.887	1.1	.472	17	.675	17	.669
10 P	.86	.326	.64	.772	.83	.711	.68	.291	05	.876	05	.894
11 P	.54	.379	.24	.909	.32	.878	.48	.312	36	.287	36	.290
12 P	.97	.127	3.5	.05	3.5	.05	.90	.127	.56	.06	.04	.849
4+3 P	12	.318	17	.470	-1.7	.479	95	.377	10	.786	01	.788
2+3+4	59	.421	-1.6	.557	-1.5	.572	02	.465	01	.977	01	.988
2+3 4+5	-2.7	.241	-4.7	.178	-4.5	.198	-1.9	.201	-4.5	.433	-4.5	.437

All Asir Region (Schist.) Table 1: Maximum temperature (Feb to May)

		Ι	Linear Regression Max.				Ordinal Logistic					
Model	Ma tempe on	ax. rature ly	Ma tempe wi seasor	ax. rature th nality	Ma tempe with correl ar seaso	ax. rature auto lation nd nality	Ma tempe on	ax. rature lly	Ma tempe wi seasor	ax. rature th nality	Ma tempe with correl ar seaso	ax. rature auto lation nd nality
	Cof.	Р	Cof.	Р	Cof.	P	Cof.	Р	Cof.	Р	Cof.	P
0 P	1.0	.51	1.2	.42	.909	.574	11	.845	05	.864	20	.521
1 P	.65	.654	80	.631	.73	.669	08	.801	09	.805	.13	.720
2 P	1.1	.569	-1.2	.657	-1.1	.697	.06	.863	.04	.948	09	.868
3 P	2.1	.258	2.5	.275	2.5	.274	.47	.291	.49	.283	.46	.323
4 P	89	.787	42	.847	35	.875	04	.921	01	.980	05	.910
5 P	28	.321	-2.9	.136	28	.418	26	.458	20	.632	26	.553
6 P	1.8	.312	1.6	.296	1.6	.316	.41	.289	.41	.201	.40	.213
7P	-1.3	.200	-1.9	.199	-1.9	.216	06	.845	06	.840	11	.725
8 P	.52	.417	.46	.378	.40	.773	.09	.635	.12	.666	.13	.635
9 P	.23	.782	.22	.868	.27	.838	.04	.997	.06	.831	.03	.907
10P	.78	.666	.56	.647	.64	.606	.08	.827	.05	.844	.04	.844
11P	25	.501	80	.550	76	.575	14	.852	09	.726	11	.697
12P	-2.2	.054	-2.6	.089	-2.7	.083	25	.355	31	.341	29	.388
4+3	1.7	.542	1.8	.558	2.0	.537	1.2	.555	1.8	.557	2.0	.537
2+3+4	1.5	.636	1.2	.768	1.5	.723	.58	.511	.55	.521	.44	.623
2+3 4+5	2.0	.489	2.1	.593	1.9	.633	.29	.795	.21	.808	.06	.947

Table 2: Maximum temperature (Oct to Jan)

		Ι	Linear Regression Mean		- <u>p • 1 • 1 • 1 • 1 • 1 • 1 • 1 • 1 • 1 • </u>			Ordinal	Logistic	stic		
Model	Me tempe on	ean rature ly	Me tempe wi seaso	ean rature th nality	Me tempe with correl ar seaso	ean rature auto lation id nality	Me tempe on	ean rature ly	Me tempe wi seasor	ean rature th nality	Me tempe with correl an season	ean rature auto lation id nality
	Cof.	Р	Cof.	Р	Cof.	P	Cof.	Р	Cof.	Р	Cof.	P
0 P	95	.402	-1.5	.529	-1.7	.492	26	.402	-2.8	.429	28	.433
1 P	1.8	.551	1.6	.534	1.7	.502	.15	.524	.13	.765	.14	.735
2 P	-1.5	.524	-1.7	.510	-1.7	.530	.05	.254	.04	.933	.06	.897
3 P	-2.2	.254	-3.1	.275	-2.9	.318	45	.101	40	.350	39	.358
4 P	-5.5	.101	-5.3	.101	-5.3	.100	25	.681	20	.681	20	.670
5P	8.2	.012	8.4	.026	8.25	.032	2.2	.025	1.9	.046	1.9	.046
6 P	-2.1	.450	-3.6	.404	-3.1	.500	07	.885	03	.959	01	.997
7P	1.7	.745	1.6	.721	1.4	.759	45	.715	48	.487	51	.468
8 P	2.2	.458	2.75	.530	2.3	.611	.75	.478	.71	.291	.71	.294
9 P	1.5	.652	1.6	.653	1.2	.744	08	.672	07	.900	09	.873
10 P	33	.956	32	.917	10	.974	35	.478	37	.451	36	.458
11 P	52	.789	42	.883	34	.907	59	.248	53	.253	53	.255
12 P	4.5	.201	4.2	.191	4.1	.117	.20	.020	1.0	.024	1.1	.023
2+4+3 P	-1.9	.658	23	.675	28	.681	.22	.587	.23	.675	.23	.627
7+8+9	4.9	.400	5.2	.433	4.5	.526	.10	.891	.12	.909	.09	.937

All Asir Region (Schist.) Table 3: Mean temperature (Feb to May)

		L	inear Re	egressio	n			(Ordinal	Logistic	ic	
Model	Me tempe on	ean rature ly	Me tempe wi seaso	ean rature th nality	Me tempe with correl ar seaso	ean rature auto lation id nality	Me tempe on	ean rature ly	Me tempe wi seasor	ean rature th nality	Me tempe with correl an season	an rature auto ation id nality
	Cof.	Р	Cof.	Р	Cof.	P	Cof.	Р	Cof.	Р	Cof.	P
0 P	.25	.895	.23	.925	.26	.916	2.0	.435	-2.8	.566	37	.454
1 P	2.7	.665	3.8	.619	3.9	.171	.77	.205	.88	.123	.87	.129
2 P	.95	.624	-1.4	.650	-1.2	.703	.19	.763	.18	.769	.12	.850
3 P	3.4	.158	4.6	.141	4.5	.153	1.3	.058	1.38	.045	1.4	.045
4 P	-1.3	.579	-1.9	.536	-1.8	.568	23	.347	52	.380	56	.348
5 P	1.5	.560	1.4	.576	1.4	.574	.65	.296	.47	.364	.45	.376
6 P	3.1	.148	3.2	.143	3.5	.126	.871	.044	.82	.084	.81	.102
7P	3.9	.118	-2.9	.148	-2.9	.162	19	.719	13	.778	18	.691
8 P	76	.732	63	.740	87	.664	04	.762	07	.854	03	.930
9 P	.83	.604	.85	.659	.83	.671	.13	.628	.18	.662	.17	.680
10 P	.27	.847	.28	.891	.36	.862	17	.837	06	.889	06	.890
11 P	-2.1	.206	-2.5	.264	-2.5	.259	21	.411	38	.402	34	.460
12 P	-4.4	.298	-32	.211	-3.4	1.88	44	.588	52	.506	61	.259
2+4+3 P	1.0	.887	.92	.848	1.2	.809	.52	.477	.66	.497	.58	.553
7+8+9	-2.1	.457	-2.5	.463	-2.7	.443	03	.984	.01	.991	.01	.982

All Asir Region (Schist.) Table 4: Mean temperature (Oct to Jan)
		L	inear Re	gressio	n	empere		00 10	Ordinal	Logistic	;	
Model	M tempe on	in rature ly	M tempe wi seaso	in rature th nality	M tempe with correl ar seaso	in erature auto lation nd nality	M tempe on	in rature ly	M tempe wi seaso	in rature th nality	M tempe with correl an season	in rature auto ation ad nality
	Cof.	Р	Cof.	Р	Cof.	P	Cof.	Р	Cof.	Р	Cof.	P
0 P	85	.582	-1.3	.591	-1.5	.539	31	.432	27	.454	28	.438
1 P	-1.7	.561	-1.1	.580	-1.0	.615	.19	.422	27	.427	26	.449
2 P	-1.7	.324	-1.6	.377	-1.6	.383	.05	.864	.01	.972	.01	.976
3 P	-2.7	.054	-3.3	.066	-3.2	.080	.65	.031	.61	.043	.62	.042
4 P	-3.5	.101	-2.5	.232	-2.4	.252	02	.881	03	.915	03	.920
5 P	-1.2	.512	-1.4	.582	-1.5	.383	42	.325	48	.243	47	.245
6 P	2.3	.480	2.5	.477	3.0	.398	.07	.725	.22	.716	.26	.674
7 P	.71	.745	67	.874	-1.1	.797	.35	.645	.36	.585	37	.573
8 P	-1.0	.858	09	.981	26	.949	.05	.868	.01	.984	.01	.987
9 P	-3.5	.462	-3.4	.364	-3.5	.365	.08	.172	.76	.185	.77	.179
10 P	-3.4	.256	-4.2	.241	-3.9	.299	.35	.168	.83	.121	.89	.114
11 P	52	.729	33	.917	22	.945	19	.801	12	.813	11	.825
12 P	.44	.781	.612	.807	.56	.826	.50	.120	.57	.163	.57	.161
2+3 P	3.8	.058	4.5	.061	5.9	.071	.52	.186	.56	.144	.57	.139
5+6	.53	.800	.04	.991	4.4	.084	.30	.491	.42	.493	.40	.510

All Asir Region (Schist.) Table 5: Minimum temperature (Feb to May)

		Ι	inear Re	egressio	n	tempe		(00110	Ordinal	Logistic	;	
Model	M tempe on	in rature ly	M tempe wi seaso	in rature th nality	M tempe with correl ar seaso	in rature auto lation id nality	M tempe on	fin erature lly	M tempe wi seaso	in rature th nality	M tempe with correl ar season	in rature auto lation id nality
	Cof.	Р	Cof.	Р	Cof.	P	Cof.	Р	Cof.	Р	Cof.	P
0 P	2.0	.164	-2.2	.165	-1.9	.250	39	.124	43	.180	32	.322
1 P	4.8	.025	4.3	.018	4.4	.021	.81	.021	.80	.046	.87	.035
2 P	11	.951	10	.970	04	.999	.09	.842	.08	.878	.07	.886
3 P	3.4	.308	3.1	.307	3.0	.330	.31	.541	.32	.587	.36	.544
4 P	-4.5	.107	-4.8	.101	-4.8	.104	1.7	.034	1.2	.044	1.2	.046
5 P	.55	.854	.59	.833	.46	.872	06	.884	05	.928	01	.985
6 P	6.4	.028	5.9	.021	6.2	.019	.90	.051	1.1	.061	1.0	.071
7P	-3.3	.157	-3.1	.187	-3.0	.202	57	.277	54	.266	56	.249
8 P	14	.924	11	.951	27	.890	04	.904	03	.945	.01	.971
9 P	.57	.625	.66	.674	.56	.681	08	.754	07	.832	08	.793
10P	.08	.921	.09	.948	.12	.935	.04	.468	.01	.468	.02	.951
11P	-1.4	.249	-1.8	.204	-1.8	.196	36	.270	35	.234	33	.257
12P	49	.754	51	.756	62	.713	.37	.210	.34	.289	.30	.354
2+3 P	2.5	.514	2.3	.544	2.3	549	.39	.647	.35	.642	.37	.621
5+6	5.2	.080	6.0	.093	6.1	.097	.88	.218	.92	.222	.91	.227

All Asir Region (Schist.) Table 6: Minimum temperature (Oct to Jan)

		L	inear Re	egressio	n			(Ordinal	Logistic	;	
Model	Rair on	nfall ly	Rain wi season	nfall th nality	Rain with correl ar seaso	nfall auto lation nd nality	Rair on	nfall ly	Rair wi seasor	nfall th nality	Rainfa au correl ar seasor	ll with to ation d nality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
0 P	1.3	.611	54	.603	55	.602	.270	.288	.044	.814	.06	.772
1 P	-2.6	.351	-1.2	.238	-1.3	.235	.481	.028	30	.149	30	.160
2 P	1.2	.485	66	.540	77	.489	172	.777	17	.415	15	.506
3 P	1.8	.258	-1.5	.204	-1.6	.171	203	.217	42	.081	42	.098
4 P	2.5	.379	.33	.776	.25	.838	.115	.837	09	.701	07	.754
5 P	-2.1	.261	.29	.803	.26	.823	.548	.046	.06	.791	.09	.704
6 P	-2.7	.875	-1.1	.369	-1.1	.357	.215	.393	25	.320	26	.301
7P	4.6	.009	08	.941	14	.899	.466	.117	07	.721	04	.839
8 P	-1.0	.532	30	.816	26	.844	.537	.038	15	.559	17	.510
9 P	-4.9	.157	.71	.560	.70	.567	094	.878	04	.885	01	.960
10 P	2.0	.489	12	.922	08	.949	229	.601	12	.605	12	.600
11 P	1.2	.441	01	.990	04	.971	.242	.213	06	.790	05	.834
12 P	-1.7	.187	88	.410	88	.418	.615	.050	29	.166	30	.158
5+6+7 P	2.6	.055	65	.716	71	.691	.820	.30	-1.7	.611	-1.4	.694
6+7	4.6	.155	.95	.528	-1.0	.511	.415	.114	.24	.404	22	.452
6+7+8	-1.4	.694	-1.1	.541	-1.1	.540	.019	.811	.34	.343	32	.366

All Asir Region (Schist.) Table 7: Rainfall (Oct to Jan)

r		T	·	· 0. ICC	iutive .	ITuilliu			$\frac{uy}{0}$	T · .·		
Model	Hum on	idity ly	Hum Hum season	idity th nality	n Hum with correl ar seaso	idity auto lation nd nality	Hum on	idity ly	Hum Wi seaso	idity th nality	Hum with correl ar seaso	idity auto lation nd nality
	Cof.	Р	Cof.	Р	Cof.	P	Cof.	Р	Cof.	Р	Cof.	P
Cur. Mon.	.055	.559	.18	.561	.15	.613	.021	.254	.034	.468	.03	.504
1 P	.021	.248	25	.434	25	.424	.025	.321	06	.308	06	.310
2 P	.045	.853	.09	.760	.07	.814	.365	.651	.04	.305	.04	.317
3 P	.078	.799	24	.387	26	.354	.024	.294	05	.237	06	.190
4 P	.032	.768	.03	.923	.04	.893	.254	.214	.01	.856	.01	.853
5 P	.213	.391	.23	.478	.20	.547	.014	.365	.04	.476	.03	.479
6 P	032	.382	.71	.066	.69	.081	.025	.065	.13	.06	.13	.06
7P	049	.562	12	.777	.12	.779	.026	.451	.05	.514	.05	.498
8 P	.211	.838	.36	.343	32	.425	.078	.328	05	.375	05	.379
9 P	064	.145	01	.967	.02	.958	.078	.426	.02	.639	.03	.616
10P	049	.566	.26	.411	17	.600	.065	.326	.05	.307	.05	.310
11P	065	.361	.15	.622	.13	.671	.062	.348	.06	.214	.06	.216
12P	.021	.780	17	.600	17	.600	.034	.512	05	.330	05	.317
4+5 P	.016	.898	.18	.622	.17	.649	.031	.254	.03	.601	.03	.601
6+7	071	.361	.49	.324	.46	.358	.054	.210	.14	.110	.14	.109

All Asir Region (Schist.) Table 8: Relative Humidity (Feb to May)

			- 1 u 0 1	<u> </u>	ciutive	Tunn	uity (5t	in to b	<u>cp)</u>	• • •		
		L	inear Re	egressio	n				Ordinal	Logistic	2	
Model	Hum on	idity ly	Hum wi seasor	idity th nality	Hum with correl ar seaso	idity auto lation 1d nality	Hum on	idity ly	Hum wi seaso	iidity ith nality	Hum with correl ar season	idity auto ation nd nality
	Cof.	Р	Cof.	Р	Cof.	P	Cof.	Р	Cof.	Р	Cof.	P
0 P	.25	.521	.13	.629	.27	.253	.03	.858	.02	.710	.07	.266
1 P	26	.274	.29	.188	.21	.294	.02	.476	.02	.716	01	.918
2 P	.15	.412	01	.984	08	.692	01	.254	04	.459	06	.254
3 P	.21	.652	.08	.707	.12	.520	02	.365	04	.379	04	.400
4 P	25	.357	25	.240	21	.292	05	.214	09	.063	.09	.069
5 P	.12	.164	.37	.099	.21	.331	.02	.061	12	.070	10	.060
6 P	.32	.254	28	.158	24	.196	15	.088	09	.084	07	.118
7P	62	.541	15	.443	17	.348	07	.074	08	.091	07	.145
8 P	.23	.325	.19	.350	20	.290	05	.365	04	.383	03	.549
9 P	05	.854	02	.921	.06	.778	02	.922	0.0	.969	.014	.764
10P	.28	.655	.46	.095	.39	.118	06	.124	09	.114	10	.092
11P	09	.985	03	.932	.04	.903	.02	.854	.01	.911	.01	.915
12P	.45	.021	.54	.041	.50	.039	.11	.054	.13	.041	.13	.043
4+5	38	.050	57	.059	40	.160	22	.094	19	.117	17	.114
6+7	35	.099	36	.167	33	.157	13	.214	15	.136	.13	.143

All Asir Region (Schist.) Table 9: Relative Humidity (Jun to Sep)

		T	incor D		n	, IIaiii		201 10 3	Ordinal	Logistic		
Model	Hum on	idity ly	Hum wi season	idity th nality	Hum with correl ar seaso	idity auto lation nd nality	Hum on	idity ly	Hum wi seaso	idity th nality	Hum with correl ar season	idity auto ation nd nality
0 P	12	.239	21	.317	-1.9	.387	.02	.653	.03	.559	02	.685
1 P	.51	.813	.41	.081	.41	.091	.03	.267	.06	.248	.07	.170
2 P	.26	.627	.24	.414	.23	.420	.02	.458	.01	.835	.02	.772
3 P	.12	.456	13	.677	12	.703	.05	.654	.01	.819	.02	.741
4 P	.26	.962	.02	.947	.02	.958	.06	.865	.02	.774	.02	.757
5 P	.45	.862	.06	.803	.06	.808	.02	.564	.00	.431	.00	.954
6 P	.26	.753	09	.693	10	.674	01	.231	04	.394	04	.433
7P	01	.851	05	.831	05	.825	05	.482	05	.333	.04	.361
8 P	.02	.963	.01	.962	.01	.979	.01	.824	0.0	.948	.01	.925
9 P	03	.852	02	.928	02	.934	08	.299	07	.192	07	.203
10P	.06	.587	.18	.416	.18	.419	06	.745	03	.518	03	.554
11P	36	.395	22	.295	25	.225	04	.365	05	.241	05	.302
12P	.24	.478	.22	.323	.21	.345	.02	.867	.01	.800	.02	.705
4+5 P	.07	.854	.06	.842	.06	.850	.02	.862	.01	.918	.01	.893
6+7	24	.564	11	.703	12	.668	08	.452	07	.241	07	.273

All Asir Region (Schist.) Table 10: Relative Humidity (Oct to Jan)

Sabt-Alalaya Sector

		Li	near Re	egressi	on	emper		0	rdinal	Logist	ic	
Model	Ma tempe on	ax. rature ly	Ma tempe wi seaso	ax. rature th nality	Ma tempe with correl ar seaso	ax. rature auto lation nd nality	Ma tempe on	ax. rature ly	Ma tempe wi seaso	ax. rature th nality	Ma tempe with correl ar seaso	ax. rature auto ation d nality
	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р	Cof.	Р
Cur. Mon.	-0.10	.888	.134	.625	.183	.417	053	.697	729	.192	.391	.715
1 P	.010	.906	192	.612	279	.366	.034	.830	-1.08	.165	435	.159
2 P	134	.477	397	.145	262	.253	096	.794	380	.485	499	.791
3 P	288	.213	215	.305	.113	.617	546	.259	527	.290	432	.652
4 P	-021	.889	135	.430	.003	.984	262	.373	563	.157	.324	.755
5 P	013	.861	195	.256	.085	.566	099	.501	627	.090	-1.08	.341
6 P	036	.571	259	.091	103	.451	182	.167	-1.28	.003	963	.261
7P	.035	.700	094	.447	086	.380	327	.107	724	.021	013	.976
8 P	033	.724	037	.785	007	.945	394	.064	442	.144	.035	.894
9 P	087	.164	070	.640	084	.476	165	.219	259	.429	.035	.939
10P	071	.153	064	.718	.060	.671	138	.191	674	.110	1.24	.058
11 P	096	.058	469	.067	349	.090	113	.295	963	.124	629	.498
12P	119	.100	112	.642	.059	.764	074	.627	.287	.560	.143	.841

Sabt-Alalaya Group Table 11: Maximum temperature (Nov to April)

		Li	near \mathbf{R}_{i}	oressi	on	temper		0	rdinal	I ogist	ic	
Model	Ma tempe on	ax. rature ly	Ma tempe wi seaso	ax. rature th nality	Ma tempe with correl ar season	ax. rature auto lation nd nality	Ma tempe on	ax. rature ly	Ma tempe wi season	ax. rature th nality	Ma tempe with correl ar season	ax. rature auto lation nd nality
	Cof.	Р	Cof.	Р	Cof.	P	Cof.	Р	Cof.	Р	Cof.	P
0 P	01	.896	.198	.596	.199	.520	075	.571	-2.71	.007	995	.606
1 P	027	.781	837	.050	952	.107	087	.645	-3.06	.008	-2.08	.112
2 P	582	.045	-1.01	.008	564	.116	-1.75	.021	-3.63	.004	-2.91	.279
3 P	466	.120	588	.106	.020	.953	-2.18	.006	-3.36	.004	-1.19	.560
4 P	013	.935	245	.335	049	.822	519	.126	-1.80	.012	694	.769
5 P	014	.866	273	.241	110	.580	154	.359	-1.38	.014	-2.92	.246
6 P	040	.612	364	.096	149	.418	218	.186	-1.83	.003	-1.02	.342
7P	.093	.358	183	311	071	.677	.240	.264	-2.01	.003	-1.40	.091
8 P	.009	.940	024	.918	.140	.452	561	.046	-1.18	.033	.644	.424
9 P	087	.192	127	.631	177	.393	248	.082	-1.88	.008	-1.18	.192
10 P	066	.167	.027	.931	.253	.306	131	.191	-2.21	.008	-1.74	.102
11 P	102	.050	634	.050	530	.141	-1.47	.808	-3.99	.081	-4.07	.114
12 P	146	.085	372	.300	.011	.970	189	.314	-2.70	.006	981	.445

Sabt-Alalaya (Schist.) Table 12: Maximum temperature (May to Oct)

Minimum Temperature

			Tuble	15.10		perutu				. .		
		Li	near Re	egressi	on			C	rdinal	Logist	ic	
Model	M tempe on	in rature ly	M tempe wi seasor	in rature th nality	M tempe with correl ar seaso	in rature auto lation nd nality	M tempe on	in rature ly	M tempe wi season	in rature th nality	M tempe with correl an season	in rature auto ation id nality
	Cof.	Р	Cof.	Р	Cof.	P	Cof.	Р	Cof.	Р	Cof.	P
Cur.	.003	.856	.018	.549	.011	.669	012	.704	025	.673	.029	.798
1 P	.018	.463	.081	.029	.061	.050	.006	.904	.015	.843	.047	.749
2 P	.043	.082	.101	.019	.044	.282	.027	.590	.040	.645	.025	.889
3 P	.021	.455	.028	.484	.000	.990	.064	.277	.122	.150	256	.276
4 P	005	.862	.012	.726	.001	.966	.077	.193	.134	.070	257	359
5P	.017	.410	.042	.167	.031	.221	.077	.089	.173	.018	.424	.115
6 P	.025	.264	.048	.113	.023	.381	.075	.126	.138	.047	.059	.667
7P	.050	.635	157	.215	033	.759	.081	.718	.124	.650	299	.488
8 P	130	.337	049	.494	027	.814	.000	.999	.011	.970	248	.567
9P	.059	.649	.052	.697	.180	.464	.039	.885	.020	.941	028	.641
10 P	.083	.549	.073	.603	.038	.736	.430	.181	.408	.219	.572	.204
11 P	008	.937	.057	.689	025	.830	041	.843	.236	.456	.464	.352
12 P	106	.277	.007	.965	077	.524	152	.450	085	.780	345	391

Sabt-Alalaya (Schist.) Table 13: Min temperature (Nov to April)

		Li	near R	egress	ion	<u>-p • 1 • • • • • • • • • • • • • • • • • </u>		0	rdinal	Logist	ic	
Model	M tempe on	in rature ly	M tempe wi seasor	in rature th nality	M tempe with corre an seaso	lin erature auto lation nd nality	M tempe on	in rature ly	M tempe wi seasor	in rature th nality	M tempe with correl ar seaso	in rature auto ation nd nality
	Cof.	Р	Cof.	Р	Cof.	P	Cof.	Р	Cof.	Р	Cof.	P
0 P	046	.621	157	.148	133	.120	312	.152	628	.034	.322	424
1 P	105	.324	168	.151	068	.477	551	.057	624	.056	202	.627
2 P	121	.178	106	.398	.001	.994	344	.115	350	.244	127	.735
3 P	093	.120	151	.342	143	.255	172	.175	445	.209	433	.377
4 P	192	.098	128	.554	.029	.870	103	.379	233	.605	.304	.621
5 P	141	.058	347	.186	193	.357	139	.389	965	.096	940	.289
6 P	255	.169	172	.526	.102	.637	217	.575	565	.316	.164	.848
7P	137	.002	.117	.264	.063	.402	.091	.224	112	.611	.209	.732
8 P	.167	.000	.053	.611	033	.656	109	.088	057	.789	.351	.550
9 P	171	.000	015	.899	069	.414	129	.034	105	.658	418	.577
10 P	.177	.000	.058	.642	.003	.976	.105	.108	041	.868	.048	.930
11 P	244	.000	011	.928	016	.863	139	.133	235	.383	351	.632
12P	072	.405	.070	.556	.070	.408	133	.349	160	.520	092	.885

Sabt-Alalaya (Schist.) Table 14: Min temperature (May to Oct)

Rainfall

		Li	near Re	egressi	on			<u>C</u>	rdinal	Logist	ic	
Model	Rair on	nfall ly	Rair wi seasor	nfall th nality	Rair with correl ar season	nfall auto lation nd nality	Rair on	nfall ly	Rair wi seasor	nfall th nality	Rainfa au correl an season	ll with to ation d nality
	Cof.	Р	Cof.	Р	Cof.	P	Cof.	Р	Cof.	Р	Cof.	P
Cur.	.124	.081	.100	.630	.288	.195	054	.670	788	.555	230	.620
1P	.121	.006	.018	.940	.187	.366	.024	.760	-1.05	.229	690	178
2P	.087	.016	194	.459	123	.584	.030	.640	-1.17	.627	896	.102
3P	.062	.085	234	.364	.123	.600	.018	.795	-1.27	.424	705	.251
4P	.056	.226	264	.305	37	.874	016	.857	-1.80	.305	-1.28	.261
5 P	038	.626	186	.404	.156	.445	152	.335	-1.01	.047	562	.304
6 P	198	.008	247	.219	086	.625	212	.179	-1.11	.018	915	.072
7P	.148	.011	.100	.721	.390	.311	.007	.946	-1.47	.210	782	.229
8 P	.096	.844	317	.835	016	.896	.026	.573	-1.25	.214	554	.139
9 P	.088	.301	291	.047	109	.363	.039	.406	-1.21	.385	731	.348
10 P	.065	.215	202	.171	019	.869	.051	.281	921	.383	378	.313
11 P	.025	.365	284	.153	157	.169	.036	.449	-1.25	.199	-1.09	.227
12P	019	.492	209	.178	089	.467	094	.248	-1.28	.425	797	.330

Sabt-Alalaya (Schist.) Table 15: Rainfall (Nov to April)

		Liı	near Re	gressi	on			C	rdinal	Logist	ic	
Model	Rainfa	ll only	Rair wi seasor	nfall th nality	Rain with correl ar seaso	nfall auto lation nd nality	Rair on	ıfall ly	Rair wi seasor	nfall th nality	Rainfa au correl an season	ll with to ation d nality
	Cof.	Р	Cof.	Р	Cof.	P	Cof.	Р	Cof.	Р	Cof.	P
0 P	104	.785	.027	.817	.068	.448	133	.154	762	.110	535	.219
1 P	113	.785	06	.956	.004	.962	155	.622	985	.602	882	.597
2 P	099	.624	075	.522	060	.513	138	.206	-1.0	.202	641	.190
3 P	045	.259	.149	.206	.215	.180	070	.152	313	.266	1.33	.073
4 P	.016	.516	.126	.286	.018	.844	.000	.990	272	.324	047	.919
5 P	.079	.2871	.156	.189	.074	.428	.066	.179	316	.253	272	.532
6P	.120	.658	.041	.731	058	.533	.119	.119	151	.585	.177	.697
7P	022	.758	116	.697	066	.790	06	.668	-1.06	.099	519	.662
8 P	015	.869	356	.462	375	.348	019	.913	-1.02	.298	469	.267
9 P	105	.590	692	.137	236	.576	234	.541	-1.76	.083	. 178	.450
10P	040	.913	178	.655	.300	.389	190	.130	-2.48	.114	118	.587
11 P	.112	.506	.175	.629	.358	.232	329	.329	-1.98	.227	321	.518
12P	.039	.707	.160	.465	.132	.666	073	.719	-1.03	.176	211	.386

Sabt-Alalaya (Schist.) Table 16: Rainfall (May to Oct)