

BMJ Open A cohort study of influences, health outcomes and costs of patients' health-seeking behaviour for minor ailments from primary and emergency care settings

M C Watson,¹ J Ferguson,² G R Barton,³ V Maskrey,⁴ A Blyth,⁴ V Paudyal,¹ C M Bond,¹ R Holland,⁴ T Porteous,¹ T H Sach,³ D Wright,⁵ S Fielding¹

To cite: Watson MC, Ferguson J, Barton GR, *et al.* A cohort study of influences, health outcomes and costs of patients' health-seeking behaviour for minor ailments from primary and emergency care settings. *BMJ Open* 2015;**5**:e006261. doi:10.1136/bmjopen-2014-006261

► Prepublication history and additional material is available. To view please visit the journal (<http://dx.doi.org/10.1136/bmjopen-2014-006261>).

Received 17 September 2014
Revised 22 December 2014
Accepted 30 December 2014



CrossMark

For numbered affiliations see end of article.

Correspondence to

Dr M C Watson;
m.c.watson@abdn.ac.uk

ABSTRACT

Objectives: To compare health-related and cost-related outcomes of consultations for symptoms suggestive of minor ailments in emergency departments (EDs), general practices and community pharmacies.

Design: Observational study; prospective cohort design.

Setting: EDs (n=2), general practices (n=6) and community pharmacies (n=10) in a mix of rural/urban and deprived/affluent areas across North East Scotland and East Anglia. Participants Adults (≥18 years) presenting between 09:00 and 18:00 (Monday–Friday) in general practices and 09:00–18:00 (Monday–Saturday) in pharmacies and EDs with ≥1 of the following: musculoskeletal pain; eye discomfort; gastrointestinal disturbance; or upper respiratory tract-related symptoms.

Interventions: Participants completed three questionnaires: baseline (prior to index consultation); satisfaction with index consultation and follow-up (2 weeks after index consultation).

Main outcome measures: Symptom resolution, quality of life, costs, satisfaction and influences on care-seeking behaviour.

Results: 377 patients participated, recruited from EDs (81), general practices (162) and community pharmacies (134). The 2-week response rate was 70% (264/377). Symptom resolution was similar across all three settings: ED (37.3%), general practice (35.7%) and pharmacy (44.3%). Mean overall costs per consultation were significantly lower for pharmacy (£29.30 (95% CI £21.60 to £37.00)) compared with general practice (£82.34 (95% CI £63.10 to £101.58)) and ED (£147.09 (95% CI £125.32 to £168.85)). Satisfaction varied across settings and by measure used. Compared with pharmacy and general practice use, ED use was significantly ($p<0.001$) associated with first episode and short duration of symptom(s), as well as higher levels of perceived seriousness and urgency for seeking care. Convenience of location was the most common reason for choice of consultation setting.

Conclusions: These results suggest similar health-related outcomes and substantially lower costs with

Strengths and limitations of this study

- This is the first study to compare health-related and cost-related outcomes across the three main healthcare settings where patients present with symptoms suggestive of minor ailments.
- Prospective data collection using the same processes and tools across all three settings reduces the risk of bias.
- While the inclusion of different types of pharmacies and a variety of locations of pharmacies and general practices increased the generalisability of the results, more rural communities were under-represented.
- A cohort study design was used rather than a randomised controlled trial because the latter was an impractical option to address the objectives of this study.

pharmacy consultations for minor ailments. Effective strategies are now needed to shift demand for minor ailment management away from EDs and general practices to the community pharmacy setting.

INTRODUCTION

Emergency departments (EDs) are reported to be in crisis¹ with 22 million attendances in EDs in England between 2012 and 2013, one third of which involved 'guidance or advice only'.² This is placing 'intense, growing and unsustainable pressure' on the service.³ In general practice, 340 million general practice consultations occurred between 2012 and 2013.³ This equates to an estimated 5.5 consultations per patient annually; 40% higher than 1995.⁴ Recent estimates suggest that 5% and 13% of consultations in ED and general practice, respectively, are for minor ailments that could be managed in community pharmacies.⁵

For the purpose of this study, minor ailments are “common or self-limiting or uncomplicated conditions which may be diagnosed and managed without medical (ie, doctor) intervention.”⁶

The above illustrates that health seeking for the management of minor ailments from high-cost settings such as EDs and general practice persists despite a wide range of alternative sources of care being widely available. Effective over-the-counter (OTC) medicines can be supplied from community pharmacies to support the management of minor ailments.⁷ Over the past 30 years, many medicines have been reclassified from ‘prescription only’ status to OTC medicines and this has increased the options available for the treatment of minor ailments. Local and national pharmacy-based minor ailment schemes have also been introduced to reduce demand on high-cost settings, with evidence from a recent systematic review suggesting that they appear to be effective.⁸ Pharmacy-based minor ailment schemes provide public access to NHS treatment and/or advice via a pharmacist or pharmacy personnel, or, where appropriate, to onward referral to other health professionals. These schemes were originally proposed by the UK health departments as part of their long-term strategy to encourage patient self-care and utilisation of pharmacies as the first port of call for minor ailments where professional support was required. However, the evidence to date is limited in terms of study quality and outcomes reported, that is, generally surrogate health outcomes have been evaluated, such as reconsultation rates. Furthermore, no formal economic evaluations have been reported.

There is an urgent need to shift demand for minor ailment management from high-cost settings to potentially cheaper alternatives, for example, community pharmacies. Reassurance, however, is needed that similar outcomes can be achieved with community pharmacy management compared with these other settings. The purpose of this study was to compare health-related and cost-related outcomes associated with the management of symptoms suggestive of minor ailments (hereafter referred to as minor ailments) in three healthcare settings: ED, general practice and community pharmacy. The following objectives were assessed:

1. Whether health-related and cost-related outcomes differ between settings.
2. Whether satisfaction with index consultation is associated with health-related outcomes.
3. What factors (triggers) influence patients’ choice of care setting.

METHODS

Study design

A prospective cohort study was conducted.

Study setting and recruitment

The study was conducted across two geographic regions (East Anglia, England and Grampian, Scotland).

Potential participating sites were identified using NHS lists of health service providers. General practices and community pharmacies were eligible if they were within a 25-mile radius of the main city in East Anglia (Norwich) and Grampian (Aberdeen). This was to ensure that participants could realistically have chosen from all three settings of interest. In Grampian, a letter of invitation was sent to all eligible practices and pharmacies. In East Anglia, the same approach was used for pharmacies. From those expressing an interest (online supplementary appendix 1), purposive sampling was used to generate a maximum variation sample to include different types and locations of pharmacies and different locations of general practices. Three general practices were purposively selected in East Anglia to achieve maximum variation.

Participant recruitment

Participant recruitment was undertaken by six researchers between July 2012 and April 2013. The same process for data collection was used across all settings. Participants were eligible for inclusion if they: were aged ≥ 18 years; requested treatment for one or more of the four included minor ailments or presented symptoms associated with these ailments; requested medicines for the treatment of one of the four minor ailments (pharmacy only); presented between 09:00 and 18:00, Monday–Saturday (ED and pharmacy) and between 09:00 and 18:00, Monday–Friday (general practice); had a face-to-face consultation (ie, interaction with member of staff). This consultation is referred to hereafter as the ‘index consultation’. These recruitment periods were selected to reflect the traditional opening hours of community pharmacies (ie, to reflect periods when patients had the option of seeking care from community pharmacies if they had chosen to do so). Patients in pharmacies were excluded if they were collecting prescription items for target minor ailments or purchasing treatment for the future management of those ailments. Patients in any setting were excluded if they did not adequately understand verbal/written information in English. The same recruitment method was used across all sites and settings. Promotional posters were displayed throughout participating sites. Receptionists (ED and general practice) and pharmacy staff could also direct potential participants to the researchers who were based within each premises. The researchers were not permitted to approach patients directly due to the NHS Ethics Committee’s decision. The researchers applied the inclusion criteria and sought informed consent from all eligible and willing patients.

The four included symptoms/ailments were:

- Musculoskeletal pain: aches or pain in arms or legs or back or hands or feet;
- Eye discomfort;
- Gastrointestinal disturbance: nausea or vomiting or diarrhoea or constipation;
- Upper respiratory tract-related: sore throat or cough or cold or sinus problems.

The above groups of symptoms were chosen because they have previously been identified as occurring frequently within EDs and general practices.⁵

Data collection

Data collection was undertaken using the same process across all settings to minimise bias. Participants were asked to self-complete three questionnaires; one at baseline prior to the consultation, one immediately after the consultation and one two weeks later. The content of the questionnaires was informed by the literature and also comprised a range of validated (EQ-5D/EQ-VAS (visual analogue scale),⁹ MISS-21 (Medical Interview satisfaction Scale),¹⁰ triggers for seeking care¹¹) and non-validated measures (global satisfaction very satisfied/dissatisfied (5-point)). The MISS-21¹⁰ was developed to use with consultations in general practice. The items were rephrased to ensure face validity for their use with participants recruited from the EDs and community pharmacies. The scale comprises four subscales: distress relief; communication; rapport and compliance intent. The urgency for seeking care and perceived severity items were derived from previous research.^{12 13}

The purpose of the baseline questionnaire was to collect basic demographics, triggers for seeking care and contact details for follow-up. Baseline information was also collected on potential predictors or confounders and included the index minor ailment, age, gender, employment status, marital status, living alone, whether it was the first time they had experienced the symptoms, how serious the symptoms were perceived to be, belief on timing of visit, how long they had had the ailment, and baseline EQ-5D visual analogue scale (EQ-VAS).

The postconsultation questionnaire asked about satisfaction (global satisfaction and MISS-21), the outcome of the consultation (eg, medicines sold or supplied) and associated costs. The 2-week follow-up questionnaire asked participants about symptom resolution, quality of life, re-consultation for the index ailment and health service utilisation since their index consultation.

One version was used for each setting (ED, general practice, community pharmacy) and differed only in site-specific questions relevant to the triggers for seeking care. Although developed for self-completion, if asked to do so, a researcher would assist participants in completing the baseline and/or satisfaction questionnaires by reading the questions and response options to the participant. Participants were permitted to take the postconsultation questionnaire away with them (with a reply-paid envelope) for later completion if necessary. Follow-up questionnaires were mailed to participants. They were given the option of self-completing these and returning them by mail or completing it with a researcher over the phone. Those opting for mailed questionnaires were sent reminders at 2 and 4 weeks after the initial mailing. The study processes and documentation (questionnaires) were piloted over a 3.5-day period and included 28 patients recruited from one pharmacy (n=3), one

general practice (n=11) and the ED (n=14) in Grampian. Changes were made to the promotional materials and questionnaires as a result of piloting.

Exposure and outcomes

The exposure of interest was setting of care: ED, general practice or pharmacy. The primary outcome was symptom resolution 2 weeks after the index consultation. Secondary outcomes (collected in the postconsultation questionnaire) were: sold or supplied medicines, satisfaction measured by MISS-21,¹⁰ participant experience, duration of consultation and quality of life (QoL) measured by the EQ-5D-3L.¹⁴ Triggers for choice of site of presentation are reported as a secondary outcome from the baseline questionnaire.

Sample size

A symptom resolution rate of 70% was assumed based on a previous estimate.¹⁵ A minimum of 81 participants were required per setting. The prevalence of minor ailments was anticipated to be higher in community pharmacies and general practices⁵ compared with EDs. As such, recruitment targets for the higher prevalence sites were set at 160 participants and from the EDs were 80. Based on these estimates, the study would have over 80% power at a 5% level to detect differences of 10% between symptom resolution rates of 70% (95% CI 60% to 80%) across settings.

Statistical methods and analysis

Data were entered into an IBM SPSS Statistics V.20 database and a 10% random sample checked for accuracy. Categorical variables were described with frequency and valid percentages. Normally distributed continuous variables were summarised using mean and SD with median and IQR provided if data were skewed. Comparison of categorical variables between sites of presentation (ED, general practice, pharmacy) was undertaken using χ^2 tests or Fisher's exact test where required. Comparison of continuous variables between sites was undertaken using analysis of variance (ANOVA) if normally distributed and Kruskal-Wallis if skewed. Logistic regression was used to ascertain predictors of binary outcomes (eg, symptom resolution). Linear regression was used to model QoL (EQ-5D) at follow-up. Variables were included in these models if significant ($p \leq 0.05$) in the univariate analysis (independent t-test, ANOVA, correlation as appropriate) or belonged to an a priori list of variables for inclusion (site, ailment, age, gender, baseline EQ-5D and duration of symptoms). All analyses were undertaken on complete cases for that outcome.

For each participant, costs were estimated for the index consultation and any actions taken in the next 2 weeks for the original minor ailment in question (health professional contacts, medication and investigations). Both NHS and any patient out of pocket costs (for medication only) were included. Costs were estimated in UK sterling (£) at 2011/2012 financial year

levels, but not discounted as the time frame was less than 1 year. A complete case approach¹⁶ was adopted at each time point. Thus, costs were only estimated if data were available for each aspect of resource use for the index consultation/follow-up questionnaire. Cost per participant was estimated by summing the index consultation cost and the costs incurred in the 2 weeks thereafter. To estimate the impact of the index consultation on health-related QoL, participants were asked to complete the EuroQoL EQ-5D-3L¹⁴ at baseline and 2-week follow-up. This enabled a cost-utility analysis to be undertaken. Quality adjusted life year (QALY) scores were only calculated for those with complete baseline and follow-up EQ-5D-3L scores, and it was assumed scores changed linearly between these two time points.¹⁷

STATA V.11 was used for all economic analyses; *p* values ≤ 0.05 were deemed statistically significant. Multiple linear regression¹⁷ with backward stepwise selection was used to estimate the mean cost difference (incremental cost) between the three settings and the mean QALY difference (incremental effect). Individuals with missing data for any of the covariates in the final model were excluded. If care in one setting did not dominate the care provided in the other two settings,¹⁸ the incremental cost per QALY gain (incremental cost-effectiveness ratio (ICER)) was calculated (mean incremental cost/mean incremental QALY gain) by comparing each remaining setting with the next cheapest. In line with National Institute for Health and Care Excellence (NICE) guidance,¹⁹ any calculated ICER would be compared with a cost-effectiveness threshold (λ) of £20 000 per QALY.

Subgroup analysis

Resolution of symptoms was compared separately across sites for each minor ailment (where numbers permitted). QoL outcomes (EQ-5D-3L utility and EQ-VAS), participant experience and MISS-21 scales at baseline and follow-up were compared separately within site for individuals whose symptoms did and did not resolve.

RESULTS

Setting characteristics

Two EDs, 6 general practices and 10 community pharmacies participated, equally distributed across the two regions. Both EDs were located in teaching hospitals. Of the 10 pharmacies, 6 were large chain, 2 small-chain and 2 independent single outlets. Four general practices were in urban areas. The pharmacies and practices were located across a range of deprivation codes (online supplementary appendix 1).

Participant characteristics

In total 377 participants were recruited, 94.3% of the target sample size ($n=400$; table 1). Of these, 81, 162 and 134 were recruited from EDs, general practices and community pharmacies, respectively. Thus target recruitment was met for general practice and ED. Participant characteristics by

setting are shown in table 1. Participants recruited from community pharmacies were significantly older than those recruited elsewhere. ED participants were significantly younger and more likely to be single compared with participants recruited in general practices and pharmacies.

Most participants presented with musculoskeletal aches and pains (46.4%, $n=175$) or upper respiratory tract symptoms (28.9%, $n=109$). Presenting condition varied across settings, for example, musculoskeletal symptoms dominated in ED (81.5%, $n=66$; table 1). In total, 39.2% ($n=147$) of participants were experiencing the symptom for the first time (64.2% ($n=52$) in ED, 38.8% ($n=62$) in general practices and 24.6% ($n=33$) in pharmacies). Participants recruited from ED were significantly more likely to have had the symptom for a short period of time prior to their index consultation (median 1 day (IQR 0.5–7.0)) compared with participants in general practice (median 14 days (IQR 3–42)). ED participants were significantly more likely to consider their symptom to be 'somewhat serious' compared with participants recruited from other sites and were significantly more likely (67.5%, ($n=52$)) to state that they needed to seek care for their symptom(s) within 24 h compared with general practice (47.2%, ($n=76$)) and pharmacy-based participants (37.9%, ($n=50$)).

Data from postconsultation questionnaires were available for 89.7% (338/377) of participants: ED (77.8%, 63/81), general practice (87%, 141/162) and pharmacy (100%). The follow-up questionnaire was completed by 263 (69.8%) participants: 63% (51/81) ED, 71.6% (116/162) general practice and 72% (97/134) pharmacy. The response rates did not differ significantly across the settings ($p=0.29$).

Health-related outcomes

Primary outcome: symptom resolution

Median follow-up was 17 days (IQR 15–21) and most (86.1%) participants were followed up within 1 month. At follow-up, the proportion of symptoms that had resolved was similar across settings with 37.3% resolving in ED, 35.7% in general practice and 44.3% in pharmacy ($p=0.415$, table 2). Univariate analysis showed that significant predictors of symptom resolution were: the type of presenting symptom, its perceived seriousness, the urgency with which the participant believed they should seek help, duration of symptoms and baseline QoL (table 3). In the adjusted model, significant predictors were: presenting symptom and belief about the urgency with which they should visit their chosen site (ie, 24 or >24 h; table 3). Participants who believed they needed to consult about their symptoms within 24 h were more likely to report symptom resolution. The odds of resolution were significantly higher for non-musculoskeletal symptoms. Setting was not a significant predictor of resolution.

Secondary outcomes: postconsultation

A significantly higher proportion ($p<0.001$) of participants was sold, prescribed or supplied medicines from

Table 1 Characteristics of participants by site including information on the symptoms per cent (n)

	Overall (N=377)	ED (N=81)	General practice (N=162)	Pharmacy (N=134)	p Value*
<i>Participant demographics</i>					
<i>Age</i>					
Mean (SD)	48.3 (17.8)	42.2 (17.9)	48.3 (17.6)	51.9 (17.1)	0.001†
Missing	(7)	(4)	(2)	(1)	
<i>Gender</i>					
Male	40.6 (151)	44.9 (35)	40.4 (65)	38.3 (51)	0.646‡
Female	59.4 (221)	55.1 (43)	59.6 (96)	61.7 (82)	
Missing	(5)	(3)	(1)	(1)	
<i>Employment</i>					
Employed full-time	60.5 (225)	66.7 (52)	64.6 (104)	51.9 (69)	0.006‡
Retired	22.0 (82)	10.3 (8)	23.6 (38)	27.1 (36)	
Other (including student, unemployed)	17.5 (65)	23.1 (18)	11.8 (19)	21.1 (28)	
Missing	(5)	(3)	(1)	(1)	
<i>Registered MAS§</i>					
Yes	10.2 (38)	10.4 (8)	6.2 (10)	15.0 (20)	0.079‡
No	81.1 (301)	84.4 (65)	82.6 (133)	77.4 (103)	
Do not know	8.6 (32)	5.2 (4)	11.2 (18)	7.5 (10)	
Missing	(6)	(4)	(1)	(1)	
<i>Marital Status</i>					
Single	24.2 (90)	37.2 (29)	22.4 (36)	18.8 (25)	0.012‡
Married/living with partner	61.6 (229)	53.8 (42)	65.2 (105)	61.7 (82)	
Divorced/separated/widowed	14.2 (53)	9.0 (7)	12.4 (20)	19.5 (26)	
Missing	(5)	(3)	(1)	(1)	
<i>Live alone</i>					
Yes	20.3 (75)	13.2 (10)	18.8 (30)	26.3 (35)	0.061‡
No	79.7 (294)	86.8 (66)	81.2 (130)	73.7 (98)	
Missing	(8)	(5)	(2)	(1)	
<i>Registered with GP in the UK</i>					
Yes	97.8 (364)	98.7 (77)	98.1 (158)	97.0 (129)	–
No	2.2 (8)	1.3 (1)	1.9 (3)	3.0 (4)	
Missing	(5)	(3)	(1)	(1)	
<i>Nationality</i>					
British	94.9 (351)	93.6 (73)	96.2 (153)	94.0 (125)	–
Other EU	3.0 (11)	2.6 (2)	3.1 (5)	3.0 (4)	
Non-EU	2.2 (8)	3.8 (3)	0.6 (1)	3.0 (4)	
Missing	(7)	(3)	(3)	(1)	
<i>Norfolk only</i>					
	(N=163)	(N=35)	(N=71)	(N=57)	
<i>Pay for prescriptions</i>					
Yes	51.2 (82)	57.6 (19)	50.0 (35)	49.1 (28)	0.713‡
No	48.8 (78)	42.4 (14)	50.0 (35)	50.9 (29)	
Do not know/missing*	(3)	(2)	(1)	(0)	
<i>Ailment characteristics</i>					
<i>Presenting Symptoms</i>					
Musculoskeletal aches and pains	46.4 (175)	81.5 (66)	36.4 (59)	37.3 (50)	<0.001‡
Eye-related	11.4 (43)	14.8 (12)	11.1 (18)	9.7 (13)	
Stomach upset	4.8 (18)	1.2 (1)	6.2 (10)	5.2 (7)	
Upper respiratory tract	28.9 (109)	0 (0)	34.0 (55)	40.3 (54)	
Multiple symptoms	8.5 (32)	2.5 (2)	12.3 (20)	7.5 (10)	
<i>First time had symptoms</i>					
Yes	39.2 (147)	64.2 (52)	38.8 (62)	24.6 (33)	<0.001‡
No	60.8 (230)	35.8 (29)	61.3 (98)	75.4 (101)	
Missing	(2)	(0)	(2)	(0)	

Continued

Table 1 Continued

	Overall (N=377)	ED (N=81)	General practice (N=162)	Pharmacy (N=134)	p Value*
How serious are the symptoms?					
Not serious at all	37.0 (136)	16.9 (13)	27.5 (43)	60.2 (80)	<0.001‡
Somewhat serious	54.1 (199)	75.3 (58)	61.4 (97)	33.1 (44)	
Very serious	9.0 (33)	7.8 (6)	11.4 (18)	6.8 (9)	
Missing	(9)	(4)	(4)	(1)	
How soon did you need to visit?					
Within 24 h	48.1 (178)	67.5 (52)	47.2 (76)	37.9 (50)	<0.001‡
24 h or more	51.9 (192)	32.5 (25)	52.8 (85)	62.1 (82)	
Missing	(7)	(4)	(1)	(2)	
Duration of symptoms					
Median (IQR) in days	7 (2–35)	1.0 (0.5–7.0)	14 (3–42)	7 (2–80)	<0.001¶
Baseline EQ-VAS	(n=370)	(n=76)	(n=160)	(n=134)	0.455†
Mean (SD)	68.0 (19.6)	68.9 (22.4)	66.6 (18.9)	69.3 (18.8)	

*Tests exclude missing data.

†ANOVA.

‡Pearson χ^2 .

§National MAS (Scotland).

¶Kruskal-Wallis test.

ANOVA, analysis of variance; ED, emergency department; EU, European Union; GP, general practitioner; MAS, Minor Ailment Scheme; VAS, visual analogue scale.

Table 2 Outcomes: resolution of symptoms, quality of life and outcome of index consultation by setting

	Overall % (n)	ED % (n)	General practice % (n)	Pharmacy % (n)	p Value
Resolution of symptoms at 2-week follow-up	(n=263)	(n=51)	(n=116)	(n=97)	0.415*
Resolved	39.2 (103)	37.3 (19)	35.7 (41)	44.3 (43)	
Follow-up EQ-VAS	(n=261)	(n=50)	(n=115)	(n=96)	0.054†
Mean (SD)	77.4 (18.2)	74.8 (19.6)	75.5 (19.5)	80.9 (15.3)	
Change in EQ-VAS	(n=259)	(n=50)	(n=113)	(n=96)	0.726‡
Mean (SD)	8.43 (18.2)	7.6 (17.9)	7.8 (18.1)	9.6 (18.5)	
Outcome of index consultation	(n=338)	(n=63)	(n=141)	(n=134)	
Sold or supplied medicines	(n=334)	(n=61)	(n=141)	(n=132)	<0.001*
Yes	66.5 (222)	31.1 (19)	61.7 (87)	87.9 (116)	
Sold or supplied items	(n=327)	(n=59)	(n=138)	(n=130)	<0.001*
Yes	9.3 (35)	37.3 (22)	2.2 (3)	7.7 (10)	
Satisfaction	(n=334)	(n=60)	(n=136)	(n=127)	0.115*‡
Satisfied	64.7 (209)	53.3 (32)	66.2 (90)	68.5 (87)	
Neither	31.9 (103)	45.0 (27)	27.9 (38)	29.9 (38)	
Dissatisfied	3.4 (11)	1.7 (1)	5.9 (8)	1.6 (2)	
Experience	(n=328)	(n=61)	(n=141)	(n=126)	0.011§
Median (IQR)	10 (8–10)	9 (7–10)	10 (8–10)	10 (8–10)	
Duration of wait (minutes)	(n=334)	(n=62)	(n=141)	(n=131)	<0.001§
Median (IQR)	5 (1–30)	47.5 (30–86.25)	15 (5–30)	1 (0–2)	
Duration of consultation (minutes)	(n=332)	(n=61)	(n=141)	(n=130)	<0.001§
Median (IQR)	5 (2–10)	10 (5–27.5)	10 (5–10)	2 (1–3)	
MISS 21 items¶	(n=268)	(n=63)	(n=141)	(n=101)	
Total score: mean (SD)	5.27 (0.97)	5.39 (0.94)	5.70 (1.01)	4.75 (0.81)	<0.001†
Distress relief: mean (SD)	4.97 (1.24)	5.29 (1.14)	5.39 (1.17)	4.31 (1.09)	<0.001†
Communication: mean (SD)	5.50 (1.19)	5.43 (1.24)	5.89 (1.16)	5.05 (1.05)	<0.001†
Rapport: mean (SD)	5.38 (1.06)	5.48 (1.02)	5.83 (1.32)	4.83 (0.91)	<0.001†
Compliance intent: mean (SD)	5.40 (1.10)	5.41 (1.18)	5.69 (1.05)	5.04 (1.02)	<0.001†

*Pearson χ^2 .

†ANOVA.

‡For 2x2 where neither and dissatisfied are combined.

§Kruskal-Wallis test.

¶n is for total score.

ANOVA, analysis of variance; ED, emergency department; MISS 21, Medicine Information Satisfaction Score; VAS, visual analogue scale.

Table 3 Logistic regression to assess factors associated with symptom resolution

Per cent (n) within variable	Resolved % (n)	Not resolved % (n)	Unadjusted		Adjusted	
			OR	95% CI	OR	95% CI
Site						
Pharmacy	44.3 (43)	55.7 (54)	1.00		1.00	
General practice	35.7 (41)	64.3 (74)	0.70	(0.40 to 1.21)	0.72	(0.35 to 1.49)
ED	37.3 (19)	62.7 (32)	0.75	(0.37 to 1.49)	0.97	(0.37 to 2.57)
Presenting symptoms						
Musculoskeletal aches/pains	22.9 (27)	77.1 (91)	1.00		1.00	
Eye pain/discomfort	73.3 (22)	26.7 (8)	9.27	(3.71 to 23.2)	7.96	(2.93 to 21.6)
Stomach upset	63.6 (7)	36.4 (4)	5.90	(1.61 to 21.7)	7.01	(1.73 to 28.5)
Upper respiratory tract	50.0 (41)	50.0 (41)	3.37	(1.83 to 6.20)	2.80	(1.28 to 6.14)
Multiple symptoms	27.3 (6)	72.7 (16)	1.26	(0.45 to 3.55)	1.70	(0.48 to 6.14)
Gender						
Male	43.5 (40)	56.5 (52)	1.00		1.00	
Female	37.3 (63)	62.7 (106)	0.77	(0.46 to 1.30)	0.67	(0.35 to 1.27)
Employment						
Employed full time	42.0 (63)	58.0 (87)	1.00			
Retired	33.8 (24)	66.2 (47)	0.71	(0.39 to 1.27)		
Other (including student, unemployed)	40.0 (16)	60.0 (24)	0.92	(0.45 to 1.87)		
Marital status						
Single	50.0 (25)	50.0 (25)	1.00			
Married/living with partner	37.4 (65)	62.6 (109)	0.60	(0.32 to 1.12)		
Divorced/separated/widowed	35.1 (13)	64.9 (24)	0.54	(0.23 to 1.30)		
Live alone						
Yes	40.1 (83)	64.2 (34)	0.84	(0.45 to 1.56)		
No	35.8 (19)	59.9 (124)	1.00			
First time had symptoms						
Yes	41.4 (41)	58.6 (58)	1.18	(0.71 to 1.97)		
No	37.4 (61)	62.6 (102)	1.00			
How serious are the symptoms?						
Not serious at all	49.0 (48)	51.0 (50)	1.00		1.00	
Somewhat/very serious	33.3 (53)	66.7 (106)	0.52	(0.31 to 0.87)	0.57	(0.28 to 1.27)
Belief on required						
Within 24 h	48.8 (60)	51.2 (63)	1.00		1.00	
Time of visit						
24 h or more	29.6 (40)	70.4 (95)	0.44	(0.27 to 0.74)	0.42	(0.23 to 0.78)
How long had symptoms?						
Median (IQR) days	3 (1–7)	14 (3.5–56)	0.99	(0.99 to 1.00)	0.99	(0.99 to 1.00)
Baseline EQ-VAS						
Mean (SD)	73.2 (17.4)	66.3 (20.6)	1.02	(1.01 to 1.03)	1.01	(0.99 to 1.02)
Age (years)						
Mean (SD)	49.8 (17.5)	53.7 (15.9)	0.99	(0.97 to 1.00)	0.99	(0.97 to 1.00)

ED, emergency department; VAS, visual analogue scale.

pharmacies (87.9%) compared with general practice (61.7%) or ED (31.1%). A higher proportion of participants was supplied other items (eg, dressings) in ED (37.3%) compared with being sold or supplied in pharmacy (7.7%) or prescribed by general practice (2.2%; [table 2](#)).

Satisfaction was assessed using two measures: MISS-21¹⁰ and global (5-point) scale (very satisfied/very dissatisfied). MISS-21 total and component scores differed significantly ($p<0.001$) across settings and were consistently highest in general practice and lowest in pharmacy ([table 2](#)). While higher satisfaction scores were achieved using the global measure in pharmacy (68.5%) and general practice (66.2%) compared with

ED (53.3%), these were not significant ($p=0.115$; [table 2](#)). The duration of wait for the index consultation was significantly longer ($p<0.001$) in ED compared with other sites and could have influenced satisfaction ratings ([table 2](#)).

Secondary outcomes at follow-up: QoL

QoL measured by the EQ-5D visual analogue scale (EQ-VAS) at follow-up was highest for those who had visited a pharmacy, but differences between sites were not statistically significant ([table 2](#)). Participants who visited a pharmacy had the greatest observed change in EQ-VAS compared with baseline ([table 2](#)), but this was not statistically different ($p=0.726$). Employment status

and participants' perceived seriousness of their symptoms were significant predictors of QoL (EQ-VAS) at follow-up in the univariate analysis (table 4). Only baseline EQ-VAS score and employment status were found to be independent predictors of follow-up QoL in the adjusted linear regression model which includes the a priori variables (table 4). Individuals with higher baseline QoL had higher follow-up QoL and those whose employment status was 'other' (ie, student, unemployed) showed lower QoL at follow-up compared with employed participants. In the adjusted model, neither site of presentation nor presenting symptoms were independent predictors of QoL, although the lower QoL for general practice was of borderline significance ($p=0.051$).

Subgroup analysis

Combining all ailments, no significant difference in proportion resolved was found between sites (table 2). However, as table 5 indicates for musculoskeletal aches and pains only, resolution was higher for ED (31.0%) and pharmacy (28.6%) than for general practice (9.8%; $p=0.045$). No difference in the proportion resolved was found for sore throat/cough cold with 51.2% resolving from pharmacy and 48.8% from general practice ($p=0.825$). There were insufficient numbers to compare for eye pain/discomfort or for stomach upset (table 5).

Individuals whose symptoms had resolved had significantly higher QoL at follow-up than for those with unresolved symptoms (table 6). Symptom resolution was also significantly associated with participant satisfaction measured by the MISS-21 (table 6) and this was driven mainly by participants recruited from general practices.

In terms of reconsultation following the index consultation, subsequent visits to general practice or pharmacies were not significantly different for site of presentation (data not presented: available from author). Participants whose index consultation occurred in ED compared with other sites were significantly more likely to return to ED within the following 2 weeks.

Cost-related outcomes

The mean (95% CI) overall cost (sum of total initial consultation cost and 2-week follow-up costs) was highest in ED £147.09 (£125.32 to £168.85, $n=48$) compared with £82.34 (£63.10 to £101.58, $n=115$) and £29.30 (£21.60 to £37.00, $n=95$) in general practice and pharmacy, respectively. The mean follow-up QALY score was estimated to be highest for pharmacy participants (a score was estimated for 67.9% of participants, $n=91$), compared with those in general practice (63.0%, $n=102$) and ED (55.6%, $n=45$), although this could partially be accounted for by the fact that they had the highest mean score at the initial visit (this is adjusted for in the subsequent regression analysis).

The number of respondents contributing to the mean incremental cost analysis was 44/81 (54.3%) for ED, 108/162 (66.7%) for general practice and 91/134 (67.9%) for pharmacy; these rates did not differ

significantly across the three settings ($p=0.10$). After adjusting for age, gender, baseline EQ-5D-3L score and type of ailment, the mean cost was estimated to be lowest for pharmacy participants; mean (95% CI) incremental costs (compared with pharmacy) were estimated to be £57.04 (£34.95 to £79.12; general practice; $p<0.001$) and £113.63 (£81.78 to £145.47; ED; $p<0.001$). After adjusting for age, gender, baseline EQ-5D-3L utility score, type of ailment, employment status and whether it was the first time they had experienced the illness, the mean QALY score was estimated to be highest for pharmacy participants, though there were no significant differences between or across settings. The mean (95% CI) incremental QALY gain for pharmacy participants compared with general practice and ED participants was estimated to be 0.001 (0.000 to 0.002) and 0.001 (−0.001 to 0.002), respectively. Looking at costs and effects together, as pharmacy was estimated to be less costly and as effective (in terms of symptom resolution) compared with ED and general practice, it can be said to dominate both of these options. It should however be noted that there were no significant differences in outcomes between any of the groups.

Triggers for seeking care

The main predefined triggers for choice of site were location and convenience of not having to travel too far (table 7). For participants who visited ED, the fact that they would have to wait longer for an appointment with a general practice ranked highly (rank 2, 37%). Those visiting a pharmacy felt their symptoms were not serious enough to consult a doctor (rank 3, 51.5%), while those visiting the general practice, 63.6% (rank 2) felt their symptoms were not serious enough for ED. The triggers differed across sites and although 'site in a convenient location' was a popular choice for participants recruited from all three types of site, it was more common in pharmacy (79.1%, $n=106$). 'Knowing the staff' was given as a reason in a higher proportion (45.7%, $n=74$) of those visiting the general practice compared with other settings. Participants were more likely to seek care from ED if they had not experienced the symptom previously or if the symptom was of recent onset (table 1). The former suggests that previous experience may affect decision-making regarding health-seeking behaviour.

DISCUSSION

Principal findings

This is the first study to compare the health-related and cost-related outcomes associated with the management of minor ailments in these three healthcare settings: EDs, general practices and community pharmacies. Symptom resolution and mean improvement in QoL was similar across participants from all settings suggesting equivalence in health outcome. While the absolute rate of symptom resolution appeared to be better in pharmacy, the overall lower than anticipated resolution rates

Table 4 Factors affecting QoL (EQ-VAS) at follow-up

Per cent (n) within variable	Mean (SD)	Univariate p Value	Adjusted linear regression (R ² =33%)		
			Coefficient	95% CI	p Value
Constant term			54.0	(38.1 to 56.6)	
Site					
Pharmacy	80.9 (15.3)	0.054*	Ref		
General practice	75.5 (19.5)		−4.58	(−9.19 to 0.02)	0.051
ED	74.8 (19.6)		−3.61	(−9.72 to 2.49)	0.245
Ailment					
Musculoskeletal aches/pains	73.1 (19.4)	0.005†	Ref		
Eye pain/discomfort	81.2 (16.5)		1.56	(−4.66 to 7.79)	0.622
Stomach upset	84.7 (11.2)		7.22	(−1.92 to 16.4)	0.121
URT	81.8 (16.5)		4.61	(−0.38 to 9.60)	0.070
Multiple ailments	74.8 (18.4)		5.33	(−2.65 to 13.3)	0.189
Gender					
Male	76.6 (18.2)	0.536‡	Ref		
Female	78.0 (18.2)		−0.12	(−4.25 to 4.02)	0.956
Employment					
Employed	80.6 (15.9)	0.010*	Ref		
Retired	74.9 (18.7)		−3.28	(−8.99 to 2.43)	0.259
Other (including student, unemployed)	70.6 (23.1)		−8.27	(−14.0 to −2.55)	0.005
Marital status					
Single	78.9 (17.1)	0.736†			
Married/living with partner	77.5 (18.6)				
Divorced/separated/widowed	75.8 (18.1)				
Live alone					
Yes	77.6 (18.7)	0.898‡			
No	77.2 (16.6)				
First time experienced illness?					
Yes	76.1 (18.9)	0.159‡			
No	79.4 (17.1)				
How serious?					
Not serious at all	82.5 (14.1)	<0.001‡	Ref		
Somewhat/very serious	74.7 (19.7)		−0.55	(−5.05 to 3.96)	0.811
Belief on required					
Within 24 h	79.4 (16.9)	0.075‡			
Time of visit					
24 h or more	75.3 (19.3)				
Correlation					
How long had ailment?					
(Days)	−0.262§	<0.001	−0.001	(−0.002 to 0.000)	0.163
Baseline EQ-VAS	0.503§	<0.001	0.47	(0.37 to 0.57)	<0.001
Age					
(Years)	−0.059§	0.343	−0.11	(−0.27 to 0.5)	0.181

Mean (SD) presented as convention for QoL, but testing may be non-parametric dependent on other assumptions.

*Kruskal-Wallis test.

†ANOVA.

‡Independent t test.

§Spearman's rank correlation.

ANOVA, analysis of variance; ED, emergency department; QoL, quality of life; URT, upper respiratory tract; VAS, visual analogue scale.

across all settings, together with reduced numbers of participants at follow-up, meant that the study was under-powered to detect whether these differences were statistically significant.

Mean overall costs were significantly lower for pharmacy consultations compared with general practice consultations and with ED consultations which were associated with the highest costs. The results suggest that substantial savings could be achieved if demand on ED

and general practice was shifted to the community pharmacy setting. Some caution is needed because the perceived seriousness and urgency with which to seek care, as well as whether the participant had experienced the symptom(s) previously, varied across settings.

Factors which influenced the patients' site of choice were explored using measures of symptom characteristics, as well as predefined triggers previously shown to influence care-seeking behaviour.¹¹ In this study, the

Table 5 Resolution of symptoms by ailment

	Overall % (n)	ED % (n)	General practice % (n)	Pharmacy % (n)	p Value
Resolution of symptoms					
Musculoskeletal aches and pains	(n=118)	(n=42)	(n=41)	(n=35)	0.045*
Resolved	22.9 (27)	31.0 (13)	9.8 (4)	28.6 (10)	
Sore throat/cough cold	(n=82)	–	(n=41)	(n=41)	0.825*
Resolved	50.0 (41)	–	48.8 (20)	51.2 (21)	
Eye pain/discomfort†	(n=30)	(n=7)	(n=12)	(n=11)	–
Resolved	73.3 (22)	71.4 (5)	83.3 (10)	63.6 (7)	
Stomach upset†	(n=11)	(n=1)	(n= 6)	(n= 4)	–
Resolved	63.6 (7)	100 (1)	50.0 (3)	75.0 (3)	

*Pearson χ^2 test.

†Insufficient power statistical test.

ED, emergency department.

majority (64.2%) of ED users were experiencing the symptom for the first time with a median onset of one day, perceived it to be somewhat/very serious (83.1%) and concluded that a visit to a health service provider was needed within 24 h. These characteristics differentiated ED users from general practice and community pharmacy users. Pharmacy users would have either presented with a direct product request or with a symptom presentation. The former scenario could suggest that the user was familiar with the condition for which they required treatment while the latter would suggest a lack of familiarity with treatment options for the symptom. As such, this may have introduced a bias into the types of user seeking care from community pharmacies. The type of presentation (direct product request/symptom presentation) was not documented during the recruitment process. This study relied on each patient's perception and self-reporting of their symptoms and did not include any qualitative exploration of these perceptions and health-seeking behaviour. Evidence is needed regarding the key determinants of patients' health-seeking behaviour for minor ailments in general and from the ED setting in particular because of the high costs associated with this setting. Theory-based studies are needed to identify key modifiable determinants of this behaviour which in turn will lead to the identification of behaviour change strategies to target these determinants.

Convenience of location and shorter travelling distances were the most important predefined triggers for the choice of healthcare setting. Among ED participants, one of the major reasons cited for choosing the ED was the expected longer wait to see a general practitioner (GP), which may support the finding that convenience was a major factor in selecting the site of treatment, or that participants may have thought it was urgent, or both. The use of EDs for non-urgent conditions has been shown to be associated with fewer available primary care providers (not pharmacies)²⁰ but the vast majority of participants in the current study were

registered with a GP so would have had access to general practice services if they had opted to do so.

The positive association between participant satisfaction and symptom resolution is an important finding. An earlier study of acute minor illness where index visits were made to EDs, family physicians or walk-in clinics, showed patient satisfaction with patient-centred care was associated with reduced reutilisation rates.¹² Doctors' communication behaviour is a major driver of patient satisfaction with GP consultations.²¹ This emphasises the need for good consultation skills among healthcare professionals. Suboptimal communication during community pharmacy consultations has been highlighted as an area for improvement^{22–23} because it is associated with suboptimal and inappropriate outcomes.^{23–25}

Other studies have compared combinations of settings, for example, walk-in clinics versus family practice versus ED,^{12–13} but no previous study included community pharmacy as a comparator. This study did not explore whether the patients were seeking care for reasons other than their minor ailment, however, the promotional materials clearly specified the four ailments of interest and as such, an assumption of this study is that the minor ailment was the primary reason for presentation. The inclusion of different types (independent single outlets, small/large chains) of pharmacies, as well as a variety of locations (urban/rural) of pharmacies and general practices, increased the generalisability of the results. The participants' choice of setting had face validity; symptoms that were perceived as somewhat serious (and therefore might be more worrying and require expert medical attention) presented more often in ED than in other settings. All participating sites were within 25 miles of a major city, therefore the results may not be generalisable to people living in more remote areas. People in these areas may behave differently when managing their symptoms, especially if convenience is a major influencing factor for them too. Both EDs were located in teaching hospitals and as such, the results may not be generalisable to EDs in other types of

Table 6 Quality of life and MISS 21 outcomes by symptom resolution

	ALL		ED		General practice		Pharmacy	
	n	Mean (SD)	n	Mean (SD)	n	Mean (SD)	n	Mean (SD)
EQ-5D-3L utility baseline								
Resolved	101	0.74 (0.22)	18	0.58 (0.28)	40	0.76 (0.22)	43	0.79 (0.17)
Not resolved	148	0.64 (0.28)	29	0.49 (0.32)	69	0.64 (0.26)	50	0.73 (0.26)
p Value*		0.002		0.362		0.01		0.195
EQ-5D-3L utility follow-up								
Resolved	99	0.93 (0.12)	18	0.94 (0.09)	39	0.95 (0.10)	42	0.91 (0.15)
Not resolved	150	0.71 (0.26)	30	0.62 (0.31)	67	0.71 (0.27)	35	0.77 (0.22)
p Value*		<0.001		<0.001		<0.001		<0.001
EQ-VAS baseline								
Resolved	103	73.2 (17.4)	19	73.1 (18.0)	41	72.6 (17.8)	43	73.7 (17.2)
Not resolved	158	66.3 (20.6)	32	64.5 (24.2)	72	64.6 (19.8)	54	70.0 (19.2)
p Value*		0.004		0.187		0.034		0.285
EQ-VAS follow-up								
Resolved	103	86.3 (11.6)	19	81.5 (14.1)	41	87.0 (10.6)	43	87.7 (11.1)
Not resolved	157	71.5 (19.4)	31	70.8 (21.6)	73	69.0 (20.5)	54	75.4 (16.1)
p Value*		<0.001		0.061		<0.001		<0.001
	n	Median (IQR)	n	Median (IQR)	n	Median (IQR)	n	Median (IQR)
Experience								
Resolved	100	10 (8–10)	18	9.5 (7.75–10)	41	10 (9–10)	41	10 (8–10)
Not resolved	155	10 (8–10)	30	9 (7.75–10)	73	10 (8–10)	52	10 (8.25–10)
p Value†		0.77		0.46		0.958		0.201
MISS 21 scores								
TOTAL score								
Resolved	90	5.38 (4.71–6.15)	15	5.48 (4.95–6.62)	37	6.05 (5.19–6.69)	38	4.74 (4.14–5.48)
Not resolved	119	5.00 (4.48–5.90)	26	5.31 (4.79–6.01)	56	5.19 (4.81–6.37)	37	4.43 (4.00–5.02)
p Value†		0.047		0.291		0.015		0.116
Distress								
Resolved	90	5.00 (4.17–6.00)	15	5.50 (5.00–7.00)	37	6.00 (5.00–7.00)	38	4.17 (4.00, 5.00)
Not resolved	119	4.67 (4.0–5.67)	26	5.33 (4.46–6.08)	56	5.19 (4.21–6.0)	37	4.00 (3.92, 4.50)
p Value†		0.009		0.074		0.004		0.046
Communication								
Resolved	90	5.00 (4.75–6.75)	15	5.50 (5.00–7.00)	37	6.50 (5.25–7.00)	38	5.00 (4.25, 6.25)
Not resolved	119	5.00 (4.50–6.25)	26	5.13 (4.69–6.50)	56	5.50 (5.00–7.00)	37	4.75 (4.13, 5.50)
p Value†		0.161		0.348		0.153		0.206
Rapport								
Resolved	90	5.38 (4.50–6.28)	15	5.75 (4.75–6.75)	37	6.00 (5.38–6.88)	38	4.56 (4.22, 5.38)
Not resolved	119	5.00 (4.75–6.00)	26	5.44 (4.84–6.31)	56	5.25 (4.88–6.88)	37	4.75 (4.00, 5.06)
p Value†		0.834		0.668		0.403		0.5
Compliance intent								
Resolved	90	5.50 (4.92–6.67)	15	6.00 (5.00–6.67)	37	6.00 (5.00–7.00)	38	5.00 (4.00, 6.00)
Not resolved	119	5.00 (4.33–6.00)	26	5.17 (4.33–6.00)	56	5.33 (5.00–6.33)	37	4.43 (4.00, 5.00)
p Value†		0.057		0.119		0.12		0.136

*Independent samples t test for resolved versus not resolved.

†Mann-Whitney test for resolved versus not resolved.

ED, emergency department; MISS 21, Medicine Information Satisfaction Score; VAS, visual analogue scale.

Table 7 Triggers for choice of site

	ED		General practice		Pharmacy		
Per cent (n) unless specified	(N=81)	Rank	(N=162)	Rank	(N=134)	Rank	p Value*
Location and time							
The 'site' is in a location convenient for me†	51.9 (42)	1	69.1 (112)	1	79.1(106)	1	<0.001
Did not have to travel far to visit the 'site'†	34.6 (28)	3	51.2 (83)	3	57.4 (77)	2	<0.001
I was already visiting 'site' for different reason†	2.5 (2)	17	6.8 (11)	16	35.8 (48)	7	<0.001
It was not convenient to go to the chemist	1.2 (1)	19	1.9 (3)	18	NA	—	Low counts
It was not convenient to go to the GP	13.6 (11)	11	NA	—	15.7 (21)	17	0.837
I can get treatment/advice at 'site' without appointment	NR	—	NA	—	36.6 (49)	6	
Knowing the staff							
I know the 'staff' here†	1.2 (1)	19	45.7 (74)	5	26.9 (36)	11	<0.001
I feel the 'staff' here know me†	3.7 (3)	14	30.9 (50)	10	19.4 (26)	14	<0.001
I am comfortable discussing my illness with 'staff' here†	23.5 (19)	8	51.2 (83)	3	47.8 (64)	4	<0.001
'staff' in 'site' in general know about this illness†	25.9 (21)	7	41.4 (67)	6	30.6 (41)	8.5	0.037
I have previously successfully used 'site' to treat this illness†	16.0 (13)	9	32.1 (52)	9	38.1 (51)	5	0.003
I could discuss my illness with 'staff' in private†	13.6 (11)	11	39.5 (64)	7	21.6 (29)	12	<0.001
I needed information about illness from 'staff' at 'site'†	34.6 (28)	3	37.0 (60)	8	20.1 (28)	13	0.009
Other options							
My illness was too serious to go to GP	30.9 (25)	5	NA	—	NR	—	—
My illness was too serious to go to chemist	27.2 (22)	6	NR	—	NA	—	—
My illness was not serious enough for ED	NA	—	63.6 (103)	2	NR	—	—
My illness was not serious enough for a doctor	NA	—	NA	—	51.5 (69)	3	—
I did not want to waste the doctor's time	NR	—	NA	—	30.6 (41)	8.5	—
I previously consulted GP about illness, but it is not better	16.0 (13)	9	NA	—	18.7 (25)	15	0.775
I previously consulted pharmacy about illness, but it is not better	3.7 (3)	14	9.9 (16)	15	NA	—	0.158
I would have to wait longer for appointment at GP	37.0 (30)	2	NA	—	29.9 (40)	10	<0.001
I would have to wait longer for appointment at other services	NR	—	14.8 (24)	13	NR	—	—
I prefer not to go to ED	NA		28.4 (46)	11	NR	—	—
I prefer not to go to GP	3.7 (3)	14	NA	—	11.2 (15)	19	0.097
I prefer not to go to a chemist	4.9 (4)	13	16.7 (27)	12	NA	—	0.019
Cost of treatment							
It was cheaper to visit than other services†	1.2 (1)	19	6.8 (11)	16	17.9 (24)	16	<0.001
Treatment/advice was cheaper for my illness†	1.2 (1)	19	11.7 (19)	14	14.9 (20)	18	0.006
Advice/referral							
I came to get advice/treatment that I had seen in media†	2.5 (2)	17	0 (0)	19	1.5(2)	20	Low counts

*Pearson χ^2 test.

†Item recorded on for all sites.

ED, emergency department; GP, general practitioner; NA, not applicable for that site; NR, item not recorded for that site version.

hospital. There was slight under-recruitment from community pharmacies. This study used a cohort design rather than randomised controlled trial design because the latter was an impractical option to address the objectives of this study. Previous research identified that perceptions of service quality influence care seeking from EDs²⁶ and that competence and credibility of staff was important and more highly rated for staff in EDs compared with general practices. The current study did not include measures of quality perception nor assess participants' perceptions of staff credibility or competence.

Meaning of the study

Despite the limitations of this study in terms of sample size, reliance on self-reported data and focusing on selected common symptoms, the results provide evidence of the value of community pharmacy as a resource from the management of these types of conditions. The Keogh report highlighted the 'intense, growing and unsustainable pressure' being placed on EDs and the need to direct demand outside of hospitals for conditions that may be 'urgent but non-life threatening'.³ The same report identified that community

pharmacies are an ‘under-utilised resource’³ which could contribute substantially to reduce the burden on other healthcare providers. Self-care is the preferred method of managing minor illness for many patients²⁷ and the need to support patients with self-care has been acknowledged as a priority.³ The results of this cohort study suggest that patients with minor ailments presenting in community pharmacies will have similar health outcomes to those in general practice and ED, but that the costs will be substantially lower with pharmacy-managed consultations. Redirection of demand needs to address accessibility, affordability, availability and most importantly, safety of care.²⁸ Earlier interventions to promote the self-management of minor illness have had limited effect;²⁹ few, if any, have adopted a theory-based approach to exploring behaviour and promoting behaviour change.

Unanswered questions and future research

Convenience of location was a major influence on health-seeking behaviour for the conditions included in this study. The evidence for co-locating community pharmacies with EDs and general practices is currently lacking, but the co-location of walk-in clinics within EDs failed to influence patient satisfaction, preference or choice.³⁰ Patients’ perceptions that obtaining a GP appointment would ‘take too long’ is one which requires further investigation to determine what urgent care arrangements should be available within general practices to manage acute conditions perceived to be serious or in need of immediate attention. The substantially lower costs associated with pharmacy management of these ailments, combined with similar health outcomes across settings, suggest there is an urgent need to identify effective interventions to promote the use of community pharmacies for the management of minor ailments. These interventions should be evidence-based and underpinned by behaviour change theory to maximise their effectiveness and reproducibility. Further research is warranted and planned regarding patients’ decision-making processes for choosing their care provider when managing minor ailments.

Author affiliations

¹Division of Applied Health Sciences, Centre of Academic Primary Care, University of Aberdeen, Aberdeen, UK

²Emergency Department, Aberdeen Royal Infirmary, Aberdeen, UK

³Health Economics Group, Norwich Medical School, Norwich, UK

⁴Norwich Medical School, University of East Anglia, Norwich, UK

⁵School of Pharmacy, University of East Anglia, Norwich, UK

Twitter Follow Margaret Watson at @MagsWatson1

Acknowledgements The authors thank the patients and healthcare providers who participated in this study. They also thank J Burr and J Inch for data collection and H Riley and N Clark for their administrative support.

Contributors MCW (guarantor) had the original idea for the study and with JF, CMB, RH, TP, THS and DW designed the trial and formed the investigator group that obtained funding. MCW, RH and SF were responsible for overseeing study implementation and AB, VM and VP were responsible for

data collection. SF, THS and GRB conducted the analyses. MCW, SF and GRB drafted the manuscript which was revised by all authors. All researchers were independent of the funders and approved the final version.

Funding This work was supported by the Pharmacy Practice Research Trust which merged with the Pharmaceutical Trust for Educational and Charitable Objects in December 2012 to form Pharmacy Research UK.

Competing interests None.

Patient consent Obtained.

Ethics approval The North of Scotland Research Ethics Committee.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

1. The College of Emergency Medicine. *The drive for quality. How to achieve safe, sustainable care in our emergency departments?* London: The College of Emergency Medicine, 2013.
2. Health and Social Care Information Centre. Focus on accident and emergency. <http://www.hscic.gov.uk/catalogue/PUB13040/acc-emer-focu-on-2013-rep-V2.pdf>. Updated 2013(accessed 26 Mar 2014).
3. NHS England. *Transforming urgent and emergency care services in England. Urgent and emergency care review end of phase 1 report. Appendix 1—Revised Evidence Base from the Urgent and Emergency Care Review*. Leeds: NHS England, 2013.
4. Hippisley-Cox J, Vinogradova Y. *Trends in consultation rates in general practice 1995/1996 to 2008/9: analysis of the QResearch database*. Final report to NHS information Centre and Department of Health. Nottingham: The NHS Information Centre for Health and Social Care, 2009.
5. Watson MC, Holland R, Ferguson J, et al. *Community pharmacy management of minor illness (the MINA study)*. London: Pharmacy Research UK, 2014.
6. Jones R, White R, Armstrong D, et al. *Managing acute illnesses: an enquiry into the quality of general practice in England*. London: The King's Fund, 2010.
7. The Royal Pharmaceutical Society of Great Britain. *RPS e-PIC references on: prescription only medicines reclassified to pharmacy only medicines*. London: RPSGB, 2008.
8. Paudyal V, Watson MC, Sach T, et al. Are pharmacy-based minor ailment schemes a substitute for other service providers? A systematic review. *Br J Gen Pract* 2013;63:e472–81.
9. EuroQol Research Foundation. EQ-5D. <http://www.euroqol.org/>
10. Meakin R, Weinman J. The ‘Medical interview satisfaction scale’ (MISS-21) adapted for British general practice. *Fam Pract* 2002;19:257–63.
11. Porteous T. *How do people choose between self-care, supported self-care and GP consultation in minor illness?* [PhD thesis]. University of Aberdeen, 2007.
12. Campbell MK, Silver RW, Hoch JS, et al. Re-utilization outcomes and costs of minor acute illness treated at family physician offices, walk-in clinics, and emergency departments. *Can Fam Physician* 2005;51:82–93.
13. Hutchison B, Østbye T, Barnsley J, et al. Patient satisfaction and quality of care in walk-in clinics, family practices and emergency departments: the Ontario walk-in clinic study. *Can Med Assoc J* 2003;168:977–83.
14. Brooks R, De Charro F. EuroQol: the current state of play. *Health Policy* 1996;37:53–72.
15. Schafheutle E, Noyce P, Sheehy C, et al. *Evaluation of a pilot scheme to allow patients exempt from prescription charges to consult community pharmacists and receive ‘over the counter’ products for minor ailments free of charge under the NHS Direct Supply by community pharmacists*. Edinburgh: Scottish Government, 2002.
16. Briggs A, Clark T, Wolstenholme J, et al. Missing...presumed at random: cost-analysis of incomplete data. *Health Econ* 2003;12:377–92.

17. Manca A, Hawkins N, Sculpher MJ. Estimating mean QALYs in trial-based cost-effectiveness analysis: the importance of controlling for baseline utility. *Health Econ* 2005;14:487–96.
18. Drummond MF, Sculpher MJ, Torrance GW, *et al.* *Methods for the economic evaluation of health care programmes*. 3rd edn. New York: Oxford University Press, 2005.
19. NICE. *Guide to the methods of technology appraisal*. London: National Institute for Health and Care Excellence, 2013.
20. Mathison DJ, Chamberlain JM, Cowan NM, *et al.* Primary care spatial density and nonurgent emergency department utilization: a new methodology for evaluating access to care. *Acad Pediatr* 2013;13:278–85.
21. Paddison C, Abel G, Roland M, *et al.* Drivers of overall satisfaction with primary care: evidence from the English general practice patient survey. *Health Expect* 2013. doi:10.1111/hex.12081
22. Which Report. Can you trust your local pharmacy's advice? *WHICH?* 20 May 2013. <http://www.which.co.uk/news/2013/05/can-you-trust-your-local-pharmacies-advice-319886/> (accessed 23 Jul 2014).
23. Watson MC, Bond CM, Grimshaw J, *et al.* Factors predicting the guideline compliant supply (or non-supply) of non-prescription medicines in the community pharmacy setting. *Br Med J* 2006;15:53–7.
24. Watson MC, Walker AE, Bond CM. Community pharmacists' views and beliefs about the treatment of symptoms suggestive of vaginal thrush in community pharmacies. *Pharm World Sci* 2000;22:130–5.
25. Watson MC, Bond CM, Grimshaw JM, *et al.* Educational strategies to promote evidence-based community pharmacy practice: a cluster randomized controlled trial (RCT). *Fam Pract* 2002;19:529–36.
26. Lo S, Mc Kechnie S. Perceptions of service quality and sacrifice in patients with minor medical conditions using emergency care. *Int J Clin Pract* 2007;61:596–602.
27. Porteous T, Ryan M, Bond CM, *et al.* Preferences for self-care or professional advice for minor illness: a discrete choice experiment. *Br J Gen Pract* 2006;56:911–17.
28. Durand AMP, Gentile S, Devictor B, *et al.* ED patients: how nonurgent are they? Systematic review of the emergency medicine literature. *Am J Emerg Med* 2011;29:333–45.
29. Little P, Somerville J, Williamson I, *et al.* Randomised controlled trial of self management leaflets and booklets for minor illness provided by post. *BMJ* 2001;322:1214–17.
30. Chalder M, Montgomery A, Hollinghurst S, *et al.* Comparing care at walk-in centres and at accident and emergency departments: an exploration of patient choice, preference and satisfaction. *Emerg Med J* 2007;24:260–4.

BMJ Open

A cohort study of influences, health outcomes and costs of patients' health-seeking behaviour for minor ailments from primary and emergency care settings

M C Watson, J Ferguson, G R Barton, V Maskrey, A Blyth, V Paudyal, C M Bond, R Holland, T Porteous, T H Sach, D Wright and S Fielding

BMJ Open 2015 5:

doi: 10.1136/bmjopen-2014-006261

Updated information and services can be found at:
<http://bmjopen.bmj.com/content/5/2/e006261>

These include:

Supplementary Material

Supplementary material can be found at:
<http://bmjopen.bmj.com/content/suppl/2015/02/18/bmjopen-2014-006261.DC1.html>

References

This article cites 16 articles, 8 of which you can access for free at:
<http://bmjopen.bmj.com/content/5/2/e006261#BIBL>

Open Access

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Emergency medicine](#) (132)
[Evidence based practice](#) (359)
[General practice / Family practice](#) (322)
[Health services research](#) (707)
[Public health](#) (1149)

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>