

Crystal structure of an eight-coordinate terbium(III) ion chelated by *N,N'*-bis(2-hydroxybenzyl)-*N,N'*-bis(pyridin-2-ylmethyl)ethylenediamine (bbpen²⁻) and nitrate

Thaiane Gregório,^a André Luis Rüdiger,^a Giovana G. Nunes,^a Jaísa F. Soares^a and David L. Hughes^{b*}

Received 24 November 2014

Accepted 5 December 2014

Edited by A. J. Lough, University of Toronto, Canada

^aDepartamento de Química, Universidade Federal do Paraná, Centro Politécnico, Jardim das Américas, 81530-900 Curitiba-PR, Brazil, and ^bSchool of Chemistry, University of East Anglia, Norwich NR4 7TJ, England. *Correspondence e-mail: d.l.hughes@uea.ac.uk

Keywords: crystal structure; lanthanide; terbium(III); *N,N'*-bis(2-hydroxybenzyl)-*N,N'*-bis(pyridin-2-ylmethyl)ethylenediamine; mononuclear; dodecahedral.


CCDC reference: 1037922

Supporting information: this article has supporting information at journals.iucr.org/e


The reaction of terbium(III) nitrate pentahydrate in acetonitrile with *N,N'*-bis(2-hydroxybenzyl)-*N,N'*-bis(pyridin-2-ylmethyl)ethylenediamine (H₂bbpen), previously deprotonated with triethylamine, produced the mononuclear compound [*N,N'*-bis(2-oxidobenzyl- κ O)-*N,N'*-bis(pyridin-2-ylmethyl- κ N)ethylenediamine- κ^2 N,N'](nitrate- κ^2 O,O')terbium(III), [Tb(C₂₈H₂₈N₄O₂)(NO₃)]. The molecule lies on a twofold rotation axis and the Tb^{III} ion is eight-coordinate with a slightly distorted dodecahedral coordination geometry. In the symmetry-unique part of the molecule, the pyridine and benzene rings are both essentially planar and form a dihedral angle of 61.42 (7)°. In the molecular structure, the N₄O₄ coordination environment is defined by the hexadentate bbpen ligand and the bidentate nitrate anion. In the crystal, a weak C—H...O hydrogen bond links molecules into a two-dimensional network parallel to (001).

1. Chemical context

As far as biological and biomedical applications are concerned, complexes of polydentate ligands with a range of metal ions in different oxidation states have been synthesized to model active sites of metalloproteins and to shed light on the consequences of heavy-metal chelation in living organisms, among many other applications (Colotti *et al.*, 2013; Nurchi *et al.*, 2013; Sears, 2013; Happe & Hemschemeier, 2014). Pyridyl and phenolate groups have been incorporated into these ligands because of their potential to mimic the coordination environments provided by the amino acids histidine and tyrosine, respectively (Hancock, 2013; Lenze *et al.*, 2013). In this context, the heterotrifunctional Lewis base *N,N'*-bis(2-hydroxybenzyl)-*N,N'*-bis(pyridin-2-ylmethyl)ethylenediamine (H₂bbpen) is suitable for the coordination of a range of *p*-, *d*- and *f*-block ions because of its versatile soft donor atoms in the pyridine rings and hard donors in the amine and phenolate groups (Neves *et al.*, 1992; Schwingel *et al.*, 1996). Electrochemical studies of the mononuclear [Mn(bbpen)]PF₆, for example, revealed that this complex mimics some of the redox features of the photosystem II (PSII) (Neves *et al.*, 1992). Complexes of bbpen²⁻ with vanadium(III) and oxido-vanadium(IV) have been obtained as models of the vanadium-modified transferrin, the probable vanadium-transporting protein in higher organisms (Neves *et al.*, 1991, 1993). Iron complexes of bbpen²⁻ modified with electron-donating and -withdrawing groups (Me, Br, NO₂), in turn, have been



synthesized to provide detailed chemical information on the enzymatic activity of iron-tyrosinate proteins (Lanzaster *et al.*, 2006). This ligand has also been employed to prepare lanthanide(III), gallium(III) and indium(III) complexes for medicinal applications such as the development of new contrast agents for magnetic resonance imaging, MRI (Wong *et al.*, 1995, 1996; Setyawati *et al.*, 2000).



More recently, lanthanide(III) chelate complexes have also attracted attention in the field of molecular magnetism due to their highly significant single-ion magnetic anisotropy (Sessoli & Powell, 2009; Luzon & Sessoli, 2012). Accordingly, a number of examples of mononuclear lanthanide complexes that exhibit single-molecule magnet (SMM) behaviour have been reported (Rinehart & Long, 2011; Chilton *et al.*, 2013;




Figure 1
View of a molecule of $[\text{Tb}(\text{bbpen})(\text{NO}_3)]$, indicating the atom-numbering scheme. H atoms have been omitted for clarity. Displacement ellipsoids are drawn at the 50% probability level [symmetry code: (i) $-x + 1, y, -z + \frac{1}{2}$].

Table 1
Selected bond lengths (\AA).

Tb1—O1	2.1947 (13)	Tb1—N3	2.5558 (16)
Tb1—O2	2.4764 (15)	Tb1—N1	2.891 (2)
Tb1—N2	2.5521 (17)		

Table 2
Hydrogen-bond geometry ($\text{\AA}, ^\circ$).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C8}-\text{H8B}\cdots\text{O3}^{\text{ii}}$	0.99	2.37	3.338 (3)	166

Symmetry code: (ii) $x + \frac{1}{2}, y + \frac{1}{2}, z$.

Ungur *et al.*, 2014; Zhang *et al.*, 2014). Our interest in the class of lanthanide complexes in which two coordination sites are occupied by relatively labile ligands, as in the title complex, comes from the possibility of using them as starting materials for the preparation of heteronuclear aggregates of d - and f -block ions that present SMM features. In this case, the replacement of the labile ligands by specific bidentate metalloligands can give rise to heteronuclear metal aggregates in which desirable ferromagnetic or ferrimagnetic exchange interactions are favoured (Totaro *et al.*, 2013; Westrup *et al.*, 2014).

2. Structural commentary

The molecular structure of the title compound is shown in Fig. 1. The Tb^{III} ion is eight-coordinate with a dodecahedral array of N and O atoms (Table 1); the four N atoms of the O_2N_4 -ligand (bbpen) form one plane, the four O atoms the other, with the phenolic O atoms in the B-sites (roughly equatorial) and the nitrate group O atoms in the A-sites (above and below the equatorial plane). The normals to the two planes are essentially perpendicular. A twofold rotation axis passes through O3 and N1 of the nitrate group, the terbium(III) atom and the mid-point of the C7—C7ⁱ bond [symmetry code (i) $1 - x, y, -z + \frac{1}{2}$]. In the symmetry-unique part of the molecule, the pyridine and benzene rings are both essentially planar and form a dihedral angle of $61.42(7)^\circ$. The eightfold coordination pattern might also be described as a distorted bicapped trigonal prism with O1 and N2 as the capping atoms. However, this ignores the symmetry of the coordination, e.g. O1 and O1ⁱ would occupy different sites in the coordination polyhedron. Also, some of the rectangular faces of the prism are difficult to identify. In contrast, the dodecahedral pattern incorporates the twofold symmetry and the distortion from the ideal geometry is minimal.

3. Supramolecular features

In the crystal, a weak $\text{C}-\text{H}\cdots\text{O}$ hydrogen bond (Table 2) links molecules into a two-dimensional network parallel to (001), Fig. 2.




Figure 2
A sheet of molecules, lying in a plane normal to the *c* axis, linked through short 'weak hydrogen bonds', as C8—H8B···O3ⁱⁱⁱ [symmetry codes: (ii) $x + \frac{1}{2}, y + \frac{1}{2}, z$; (iii) $x - \frac{1}{2}, y - \frac{1}{2}, z$].

4. Database survey

Some examples of complexes with bbpen^{2-} and related ligands with *d*-block metal ions appear in the literature (Xu *et al.*, 2000; dos Anjos *et al.*, 2006; Lanznaster *et al.*, 2006; Golchoubian & Gholamnezhad, 2009; Thomas *et al.*, 2010) as well as *p*-block metal(III) compounds (Wong *et al.*, 1995, 1996) and related yttrium(III) and lanthanide(III) complexes (Setyawati *et al.*, 2000; Yamada *et al.*, 2010).

5. Synthesis and crystallization

$\text{Tb}(\text{NO}_3)_3 \cdot 5\text{H}_2\text{O}$, ethylenediamine, salicylaldehyde, sodium borohydride, 2-picoyl-chloride hydrochloride and triethylamine were purchased from Aldrich and used without purification. *N,N'*-bis(salicylidene)ethylenediamine (H_2salen) (Diehl *et al.*, 2007), *N,N'*-bis(2-hydroxybenzyl)ethylenediamine (H_2bbsen) and *N,N'*-bis(2-hydroxybenzyl)-*N,N'*-bis(2-pyridylmethyl)ethylenediamine (H_2bbpen) (Neves *et al.*, 1992) were prepared as described in the literature. The preparation of the title complex was carried out under $\text{N}_2(\text{g})$ using standard Schlenk and glove-box techniques. Acetonitrile was dried with CaH_2 and distilled prior to use. A solution containing triethylamine (300 μl , 2.15 mmol) in acetonitrile (10 ml) was added to a suspension of H_2bbpen (0.454 g, 1.00 mmol) in acetonitrile (25 ml) under stirring, giving a clear light-orange solution. After 15 min, this solution was added to a colourless solution of $\text{Tb}(\text{NO}_3)_3 \cdot 5\text{H}_2\text{O}$ (0.434 g, 0.998 mmol) in acetonitrile (25 ml). A pale-yellow solution was obtained, which gave a 65% yield of the solid of the title compound upon cooling at 253 K for 2–3 days. Recrystallization of this

Table 3
Experimental details.

Crystal data	
Chemical formula	$[\text{Tb}(\text{C}_{28}\text{H}_{28}\text{N}_4\text{O}_2)(\text{NO}_3)]$
M_r	673.47
Crystal system, space group	Orthorhombic, $C22_1$
Temperature (K)	100
<i>a</i> , <i>b</i> , <i>c</i> (Å)	8.5947 (6), 18.2401 (17), 16.9272 (13)
<i>V</i> (Å ³)	2653.6 (4)
<i>Z</i>	4
Radiation type	Mo <i>K</i> α
μ (mm ⁻¹)	2.71
Crystal size (mm)	0.43 × 0.20 × 0.20
Data collection	
Diffractometer	Bruker D8 Venture/Photon 100 CMOS
Absorption correction	Multi-scan (<i>SADABS2014/2</i> ; Bruker, 2014)
T_{min} , T_{max}	0.581, 0.746
No. of measured, independent and observed [$I > 2\sigma(I)$] reflections	75009, 3320, 3289
R_{int}	0.020
$(\sin \theta/\lambda)_{\text{max}}$ (Å ⁻¹)	0.668
Refinement	
$R[F^2 > 2\sigma(F^2)]$, $wR(F^2)$, <i>S</i>	0.010, 0.027, 1.15
No. of reflections	3320
No. of parameters	178
H-atom treatment	H-atom parameters constrained
$\Delta\rho_{\text{max}}$, $\Delta\rho_{\text{min}}$ (e Å ⁻³)	0.87, −0.30
Absolute structure	Flack <i>x</i> determined using 1431 quotients $[(I^+) - (I^-)] / [(I^+) + (I^-)]$ (Parsons & Flack, 2004)
Absolute structure parameter	−0.0107 (19)

Computer programs: *APEX2* and *SAINT* (Bruker, 2010), *SHELXS97* and *SHELXL2013* (Sheldrick, 2008), *ORTEPII* (Johnson, 1976), *ORTEP-3 for Windows* and *WinGX* (Farrugia, 2012).

solid by vapor diffusion of dimethoxyethane into the reaction mixture gave pale-pink crystals after two weeks at room temperature. These crystals are air-stable and insoluble in all common organic solvents.

6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 3. Hydrogen atoms were included in idealized positions (with C—H distances set at 0.97 and 0.93 Å for the methylene and trigonal-planar groups, respectively) and their U_{iso} values were set to ride (1.2×) on the U_{eq} values of the parent carbon atoms.

Acknowledgements

Financial support from the Brazilian agencies CNPq (grant No. 307592/2012–0) and CAPES (grant PVE A099/2013) is gratefully acknowledged. The authors also thank CNPq, CAPES and Fundação Araucária (Brazil) for fellowships.

References

- Anjos, A. dos, Bortoluzzi, A. J., Caro, M. S. B., Peralta, R. A., Friedermann, G. R., Mangrich, A. S. & Neves, A. (2006). *J. Braz. Chem. Soc.* **17**, 1540–1550.

- Bruker (2010). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2014). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chilton, N. F., Langley, S. K., Moubaraki, B., Soncini, A., Batten, S. R. & Murray, K. S. (2013). *Chem. Sci.* **4**, 1719–1730.
- Colotti, G., Ilari, A., Boffi, A. & Morea, V. (2013). *Mini Rev. Med. Chem.* **13**, 211–221.
- Diehl, H., Hach, C. C. & Bailar, J. C. (2007). *Inorganic Synthesis*, pp. 196–201. New York: John Wiley & Sons Inc.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Golchoubian, H. & Gholamnezhad, P. (2009). *X-Ray Struct. Anal. Online*, **25**, 95–96.
- Hancock, R. D. (2013). *Chem. Soc. Rev.* **42**, 1500–1524.
- Happe, T. & Hemschemeier, A. (2014). *Trends Biotechnol.* **32**, 170–176.
- Johnson, C. K. (1976). *ORTEP II*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
- Lanznaster, M., Neves, A., Bortoluzzi, A. J., Assumpção, A. M. C., Vencato, I., Machado, S. P. & Drechsel, S. M. (2006). *Inorg. Chem.* **45**, 1005–1011.
- Lenze, M., Sedinkin, S. L. & Bauer, E. B. (2013). *J. Mol. Catal. A Chem.* **373**, 161–171.
- Luzon, J. & Sessoli, R. (2012). *Dalton Trans.* **41**, 13556–13567.
- Neves, A., Ceccato, A. S., Erthal, S. M. D., Vencato, I., Nuber, B. & Weiss, J. (1991). *Inorg. Chim. Acta*, **187**, 119–121.
- Neves, A., Ceccato, A. S., Erasmus-Buhr, C., Gehring, S., Haase, W., Paulus, H., Nascimento, O. R. & Batista, A. A. (1993). *J. Chem. Soc. Chem. Commun.* pp. 1782–1784.
- Neves, A., Erthal, S. M. D., Vencato, I., Ceccato, A. S., Mascarenhas, Y. P., Nascimento, O. R., Horner, M. & Batista, A. A. (1992). *Inorg. Chem.* **31**, 4749–4755.
- Nurchi, V. M., Crespo-Alonso, M., Toso, L., Lachowicz, J. I. & Crisponi, G. (2013). *Mini Rev. Med. Chem.* **13**, 1541–1549.
- Parsons, S. & Flack, H. (2004). *Acta Cryst. A* **60**, s61.
- Rinehart, J. D. & Long, J. R. (2011). *Chem. Sci.* **2**, 2078–2085.
- Schwingel, E. W., Arend, K., Zarling, J., Neves, A. & Szpoganicz, B. (1996). *J. Braz. Chem. Soc.* **7**, 31–37.
- Sears, M. E. (2013). *Sci. World J.*, article no. 219840.
- Sessoli, R. & Powell, A. K. (2009). *Coord. Chem. Rev.* **253**, 2328–2341.
- Setyawati, I. A., Liu, S., Rettig, S. J. & Orvig, C. (2000). *Inorg. Chem.* **39**, 496–507.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Thomas, F., Arora, H., Philouze, C. & Jarjays, O. (2010). *Inorg. Chim. Acta*, **363**, 3122–3130.
- Totaro, P., Westrup, K. C. M., Boulon, M.-E., Nunes, G. G., Back, D. F., Barison, A., Ciattini, S., Mannini, M., Sorace, L., Soares, J. F., Cornia, A. & Sessoli, R. (2013). *Dalton Trans.* **42**, 4416–4426.
- Ungur, L., Le Roy, J. J., Korobkov, I., Murugesu, M. & Chibotaru, L. F. (2014). *Angew. Chem. Int. Ed.* **53**, 4413–4417.
- Westrup, K. C. M., Boulon, M.-E., Totaro, P., Nunes, G. G., Back, D. F., Barison, A., Jackson, M., Paulsen, C., Gatteschi, D., Sorace, L., Cornia, A., Soares, J. F. & Sessoli, R. (2014). *Chem. Eur. J.* **20**, 13681–13691.
- Wong, E., Caravan, P., Liu, S., Rettig, S. J. & Orvig, C. (1996). *Inorg. Chem.* **35**, 715–724.
- Wong, E., Liu, S., Rettig, S. & Orvig, C. (1995). *Inorg. Chem.* **34**, 3057–3064.
- Xu, L., Setyawati, I. A., Pierrero, J., Pink, M., Young, V. G., Patrick, B. O., Rettig, S. J. & Orvig, C. (2000). *Inorg. Chem.* **39**, 5958–5963.
- Yamada, Y., Takenouchi, S. I., Miyoshi, Y. & Okamoto, K. I. (2010). *J. Coord. Chem.* **63**, 996–1012.
- Zhang, P., Zhang, L., Wang, C., Xue, S. F., Lin, S. Y. & Tang, J. K. (2014). *J. Am. Chem. Soc.* **136**, 4484–4487.

supporting information

Acta Cryst. (2015). E71, 65-68 [doi:10.1107/S2056989014026826]

Crystal structure of an eight-coordinate terbium(III) ion chelated by *N,N'*-bis(2-hydroxybenzyl)-*N,N'*-bis(pyridin-2-ylmethyl)ethylenediamine (bbpen²⁻) and nitrate

Thaiane Gregório, André Luis Rüdiger, Giovana G. Nunes, Jaísa F. Soares and David L. Hughes

Computing details

Data collection: *APEX2* (Bruker, 2010); cell refinement: *SAINTE* (Bruker, 2010); data reduction: *SAINTE* (Bruker, 2010); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2013* (Sheldrick, 2008); molecular graphics: *ORTEPII* (Johnson, 1976) and *ORTEP-3 for Windows* (Farrugia, 2012); software used to prepare material for publication: *SHELXL2013* (Sheldrick, 2008) and *WinGX* (Farrugia, 2012).

[*N,N'*-Bis(2-oxidobenzyl- κ O)-*N,N'*-bis(pyridin-2-ylmethyl- κ N)ethylenediamine- κ^2 N,N'](nitrate- κ^2 O,O')terbium(III)

Crystal data

[Tb(C₂₈H₂₈N₄O₂)(NO₃)]
M_r = 673.47
 Orthorhombic, C222₁
a = 8.5947 (6) Å
b = 18.2401 (17) Å
c = 16.9272 (13) Å
V = 2653.6 (4) Å³
Z = 4
F(000) = 1344

D_x = 1.686 Mg m⁻³
 Mo *K*α radiation, λ = 0.71073 Å
 Cell parameters from 9558 reflections
 θ = 2.9–28.3°
 μ = 2.71 mm⁻¹
T = 100 K
 Prism, pale pink
 0.43 × 0.20 × 0.20 mm

Data collection

Bruker D8 Venture/Photon 100 CMOS
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 Detector resolution: 10.4167 pixels mm⁻¹
 φ and ω scans
 Absorption correction: multi-scan
 (*SADABS2014/2*; Bruker, 2014)
T_{min} = 0.581, *T_{max}* = 0.746

75009 measured reflections
 3320 independent reflections
 3289 reflections with *I* > 2σ(*I*)
R_{int} = 0.020
 θ_{max} = 28.4°, θ_{min} = 2.9°
h = -11→11
k = -24→24
l = -22→22

Refinement

Refinement on *F*²
 Least-squares matrix: full
R[*F*² > 2σ(*F*²)] = 0.010
wR(*F*²) = 0.027
S = 1.15
 3320 reflections

178 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods
 Hydrogen site location: inferred from
 neighbouring sites

H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0139P)^2 + 1.0428P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.004$
 $\Delta\rho_{\max} = 0.87 \text{ e } \text{Å}^{-3}$

$\Delta\rho_{\min} = -0.30 \text{ e } \text{Å}^{-3}$
 Absolute structure: Flack x determined using
 1431 quotients $[(F^+)-(F^-)]/[(F^+)+(F^-)]$ (Parsons &
 Flack, 2004)
 Absolute structure parameter: -0.0107 (19)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Tb1	0.5000	0.51826 (2)	0.2500	0.01172 (4)
O1	0.59854 (16)	0.54526 (8)	0.13398 (8)	0.0167 (3)
O2	0.4357 (2)	0.39605 (9)	0.30473 (10)	0.0312 (4)
O3	0.5000	0.29285 (11)	0.2500	0.0586 (9)
N1	0.5000	0.35974 (11)	0.2500	0.0293 (6)
N2	0.7540 (2)	0.49068 (9)	0.32221 (10)	0.0183 (3)
N3	0.66305 (19)	0.63257 (8)	0.27777 (9)	0.0130 (3)
C1	0.8272 (3)	0.42575 (12)	0.32538 (14)	0.0249 (4)
H1	0.7809	0.3851	0.2993	0.030*
C2	0.9672 (2)	0.41490 (13)	0.36483 (14)	0.0275 (5)
H2	1.0155	0.3681	0.3656	0.033*
C3	1.0342 (2)	0.47389 (13)	0.40282 (14)	0.0253 (5)
H3	1.1304	0.4685	0.4298	0.030*
C4	0.9590 (2)	0.54124 (13)	0.40111 (12)	0.0203 (4)
H4	1.0022	0.5823	0.4278	0.024*
C5	0.8200 (2)	0.54783 (11)	0.35990 (10)	0.0151 (3)
C6	0.7349 (2)	0.62034 (12)	0.35643 (12)	0.0160 (4)
H6A	0.6528	0.6212	0.3975	0.019*
H6B	0.8088	0.6606	0.3679	0.019*
C7	0.5652 (2)	0.69987 (10)	0.28069 (12)	0.0156 (3)
H7A	0.6323	0.7432	0.2719	0.019*
H7B	0.5190	0.7043	0.3340	0.019*
C8	0.7933 (2)	0.64051 (11)	0.21953 (12)	0.0162 (4)
H8A	0.8566	0.5952	0.2212	0.019*
H8B	0.8607	0.6814	0.2372	0.019*
C9	0.7479 (2)	0.65450 (12)	0.13510 (12)	0.0161 (4)
C10	0.6559 (2)	0.60220 (10)	0.09514 (11)	0.0155 (4)
C11	0.6273 (2)	0.61288 (12)	0.01400 (12)	0.0188 (4)
H11	0.5663	0.5782	-0.0142	0.023*
C12	0.6873 (3)	0.67359 (13)	-0.02515 (12)	0.0232 (4)
H12	0.6664	0.6801	-0.0798	0.028*
C13	0.7777 (3)	0.72492 (13)	0.01476 (14)	0.0251 (4)
H13	0.8186	0.7663	-0.0123	0.030*
C14	0.8073 (2)	0.71496 (11)	0.09491 (12)	0.0199 (4)

H14 0.8688 0.7498 0.1225 0.024*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Tb1	0.01260 (5)	0.01031 (5)	0.01225 (5)	0.000	0.00008 (7)	0.000
O1	0.0177 (6)	0.0180 (6)	0.0145 (6)	-0.0021 (5)	0.0023 (5)	-0.0009 (5)
O2	0.0439 (9)	0.0209 (7)	0.0287 (8)	-0.0086 (7)	-0.0077 (7)	0.0083 (6)
O3	0.096 (2)	0.0096 (8)	0.0704 (19)	0.000	-0.046 (3)	0.000
N1	0.0418 (14)	0.0119 (9)	0.0343 (13)	0.000	-0.028 (2)	0.000
N2	0.0170 (8)	0.0191 (8)	0.0188 (8)	0.0028 (7)	-0.0021 (6)	0.0002 (6)
N3	0.0124 (7)	0.0136 (7)	0.0128 (6)	0.0003 (6)	-0.0002 (5)	0.0006 (5)
C1	0.0244 (10)	0.0208 (10)	0.0294 (11)	0.0046 (8)	-0.0059 (9)	-0.0027 (8)
C2	0.0250 (14)	0.0275 (10)	0.0298 (10)	0.0106 (8)	-0.0037 (8)	0.0042 (8)
C3	0.0180 (13)	0.0364 (11)	0.0214 (9)	0.0023 (8)	-0.0036 (7)	0.0102 (8)
C4	0.0176 (11)	0.0282 (10)	0.0153 (8)	-0.0033 (7)	-0.0028 (6)	0.0061 (8)
C5	0.0147 (8)	0.0196 (9)	0.0110 (8)	0.0001 (7)	0.0013 (6)	0.0032 (7)
C6	0.0162 (9)	0.0181 (9)	0.0138 (9)	-0.0009 (8)	-0.0022 (7)	-0.0010 (8)
C7	0.0161 (8)	0.0110 (8)	0.0198 (8)	-0.0006 (7)	-0.0002 (7)	-0.0012 (7)
C8	0.0124 (8)	0.0210 (9)	0.0153 (8)	-0.0023 (7)	0.0007 (7)	0.0021 (7)
C9	0.0136 (9)	0.0209 (10)	0.0139 (9)	0.0017 (8)	0.0009 (7)	0.0012 (8)
C10	0.0130 (8)	0.0175 (9)	0.0158 (8)	0.0028 (7)	0.0024 (7)	0.0001 (7)
C11	0.0181 (9)	0.0231 (10)	0.0153 (9)	0.0024 (8)	0.0000 (7)	-0.0021 (8)
C12	0.0266 (11)	0.0286 (11)	0.0142 (9)	0.0029 (9)	-0.0004 (8)	0.0042 (8)
C13	0.0303 (11)	0.0238 (10)	0.0210 (11)	-0.0032 (9)	0.0013 (9)	0.0080 (8)
C14	0.0196 (9)	0.0208 (9)	0.0194 (10)	-0.0012 (8)	0.0002 (8)	0.0022 (8)

Geometric parameters (Å, °)

Tb1—O1 ⁱ	2.1947 (13)	C3—H3	0.9500
Tb1—O1	2.1947 (13)	C4—C5	1.389 (3)
Tb1—O2 ⁱ	2.4764 (15)	C4—H4	0.9500
Tb1—O2	2.4764 (15)	C5—C6	1.513 (3)
Tb1—N2 ⁱ	2.5521 (17)	C6—H6A	0.9900
Tb1—N2	2.5521 (17)	C6—H6B	0.9900
Tb1—N3 ⁱ	2.5558 (16)	C7—C7 ⁱ	1.529 (4)
Tb1—N3	2.5558 (16)	C7—H7A	0.9900
Tb1—N1	2.891 (2)	C7—H7B	0.9900
O1—C10	1.324 (2)	C8—C9	1.503 (3)
O2—N1	1.266 (2)	C8—H8A	0.9900
O3—N1	1.220 (3)	C8—H8B	0.9900
N1—O2 ⁱ	1.266 (2)	C9—C14	1.393 (3)
N2—C1	1.342 (3)	C9—C10	1.412 (3)
N2—C5	1.347 (3)	C10—C11	1.409 (3)
N3—C6	1.485 (2)	C11—C12	1.390 (3)
N3—C7	1.489 (2)	C11—H11	0.9500
N3—C8	1.498 (2)	C12—C13	1.391 (3)
C1—C2	1.391 (3)	C12—H12	0.9500

C1—H1	0.9500	C13—C14	1.392 (3)
C2—C3	1.380 (3)	C13—H13	0.9500
C2—H2	0.9500	C14—H14	0.9500
C3—C4	1.388 (3)		
O1 ⁱ —Tb1—O1	154.07 (7)	C8—N3—Tb1	111.56 (11)
O1 ⁱ —Tb1—O2 ⁱ	128.52 (6)	N2—C1—C2	123.4 (2)
O1—Tb1—O2 ⁱ	77.36 (6)	N2—C1—H1	118.3
O1 ⁱ —Tb1—O2	77.36 (6)	C2—C1—H1	118.3
O1—Tb1—O2	128.52 (6)	C3—C2—C1	118.3 (2)
O2 ⁱ —Tb1—O2	51.64 (9)	C3—C2—H2	120.9
O1 ⁱ —Tb1—N2 ⁱ	98.20 (5)	C1—C2—H2	120.9
O1—Tb1—N2 ⁱ	86.89 (5)	C2—C3—C4	119.11 (19)
O2 ⁱ —Tb1—N2 ⁱ	80.45 (6)	C2—C3—H3	120.4
O2—Tb1—N2 ⁱ	79.10 (6)	C4—C3—H3	120.4
O1 ⁱ —Tb1—N2	86.89 (5)	C3—C4—C5	119.2 (2)
O1—Tb1—N2	98.20 (5)	C3—C4—H4	120.4
O2 ⁱ —Tb1—N2	79.10 (6)	C5—C4—H4	120.4
O2—Tb1—N2	80.45 (6)	N2—C5—C4	122.21 (19)
N2 ⁱ —Tb1—N2	157.26 (8)	N2—C5—C6	117.05 (16)
O1 ⁱ —Tb1—N3 ⁱ	76.70 (5)	C4—C5—C6	120.74 (19)
O1—Tb1—N3 ⁱ	82.18 (5)	N3—C6—C5	111.54 (16)
O2 ⁱ —Tb1—N3 ⁱ	141.99 (5)	N3—C6—H6A	109.3
O2—Tb1—N3 ⁱ	132.91 (6)	C5—C6—H6A	109.3
N2 ⁱ —Tb1—N3 ⁱ	66.65 (5)	N3—C6—H6B	109.3
N2—Tb1—N3 ⁱ	135.88 (5)	C5—C6—H6B	109.3
O1 ⁱ —Tb1—N3	82.18 (5)	H6A—C6—H6B	108.0
O1—Tb1—N3	76.70 (5)	N3—C7—C7 ⁱ	113.05 (13)
O2 ⁱ —Tb1—N3	132.91 (6)	N3—C7—H7A	109.0
O2—Tb1—N3	141.99 (5)	C7 ⁱ —C7—H7A	109.0
N2 ⁱ —Tb1—N3	135.88 (5)	N3—C7—H7B	109.0
N2—Tb1—N3	66.65 (5)	C7 ⁱ —C7—H7B	109.0
N3 ⁱ —Tb1—N3	70.67 (7)	H7A—C7—H7B	107.8
O1 ⁱ —Tb1—N1	102.97 (4)	N3—C8—C9	116.63 (16)
O1—Tb1—N1	102.97 (4)	N3—C8—H8A	108.1
O2 ⁱ —Tb1—N1	25.82 (4)	C9—C8—H8A	108.1
O2—Tb1—N1	25.82 (4)	N3—C8—H8B	108.1
N2 ⁱ —Tb1—N1	78.63 (4)	C9—C8—H8B	108.1
N2—Tb1—N1	78.63 (4)	H8A—C8—H8B	107.3
N3 ⁱ —Tb1—N1	144.67 (4)	C14—C9—C10	120.42 (18)
N3—Tb1—N1	144.67 (4)	C14—C9—C8	120.25 (19)
C10—O1—Tb1	139.80 (12)	C10—C9—C8	119.08 (19)
N1—O2—Tb1	95.73 (12)	O1—C10—C11	121.83 (18)
O3—N1—O2	121.55 (11)	O1—C10—C9	120.05 (17)
O3—N1—O2 ⁱ	121.55 (11)	C11—C10—C9	118.13 (18)
O2—N1—O2 ⁱ	116.9 (2)	C12—C11—C10	120.67 (19)
O3—N1—Tb1	180.0	C12—C11—H11	119.7
O2—N1—Tb1	58.45 (11)	C10—C11—H11	119.7

O2 ⁱ —N1—Tb1	58.45 (11)	C11—C12—C13	120.8 (2)
C1—N2—C5	117.83 (17)	C11—C12—H12	119.6
C1—N2—Tb1	126.43 (14)	C13—C12—H12	119.6
C5—N2—Tb1	115.74 (12)	C12—C13—C14	119.2 (2)
C6—N3—C7	109.19 (15)	C12—C13—H13	120.4
C6—N3—C8	107.09 (15)	C14—C13—H13	120.4
C7—N3—C8	111.34 (15)	C13—C14—C9	120.8 (2)
C6—N3—Tb1	105.69 (12)	C13—C14—H14	119.6
C7—N3—Tb1	111.67 (11)	C9—C14—H14	119.6
Tb1—O2—N1—O3	180.000 (1)	Tb1—N3—C7—C7 ⁱ	-38.2 (2)
Tb1—O2—N1—O2 ⁱ	0.000 (1)	C6—N3—C8—C9	-179.19 (19)
C5—N2—C1—C2	-0.5 (3)	C7—N3—C8—C9	-59.9 (2)
Tb1—N2—C1—C2	178.82 (17)	Tb1—N3—C8—C9	65.61 (19)
N2—C1—C2—C3	0.2 (4)	N3—C8—C9—C14	125.3 (2)
C1—C2—C3—C4	0.8 (3)	N3—C8—C9—C10	-60.3 (3)
C2—C3—C4—C5	-1.3 (3)	Tb1—O1—C10—C11	-142.18 (16)
C1—N2—C5—C4	0.0 (3)	Tb1—O1—C10—C9	37.5 (3)
Tb1—N2—C5—C4	-179.43 (14)	C14—C9—C10—O1	-179.29 (18)
C1—N2—C5—C6	-179.33 (18)	C8—C9—C10—O1	6.3 (3)
Tb1—N2—C5—C6	1.2 (2)	C14—C9—C10—C11	0.4 (3)
C3—C4—C5—N2	0.9 (3)	C8—C9—C10—C11	-174.00 (18)
C3—C4—C5—C6	-179.80 (18)	O1—C10—C11—C12	179.21 (19)
C7—N3—C6—C5	173.14 (16)	C9—C10—C11—C12	-0.5 (3)
C8—N3—C6—C5	-66.2 (2)	C10—C11—C12—C13	0.3 (3)
Tb1—N3—C6—C5	52.88 (17)	C11—C12—C13—C14	-0.2 (4)
N2—C5—C6—N3	-38.3 (2)	C12—C13—C14—C9	0.1 (3)
C4—C5—C6—N3	142.41 (18)	C10—C9—C14—C13	-0.2 (3)
C6—N3—C7—C7 ⁱ	-154.7 (2)	C8—C9—C14—C13	174.1 (2)
C8—N3—C7—C7 ⁱ	87.2 (2)		

Symmetry code: (i) $-x+1, y, -z+1/2$.

Hydrogen-bond geometry (Å, °)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C8—H8B ⁱⁱ ⋯O3 ⁱⁱ	0.99	2.37	3.338 (3)	166

Symmetry code: (ii) $x+1/2, y+1/2, z$.