

**Cognitive vulnerabilities for depression and anxiety in childhood: Specificity of anxiety sensitivity and rumination**

Hannah M. Brown<sup>1</sup>, Richard Meiser-Stedman<sup>2</sup>, Harriet Woods<sup>3</sup> & Kathryn J. Lester<sup>1</sup>

<sup>1</sup> MRC Social, Genetic and Developmental Psychiatry Centre, Institute of Psychiatry, London, SE5 8AF, England; [hannah.brown@kcl.ac.uk](mailto:hannah.brown@kcl.ac.uk), [kathryn.lester@kcl.ac.uk](mailto:kathryn.lester@kcl.ac.uk)

<sup>2</sup> MRC Cognition & Brain Sciences Unit, Cambridge University, Cambridge CB3 9BB, England; [Richard.Meiser-Stedman@mrc-cbu.cam.ac.uk](mailto:Richard.Meiser-Stedman@mrc-cbu.cam.ac.uk)

<sup>3</sup> Department of Psychosis Studies, Institute of Psychiatry, London, SE5 8AF, England; [harriet.woods@kcl.ac.uk](mailto:harriet.woods@kcl.ac.uk)

**Corresponding authors:**

Hannah Brown, Social, Genetic and Developmental Psychiatry Centre, Institute of Psychiatry, London, SE5 8AF, England; [hannah.brown@kcl.ac.uk](mailto:hannah.brown@kcl.ac.uk); phone: +44 (0) 20 7848 0074; fax: +44(0)20 7848 0866

Kathryn Lester, Social, Genetic and Developmental Psychiatry Centre, Institute of Psychiatry, London, SE5 8AF, England; [kathryn.lester@kcl.ac.uk](mailto:kathryn.lester@kcl.ac.uk); phone: +44 (0) 20 7848 5414; fax: +44(0)20 7848 0866

**Running Head:** Cognitive specificity anxiety depression

**Key words:** Anxiety, Depression, Cognition, Specificity

## **Cognitive vulnerabilities for depression and anxiety in childhood: Specificity of anxiety sensitivity and rumination**

**Background:** Childhood anxiety and depression frequently co-occur. Exploring specificity in cognitive processes for anxiety and depression in childhood can provide insight into cognitive vulnerabilities contributing to the development of anxiety and depressive disorders and inform targeted psychological interventions. Anxiety sensitivity and rumination are robust cognitive vulnerabilities for anxiety and depression, respectively. However, despite conceptual similarities, they are rarely considered together within a single study. **Aims:** The current study explored specific and shared associations between anxiety sensitivity subscales and rumination and anxiety and depressive symptoms in unselected children. **Methods:** Multiple regression analyses explored to what extent specific self-reported anxiety sensitivity subscales (Physical, Social and Mental concerns) and rumination predicted anxiety and depressive symptoms in 147 unselected children, aged 7 – 11 years. **Results:** Physical and Social concern subscales of anxiety sensitivity were specifically associated with anxiety whilst rumination was specifically associated with depressive symptoms. The Mental concerns subscale of anxiety sensitivity was independently associated with both anxiety and depressive symptoms. These associations were only partially mediated by rumination. **Conclusions:** Anxiety and depression in young people are characterised by specific and shared cognitions. Evidence for shared and specific associations between the cognitive vulnerabilities of anxiety sensitivity and rumination, and anxiety and depression highlight the utility of transdiagnostic research and confirm that cognitive therapies may benefit from targeting cognitive concerns relating specifically to patient's presenting symptoms.

## Background

Anxiety and depression are common in young people and frequently co-occur (Ford, Goodman, & Meltzer, 2003). Between 25 – 50% of young people with an anxiety *disorder* are also diagnosed with depression (Angold, Costello, & Erkanli, 1999) and correlations between self-reported anxiety and depressive symptoms frequently range from .40–.80 in clinical and community youth samples (Axelson & Birmaher, 2001); indicating that many youth experiencing depressive symptoms also experience elevated anxiety symptoms and vice versa.

Substantial co-occurrence of anxiety and depression in youth highlight the importance of transdiagnostic theoretical frameworks to understand their development in childhood.

Limited transdiagnostic models have proposed both shared and specific components of anxiety and depression. The tripartite model proposes that whilst anxiety and depression are both characterised by elevated negative affect, physiological hyperarousal is specific to anxiety and low positive affect is specific to depression (Clark & Watson, 1991). Similarly, the cognitive content-specificity hypothesis proposes that anxiety and depression are both characterised by cognitive distortions for processing emotional information but can be differentiated by divergent cognitive themes. Specifically, anxious cognitions are thought to be characterised by threat and danger whilst depressive cognitions focus on loss and negative self-evaluation (Clark & Beck, 1989).

Identifying shared and specific cognitions in anxiety and depressive symptoms in youth can provide insight into possible mechanisms that explain their co-occurrence. Additionally, identifying converging and diverging cognitive themes can inform cognitive behavioural therapies (CBT). CBT for both anxiety and depression focuses on modifying maladaptive cognitions maintaining emotional symptoms. Identifying converging cognitive themes

suggests a transdiagnostic treatment protocol would be effective whereas differentiated cognitive themes imply interventions specifically targeted to either anxiety or depression.

Despite important implications, research examining specificity between cognitive vulnerability factors and anxiety and depression within a single study is limited; largely separate literature bases focusing on either anxiety- or depression-related cognitions have evolved. Many cognitive vulnerabilities have been implicated in the aetiology of anxiety and depression (Clark & Beck, 1989). The cognitive of anxiety sensitivity has received considerable attention as a cognitive vulnerability for anxiety and more recently depression (Naragon-Gainey, 2010). Building on previous research, our team identified a three-factor model of anxiety sensitivity (Brown, Trzaskowski, Zavos, Rijdsdijk, Gregory, & Eley, 2012) with each dimension differentially associated with anxiety and depressive symptoms in young people (Brown, Waszczuk, Zavos, Trzaskowski, Gregory, & Eley, 2014). In the current study we extend this examination by including rumination, a robust cognitive vulnerability for depression (Nolen-Hoeksema, 1991), and examining independent associations between these cognitive constructs and anxiety and depressive symptoms.

### **Anxiety sensitivity**

Anxiety sensitivity (AS) refers to the tendency to be fearful of future anxiety symptoms with a belief they are harmful (Reiss, Peterson, Gursky, & McNally, 1986). Meta-analyses show higher AS in youth with anxiety disorders and robust associations between AS and anxiety symptoms in unselected youth (Noel & Francis, 2011). Furthermore, prospective associations between AS and later anxiety symptoms beyond current anxiety symptoms in childhood implicate AS as a vulnerability for anxiety in young people (McLaughlin, Stewart, & Taylor, 2007; Schmidt et al., 2010). However, AS has also been associated with depressive symptoms in young people (Muris, Schmidt, Merckelbach, & Schouten, 2001; Weems,

Hammond-Laurence, Silverman, & Ferguson, 1997), suggesting AS may represent a transdiagnostic cognitive vulnerability for both anxiety and depression.

Although AS was originally developed as a unitary construct, contemporary conceptualisations propose a number of AS dimensions which may each be differentially associated with anxiety and depression (Olatunji & Wolitzky-Taylor, 2009). Factor analyses generally support a multidimensional structure consisting of three subscales representing *Physical*, *Social* and *Mental* symptom-related concerns (Brown et al., 2012; Wright et al., 2010). The *Physical concerns* subscale pertains to fear of biological symptoms of anxiety (e.g. Funny feelings in my body scare me), whilst the *Social concerns* facet represents fear of publically observable symptoms (e.g. I don't want other people to know when I'm afraid) and the *Mental concerns* subscale depicts worries regarding cognitive control over distress (e.g. When I am afraid, I worry I might be going crazy). Limited studies with adults (Schmidt, Lerew, & Joiner, 1998; Taylor, Koch, Woody, & McLean, 1996) suggest that the *Physical* and *Social* concerns subscales of AS tends to be associated with anxiety but not depressive symptoms. Conversely, *Mental* AS seems more strongly associated with depression. Studies with unselected youth are scarce and reveal a mixed pattern of associations (Dehon, Weems, Stickle, Costa, & Berman, 2005; Joiner et al., 2002). For example, one study of unselected adolescents found *Physical* AS was independently associated with anxiety but not depression whilst *Social* and *Mental* AS were associated with anxiety and depression (Brown et al., 2014; Dehon et al., 2005). Similarly, a recent twin study of childhood, adolescence and early adulthood by our group showed that *Physical* AS was independently associated with anxiety at all time points (Brown et al., 2014). *Social* AS was not associated with anxiety or depression in childhood but was associated with both anxiety and depression in adolescence, whereas *Mental* AS was associated with both anxiety and depressive symptoms to a similar extent at all ages.

## **Rumination**

Rumination, the tendency to recurrently think about the causes and consequences of depressive symptoms is consistently implicated in the development and maintenance of depression in adults (Nolen-Hoeksema, 1991, 2000). Similarly, meta-analytic findings from studies with unselected children show robust associations with both concurrent and future depressive symptoms (Rood, Roelofs, Bogels, Nolen-Hoeksema, & Schouten, 2009). However, similar to studies of AS, limited recent studies have also identified prospective associations with anxiety in children and adolescents (Muris, Fokke, & Kwik, 2009; Roelofs et al., 2009), indicating rumination may also represent a transdiagnostic cognitive vulnerability for anxiety and depression in youth.

### **Associations between AS, rumination, anxiety and depression**

At a conceptual level, AS and rumination are similar constructs; both describing repetitive negative thoughts. However, AS reflects fear of *future* distressing symptoms whilst rumination is concerned with negative reflections on *past* symptoms of distress. Despite theoretical similarities, few studies have considered the possible interplay between AS and rumination and their associations with anxiety and depressive symptoms.

A small number of studies have demonstrated associations between rumination and the more general construct of worry in unselected adults (Fresco, Frankel, Mennin, Turk, & Heimberg, 2002; Segerstrom, Tsao, Alden, & Craske, 2000) and children (Muris, Roelofs, Meesters, & Boomsma, 2004; Verstraeten, Bijttebier, Vasey, & Raes, 2011). However, conclusions regarding their specific roles in anxiety and depression are mixed. For example, one study showed specific associations between worry and anxiety and rumination and depression (Verstraeten et al., 2011) whilst others find similar associations with anxiety and depression (Fresco et al., 2002).

With regards to AS, one study of unselected girls (aged 9–12) showed a substantial correlation between *total* AS and rumination and between them both and anxiety and depressive symptoms (Epkins, Gardner, & Scanlon, 2013). However, when controlling for covariance between anxiety and depression, AS was specifically associated with anxiety whilst rumination was specific to depression; suggesting AS and rumination may represent specific cognitive vulnerabilities for anxiety and depression, respectively. To date, no studies have examined specificity between AS *subscales* and rumination in relation to anxiety and depression in youth. One study of depressed adults found that only the *Mental concerns* AS subscale was associated with depressive symptoms but was fully mediated by rumination (Cox, Enns, & Taylor, 2001). The authors concluded that *Mental* AS may represent a broader fear of depressive symptoms ('depression sensitivity'), driving individuals to ruminate on depressive symptoms, serving to increase depressive symptom severity.

### **The current study**

The current study extends these limited studies, exploring specificity in associations between AS (total and subscales) and rumination and anxiety and depressive symptoms in a sample of unselected children. A number of hypotheses were tested. First, given substantial co-occurrence of anxiety and depressive symptoms, we expected AS and rumination to be correlated with both sets of symptoms. However, when controlling for covariance between anxiety and depression, we expected AS to be more strongly associated with anxiety and rumination to depression, in agreement with limited previous studies examining specificity of associations between AS and rumination and anxiety and depression separately (hypothesis 1). Furthermore, when controlling for covariance between AS subscales and rumination, we anticipated, in line with the cognitive content-specificity hypothesis (Beck, Brown, Steer, Eidelson, & Riskind, 1987) and previous limited studies (Brown et al., 2014; Verstraeten et al., 2011), that rumination would show a specific relation with depression whilst AS

subscales would be solely associated with anxiety; with the exception of Mental AS which we more tentatively expected to be associated with both anxiety and depression (hypothesis 2). Finally, in line with the only other study to examine specificity in associations between AS subscales, rumination and depression (Cox et al., 2001), we explored whether the association between Mental AS and depression was mediated by rumination in our child sample (hypothesis 3). We also extended this to examine whether the mediating effect of rumination was specific to depression or common across both anxiety and depressive symptoms.

## Methods

### Participants

Ethical approval was granted by the Psychiatry, Nursing and Midwifery Research Ethics Subcommittee of King's College London (ref no: PNM/10/12-10). Children aged 7 to 11 years were recruited from a mainstream primary school in Cambridgeshire. A passive consent procedure was used. Parents were sent detailed study information and given a minimum of a week to decline. Reminders were sent via e-mail by school staff. In total, 147 children (69 boys; 78 girls) were included. The mean age of the sample was 9yrs 2months (Range = 7 – 11 years). The sample described themselves as white British (59%), Indian (11%), Pakistani (4%), Chinese (2%) or 'other' (20%).

### Measures

*Anxiety Sensitivity* AS was measured using the Children's Anxiety Sensitivity Index (CASI; Silverman, Fleisig, Rabian, & Peterson, 1991); an 18-item self-report questionnaire assessing fear of anxiety sensations (e.g., "It scares me when I feel shaky"). Children rated item agreement on a 3-point scale. Responses were summed to calculate total AS scores. The

CASI shows adequate test-retest reliability and validity in unselected children (Silverman et al., 1991). AS subscales representing Physical, Social and Mental concerns were created following the most replicated factor structure of the CASI (Brown et al., 2012; Wright et al., 2010).

*Rumination* Rumination was measured using the rumination subscale of the Children's Response Styles Questionnaire (CRSQ; Abela, Brozina, & Haigh, 2002); a 13-item self-report questionnaire describing self-focused responses to depressed moods (e.g. "When I am sad, I think about how alone I feel"). Children indicated how often they experienced each item using a four-point scale. Responses were summed to create a total rumination score. The CRSQ rumination subscale shows adequate reliability and validity with children (Abela et al., 2002).

*Anxiety* Anxiety symptoms were measured using the Spence Children's Anxiety Scale (SCAS; Spence, 1998); a 38-item self-report questionnaire. Responses were measured on a four-point Likert scale and summed to create total anxiety scores. The SCAS shows good reliability and concurrent validity (Spence, 1998).

*Depression* Depressive symptoms were measured using the Short Mood and Feelings Questionnaire (SMFQ; Angold, Costello, Messer, Pickles, Winder, & Silver, 1995); a 13-item self-report measure assessing how often children experienced depressive symptoms in the previous two weeks using a three-point scale. Total depression scores were created by summing all items. The SMFQ has adequate reliability and validity in children as young as 8 (Angold et al., 1995).

## **Procedure**

Children completed the questionnaires in groups of 8 or less and were supervised by researchers and members of staff to ensure comprehension. Any children requiring supervision undertook the study individually. Questionnaires were presented in the same order for all participants (anxiety, anxiety sensitivity, depression, rumination). The school were given a monetary voucher for participating.

### **Data analysis plan**

Descriptive statistics were used to examine the means and possible sex differences in self-reported emotional symptoms and cognitive constructs.

In order to identify specificity in relations between the cognitive constructs and anxiety and depression, and account for sex differences in some variables, separate multiple regression models predicted (i) anxiety and (ii) depressive symptoms from; sex, anxiety/depressive symptoms, rumination and total AS (hypothesis 1) or AS subscales (hypothesis 2). Age was not included owing to the narrow range and lack of associations with the constructs of interest. Standardised beta co-efficients and semi-partial correlation co-efficients were calculated to estimate the magnitude of independent effects.

Finally, Preacher and Hayes' (2008) SPSS 'PROCESS' macro for mediation was used to examine whether associations between *Mental* AS and anxiety and depressive symptoms were mediated by rumination (hypothesis 3), One thousand re-samples were taken and 95% bias corrected accelerated confidence intervals were calculated. The magnitude of indirect effects was computed using bootstrapped re-sampling following Preacher and Hayes' recommendations. The proportion of the total effect mediated by rumination was calculated according to MacKinnon and Colleagues (1995).

## Results

Descriptive statistics are shown in Table 1. Internal consistency was substantial for the majority of scales. Estimates were somewhat lower for *Social* and *Mental* AS, reflecting fewer items within those scales. This is similar to other studies with unselected youth (Brown et al., 2014; Dehon et al., 2005). Mean scores for all measures were comparable to those reported in previous unselected child samples (Brown et al., 2014). Anxiety, total AS and *Physical* AS were significantly higher for females than males ( $p < .001$ ). Pearson's correlations revealed no associations with age (in months).

### [Table 1 here]

**Hypothesis 1** There were substantial associations between total AS, rumination, anxiety and depression (all  $r_s > .56$ ,  $p_s < .001$ ). Regression co-efficients for models predicting anxiety and depressive symptoms are shown in columns 1 and 2 (respectively) of Table 2. Sex significantly predicted anxiety ( $\beta$  (standardised beta) = .88,  $p < .01$ ) but not depression ( $\beta = -.08$ ,  $ns$ ). Total AS and rumination were significant predictors of both anxiety and depression ( $\beta_s = .25 - .56$ ,  $p_s < .01$ ). However, semi-partial correlation coefficients indicated that anxiety was more strongly correlated with AS than rumination, whilst depression was more strongly associated with rumination. Interestingly, anxiety and depression were not related when controlling for AS and rumination.

**Hypothesis 2** Next AS *subscales* were included to examine specificity between AS subscales and rumination in predicting anxiety and depression (columns 3 and 4 of Table 2). *Physical* and *Social* AS showed specific relations to anxiety symptoms ( $\beta_s = .27$  and  $.16$ , respectively). Rumination was specific to depressive symptoms ( $\beta = .40$ ). *Mental* AS showed independent associations with both anxiety and depression ( $\beta_s = .34$  and  $.24$ , respectively). Again, anxiety and depression were unrelated.

[Table 2 here]

**Hypothesis 3** Mediation analyses revealed that *Mental AS* significantly predicted rumination (*a* paths); explaining 32.34% of the variance in rumination (figure 1)<sup>1</sup>. The direct effects of rumination (*b* paths) and *Mental AS* (*c* path) significantly predicted anxiety and depression; explaining 55.44% and 57.04% of the variance, respectively. The significant effect of *Mental AS* on anxiety and depressive symptoms (*c* paths) reduced but remained significant when controlling for rumination (*c'* paths), indicating partial mediation. There were significant, large indirect effects of *Mental AS*, via rumination on depression ( $b = 1.17$  [.69, 1.77],  $p < .001$ ;  $\kappa^2 = .32$  [.21, .45]) and anxiety ( $b = 2.52$  [1.50, 3.74],  $p < .001$ ;  $\kappa^2 = .26$  [.16, .37]). Children's rumination scores accounted for 25% of the association between *Mental AS* and anxiety scores (34.95% of the total effect) and 31% of the association with depression scores (52.5% of the total effect).

[Figure 1 here]

### Discussion

To our knowledge this is the first study to combine AS subscales and rumination and examine specificity of associations with both anxiety and depressive symptoms in children. As expected, we found substantial inter-correlations between all measures, in line with the high co-occurrence of anxiety and depression (Axelson & Birmaher, 2001) and theoretical similarities between anxiety- and depression-related cognitive vulnerabilities (Clark & Beck, 1989). However, in line with previous studies exploring these cognitive vulnerabilities

---

<sup>1</sup> Mediation analyses were repeated with sex and either anxiety or depressive symptoms included as covariates. This showed the same pattern of results with reduced, but still significant beta coefficients. Unfortunately, at the time of writing the SPSS PROCESS macro cannot calculate the effect size of the indirect effect when covariates are included in the model, therefore only the full model is reported. Further details are available on request from the first author.

separately (Muris et al., 2009; Weems et al., 1997), regression analyses revealed stronger associations between AS and anxiety, and between rumination and depression.

The current findings extend previous studies by exploring specificity in associations between AS *subscales* and rumination and anxiety and depression, controlling for their covariance, within a single study. Rumination was associated with depression but not anxiety. The *Physical* and *Social* AS subscales were specifically associated with anxiety whilst *Mental* AS was independently associated with anxiety and depression. Rumination partially mediated associations between *Mental* AS and anxiety and particularly depression. Anxiety was not correlated with depression when controlling for covariance with rumination and AS subscales, suggesting that shared cognitive vulnerabilities explained much of the association between anxiety and depression in the current study.

### **Content-specificity in anxiety and depression**

Results were largely in line with a recent study examining specificity between AS subscales and anxiety and depressive symptom reports in children and adolescents (Brown et al., 2014) and provided partial support for the content-specificity hypothesis (Beck et al., 1987). Specifically, anxiety but not depression was significantly associated with fear of biological and social symptoms of distress and their perceived threatening consequences (e.g. “When my stomach hurts, I worry that I might be really sick”). Conversely, depression but not anxiety was characterised by repetitive negative reflection on the causes and consequences of symptoms of distress (e.g. “when I am sad, I think about all my failures”). However, both anxiety and depression were independently associated with the *Mental* AS subscale, suggesting that worries relating to loss of cognitive control (e.g. lack of concentration, going crazy) are characteristic of both anxiety and depression. Evidence for an independent association between *Mental* AS and depression extends previous studies showing associations

between total AS and depression, suggesting associations are driven by fears relating to losing cognitive control rather than worries regarding physical or social symptoms.

Furthermore, this association was only partially mediated by rumination, suggesting depression is not only characterised by ruminative thoughts about *past* distress but also fear of cognitive symptoms of distress in the *future*.

It has previously been suggested that the *Mental* concerns subscale of AS may reflect a broader ‘depression sensitivity’ whereby normal, although unpleasant, experiences such as fatigue or difficulty concentrating are interpreted as signs of future depressive episodes (Cox et al., 2001). Sensitivity to symptoms of depression and the belief they will cause future distress is in line with a ‘scar hypothesis’ positing that chronic depression results in enduring cognitive changes that leave an individual prone to future depressive episodes (Lewinsohn, Steinmetz, Larson, & Franklin, 1981). It could be that the tendency to ruminate on depressive symptoms could reinforce cognitive distortions about the impact of depressed mood and increase sensitivity to future symptoms.

Results of the mediation analysis are somewhat at odds with the only other study to explore associations between rumination and *Mental* AS in clinically depressed adults which showed complete mediation of the association between *Mental* AS and depression (Cox et al., 2001). Discrepancies are not easily explained given the paucity of relevant research. It is possible that these cognitive vulnerabilities play different roles in children or non-clinical samples or perhaps AS acts through other mediators to influence depression (e.g. memory or attentional biases which guide attention to negative stimuli). Extending previous analyses (Cox et al., 2001), we found that rumination also partially mediated the association between *Mental* AS and anxiety, suggesting broader interplay between AS subscales and rumination and their role in childhood anxiety and depression. Prospective research is needed to examine the causal relations between *Mental* AS and rumination and anxiety and depression.

## Limitations

The main study limitation is the cross-sectional design. Although AS and rumination are hypothesised as cognitive vulnerabilities for anxiety and depression and prospective studies support their causal role, causal associations could not be examined in the current study. Mediation analyses should be interpreted with caution and future longitudinal research is needed to clarify causal relationships. Second, the sample focused on a relatively narrow age range, precluding analysis of age-effects. Future research with broader age ranges and/or larger samples are needed to assess the generalisability of current findings. Third, the study relied on children's self-report of emotional symptoms and cognition. There is some debate regarding the validity of children's reports of such internal processes. However, experimental studies have shown that children as young as 7 years can make valid interpretations of anxiety symptoms (Mattis & Ollendick, 1997; Muris, Hovee, Meesters, & Mayer, 2004) and can reliably report their emotional experiences over time (Michael & Merrell, 1998). Furthermore, all the selected measures have been validated in children in the current age range and, with the exception of the Mental and Social AS subscales, all measures demonstrated adequate internal consistency in the current, suggesting children were responding consistently within them. Reliance on self-report measures however, may inflate correlations due to common method variance. Future research may benefit from combining self- and/or parent-report with experimental measures. Finally, the lower internal consistency of the *Social* and *Mental* AS subscales is in line with other studies using AS subscales and reflects the small number of items in each scale but could reduce power to detect associations. Future research would benefit from expanded measures of AS, better capturing AS dimensions. Additionally, rumination has been conceptualised as two dimensions; 'brooding' and 'reflective' rumination (Verstraeten et al., 2011). The CRSQ rumination

subscale largely reflects brooding rumination. It would be interesting to explore specificity with regards to both domains.

### **Implications**

Substantial covariation between AS and rumination and anxiety and depressive symptoms highlights the importance of integrative cognitive models of anxiety and depression and transdiagnostic research to examine their differential development (Harvey, Watkins, Mansell, & Shafran, 2004). Identifying disorder-specific cognitions, as well as those shared between anxiety and depression could help to tailor CBT programmes to a given diagnosis. Specifically, reflecting on and reappraising past symptoms of distress is likely to be more applicable in treating depression whilst targeting fears surrounding future symptoms may be more important in anxiety treatment. However, anxiety is heterogeneous. Although not possible in the current study owing to the relatively small sample, future research would benefit from exploring associations between these cognitive vulnerabilities and anxiety symptom subtypes to tailor cognitive interventions more precisely to specific anxiety disorders.

### **Conclusions**

The current findings extend those from previous studies examining associations with AS and rumination separately. Findings suggested that childhood anxiety and depressive symptoms are characterised by both shared and symptom-specific cognitions. Specifically, anxiety but not depression was characterised by concerns relating to physical and social symptoms of distress whilst depression was specifically associated with rumination. Conversely, concerns relating to cognitive symptoms of distress were common to both depression and anxiety. The degree of covariance between rumination, AS and both anxiety and depression highlights the importance of transdiagnostic research, combining multiple measures of cognitive

vulnerabilities and examining specificity to anxiety and depression to inform theoretical frameworks and clinical interventions for these common problems.

- Abela, J. Z., Brozina, K., & Haigh, E. (2002). An Examination of the Response Styles Theory of Depression in Third- and Seventh-Grade Children: A Short-Term Longitudinal Study. *Journal of Abnormal Child Psychology, 30*(5), 515-527.
- Angold, A., Costello, E. J., & Erkanli, A. (1999). Comorbidity. *Journal of Child Psychology and Psychiatry, 40*(1), 57-87.
- Angold, A., Costello, E. J., Messer, S. C., Pickles, A., Winder, F., & Silver, D. (1995). The development of a short questionnaire for use in epidemiological studies of depression in children and adolescents. *International journal of methods in psychiatric research, 5*, 1-12.
- Axelson, D. A., & Birmaher, B. (2001). Relation between anxiety and depressive disorders in childhood and adolescence. *Depression and Anxiety, 14*(2), 67-78.
- Beck, A. T., Brown, G., Steer, R. A., Eidelson, J. I., & Riskind, J. H. (1987). Differentiating anxiety and depression: a test of the cognitive content-specificity hypothesis. *Journal of Abnormal Psychology, 96*(3), 179.
- Brown, H. M., Trzaskowski, M., Zavos, H. M. S., Rijdsdijk, F. V., Gregory, A. M., & Eley, T. C. (2012). Phenotypic and genetic structure of anxiety sensitivity in adolescence and early adulthood. *Journal of Anxiety Disorders, 26*, 680-688.
- Brown, H. M., Waszczuk, M. A., Zavos, H. M. S., Trzaskowski, M., Gregory, A. M., & Eley, T. C. (2014). Cognitive content specificity in anxiety and depressive disorder symptoms: a twin study of cross-sectional associations with anxiety sensitivity dimensions across development. *Psychological Medicine, FirstView*, 1-12.
- Clark, D. A., & Beck, A. T. (1989). Cognitive theory and therapy of anxiety and depression. In P. C. Kendall & D. Watson (Eds.), *Anxiety and depression: Distinctive and overlapping features* (pp. 379-411). San Diego: Academic Press.
- Clark, L. A., & Watson, D. (1991). Tripartite model of anxiety and depression: psychometric evidence and taxonomic implications. *Journal of Abnormal Psychology, 100*(3), 316.
- Cox, B. J., Enns, M. W., & Taylor, S. (2001). The effect of rumination as a mediator of elevated anxiety sensitivity in major depression. *Cognitive therapy and research, 25*(5), 525-534.

- Dehon, C., Weems, C. F., Stickle, T. R., Costa, N. M., & Berman, S. L. (2005). A cross-sectional evaluation of the factorial invariance of anxiety sensitivity in adolescents and young adults. *Behaviour Research and Therapy, 43*(6), 799-810.
- Epkins, C. C., Gardner, C., & Scanlon, N. (2013). Rumination and Anxiety Sensitivity in Preadolescent Girls: Independent, Combined, and Specific Associations with Depressive and Anxiety Symptoms. *Journal of psychopathology and behavioral assessment, 1-12*.
- Ford, T., Goodman, R., & Meltzer, H. (2003). The British Child and Adolescent Mental Health Survey 1999: The Prevalence of DSM-IV Disorders. *Journal of the American Academy of Child and Adolescent Psychiatry, 42*(10), 1203-1211.
- Fresco, D., Frankel, A., Mennin, D., Turk, C., & Heimberg, R. (2002). Distinct and Overlapping Features of Rumination and Worry: The Relationship of Cognitive Production to Negative Affective States. *Cognitive therapy and research, 26*(2), 179-188.  
doi:10.1023/a:1014517718949
- Harvey, A., Watkins, E., Mansell, W., & Shafran, R. (2004). *Cognitive behavioural processes across psychological disorders: A transdiagnostic approach to research and treatment*. Oxford: Oxford University Press.
- Joiner, T. E., Schmidt, N. B., Schmidt, K. L., Laurent, J., Catanzaro, S. J., Perez, M., et al. (2002). Anxiety sensitivity as a specific and unique marker of anxious symptoms in youth psychiatric inpatients. *Journal of Abnormal Child Psychology, 30*(2), 167-175.  
doi:http://dx.doi.org/10.1023/A:1014757300294
- Lewinsohn, P. M., Steinmetz, J. L., Larson, D. W., & Franklin, J. (1981). Depression-related cognitions: Antecedent or consequence? *Journal of Abnormal Psychology, 90*(3), 213.
- MacKinnon, D. P., Warsi, G., & Dwyer, J. H. (1995). A simulation study of mediated effect measures. *Multivariate behavioral research, 30*(1), 41-62.
- Mattis, S. G., & Ollendick, T. H. (1997). Children's cognitive responses to the somatic symptoms of panic. *Journal of Abnormal Child Psychology, 25*(1), 47-57.

- McLaughlin, E. N., Stewart, S. H., & Taylor, S. (2007). Childhood Anxiety Sensitivity Index Factors Predict Unique Variance in DSM-IV Anxiety Disorder Symptoms. *Cognitive Behaviour Therapy, 36*(4), 210-219.
- Michael, K. D., & Merrell, K. W. (1998). Reliability of children's self-reported internalizing symptoms over short to medium-length time intervals. *J Am Acad Child Adolesc Psychiatry, 37*(2), 194-201.
- Muris, P., Dreessen, L., Bögels, S., Weckx, M., & van Melick, M. (2004). A questionnaire for screening a broad range of DSM-defined anxiety disorder symptoms in clinically referred children and adolescents. *Journal of Child Psychology and Psychiatry, 45*(4), 813-820.
- Muris, P., Fokke, M., & Kwik, D. I. (2009). The ruminative response style in adolescents: An examination of its specific link to symptoms of depression. *Cognitive therapy and research, 33*(1), 21-32.
- Muris, P., Hovee, I., Meesters, C., & Mayer, B. (2004). Children's perception and interpretation of anxiety-related physical symptoms. *Journal of Behavior Therapy and Experimental Psychiatry, 35*(3), 233-244. doi:<http://dx.doi.org/10.1016/j.jbtep.2004.03.008>
- Muris, P., Roelofs, J., Meesters, C., & Boomsma, P. (2004). Rumination and worry in nonclinical adolescents. *Cognitive therapy and research, 28*(4), 539-554.
- Muris, P., Schmidt, H., Merckelbach, H., & Schouten, E. (2001). Anxiety sensitivity in adolescents: factor structure and relationships to trait anxiety and symptoms of anxiety disorders and depression. *Behaviour Research and Therapy, 39*(1), 89-100. doi:S0005-7967(99)00179-5
- Naragon-Gainey, K. (2010). Meta-analysis of the relations of anxiety sensitivity to the depressive and anxiety disorders. *Psychological Bulletin, 136*(1), 128.
- Noel, V. A., & Francis, S. E. (2011). A meta-analytic review of the role of child anxiety sensitivity in child anxiety. *Journal of Abnormal Child Psychology, 39*(5), 721-733.
- Nolen-Hoeksema, S. (1991). Responses to depression and their effects on the duration of depressive episodes. *Journal of Abnormal Psychology, 100*(4), 569. Nolen-Hoeksema, S. (2000). The role of rumination in depressive disorders and mixed anxiety/depressive symptoms. *Journal of Abnormal Psychology, 109*(3), 504-511.

- Olatunji, B. O., & Wolitzky-Taylor, K. B. (2009). Anxiety sensitivity and the anxiety disorders: a meta-analytic review and synthesis. *Psychological Bulletin*, *135*(6), 974-999.
- Preacher, K. J., & Hayes, A. F. (2008). Asymptotic and resampling strategies for assessing and comparing indirect effects in multiple mediator models. *Behavior Research Methods*, *40*(3), 879-891
- Reiss, S., Peterson, R., Gursky, D., & McNally, R. (1986). Anxiety sensitivity, anxiety frequency and the prediction of fearfulness. *Behaviour Research and Therapy*, *24*(1), 1-8.
- Roelofs, J., Rood, L., Meesters, C., te Dorsthorst, V., Bogels, S., Alloy, L. B., et al. (2009). The influence of rumination and distraction on depressed and anxious mood: A prospective examination of the response styles theory in children and adolescents. *European Child and Adolescent Psychiatry*, *18*(10), 635-642.
- Rood, L., Roelofs, J., Bogels, S. M., Nolen-Hoeksema, S., & Schouten, E. (2009). The influence of emotion-focused rumination and distraction on depressive symptoms in non-clinical youth: A meta-analytic review. *Clinical Psychology Review*, *29*(7), 607-616.
- Schmidt, N. B., Keough, M. E., Mitchell, M. A., Reynolds, E. K., Macpherson, L., Zvolensky, M. J., et al. (2010). Anxiety sensitivity: prospective prediction of anxiety among early adolescents. *Journal of Anxiety Disorders*, *24*(5), 503-508.
- Schmidt, N. B., Lerew, D. R., & Joiner, T. E. (1998). Anxiety sensitivity and the pathogenesis of anxiety and depression: Evidence for symptom specificity. *Behaviour Research and Therapy*, *36*(2), 165-177.
- Segerstrom, S., Tsao, J. I., Alden, L., & Craske, M. (2000). Worry and Rumination: Repetitive Thought as a Concomitant and Predictor of Negative Mood. *Cognitive therapy and research*, *24*(6), 671-688.
- Silverman, W., Fleisig, W., Rabian, B., & Peterson, R. (1991). Childhood anxiety sensitivity index. *Journal of Clinical Child & Adolescent Psychology*, *20*(2), 162-168.
- Spence, S. H. (1998). A measure of anxiety symptoms among children. *Behaviour Research and Therapy*, *36*(5), 545-566.

- Taylor, S., Koch, W. J., Woody, S., & McLean, P. (1996). Anxiety sensitivity and depression: how are they related? *Journal of Abnormal Psychology, 105*(3), 474-479.
- Verstraeten, K., Bijttebier, P., Vasey, M. W., & Raes, F. (2011). Specificity of worry and rumination in the development of anxiety and depressive symptoms in children. *British Journal of Clinical Psychology, 50*(4), 364-378. doi:<http://dx.doi.org/10.1348/014466510X532715>
- Weems, C. F., Hammond-Laurence, K., Silverman, W. K., & Ferguson, C. (1997). The relation between anxiety sensitivity and depression in children and adolescents referred for anxiety. *Behaviour Research and Therapy, 35*(10), 961-966.
- Wright, K., Asmundson, G., McCreary, D., Stewart, S., McLaughlin, E., Comeau, M., et al. (2010). Confirmatory Factor Analysis of the Childhood Anxiety Sensitivity Index: A Gender Comparison. *Cognitive Behaviour Therapy, 39*(3), 225-235.

**Table 1.** Descriptive statistics for self-reported anxiety sensitivity, rumination, anxiety and depressive symptoms in unselected children (7 – 11 years)


	$\alpha$	Total	Mean (SD)	
			Females	Males
Anxiety sensitivity <sup>Δ</sup>	.89	29.13 (7.37)	30.74 (7.45)	27.31 (6.90)
<i>Physical</i> <sup>Δ</sup>	.86	19.11 (5.47)	20.35 (5.61)	17.71 (4.99)
<i>Social</i>	.55	5.84 (1.59)	6.03 (1.49)	5.63(1.68)
<i>Mental</i>	.72	4.18 (1.51)	4.36 (1.59)	3.97 (1.40)
Rumination	.90	9.99 (8.41)	10.64 (8.24)	9.23 (8.60)
Anxiety <sup>Δ</sup>	.86	28.27 (15.90)	32.23 (15.73)	23.72 (14.95)
Depression	.90	5.95 (5.70)	6.32 (5.52)	5.53 (5.92)

Δ females significantly higher than males ( $p < .05$ )

**Table 2.** Multiple regression predicting anxiety and depressive symptoms in unselected children (7 – 11 years)

	<u>Total anxiety sensitivity</u>				<u>Anxiety sensitivity dimensions</u>			
	<b>Anxiety</b>		<b>Depression</b>		<b>Anxiety</b>		<b>Depression</b>	
	<i>b</i>	<i>sr</i>	<i>b</i>	<i>sr</i>	<i>b</i>	<i>sr</i>	<i>b</i>	<i>sr</i>
Gender	.12*	--	-.03	--	.13*	--	-.02	--
Anxiety sensitivity	.56**	.36**	.31*	.18*	-	-	-	-
<i>Physical</i>	-	-	-	-	.27**	.19**	.10	.07
<i>Social</i>	-	-	-	-	.16*	.13*	.14	.11
<i>Mental</i>	-	-	-	-	.34**	.25**	.24*	.17*
Rumination	.25*	.15*	.48**	.32**	.14	.10	.40**	.27**
Anxiety	-	-	.01	.01	-	-	.02	.01
Depression	.01	.01	-	-	.01	.00	-	-
adj. R <sup>2</sup>	.62		.53		.65		.56	
F (df)	59.14**		41.21 **		45.39 **		31.30 **	
	(4, 138)		(4, 138)		(6, 136)		(6,136)	

\*  $p < .01$ , \*\*  $p < .001$  *b* unstandardised beta co-efficient *sr* semi-partial correlation coefficient *adj. R<sup>2</sup>* adjusted proportion of variance explained



**Figure 1.** Mediation model of the direct and indirect (via rumination) associations between Mental concerns of anxiety sensitivity (AS) and depression (a) and anxiety (b)

*b* unstandardised beta coefficient *p* significance value