

# Isolation of isoprene degrading bacteria from soils, development of *isoA* gene probes and identification of the active isoprene-degrading soil community using DNA-stable isotope probing

Myriam El Khawand,<sup>1</sup> Andrew T. Crombie,<sup>1\*</sup> Antonia Johnston,<sup>1</sup> Dmitrii V. Vavlline,<sup>2</sup> Joseph C. McAuliffe,<sup>2</sup> Jacob A. Latone,<sup>2</sup> Yuliya A. Primak,<sup>2</sup> Sang-Kyu Lee,<sup>2</sup> Gregg M. Whited,<sup>2</sup> Terry J. McGenity<sup>3</sup> and J. Colin Murrell<sup>1</sup>

<sup>1</sup>University of East Anglia, Norwich Research Park, Norwich, UK.

<sup>2</sup>DuPont Industrial Biosciences, 925 Page Mill Road, Palo Alto, CA 94304, USA.

<sup>3</sup>University of Essex, Wivenhoe Park, Colchester, UK.

## Summary

Emissions of biogenic volatile organic compounds (bVOCs), are an important element in the global carbon cycle, accounting for a significant proportion of fixed carbon. They contribute directly and indirectly to global warming and climate change and have a major effect on atmospheric chemistry. Plants emit isoprene to the atmosphere in similar quantities to emissions of methane from all sources and each accounts for approximately one third of total VOCs. Although methanotrophs, capable of growth on methane, have been intensively studied, we know little of isoprene biodegradation. Here, we report the isolation of two isoprene-degrading strains from the terrestrial environment and describe the design and testing of polymerase chain reaction (PCR) primers targeting *isoA*, the gene encoding the active-site component of the conserved isoprene monooxygenase, which are capable of retrieving *isoA* sequences from isoprene-enriched environmental samples. Stable isotope probing experiments, using biosynthesized <sup>13</sup>C-labelled isoprene, identified the active isoprene-degrading bacteria in soil. This study identifies novel isoprene-degrading strains using both culture-dependent and, for the first time, culture-independent methods and provides the tools and foundations for con-


tinued investigation of the biogeography and molecular ecology of isoprene-degrading bacteria.

## Introduction

Isoprene accounts for approximately one third of the total flux of volatile organic compounds to the atmosphere, an amount that is similar to the methane flux (Guenther *et al.*, 2012; Kirschke *et al.*, 2013). In the atmosphere, isoprene is rapidly photochemically oxidized resulting in a short lifetime (of the order of hours) and consequent low concentrations. Attack by hydroxyl or nitrate radicals or ozone leads to a variety of products depending on temperature and pollutant (nitrogen oxides, NO<sub>x</sub>) levels (Atkinson and Arey, 2003). Overall, isoprene has a significant effect on atmospheric chemistry and hence climate change, due both to the production of greenhouse gases (principally ozone) and by reducing the hydroxyl radical-mediated oxidizing capacity of the atmosphere, which increases the lifetime of methane (Pacifico *et al.*, 2009). In addition, isoprene oxidation-products form secondary organic aerosols and cloud condensation nuclei, with implications for air quality and climate (Fiore *et al.*, 2012).

About 600 Tg y<sup>-1</sup> isoprene is emitted to the atmosphere by terrestrial plants, although not all plant species produce isoprene (Sharkey, 2013; Loreto and Fineschi, 2015). Isoprene synthesis occurs in the chloroplast, via the enzyme isoprene synthase, which converts dimethylallyl pyrophosphate to isoprene (Logan *et al.*, 2000). Isoprene protects plants against heat stress by reducing heat-induced cell-membrane damage, enhances tolerance of reactive oxygen species and may affect plant-insect interactions (Loivamäki *et al.*, 2008; Sharkey *et al.*, 2008; Vickers *et al.*, 2009; Sharkey, 2013). In the marine environment isoprene is released by phytoplankton and macroalgae (Broadgate *et al.*, 2004; Exton *et al.*, 2015). Some bacteria, including soil-dwelling species such as *Bacillus subtilis*, release isoprene, as do some fungi (Kuzma *et al.*, 1995; Julsing *et al.*, 2007; Bäck *et al.*, 2010), although we lack a clear understanding of why these organisms

\*For correspondence. E-mail a.crombie@uea.ac.uk; Tel. +44 (0)1603 592239; Fax +44 (0)1603 259492.



**Fig. 1.** Isoprene metabolism in *Rhodococcus* sp. AD45. Enzymes: IsoABCDEF, isoprene monooxygenase; IsoI, glutathione-S-transferase; IsoH, dehydrogenase. HGMB, 1-hydroxy-2-glutathionyl-2-methyl-3-butene; GMB, 2-glutathionyl-2-methyl-3-butenal; GMBA, 2-glutathionyl-2-methyl-3-butenic acid; SG, glutathione; GSH, reduced glutathione.

produce isoprene. A non-enzymatic reaction resulting in isoprene was reported in humans and other animals, associated with the mevalonate pathway of cholesterol synthesis (Gelmont *et al.*, 1981).

Although atmospheric isoprene concentrations are low [ $< 1\text{--}4$  ppbv in one wide-ranging study (Greenberg *et al.*, 1999)], in the vicinity of isoprene sources (around or below tree canopy level) concentrations are significantly higher. For example, Wiedinmyer *et al.* measured ground level isoprene concentrations of 11 and 36 ppbv at sites in Texas and Missouri respectively (Wiedinmyer *et al.*, 2001; Wiedinmyer *et al.*, 2005). Soils can act as a biological sink for isoprene, at or below these concentrations. In field chambers set up in temperate forest soils, isoprene was rapidly depleted to below the 5 ppbv limit of detection (Cleveland and Yavitt, 1997; Cleveland and Yavitt, 1998). In continuous flow experiments conducted by Gray *et al.* (2015), soils supplied with isoprene at concentrations of 2–200 ppbv consumed isoprene at all concentrations, with a rate of  $62 \text{ pmol g}^{-1} \text{ h}^{-1}$  at 20 ppbv. These data demonstrate both the potential of soils to consume isoprene released locally in soils and also to take up atmospheric isoprene, conclusions also reached in mesocosm experiments by Pegoraro *et al.* (2005). Several bacterial strains, tentatively assigned to the genera *Nocardia*, *Rhodococcus* (Actinobacteria) and *Alcaligenes* (Betaproteobacteria), were isolated from isoprene enrichment cultures and shown to grow on isoprene as sole growth substrate, (van Ginkel *et al.*, 1987a,b; Ewers *et al.*, 1990; Cleveland and Yavitt, 1997) and, more recently, *Pseudomonas*, *Alcaligenes* and *Klebsiella* isoprene-degrading strains were isolated from rubber-contaminated soil (Srivastva *et al.*, 2015). Strains were also obtained from the marine environment, including representatives of Actinobacteria, Bacteroidetes and Alpha- and Gammaproteobacteria (Acuña Alvarez *et al.*, 2009). None of these terrestrial isolates was extensively characterized and the best documented isoprene degrader to date is *Rhodococcus* sp. AD45, a Gram-positive actinobacterium isolated nearly 20 years ago from freshwater sediment by the group of Dick Janssen (van Hylckama Vlieg *et al.*, 1998).

In *Rhodococcus* sp. AD45, isoprene is oxidized to epoxyisoprene (1,2-epoxy-2-methyl-3-butene) by a four-component

soluble diiron centre monooxygenase (SDIMO) with homology to enzymes including the soluble methane monooxygenase (sMMO) and alkene/aromatic monooxygenases (van Hylckama Vlieg *et al.*, 2000; Leahy *et al.*, 2003). The epoxide is then conjugated with glutathione, catalysed by glutathione-S-transferase (GST) (IsoI) and oxidized in two steps by a dehydrogenase (IsoH) resulting in 2-glutathionyl-2-methylbutenoic acid (Fig. 1) (van Hylckama Vlieg *et al.*, 1998; van Hylckama Vlieg *et al.*, 1999). Interestingly, conjugation with glutathione in *Rhodococcus* sp. AD45 contrasts with other alkene utilizers, which often overcome the toxicity of epoxides by forming coenzyme M conjugates or by hydrolysis (Ensign, 2001; Kottegoda *et al.*, 2015). The genes encoding the monooxygenase (*isoABCDEF*) and two subsequent enzymes, together with two additional genes of unknown function, were cloned and sequenced (van Hylckama Vlieg *et al.*, 2000). Recently, we sequenced the genome of *Rhodococcus* sp. AD45 and showed, by mutagenesis, that isoprene monooxygenase (*isoMO*) was essential for isoprene metabolism. Using RNAseq, a cluster of 22 genes was identified, all of which were induced by isoprene or the immediate product of isoprene oxidation, epoxyisoprene (Crombie *et al.*, 2015).

DNA stable isotope probing (DNA-SIP) is a cultivation-independent technique with the ability to identify active substrate-consuming organisms in environmental samples (Radajewski *et al.*, 2000; Dumont and Murrell, 2005). The method relies on incubation of samples with stable-isotope-labelled growth substrate. The incorporation of isotope (typically  $^{13}\text{C}$  or  $^{15}\text{N}$ ) into biomass (including DNA), enables the identification of active microorganisms following separation of labelled and unlabelled DNA by isopycnic (density gradient) centrifugation.

Despite its abundance and climatic importance, our knowledge of isoprene in the environment is heavily skewed towards production in plants and atmospheric oxidation, with only a few studies investigating isoprene biodegradation. Isoprene is an abundant plant secondary metabolite, also produced in soils, and would provide a good source of carbon and energy for bacteria. Strains capable of growth on isoprene have frequently been isolated from diverse environments, albeit generally not characterized at the molecular level. Our hypothesis was

that isoprene degraders are widely distributed and may play an important role in the biogeochemistry of this environmentally important trace gas. Our aim was to isolate and sequence isoprene degraders, and identify putative isoprene metabolic genes. We aimed to develop gene probes to target key diagnostic markers of isoprene degradation and to identify the active isoprene-assimilating organisms in soil enrichments, using DNA-SIP.

## Results and discussion

### *Enrichment and isolation of two novel terrestrial isoprene-utilizing bacteria*

Our initial aim was to isolate isoprene degraders from contrasting environments in order to provide sequence data for later cultivation-independent approaches. As isoprene consumption has previously been observed in soils and since isoprene is emitted from tree leaves, we used these as source material. Separate enrichments were set up, using either garden soil or Horse Chestnut (*Aesculus hippocastanum*) leaves, in minimal medium and incubated with isoprene. Both enrichments consumed isoprene and two isolates (designated SC4 and LB1), capable of growing on isoprene as sole source of carbon and energy, were obtained from soil and leaf samples respectively. The nearly complete (1521 nucleotides) 16S rRNA gene sequences of both strains SC4 and LB1 were identical to strains of both *Rhodococcus opacus* and *R. wratislaviensis* (Supporting Information Fig. S1). Strains SC4 and LB1 also grew on acetate, succinate, glucose, fructose, propane and butane as sole source of carbon and energy (Supporting Information Table S1), in contrast to *Rhodococcus* sp. AD45, which does not grow on propane or butane (Crombie *et al.*, 2015).

### *Genome sequencing*

To identify isoprene-related gene sequences, we sequenced the genomes of strains SC4 and LB1. Their genomes, 10.6 and 10.7 Mbp, are considerably larger than that of *Rhodococcus* sp. AD45 (6.9 Mbp) and closer to that of *R. jostii* RHA1 (9.7 Mbp) (McLeod *et al.*, 2006; Crombie *et al.*, 2015), whereas the GC contents (66.7% and 66.6% respectively) are typical of the genus (Supporting Information Table S2).


### *Identification of isoprene related genes in isolates*

Our previous work (Crombie *et al.*, 2015) detected a cluster of 22 genes induced by isoprene, which appeared to be the complete inventory of isoprene-responsive genes. In addition to *isoABCDEF*, encoding the monooxygenase, and *isoGHIJ* that encode a protein of unknown function, a dehydrogenase and two glutathione-S-transferases previously described (van Hylckama Vlieg *et al.*, 2000), the cluster includes glutathione

biosynthesis genes, predicted aldehyde dehydrogenases and a coenzyme-A disulfide reductase (Crombie *et al.*, 2015). We therefore searched for homologous sequences in the genomes of strains SC4 and LB1, which are highly similar to each other in this region (over 99% nucleotide identity). Using the isoprene-responsive gene products previously identified in *Rhodococcus* sp. AD45 as query sequences in tBLASTn searches, we identified homologues of all of these 22 genes, with amino acid sequence identity ranging from 50% to 96% (Supporting Information Table S3). The most highly conserved were *isoABCDEF* encoding the multicomponent IsoMO (81–96%), whereas a predicted protein of unknown function (SZ00\_06083), highly induced by isoprene in *Rhodococcus* sp. AD45, shared 50% amino acid identity with sequences from these strains. We observed the same duplication of *isoGHIJ* (Fig. 2, 77–88% amino acid identity between copies) as is present in *Rhodococcus* sp. AD45. Between *isoA* and *isoJ*, both of the new strains contain a second copy (81% amino acid identity between copies), not present in *Rhodococcus* sp. AD45, of an aldehyde dehydrogenase (*aldh1*) which is located approximately 10 000 nucleotides (nt) upstream of the monooxygenase in *Rhodococcus* sp. AD45. A gene encoding a predicted coenzyme-A disulfide reductase is also present in two copies (69% amino acid identity between copies), although in strain LB1 (but not in strain SC4) one copy has a nucleotide insertion approximately 255 nt from the end, resulting in a frameshift mutation, suggesting this may not encode a functional protein. In comparison with *Rhodococcus* sp. AD45, five additional genes are present, in both strains, in the middle of the cluster. These are predicted to encode two hypothetical proteins and an alpha/beta hydrolase domain-containing protein of unknown function, an acetyl-CoA acetyltransferase and a 3-hydroxyacyl-CoA dehydrogenase. Interestingly, in strain LB1, another insertion in the acetyl-CoA acetyltransferase has resulted in a frameshift mutation. None of these five genes are present in this region of the *Rhodococcus* sp. AD45 genome, nor were remotely located homologous sequences induced by isoprene (Crombie *et al.*, 2015), implying that they are not essential for isoprene metabolism. In comparison with *Rhodococcus* sp. AD45, strains LB1 and SC4 are more similar in this region of the genome, in terms of both sequence identity and gene layout, with *R. opacus* PD630 (also included in Fig. 2). Previously, using sequence data, we predicted, and confirmed, that *R. opacus* PD630 could grow on isoprene (Crombie *et al.*, 2015), although, beyond this, we have no direct experimental data regarding isoprene-related gene function in this strain.

### *Development and validation of functional gene markers targeting isoA*

Using the sequence data from these isoprene-degrading strains, we designed probes to detect isoprene-related



**Fig. 2.** The isoprene metabolic gene cluster from *Rhodococcus* sp. AD45 (Crombie *et al.*, 2015), together with similar regions from *R. opacus* PD630 and strains SC4 and LB1. The contigs containing the genes are identified by horizontal lines and numbers below. The monooxygenase genes are shown in red, and other colours indicate genes of the corresponding predicted function between strains. Locus tags and gene names are indicated with angled text. Locus tag prefixes: *R. sp. AD45*, SZ00\_; *R. opacus* PD630, Pd630\_LPD; *R. sp. SC4*, AXA44\_; *R. sp. LB1*, AZG88\_.


genes in environmental samples. *isoA* codes for the alpha-subunit of the hydroxylase of IsoMO and contains the diiron centre active site and was shown to be essential for isoprene degradation in *Rhodococcus* sp. AD45 (Crombie *et al.*, 2015). We, therefore, selected *isoA* as target for a functional gene probe. To expand the diversity of *isoA* sequences, we also retrieved *isoA* genes from draft sequences of a *Gordonia* strain and a *Mycobacterium* strain previously isolated from surface sediment from the Colne estuary (Essex, UK), identified as *Gordonia* i37 and *Mycobacterium* AT1 (Acuña Alvarez *et al.*, 2009; Johnston, 2014). The *isoA* sequences of strains SC4, LB1, i37 and AT1, together with those of *Rhodococcus* sp. AD45 (SZ00\_06091) and *R. opacus* PD630 (Pd630\_LPD03572), were aligned at the amino acid level, and conserved regions were used to design primers targeting *isoA*. To exclude other non-isoprene-degrading members of the SDIMO family, sequences of *mmoX* and *xamoA*, encoding the alpha subunit of sMMO from *Methylosinus trichosporium* OB3b (Cardy *et al.*, 1991), and alkene monooxygenase from *Xanthobacter autotrophicus* Py2 (Zhou *et al.*, 1996), respectively, were also included in the alignment.

The *isoA* primers were validated by PCR amplification of template DNA from contrasting sources (Supporting Information Table S4): (i) genomic DNA from 15 isoprene-degrading isolates, (ii) DNA extracted from three isoprene-enriched soils and four isoprene enrichments of marine and estuarine water and sediment and (iii) control DNA extracted from eight non-isoprene-degrading isolates able to grow on alkanes, alkenes or aromatic compounds. We

obtained PCR products of the expected size using DNA extracted from all the isoprene-degrading isolates and enrichments, but not from any of the non-isoprene degraders. PCR products from enrichments were cloned and analysed by restriction fragment length polymorphism (RFLP) (Supporting Information Table S5). Representatives of each operational taxonomic unit (OTU) were sequenced, (all of which appeared to be *isoA* sequences) and aligned at the amino acid level with *isoA* sequences obtained from the sequenced genomes. A phylogenetic tree of the *isoA* nucleotide sequences (1011 nt) was constructed from the alignment (Fig. 3). All the sequences, although from diverse phylogenetic groups including both Gram-positive and Gram-negative strains, were relatively similar (> 86% amino acid identity between sequences), but could be broadly separated into two groups in which the terrestrial sequences and those from the low-salinity environment of Hythe, on the Colne estuary, were distinct from marine and other estuarine sequences similar to *isoA* of *Gordonia* i37.


#### Active isoprene-assimilating bacteria identified by DNA-stable isotope probing

As all four of our sequenced terrestrial isolates were rhodococci and all *isoA* sequences retrieved were relatively similar, we used cultivation-independent methods to test whether a greater diversity of isoprene degraders existed in soils in a DNA-stable isotope experiment (DNA-SIP) using  $^{13}\text{C}$ -labelled isoprene, biosynthesized as described in




**Fig. 3.** Phylogenetic tree of *isoA* genes from isolates and environmental clones, constructed using the Maximum Likelihood method in MEGA6 (Tamura *et al.*, 2013). All positions containing gaps and missing data were eliminated and there were 1011 nucleotide positions in the final dataset. The scale bar shows nucleotide substitutions per site. Bootstrap values (500 replications) greater than 50% are shown at the nodes. Isoprene-degrading isolates are shown in bold. Cloned *isoA* sequences (Supporting Information Tables S4 and S5) are labelled with the sample site followed by clone identification number. WCO\_L4, Western Channel Observatory station L4; FW, freshwater.

**Experimental Procedures.** Soil microcosms (in triplicate for labelled substrate and duplicate for unlabelled-substrate controls) were incubated with 0.5% (v/v) isoprene, without any other amendments. Isoprene was consumed without an appreciable lag phase, and incubations were terminated after consumption of 10  $\mu\text{mol}$  substrate  $\text{g}^{-1}$  soil (15 days). DNA was extracted from  $^{13}\text{C}$ - and control  $^{12}\text{C}$ -isoprene incu-



**Fig. 4.** Soil bacterial community (based on 16S rRNA gene amplicons) of labelled isoprene DNA-SIP microcosms. The pie charts show (left) the unenriched soil community from timepoint zero and (right) the  $^{13}\text{C}$ -isoprene-enriched total community (prior to isopycnic centrifugation and DNA fractionation). For the right hand chart, DNA from triplicate enrichments was pooled prior to analysis.

bations and separated into heavy and light fractions and used for 454 pyrosequencing of 16S rRNA genes. The unenriched soil at timepoint zero displayed a typically diverse community, comprising 50% Proteobacteria, with Planctomycetes, Actinobacteria, Bacteroidetes, Chloroflexi and Firmicutes contributing an additional 40% (Fig. 4). The major effect of the incubations was to greatly increase the relative abundance of Actinobacteria, while having a relatively minor effect on the remaining phyla (Fig. 4). Multivariate analysis (Supporting Information Fig. S2) shows that the unlabelled bacterial community of the  $^{13}\text{C}$ -isoprene incubations (i.e. light DNA fraction) was extremely similar to that of the timepoint zero community, suggesting that the change in community profile during the incubations was due to an enrichment of isoprene degraders. As expected, the community represented by the light fraction of the  $^{12}\text{C}$ -isoprene incubations, derived from both isoprene-consumers and non-consumers, was extremely similar to the total (unfractionated) DNA from  $^{13}\text{C}$ -incubations. The heavy fraction of  $^{12}\text{C}$ -isoprene incubations, which contained only a very small fraction of the total DNA of  $^{12}\text{C}$ -incubations, was less similar. This DNA results mainly from incomplete separation, with minor effects due to factors such as GC content of DNA. In contrast to all of these, the active isoprene degraders represented by the heavy fractions of the  $^{13}\text{C}$  incubations formed a distinct community (Supporting Information Fig. S2) and displayed a dramatically altered taxonomic profile (Fig. 5), becoming dominated by *Rhodococcus* sequences, which comprised  $88\% \pm 5\%$  (mean  $\pm$  s.d.) of the heavy fraction of  $^{13}\text{C}$  incubations. These sequences were predominantly those of *Rhodococcus wratislaviensis*, *R. koreensis* and *R. globerulus* (Fig. 5) and were 147- to 161-fold more abundant in the heavy



**Fig. 5.** Bacterial communities derived from labelled isoprene DNA-SIP enrichments and fractionation. The bar chart shows 16S rRNA gene-based composition of the unenriched soil community (timepoint zero), the isoprene-enriched total community (unfractionated) and labelled (heavy) and unlabelled (light) components separated by isopycnic centrifugation and fractionation. The symbols on the x-axis correspond with those shown in Supporting Information Fig. S2. The isoprene-assimilating community is represented by the heavy fractions of the three replicate  $^{13}\text{C}$  incubations (solid red diamonds). T-0, timepoint zero; U-F, unfractionated; H, heavy fraction; L, light fraction. The inset (b) shows the species composition of the rhodococci in the heavy fractions (mean of three replicates)

fractions compared to light fractions of  $^{13}\text{C}$  incubations, but without any corresponding enrichment in the heavy fraction of  $^{12}\text{C}$ -incubations (Supporting Information Fig. S3). Also enriched in the heavy fractions of  $^{13}\text{C}$ -incubations were members of the Betaproteobacteria, *Comamonas* spp. and *Variovorax* spp., which together comprised  $6.5\% \pm 1.2\%$  of the  $^{13}\text{C}$  heavy fractions (but were not detected in the  $^{13}\text{C}$  light fractions), and were also enriched during the incubations, increasing from 0.3% to 0.7% of the total community.

These data clearly demonstrate that *Rhodococcus* and, to a lesser extent, *Comamonas* and *Variovorax* spp. had assimilated carbon, directly or indirectly, from labelled isoprene. Searches of the publicly available databases did not identify high-similarity putative isoprene-related genes in the family Comamonadaceae (which encompasses both *Comamonas* and *Variovorax*) (see Experimental Procedures). Comparison with the PCR-based *isoA* analysis, which generated similar sequences from diverse phylogenetic groups,

suggests that the isoprene-degrading members of the Comamonadaceae implicated in the SIP experiment are not represented in the NCBI databases, or that they use genes or possibly pathways dissimilar to those of the characterized strains.

Previous studies of terrestrial environments have isolated *Alcaligenes*, *Klebsiella* and *Pseudomonas* isoprene-degrading strains in addition to the Actinobacteria *Nocardia* and *Rhodococcus*, although some of these identifications were not based on molecular data and most strains were not extensively characterized (van Ginkel *et al.*, 1987a,b; Ewers *et al.*, 1990; Cleveland and Yavitt, 1997; van Hylckama Vlieg *et al.*, 1998; Srivastva *et al.*, 2015). In our SIP incubations, 16S rRNA gene sequences of *Alcaligenaceae* spp. were slightly enriched during the incubations, increasing from 0.3% to 0.6% of the total community, but were concentrated (14:1) in the non-labelled light DNA fractions. *Pseudomonas* spp., present at 0.9% of the initial community, decreased to 0.1% following isoprene incubations and were also not labelled, whereas *Klebsiella* were not detected at any point. These data indicate that these taxa had not assimilated carbon from isoprene under our experimental conditions.

### Conclusions

The isoprene concentrations used here are considerably in excess of those normally encountered in the environment and the aim was not to replicate environmental conditions, but rather to expand the diversity of known isoprene degraders. Draft genome sequences showed that isoprene monooxygenase was present in all our isoprene-degrading isolates, which enabled the development of *isoA* primers to effectively target this enzyme with high specificity. While we cannot exclude the possibility that some sequences may be missed, or that other isoprene-degrading enzymes or pathways exist, these primers constitute an effective tool to identify isoprene-related gene sequences in environmental samples. DNA-SIP, to our knowledge the first time this technique has been used to identify isoprene assimilators, showed that the major isoprene utilisers in the microcosms were *Rhodococcus* strains and that members of the Comamonadaceae were also active in isoprene degradation. The fact that we did not obtain isolates from this family suggests that they may be resistant to cultivation under our laboratory conditions, emphasizing the importance of cultivation-independent techniques. This study confirms that soils readily consume isoprene and harbor a diverse community of isoprene degraders. Investigation of their diversity, abundance and mechanisms of isoprene degradation is essential to assess the environmental relevance of the global biological isoprene sink and the extent to which biodegradation moderates the effect on the atmosphere of this abundant and climate-active trace

gas. This study provides the tools and foundations to further investigate these topics. Future experiments should search for novel genes and pathways involved in isoprene degradation, perhaps using SIP coupled with metagenomics. The isoprene-degrading community in the phyllosphere is also worthy of investigation, as our isolation of a strain from this environment, to our knowledge the first published example, suggests that isoprene degraders may be present or abundant on leaves, close to the major source of isoprene to the atmosphere.

### Experimental procedures

#### Cultivation of bacterial strains

The terrestrial isoprene-degrading strains were routinely grown in minimal medium (CBS medium) which contained (per 1 l): 0.1 g MgSO<sub>4</sub>·7H<sub>2</sub>O, 0.8 g NH<sub>4</sub>Cl, 1.5 g KH<sub>2</sub>PO<sub>4</sub>, 6.3 g Na<sub>2</sub>HPO<sub>4</sub> and 10 ml of trace element solution (Tuovinen and Kelly, 1973) (pH 7.0). Marine isoprene degrading-bacteria were grown in mineral salts medium prepared according to Schaefer *et al.* (2002), except that it contained 20 g l<sup>-1</sup> NaCl and was supplemented with Na<sub>3</sub>VO<sub>4</sub> and Na<sub>2</sub>SeO<sub>3</sub> (5 ng l<sup>-1</sup> each). Cultures were set up in serum vials (120 ml) sealed with grey butyl rubber seals or in Quickfit flasks (250 ml or 2 l) fitted with SubaSeal stoppers (Sigma-Aldrich) and isoprene was added (as gas) to a final concentration of 0.6–1% (v/v) by injection through the septum. The cultures were incubated at 30°C, shaking at 150 r.p.m.

For growth tests on other gaseous substrates, 25 ml of CBS medium in serum vials (120 ml) was inoculated with isoprene-grown culture (5% inoculum) and incubated with 10% (v/v) substrate. Cultures with succinate, glucose, fructose or acetate (10 mM) as growth substrates were prepared in universal bottles (20 ml) containing 5 ml CBS medium inoculated with 5% of isoprene-grown culture. Isoprene (catalogue no. I19551) was obtained from Sigma Aldrich.

#### Isolation of isoprene-degrading strains

Isoprene enrichment cultures were set up using garden soil from Leamington Spa (UK) or leaves of a Horse Chestnut tree from the campus of the University of Warwick (Coventry, UK). Isoprene (0.6% v/v) was added to 50 ml CBS minimal medium in flasks (250 ml) and inoculated with either 0.3 g soil or one leaf, cut into small pieces. The optical density of enrichment cultures was followed spectrophotometrically at 540 nm and isoprene uptake was monitored with a gas chromatograph fitted with a flame ionization detector (GC-FID) as described previously (Crombie *et al.*, 2015). Enrichment cultures were streaked on CBS agar plates and incubated at 30°C in a desiccator with isoprene vapour (approximately 5% v/v). Colonies were subcultured until pure, confirmed by phase contrast microscopy (Zeiss Axioscop). Marine strains for primer design and validation were isolated as described previously (Acuña Alvarez *et al.*, 2009; Johnston, 2014).

### DNA extraction, amplification and sequencing of 16S rRNA genes

DNA was extracted from isolates and enrichment cultures using the FastDNA Spin Kit for Soil (MP Biomedicals), following the manufacturer's instructions. For identification of strains, 16S rRNA genes were amplified using 27f/1492R primers (Lane, 1991). Amplicons were purified, cloned into pGEMT-easy vector (Promega) and sequenced with M13 primers (Invitrogen).

### Genome sequencing, annotation and mining

High molecular-mass genomic DNA was extracted from 500 ml mid-late exponential cultures of isoprene-degrading strains following the Marmur extraction method (Marmur, 1961) except that the sodium dodecyl sulfate (SDS) concentration was increased to 2% (w/v) and the incubation period at 55°C in sucrose/ethylenediaminetetraacetic acid/Tris buffer and SDS extended to 5 h to achieve better cell lysis.

The genome of *Gordonia* i37 was sequenced at Oregon State University (USA) using a Roche 454 pyrosequencing platform. The genomes of *Mycobacterium* AT1, *Rhodococcus* SC4 and *Rhodococcus* LB1 were sequenced using Illumina GAIIx at the University of Warwick Genomics Facility (Coventry, UK). Reads were assembled into contigs using CLC Genomics Workbench for de novo assembly (CLC bio, Aarhus, Denmark). The genome sequences were uploaded to RAST (Rapid Annotation using Subsystem Technology) for annotation. Local nucleotide databases were constructed using NCBI BLAST in BioEdit. The database was mined using tBLASTn with the amino acid sequences of genes shown to be important in isoprene metabolism in *Rhodococcus* sp. AD45 as query sequences. These Whole Genome Shotgun projects have been deposited at DDBJ/EMBL/GenBank under the accession numbers LTCZ00000000 (SC4) and LSBM00000000 (LB1). Versions described in this paper are LTCZ01000000 and LSBM01000000. Cloned *isoA* sequences and *isoA* gene sequences of *Mycobacterium* AT1 and *Gordonia* i37 have been deposited under accession numbers KU870702 – KU870744 and KU870745 and KU870746 respectively.

### *isoA* Primer design and PCR

Deduced amino acid sequences of *isoA* and other related enzymes were aligned using ClustalW in Mega6 (Tamura et al., 2013). The alignment was visually inspected for conserved regions specific to isoprene degraders, which were not conserved among homologous sequences from related SDIMO enzymes. Primers for *isoA* amplification, containing a maximum of three degenerate bases, were designed based on these regions (5'-TGCATGGTCGARCAYATG-3' and 5'-GRTCYTYGTCGAAGCACCACCTT-3'), yielding a predicted amplicon of 1015 bp (*Rhodococcus* sp. AD45 template). These primers were used with a touchdown PCR protocol: an initial step at 94°C for 3 min was followed by 19 cycles of denaturation at 94°C for 30 s, annealing at variable temperatures for 45 s, and extension at 72°C for 60 s. The annealing temperature, initially 72°C, was decreased by 1°C per cycle

until 54°C and maintained at this temperature for a further 25 cycles, followed by a final extension at 72°C for 5 min.

### Clone libraries

Clone libraries were constructed from *isoA* amplicons using DNA extracted from isoprene enrichments of soil, sediment or water. The purified *isoA* amplicons were cloned into pGEMT Easy vector (Promega) prior to transformation into *Escherichia coli* TOP10 cells (Invitrogen) following the manufacturers' instructions. Clones were screened by PCR using M13 primers and RFLP using *EcoRI* and *MspI* restriction enzymes (Fermentas). Clones yielding identical restriction fragment profiles on agarose gels were designated OTUs and representatives of each OTU were sequenced using M13 primers.

### Search for *isoA* genes in the Comamonadaceae

Tblastn (Altschul et al., 1990) was used to search the NCBI nr and genome databases and 153 whole genome shotgun projects deposited in NCBI, for isoprene-degrading gene sequences among the Comamonadaceae. Using *IsoA* from *Rhodococcus* sp. AD45 as query, genes with 46–48% amino acid identity were identified in the genomes of *Hydrogenophaga* sp. T4, *Comamonas badia* DSM 17552 and *Variovorax paradoxus* ZNC0006. Genes encoding the other subunits of the monooxygenase were identified in the *Hydrogenophaga* and *Comamonas* genomes, but no identifiable additional isoprene metabolic genes were found nearby, and no hits to *IsoI* were found in any of the genomes in any location (evalue 0.0001). The *isoA* homologues were more similar to characterized toluene monooxygenase genes, such as *touA* from *Pseudomonas stutzeri* OX1 (67–73% amino acid identity) (Bertoni et al., 1998), suggesting that isoprene may not be their natural substrate.

### Biosynthesis of [1,2,3,4,5-<sup>13</sup>C]-2-methyl-1,3-butadiene ([U-<sup>13</sup>C]-isoprene) from D-[U-<sup>13</sup>C]-glucose

Uniformly labelled <sup>13</sup>C-isoprene was prepared utilizing an isoprene-producing engineered variant of *E. coli* BL21, expressing a functional mevalonate pathway and an isoprene synthase, as biocatalyst. Specifically, the genes encoding thiolase, mevalonate synthase and 3-hydroxy-3-methyl-glutaryl-(HMG-) CoA reductase (*mvaE*, *mvaS*) were derived from *Enterococcus faecalis*, the gene encoding mevalonate kinase was derived from *Methanosarcina mazei*, and the genes encoding phosphomevalonate kinase, mevalonate pyrophosphate decarboxylase and isopentyl-PP isomerase were from *Saccharomyces cerevisiae*. This recombinant pathway serves to channel carbon from glucose to the isoprene precursor dimethylallyl pyrophosphate (DMAPP). The final intracellular enzymatic step to convert DMAPP to isoprene was catalyzed by recombinant isoprene synthase from *Populus alba*. This strain was designated *E. coli* CMP1082 and has been described in detail elsewhere (McAuliffe et al., 2015). Details of fermentation and isoprene recovery and its <sup>1</sup>H NMR spectrum are included in Supporting Information.



### DNA stable isotope probing experiments

For SIP incubations, 5 g soil (pH 7.4), collected from the upper 5 cm (after removal of vegetation and leaf litter), in the vicinity of Willow (*Salix fragilis*) trees on the University of East Anglia campus, was incubated in serum vials (120 ml volume) with 0.5% (v/v) isoprene in triplicate (labelled substrate) or duplicate (unlabelled substrate). Vials were incubated at room temperature (22°C) in the dark. Headspace isoprene concentration was monitored by GC-FID. When substrate was depleted, additional isoprene was injected through the septum to the original concentration. Incubations were terminated and DNA extracted from the soil when 10  $\mu\text{mol g}^{-1}$  had been consumed. Control incubations with autoclaved soil were also conducted. DNA extracted from samples was separated into heavy ( $^{13}\text{C}$ -labelled) and light (unlabelled) fractions as previously described (Neufeld *et al.*, 2007). Briefly, 1–2  $\mu\text{g}$  DNA was added to caesium chloride solution (final density of 1.725  $\text{g ml}^{-1}$ ) and subjected to density gradient ultracentrifugation (177 000  $\times g$ , 40 h, 20°C, Beckman Vti 65.2 rotor). The contents of each tube were separated into 12–15 fractions and the density of each fraction measured by refractometry (AR200 digital refractometer, Reichert, Buffalo, USA). DNA from each fraction was precipitated, resuspended in nuclease-free water and characterized by 16S rRNA gene analysis using denaturing gradient gel electrophoresis (DGGE). DGGE showed unique profiles in heavy, compared to light, fractions of  $^{13}\text{C}$ -incubations while the  $^{12}\text{C}$  incubations exhibited similar profiles between fractions, indicating that heavy fractions from  $^{13}\text{C}$  incubations contained the DNA of isoprene degraders (data not shown). Based on fraction density and DGGE profiles, fractions 7 and 11 were identified as containing labelled and unlabelled DNA, respectively, designated 'heavy' and 'light' and used for downstream analysis. DNA was obtained from six different treatments: extracted from timepoint zero (one sample); extracted following  $^{13}\text{C}$ -isoprene incubation but prior to fractionation (unfractionated); and after separation into heavy and light fractions for both  $^{13}\text{C}$ - and  $^{12}\text{C}$ -isoprene incubations. For  $^{13}\text{C}$  heavy fractions triplicate samples were analysed individually, but for the other treatments the DNA was pooled prior to analysis.

### Denaturing gradient gel electrophoresis

Bacterial 16S rRNA genes were amplified using primers 341F-GC (Muyzer *et al.*, 1993) and 907R (Muyzer *et al.*, 1998). The PCR products were run on an 8% (w/v) polyacrylamide gel with a 30–70% linear denaturant gradient. Electrophoresis was carried out for 16 h, at 80 V, 60°C using the DCode™ Universal Mutation Detection System (Bio-Rad), stained with SYBR® Gold Nucleic Acid Gel Stain (Invitrogen) and imaged using a BioRad GelDoc system.

### Bacterial 16S rRNA gene analysis by 454 pyrosequencing

Labelled (heavy) and unlabelled (light) DNA from SIP incubations was characterized by sequencing of 16S rRNA gene amplicons generated by PCR using the primers 27Fmod (5'-AGRGTGGATCMTGGCTCAG-3') and 519Rmodbio (5'-GTNTTACNGCGGCKGCTG-3') using a Roche 454 FLX tita-

nium instrument at MRDNA (Molecular Research LP), Shallowater, USA. Sequence data were processed at MRDNA using a published pipeline (Dowd *et al.*, 2008; Capone *et al.*, 2011). Briefly, the Q25 reads were stripped of barcodes and primers. Short sequences (< 200 bp), sequences with ambiguous base calls and those with > 6 bp homopolymer runs were removed. Remaining sequences were denoised using a custom pipeline, OTUs clustered at 97% sequence identity, chimeric sequences were removed using Uchime (Edgar *et al.*, 2011) and taxonomy was assigned using BLASTn against the RDP/NCBI database (v 11.1) (Cole *et al.*, 2014). An average of 3320 sequences per sample were used for analysis.

### Statistical analysis

The Bray-Curtis distance measure was applied to log-transformed relative abundance and used to generate non-metric multidimensional scaling (NMDS) plots using Primer 6 (Primer-E, Plymouth, UK).

### Acknowledgements

We thank Dan Arp, Luis Sayavedra-Soto and colleagues at Oregon State University for *Gordonia* sequence information. This research was funded by a Natural Environment Research Council (NERC) grant to JCM (NE/J009725/1) and TJM (NE/J009555/1) entitled *Microbial degradation of isoprene in the terrestrial environment* and by the Earth and Life Systems Alliance (ELSA) at the University of East Anglia. We acknowledge the support of a University of Warwick Postgraduate Research Scholarship awarded to MK and a NERC studentship awarded to AJ. We thank Hendrik Schäfer for help with genome assembly.

### References

- Acuña Alvarez, L., Exton, D.A., Timmis, K.N., Suggett, D.J., and McGenity, T.J. (2009) Characterization of marine isoprene-degrading communities. *Environ Microbiol* **11**: 3280–3291.
- Altschul, S.F., Gish, W., Miller, W., Myers, E.W., and Lipman, D.J. (1990) Basic local alignment search tool. *J Mol Biol* **215**: 403–410.
- Atkinson, R., and Arey, J. (2003) Atmospheric degradation of volatile organic compounds. *Chem Rev* **103**: 4605–4638.
- Bäck, J., Aaltonen, H., Hellén, H., Kajos, M.K., Patokoski, J., Taipale, R. *et al.* (2010) Variable emissions of microbial volatile organic compounds (MVOCs) from root-associated fungi isolated from Scots pine. *Atmos Environ* **44**: 3651–3659.
- Bertoni, G., Martino, M., Galli, E., and Barbieri, P. (1998) Analysis of the gene cluster encoding toluene/o-xylene monooxygenase from *Pseudomonas stutzeri* OX1. *Appl Environ Microbiol* **64**: 3626–3632.
- Broadgate, W.J., Malin, G., Küpper, F.C., Thompson, A., and Liss, P.S. (2004) Isoprene and other non-methane hydrocarbons from seaweeds: a source of reactive hydrocarbons to the atmosphere. *Mar Chem* **88**: 61–73.

- Capone, K.A., Dowd, S.E., Stamatas, G.N., and Nikolovski, J. (2011) Diversity of the human skin microbiome early in life. *J Invest Dermatol* **131**: 2026–2032.
- Cardy, D.L., Laidler, V., Salmond, G.P., and Murrell, J.C. (1991) Molecular analysis of the methane monooxygenase (MMO) gene cluster of *Methylosinus trichosporium* OB3b. *Mol Microbiol* **5**: 335–342.
- Cleveland, C.C., and Yavitt, J.B. (1997) Consumption of atmospheric isoprene in soil. *Geophys Res Lett* **24**: 2379–2382.
- Cleveland, C.C., and Yavitt, J.B. (1998) Microbial consumption of atmospheric isoprene in a temperate forest soil. *Appl Environ Microbiol* **64**: 172–177.
- Cole, J.R., Wang, Q., Fish, J.A., Chai, B., McGarrell, D.M., Sun, Y. *et al.* (2014) Ribosomal Database Project: data and tools for high throughput rRNA analysis. *Nucleic Acids Res* **42**: D633–642.
- Crombie, A.T., Khawand, M.E., Rhodius, V.A., Fengler, K.A., Miller, M.C., Whited, G.M. *et al.* (2015) Regulation of plasmid-encoded isoprene metabolism in *Rhodococcus*, a representative of an important link in the global isoprene cycle. *Environ Microbiol* **17**: 3314–3329.
- Dowd, S.E., Callaway, T.R., Wolcott, R.D., Sun, Y., McKeethan, T., Hagevoort, R.G., and Edrington, T.S. (2008) Evaluation of the bacterial diversity in the feces of cattle using 16S rDNA bacterial tag-encoded FLX amplicon pyrosequencing (bTEFAP). *BMC Microbiol* **8**: 125.
- Dumont, M.G., and Murrell, J.C. (2005) Stable isotope probing – linking microbial identity to function. *Nat Rev Microbiol* **3**: 499–504.
- Edgar, R.C., Haas, B.J., Clemente, J.C., Quince, C., and Knight, R. (2011) UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics* **27**: 2194–2200.
- Ensign, S.A. (2001) Microbial metabolism of aliphatic alkenes. *Biochemistry* **40**: 5845–5853.
- Ewers, J., Freier-Schroder, D., and Knackmuss, H.J. (1990) Selection of trichloroethene (TCE) degrading bacteria that resist inactivation by TCE. *Arch Microbiol* **154**: 410–413.
- Exton, D.A., McGenity, T.J., Steinke, M., Smith, D.J., and Suggett, D.J. (2015) Uncovering the volatile nature of tropical coastal marine ecosystems in a changing world. *Glob Change Biol* **21**: 1383–1394.
- Fiore, A.M., Naik, V., Spracklen, D.V., Steiner, A., Unger, N., Prather, M. *et al.* (2012) Global air quality and climate. *Chem Soc Rev* **41**: 6663–6683.
- Gelmont, D., Stein, R.A., and Mead, J.F. (1981) Isoprene – the main hydrocarbon in human breath. *Biochem Biophys Res Commun* **99**: 1456–1460.
- van Ginkel, C.G., Welten, H.G.J., and de Bont, J.A.M. (1987a) Oxidation of gaseous and volatile hydrocarbons by selected alkene-utilizing bacteria. *Appl Environ Microbiol* **53**: 2903–2907.
- van Ginkel, C.G., de Jong, E., Tilanus, J.W.R., and de Bont, J.A.M. (1987b) Microbial oxidation of isoprene, a biogenic foliage volatile and of 1,3-butadiene, an anthropogenic gas. *FEMS Microbiol Lett* **45**: 275–279.
- Gray, C.M., Helmig, D., and Fierer, N. (2015) Bacteria and fungi associated with isoprene consumption in soil. *Elem Sci Anth* **3**: 000053.
- Greenberg, J.P., Guenther, A., Zimmerman, P., Baugh, W., Geron, C., Davis, K. *et al.* (1999) Tethered balloon measurements of biogenic VOCs in the atmospheric boundary layer. *Atmos Environ* **33**: 855–867.
- Guenther, A.B., Jiang, X., Heald, C.L., Sakulyanontvittaya, T., Duhl, T., Emmons, L.K., and Wang, X. (2012) The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions. *Geosci Model Dev* **5**: 1471–1492.
- van Hylckama Vlieg, J.E.T., Kingma, J., van den Wijngaard, A.J., and Janssen, D.B. (1998) A glutathione S-transferase with activity towards cis-1,2-dichloroepoxyethane is involved in isoprene utilization by *Rhodococcus* sp. strain AD45. *Appl Environ Microbiol* **64**: 2800–2805.
- van Hylckama Vlieg, J.E.T., Kingma, J., Kruijzinga, W., and Janssen, D.B. (1999) Purification of a glutathione S-transferase and a glutathione conjugate-specific dehydrogenase involved in isoprene metabolism in *Rhodococcus* sp. strain AD45. *J Bacteriol* **181**: 2094–2101.
- van Hylckama Vlieg, J.E.T., Leemhuis, H., Spelberg, J.H.L., and Janssen, D.B. (2000) Characterization of the gene cluster involved in isoprene metabolism in *Rhodococcus* sp. strain AD45. *J Bacteriol* **182**: 1956–1963.
- Johnston, A. (2014) Molecular ecology of marine isoprene degradation. PhD Thesis. School of Environmental Sciences, Norwich, UK: University of East Anglia.
- Julsing, M., Rijpkema, M., Woerdenbag, H., Quax, W., and Kayser, O. (2007) Functional analysis of genes involved in the biosynthesis of isoprene in *Bacillus subtilis*. *Appl Microbiol Biotechnol* **75**: 1377–1384.
- Kirschke, S., Bousquet, P., Ciais, P., Saunoy, M., Canadell, J.G., Dlugokencky, E.J. *et al.* (2013) Three decades of global methane sources and sinks. *Nat Geosci* **6**: 813–823.
- Kottegoda, S., Waligora, E., and Hyman, M. (2015) Metabolism of 2-methylpropene (isobutylene) by the aerobic bacterium *Mycobacterium* sp. strain ELW1. *Appl Environ Microbiol* **81**: 1966–1976.
- Kuzma, J., Nemecek-Marshall, M., Pollock, W., and Fall, R. (1995) Bacteria produce the volatile hydrocarbon isoprene. *Curr Microbiol* **30**: 97–103.
- Lane, D.J. (1991) 16S/23S rRNA sequencing. In *Nucleic Acid Techniques in Bacterial Systematics*. Stackebrandt, E., and Goodfellow, M. (eds). New York, USA: Wiley, pp. 115–175.
- Leahy, J.G., Batchelor, P.J., and Morcomb, S.M. (2003) Evolution of the soluble diiron monooxygenases. *FEMS Microbiol Rev* **27**: 449–479.
- Logan, B.A., Monson, R.K., and Potosnak, M.J. (2000) Biochemistry and physiology of foliar isoprene production. *Trends Plant Sci* **5**: 477–481.
- Loivamäki, M., Mumm, R., Dicke, M., and Schnitzler, J.-P. (2008) Isoprene interferes with the attraction of bodyguards by herbaceous plants. *Proc Natl Acad Sci USA* **105**: 17430–17435.
- Loreto, F., and Fineschi, S. (2015) Reconciling functions and evolution of isoprene emission in higher plants. *New Phytol* **206**: 578–582.
- Marmur, J. (1961) A procedure for the isolation of deoxyribonucleic acid from micro-organisms. *J Mol Biol* **3**: 208–IN201.
- McAuliffe, J.C., Muir, R.E., Nielsen, A.T., Peres, C.M., Vaviline, D.V., and Wells, D.H. Enhanced production of isoprene using host cells having decreased ispA activity. US patent 8,975,051 B2, March 10, 2015.

- McLeod, M.P., Warren, R.L., Hsiao, W.W.L., Araki, N., Myhre, M., Fernandes, C. *et al.* (2006) The complete genome of *Rhodococcus* sp. RHA1 provides insights into a catabolic powerhouse. *Proc Natl Acad Sci USA* **103**: 15582–15587.
- Muyzer, G., de Waal, E.C., and Uitterlinden, A.G. (1993) Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified genes coding for 16S rRNA. *Appl Environ Microbiol* **59**: 695–700.
- Muyzer, G., Brinkhoff, T., Nuebel, U., Santegoeds, C., Schaefer, H., and Waver, C. (eds) (1998) *Denaturing Gradient Gel Electrophoresis (DGGE) in Microbial Ecology*. Dordrecht, The Netherlands: Kluwer Academic Publishers.
- Neufeld, J.D., Vohra, J., Dumont, M.G., Lueders, T., Manfield, M., Friedrich, M.W., and Murrell, J.C. (2007) DNA stable-isotope probing. *Nat Protocols* **2**: 860–866.
- Pacifico, F., Harrison, S.P., Jones, C.D., and Sitch, S. (2009) Isoprene emissions and climate. *Atmos Environ* **43**: 6121–6135.
- Pegoraro, E., Abrell, L., Van Haren, J., Barron-Gafford, G., Grieve, K.A., Malhi, Y. *et al.* (2005) The effect of elevated atmospheric CO<sub>2</sub> and drought on sources and sinks of isoprene in a temperate and tropical rainforest mesocosm. *Glob Change Biol* **11**: 1234–1246.
- Radajewski, S., Ineson, P., Parekh, N.R., and Murrell, J.C. (2000) Stable-isotope probing as a tool in microbial ecology. *Nature* **403**: 646–649.
- Schaefer, J.K., Goodwin, K.D., McDonald, I.R., Murrell, J.C., and Oremland, R.S. (2002) *Leisingera methylohalidivorans* gen. nov., sp. nov., a marine methylotroph that grows on methyl bromide. *Int J Syst Evol Microbiol* **52**: 851–859.
- Sharkey, T.D. (2013) Is it useful to ask why plants emit isoprene? *Plant Cell Environ* **36**: 517–520.
- Sharkey, T.D., Wiberley, A.E., and Donohue, A.R. (2008) Isoprene emission from plants: why and how. *Ann Bot* **101**: 5–18.
- Srivastva, N., Shukla, A.K., Singh, R.S., Upadhyay, S.N., and Dubey, S.K. (2015) Characterization of bacterial isolates from rubber dump site and their use in biodegradation of isoprene in batch and continuous bioreactors. *Bioresour Technol* **188**: 84–91.
- Tamura, K., Stecher, G., Peterson, D., Filipski, A., and Kumar, S. (2013) MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Mol Biol Evol* **30**: 2725–2729.
- Tuovinen, O.H., and Kelly, D.P. (1973) Studies on the growth of *Thiobacillus ferrooxidans*. *Arch Mikrobiol* **88**: 285–298.
- Vickers, C.E., Gershenson, J., Lerda, M.T., and Loreto, F. (2009) A unified mechanism of action for volatile isoprenoids in plant abiotic stress. *Nat Chem Biol* **5**: 283–291.
- Wiedinmyer, C., Friedfeld, S., Baugh, W., Greenberg, J., Guenther, A., Fraser, M., and Allen, D. (2001) Measurement and analysis of atmospheric concentrations of isoprene and its reaction products in central Texas. *Atmos Environ* **35**: 1001–1013.
- Wiedinmyer, C., Greenberg, J., Guenther, A., Hopkins, B., Baker, K., Geron, C. *et al.* (2005) Ozarks Isoprene Experiment (OZIE): measurements and modeling of the “isoprene volcano”. *J Geophys Res Atmos* **110**: D18307.
- Zhou, N.Y., Chan Kwo Chion, C.K., and Leak, D.J. (1996) Cloning and expression of the genes encoding the propene monooxygenase from *Xanthobacter* Py2. *Appl Microbiol Biotechnol* **44**: 582–588.

## Supporting information

Additional Supporting Information may be found in the online version of this article at the publisher's web-site:

**Table S1.** Growth of *Rhodococcus* sp. AD45, SC4 and LB1 (as sole source of carbon and energy).

**Table S2.** Summary of *Rhodococcus* strain SC4 and *Rhodococcus* strain LB1 genomes.

**Table S3.** Isoprene metabolic genes identified in strain LB1 and SC4 using coding sequences from *Rhodococcus* sp. AD45 as query sequence in tBLASTn.

**Table S4.** Strains and enrichments used for *isoA* primer validation.

**Table S5.** RFLP analysis of *isoA* clones.

**Fig. S1.** Neighbour-joining phylogenetic tree based on an alignment of 16S rRNA gene sequences of leaf and soil isolate strains LB1 and SC4, *Rhodococcus* sp. AD45 (shown in bold) and other representative strains. The tree was constructed using MEGA6 (Tamura *et al.*, 2013). All positions containing gaps and missing data were eliminated and there were 1360 nucleotide positions in the final dataset. The scale bar shows nucleotide substitutions per site. Bootstrap values (1000 replications) greater than 50% are shown at the nodes.

**Fig. S2.** Similarity of the bacterial communities derived from fully-labelled isoprene DNA-SIP enrichments and fractionation. The NMDS plot shows 16S rRNA gene-based similarity between the un-enriched soil community (time-point zero), the isoprene-enriched total community (unfractionated) and labelled (heavy) and unlabelled (light) components separated by isopycnic centrifugation and fractionation. The heavy fractions of three replicates from <sup>13</sup>C incubations were analysed separately, whereas the light fractions, unfractionated (<sup>13</sup>C) DNA and fractionated DNA from <sup>12</sup>C incubations was pooled prior to analysis. The data are based on Bray-Curtis similarity of log-transformed relative-abundance of 178 genera.

**Fig. S3.** The ratio of *Rhodococcus* 16S rRNA gene-sequence relative-abundances in heavy and light fractions of DNA extracted from <sup>12</sup>C and <sup>13</sup>C incubations. For the <sup>13</sup>C heavy fraction *n* = 3; other fractions, pooled DNA from duplicate (<sup>12</sup>C) or triplicate (<sup>13</sup>C) incubations. The bar chart shows that *Rhodococcus* sequences were not enriched in the heavy fraction of <sup>12</sup>C-isoprene incubations, but were highly enriched (over 100-fold) in the heavy fractions, compared to light fractions, of <sup>13</sup>C-isoprene incubations.

**Fig. S4.** <sup>1</sup>H-NMR spectrum of U-<sup>13</sup>C-isoprene biosynthesized by recombinant *E. coli* strain CMP1082. Supplementary Methods.