

# A mechanism for bacterial transformation of dimethylsulfide to dimethylsulfoxide: a missing link in the marine organic sulfur cycle

Ian Lidbury,<sup>1†</sup> Eileen Kröber,<sup>1†</sup> Zhidong Zhang,<sup>2</sup>  
Yijun Zhu,<sup>1</sup> J. Colin Murrell,<sup>3</sup> Yin Chen<sup>1\*</sup> and  
Hendrik Schäfer<sup>1\*\*</sup>

<sup>1</sup>*School of Life Sciences, Gibbet Hill Campus, University of Warwick, Coventry CV4 7AL, UK.*

<sup>2</sup>*Institute of Microbiology, Xinjiang Academy of Agriculture Sciences, Urumqi 830091, China.*

<sup>3</sup>*School of Environmental Sciences, University of East Anglia, Norwich Research Park, NR4 7TJ, Norwich, UK.*

## Summary

**The volatile organosulfur compound, dimethylsulfide (DMS), plays an important role in climate regulation and global sulfur biogeochemical cycles. Microbial oxidation of DMS to dimethylsulfoxide (DMSO) represents a major sink of DMS in surface seawater, yet the underlying molecular mechanisms and key microbial taxa involved are not known. Here, we reveal that *Ruegeria pomeroyi*, a model marine heterotrophic bacterium, can oxidize DMS to DMSO using trimethylamine monooxygenase (Tmm). Purified Tmm oxidizes DMS to DMSO at a 1:1 ratio. Mutagenesis of the *tmm* gene in *R. pomeroyi* completely abolished DMS oxidation and subsequent DMSO formation. Expression of Tmm and DMS oxidation in *R. pomeroyi* is methylamine-dependent and regulated at the post-transcriptional level. Considering that Tmm is present in approximately 20% of bacterial cells inhabiting marine surface waters, particularly the marine *Roseobacter* clade and the SAR11 clade, our observations contribute to a mechanistic understanding of biological DMSO production in surface seawater.**

## Introduction

Approximately 300 Tg of the volatile organosulfur compound, dimethylsulfide (DMS) is produced in the marine environment annually (Curson *et al.*, 2011), making the oceans the primary contributor to DMS in the atmosphere with an estimated flux of  $\sim 20.7$  Tg year<sup>-1</sup> (Watts, 2000). DMS production occurs through transformation of the multifunctional metabolite dimethylsulfoniopropionate (DMSP) by a variety of algal and bacterial lyases (Curson *et al.*, 2011; Todd *et al.*, 2012). Interest in the cycling of DMS increased after the proposal that it can lead to the formation of cloud condensation nuclei, increasing cloud albedo and thereby acting as a global coolant (Charlson *et al.*, 1987). Therefore, any process contributing to the removal of DMS from surface seawater will have a direct effect on the attenuation of this flux, which may in turn have significant impacts on the global sulfur cycle, specifically its deposition to the terrestrial biome. DMS loss in surface seawater is mediated through different pathways including photochemical oxidation (Brimblecombe and Shooter, 1986) and biological consumption, which includes its assimilation into biomass, dissimilation to CO<sub>2</sub> and inorganic soluble sulfur compounds, or oxidation to dimethylsulfoxide (DMSO) (Schäfer *et al.*, 2010). Importantly, DMSP, DMS and DMSO are frequently detected in the water column (Hatton *et al.*, 1998; Simó *et al.*, 2000; Vila-Costa *et al.*, 2008) and also sea ice (Asher *et al.*, 2011) at comparable values with concentrations of dissolved DMSO tending to increase towards the end of phytoplankton blooms as cells undergo senescence (Simó *et al.*, 1998; Hatton and Wilson, 2007) when DMSO concentrations can even exceed those of DMSP (Hatton *et al.*, 1998).

In surface seawater, biological consumption of DMS was shown to be a major component of the global sink for DMS, accounting for up to 99% of the DMS produced (Kiene and Bates, 1990). Tracer studies using <sup>35</sup>S-DMS suggest that only a small fraction of DMS ( $\sim 2$ –7%) is assimilated as a sulfur source by marine microorganisms whilst the majority of DMS is transformed into either DMSO or sulfate (Zubkov *et al.*, 2001; Vila-Costa *et al.*, 2006; del Valle *et al.*, 2007a), potentially *via* other inorganic sulfur compounds such as thiosulfate and tetrathionate

Received 24 February, 2016; accepted 19 April, 2016. For correspondence. \*E-mail Y.chen.25@warwick.ac.uk; Tel. 00442476528976; Fax +44 (0)24 7652 2052. \*\*E-mail H.schaefer@warwick.ac.uk; Tel. 00442476575052; Fax +44 (0)24 7652 2052. †These authors contributed equally to this study.

which are known end-products of DMS catabolism in marine DMS-degrading bacteria (De Zwart *et al.*, 1996; Boden *et al.*, 2012). In marine surface waters, oxidation to DMSO is the major fate of DMS, accounting for ~70% and 50–70% of the total oxidized DMS in the Sargasso Sea (del Valle *et al.*, 2007a) and in the Ross Sea (del Valle *et al.*, 2009) respectively. Although it has previously been suggested that DMSO production is carried out by phytoplankton (Simó *et al.*, 1998; del Valle *et al.*, 2007b), DMSO production from DMS oxidation can be stimulated by the availability of organic carbon suggesting a role of marine heterotrophic bacteria in DMSO formation. The underpinning molecular and biochemical mechanisms, however, remain poorly understood (González *et al.*, 1999; Green *et al.*, 2011; Hatton *et al.*, 2012).

Marine surface waters are frequently dominated by two groups of heterotrophic bacteria of the *Alphaproteobacteria* phylum, the marine *Roseobacter* clade (MRC) and the SAR11 clade (Morris *et al.*, 2002; Buchan *et al.*, 2005; Giebel *et al.* 2011; Gilbert *et al.*, 2012). The SAR11 clade are oligotrophic microbes that are adapted to low nutrient environments and are found throughout the world's oceans, dominating bacterial communities in the open ocean gyres (Morris *et al.*, 2002; Giovannoni *et al.*, 2005). In contrast, the MRC can account for up to 20% of bacterial cells in coastal waters during periods of elevated primary production (González *et al.*, 2000; Buchan *et al.*, 2005; Gilbert *et al.*, 2012). Combined, these two clades are major players in the uptake and metabolism of phytoplankton-derived organic carbon and nitrogen and reduced sulfur compounds in marine surface waters (Miller and Belas, 2004; Alonso and Pernthaler, 2006; Sowell *et al.*, 2008; Sun *et al.*, 2011; Williams *et al.*, 2012; Gifford *et al.*, 2013; Ottesen *et al.*, 2013; Aylward *et al.*, 2015; Lidbury *et al.*, 2015a). The abundance of MRC in surface seawater is often positively correlated with elevated levels of DMSP, DMS and DMSO (González *et al.*, 2000; Zubkov *et al.*, 2001; Hatton *et al.*, 2012; Nelson *et al.*, 2014; Cui *et al.*, 2015).

We have previously shown that bacterial trimethylamine monooxygenase (Tmm), which is responsible for the oxidation of trimethylamine (TMA) into trimethylamine *N*-oxide (TMAO), can also oxidize DMS at comparable rates to that of TMA (Chen *et al.*, 2011). Specifically, purified Tmm from both the SAR11 clade and MRC can efficiently oxidize DMS. The gene encoding Tmm occurs in ~20% of bacterial cells inhabiting the surface waters of the oceans (Chen *et al.*, 2011), and is particularly common in bacteria of the MRC and SAR11 clade. In the Sargasso Sea, a region where biological DMSO production occurs (del Valle *et al.*, 2007a), TMA was rapidly oxidized to CO<sub>2</sub> (Sun *et al.*, 2011) and in the Antarctic Ocean, both concentrations of TMAO and DMSO were significantly higher than either TMA or DMS, suggesting that cooxidation of these compounds was taking place at the time of sampling (Gibb and


Hatton, 2004). We, therefore, hypothesized that a significant proportion of the observed DMSO formation in surface seawater is due to the oxidation of DMS by Tmm-containing heterotrophic bacteria. In this study, we describe for the first time DMSO formation in the model marine bacterium, *Ruegeria pomeroyi* DSS-3 (González *et al.*, 2003) and demonstrate that methylated amine-dependent DMSO formation is a common trait of the MRC.

## Results

### *Oxidation of DMS to DMSO by Ruegeria pomeroyi* DSS-3

We previously showed that purified Tmm from selected MRC and SAR11 bacteria can oxidize DMS with an affinity similar to that of its native substrate, TMA (Chen *et al.*, 2011). In order to determine the product of DMS oxidation by Tmm, we purified Tmm of *R. pomeroyi* from a recombinant *E. coli* expressing Tmm and showed that DMS oxidation resulted in the formation of DMSO at a 1:1 ratio (Fig. 1A). Unlike TMAO, which can be further degraded in this bacterium through a TMAO demethylase (Tdm, E.C.4.1.2.32) (Lidbury *et al.*, 2014; Zhu *et al.*, 2014), DMSO is not a substrate for Tdm (Fig. 1B), suggesting that DMSO is the end-product of DMS co-oxidation by Tmm (Fig. 1C).

We then investigated whether DMSO could be formed from DMS oxidation by *R. pomeroyi*. It has previously been reported that *R. pomeroyi* can oxidize DMS (González *et al.*, 1999), however, no significant DMS oxidation was observed when it was grown on glucose and ammonium as the sole carbon and nitrogen source, respectively, (Fig. 2A). Under these growth conditions, Tmm activity was not detectable (Chen *et al.*, 2011). TMA has previously been shown to be both a nitrogen and energy, but not a carbon source for *R. pomeroyi* (Chen *et al.*, 2011; Lidbury *et al.*, 2014). In order to induce Tmm expression, we incubated a *R. pomeroyi* starter culture supplemented with 0.25 mM TMA in addition to glucose and ammonium (Chen *et al.*, 2011) and used this as the inoculum for induction experiments. Subsequently, complete DMS oxidation occurred within 42 h (Fig. 2B). All the DMS (0.8 mM) was transformed into DMSO (0.81 ± 0.03 mM) indicating that *R. pomeroyi* does not further catabolize DMSO under these growth conditions. To further confirm that DMS oxidation was due to the activity of Tmm *in vivo*, we tested the *tmm* knockout mutant ( $\Delta tmm::Gm$ ) (Lidbury *et al.*, 2015b), which can no longer utilize TMA, under the same conditions. As expected, no DMS oxidation was detected throughout the growth cycle (Fig. 2C). The essential role of Tmm in DMS oxidation to DMSO was further validated by complementation of the *tmm* knockout mutant with either *tmm* from *R. pomeroyi* or *Roseovarius* sp. 217 (Schäfer *et al.*, 2005). Complementation of the  $\Delta tmm::Gm$  mutant with either *tmm* homolog restored both growth on TMA as




**Fig. 1.** Co-oxidation of DMS to DMSO by Tmm.

A. Trimethylamine monoxygenase (Tmm) of *R. pomeroyi* DSS-3 oxidizes DMS to DMSO at a 1:1 ratio.

B. The substrate specificity of purified trimethylamine *N*-oxide demethylase (Tdm) of *R. pomeroyi*. Error bars denote s.d. from triplicate reactions.

C. TMA oxidation and subsequent metabolism by Tmm, Tdm and the enzymes involved in the gammaglutamylmethylamide/*N*-methylglutamate pathway (Chen, 2012) results in the release of ammonium as a nitrogen source for this bacterium. However, DMSO, the product from the co-oxidation of DMS by Tmm, appears to be the end product of this pathway because it is not a substrate for Tdm.

TMA, trimethylamine; TMAO, trimethylamine *N*-oxide; DMA, dimethylamine; MMA, monomethylamine,  $\text{NH}_4^+$ , ammonium.



**Fig. 2.** DMS oxidation in the cultures of *R. pomeroyi* DSS-3 wild type and a *tmm* knockout mutant. (A) No decrease in DMS concentration (white diamonds) was observed in cultures of wild-type cells pre-incubated with glucose and ammonium only. (B) Complete oxidation of DMS (white diamonds) to DMSO occurred when wild-type cells were pre-incubated with TMA (250  $\mu\text{M}$ ) as well as glucose and ammonium. (C) DMS oxidation (white diamonds) did not occur in the mutant,  $\Delta tmm::Gm$ , despite pre-incubation with TMA, glucose and ammonium (250  $\mu\text{M}$ ).

Complementation of the  $\Delta tmm::Gm$  mutant with either native *tmm* of *R. pomeroyi* DSS-3 or *tmm* from *Roseovarius* sp. 217 restored its growth on TMA as a sole nitrogen source (D) and TMA-dependent DMS oxidation (E). No effect on the growth rate or final growth yield was observed for *R. pomeroyi* cultures grown in the presence (white circles) or absence (grey circles) of DMS (0.8 mM). Results are the mean of triplicate cultures. Error bars denote s.d.


a sole nitrogen source (Fig. 2D) and TMA-dependent DMS oxidation (Fig. 2E). Collectively, our data confirm that DMS can be oxidized to DMSO in *Ruegeria pomeroyi* through Tmm, which can be induced by TMA.

#### TMA attenuates DMS production during growth of *R. pomeroyi* on DMSP

Next, we investigated whether *R. pomeroyi* can also produce DMSO during growth on DMSP. DMSP is an important osmolyte and a precursor for DMS, which is common in seawater (Kiene *et al.*, 2000). *R. pomeroyi* was isolated using DMSP as the sole carbon source from seawater collected off the coast of Georgia during an algal bloom (González *et al.*, 1999; 2003). *R. pomeroyi* can use DMSP as the sole carbon source and it is known to produce DMS directly from the cleavage of DMSP using various DMSP lyases (Curson *et al.*, 2011; Todd *et al.*, 2012). When the wild type and the  $\Delta tmm$  mutant were grown on ammonium and DMSP (5 mM) as the sole nitrogen and carbon sources, respectively,  $\sim 1.4$  mM of DMS accumulated in the culture and was not subjected to further degradation (Fig. 3A and B). To determine whether TMA had an impact on the production and subsequent conversion of DMS to DMSO, *R. pomeroyi* was grown on DMSP (5 mM) as the sole carbon source with TMA (0.25 mM) as the sole nitrogen source. The  $\Delta tmm$  mutant failed to grow on TMA as the sole nitrogen source (Fig. 3B); consequently, only 0.2 mM DMS was produced (Fig. 3B). Given the fact that no DMS oxidation occurred when this mutant was grown on glucose and ammonium after pre-incubation with TMA (Fig. 2C) it was not surprising that none of the DMS produced was converted to DMSO. Wild-type TMA-grown cells showed a 27% reduction of maximal DMS concentrations in the headspace during initial growth (within 100 h) compared to that in the ammonium-grown cultures (Fig. 3A). This was followed by near complete oxidation of DMS to DMSO in TMA-grown cells (Fig. 3A). These data therefore suggest that in the presence of TMA, DMSO represents a major metabolic end product of DMSP catabolism in *R. pomeroyi*.

#### Co-oxidation of DMS can also be induced by the metabolites of TMA degradation in *R. pomeroyi*


Considering that individual methylated amines species [monomethylamine (MMA), dimethylamine (DMA), TMA or TMAO] rarely exist in isolation in the marine environment (Gibb *et al.*, 1999; Gibb and Hatton, 2004), we investigated whether Tmm activity and hence DMS oxidation occurred in the presence of other methylated amines e.g. MMA, DMA or TMAO. The wild-type and the  $\Delta tmm$  mutant of *R. pomeroyi* were grown on methylated amines which are metabolites of TMA degradation (TMAO, DMA and MMA) as a source of



**Fig. 3.** DMS oxidation to DMSO in *R. pomeroyi* DSS-3 grown on DMSP as the main carbon substrate. Quantification of DMS (diamonds) during growth of *R. pomeroyi* (circles) on either ammonium (grey) or TMA (white) as the sole nitrogen source (0.25 mM) and DMSP (5 mM) as the main carbon substrate. Either wild-type (A) or the mutant,  $\Delta tmm::Gm$  (B), cells were used for incubations. Results are the mean of triplicate cultures. Error bars denote s.d.

nitrogen and DMS consumption was quantified. For the wild-type cells, in addition to TMA, growth on TMAO and DMA as a nitrogen source also led to the oxidation of DMS to DMSO (Fig. 4A). As expected, the  $\Delta tmm$  mutant failed to oxidize DMS despite growth on these metabolites of TMA metabolism (Fig. 4B) further confirming that Tmm is solely responsible for DMS oxidation in *R. pomeroyi*.

To investigate the transcriptional regulation of *tmm*, and hence DMS oxidation to DMSO by Tmm, the sensitivity of the promoter of *tmm* ( $P_{tmm}$ ) in response to methylated compounds was determined using a *lacZ* reporter assay (Todd *et al.*, 2012). A *R. pomeroyi* transconjugant, containing the  $P_{tmm}$ -*lacZ* fusion plasmid, was grown overnight on a minimal medium containing glucose and ammonium and



**Fig. 4.** Methyamine-dependent DMS oxidation in *R. pomeroyi* DSS-3. Quantification of DMS during growth of *R. pomeroyi* wild type (A) or the mutant  $\Delta tmm::Gm$  (B) on different methylated amines as a sole nitrogen source. Results are mean of triplicate cultures. Error bars denote s.d.


supplemented with TMA, TMAO, DMA, MMA, DMS or DMSO (0.5 mM) before assaying for the reporter,  $\beta$ -galactosidase activity (Lidbury *et al.*, 2014). Compared with controls with NH<sub>4</sub><sup>+</sup>, incubation with TMA and DMA led to the induction of the  $P_{tmm}$ -lacZ fusion giving twofold and threefold levels of  $\beta$ -galactosidase activity (Fig. 5A). Strikingly, DMS alone resulted in a  $\sim$ 7-fold induction of the  $P_{tmm}$ -lacZ fusion, however, *R. pomeroyi* does not oxidize DMS without methylated amines in the defined medium (Fig. 4A). RT-PCR assays targeting *tmm* confirmed that the presence of DMS does indeed result in the upregulation of *tmm* transcription (Fig. 5C) despite there being no detectable enzyme activity of Tmm (Fig. 5B). No apparent difference in the expression of the 16S rRNA gene was detected between either the control cultures or those cultures supplemented with either TMA or DMS (Fig. 5C). Furthermore, no detectable Tmm activity was observed in cells grown on TMAO as a nitrogen source, but the presence of both TMAO and DMS resulted in the induction of Tmm activity (Fig. 5B). These results confirm that TMA, DMA, as well as DMS are capable of inducing *tmm* transcription. However this induction of *tmm* at the transcriptional level does not necessarily result in a functional Tmm.

*The transcriptional regulator TmoR is required for tmm transcription, however production of a functional Tmm is regulated post-transcriptionally in R. pomeroyi*

The lacZ reporter assays suggested that *tmm* transcription in wild type cells of *R. pomeroyi* is induced by TMA and other methylated amine intermediates, such as DMA. Transcriptional regulation of *tmm* is predicted to be regulated

by a putative GntR repressor encoded by SPO1553, containing a conserved pfam00392 domain, which is found in GntR-like proteins. This putative GntR-like repressor (hereafter named as TmoR) was therefore targeted for mutagenesis, generating the mutant,  $\Delta tmoR::Gm$ . Results from  $P_{tmm}$ -lacZ reporter assays in the  $\Delta tmoR::Gm$  mutant (Fig. 5A) demonstrate that *tmm* is constitutively expressed in this mutant, confirming that TmoR functions as a repressor for *tmm*. RT-PCR assays confirmed that *tmm* is constitutively expressed even in the absence of methylated amines or DMS (Fig. 5C). Despite constitutive expression of *tmm* in the  $\Delta tmoR::Gm$  mutant, this strain still could not oxidize DMS in the absence of methylated amines (Fig. 5D), but could oxidize DMS in their presence. These results, therefore, point to post-transcriptional regulation of Tmm activity, and hence mediation of DMS oxidation to DMSO by methylated amines.

To better understand which methylated amines are responsible for post-transcriptional regulation of Tmm activity, we performed DMS oxidation experiments using the  $\Delta tdm::Gm$  mutant, which is unable to demethylate TMAO to DMA (Lidbury *et al.*, 2014). When grown under similar conditions to the wild-type, i.e. the addition of TMA to the starter culture, no DMS oxidation was observed (Fig. 6A). When the same experiment was performed using the *tdm*-complemented strain, which has a restored ability to demethylate TMAO to DMA (Lidbury *et al.*, 2014), complete oxidation of DMS occurred (Fig. 6A). This suggests that formation of a fully functional Tmm is probably dependent on the presence of DMA. To test this hypothesis, we grew the  $\Delta tdm$  mutant (using a starter culture



**Fig. 5.** Transcriptional regulation of *tmm* in *R. pomeroyi*.

A. *lacZ* reporter assays using the promoter of *tmm* ( $P_{tmm}$ ) show that  $P_{tmm}$  is activated by DMS as well as TMA and that knocking out the repressor gene (*tmoR*) leads to constitutive transcription of  $P_{tmm}$ .

B. No Tmm activity was detected when *R. pomeroyi* was exposed to DMS without methylated amines present.

C. Reverse transcription-PCR results show that both DMS and TMA led to the up-regulation of transcription of *tmm* (compared to the levels of 16S rRNA transcription) in the wild type strain whereas in the *tmoR* mutant, *tmm* transcription appears to be constitutive. Minus sign indicates negative control without reverse transcriptase.



D. DMS consumption in the  $\Delta tmoR$  mutant in the presence of various methylamines. Results presented in panels A, B and D are mean of triplicate cultures. Error bars denote s.d.

supplemented with TMA to induce *tmm*) using ammonium as a nitrogen source and supplemented it with TMA, TMAO or DMA (0.5 mM). Indeed, the rate of DMS oxidation was significantly greater in the presence of DMA (1.38 mM d<sup>-1</sup>) compared with that of cultures grown in the presence of either TMA (0.32 mM d<sup>-1</sup>) or TMAO (0.22 mM d<sup>-1</sup>) or ammonium alone (0 mM d<sup>-1</sup>) (Fig. 6B), suggesting that, out of the compounds tested in this study, DMA is mainly responsible for post-transcriptional regulation of Tmm activity.

#### Methylamine-dependent DMS co-oxidation is a common trait of MRC bacteria

In order to investigate whether methylamine-dependent DMS co-oxidation is unique in *Ruegeria pomeroyi* DSS-3,

we further tested a number of cultivated bacteria of the MRC clade. The data presented in Table 1 showed that several members of the MRC that are capable of utilizing TMA, including *Roseovarius* sp. TM1035, *Roseobacter litoralis* Och149, also displayed methylamine-dependent DMS oxidation. No DMS oxidation occurred unless cultures were supplemented with methylated amines such as TMA and TMAO. Those isolates containing *tmm* also oxidized DMS when grown in a complex medium (1/2 YTSS), probably due to the presence of amines in this complex medium. Furthermore, two new strains, *Phaeobacter* sp. TMAL401 and *Roseobacter* sp. TMAL402, which were isolated during this study on TMA as a sole carbon source from coastal seawater off the coast of Plymouth also showed methylamine-dependent DMS oxidation (Table 1). Together these data suggest that methylamine-dependent



**Fig. 6.** Methylated amine-dependent DMS oxidation in the *R. pomeroyi* mutant,  $\Delta tdm::Gm$ . The  $\Delta tdm$  mutant (white) and the complemented  $\Delta tdm$  mutant (grey) grown on glucose as a sole carbon and energy source and ammonium as a sole nitrogen source with TMA (0.25 mM) added to the starter culture (A). DMS oxidation rates calculated for the  $\Delta tdm$  mutant when grown on glucose and ammonium and supplemented with either TMA, TMAO or DMA (B). Results are mean of triplicate cultures. Error bars denote s.d.

co-oxidation of DMS may be a ubiquitous trait of TMA-utilising heterotrophic bacteria.

## Discussion

Here we present evidence that the oxidation of DMS by Tmm is responsible for DMSO formation in the heterotrophic bacterium, *R. pomeroyi* and other closely-related members of the MRC that harbour the *tmm* gene. The MRC is frequently associated with phytoplankton blooms and often represents a major proportion of the active community involved in the turnover of algal-derived organic matter, including DMSP (González *et al.*, 2000; Vila *et al.*, 2004; Buchan *et al.*, 2005; Alonso and Pernthaler, 2006). Our results reveal that in *R.*

*pomeroyi* the DMS produced by DMSP lyases (Curson *et al.*, 2011; Todd *et al.*, 2012) during growth on DMSP can be further oxidized to DMSO by Tmm in the presence of methylated amines. We therefore speculate that this molecular mechanism of DMS oxidation likely explains the frequently observed link between increased phytoplankton activity, dissolved organic matter including DMSP, abundance of MRC cells and elevated concentrations of DMSO (Hatton *et al.*, 1998; 1999; 2012; Zubkov *et al.*, 2001; 2004; Hatton and Wilson, 2007; Green *et al.*, 2011). Further research measuring the concentrations of methylated amines during periods of elevated primary production and phytoplankton blooms is required to better understand the role of Tmm-containing heterotrophs in the production of DMSO.

**Table 1.** A comparison of dimethylsulfide consumption among different members of the MRC when grown on different media.

Strain <sup>1</sup>	<i>tmm</i> <sup>2</sup>	NH <sub>4</sub> <sup>+</sup> <sup>3</sup>	TMA <sup>3</sup>	TMAO <sup>3</sup>	1/2 YTSS
<i>Ruegeria pomeroyi</i> DSS-3 wild type	+	–	+	+	a
$\Delta tmm::Gm$	–	–	–	+	–
$\Delta tmm::Gm + DSS-3:tmm$	+	–	+	NT	NT
$\Delta tmm::Gm + 217:tmm$	+	–	+	NT	NT
$\Delta tmoR::Gm$	+	–	+	+	NT
$\Delta tdm::Gm$	+	–	Reduced	Reduced	NT
$\Delta tdm::Gm + DSS-3:tdm$	+	–	+	+	NT
<i>Roseovarius</i> sp. 217	+	–	+	+	+
<i>Roseovarius</i> sp. TM1035	+	–	+	+	+
<i>Roseovarius nubinihibens</i> ISM	+	–	+	+	+
<i>Roseobacter litoralis</i> Och 149	+	–	+	+	+
<i>Phaeobacter</i> sp. TMAL401	+	a	+	+	+
<i>Roseobacter</i> sp. TMAL402	+	–	+	+	+
<i>Roseobacter</i> sp. MED193	–	–	–	–	–

a. Indicates variable DMS oxidation.

<sup>1</sup>Incubations were performed in triplicate supplemented with 1 mM DMS.

<sup>2</sup>Presence (+) or absence (–) of the *tmm* gene in the sequenced genome.

<sup>3</sup>Defined medium supplemented with different nitrogen sources.

NH<sub>4</sub><sup>+</sup>, ammonium; TMA, trimethylamine; TMAO, trimethylamine *N*-oxide; 1/2 YTSS, half strength yeast extract-tryptone-sea salt medium.


It was previously shown that *tmm* is abundant in metagenomes of marine surface waters, due to the occurrence of members of the MRC and the SAR11 clade (Chen *et al.*, 2011). Whether or not SAR11 clade bacteria can perform methylated amine-dependent DMS oxidation remains to be established, however purified recombinant Tmm from strains of the SAR11 clade can oxidize DMS in a similar manner to that of Tmm from MRC (Chen *et al.*, 2011). It is also known that *Candidatus Pelagibacter ubique* HTCC1062, a Tmm-containing representative of the SAR11 clade, can oxidize TMA to stimulate ATP production (Sun *et al.*, 2011). Furthermore, addition of TMA to natural Sargasso Sea water resulted in its complete oxidation (Sun *et al.*, 2011). In oligotrophic waters, such as the Sargasso Sea, SAR11 clade bacteria represent the numerically dominant proportion of the heterotrophic community (Morris *et al.*, 2002, Sjöstedt *et al.*, 2014) and peptides mapping to Tmm from SAR11 have been detected in this region (Sowell *et al.*, 2008). Therefore, this clade may be largely responsible for the organic carbon-stimulated increase in DMSO production (~70–90%) from DMS oxidation in the Sargasso Sea (Vila-Costa *et al.*, 2006).

As Tmm can oxidize DMS to DMSO, understanding the regulation of this enzyme is vital in order to determine the factors that control the loss of DMS from marine surface waters. Our data reveal that Tmm expression is controlled at both the transcriptional (through the repressor, TmoR) and post-transcriptional level because the TmoR knockout mutant did not have a methylamine-independent DMS oxidation phenotype (Fig. 5D). Post-transcriptional regulation in bacteria can be mediated by the occurrence of *cis*-acting antisense RNA molecules (asRNA) which can modulate translation of a given target gene through a variety of mechanisms (Georg and Hess, 2011; Sesto *et al.*, 2013). In the cyanobacterial strains, *Synechocystis* sp. PCC6803 and *Prochlorococcus* sp. MED4, asRNA can help promote translation through interactions with the primary gene transcript (Stazic *et al.*, 2011; Sakurai *et al.*, 2012). Interestingly, the diazotrophic cyanobacterium, *Trichodesmium erythraeum* IMS101, has a *tmm* homolog (53.6% identity to that of *R. pomeroyi*) and an associated asRNA molecule (Pfreundt *et al.*, 2014) and we also detected a putative promoter ( $P_{astmm}$ ) located on the antisense strand of *tmm* within the coding region (Fig. 7). We therefore propose a model for Tmm regulation in *R. pomeroyi* (Fig. 7), which includes the presence of an asRNA molecule that helps to stabilize the primary transcript of *tmm*, thus protecting it from degradation by RNases. Based on our experimental data, we hypothesize that transcription of the asRNA molecule (*astmm*) is induced by DMA through an as yet unknown regulator. In support of this model, RT-PCR assays targeting the antisense strand of *tmm* revealed the presence of a potential

asRNA molecule that is upregulated in the presence of TMA (Supporting Information Fig. S1). Although other posttranscriptional and posttranslational regulation mechanisms of Tmm activity may also occur in this bacterium, such as protein modification and bacterial riboswitches (Bastet *et al.*, 2011), further research regarding the possible involvement of asRNA on Tmm activity is certainly warranted to better understand the regulation of this ecologically important enzyme.

DMS oxidation to DMSO can enhance chemoorganoheterotrophic growth in some marine bacteria (Boden *et al.*, 2011; Green *et al.*, 2011), but the molecular mechanism for the increase in growth yield in these isolates is unknown. The phototrophic bacterium *Rhodovulum sulfidophilum* can utilize DMS to DMSO (performed by DMS dehydrogenase) as an electron donating reaction for photoautotrophic growth (McDevitt *et al.*, 2002), however oxidation of DMS by Tmm does not result in any net gain of electrons or reducing equivalents. Indeed, Tmm actually consumes NADPH for the conversion of TMA to TMAO and DMS to DMSO (Chen *et al.*, 2011). This begs the question ‘why do these bacteria possess an enzyme that shares a similar high affinity for two differing substrates?’ The FAD-binding site of Tmm and closely related flavin-containing monooxygenases (FMOs) has been likened to a ‘cocked gun’ often present in a reduced state within the cell, awaiting a suitable substrate (Krueger and Williams, 2005). Bacterial and mammalian FMOs produce both superoxide anion radicals and hydrogen peroxide through spontaneous NADPH oxidation in the absence of these substrates (Williams *et al.*, 1985; Krueger and Williams, 2005; Alfieri *et al.*, 2008). Indeed, in FMO purified from rabbit liver, up to 41% of the total NADPH oxidized resulted in hydrogen peroxide production (Tynes *et al.*, 1986). Co-oxidation of DMS by Tmm may therefore have a role in the prevention of free radical formation and subsequent oxidative stress when TMA stocks are depleted. In addition, DMSO can accumulate to higher concentration within the cell compared with DMS (Simó *et al.*, 2000) and does indeed act as a highly effective antioxidant in a number of algal species during times of physiological stress (Lee and De Mora, 1999; Sunda *et al.*, 2002; Riseman and DiTullio, 2004; Bucciarelli *et al.*, 2013).

In conclusion, we show that marine heterotrophic bacteria of the MRC can rapidly oxidize DMS to DMSO using the enzyme Tmm. The widespread occurrence of this enzyme in key microbial taxa that are ubiquitous in the marine environment represents a significant molecular mechanism for the oxidation of this volatile compound to DMSO in the marine surface waters. We speculate that this methylamine-dependent co-oxidation pathway of DMS may be a significant route for DMSO production in the oceans.



**Fig. 7.** Proposed model of transcriptional and post-transcriptional regulation of trimethylamine monoxygenase (Tmm) in *R. pomeroyi*. A. The transcriptional regulator TmoR acts as a repressor of *tmm*. In the presence of TMA and DMS, TmoR is released from the promoter ( $P_{tmm}$ ), allowing transcription of *tmm*. A promoter on the antisense strand of *tmm* ( $P_{astmm}$ ) is predicted. Transcription of the antisense regulatory RNA (asRNA) is directly regulated by DMA. B. The role of the putative asRNA is likely to stabilize *tmm* transcript by preventing its degradation by RNases. In the absence of methylated amines, DMS leads to the transcription of *tmm*, but not asRNA, resulting in the degradation of *tmm* prior to translation. However, in the presence of methylated amines and DMS, both *tmm* and *astmm* are transcribed which subsequently form a duplex that protects *tmm* transcript from nuclease degradation. TmoXWV encodes an ABC-type transporter for TMAO. SPO1552 is annotated as an unknown periplasmic protein. SPO1554 encodes a putative ammonium transporter.

## Experimental procedures

### Cultivation of *Ruegeria pomeroyi* DSS-3 and other MRC isolates

A complete list of *R. pomeroyi* DSS-3 strains used in this study can be found in Tables 1 and Supporting Information S1. All bacterial strains were maintained on 2216 marine broth (Difco) agar plates (1.5% w/v). For growth experiments, *R. pomeroyi* was grown at 30°C in 125-ml serum vials in triplicate using the defined marine ammonium mineral salts (MAMS) medium using a 5% (v/v) inoculum. Vitamins were added as described previously (Chen, 2012). Glucose (10 mM) or DMSP (5 mM) was used as the main carbon substrate and either ammonium or TMA as a sole nitrogen source (0.5 mM). Starter cultures were grown in glucose and ammonium in the presence or absence (+/–) of TMA (0.5 mM). Other MRC isolates were grown in triplicate at either 30°C or 18°C in 125-ml serum vials with rubber stoppers in MAMS using a 5% inoculum (v/v). Ammonium ( $\text{NH}_4^+$ ), TMA, TMAO, DMA or MMA (0.5 mM) was used as the sole N source. Succinate (5–10 mM) was used as the sole carbon source +/- DMS (0.5 mM). Two MRC isolates (*Phaeobacter* sp. TMAL401,

*Roseobacter* sp. TMAL402) isolated from seawater collected from the L4 sampling station, off the coast of Plymouth (Devon, UK) in June 2012, were also used in this study. The seawater samples were enriched using a defined medium as described previously (Chen, 2012) with TMA (0.5 mM) as the sole carbon source. These isolates were obtained by plating out liquid enrichments in increasing dilutions onto MAMS agar plates with TMA (3 mM) as the sole carbon source.

### Quantification of methylated amines, DMS, DMSP and DMSO

Quantification of methylated amines and ammonium was achieved by cation-exchange ion chromatography with a Metrosep C4/250-mm separation column and a conductivity detector (Metrohm) as described previously (Lidbury *et al.*, 2014). The eluent used for separation was prepared as follows (10× stock solution): double distilled water ( $\text{ddH}_2\text{O}$ ) up to 1 L, nitric acid (1.4 M), 500 ml acetone, 350 ml 2, 4-pyridinedicarboxylic acid monohydrate. Eluent stock solution was stored at 4°C and diluted with  $\text{ddH}_2\text{O}$  prior to use.

*R. pomeroyi* cultures incubated with DMS were grown in 125-ml serum vials and sealed with rubber stoppers. Quantification of DMS in headspace gas was measured by injecting 100 µl of a headspace gas sample into a Shimadzu GC-2010 Plus gas chromatograph (Shimadzu Corporation, Columbia, USA) fitted with a 30 m × 0.32 mm ID SHIM-1 × 3.0 µm capillary dimethylpolysiloxane column (Shimadzu Corporation, Columbia, USA.). Helium was used as the carrier gas (column flow rate, 2 ml min<sup>-1</sup>) and the column temperature was 180°C. A flame photometric detector was used to detect DMS. DMS concentrations were calculated by regression analysis based on a five-point calibration with standard DMS solutions in MAMS.

DMSP concentrations in the cultures were measured as DMS following alkaline hydrolysis as reported previously (Miller and Belas, 2004). An aliquot of the culture sample was added to a 25-ml serum vial with the addition of the same volume of either 5 M NaOH or distilled water. Solutions of pure DMSP at 0.1–1 mM dissolved in distilled water were prepared along with each experiment. After overnight incubation, DMS resulting from alkaline hydrolysis of DMSP was measured as described above. DMSO was determined according to a minor modification of the method of Jonkers (Jonkers *et al.*, 1996): DMSO was measured as DMS after reduction with acidified stannous chloride (20 g SnCl<sub>2</sub> in 100 ml 37% HCl, v/v) for 90 min at 55°C.

#### Activity assay of DMS oxidation to DMSO by *Tmm*, DMSO demethylation by *Tdm* and TMA oxidation to TMAO by *Tmm*

*E. coli* BLR (DE3) expressing either *Tmm* from *R. pomeroyi* DSS-3 or *Tdm* from *R. pomeroyi* was used for activity assays (Chen *et al.*, 2011; Lidbury *et al.*, 2014). *E. coli* cells were grown at 37°C to an OD<sub>540</sub> of 0.6, and isopropyl β-D-1-thiogalactopyranoside (IPTG) was then added to a final concentration of 0.2 mM. Cells were broken by passing three times through a French pressure cell (American Instrument) at 110 MPa. Cell debris was removed by centrifugation at (21 000 × g) for 15 min. Overexpressed *Tmm* was purified using a His-tag protein purification kit (Novagen) as described in the manufacturer's protocol (Merck KGaA, Darmstadt, Germany). *Tmm* activity was measured by following the decrease in absorbance at 340 nm of NADPH (Sigma-Aldrich). Enzyme assays were performed in triplicate at 22°C. A 2-ml mixture contained 0.26 mg purified enzyme, 10 mM PIPES (pH 7.6), and 0.25 mM NADPH. The reaction was initiated by adding the substrate, and the decrease in NADPH and DMS was measured via UV-visible spectrophotometer and gas chromatography respectively. Enzyme assays were carried out with varying concentrations of DMS (0.05 mM, 0.1 mM, 0.15 mM, 0.2 mM, 0.25 mM, 0.3 mM, 0.35 mM, 0.4 mM). DMSO concentrations were then determined as described above. *Tdm* activity assays were performed as previously described (Zhu *et al.*, 2014).

#### Construction of the mutant, $\Delta tmoR::Gm$

To construct a *tmoR* mutant, a region towards the 5' end (with *Pst*I and *Xba*I sites engineered in) and a region towards the 3'

end (with a *Hind*III and *Xba*I engineered in) of the target gene (*Spo1553*) was amplified. The two regions, along with a gentamicin gene cassette inserted at an *Xba*I site between the two regions, were cloned into the cloning vector, pGEM-T (Promega). Primers used for PCR amplification are listed in Supporting Information Table S2. The entire construct was ligated into the suicide vector pK18mobsacB, harbouring a kanamycin resistance cassette, at sites *Pst*I and *Hind*III. The resulting plasmid was transformed into *E. coli* S17-1 via electroporation and mobilized into *R. pomeroyi* via conjugation, using 1/2 YTSS as the medium (DSMZ). Transconjugants were selected for on the sea salts minimal medium with gentamicin (10 µg ml<sup>-1</sup>) and MMA (3 mM) as a sole nitrogen source. Double crossover mutants were selected by their sensitivity to kanamycin (80 µg ml<sup>-1</sup>) and homologous recombination was confirmed by PCR and subsequent DNA sequencing.

#### Construction of the $P_{tmm}::lacZ$ fusion report probe and quantification of β-galactosidase activity

The 250 bp 5' untranslated region of *tmm* (SPO1551) of *R. pomeroyi* was cloned into the reporter probe plasmid, pBIO1878 (Todd *et al.*, 2012) using the restriction sites *Kpn*I and *Pst*I. The plasmid was mobilized into *R. pomeroyi* wild type and the  $\Delta tmoR::Gm$  mutant by conjugation with *E. coli* S17-1. A mixed cell suspension was plated onto minimal medium plates containing MMA as the sole nitrogen source plus spectinomycin (175 µg ml<sup>-1</sup>) to select against *E. coli* S17-1. Activity assay for β-galactosidase was performed as described in the Supporting Information (Lidbury *et al.*, 2014).

#### Complementation of the mutant, $\Delta tmm::Gm$ , with *tmm* of *R. pomeroyi* or *Roseovarius sp. 217*

The  $\Delta tmm::Gm$  mutant was constructed previously (Lidbury *et al.*, 2014). To complement this mutant, the promoter of *tmm* from *R. pomeroyi* DSS-3 was amplified with an *Xba*I and an *Nde*I site engineered into the 5' and 3' end, respectively, and sub-cloned into pGEM-T. The promoter sequence was then released via enzymatic digestion and inserted into the pET28a plasmid, containing the *tmm* from *R. pomeroyi* or *Roseovarius sp. 217* (Chen *et al.*, 2011). The combined construct was released from pET28a and inserted into the broad-range plasmid, pBBR1MCS-km at sites *Xba*I/*Eco*RI. The resulting plasmids were transformed via electroporation into *E. coli* S17-1 and then mobilized into the *R. pomeroyi*  $\Delta tmm::Gm$  mutant via conjugation. Transconjugants were selected for using kanamycin (80 µg ml<sup>-1</sup>) as described previously (Lidbury *et al.*, 2014). The  $\Delta tdm::Gm$  mutant and the complemented *tdm* mutant were constructed previously (Lidbury *et al.*, 2014).

#### Extraction and amplification of nucleic acids

All DNA extractions were performed using the FastDNA™ SPIN Kit for Soil (MP Biomedicals, LLC, CA, USA) according to the manufacturer's instructions. For cultivated bacterial isolates, 1–5 ml of liquid culture was centrifuged (8000 × g for 5 min) to generate a cell pellet. For RNA work, all glassware, water and solutions were treated with diethylpyrocarbonate

(DEPC) (or prepared with DEPC-treated water where appropriate) by shaking overnight at 37°C in a 0.1% (v/v) solution prior to autoclaving. Total RNA was isolated from *R. pomeroyi* using the hot acid-phenol method of Gilbert and colleagues. (Gilbert *et al.*, 2000). The quality of the RNA was analysed by running 1–5 µl on a 1% (w/v) TBE-agarose gel. DNA was removed by one treatment using RNase-free DNase (Promega) and the DNA-free RNA was purified using an RNeasy spin column (Qiagen, Crawley, UK) following the manufacturer's instructions (RNA-clean up protocol). Removal of DNA was confirmed by the absence of a PCR product of the 16S rRNA gene, using the primer set 341F/518R (Supporting Information Table S2) in reactions using 1 µl or 4 µl (150–200 ng of RNA equivalent) of RNA template undergoing 35 cycles of PCR. In addition, minus (–) reverse transcriptase (RT) controls were performed in parallel to cDNA library preparation to ensure no DNA contamination was affecting the results from RT-PCR. PCR amplification of *tmm* cDNA was performed using the primers *tmm*\_RTF: 5'-CCGGCTACAAG-CATTTCTTC-3'/*tmm*\_RTR: 5'-GATGTCTTCGCCCTTGTGTT-3'. 150 ng of cDNA was used as a template. Conditions for PCR amplification were as follows: 3 min denaturation step at 95°C, followed by 30 cycles of, 95°C for 1 min, an annealing step (45–60°C dependent on primers) of 30 s, an elongation step at 72°C for an appropriate length of time (30 s for every 500 bp), followed by a final elongation step at 72°C for 5 min.

### Acknowledgements

We thank the Natural Environment Research Council of the UK for supporting this work through PhD studentships to IL and EK and research grants to HS and YC (NE/L006448/1, NE/H008918/1, NE/H016236/1). We are also grateful for a DAAD studentship to EK, a Warwick Chancellor's International Scholarship to YZ and a China Scholarship Council award to ZZ.

### References

- Alfieri, A., Malito, E., Orru, R., Fraaije, M.W., and Mattevi, A. (2008) Revealing the moonlighting role of NADP in the structure of a flavin-containing monooxygenase. *Proc Natl Acad Sci* **105**: 6572–6577.
- Alonso, C., and Pernthaler, J. (2006) *Roseobacter* and SAR11 dominate microbial glucose uptake in coastal North Sea waters. *Environ Microbiol* **8**: 2022–2030.
- Asher, E.C., Dacey, J.W.H., Mills, M.M., Arrigo, K.R., and Tortell, P.D. (2011) High concentrations and turnover rates of DMS, DMSP and DMSO in Antarctic sea ice. *Geophys Res Lett* **38**: 1–5.
- Aylward, F.O., Eppley, J.M., Smith, J.M., Chavez, F.P., Scholin, C.A., and DeLong, E.F. (2015) Microbial community transcriptional networks are conserved in three domains at ocean basin scales. *Proc Natl Acad Sci* **112**: 5443–5448.
- Bastet, L., Dubé, A., Massé, E., and Lafontaine, D.A. (2011) New insights into riboswitch regulation mechanisms. *Mol Microbiol* **80**: 1148–1154.
- Brimblecombe, P., and Shooter, D. (1986) Photo-oxidation of dimethylsulphide in aqueous solution. *Mar Chem* **19**: 343–353.
- Boden, R., Murrell, J.C., and Schäfer, H. (2011) Dimethylsulfide is an energy source for the heterotrophic marine bacterium *Sagittula stellata*. *FEMS Microbiol Lett* **322**: 188–193.
- Boden, R., Kelly, D.P., Murrell, J.C., and Schäfer, H. (2012) Oxidation of dimethylsulfide to tetrathionate by *Methylophaga thiooxidans* sp. nov.: a new link in the sulfur cycle. *Environ Microbiol* **12**: 2688–2699.
- Bucciarelli, E., Ridame, C., Sunda, W.G., Dimier-Hugueney, C., Cheize, M., and Belviso, S. (2013) Increased intracellular concentrations of DMSP and DMSO in iron-limited oceanic phytoplankton *Thalassiosira oceanica* and *Trichodesmium erythraeum*. *Limnol Oceanogr* **58**: 1667–1679.
- Buchan, A., González, J.M., and Moran, M.A. (2005) Overview of the marine *Roseobacter* lineage. *Appl Environ Microbiol* **71**: 5665–5677.
- Charlson, R.J., Lovelock, J.E., Andreae, M.O., and Warren, S.G. (1987) Oceanic phytoplankton, atmospheric sulphur, cloud albedo and climate. *Nature* **326**: 655–661.
- Chen, Y. (2012) Comparative genomics of methylated amine utilization by marine *Roseobacter* clade bacteria and development of functional gene markers (*tmm*, *gmaS*). *Environ Microbiol* **14**: 2308–2322.
- Chen, Y., Patel, N.A., Crombie, A., Scrivens, J.H., and Murrell, J.C. (2011) Bacterial flavin-containing monooxygenase is trimethylamine monooxygenase. *Proc Natl Acad Sci* **108**: 17791–17796.
- Cui, Y., Suzuki, S., Omori, Y., Wong, S.-K., Ijichi, M., Kaneko, R., *et al.* (2015) Abundance and distribution of dimethylsulfonylpropionate degradation genes and the corresponding bacterial community structure at dimethyl sulfide hot spots in the tropical and subtropical Pacific Ocean. *Appl Environ Microbiol* **81**: 4184–4194.
- Curson, A.R.J., Todd, J.D., Sullivan, M.J., and Johnston, A.W.B. (2011) Catabolism of dimethylsulfonylpropionate: microorganisms, enzymes and genes. *Nat Rev Micro* **9**: 849–859.
- del Valle, D.A., Kieber, D.J., and Kiene, R.P. (2007a) Depth-dependent fate of biologically-consumed dimethylsulfide in the Sargasso Sea. *Mar Chem* **103**: 197–208.
- del Valle, D.A., Kieber, D.J., Bisgrove, J., and Kiene, R.P. (2007b) Light-stimulated production of dissolved DMSO by a particle-associated process in the Ross Sea, Antarctica. *Limnol Oceanogr* **52**: 2456–2466.
- del Valle, D.A., Kieber, D.J., Toole, D.A., Bisgrove, J., and Kiene, R.P. (2009) Dissolved DMSO production via biological and photochemical oxidation of dissolved DMS in the Ross Sea, Antarctica. *Deep Sea Res Part 1 Oceanogr Res Pap* **56**: 166–177.
- De Zwart, J.M.M., Nelisse, P.N., and Kuenen, J.G. (1996) Isolation and characterization of *Methylophaga sulfidovorans* sp. nov.: an obligately methylotrophic, aerobic, demethylsulfide oxidizing bacterium from a microbial mat. *Microbiol Ecol* **20**: 261–270.
- Georg, J., and Hess, W.R. (2011) *cis*-Antisense RNA, another level of gene regulation in bacteria. *Microbiol Mol Biol Rev* **75**: 286–300.
- Gibb, S.W., and Hatton, A.D. (2004) The occurrence and distribution of trimethylamine-N-oxide in Antarctic coastal waters. *Mar Chem* **91**: 65–75.
- Gibb, S.W., Mantoura, R.F.C., Liss, P.S., and Barlow, R.G. (1999) Distributions and biogeochemistries of methylamines

- and ammonium in the Arabian Sea. *Deep Sea Res Part 2 Top Stud Oceanogr* **46**: 593–615.
- Giebel, H.-A., Kalhoefer, D., Lemke, A., Thole, S., Gahl-Janssen, R., Simon, M., et al. (2011) Distribution of *Roseobacter* RCA and SAR11 lineages in the North Sea and characteristics of an abundant RCA isolate. *ISME J* **5**: 8–19.
- Gifford, S.M., Sharma, S., Booth, M., and Moran, M.A. (2013) Expression patterns reveal niche diversification in a marine microbial assemblage. *ISME J* **7**: 281–298.
- Gilbert, B., McDonald, I.R., Finch, R., Stafford, G.P., Nielsen, A.K., and Murrell, J.C. (2000) Molecular analysis of the *pmo* (particulate methane monooxygenase) operons from two type II methanotrophs. *Appl Environ Microbiol* **66**: 966–975.
- Gilbert, J.A., Steele, J.A., Caporaso, J.G., Steinbruck, L., Reeders, J., Temperton, B., et al. (2012) Defining seasonal marine microbial community dynamics. *ISME J* **6**: 298–308.
- Giovannoni, S.J., Tripp, H.J., Givan, S., Podar, M., Vergin, K.L., Baptista, D., et al. (2005) Genome streamlining in a cosmopolitan oceanic bacterium. *Science* **309**: 1242–1245.
- González, J.M., Kiene, R.P., and Moran, M.A. (1999) Transformation of sulfur compounds by an abundant lineage of marine bacteria in the  $\alpha$ -subclass of the class *Proteobacteria*. *Appl Environ Microbiol* **65**: 3810–3819.
- González, J.M., Simó, R., Massana, R., Covert, J.S., Casamayor, E.O., Pedrós-Alió, C., et al. (2000) Bacterial community structure associated with a dimethylsulfoniopropionate-producing North Atlantic algal bloom. *Appl Environ Microbiol* **66**: 4237–4246.
- González, J.M., Covert, J.S., Whitman, W.B., Hendriksen, J.R., Mayer, F., Scharf, B., et al. (2003) *Silicibacter pomeroyi* sp. nov. and *Roseovarius nubinhibens* sp. nov., dimethylsulfoniopropionate-demethylating bacteria from marine environments. *Int J Syst Evol Microbiol* **53**: 1261–1269.
- Green, D.H., Shenoy, D.M., Hart, M.C., and Hatton, A.D. (2011) Coupling of dimethylsulfide oxidation to biomass production by a marine *Flavobacterium*. *Appl Environ Microbiol* **77**: 3137–3140.
- Hatton, A., and Wilson, S. (2007) Particulate dimethylsulphoxide and dimethylsulphoniopropionate in phytoplankton cultures and Scottish coastal waters. *Aquat Sci* **69**: 330–340.
- Hatton, A.D., Malin, G., and Liss, P.S. (1999) Distribution of biogenic sulphur compounds during and just after the southwest monsoon in the Arabian Sea. *Deep Sea Res Part 2 Top Stud Oceanogr* **46**: 617–632.
- Hatton, A.D., Turner, S.M., Malin, G., and Liss, P.S. (1998) Dimethylsulphoxide and other biogenic sulphur compounds in the Galapagos Plume. *Deep Sea Res Part 2 Top Stud Oceanogr* **45**: 1043–1053.
- Hatton, A., Shenoy, D., Hart, M., Mogg, A., and Green, D. (2012) Metabolism of DMSP, DMS and DMSO by the cultivable bacterial community associated with the DMSP-producing dinoflagellate *Scrippsiella trochoidea*. *Biogeochemistry* **110**: 131–146.
- Jonkers, H.M., Maarel van der M.J.E.C., van Gemerden, H., and Hansen, T.A. (1996) Demethylsulfoxide reduction by marine sulfate-reducing bacteria. *Microbiol Ecol* **136**: 283–287.
- Kiene, R.P., and Bates, T.S. (1990) Biological removal of dimethyl sulphide from sea water. *Nature* **345**: 702–705.
- Kiene, R.P., Linn, L.J., and Bruton, J.A. (2000) New and important roles for DMSP in marine microbial communities. *J Sea Res* **43**: 209–224.
- Krueger, S.K., and Williams, D.E. (2005) Mammalian flavin-containing monooxygenases: structure/function, genetic polymorphisms and role in drug metabolism. *Pharmacol Therapeut* **106**: 357–387.
- Lee, P.A., and De Mora, S.J. (1999) Intracellular dimethylsulfoxide (DMSO) in unicellular marine algae: speculations on its origin and possible biological role. *J Phycol* **35**: 8–18.
- Lidbury, I., Murrell, J.C., and Chen, Y. (2014) Trimethylamine N-oxide metabolism by abundant marine heterotrophic bacteria. *Proc Natl Acad Sci* **111**: 2710–2715.
- Lidbury, I., Kimberley, G., Scanlan, D.J., Murrell, J.C., and Chen, Y. (2015a) Comparative genomics and mutagenesis analyses of choline metabolism in the marine *Roseobacter* clade. *Environ Microbiol* **17**: 5048–5062.
- Lidbury, I., Murrell, J.C., and Chen, Y. (2015b) Trimethylamine and trimethylamine N-oxide are supplementary energy sources for a marine heterotrophic bacterium: implications for marine carbon and nitrogen cycling. *ISME J* **9**: 760–769.
- McDevitt, C.A., Hugenholtz, P., Hanson, G.R., and McEwan, A.G. (2002) Molecular analysis of dimethyl sulphide dehydrogenase from *Rhodovulum sulfidophilum*: its place in the dimethyl sulphoxide reductase family of microbial molybdopterin-containing enzymes. *Mol Microbiol* **44**: 1575–1587.
- Miller, T.R., and Belas, R. (2004). Dimethylsulfoniopropionate metabolism by *Pfiesteria*-associated *Roseobacter* spp. *Appl Environ Microbiol* **70**: 3383–3391.
- Morris, R.M., Rappe, M.S., Cannon, S.A., Vergin, K.L., Siebold, W.A., Carlson, C.A., et al. (2002) SAR11 clade dominates ocean surface bacterioplankton communities. *Nature* **420**: 806–810.
- Nelson, C.E., Carlson, C.A., Ewart, C.S., and Halewood, E.R. (2014) Community differentiation and population enrichment of Sargasso Sea bacterioplankton in the euphotic zone of a mesoscale mode-water eddy. *Environ Microbiol* **16**: 871–887.
- Ottesen, E.A., Young, C.R., Eppley, J.M., Ryan, J.P., Chavez, F.P., Scholin, C.A., et al. (2013) Pattern and synchrony of gene expression among sympatric marine microbial populations. *Proc Natl Acad Sci* **110**: E488–497.
- Pfreundt, U., Kopf, M., Belkin, N., Berman-Frank, I., and Hess, W.R. (2014) The primary transcriptome of the marine diazotroph *Trichodesmium erythraeum* IMS101. *Sci Rep* **4**. doi: 10.1038/srep06187.
- Riseman, S.F., and DiTullio, G.R. (2004) Particulate dimethylsulfoniopropionate and dimethylsulfoxide in relation to iron availability and algal community structure in the Peru upwelling system. *Can J Fish Aquat Sci* **61**: 721–735.
- Sakurai, I., Stazic, D., Eisenhut, M., Vuorio, E., Steglich, C., Hess, W.R., and Aro, E.-M. (2012) Positive regulation of *psbA* gene expression by *cis*-encoded antisense RNAs in *Synechocystis* sp. PCC 6803. *Plant Physiol* **160**: 1000–1010.
- Schäfer, H., McDonald, I.R., Nightingale, P.D., and Murrell, J.C. (2005) Evidence for the presence of a CmuA methyltransferase pathway in novel marine methyl halide oxidising bacteria. *Environ Microbiol* **7**: 839–852.
- Schäfer, H., Myronova, N., and Boden, R. (2010) Microbial degradation of dimethylsulphide and related C1-sulphur compounds: organisms and pathways controlling fluxes of sulphur in the biosphere. *J Exp Bot* **61**: 315–334.

- Sesto, N., Wurtzel, O., Archambaud, C., Sorek, R., and Cossart, P. (2013) The excludon: a new concept in bacterial antisense RNA-mediated gene regulation. *Nat Rev Micro* **11**: 75–82.
- Simó, R., Hattton, A., Gillian Malin, G., and Liss, P. (1998) Particulate dimethyl sulphoxide in seawater: production by microplankton. *Mar Ecol Prog Ser* **167**: 291–296.
- Simó, R., Pedrós-Alió, C., Malin, G., and Grimalt, J.O. (2000) Biological turnover of DMS, DMSP and DMSO in contrasting open-sea waters. *Mar Ecol Prog Ser* **203**: 1–11.
- Sjöstedt, J., Martiny, J.B.H., Munk, P., and Riemann, L. (2014) Abundance of broad bacterial taxa in the Sargasso Sea explained by environmental conditions but not water mass. *Appl Environ Microb* **80**: 2786–2795.
- Sowell, S.M., Wilhelm, L.J., Norbeck, A.D., Lipton, M.S., Nicora, C.D., Barofsky, D.F., *et al.* (2008). Transport functions dominate the SAR11 metaproteome at low-nutrient extremes in the Sargasso Sea. *ISME J* **3**: 93–105.
- Stazic, D., Lindell, D., and Steglich, C. (2011) Antisense RNA protects mRNA from RNase E degradation by RNA–RNA duplex formation during phage infection. *Nucleic Acids Res* **39**: 4890–4899.
- Sun, J., Steindler, L., Thrash, J.C., Halsey, K.H., and Smith, D.P., Carter, A.E., *et al.* (2011) One carbon metabolism in SAR11 pelagic marine bacteria. *PLoS One* **6**: e23973.
- Sunda, W., Kieber, D.J., Kiene, R.P., and Huntsman, S. (2002) An antioxidant function for DMSP and DMS in marine algae. *Nature* **418**: 317–320.
- Todd, J.D., Kirkwood, M., Newton-Payne, S., and Johnston, A.W.B. (2012) DddW, a third DMSP lyase in a model *Roseobacter* marine bacterium, *Ruegeria pomeroyi* DSS-3. *ISME J* **6**: 223–226.
- Tynes, R.E., Sabourin, P.J., Hodgson, E., and Philpot, R.M. (1986) Formation of hydrogen peroxide and N-hydroxylated amines catalyzed by pulmonary flavin-containing monooxygenases in the presence of primary alkylamines. *Arch Biochem Biophys* **251**: 654–664.
- Vila, M., Simó, R., Kiene, R.P., Pinhassi, J., González, J.M., Moran, M.A., *et al.* (2004) Use of microautoradiography combined with fluorescence in situ hybridization to determine dimethylsulfoniopropionate incorporation by marine bacterioplankton taxa. *Appl Environ Microbiol* **70**: 4648–4657.
- Vila-Costa, M., Kiene, R.P., and Simó, R. (2008) Seasonal variability of the dynamics of dimethylated sulfur compounds in a coastal northwest Mediterranean site. *Limnol Oceanogr* **53**: 198–211.
- Vila-Costa, M., del Valle, D.A., González, J.M., Slezak, D., Kiene, R.P., Sanchez, O., *et al.* (2006). Phylogenetic identification and metabolism of marine dimethylsulfide-consuming bacteria. *Environ Microbiol* **8**: 2189–2200.
- Watts, S.F. (2000) The mass budgets of carbonyl sulfide, dimethyl sulfide, carbon disulfide and hydrogen sulfide. *Atmos Environ* **34**: 761–779.
- Williams, D.E., Hale, S.E., Muerhoff, A.S., and Masters, B.S. (1985) Rabbit lung flavin-containing monooxygenase. Purification, characterization, and induction during pregnancy. *Mol Pharmacol* **28**: 381–390.
- Williams, T.J., Long, E., Evans, F., DeMaere, M.Z., Lauro, F.M., Raftery, M.J., *et al.* (2012) A metaproteomic assessment of winter and summer bacterioplankton from Antarctic Peninsula coastal surface waters. *ISME J* **6**: 1883–1900.
- Zhu, Y., Jameson, E., Parslow, R.A., Lidbury, I., Fu, T., Dafforn, T.R. *et al.* (2014) Identification and characterisation of trimethylamine N-oxide (TMAO) demethylase and TMAO permease in *Methylocella silverstris* BL2. *Environ Microbiol* **16**: 3318–3330.
- Zubkov, M.V., Fuchs, B.M., Archer, S.D., Kiene, R.P., Amann, R., and Burkhill, P.H. (2001) Linking the composition of bacterioplankton to rapid turnover of dissolved dimethylsulphoniopropionate in an algal bloom in the North Sea. *Environ Microbiol* **3**: 304–311.
- Zubkov, M., Linn, L.J., Amann, R., and Kiene, R.P. (2004) Temporal patterns of biological dimethylsulfide (DMS) consumption during laboratory-induced phytoplankton bloom cycles. *Mar Ecol Prog Ser* **271**: 77–86.

## Supporting information

Additional supporting information may be found in the online version of this article at the publisher's web-site:

### Quantification of $\beta$ -galactosidase activity

**Table S1.** A list of *R. pomeroyi* DSS-3 strains and plasmids used in this study

**Table S2.** Primers used in this study. Bases underlined represent the restriction sites engineered in for enzymatic digestion.

**Fig. S1.** RT-PCR assays targeting the antisense strand of *tmm* of *R. pomeroyi* cells (duplicate cultures, 1 and 2) grown on either ammonium ( $\text{NH}_4^+$ ) or trimethylamine (TMA) as the sole nitrogen source respectively. The presence of an antisense RNA was observed in TMA-grown cultures. Controls were performed for RT-PCR where no reverse transcriptase was added (-) in the reverse transcription reactions. Positive and negative PCR controls were also set up where genomic DNA of *R. pomeroyi* or ddH<sub>2</sub>O was used as the template, respectively.