

Extinction debt on reservoir land-bridge islands

Isabel L. Jones^{a*}, Nils Bunnefeld^a, Alistair S. Jump^a, Carlos A. Peres^b, Daisy H. Dent^{a,c}

^aBiological and Environmental Sciences, University of Stirling, Stirling, FK9 4LA, UK.

*Corresponding author. Phone +44 (0)7738 009612, email <i.l.jones@stir.ac.uk>

5 ^bSchool of Environmental Sciences, University of East Anglia, Norwich, NR4 7TJ, UK.

^cSmithsonian Tropical Research Institute, Apartado 0843–03092, Balboa, Panama.

KEYWORDS:

Conservation policy; dams; hydropower; fragmentation; isolation; species richness

10

ABSTRACT

Large dams cause extensive inundation of terrestrial habitats, whereby hilltops become land-bridge islands: habitat is fragmented and isolated, inducing local extinctions and degradation of remnant biological communities. “Good practice” dam development guidelines propose using reservoir islands for species conservation, mitigating some of the detrimental impacts associated with flooding terrestrial habitats. The degree of species retention on islands, and hence, whether they are effective for long-term conservation is currently unknown. Here, we quantitatively review species’ responses to isolation on reservoir islands. We specifically investigate island species richness in relation to neighbouring continuous habitat, and relationships between species richness and island area, isolation time, distance to mainland and other islands. Species’ responses to isolation on reservoir islands have been investigated in only 15 of the >58 000 large-dam reservoirs (dam height >15 m) operating globally. Research predominantly originates from wet tropical forest habitats and focuses on mammals, with species richness being the most widely-reported ecological metric. Terrestrial taxa are, overall, negatively impacted by isolation on reservoir islands. Reservoir island species richness declines with isolation time, and though it increases with area, all islands exhibit depauperate species richness <100 years after isolation compared to continuous mainland habitats. Such a pattern of

15
20
25

sustained and delayed species loss following large-scale habitat disturbance is indicative of an extinction debt existing for reservoir island species, and is evident across all taxonomic groups and dams studied. Thus, reservoir islands cannot reliably be used for species conservation as part of impact mitigation measures, and should instead be included in area calculations for land impacted by dam creation. Environmental licensing assessments as a precondition for future dam development should explicitly consider the long-term fate of island communities when assessing biodiversity loss vs energy output.

35

1. INTRODUCTION

There are 58 402 large dams (dam height >15 m) operating globally, constructed predominantly for irrigation and hydropower generation (ICOLD, 2016). A growing human population is predicted to increase the demand for water by 2-3% per year, and the demand for energy by >56% globally, and by 90% in increasingly industrialised countries with emerging economies between 2010-2040 (EIA, 2013; WCD, 2000). Concurrently, changing climatic and precipitation patterns, including severe droughts, will likely further increase demand for water and reduce hydropower generation from large reservoirs (Oki and Kanae, 2006).

45

Hydropower is regarded as a renewable “green” energy source, and dams constructed in areas with steep topography and high rainfall produce the most energy per unit area (Finer and Jenkins, 2012). However dams are often constructed in low-lying high conservation value areas: for example 154 dams operate in the Amazon basin with a further 277 planned (Castello et al., 2013; Lees et al., 2016).

50 The construction of dams directly impacts both terrestrial and freshwater ecosystems through inundation of habitat, compositional changes in aquatic communities, and the loss of structural and functional connectivity between upper and lower reaches of watersheds (Finer et al., 2008; Nilsson et al., 2005; Lees et al., 2016; Palmeirim et al., 2014; Sá-Oliveira et al., 2015).

55 Over 50% of the world's large river systems and >60% of the combined habitat area of tropical, subtropical and boreal forests, tropical and subtropical grasslands, savannahs and shrublands have been estimated to be impacted by dams (Nilsson et al., 2005). Inundation of terrestrial habitats, and tropical forests in particular, can also result in significant carbon emissions from reservoirs in the form of CO₂ and CH₄, which can persist for many years after inundation and often over the lifetime of
60 the reservoir (Abril et al., 2005; Demarty & Bastien, 2011; Fearnside, 2002; Fearnside & Pueyo, 2012). Direct social impacts arise from the loss of indigenous lands, displacement of communities, and disruption to local economies reliant on fisheries often concurrently affected by heavy metal accumulation (Boudou et al., 2005; Fearnside, 1999). Additionally, increased access to previously undisturbed habitat can elevate levels of hunting and deforestation in areas surrounding reservoirs
65 (Kirby et al., 2006; Peres and Lake, 2003).

When dams are built, habitat is lost through inundation, with remnants of previously continuous terrestrial habitat confined to highly fragmented land-bridge island archipelagos comprised of former hilltops. "Good practice" guidelines (International Energy Agency, 2006) for dam developers to
70 mitigate ecological impacts from dam construction include implementing protected areas covering land-bridge islands and habitat surrounding reservoirs. For example, the REBIO Uatumã (the largest Biological Reserve in Brazil) encompasses approximately half of the Balbina hydroelectric reservoir, including all islands east of the former left bank of the Uatumã river and mainland areas extending away from the eastern edge of the reservoir. Strictly-protected area status has largely deterred small-
75 scale slash-and-burn agriculture and extraction of resources within the REBIO Uatumã, on both islands and within surrounding continuous forest (Benchimol & Peres, 2015a, 2015b). However, we do not know whether protecting reservoir islands is effective for biodiversity conservation, due to a lack of long-term monitoring. The International Energy Agency highlights the dearth of systematic evaluation of any mitigation, enhancement, and compensation measures currently being
80 recommended to developers (International Energy Agency, 2000; Trussart et al., 2002).

Fragmentation of habitat causes a number of impacts to species, such as population reductions and local extinctions; the strength of fragmentation impacts differ depending on the taxonomic group and life-history traits of species (Bender et al., 1998; Fahrig 2003; Forman, 1995). Previous studies of
85 reservoir island archipelagos have shown that island taxa typically experience a novel hyper-disturbance regime, resulting in drastic shifts in species diversity and community composition through species turnover and altered carrying capacity of remaining habitat (Benchimol & Peres, 2015a; Cosson et al., 1999a; Ferreira et al., 2012; Hanski & Ovaskainen, 2000; Terborgh et al., 2001). Local species extinctions on reservoir islands have been observed for plants (Benchimol & Peres, 2015a; Yu
90 et al., 2012), invertebrates (Emer et al., 2013; Feer & Hingrat, 2005), birds (Yu et al., 2012), bats (Cosson et al., 1999b), small-mammals (Gibson et al., 2013; Lambert et al., 2003), and midsized to large-bodied vertebrates (Benchimol & Peres, 2015b, 2015c). In contrast, populations of some species can become hyper-abundant on islands, further impacting other taxa (Chauvet & Forget, 2005; Feeley & Terborgh, 2006; Lopez & Terborgh, 2007), as can the establishment of invasive species (Gibson et
95 al., 2013).

Changes in island communities may not occur immediately after inundation; instead, species may be subject to an “extinction debt” whereby a portion of species are initially lost, followed, potentially multiple generations later, by further species extinctions (Halley et al., 2014; Kitzes and Harte, 2015;
100 Kuussaari et al., 2009; Tilman et. al., 1994). Thus, the effects of fragmentation and isolation can persist for years after initial habitat loss, as communities undergo “relaxation” towards a new equilibrium community (Diamond, 1972; Diamond, 2001; Ewers and Didham, 2006; Feeley et al., 2007; Terborgh et al., 1997; Wang et al., 2009). The “relaxation” process is likely mediated by island area, with species losses faster on smaller islands, and a greater time-lag for species loss on larger
105 islands (Diamond, 1972; Gonzalez, 2000). There are a number of empirical methods for calculating extinction debt (Kitzes and Harte, 2015; Wearn et al., 2012), and here we consider a decline in species richness on islands over time, compared to continuous mainland habitat, as evidence of extinction debt. In the absence of extinction debt, we assume that all species extinctions would happen

immediately, with no evidence of further degradation of insular biological communities through time
110 (Kitzes and Harte, 2015).

In the long-term it is unknown how island communities will continue to respond with increasing
island isolation time, as the creation of artificial archipelagos from dam construction has only
occurred over the past century. Our present knowledge of ecological communities within artificial
115 archipelagos comes from multiple snapshot studies from different countries, dams, habitats and taxa,
at different time points since the originally continuous habitat was fragmented. Bringing these
snapshots together enables identification of general trends across disparate studies, aiding develop of
policy-relevant recommendations in terms of the conservation value of reservoir islands.

120 Here, we quantitatively review peer-reviewed research detailing responses of terrestrial taxa to habitat
fragmentation, and subsequent isolation, on reservoir land-bridge islands. We then analyse species
richness data from 249 islands and adjacent continuous habitats through time. In particular, we ask:
(1) is there evidence of an extinction debt existing for reservoir island species; i.e. compared to
continuous habitat, does island species richness decrease with increasing island isolation time? and (2)
125 how does island size, distance to continuous habitat and distance to other islands relate to patterns of
species richness and rates of species loss?

2. METHODS

130 2.1. Literature summary

2.1.1 Dataset collation

We conducted a literature search using Web of Knowledge and Google Scholar search engines
between January 2014 and June 2015 using the key words: hydropower or hydroelectric, reservoir or
135 dam, and island or land-bridge, forest islands or fragments. Only full-text, peer-reviewed articles in
English were retained; unpublished or grey literature was not included. Studies researching terrestrial

species, guilds, taxonomic groups or communities on reservoir islands, attributing ecological responses observed to reservoir creation were retained. Experimental studies or those not explicitly stating an aspect of reservoir creation as a causal factor for the response observed were excluded.

140 Studies which met the inclusion criteria were entered into a dataset (henceforth referred to as “dataset studies”). Literature cited in dataset studies was also screened for inclusion, and searches for names of dams in dataset studies performed. A total of 129 studies were assessed for inclusion in our study, 100 of which met the criteria to be retained.

145 2.1.2 Data extraction

Data such as the number of islands surveyed and island area, taxonomic groups investigated, and time since island isolation were extracted from studies (see Database A). Each study was assigned a broad habitat type (wet tropical forest, tropical grassland e.g. *cerrado*, subtropical forest, Mediterranean forest, boreal forest). Taxa investigated were broadly grouped into mammals, birds, invertebrates, 150 herptiles, plants, and fungi. If multiple taxa were included within a study, data were extracted for each group separately due to the potential for different responses. The precise isolation time of islands is seldom reported, thus we estimated island isolation time as the year of dam closure minus the year of field data collection. In six studies field data collection dates were not reported, thus, data collection date was conservatively estimated as two years prior to publication date.

155

2.1.3. Assigning study response directions

For each study the authors’ key results and conclusions were used to assign an overall response of the study taxa to isolation on islands (response: positive, negative, variable, or neutral; see column ‘L’, Database A). For example, a study reporting declining species richness on islands would be assigned 160 an overall negative response. An overall positive response would be assigned if, for example, recorded sightings (e.g. presence/absence data) were higher on islands. Overall variable responses could result from research involving different species within the same taxon, e.g. two species of bat exhibiting divergent responses to isolation. Neutral responses would result if no differences or alterations in taxa on islands compared to mainland sites were reported. If authors did not draw a

165 conclusion as to the response directions observed, we examined the data reported and assigned a
response direction accordingly. If multiple response directions for the same taxa were observed over
time, the predominant response direction (i.e. over most years) was used as the overall direction.

To account for within-study complexity i.e. inclusion of multiple taxonomic groups and/or ecological
170 metrics, response directions were derived for each taxonomic group and ecological metric
investigated (see columns 'M-P', Database A). Ecological metrics included species richness,
population density, behaviour (e.g. foraging behaviour), community composition, presence/absence,
fitness/recruitment (e.g. breeding output), genetic diversity, and functional diversity.

175 **2.2 Species richness analysis**

Estimates of species richness were the most widely-reported and accessible data available in the
collated studies, and therefore we selected this ecological metric for in-depth analysis.

180 *2.2.1. Data collection*

Dataset studies presenting species richness data for islands and nearby continuous (control) habitat, as
well as island areas and isolation time, were used to assess variation in species richness on reservoir
islands compared to control habitat (Table B1). These data also allowed investigation of the
relationship between species richness and island area, isolation time, distance to mainland and
185 distance to nearest island. Of the 100 dataset studies, 17 presented species richness data for islands (n
= 249; size range <1-1690 ha; isolation time <1-92 years) and control sites (n = 84), and were used for
the in-depth analysis of species richness data (Table B1; Database B). If data for the distance to
mainland or nearest island were not presented, then if possible these data were calculated from
satellite imagery using Google Earth Pro (Google, 2015). Geographically, the 17 studies suitable for
190 species richness analysis originated from nine dams, located on three continents in three broad habitat
types (wet tropical forest, subtropical forest, and tropical grassland).

2.2.2. Data analysis

For each study the ratio of island species richness to average control species richness (S_{RICH}) was
195 calculated (see Database B). If a study contained data over multiple years, and thus, multiple isolation
times, then species richness for control sites over the same isolation time period was averaged. If a
study had multiple species richness values for the same island size, taxon, and isolation time, species
richness values were averaged to avoid pseudo-replication.

200 To normalise data, all data were logged (natural logarithm) prior to analysis. S_{RICH} values were
modelled using linear mixed effects models (lmer using lme4; Bates et al., 2014), as a function of
island isolation time (T_{ISO}), island area (AREA), distance to mainland (D_{MAIN}) and distance to nearest
island (D_{ISLAND}) as fixed effects, with taxonomic group (TAXA), dam identity (DAM; a surrogate for
location), and study (STUDY; to account for differing survey methods and survey intensity between
205 studies) as random effects (Bunnfeld and Phillimore, 2012; see Database B). Interaction terms were
included between AREA, T_{ISO} , D_{MAIN} and D_{ISLAND} , as well as between TAXA, DAM and STUDY;
quadratic terms were also tested for.

Due to missing values for D_{MAIN} and D_{ISLAND} we reduced the dataset to only those data rows
210 containing values for all variables being tested (n islands = 178) and used this dataset for linear
regression and model selection in R (R Core Team, 2015). Models were simplified following stepwise
deletion of non-significant terms i.e. those with a t-value < 2 and models compared using Chi-square
tests in ANOVA (Crawley, 2005; Table B2). Following model simplification, the final model did not
include variables with missing values, thus, the final model was fitted to the whole dataset (n islands
215 = 249). The best linear unbiased predictors (BLUPs) for each dam were extracted using the ‘ranef’
function within the lme4 R package (Pinheiro and Bates, 2000). Each dam has a different intercept,
which can fall above or below that of the overall model: positive BLUPs indicate that the dam has
higher than expected levels of species richness estimated from the fixed effects, and those falling
below the model average indicate that species richness is lower than expected. A variance components
220 analysis was carried out for the random effects (Crawley, 2005).

3. RESULTS

3.1. Literature summary

225

The 100 dataset studies examined here were predominantly from Neotropical wet tropical forest habitats (Fig. 1; 2a). Mammals were the best-studied taxonomic group (Fig. 2b); responses of terrestrial taxa isolated on reservoir islands were most often expressed in terms of species richness and presence/absence, and rarely with regards to behaviour, genetic or functional diversity (Fig. 2c). An overall negative response of terrestrial taxa to dam creation was reported in >75% of studies, and these negative responses were seen across all habitat types, ecological metrics, and taxonomic groups investigated (Fig. 2a-c). Overall positive responses were confined to only two of the 100 studies (Fig. 2a), of which one reported increased and more stable population densities of small mammals (Adler, 1996), and the second, increased food resources for a raptor due to prey being ‘captive’ on isolated islands (Benchimol and Venticinque, 2010). Studies report results for islands isolated from <1 to 92 years, with the mean island isolation age of ~33 years (Fig. 2d).

230

235

3.2. Species richness analysis

240

The final model for analysis of S_{RICH} included T_{ISO} and AREA as fixed effects, and TAXA, DAM and STUDY as random effects (Table 1); D_{MAIN} and D_{ISLAND} had no significant effect on island S_{RICH} , and no interaction terms were significant (Table B2). Of the fixed effects, 36% of variation was explained by STUDY, 17% explained by DAM, with 47% residual variance; TAXA did not explain any variance.

245

For all taxonomic groups and dams, species richness declined with island isolation time, with larger islands retaining more species than smaller islands (Fig. 3): predicted S_{RICH} on the largest island (1690 ha, Balbina hydroelectric dam, Brazilian Amazon) is predicted to be 3.2 at the mean isolation time of islands in the analysis compared to a predicted S_{RICH} of 1.2 on the smallest island (0.17 ha, Cabra

Corral, Argentina). Even the largest island (1690 ha) exhibits reduced species richness compared to
250 mainland continuous habitat in less than 30 years of isolation; Barro Colorado Island (~1500 ha,
Gatun Lake, Panama) isolated for the longest period in our study (~92 years) similarly shows
sustained species richness declines.

The estimates for the random effect of DAM (BLUPs) showed for the majority of dams (66%) lower
255 species richness levels than the overall intercept estimated by the fixed effects; only islands in Gatun
Lake, Balbina, and Thousand Island Lake maintain higher species richness than the modelled fixed
effects intercept (Fig. 3; Table B3). Using our model we can predict S_{RICH} values for islands of mean
area at a given isolation time, and islands of different areas at the mean isolation time for each
reservoir. For example the S_{RICH} for mean island size within Gatun Lake reduces from 2.24 at five
260 years of isolation to 1.49 after 90 years of isolation; in Lake Kenyir, which maintains the lowest
expected species richness values, a small island of 5ha (at mean island isolation time) has a predicted
 S_{RICH} value of 1.35, which is increased to just 2.23 on an island of 1000 ha. There was no evidence that
islands located nearer other terrestrial habitat or mainland continuous habitat had reduced levels of
species loss.

265

4. DISCUSSION

Our study finds that terrestrial taxa isolated on reservoir islands experience significant reductions in
species richness in less than a century of isolation. Such sustained local species losses since the initial
270 loss of habitat indicates that reservoir island species are subject to an extinction debt, which is evident
across all dams, habitats, and taxa. All islands showed depauperate levels of species richness
compared to continuous habitats, with smaller islands maintaining lower species richness than larger
islands. Island isolation time and area, but not distance from other terrestrial habitat or the mainland,
were the drivers of species richness patterns observed.

275

More broadly, we show that the majority of taxa are negatively impacted by reservoir creation across a range of other ecological metrics including behaviour and genetic diversity. Our current knowledge of the impacts of reservoir creation is disproportionately focussed on mammals, and originates predominantly from evergreen Neotropical forest habitats. While not all dams create archipelagic
280 landscapes, research within our synthesis covers just 15 of the >58 000 large dams operating globally, representing a small and potentially biased sample of possible island systems. However, even with such limited data we clearly demonstrate the negative impact of dam creation on island species richness. Furthermore, we highlight the shortfalls in current conservation and impact mitigation strategies for dam development, particularly in terms of long-term biological costs, in addition to the
285 immediate direct loss of lowland habitat during flooding.

4.1. Island species richness, area, and isolation time

Classic island biogeography theory (IBT, MacArthur & Wilson, 1967) explains variation in island
290 species richness through a balance of species immigration and distance from species source pools. In the artificial archipelagic systems we investigate in our analysis, rather than a process of species accumulation on islands, remnant communities of formerly continuous habitat undergo species loss (“relaxation”) until a new equilibrium community is reached (Diamond, 1972; Gonzalez, 2000; Lomolino, 2000).

295 Area was a significant predictor of species richness on islands within our analysis as expected from the species-area relationship and IBT (Connor & McCoy, 1979; MacArthur & Wilson, 1967; Triantis et al., 2012). However distance, both to the mainland and other islands, was not a significant predictor of island species richness: this represents a departure from the IBT, and reduced importance of
300 metapopulation dynamics (Hanski and Gilpin, 1991; With and King, 2001) and the “rescue effect” (Brown and Kodric-Brown, 1977) for maintaining insular populations in artificial archipelagic systems.

In the case of reservoir islands, remnant terrestrial habitat fragments are surrounded by a high-contrast, inhospitable matrix, presenting a prohibitive dispersal barrier for certain taxa. Such an extreme dispersal barrier effectively renders all islands as too isolated for any “rescue effect” from wider species source pools to maintain island communities and species richness, and explains the lack of distance effects we find in our analysis (Watson, 2002). The evolutionary history and traits of species resident in continuous habitats make many incapable of dispersing through open habitats, across large distances, or through a high-contrast matrix such as land and water (see Fig. 2 in Ewers & Didham, 2006). The ability of tropical understorey bird species to disperse across a water matrix between islands was tested in Gatun Lake, Panama, where some species were limited to <100m of flight (Moore et al., 2008). Species reliant on continuous habitats can be averse to crossing even small clearings, such as logging roads, even when the forest canopy is closed (Develey & Stouffer, 2001; Laurance et al., 2004).

Habitat fragments surrounded by water therefore represent a worse-case scenario in terms of fragmentation effects: aside from extreme dispersal barriers preventing species migration, islands are subject to extreme edge effects from increased UV and wind damage, often penetrating deep into islands leading to further degradation of island biota (Benchimol & Peres, 2015b; Laurance, 2008; Murcia, 1995). Habitat fragments embedded within a more similar and potentially hospitable but low-quality terrestrial habitat matrix (e.g. forest fragments within an agricultural landscape) can retain higher levels of species diversity with reduced local extinction rates (Mendenhall et al., 2011), when compared to reservoir islands of a similar size (Mendenhall et al., 2014).

We find a reduction in species richness on all islands with increasing time since initial habitat loss. Such a pattern of sustained and delayed species loss is indicative of extinction debt (Tilman et al., 1994; Kitzes & Harte, 2015; Kuussaari et al., 2009). Extinction debts are especially high in areas subject to recent large-scale habitat loss, such as islands created by rapid flooding of terrestrial habitats (Hanski and Ovaskainen, 2002). Our analysis illustrates that reservoir islands are of limited long-term conservation value due to evidence of an extinction debt: species loss appears most rapid on

smaller islands and even the largest islands studied (~1690 ha) exhibited lowered species richness in under 30 years of isolation. Ongoing species losses have been reported on another large island in our synthesis: Barro Colorado Island (BCI, ~1500 ha) has been isolated for 92 years since the formation
335 of the Gatun Lake, Panama. In less than a century of island isolation, and despite strict environmental protection of BCI and surrounding peninsulas, 65 bird species (Robinson, 1999) and 23 butterfly species (Basset et al., 2015) have become locally extinct, alongside long-term degradation of the tree community (Leigh et al., 1993). In the Balbina hydroelectric megadam system in Amazonia, to conserve >80% of terrestrial and arboreal vertebrates on islands, a threshold island size of 475 ha was
340 identified by Benchimol & Peres (2015b). However, only 25 out of 3546 islands in the Balbina archipelago meet this size criterion. Balbina is protected by the largest biological reserve in Brazil, and thus represents a best case scenario for biodiversity conservation within an artificial archipelago system. Species inhabiting other such systems, without protection, will therefore likely suffer not only from direct habitat loss through flooding and potential extinction debt, but additional human-mediated
345 impacts such as deforestation, agriculture, hunting, and fire (Laurance, 2008; Peres, 2001).

The data we use for analysis of species richness on reservoir land-bridge islands originate from 249 islands within 9 of the 15 dams presented in Fig. 1 and allow us to show patterns applicable to all dams and taxonomic groups, although we acknowledge that publication bias towards negative impacts
350 of reservoir creation could influence the response patterns presented. While the data do not allow us to disentangle species richness patterns for individual taxonomic groups, dams and habitat types, we addressed this shortcoming by using random effects in linear mixed effects models (Bunnefeld and Phillimore, 2012). Similarly we cannot calculate the magnitude of extinction debts for individual taxonomic groups and/or habitat types, and instead highlight evidence that all reservoir islands are
355 subject to an extinction debt, and therefore cannot be relied upon for long-term species conservation.

The observed patterns of depauperate island species richness could be shaped by landscape attributes prior to inundation and non-random loss of more species-rich lowland habitat during flooding (Seabloom et al., 2002). Mainland species richness levels may have been elevated through surveying

360 lowland habitats; such a potential sampling effect should be accounted for during survey site selection
(e.g. Benchimol and Peres, 2015a). In continuous habitats the greater availability of resources allows
more species to inhabit a given area, compared to the same area of isolated habitat (Ewers and
Didham, 2006). Thus, sampling islands can inherently give lower species richness values than an
equal area of continuous habitat (Crawley & Hurrall, 2001; Gonzalez, 2000; Halley et al., 2014;
365 MacArthur & Wilson, 1963).

Data for island taxa in artificial archipelagos come from snapshots of responses to isolation in <100
years of reservoir lifetime, across multiple taxa and habitat types. In addition, no studies monitored
changes in insular community dynamics over a significant post-isolation time. Consequently, we
370 cannot currently determine if the rates of local species loss are predictable beyond the relatively short
time frame analysed here. Nor can we accurately quantify extinction debt to predict the eventual
number of species able to persist in the artificial archipelago systems created due to the assumptions
that would be required to do so. Further long-term monitoring of reservoir island biota is needed to
allow these more detailed assessments to be made, since at present only Gatun Lake, Panama,
375 provides data for a reservoir >90 years of age.

4.2. Conservation implications

Our study strongly suggests that islands within reservoir systems do not sustain full complements of
380 flora and fauna in the long term; larger islands retain species for longer than smaller islands, but all
island communities likely face an extinction debt. Given that degradation of island communities can
be predicted to occur in all artificial archipelagic systems created by dam development, we emphasise
that reservoir islands cannot be used for species conservation as part of impact mitigation strategies.
The combined area of reservoir islands should be explicitly included in environmental impact
385 assessments, in addition to the area of habitat directly lost through inundation.

Current policy to mitigate the negative impacts of dam creation on terrestrial environments consists of “good practice” guidelines with no statutory legislation requiring specific actions by developers (International Energy Agency, 2006). Environmental legislation is highly variable among countries, and there is no signatory international agreement on how to forecast, prevent or mitigate the effects of large dams. Mitigation measures can take a multitude of forms, ranging from conducting wildlife inventories and environmental impact assessments before reservoir filling, creating new habitats such as wetland zones within the reservoir system, and conservation offsets such as strictly protecting land both within and surrounding reservoirs. There is however no long-term monitoring of such practices to assess whether these mitigation measures are effective (International Energy Agency, 2000).

A great many dams planned to meet future water and electricity needs, especially in developing countries. We call for better trade-off calculations (Kareiva, 2012) to be made for future dams, accounting for long-term species loss on islands created by flooding. In addition, enhanced protection of larger islands and surrounding non-fragmented habitats is essential to avoid biological collapse in artificial archipelagic systems. We highlight the potential for additional impacts from long-term degradation of high carbon-storing habitats such as tropical forests: erosion of island tree communities (Benchimol and Peres, 2015a) could lead to a future carbon loss from tropical dams, exacerbating the greenhouse gas emissions already documented from this “green” energy source (Demarty and Bastien, 2011; Fearnside, 2009).

4.3. Conclusions

We have shown that there is an overall negative response of terrestrial species and communities to isolation on reservoir land-bridge islands. These trends are seen across a broad spectrum of taxonomic groups and ecological metrics. Species isolated on reservoir islands will likely experience extinction debt, and the rate of local extinctions is driven by island size and island isolation time, independently of distance from potential source populations within the landscape. Our synthesis of current literature allows broad conclusions about the ecological impacts of reservoirs through time, and highlights the

415 need for further research from a greater number of reservoirs over the duration of their lifetime.
Building upon the findings that we present here, investigation of the many other direct and indirect
ecological impacts of reservoirs such as loss of river habitats and connectivity, land tenure rights, and
the impacts of wider infrastructure development on surrounding habitats, should be a priority for
future research.

420

Acknowledgements

We thank the Tropical Ecology and Conservation (T.E.A.C.) research group at the University of
Stirling, UK, for valuable discussion when designing this study, and when interpreting the findings.

425 We also thank Faulk Huettmann and an anonymous reviewer, for providing valuable comments that
greatly improved the manuscript.

Role of the funding source

430 This study was funded by the University of Stirling (UoS). UoS was not involved with the study
design, data collection, analysis or interpretation of data, writing or decision to submit this
manuscript.

REFERENCES

- 435 Abril, G., Guérin, F., Richard, S., Delmas, R., Galy-Lacaux, C., Gosse, P., Tremblay, A., Varfalvy, L.,
Dos Santos, M.A., Matvienko, B., 2005. Carbon dioxide and methane emissions and the carbon
budget of a 10-year old tropical reservoir (Petit Saut, French Guiana). *Global Biogeochem.*
Cycles 19, GB4007. doi:10.1029/2005GB002457
- Adler, G.H., 1996. The Island Syndrome in Isolated Populations of a Tropical Forest Rodent.
Oecologia 108, 694–700
- 440 Basset, Y., Barrios, H., Segar, S., Srygley, R.B., Aiello, A., Warren, A.D., Delgado, F., Coronado, J.,
Lezcano, J., Arizala, S., Rivera, M., Perez, F., Bobadilla, R., Lopez, Y., Ramirez, J.A., 2015.
The Butterflies of Barro Colorado Island, Panama: Local Extinction since the 1930s. *PLoS One*
10, e0136623. doi:10.1371/journal.pone.0136623
- 445 Bates, D., Maechler, M., Bolker, B., Walker, S., 2014. lme4: Linear mixed-effects models using Eigen
and S4. URL: <http://cran.r-project.org/package=lme4> [Accessed 21/12/15]

- Benchimol, M., Peres, C. A., 2015a. Edge-mediated compositional and functional decay of tree assemblages in Amazonian forest islands after 26 years of isolation. *J. Ecol.* 103, 408–420. doi:10.1111/1365-2745.12371
- 450 Benchimol, M., Peres, C. A., 2015b. Predicting local extinctions of Amazonian vertebrates in forest islands created by a mega dam. *Biol. Conserv.* 187, 61–72. doi:10.1016/j.biocon.2015.04.005
- Benchimol, M., Peres, C.A., 2015c. Widespread Forest Vertebrate Extinctions Induced by a Mega Hydroelectric Dam in Lowland Amazonia. *PLoS ONE* 10(7): e0129818. doi: 10.1371/journal.pone.0129818
- 455 Benchimol, M., Venticinque, E.M., 2010. Harpy Eagle (*Harpia harpyja*) predation on an Infant brown Capuchin Monkey (*Cebus apella*) in the Brazilian Amazon. *Rev. Bras. Ornitol.* 18, 352–354
- Brown, J.H., Kodric-Brown, A., 1977. Turnover Rates in Insular Biogeography: Effect of Immigration on Extinction. *Ecology* 58, 445–449
- 460 Bunnefeld, N., Phillimore, A.B., 2012. Island, archipelago and taxon effects: mixed models as a means of dealing with the imperfect design of nature’s experiments. *Ecography* 35, 15–22. doi: 10.1111/j.1600-0587.2011.07078.x
- Canale, G.R., Peres, C.A., Guidorizzi, C.E., Ferreira Gatto, C.A., Kierulff, M.C.M., 2012. Pervasive defaunation of forest remnants in a tropical biodiversity hotspot. *PLoS One* 7, e41671. doi:10.1371/journal.pone.0041671
- 465 Castello, L., McGrath, D.G., Hess, L.L., Coe, M.T., Lefebvre, P. A., Petry, P., Macedo, M.N., Renó, V.F., Arantes, C.C., 2013. The vulnerability of Amazon freshwater ecosystems. *Conserv. Lett.* 6, 217–229. doi:10.1111/conl.12008
- Chauvet, S., Forget, P.-M., 2005. Edge effects on post-dispersal seed removal in a fragmented rain forest in French Guiana. *J. Trop. Ecol.* 21, 113–116. doi:10.1017/S0266467404001944
- 470 Connor, E.F., McCoy, E.D., 1979. The Statistics and Biology of the Species-Area Relationship. *Am. Nat.* 113, 791–833
- Cosson, J.F., Ringuet, S., Claessens, O., de Massary, J.C., Daleky, A., Villiers, J.F., Granjon, L., Pons, J.M., 1999a. Ecological changes in recent land-bridge islands in French Guiana, with emphasis on vertebrate communities. *Biol. Conserv.* 91, 213–222
- 475 Cosson, J., Pons, J.-M., Masson, D., 1999b. Effects of forest fragmentation on frugivorous and nectarivorous bats in French Guiana. *J. Trop. Ecol.* 15, 515–534
- Crawley, M.J., 2005. *Statistics: an introduction using R.* John Wiley & Sons Ltd, Chichester
- Crawley, M.J., Harral, J.E., 2001. Scale dependence in plant biodiversity. *Science* 291, 864–868. doi:10.1126/science.291.5505.864
- 480 Demarty, M., Bastien, J., 2011. GHG emissions from hydroelectric reservoirs in tropical and equatorial regions - review of 20 years of CH₄ emission measurements. *Energy Policy* 39, 4197–4206. doi:10.1016/j.enpol.2011.04.033

- Develey, P.F., Stouffer, P.C., 2001. Effects of Roads on Movements by Understory Birds in Mixed-Species Flocks in Central Amazonian Brazil. *Conserv. Biol.* 15, 1416–1422. doi:10.1111/j.1523-1739.2001.00170.x
- 485 Diamond, J., 2001. Dammed Experiments! *Science* 294, 1847–1848
- Diamond, J.M., 1972. Biogeographic Kinetics: Estimation of Relaxation Times for Avifaunas of Southwest Pacific Islands. *Proc. Natl. Acad. Sci. USA* 69, 3199–3203. doi:10.1073/pnas.69.11.3199
- 490 EIA, 2013. International Energy Outlook 2013. U.S. Energy Information Administration, Office of Energy Analysis, U.S. Department of Energy, Washington, DC 20585
- Emer, C., Venticinque, E.M., Fonseca, C.R., 2013. Effects of dam-induced landscape fragmentation on Amazonian ant-plant mutualistic networks. *Conserv. Biol.* 27, 763–73. doi:10.1111/cobi.12045
- 495 Ewers, R.M., Didham, R.K., 2006. Confounding factors in the detection of species responses to habitat fragmentation. *Biol. Rev.* 81, 117–42. doi:10.1017/S1464793105006949
- Fahrig, L., 2003. Effects of Habitat Fragmentation on Biodiversity. *Annu. Rev. Ecol. Syst.* 34, 487–515. doi:10.1146/132419
- Fearnside, P.M., 2002. Greenhouse Gas Emissions from a Hydroelectric Reservoir (Brazil's Tucuruí Dam) and the Energy Policy Implications. *Water. Air. Soil Pollut.* 133, 69–96
- 500 Fearnside, P.M., 1999. Social Impacts of Brazil's Tucuruí Dam. *Environ. Manage.* 24, 483–495
- Fearnside, P.M., Pueyo, S., 2012. Greenhouse-gas emissions from tropical dams. *Nat. Clim. Chang.* 2, 382–384. doi:10.1038/nclimate1540
- 505 Feeley, K.J., Gillespie, T.W., Lebbin, D.J., Walter, H.S., 2007. Species characteristics associated with extinction vulnerability and nestedness rankings of birds in tropical forest fragments. *Anim. Conserv.* 10, 493–501. doi:10.1111/j.1469-1795.2007.00140.x
- Feeley, K.J., Terborgh, J.W., 2006. Habitat Fragmentation and Effects of Herbivore (Howler Monkey) Abundances on Bird Species Richness. *Ecology* 87, 144–150
- Feer, F., Hingrat, Y., 2005. Effects of Forest Fragmentation on a Dung Beetle Community in French Guiana. *Conserv. Biol.* 19, 1103–1112. doi:10.1111/j.1523-1739.2005.00087.x
- 510 Ferreira, L. V, Neckel-Oliveira, S., Galatti, U., Fáveri, S.B., Parolin, P., 2012. Forest structure of artificial islands in the Tucuruí dam reservoir in northern Brazil: a test core-area model. *Acta Amaz.* 42, 221–226
- 515 Finer, M., Jenkins, C.N., 2012. Proliferation of hydroelectric dams in the Andean Amazon and implications for Andes-Amazon connectivity. *PLoS One* 7, e35126. doi:10.1371/journal.pone.0035126
- Finer, M., Jenkins, C.N., Pimm, S.L., Keane, B., Ross, C., 2008. Oil and gas projects in the Western Amazon: Threats to Wilderness, Biodiversity and Indigenous Peoples. *PLoS One* 3, e2932. doi:10.1371/journal.pone.0002932


- 520 Gibson, L., Lynam, A.J., Bradshaw, C.J.A, He, F., Bickford, D.P., Woodruff, D.S., Bumrungsri, S.,
Laurance, W.F., 2013. Near-complete extinction of native small mammal fauna 25 years after
forest fragmentation. *Science* 341, 1508–10. doi:10.1126/science.1240495
- Gonzalez, A., 2000. Community relaxation in fragmented landscapes: the relation between species
richness, area and age. *Ecol. Lett.* 3, 441–448. doi:10.1046/j.1461-0248.2000.00171.x
- 525 Google, 2015. Google Earth Pro. URL: <http://www.google.co.uk/earth/download/gep/agree.html>
[Accessed 21/12/15]
- Haddad, N.M., Brudvig, L.A., Clobert, J., Davies, K.F., Gonzalez, A., Holt, R.D., Lovejoy, T.E.,
Sexton, J.O., Austin, M.P., Collins, C.D., Cook, W.M., Damschen, E.I., Ewers, R.M., Foster,
B.L., Jenkins, C.N., King, A.J., Laurance, W.F., Levey, D.J., Margules, C.R., Melbourne, B.A.,
530 Nicholls, A.O., Orrock, J.L., Song, D.-X., Townshend, J.R., 2015. Habitat fragmentation and its
lasting impact on Earth’s ecosystems. *Sci. Adv.* e1500052. doi: 10.1126/sciadv.1500052
- Halley, J.M., Sgardeli, V., Triantis, K.A., 2014. Extinction debt and the species-area relationship: a
neutral perspective. *Glob. Ecol. Biogeogr.* 23, 113–123. doi:10.1111/geb.12098
- Hanski, I., Gilpin, M., 1991. Metapopulation dynamics: brief history and conceptual domain. *Biol. J.
Linn. Soc.* 42, 3–16
- 535 Hanski, I., Ovaskainen, O., 2002. Extinction Debt at Extinction Threshold. *Conserv. Biol.* 16, 666–
673
- Haya, B., McCully, P., Pearson, B., 2002. Damming The CDM: Why Big Hydro Is Ruining the Clean
Development Mechanism. International Rivers Network/CDM Watch
- Hortal, J., Triantis, K.A., Meiri, S., Thébault, E., Sfenthourakis, S., 2009. Island Species Richness
540 Increases with Habitat Diversity. *Am. Nat.* 174, E205–E217. doi:10.1086/645085
- ICOLD, 2015. International Commission on Large Dams. URL: <http://www.icold-cigb.org/> [Accessed
21/12/15]
- International Energy Agency, 2006. Annex VIII, Hydropower Good Practices: Environmental
Mitigation Measures and Benefits. New Energy Foundation, Japan
- 545 International Energy Agency, 2000. Hydropower and the Environment: Effectiveness of Mitigation
Measures. IEA Technical Report, Annex III – Subtask 6
- Kareiva, P.M., 2012. Dam choices: Analyses for multiple needs. *Proc. Natl. Acad. Sci.* 109, 5553–
5554. doi:10.1073/pnas.1203263109
- 550 Kirby, K.R., Laurance, W.F., Albernaz, A.K., Schroth, G., Fearnside, P.M., Bergen, S., Venticinque,
E.M., da Costa, C., 2006. The future of deforestation in the Brazilian Amazon. *Futures* 38, 432–
453. doi:10.1016/j.futures.2005.07.011
- Kuussaari, M., Bommarco, R., Heikkinen, R.K., Helm, A., Krauss, J., Lindborg, R., Öckinger, E.,
Pärtel, M., Pino, J., Rodà, F., Stefanescu, C., Teder, T., Zobel, M., Steffan-Dewenter, I., 2009.
555 Extinction debt: a challenge for biodiversity conservation. *Trends Ecol. Evol.* 24, 564–571.
doi:10.1016/j.tree.2009.04.011

- Lambert, T.D., Adler, G.H., Riveros, C.M., Lopez, L., Ascanio, R., Terborgh, J., 2003. Rodents on tropical land-bridge islands. *J. Zool.* 260, 179–187. doi:10.1017/S0952836903003637
- Laurance, S.G.W., Stouffer, P.C., Laurance, W.F., 2004. Effects of road clearings on movement patterns of understory rainforest birds in central Amazonia. *Conserv. Biol.* 18, 1099–1109. doi:10.1111/j.1523-1739.2004.00268.x
- 560 Laurance, W., 2008. Theory meets reality: How habitat fragmentation research has transcended island biogeographic theory. *Biol. Conserv.* 141, 1731–1744. doi:10.1016/j.biocon.2008.05.011
- Lomolino, M. V., 2000. A call for a new paradigm of island biogeography. *Glob. Ecol. Biogeogr.* 9, 1–6. doi:10.1046/j.1365-2699.2000.00185.x
- 565 Lopez, L., Terborgh, J., 2007. Seed predation and seedling herbivory as factors in tree recruitment failure on predator-free forested islands. *J. Trop. Ecol.* 23, 129. doi:10.1017/S0266467406003828
- MacArthur, R.H., Wilson, E.O., 1963. An equilibrium theory of insular zoogeography. *Evolution* 17, 373–387
- 570 MacArthur, R.H., Wilson, E.O., 1967. *The theory of island biogeography*. Princeton University Press, Princeton
- Mace, G., Masundire, H., Baillie, J., 2005. Biodiversity, in: Ceballos, G., Lavorel, S., Orians, G., Pacala, S. (Eds.), *Ecosystems and Human Well-Being: Current State and Trends*
- 575 Mendenhall, C.D., Karp, D.S., Meyer, C.F.J., Hadly, E. A, Daily, G.C., 2014. Predicting biodiversity change and averting collapse in agricultural landscapes. *Nature* 509, 213–7. doi:10.1038/nature13139
- Mendenhall, C.D., Sekercioglu, C.H., Oviedo Brenes, F., Ehrlich, P.R., Daily, G.C., 2011. Predictive model for sustaining biodiversity in tropical countryside. *Proc. Natl. Acad. Sci. U. S. A.* 108, 16313–6. doi:10.1073/pnas.1111687108
- 580 Moore, R.P., Robinson, W.D., Lovette, I.J., Robinson, T.R., 2008. Experimental evidence for extreme dispersal limitation in tropical forest birds. *Ecol Lett* 11, 960–968. doi:10.1111/j.1461-0248.2008.01196.x
- Murcia, C., 1995. Edge effects in fragmented forests: implications for conservation. *Trends Ecol. Evol.* 10, 58–62. doi:10.1016/S0169-5347(00)88977-6
- 585 Nilsson, C., Reidy, C. A., Dynesius, M., Revenga, C., 2005. Fragmentation and flow regulation of the world's large river systems. *Science* 308, 405–8. doi:10.1126/science.1107887
- Oki, T., Kanae, S., 2006. Global Hydrological Cycles and World Water Resources. *Science* 313, 1068–1072. doi:10.1126/science.1128845
- 590 Oksanen, J., 2013. *Multivariate analysis of ecological communities in R: vegan tutorial*. doi:10.1016/0169-5347(88)90124-3
- Orrock, J.L., Watling, J.I., 2010. Local community size mediates ecological drift and competition in metacommunities. *Proc. Biol. Sci.* 277, 2185–91. doi:10.1098/rspb.2009.2344

- 595 Palmeirim, A.F., Peres, C. A., Rosas, F.C.W., 2014. Giant otter population responses to habitat expansion and degradation induced by a mega hydroelectric dam. *Biol. Conserv.* 174, 30–38. doi:10.1016/j.biocon.2014.03.015
- Peres, C. A., 2001. Synergistic Effects of Subsistence Hunting and Habitat Fragmentation on Amazonian Forest Vertebrates. *Conserv. Biol.* 15, 1490–1505. doi:10.1046/j.1523-1739.2001.01089.x
- 600 Peres, C. A., Lake, I.R., 2003. Extent of Nontimber Resource Extraction in Tropical Forests: Accessibility to Game Vertebrates by Hunters in the Amazon Basin. *Conserv. Biol.* 17, 521–535. doi:10.1046/j.1523-1739.2003.01413.x
- Pinheiro, J.C., Bates, D.M., 2000. *Mixed-effects Models in S and S-Plus*. Springer, New York. doi:10.1007/b98882
- 605 R Core Team, 2015. R: A Language and Environment for Statistical Computing. URL: <https://www.r-project.org/> [Accessed 21/12/15]
- Robinson, W.D., 1999. Long-Term Changes in the Avifauna of Barro Colorado Island, Panama, a Tropical Forest Isolate. *Conserv. Biol.* 13, 85–97. doi:10.1046/j.1523-1739.1999.97492.x
- 610 Sá-Oliveira, J.C., Hawes, J.E., Isaac-Nahum, V.J., Peres, C. A., 2015. Upstream and downstream responses of fish assemblages to an eastern Amazonian hydroelectric dam. *Freshw. Biol.* 60, 2037–2050. doi:10.1111/fwb.12628
- Seabloom, E.W., Dobson, A.P., Stoms, D.M., 2002. Extinction rates under nonrandom patterns of habitat loss. *Proc. Natl. Acad. Sci.* 99, 11229–11234. doi:10.1073/pnas.162064899
- St. Louis, V., Kelly, C.A., Duchemin, É., Rudd, J.W.M., Rosenberg, D.M., 2013. Reservoir Surfaces as Sources of Greenhouse Gases to the Atmosphere: A Global Estimate. *Bioscience* 50, 766–775
- 615 Terborgh, J., Lopez, L., Nuñez, P., Rao, M., Shahabuddin, G., Orihuela, G., Riveros, M., Ascanio, R., Adler, G.H., Lambert, T.D., Balbas, L., 2001. Ecological meltdown in predator-free forest fragments. *Science* 294, 1923–6. doi:10.1126/science.1064397
- 620 Terborgh, J., Lopez, L., Tello, J., Yu, D., Bruni, A.R., 1997. Transitory states in Relaxing Ecosystems of Land Bridge Islands, in: Laurance, W.F., Bierregaard, R.O. (Eds.), *Tropical Forest Remnants: Ecology, Management, and Conservation of Fragmented Communities*. University of Chicago Press, London, pp. 256–274
- Tilman, D., May, R.M., Lehman, C.L., Nowak, M.A., 1994. Habitat destruction and the extinction debt. *Nature* 371, 65–66
- 625 Triantis, K.A., Guilhaumon, F., Whittaker, R.J., 2012. The island species-area relationship: biology and statistics. *J. Biogeogr.* 39, 215–231. doi:10.1111/j.1365-2699.2011.02652.x
- Trussart, S., Messier, D., Roquet, V., Aki, S., 2002. Hydropower projects: A review of most effective mitigation measures. *Energy Policy* 30, 1251–1259. doi:10.1016/S0301-4215(02)00087-3
- 630 Van Houtan, K.S., Pimm, S.L., Halley, J.M., Bierregaard, R.O., Lovejoy, T.E., 2007. Dispersal of Amazonian birds in continuous and fragmented forest. *Ecol. Lett.* 10, 219–229. doi:10.1111/j.1461-0248.2007.01004.x

- Wang, Y., Zhang, J., Feeley, K.J., Jiang, P., Ding, P., 2009. Life-history traits associated with fragmentation vulnerability of lizards in the Thousand Island Lake, China. *Anim. Conserv.* 12, 329–337. doi:10.1111/j.1469-1795.2009.00257.x
- 635 Watson, D.M., 2002. A conceptual framework for studying species composition in fragments, islands and other patchy ecosystems. *J. Biogeogr.* 29, 823–834
- With, K. A., King, A.W., 2001. Analysis of landscape sources and sinks: The effect of spatial pattern on avian demography. *Biol. Conserv.* 100, 75–88. doi:10.1016/S0006-3207(00)00209-3
- WCD, 2000. Dams and Development: A new framework for decision-making. World Commission on Dams. Earthscan Publications Ltd, London and Sterling, VA
- 640 Yu, M., Hu, G., Feeley, K.J., Wu, J., Ding, P., 2012. Richness and composition of plants and birds on land-bridge islands: effects of island attributes and differential responses of species groups. *J. Biogeogr.* 39, 1124-1133. 1124. doi:10.1111/j.1365-2699.2011.02676.x

FIGURES







645

Figure 1: Geography of research detailing responses of terrestrial taxa to isolation on reservoir land-bridge islands. Dam names and the percentage of total dataset studies (n = 100) originating from each are presented. Broad habitat type is indicated by colour: dark green = wet tropical forest; light green = subtropical forest; yellow = tropical grassland (e.g. *cerrado*); cream = Mediterranean forest; blue = boreal forest.

650

[2 column fitting]



655 **Figure 2:** Overview of research presented within dataset studies (n=100). 2a-c) present the proportion of total studies (black bars) for habitat type, ecological metric and taxonomic group investigated respectively; pie charts represent overall response directions (red = negative; blue = positive; green = neutral; yellow = variable). 2d) presents the distribution of studies through island isolation time (red dashed line represents mean island isolation time, ~33 years).

660 **[Single column fitting]**




Figure 3: Analysis of species richness (S_{RICH}) data from 249 islands and 84 control sites available from nine dams in three broad habitat types (wet tropical forest, subtropical forest, and tropical grassland), modelled with time since island isolation (T_{ISO}) and island area ($AREA$). Bold black lines represent the slope for the overall model, with individual lines for each dam fitted using the BLUPs extracted from random effects. Colour indicates dam identity: grey = Petit Saut; green = Chiew Larn; magenta = Lago Guri; brown = Randenigala; light pink = Cabra Corral; orange = Lake Kenyir; purple = Balbina; red = Thousand Island Lake; blue = Gatun Lake. Axes are on a natural log scale.

[2 column fitting]

670

Table 1: Coefficient estimates for fixed effects in the most parsimonious model used for species richness analysis, with TAXA, DAM and STUDY as random effects; t-values >2 were treated as significant.

	Estimate	Standard Error	t-value
Intercept	-0.514	0.237	-2.168
AREA	0.185	0.015	11.944
T_{ISO}	-0.244	0.067	-3.641

[Single column fitting]

679 **APPENDIX A**

680 **Full reference list for Database A**

- 681 Adler, G.H., 1994. Tropical forest fragmentation and isolation promote asynchrony among
682 populations of a frugivorous rodent. *J. Anim. Ecol.* 63, 903–911
- 683 Adler, G.H., Beatty, R.P., 1997. Changing reproductive rates in a Neotropical forest rodent,
684 *Proechimys semispinosus*. *J. Anim. Ecol.* 66, 472–480
- 685 Adler, G.H., Seamon, J.O., 1991. Distribution and abundance of a tropical rodent, the spiny
686 rat, on islands in Panama. *J. Trop. Ecol.* 7, 349–360. doi:10.1017/S0266467400005617
- 687 Aponte, C., Barreto, G.R., Terborgh, J., 2003. Consequences of Habitat Fragmentation on
688 Age Structure and Life History in a Tortoise Population. *Biotropica* 35, 550–555
- 689 Arnold, A.E., Asquith, N.M., 2002. Herbivory in a fragmented tropical forest: patterns from
690 islands at Lago Gatun, Panama. *Biodivers. Conserv.* 11, 1663–1680
- 691 Asquith, N.M., Mejía-Chang, M., 2005. Mammals, Edge Effects, and the Loss of Tropical
692 Forest Diversity. *Ecology* 86, 379–390
- 693 Asquith, N.M., Terborgh, J., Arnold, A.E., Riveros, C.M., 1999. The fruits the agouti ate:
694 *Hymenaea courbaril* seed fate when its disperser is absent. *J. Trop. Ecol.* 15, 229–235
- 695 Asquith, N.M., Wright, S.J., Clauss, M.J., 1997. Does mammal community composition
696 control recruitment in neotropical forests? Evidence from Panama. *Ecology* 78, 941–946
- 697 Badano, E.I., Regidor, H.A., Nunez, H.A., Acosta, R., Gianoli, E., 2005. Species richness and
698 structure of ant communities in a dynamic archipelago: effects of island area and age. *J.*
699 *Biogeogr.* 32, 221–227
- 700 Bastos, H.B., Gonçalves, E.C., Ferrari, S.F., Silva, A., Schneider, M.P.C., 2010. Genetic
701 structure of red-handed howler monkey populations in the fragmented landscape of
702 Eastern Brazilian Amazonia. *Genet. Mol. Biol.* 33, 774–80
- 703 Bégin, Y., Sirois, L., Meunier, C., 2010. The effects of hydroelectric flooding on a reservoir's
704 peripheral forests and newly created forest islands, in: Stoffel, M., Bollschweiler, M.,
705 Butler, D.R., Luckman, B.H. (Eds.), *Tree Rings and Natural Hazards: A State-of-the-*
706 *Art, Advances in Global Change Research.* Springer Netherlands, Dordrecht, pp. 241–
707 256. doi:10.1007/978-90-481-8736-2
- 708 Benchimol, M., Peres, C.A., 2015a. Edge-mediated compositional and functional decay of
709 tree assemblages in Amazonian forest islands after 26 years of isolation. *J. Ecol.* 103,
710 408–420. doi:10.1111/1365-2745.12371
- 711 Benchimol, M., Peres, C.A., 2015b. Predicting local extinctions of Amazonian vertebrates in
712 forest islands created by a mega dam. *Biol. Conserv.* 187, 61–72.
713 doi:10.1016/j.biocon.2015.04.005

- 714 Benchimol, M., Peres, C.A., 2015c. Widespread Forest Vertebrate Extinctions Induced by a
715 Mega Hydroelectric Dam in Lowland Amazonia. PLoS One 1–15.
716 doi:10.5061/dryad.c301h
- 717 Benchimol, M., Venticinque, E.M., 2014. Responses of Primates to Landscape Change in
718 Amazonian Land-bridge islands-a Multi-scale Analysis. Biotropica 0, 1–9.
719 doi:10.1111/btp.12122
- 720 Benchimol, M., Venticinque, E.M., 2010. Harpy Eagle (*Harpia harpyja*) predation on an
721 Infant brown Capuchin Monkey (*Cebus apella*) in the Brazilian Amazon. Rev. Bras.
722 Ornitol. 18, 352–354
- 723 Boyett, W.D., Endries, M.J., Adler, G.H., 2000. Colonization-extinction dynamics of
724 opossums on small islands in Panama. Can. J. Zool. 78, 1972–1979
- 725 Brandão, R., Araújo, A., 2008. Changes in Anuran Species Richness and Abundance
726 Resulting from Hydroelectric Dam Flooding in Central Brazil. Biotropica 40, 263–266
- 727 Carvalho, L.S., Gomes, J.O., Neckel-Oliveira, S., Lo-Man-Hung, N.F., 2012. Microhabitat
728 use and intraspecific associations in the whip spider *Heterophrynus longicornis*
729 (Arachnida: Amblypygi) in forest fragments formed by the Tucuruí Dam lake , Pará ,
730 Brazil 46, 1263–1272
- 731 Cosson, J., Pons, J.M., Masson, D., 1999. Effects of forest fragmentation on frugivorous and
732 nectarivorous bats in French Guiana. J. Trop. Ecol. 15, 515–534
- 733 Cosson, J.F., Ringuet, S., Claessens, O., de Massary, J.C., Dalecky, A., Villiers, J.F.,
734 Granjon, L., Pons, J.M., 1999. Ecological changes in recent land-bridge islands in
735 French Guiana, with emphasis on vertebrate communities. Biol. Conserv. 91, 213–222.
736 doi:10.1016/S0006-3207(99)00091-9
- 737 Crete, M., Huot, J., Fortin, M.-J., Doucet, G., 1997. Comparison of plant and animal diversity
738 on new reservoir islands and established lake islands in Northern Boreal Forest of
739 Quebec. Can. field-naturalist 111, 407–416
- 740 Daleky, A., Chauvet, S., Ringuet, S., Claessens, O., Juda, J., Larue, M., Cosson, J.F., 2002.
741 Large mammals on small islands: short-term effects of forest fragmentation on the large
742 mammal fauna in French Guiana. Rev. d'écologie 57, 145–164
- 743 Davies, R., 2002. Feeding group responses of a Neotropical termite assemblage to rain forest
744 fragmentation. Oecologia 133, 233–242. doi:10.1007/s00442-002-1011-8
- 745 Davies, R.G., Hernández, L.M., Eggleton, P., Didham, R.K., Fagan, L.L., Winchester, N.N.,
746 2003. Environmental and spatial influences upon species composition of a termite
747 assemblage across neotropical forest islands. J. Trop. Ecol. 19, 509–524.
748 doi:10.1017/S0266467403003560
- 749 Ding, Z., Feeley, K.J., Wang, Y., Pakeman, R.J., Ding, P., 2013. Patterns of bird functional
750 diversity on land-bridge island fragments. J. Anim. Ecol. 781–790. doi:10.1111/1365-
751 2656.12046

- 752 Emer, C., Venticinque, E.M., Fonseca, C.R., 2013. Effects of dam-induced landscape
753 fragmentation on amazonian ant-plant mutualistic networks. *Conserv. Biol.* 27, 763–73.
754 doi:10.1111/cobi.12045
- 755 Ennen, J.R., Birkhead, R.D., Kreiser, B.R., Gaillard, D.L., Qualls, C.P., Lovich, J.E., 2011.
756 The effects of isolation on the demography and genetic diversity of long-lived species:
757 implications for conservation and management of the gopher tortoise (*Gopherus*
758 *polyphemus*). *Herpetol. Conserv. Biol.* 6, 202–214.
- 759 Estrada-Villegas, S., Meyer, C.F.J., Kalko, E.K.V., 2010. Effects of tropical forest
760 fragmentation on aerial insectivorous bats in a land-bridge island system. *Biol. Conserv.*
761 143, 597–608. doi:10.1016/j.biocon.2009.11.009
- 762 Feeley, K., 2003. Analysis of avian communities in Lake Guri, Venezuela, using multiple
763 assembly rule models. *Oecologia* 137, 104–13. doi:10.1007/s00442-003-1321-5
- 764 Feeley, K.J., Gillespie, T.W., Lebbin, D.J., Walter, H.S., 2007. Species characteristics
765 associated with extinction vulnerability and nestedness rankings of birds in tropical
766 forest fragments. *Anim. Conserv.* 10, 493–501. doi:10.1111/j.1469-1795.2007.00140.x
- 767 Feeley, K.J., Terborgh, J.W., 2008. Direct versus indirect effects of habitat reduction on the
768 loss of avian species from tropical forest fragments. *Anim. Conserv.* 11, 353–360.
769 doi:10.1111/j.1469-1795.2008.00182.x
- 770 Feeley, K.J., Terborgh, J.W., 2006. Habitat Fragmentation and Effects of Herbivore (Howler
771 Monkey) Abundances on Bird Species Richness. *Ecology* 87, 144–150
- 772 Feeley, K.J., Terborgh, J.W., 2005. The Effects of Herbivore Density on Soil Nutrients and
773 Tree Growth in Tropical Forest Fragments. *Ecology* 86, 116–124
- 774 Feer, F., Hingrat, Y., 2005. Effects of Forest Fragmentation on a Dung Beetle Community in
775 French Guiana. *Conserv. Biol.* 19, 1103–1112. doi:10.1111/j.1523-1739.2005.00087.x
- 776 Ferreira, L. V., Neckel-Oliveira, S., Galatti, U., Fáveri, S.B., Parolin, P., 2012. Forest
777 structure of artificial islands in the Tucuruí dam reservoir in northern Brazil: a test core-
778 area model. *Acta Amaz.* 42, 221–226.
- 779 Gibson, L., Lynam, A.J., Bradshaw, C.J. A, He, F., Bickford, D.P., Woodruff, D.S.,
780 Bumrungsri, S., Laurance, W.F., 2013. Near-complete extinction of native small
781 mammal fauna 25 years after forest fragmentation. *Science* 341, 1508–10.
782 doi:10.1126/science.1240495
- 783 Granjon, L., Cosson, J., Judas, J., Ringlet, S., 1996. Influence of tropical rainforest
784 fragmentation on mammal communities in French Guiana: short-term effects. *Acta*
785 *Oecologica* 17, 673–672
- 786 Granjon, L., Ringlet, S., Cheylan, G., 2002. Evolution of small terrestrial mammal species
787 richness on newly formed islands in primary tropical forest of French Guiana: a 6 year
788 study. *Rev. d'écologie* 8, 131–144.

- 789 Henry, M., Cosson, J.F., Pons, J.M., 2010. Modelling multi-scale spatial variation in species
790 richness from abundance data in a complex neotropical bat assemblage. *Ecol. Modell.*
791 221, 2018–2027. doi:10.1016/j.ecolmodel.2010.05.011
- 792 Henry, M., Cosson, J.-F., Pons, J.-M., 2007a. Abundance may be a misleading indicator of
793 fragmentation-sensitivity: The case of fig-eating bats. *Biol. Conserv.* 139, 462–467.
794 doi:10.1016/j.biocon.2007.06.024
- 795 Henry, M., Pons, J.-M., Cosson, J.-F., 2007b. Foraging behaviour of a frugivorous bat helps
796 bridge landscape connectivity and ecological processes in a fragmented rainforest. *J.*
797 *Anim. Ecol.* 76, 801–13. doi:10.1111/j.1365-2656.2007.01258.x
- 798 Hu, G., Feeley, K.J., Wu, J., Xu, G., Yu, M., 2011. Determinants of plant species richness
799 and patterns of nestedness in fragmented landscapes: evidence from land-bridge islands.
800 *Landsc. Ecol.* 26, 1405–1417. doi:10.1007/s10980-011-9662-7
- 801 Hu, G., Wu, J., Feeley, K.J., Xu, G., Yu, M., 2012. The effects of landscape variables on the
802 species-area relationship during late-stage habitat fragmentation. *PLoS One* 7, e43894.
803 doi:10.1371/journal.pone.0043894
- 804 Karr, J.R., 1990. Avian Survival Rates and the Extinction Process on Barro Colorado Island,
805 Panama Island. *Conserv. Biol.* 4, 391–397
- 806 Karr, J.R., 1982a. Population Variability and Extinction in the Avifauna of a Tropical Land
807 Bridge Island. *Ecology* 63, 1975–1978
- 808 Karr, J.R., 1982b. Avian Extinction on Barro Colorado Island, Panama: A Reassessment.
809 *Am. Nat.* 119, 220–239
- 810 Lambert, T.D., Adler, G.H., Riveros, C.M., Lopez, L., Ascanio, R., Terborgh, J., 2003.
811 Rodents on tropical land-bridge islands. *J. Zool.* 260, 179–187.
812 doi:10.1017/S0952836903003637
- 813 Larsen, T.H., Lopera, A., Forsyth, A., 2008. Understanding trait-dependent community
814 disassembly: dung beetles, density functions, and forest fragmentation. *Conserv. Biol.*
815 22, 1288–98. doi:10.1111/j.1523-1739.2008.00969.x
- 816 Larsen, T.H., Williams, N.M., Kremen, C., 2005. Extinction order and altered community
817 structure rapidly disrupt ecosystem functioning. *Ecol. Lett.* 8, 538–47.
818 doi:10.1111/j.1461-0248.2005.00749.x
- 819 Leigh, E.G., Wright, S.J., Herre, E.A., Putz, F.E., 1993. The decline of tree diversity on
820 newly isolated tropical islands: a test of a null hypothesis and some implications. *Evol.*
821 *Ecol.* 7, 76–99.
- 822 Lima, J., Galatti, U., Lima, C., Faveri, S., Vasconcelos, H., Neckel-Oliveira, S., 2015.
823 Amphibians on Amazonian Land-Bridge Islands are Affected More by Area Than
824 Isolation. *Biotropica* 47, 369–376

- 825 Lopez, G.O., Terborgh, J., Ceballos, N., 2005. Food selection by a hyperdense population of
826 red howler monkeys (*Alouatta seniculus*). *J. Trop. Ecol.* 21, 445–450.
827 doi:10.1017/S0266467405002415
- 828 Lopez, L., Terborgh, J., 2007. Seed predation and seedling herbivory as factors in tree
829 recruitment failure on predator-free forested islands. *J. Trop. Ecol.* 23, 129.
830 doi:10.1017/S0266467406003828
- 831 Lu, J., Jiang, L., Yu, L., Sun, Q., 2011. Local factors determine plant community structure on
832 closely neighbored islands. *PLoS One* 6, e19762. doi:10.1371/journal.pone.0019762
- 833 Lynam, A.J., 1997. Rapid decline of small mammal diversity in Monsoon evergreen
834 fragments in Thailand, in: *Tropical Forest Remnants: Ecology, Management, and*
835 *Conservation of Fragmented Communities*. pp. 222–240
- 836 Lynam, A.J., Billick, I., 1999. Differential responses of small mammals to fragmentation in a
837 Thailand tropical forest. *Biol. Conserv.* 91, 191–200
- 838 Mangan, S.A., Eom, A.-H., Adler, G.H., Yavitt, J.B., Herre, E.A., 2004. Diversity of
839 arbuscular mycorrhizal fungi across a fragmented forest in Panama: insular spore
840 communities differ from mainland communities. *Oecologia* 141, 687–700.
841 doi:10.1007/s00442-004-1684-2
- 842 Meyer, C.F.J., Kalko, E.K. V., 2008a. Assemblage-level responses of phyllostomid bats to
843 tropical forest fragmentation: land-bridge islands as a model system. *J. Biogeogr.* 35,
844 1711–1726. doi:10.1111/j.1365-2699.2008.01916.x
- 845 Meyer, C.F.J., Kalko, E.K. V., 2008b. Bat assemblages on Neotropical land-bridge islands:
846 nested subsets and null model analyses of species co-occurrence patterns. *Divers.*
847 *Distrib.* 14, 644–654. doi:10.1111/j.1472-4642.2007.00462.x
- 848 Meyer, C.F.J., Kalko, E.K. V., Kerth, G., 2009. Small-Scale Fragmentation Effects on Local
849 Genetic Diversity in Two Phyllostomid Bats with Different Dispersal Abilities in
850 Panama. *Biotropica* 41, 95–102. doi:10.1111/j.1744-7429.2008.00443.x
- 851 Milton, K., Lozier, J.D., Lacey, E. A., 2008. Genetic structure of an isolated population of
852 mantled howler monkeys (*Alouatta palliata*) on Barro Colorado Island, Panama.
853 *Conserv. Genet.* 10, 347–358. doi:10.1007/s10592-008-9584-3
- 854 Norconk, M.A., Grafton, B.W., 2003. Changes in forest composition and potential feeding
855 tree availability on a small land-bridge island in Lago Guri, Venezuela , in: Marsh, L.K.
856 (Ed.), *Primates in Fragments*. Springer US, pp. 211–227
- 857 Peetz, A., Norconk, M.A., Kinzey, W.G., 1992. Predation by jaguar on howler monkeys
858 (*Alouatta seniculus*) in Venezuela. *Am. J. Primatol.* 28, 223–228
- 859 Pons, J.-M., Cosson, J.-F., 2002. Use of forest fragments by animalivorous bats in French
860 Guiana. *Rev. d'ecologie* 57, 117–130

- 861 Qie, L., Lee, T.M., Sodhi, N.S., Lim, S.L.-H., 2011. Dung beetle assemblages on tropical
862 land-bridge islands: small island effect and vulnerable species. *J. Biogeogr.* 38, 792–
863 804. doi:10.1111/j.1365-2699.2010.02439.x
- 864 Rao, M., 2000. Variation in leaf-cutter ant (*Atta* sp.) densities in forest isolates: the potential
865 role of predation. *J. Trop. Ecol.* 16, 209–225.
- 866 Rao, M., Terborgh, J., Nunez, P., 2001. Increased Plant Herbivory in Forest Isolates:
867 Implications for Plant Community Structure and Composition. *Conserv. Biol.* 15, 624–
868 633.
- 869 Rebelo, H., Rainho, A., 2009. Bat conservation and large dams: spatial changes in habitat use
870 caused by Europe’s largest reservoir. *Endanger. Species Res.* 8, 61–68.
871 doi:10.3354/esr00100
- 872 Ringuet, S., 2000. An assessment of the potential influence of rainforest fragmentation on
873 small terrestrial mammal predation in French Guiana. *Rev. d’ecologie* 55, 101–116
- 874 Robinson, W.D., 1999. Long-Term Changes in the Avifauna of Barro Colorado Island,
875 Panama, a Tropical Forest Isolate. *Conserv. Biol.* 13, 85–97. doi:10.1046/j.1523-
876 1739.1999.97492.x
- 877 Shahabuddin, G., Herzner, G.A., Aponte, C., Gomez, M. del C., 2000. Persistence of a
878 frugivorous butterfly species in Venezuelan forest fragments: the role of movement and
879 habitat quality. *Biodivers. Conserv.* 9, 1623–1641
- 880 Shahabuddin, G., Ponte, C. A., 2005. Frugivorous butterfly species in tropical forest
881 fragments: correlates of vulnerability to extinction. *Biodivers. Conserv.* 14, 1137–1152.
882 doi:10.1007/s10531-004-7842-3
- 883 Shahabuddin, G., Terborgh, J.W., 2000. Frugivorous butterflies in Venezuelan forest
884 fragments: abundance, diversity and the effects of isolation. *J. Trop. Ecol.* 15, 703–722
- 885 Silva, S.S.B., Ferrari, S.F., 2009. Behavior patterns of southern bearded sakis (*Chiropotes*
886 *satanas*) in the fragmented landscape of eastern Brazilian Amazonia. *Am. J. Primatol.*
887 71, 1–7. doi:10.1002/ajp.20624
- 888 Terborgh, J., Feeley, K., Silman, M., Nunez, P., Balukjian, B., 2006. Vegetation dynamics of
889 predator-free land-bridge islands. *J. Ecol.* 94, 253–263. doi:10.1111/j.1365-
890 2745.2006.01106.x
- 891 Terborgh, J., Lopez, L., Nuñez, P., Rao, M., Shahabuddin, G., Orihuela, G., Riveros, M.,
892 Ascanio, R., Adler, G.H., Lambert, T.D., Balbas, L., 2001. Ecological meltdown in
893 predator-free forest fragments. *Science* 294, 1923–6. doi:10.1126/science.1064397
- 894 Terborgh, J., Lopez, L., Tello, J.S., 1997a. Bird Communities in Transition: The Lago Guri
895 Islands. *Ecology* 78, 1494–1501
- 896 Terborgh, J., Lopez, L., Tello, J.S., Yu, D., Bruni, A.R., 1997b. Transitory states in Relaxing
897 Ecosystems of Land Bridge Islands, in: Laurance, W.F., Bierregaard, R.O. (Eds.),

- 898 Tropical Forest Remnants: Ecology, Management, and Conservation of Fragmented
899 Communities. University of Chicago Press, London, pp. 256–274
- 900 Tremblay, A., Begin, Y., 2005. The effects of snow packing on tree growth forms on an
901 island in a recently created reservoir in northern Quebec. *Ecoscience* 12, 530–539
- 902 Tremblay, A., Begin, Y., 2000. The response of black spruce to the climatic influence of
903 Robert-Bourassa Reservoir, in northern Quebec. *Ecoscience* 7, 228–236
- 904 Veiga, L.M., Ferrari, S.F., 2006. Predation of Arthropods by Southern Bearded Sakis
905 (*Chiropotes satanas*) in Eastern Brazilian Amazonia. *Am. J. Primatol.* 68, 209–215.
906 doi:10.1002/ajp
- 907 Wang, Y., Zhang, J., Feeley, K.J., Jiang, P., Ding, P., 2009. Life-history traits associated with
908 fragmentation vulnerability of lizards in the Thousand Island Lake, China. *Anim.*
909 *Conserv.* 12, 329–337. doi:10.1111/j.1469-1795.2009.00257.x
- 910 Wang, Y., Bao, Y., Yu, M., Xu, G., Ding, P., 2010. Nestedness for different reasons: the
911 distributions of birds, lizards and small mammals on islands of an inundated lake.
912 *Divers. Distrib.* 16, 862–873. doi:10.1111/j.1472-4642.2010.00682.x
- 913 Wang, Y., Chen, S., Ding, P., 2011. Testing multiple assembly rule models in avian
914 communities on islands of an inundated lake, Zhejiang Province, China. *J. Biogeogr.* 38,
915 1330–1344. doi:10.1111/j.1365-2699.2011.02502.x
- 916 Wang, Y., Zhang, M., Wang, S., Ding, Z., Zhang, J., Sun, J., Li, P., Ding, P., 2012a. No
917 evidence for the small-island effect in avian communities on islands of an inundated
918 lake. *Oikos* 121, 1945–1952. doi:10.1111/j.1600-0706.2012.20322.x
- 919 Wang, Y., Wang, X., Ding, P., 2012b. Nestedness of snake assemblages on islands of an
920 inundated lake. *Curr. Zool.* 58, 828–836
- 921 Wang, Y., Zhu, Y., Wang, Y., 2012c. Differences in spatial genetic structure and diversity in
922 two mosses with different dispersal strategies in a fragmented landscape. *J. Bryol.* 34, 9–
923 16. doi:10.1179/1743282011Y.0000000035
- 924 Wang, Y., Lane, A., Ding, P., 2012d. Sex-biased dispersal of a frog (*Odorrana schmackeri*)
925 is affected by patch isolation and resource limitation in a fragmented landscape. *PLoS*
926 *One* 7, e47683. doi:10.1371/journal.pone.0047683
- 927 Weerakoon, W.M.S.K., 2001. Comparative richness of amphibians inhabiting the islands at
928 Randenigala reservoir with mainland of the Victoriya Randenigala Ranambe (VRR)
929 sanctuary. *Lyriocephalus* 44, 124–129
- 930 Willis, E.O., 1974. Populations and Local Extinctions of Birds on Barro Colorado Island,
931 Panama. *Ecol. Monogr.* 44, 153–169
- 932 Wright, S.J., Gompper, M.E., DeLeon, B., 1994. Are large predators keystone species in
933 Neotropical forests? The evidence from Barro Colorado Island. *Oikos* 71, 279–294

- 934 Yong, D.L., Lohman, D.J., Gan, C.W., Qie, L., Lim, S.L.-H., 2012. Tropical butterfly
935 communities on land-bridge islands in Peninsular Malaysia. *Raffles Bull. Zool.* 25, 161–
936 172
- 937 Yong, D.L., Qie, L., Sodhi, N.S., Koh, L.P., Peh, K.S.-H., Lee, T.M., Lim, H.C., Lim, S.L.-
938 H., 2010. Do insectivorous bird communities decline on land-bridge forest islands in
939 Peninsular Malaysia? *J. Trop. Ecol.* 27, 1–14. doi:10.1017/S0266467410000520
- 940 Yu, L., Lu, J., 2011. Does landscape fragmentation influence sex ratio of dioecious plants? A
941 case study of *Pistacia chinensis* in the Thousand-Island Lake region of China. *PLoS One*
942 6, e22903. doi:10.1371/journal.pone.0022903
- 943 Yu, M., Hu, G., Feeley, K.J., Wu, J., Ding, P., 2012. Richness and composition of plants and
944 birds on land-bridge islands: effects of island attributes and differential responses of
945 species groups. *J. Biogeogr.* 39, 1124–1133. doi:10.1111/j.1365-2699.2011.02676.x
- 946 Yuan, N., Comes, H.P., Mao, Y.-R., Qi, X.-S., Qiu, Y.-X., 2012. Genetic effects of recent
947 habitat fragmentation in the Thousand-Island Lake region of southeast China on the
948 distylous herb *Hedyotis chrysotricha* (Rubiaceae). *Am. J. Bot.* 99, 1715–25.
949 doi:10.3732/ajb.1200054
- 950 Zhang, X., Xu, G., Shen, D., Gu, Y., Gao, H., Luo, X., Chen, X., 2007. Maintenance and
951 natural regeneration of *Castanopsis sclerophylla* populations on islands of Qiandao Lake
952 Region, China. *Acta Ecol. Sin.* 27, 424–430. doi:10.1016/S1872-2032(07)60014-6
- 953

954 **APPENDIX B**955 **Table B1**

956 Summary of research articles used in the species richness analysis. For full references, see
 957 reference list below Table A1

Reference	Habitat type	Region	Country	Dam	Isolation time (years)	Number of islands studied	Number of control sites	Island areas (ha)
Badano et al. (2005)	Subtropical forest	South America	Argentina	Cabra Corral	15	9	1	0.16-62.5
Benchimol & Peres (2015)	Wet tropical forest	South America	Brazil	Balbina	26	34	12	<1-1690
Cosson et al. (1999)b	Wet tropical forest	South America	French Guiana	Petit Saut	1	6	3	2-40
Estrada-Villegas et al. (2010)	Wet tropical forest	Central America	Panama	Gatun Lake	92	8	6	2.5-50
Feer & Hingrat (2005)	Wet tropical forest	South America	French Guiana	Petit Saut	4	7	3	1.1-25.5
Gibson et al. (2013)	Wet tropical forest	Asia	Thailand	Chiew Larn	26	16	1	0.3-56.3
Granjon et al. (1996)	Wet tropical forest	South America	French Guiana	Petit Saut	1	10	1	0.35-30
Karr (1982)b	Wet tropical forest	Central America	Panama	Gatun Lake	66	1	2	1500
Leigh et al. (1993)	Wet tropical forest	Central America	Panama	Gatun Lake	66	7	4	0.6-1500
Meyer & Kalko (2008)a	Wet tropical forest	Central America	Panama	Gatun Lake	91	11	6	2.5-50
Pons & Cosson (2002)	Wet tropical forest	South America	French Guiana	Petit Saut	2	16	1	<6-28
Qui et al. (2011)	Wet tropical forest	Asia	Malaysia	Lake Kenyir	23	24	3	<1-383.3
Terborgh et al. (1997)	Wet tropical forest	South America	Venezuela	Lago Guri	9	12	1	1-350
Wang et al. (2009)	Subtropical forest	Asia	China	Thousand Island Lake	49	42	7	0.67-1289.23
Weerakoon (2009)	Subtropical forest	Asia	Sri Lanka	Randenigalla	12	6	5	2-167
Yong et al. (2010)	Wet tropical forest	Asia	Malaysia	Lake Kenyir	22	6	2	<20->100
Yong et al. (2012)	Wet tropical forest	Asia	Malaysia	Lake Kenyir	22	6	2	<20->100

958

959

960 **Table B2**

961 Coefficients for the fixed effects of models that treat study identity, dam, and taxonomic
 962 group as random effects. The Chi-square (x^2) value and p-value from model comparison by
 963 ANOVA is given. The final model used in analysis only included significant fixed effects:
 964 AREA and T_{ISO} . Values presented in this table are from model comparisons using a reduced dataset (n
 965 islands = 148) to account for missing values. Following model comparison, the final model was used
 966 on the full dataset (n islands = 249) which did not have missing values for the variables included in
 967 the model.

Fixed effects	Estimate	SE	t-value	df	x^2	p-value
Intercept	-0.514	0.237	-2.168			
AREA	0.237	0.02	11.958	1	94.744	<0.001
T_{ISO}	-0.328	0.069	-4.720	1	16.136	<0.001
D_{MAIN}	-0.037	0.039	-0.951	1	0.894	0.344
D_{ISLAND}	-0.062	0.043	-1.434	1	1.991	0.158

968

969 **Table B3**

970 Intercepts for the best linear unbiased predictors (BLUPs) for each dam generated using the
 971 ‘ranef’ function in lme4 (Bates et al., 2014).

Dam	Intercept (Dam)
Balbina	0.0367
Cabra Corral	-0.131
Chiew Larn	-0.071
Gatun Lake	0.361
Lago Guri	-0.059
Lake Kenyir	-0.247
Petit Saut	-0.079
Randenigala	-0.007
Thousand Island Lake	0.196

972