

Table 2. Overview of anatomical origin of isolates or specimen type.

Specimen	<i>K. pneumoniae</i>		<i>E. coli</i>	
	Non-susceptible to carbapenems (%)	Comparator isolates (%)	Non-susceptible to carbapenems (%)	Comparator isolates (%)
Urine	540 (44.9)	549 (50.0)	109 (56.2)	117 (56.2)
Blood	238 (19.8)	234 (21.3)	23 (11.9)	55 (26.4)
Lower respiratory tract	189 (15.7)	127 (11.6)	10 (5.1)	13 (6.2)
Wound	141 (11.7)	85 (7.7)	28 (14.4)	13 (6.2)
Puncture fluid	26 (2.2)	32 (2.9)	6 (3.1)	2 (1.0)
Abdominal	11 (0.9)	6 (0.5)	4 (2.1)	1 (0.5)
Soft tissue	10 (0.8)	9 (0.8)	6 (3.1)	1 (0.5)
Catheter exit point	5 (0.4)	16 (1.5)	3 (1.5)	2 (1.0)
Bone & Joint	1 (0.1)	0	1 (0.5)	0
Cerebral spinal fluid	0	1 (0.1)	0	0
Gynaecology	0	5 (0.5)	0	1 (0.5)
Other ¹	42 (3.5)	34 (3.1)	4 (2.1)	3 (1.4)
Total	1,203 (100)	1,098 (100)	194 (100)	208 (100)

¹ Other = specimen not attributed to any of the above.