

DIAGRAM MONOIDS AND GRAHAM–HOUGHTON GRAPHS: IDEMPOTENTS AND GENERATING SETS OF IDEALS

James East¹ and Robert D. Gray²

August 15, 2016

ABSTRACT. We study the ideals of the partition, Brauer, and Jones monoid, establishing various combinatorial results on generating sets and idempotent generating sets via an analysis of their Graham–Houghton graphs. We show that each proper ideal of the partition monoid \mathcal{P}_n is an idempotent generated semigroup, and obtain a formula for the minimal number of elements (and the minimal number of idempotent elements) needed to generate these semigroups. In particular, we show that these two numbers, which are called the rank and idempotent rank (respectively) of the semigroup, are equal to each other, and we characterize the generating sets of this minimal cardinality. We also characterize and enumerate the minimal idempotent generating sets for the largest proper ideal of \mathcal{P}_n , which coincides with the singular part of \mathcal{P}_n . Analogous results are proved for the ideals of the Brauer and Jones monoids; in each case, the rank and idempotent rank turn out to be equal, and all the minimal generating sets are described. We also show how the rank and idempotent rank results obtained, when applied to the corresponding twisted semigroup algebras (the partition, Brauer, and Temperley–Lieb algebras), allow one to recover formulae for the dimensions of their cell modules (viewed as cellular algebras) which, in the semisimple case, are formulae for the dimensions of the irreducible representations of the algebras. As well as being of algebraic interest, our results relate to several well-studied topics in graph theory including the problem of counting perfect matchings (which relates to the problem of computing permanents of $\{0, 1\}$ -matrices and the theory of Pfaffian orientations), and the problem of finding factorizations of Johnson graphs. Our results also bring together several well-known number sequences such as Stirling, Bell, Catalan and Fibonacci numbers.

1. INTRODUCTION

There has been a lot of interest recently in algebras with a basis consisting of diagrams that are multiplied in some natural diagrammatic way. Examples of such “diagram algebras” include the Brauer algebra [8], Temperley–Lieb algebra [60], and the Jones algebra [90]. All of these examples arise in a natural way as subalgebras of the partition algebra [102], whose basis consists of all set-partitions of a $2n$ -element set (see below for a formal definition). The partition algebra first appeared independently in the work of Martin [101, 102] and Jones [89]. In both cases, their motivation for studying this algebra was as a generalization of the Temperley–Lieb algebra and the Potts model in statistical mechanics. Since its introduction, the partition algebra has received a great deal of attention in the literature; see for example [47, 71, 74, 76–80, 82, 92, 103, 105, 106, 110, 136, 138].

All of the diagram algebras mentioned above are examples of cellular algebras, an important class of algebras introduced by Graham and Lehrer in [62]. The fact that these algebras are cellular allows one to obtain information about the semisimplicity of the algebra and about its representation theory, even in the non-semisimple case. In

2010 *Mathematics Subject Classification.* 20M20 (20M10, 20M17, 05E15, 05A18).

Key words and phrases. Partition monoid, Brauer monoid, Jones monoid, diagram algebra, generating sets, idempotents, rank, idempotent rank, Graham–Houghton graph.

¹Centre for Research in Mathematics; School of Computing, Engineering and Mathematics; Western Sydney University; Locked Bag 1797, Penrith, NSW, 2751, Australia. J.East@WesternSydney.edu.au

²School of Mathematics; University of East Anglia; Norwich NR4 7TJ, UK. Robert.D.Gray@uea.ac.uk
This author was partially supported by the EPSRC grant EP/N033353/1.

the partition, Brauer, and Temperley–Lieb algebras, the product of two diagram basis elements is always a scalar multiple of another basis element. Using this observation as a starting point, Wilcox [136], showed that these algebras are isomorphic to certain twisted semigroup algebras. By realizing the algebras in this way, many questions concerning the algebras can be related to questions for the corresponding semigroups. For instance, cellularity of the algebra can be deduced from various aspects of the structure of the monoid. The original study of cellular semigroup algebras may be found in [35]; see also [73, 74, 112, 113] for some recent developments. Another example of how the study of these semigroups can give information about the associated algebras may be found in work of the first author [36], who gives presentations for the partition monoid and shows how these presentations give rise to presentations for the partition algebra; see also [37, 43, 44]. A further example is given in the paper [27] where idempotents in the partition, Brauer and partial Brauer monoids are described and enumerated, and then the results are applied to determine the number of idempotent basis elements in the finite dimensional partition, Brauer and partial Brauer algebras; see also [28].

The corresponding semigroups—the partition, Brauer, and Jones monoids, and other related semigroups—have been studied, for instance, in [2–5, 25–27, 29, 36, 37, 41, 43, 44, 46, 51, 94, 96, 100, 104, 114, 115]. Recently, the first author [37] considered the subsemigroup generated by the set of idempotents in the partition monoid \mathcal{P}_n , showing in particular that every non-invertible element is expressible as a product of idempotents (we shall see in Theorem 7.5 below that this result holds more generally for any proper two-sided ideal of the partition monoid); presentations were also obtained in [37], and the infinite case was considered in [40, 46]. So the singular part of \mathcal{P}_n is an idempotent generated semigroup, and this is a property that \mathcal{P}_n has in common with several other naturally arising monoids. For instance, every non-invertible matrix from the full linear monoid $M_n(Q)$ of $n \times n$ matrices over an arbitrary division ring Q is expressible as a product of idempotent matrices [49, 95], and the same result is true for the full transformation semigroup of all maps from a finite set to itself [84]. Presentations for certain idempotent generated semigroups may be found in [37–39, 43, 44, 100]. More recently, in a significant extension of Erdos’s result from [49], Putcha [125] gave necessary and sufficient conditions for a reductive linear algebraic monoid to have the property that every non-unit is a product of idempotents. Another reason idempotent generated semigroups have received considerable attention in the literature is that they possess a universal property: every semigroup embeds into an idempotent generated semigroup [84] (indeed, in an idempotent generated regular $*$ -semigroup [46]), and if the semigroup is (finite) countable it can be embedded in a (finite) semigroup generated by three idempotents [11].

The Graham–Houghton graph of a semigroup is a bipartite graph with one part indexed by the \mathcal{R} -classes of the semigroup, the other part indexed by the \mathcal{L} -classes, and edges corresponding precisely to those \mathcal{H} -classes that contain idempotents (see Section 3 for the definition of Green’s relations). Graham introduced these graphs in [64] to study the idempotent generated subsemigroup of a 0-simple semigroup. Graham’s results were later rediscovered by Houghton who gave them a topological interpretation [83]. In the case that the semigroup is regular and idempotent generated, the connected components of this graph are in natural bijective correspondence with the \mathcal{D} -classes of the semigroup. Graham’s results show that these graphs are important tools for studying idempotent generated semigroups. More background on these graphs and their applications in semigroup theory may be found in [126, Section 4.13].

In addition to being a fundamental tool for studying products of idempotents, there are several other reasons that motivate the problem of obtaining a better understanding of Graham–Houghton graphs. Firstly, these graphs are important in the study of *free idempotent generated semigroups*. There has been a recent resurgence of interest in the study of these semigroups [9, 17, 18, 21, 31, 33, 34, 61, 68, 69, 119], with a particular focus

on describing their maximal subgroups. The theory developed in [9] shows that maximal subgroups of free idempotent generated semigroups are precisely the fundamental groups of Graham–Houghton complexes. The Graham–Houghton complex of an idempotent generated semigroup is a 2-complex, whose 1-skeleton is the Graham–Houghton graph of the semigroup, and which has 2-cells glued in for each *singular square* of idempotents (in the sense of Nambooripad [119]). Thus a necessary first step in determining these maximal subgroups is to obtain a description of the underlying Graham–Houghton graph. This has been done with success, and maximal subgroups have been computed, for certain fundamental examples such as the full transformation monoid [68] and the full linear monoid [31]. In contrast, currently nothing is known about the maximal subgroups of free idempotent generated semigroups arising from the partition, Brauer or Jones monoids. We hope the descriptions we obtain here of the Graham–Houghton graphs of the \mathcal{D} -classes of these semigroups will help with this research program.

Graham–Houghton graphs also play an important role in the representation theory of finite semigroups. Fundamental results of Clifford [12,13], Munn [117,118] and Ponizovskii [124] show that the irreducible representations of any finite semigroup can be parametrised in terms of the irreducible representations of its maximal subgroups. A full account of this theory may be found in [14, Chapter 5]. A short modern proof of the Clifford–Munn–Ponizovskii theorem is given in [54, Theorem 7]. This result gives a bijection between the irreducible representations of S and irreducible representations of the maximal subgroups G_{e_J} of S , where $\{e_J : J \in \mathcal{U}(S)\}$ is an idempotent transversal of the set $\mathcal{U}(S)$ of regular \mathcal{J} -classes of S . The work of Munn and Ponizovskii also gives necessary and sufficient conditions for the semigroup algebra KS over a field K to be semisimple. These conditions use Rees’s theorem and the notion of a Rees matrix semigroup. Associated to a regular \mathcal{J} -class J of a finite semigroup S is a (so-called) *Rees (0-)matrix semigroup* whose structure is governed by a group G_J and a “structure matrix” P with entries from $G_J \cup \{0\}$ (see Section 3 for more precise definitions). The locations of the non-zero entries of P are determined by the edges of the Graham–Houghton graph. The semigroup algebra KS is semisimple if and only if S is a regular semigroup and for each \mathcal{J} -class J , each KG_J is semisimple and the structure matrix corresponding to J is square and invertible. (Note that $\mathcal{J} = \mathcal{D}$ here since S is finite.)

These general results of course apply to the semigroup algebras of the monoids studied in this paper. In fact, as mentioned above, the algebras associated with the semigroups studied here actually all belong to a special class called cellular algebras. It was shown by Wilcox [136] that the partition, Brauer, and Temperley–Lieb algebras can be realised as twisted semigroup algebras of, respectively, the partition, Brauer and Jones monoids. There are twisted semigroup algebra analogues of the general results mentioned above: e.g., an analogous result giving necessary and sufficient conditions for semisimplicity; see [136, Theorem 15]. In the particular case of the diagram algebras we consider here, the results of Wilcox show how a cellular structure for the twisted semigroup algebra can be built from cellular structures of the group algebras of the maximal subgroups of the semigroup. Both the size and the structure of the Graham–Houghton graphs then encodes important information about these cellular structures, and thus about the corresponding diagram algebras and their representation theory. For example, the sizes of the Graham–Houghton graphs relate to the dimensions of the cell modules of the algebras. Indeed, in each case the Graham–Houghton graph is a balanced bipartite graph with each part of the bipartition having size equal to the number of \mathcal{R} -classes in the corresponding \mathcal{J} -class. These numbers can be used to relate the dimensions of cell modules for the twisted semigroup algebra to the dimensions of the cell modules corresponding to the group algebras of the maximal subgroups (see Equation 10.3). In this way, our results (specifically, Theorems 7.5, 8.4 and 9.5, which count the number of \mathcal{R} - and \mathcal{L} -classes in the \mathcal{J} -classes of the partition, Brauer and Jones monoids) allow for alternative derivations of (and in

some cases, alternative formulae for) the dimensions of the cell modules of the partition, Brauer and Temperley-Lieb algebras; see [7, 48, 62, 63, 72, 79, 91, 93, 107–109, 122, 134] for earlier derivations. Full details of this will be given in Section 10. The structure of the Graham–Houghton graph is also relevant when considering products of basis elements of the cellular basis of these twisted semigroup algebras. Indeed, the elements $C_{\mathfrak{s}, \mathfrak{t}}^\lambda$ of the cellular bases of these diagram algebras are all sums over elements from certain \mathcal{H} -classes in a corresponding diagram semigroup. A key step in understanding the representation theory of these cellular algebras is to investigate the bilinear form $\phi_\lambda : W(\lambda) \times W(\lambda) \rightarrow R$, $\phi_\lambda(C_{\mathfrak{s}}, C_{\mathfrak{t}}) = \phi(\mathfrak{s}, \mathfrak{t})$, which is defined by considering products of the form $C_{\mathfrak{s}_1, \mathfrak{t}_1}^\lambda C_{\mathfrak{s}_2, \mathfrak{t}_2}^\lambda$ (see Section 10 for details). In order to describe this bilinear form it is necessary to know when such a product “moves down” in the algebra, and in particular whether this sum moves down in the \mathcal{J} -order of the semigroup. This is determined precisely by the location of the idempotents in the \mathcal{J} -class of the elements arising in the sums defining $C_{\mathfrak{s}_1, \mathfrak{t}_1}^\lambda$ and $C_{\mathfrak{s}_2, \mathfrak{t}_2}^\lambda$. The Graham–Houghton graphs of these \mathcal{J} -classes record exactly this information.

In this paper we shall primarily be concerned with using Graham–Houghton graphs as a tool to study generating sets, idempotents, and subsemigroups generated by certain sets of idempotents, in the partition monoid and some of its key submonoids. Idempotent generated semigroups have been investigated using this approach in, for example [70, 85, 87, 97, 98]. There are numerous results in the literature concerning the problem of finding small generating sets (and generating sets of idempotents) for certain naturally arising semigroups, most often semigroups of transformations, matrix semigroups, and more generally semigroups of endomorphisms of various combinatorial or algebraic structures. One of the earliest results of this kind may be found in the work of Howie [85], where minimal idempotent generating sets of Sing_n , the singular part of the full transformation semigroup, were classified (by associating a tournament in a natural way with certain sets of idempotents); in subsequent work with McFadden [87], the ranks and idempotent ranks of arbitrary ideals of the full transformation monoid were calculated. Since then, many more results of this flavour have appeared in the literature [19, 20, 22–24, 30, 52, 53, 55, 57, 58, 65, 66, 97, 98]. Motivated in part by this work, for each of the proper ideals of the subsemigroups of the partition monoid we consider, we shall prove that the ideal is idempotent generated, and then investigate its generating sets, and idempotent generating sets, with a particular emphasis on describing small generating sets (and idempotent generating sets). We shall establish formulae for the smallest number of elements needed to generate the ideal (called the *rank*) and also the smallest number of idempotents required to generate the ideal (the *idempotent rank*). In all cases we will show that these two numbers coincide. This fact, together with some general results given in Section 4, can then be used to completely describe all the minimal generating sets for these ideals. We then go on to use Graham–Houghton graphs (and certain quotients of them) to study the idempotent generating sets in more detail. Specifically we investigate the problem of whether we can count all the idempotent generating sets of minimal cardinality. We shall see that this problem is equivalent to counting the number of perfect matchings in the corresponding Graham–Houghton graph. Counting perfect matchings in bipartite graphs is a well-studied problem in combinatorics, and relates to the problem of computing permanents of $\{0, 1\}$ -matrices and the theory of Pfaffian orientations; see [128]. We note that throughout the paper by a *minimal (idempotent) generating set* we will always mean a generating set (idempotent generating set) of minimal possible cardinality, as opposed to simply a generating set that is minimal with respect to set-theoretic inclusion (these two notions of minimality do not always coincide; see Example 7.16).

Including this introduction, the article comprises ten sections and is structured as follows. In Section 2, we present some results of Howie about idempotent generators in the full transformation monoid \mathcal{T}_n . We also take the opportunity to correct a mistake in the

formula given by Howie [85] for the number of distinct minimal idempotent generating sets for the singular part of \mathcal{T}_n . In Section 3, we present some general theory showing how (idempotent) generating sets for finite semigroups can be related to (idempotent) generating sets of their principal factors. In Section 4, we give some background on the theory of Graham–Houghton graphs, and how they can be used to investigate (idempotent) generating sets. In Section 5, we develop the theory introduced in earlier sections, tailoring it to the study of regular $*$ -semigroups. This is a class of regular semigroups that includes the partition monoid and all of the submonoids of the partition monoid considered in this article. In Section 6, we relate the ideas of Sections 4 and 5, showing how certain graphs defined in Section 5 are isomorphic to natural quotient graphs of the Graham–Houghton graphs discussed in Section 4. Specifically, we present a result that gives necessary and sufficient conditions for a set of idempotents to generate a given regular $*$ -semigroup, given in terms of an associated two-coloured directed graph. In Section 7, we turn our attention to the partition monoid \mathcal{P}_n . We prove that the proper two-sided ideals of \mathcal{P}_n are idempotent generated, and we give a formula for the rank and idempotent rank, showing that these are equal. We completely characterize the minimal generating sets, and minimal idempotent generating sets, and for the singular part of \mathcal{P}_n we also enumerate the minimal idempotent generating sets. We also apply the results of Section 6 to give necessary and sufficient conditions for a set of idempotents to generate the singular part of \mathcal{P}_n . The Brauer monoid is the subject of Section 8, where, as for the partition monoid, we consider its proper two sided ideals, showing they are idempotent generated, computing the idempotent rank, showing that it is equal to the rank, and characterizing the minimal generating sets. We also establish a bijection between the minimal idempotent generating sets of its singular part with certain factorizations of particular Johnson graphs. In Section 9, we consider the Jones monoid, where we establish analogous results for (idempotent) generating sets of its two-sided ideals, and also show that the number of distinct minimal idempotent generating sets is given by the n th Fibonacci number. Finally, in Section 10 we explain the connection between ranks (and idempotent ranks) of ideals of partition, Brauer and Jones monoids, and dimensions of cell modules (and irreducible representations) of the corresponding partition, Brauer, and Temperley–Lieb algebras.

2. IDEMPOTENT GENERATORS IN THE FULL TRANSFORMATION MONOID

In this section, we summarize some results of Howie and his collaborators [57, 84, 85, 87] on idempotent generators in the full transformation monoid. We do this in part because it will give a flavour of the kind of results we aim to obtain later on for ideals of the partition monoid and related monoids. We have also included this material so that we can correct a mistake in a formula given by Howie in [85]. Denote by \mathcal{T}_n and \mathcal{S}_n the full transformation semigroup and symmetric group on the set $[n] = \{1, \dots, n\}$, respectively. For $\alpha \in \mathcal{T}_n$ and $i \in [n]$, we write $i\alpha$ for the image of i under α ; in this way, transformations compose from left to right.

As mentioned in the introduction, the study of idempotent generated semigroups dates back to the paper [84], where Howie shows that the semigroup $\text{Sing}_n = \mathcal{T}_n \setminus \mathcal{S}_n$ is idempotent generated. In particular, he shows that Sing_n is generated by the set of idempotents with image size $n - 1$. In a later paper [85], he went on to study generating sets of idempotents in Sing_n in more detail, giving a combinatorial characterization of the minimal idempotent generating sets of the semigroup, and counting the number of such idempotent generating sets. In a subsequent work with McFadden [87], the ranks and idempotent ranks of the proper ideals of \mathcal{T}_n were obtained. See [59] for a historical overview of Howie’s work, including this particular research program.

In more detail, let $S = \text{Sing}_n = \mathcal{T}_n \setminus \mathcal{S}_n$. Set

$$F = \{\alpha \in E(S) : |\text{im } \alpha| = n - 1\},$$

where $E(S)$ denotes the set of idempotents in the semigroup S . In [84], it is shown that $\text{Sing}_n = \langle F \rangle$. Each $\beta \in F$ has the property that $i\beta = i$ for all $i \in \text{im } \beta$ (since $\beta^2 = \beta$) and since $|\text{im } \beta| = n - 1$ there is exactly one $i \in [n]$ such that $i\beta = j$ where $j \neq i$. In this case, we denote β by $\binom{i}{j}$, meaning β maps i to j and fixes every other point. Defining relations were given for Sing_n with respect to the generating set F in [38]; see also [37, 39, 42, 100] for presentations of other singular semigroups.

Let $X \subseteq F$. Define a digraph $\Gamma(X)$ with vertex set $[n]$ and an arc $j \rightarrow i$ if and only if $\binom{i}{j} \in X$. With this notation, in [85] the following result is obtained (recall that a digraph is a *tournament* if for each pair of distinct vertices x, y , the graph contains exactly one of the edges $x \rightarrow y$ or $y \rightarrow x$).

Theorem 2.1. *Let X be a set of idempotents from F in \mathcal{T}_n , where $n \geq 3$. Then X is a minimal (idempotent) generating set for Sing_n if and only if $\Gamma(X)$ is a strongly connected tournament.*

In particular, counting the number of arcs in a strongly connected tournament, one sees that

$$\text{rank}(\text{Sing}_n) = \text{idrank}(\text{Sing}_n) = \binom{n}{2} = \frac{n(n-1)}{2}.$$

It follows from the correspondence given by Theorem 2.1 that the number of minimal idempotent generating sets for Sing_n is precisely the number of strongly connected labeled tournaments on n vertices. The formula for this number stated in the paper [85] is actually incorrect. The correct formula may be found in a paper of Wright [137]; it may also be found as number sequence A054946 on [1]. Let w_n denote the number of strongly connected labeled tournaments on n vertices. Then w_n is given by the recurrence

$$w_1 = 1 \quad \text{and} \quad w_n = F_n - \sum_{s=1}^{n-1} \binom{n}{s} w_s F_{n-s} \quad \text{for } n \geq 2,$$

where $F_k = 2^{\binom{k}{2}} = 2^{k(k-1)/2}$. The first few terms in this sequence are displayed in Table 1. So, via the correspondence established by Theorem 2.1, for $n \geq 3$, w_n is the number of distinct minimal idempotent generating sets for Sing_n . Arbitrary minimal generating sets of Sing_n were characterized in [6]. Arbitrary idempotent generating sets of Sing_n were classified and enumerated in [23].

n	1	2	3	4	5	6	7	8	9	10
w_n	1	0	2	24	544	22320	1677488	236522496	64026088576	33832910196480

TABLE 1. The sequence w_n . For $n \geq 3$, w_n is equal to the number of minimal idempotent generating sets for $\text{Sing}_n = \mathcal{T}_n \setminus \mathcal{S}_n$.

In subsequent work, Howie and McFadden [87] investigated the ideals of \mathcal{T}_n . These are the sets

$$I_r(\mathcal{T}_n) = \{\alpha \in \mathcal{T}_n : |\text{im } \alpha| \leq r\}, \quad 1 \leq r \leq n.$$

Typically, the ideal $I_r(\mathcal{T}_n)$ is denoted $K(n, r)$, but we use the current notation for consistency with later usage.

Theorem 2.2. *For $1 \leq r \leq n - 1$, the ideal $I_r(\mathcal{T}_n)$ is idempotent generated, and*

$$\text{rank}(I_r(\mathcal{T}_n)) = \text{idrank}(I_r(\mathcal{T}_n)) = \begin{cases} n & \text{if } r = 1 \\ S(n, r) & \text{if } 2 \leq r \leq n - 1, \end{cases}$$

where $S(n, r)$ denotes the Stirling number of the second kind.

These (idempotent) ranks are given in Table 2.

$n \setminus r$	1	2	3	4	5	6	7	8	9	10
1	1									
2	1	1								
3	1	3	1							
4	1	7	6	1						
5	1	15	25	10	1					
6	1	31	90	65	15	1				
7	1	63	301	350	140	21	1			
8	1	127	966	1701	1050	266	28	1		
9	1	255	3025	7770	6951	2646	462	36	1	
10	1	511	9330	34105	42525	22827	5880	750	45	1

TABLE 2. The Stirling numbers of the second kind, $S(n, r)$. For $2 \leq r \leq n - 1$, $S(n, r)$ is equal to $\text{idrank}(I_r(\mathcal{T}_n)) = \text{rank}(I_r(\mathcal{T}_n))$.

3. SMALL GENERATING SETS: IDEALS AND PRINCIPAL FACTORS

We begin this section with some general observations about generating sets of semigroups that will explain why, in many situations, the problem of finding small generating sets for a given semigroup S reduces to the problem of finding small generating sets for a certain principal factor of S .

Let S be a semigroup, let S^1 denote S with an identity element adjoined (if necessary), and let $u, v \in S$. Recall that Green's relations \mathcal{R} , \mathcal{L} , \mathcal{J} , \mathcal{H} and \mathcal{D} are equivalence relations on S that reflect the ideal structure of the semigroup S and are defined by

$$u\mathcal{R}v \Leftrightarrow uS^1 = vS^1, \quad u\mathcal{L}v \Leftrightarrow S^1u = S^1v, \quad u\mathcal{J}v \Leftrightarrow S^1uS^1 = S^1vS^1,$$

$$\mathcal{H} = \mathcal{R} \cap \mathcal{L}, \quad \mathcal{D} = \mathcal{R} \vee \mathcal{L} = \mathcal{R} \circ \mathcal{L} = \mathcal{L} \circ \mathcal{R}.$$

All of the semigroups we consider in this paper will be finite, and in finite semigroups the relations \mathcal{J} and \mathcal{D} coincide. The \mathcal{J} relation gives rise to a natural preorder on S given by

$$u \leq_{\mathcal{J}} v \Leftrightarrow S^1uS^1 \subseteq S^1vS^1.$$

This preorder induces in a natural way a partial order on the set S/\mathcal{J} of \mathcal{J} -classes of S . By a maximal \mathcal{J} -class of a semigroup S we mean a \mathcal{J} -class that is maximal in this partially ordered set of \mathcal{J} -classes of S . An element $s \in S$ is called (von Neumann) *regular* if $s \in sSs$. If every element of S is regular we say that S is a *regular semigroup*. All the semigroups considered in this paper will be regular.

Recall that the principal factors of a semigroup S are the basic building blocks of the semigroup, and are obtained by taking a \mathcal{J} -class J and forming a semigroup $J^* = J \cup \{0\}$, where 0 is a symbol that does not belong to J , with multiplication given, for $s, t \in J \cup \{0\}$, by

$$s * t = \begin{cases} st & \text{if } s, t \text{ and } st \text{ belong to } J \\ 0 & \text{otherwise.} \end{cases}$$

If S is finite, then every principal factor J^* is either a semigroup with zero multiplication or a completely 0-simple semigroup (see the next paragraph). If, in addition, S is regular then every principal factor J^* is completely 0-simple.

We recall that a subset I of a semigroup is an *ideal* if $SI \subseteq S$ and $IS \subseteq S$. A semigroup S with zero is called *0-simple* if $\{0\}$ and S are its only ideals (and $S^2 \neq \{0\}$). It is said to be *completely 0-simple* if it is 0-simple and has 0-minimal left and right ideals. In particular

every finite 0-simple semigroup is completely 0-simple. Howie’s monograph [86] may be consulted for more information on completely 0-simple semigroups and other background on semigroup theory. An important result for our purposes is the Rees Theorem [86, Theorem 3.2.3] which says that any finite completely 0-simple semigroup is isomorphic to a (so-called) *regular Rees (0-)matrix semigroup* $\mathcal{M}^0[G; I, \Lambda; P]$ defined as follows. First, G is a finite group, I and Λ finite sets, and $P = (p_{\lambda,i})$ a $\Lambda \times I$ matrix with entries in $G \cup \{0\}$, where 0 is assumed not to belong to G , and such that each row and each column of P contains at least one non-zero entry. The semigroup $\mathcal{M}^0[G; I, \Lambda; P]$ is then defined on the set $(I \times G \times \Lambda) \cup \{0\}$, and with product defined by setting

$$(i, g, \lambda)(j, h, \mu) = (i, gp_{\lambda,j}h, \mu) \quad \text{if } p_{\lambda,j} \neq 0$$

and with all other products defined to be 0.

Recall that the *Rees quotient* of S by an ideal I , denoted S/I , is defined on underlying set $(S \setminus I) \cup \{0\}$, where 0 is a symbol not belonging to S , and with product \circ defined, for $s, t \in (S \setminus I) \cup \{0\}$, by

$$s \circ t = \begin{cases} st & \text{if } s, t \text{ and } st \text{ belong to } S \setminus I \\ 0 & \text{otherwise.} \end{cases}$$

A principal factor J^* (as defined above) is of course (isomorphic to) a special case of this construction: we take $S = \langle J \rangle$ and $I = \langle J \rangle \setminus J$.

The following observation lies at the heart of what is to follow: it is similar to [88, Lemma 3.2] (which is proved for monoids but may easily be adapted for semigroups); its proof is routine, and is omitted.

Lemma 3.1. *Let S be a finite semigroup and let I be an ideal of S . Moreover, suppose that S is generated by $U = S \setminus I$. Let A be a generating set of minimal cardinality for S . Then $A \subseteq U$ and*

$$\text{rank}(S) = \text{rank}(S/I).$$

Furthermore, S is idempotent generated if and only if S/I is idempotent generated, in which case

$$\text{idrank}(S) = \text{idrank}(S/I).$$

A special case of the above situation is when U is a maximal \mathcal{J} -class of S .

Lemma 3.2. *Let S be a semigroup with a maximal \mathcal{J} -class J such that $\langle J \rangle = S \neq J$. Then*

$$\text{rank}(S) = \text{rank}(J^*).$$

Furthermore, S is idempotent generated if and only if J^ is idempotent generated, in which case*

$$\text{idrank}(S) = \text{idrank}(J^*).$$

Proof. This is a straightforward consequence of the preceding lemma since $S \setminus J$ is an ideal of S , and the quotient $S/(S \setminus J)$ is isomorphic to the principal factor J^* . \square

Remark 3.3. Throughout this article, when we consider semigroups with a zero element (e.g., completely 0-simple semigroups) we will use $\langle X \rangle$ to denote all the elements that can be written as products of elements of X , together with the zero element if it is not already generated. In almost all cases where we apply results for completely 0-simple semigroups to principal factors of semigroups, including the partition monoid and related semigroups, this convention will not make any difference since all the completely 0-simple semigroups that arise will have the property that the zero element is generated by the other elements of the semigroup; the only exception is in the bottom ideal (which consists of a single \mathcal{J} -class). With this convention, we need not require that $S \neq J$ in the statement of Lemma 3.2.

Principal factors naturally come into play when analyzing idempotent generated semigroups, in part because of the following classical result of FitzGerald, which is distilled from the proof of [50, Lemma 2]; see also [126, Lemma 4.13.1].

Theorem 3.4. *Let S be a semigroup and let $e_1, \dots, e_m \in E(S)$. If $e_1 \cdots e_m$ is regular, then there is a sequence of (not necessarily distinct) idempotents $f_1, \dots, f_m \in E(S)$ such that $f_i \mathcal{J} e_1 \cdots e_m$ for all i and*

$$f_1 \cdots f_m = e_1 \cdots e_m.$$

The next result quickly follows.

Corollary 3.5. *Let S be a finite regular idempotent generated semigroup. Then for every \mathcal{J} -class J of S , the principal factor J^* is an idempotent generated completely 0-simple semigroup.*

Example 3.6 (Full transformation monoid). As before, let \mathcal{T}_n denote the full transformation monoid on n points. The proper two-sided ideals of \mathcal{T}_n are the sets

$$I_r = I_r(\mathcal{T}_n) = \{\alpha \in \mathcal{T}_n : \text{rank}(\alpha) \leq r\}, \quad 1 \leq r < n.$$

(The *rank*, $\text{rank}(\alpha)$, of a transformation $\alpha \in \mathcal{T}_n$ is equal to $|\text{im}(\alpha)|$.) Let

$$J_r = J_r(\mathcal{T}_n) = \{\alpha \in \mathcal{T}_n : \text{rank}(\alpha) = r\}.$$

Then it follows from [52, Lemma 2.2] that I_r is an idempotent generated semigroup with a unique maximal \mathcal{J} -class J_r , and $I_r = \langle J_r \rangle$. It follows that J_r^* is idempotent generated, with

$$\text{rank}(I_r) = \text{rank}(J_r^*) \quad \text{and} \quad \text{idrank}(I_r) = \text{idrank}(J_r^*),$$

where, since \mathcal{T}_n is regular, J_r^* is a completely 0-simple semigroup.

Example 3.7 (Full linear monoid). Let $M_n(F)$ denote the full linear monoid of all $n \times n$ matrices over a finite field F . The proper two-sided ideals of $M_n(F)$ are the sets

$$I_r = I_r(M_n(F)) = \{A \in M_n(F) : \text{rank}(A) \leq r\}, \quad 0 \leq r < n.$$

(The *rank*, $\text{rank}(A)$, of a matrix $A \in M_n(F)$ is equal to the dimension of its row or column space.) Let

$$J_r = J_r(M_n(F)) = \{A \in M_n(F) : \text{rank}(A) = r\}.$$

Then it also follows from [52, Lemma 2.2] that I_r is an idempotent generated semigroup with a unique maximal \mathcal{J} -class J_r and $I_r = \langle J_r \rangle$. It follows that J_r^* is idempotent generated, with

$$\text{rank}(I_r) = \text{rank}(J_r^*) \quad \text{and} \quad \text{idrank}(I_r) = \text{idrank}(J_r^*),$$

where, since $M_n(F)$ is regular, J_r^* is a completely 0-simple semigroup.

Remark 3.8. The two examples above admit a common generalization to endomorphism monoids of finite dimensional *independence algebras* [52]; see also [53] for a study of the infinite dimensional case.

Example 3.9 (Partition monoid). Let \mathcal{P}_n denote the partition monoid. There is a natural notion of *rank* for the elements of \mathcal{P}_n (see below for the definition). The proper two-sided ideals of \mathcal{P}_n are the sets

$$I_r = I_r(\mathcal{P}_n) = \{\alpha \in \mathcal{P}_n : \text{rank}(\alpha) \leq r\}, \quad 0 \leq r < n.$$

Let

$$J_r = J_r(\mathcal{P}_n) = \{\alpha \in \mathcal{P}_n : \text{rank}(\alpha) = r\}.$$

Then I_r is an idempotent generated semigroup with unique maximal \mathcal{J} -class J_r and $I_r = \langle J_r \rangle$ (see Lemma 7.4 below). Since \mathcal{P}_n is regular, it follows that J_r^* is an idempotent generated completely 0-simple semigroup satisfying

$$\text{rank}(I_r) = \text{rank}(J_r^*) \quad \text{and} \quad \text{idrank}(I_r) = \text{idrank}(J_r^*).$$

We will also see that similar statements are true of the Brauer and Jones monoids \mathcal{B}_n and \mathcal{J}_n .

Thus, in all the examples above— $I_r(\mathcal{T}_n)$, $I_r(M_n(F))$ and $I_r(\mathcal{P}_n)$ —the problem of determining rank and idempotent rank reduces to answering the same question for some idempotent generated completely 0-simple semigroup, so let us now turn our attention to this class.

4. IDEMPOTENT GENERATED COMPLETELY 0-SIMPLE SEMIGROUPS

The study of idempotent generated completely 0-simple semigroups is a classical topic, with early results by Graham, Houghton, Howie, etc. Here we summarize some results from [66].

Let $S = \mathcal{M}^0[G; I, \Lambda; P]$ be a finite completely 0-simple semigroup represented as a Rees matrix semigroup over a group G with I indexing the \mathcal{R} -classes of S , Λ indexing the \mathcal{L} -classes, and structure matrix P with entries from $G \cup \{0\}$. Our interest is in the case that S is idempotent generated. In this situation the problem of determining the rank has a straightforward answer. The next result has an easy inductive proof; see [67, Theorem 2.4] and [66, Lemma 2.2, Lemma 2.3].

Lemma 4.1. *Let $S = \mathcal{M}^0[G; I, \Lambda; P]$ be a finite completely 0-simple semigroup. If S is idempotent generated, then*

$$\text{rank}(S) = \max\{|I|, |\Lambda|\}.$$

In particular, in all of the examples discussed in Section 3, the rank is given simply by counting the number of \mathcal{R} -classes and \mathcal{L} -classes in each $\mathcal{D} = \mathcal{J}$ -class and choosing the larger of the two numbers.

Now, of course, in general for an idempotent generated semigroup S , we have

$$\text{idrank}(S) \geq \text{rank}(S). \tag{4.2}$$

It is easy to construct examples (even when S is a finite completely 0-simple semigroup) where $\text{rank}(S)$ and $\text{idrank}(S)$ are not equal; see for example [66, Example 2.7]. A natural question, therefore, is that of when they are equal. The main result of [66, Section 2] gives necessary and sufficient for equality in equation (4.2) where S is an arbitrary idempotent generated completely 0-simple semigroup; see [66, Theorem 2.16]. We shall not need the full generality of those results here but shall concentrate on a particularly nice special situation: namely, the case when $|I| = |\Lambda|$. As we shall see below, this condition is satisfied for the principal factors of the partition, Brauer and Jones monoids.

Before stating the results, we first need to introduce some definitions. The key idea is that the question of whether $\text{rank}(S) = \text{idrank}(S)$ comes down to a consideration of the way that the idempotents are distributed in S ; that is, the position of the non-zero elements in the structure matrix P .

Definition 4.3 (Graham–Houghton graph). Let $S = \mathcal{M}^0[G; I, \Lambda; P]$ be a Rees matrix semigroup. Let $\Delta(S)$ denote the bipartite graph with vertex set $I \cup \Lambda$ and an edge connecting $i \in I$ to $\lambda \in \Lambda$ if and only if $p_{\lambda i} \neq 0$. We call $\Delta(S)$ the *Graham–Houghton graph* of S .

Thus, the edges in $\Delta(S)$ record the positions of the idempotents in S . Graham–Houghton graphs will be discussed in more detail in Section 6 below.

Let Γ be a graph and let $v \in V(\Gamma)$ be a vertex of Γ . Define the neighbourhood of v to be the set

$$\Gamma(v) = \{w \in V(\Gamma) : w \sim v\}.$$

Here we use \sim for the adjacency relation in the graph. More generally, for a subset W of $V(\Gamma)$, the neighbourhood of W is the set

$$\Gamma(W) = \bigcup_{v \in W} \Gamma(v) = \{w \in V(\Gamma) : w \sim v (\exists v \in W)\}.$$

In the case that $S = \mathcal{M}^0[G; I, \Lambda; P]$ is an idempotent generated completely 0-simple semigroup with $|I| = |\Lambda|$, we have $\text{rank}(S) = |I| = |\Lambda|$. If $\text{idrank}(S) = \text{rank}(S)$, then there is an idempotent generating set with size $|I| = |\Lambda|$ which, since every generating set must clearly intersect every non-zero \mathcal{R} -class and every non-zero \mathcal{L} -class of S , must correspond to a perfecting matching in the bipartite graph $\Delta(S)$.

Thus, a necessary condition for equality in equation (4.2) in the case $|I| = |\Lambda|$ is that $\Delta(S)$ admits a perfect matching. It is a classical result from combinatorics, due to Hall [75], that a balanced bipartite graph admits a perfect matching if and only if it satisfies the so-called *Hall's condition*. In what follows, when we refer to a bipartite graph $\Gamma = X \sqcup Y$, we mean that the vertex set of Γ is the disjoint union of X and Y , and all edges of Γ join a vertex from X and a vertex from Y .

Theorem 4.4 (Hall [75]). *A bipartite graph $\Gamma = X \sqcup Y$ with $|X| = |Y|$ has a perfect matching if and only if the following condition is satisfied:*

$$|\Gamma(A)| \geq |A| \text{ for all } A \subseteq X. \tag{HC}$$

Although necessary, this property is not sufficient to guarantee $\text{idrank}(S) = \text{rank}(S)$ here, and what we actually require is a slight strengthening of Hall's condition:

Definition 4.5. A bipartite graph $\Gamma = X \sqcup Y$ with $|X| = |Y|$ is said to satisfy the *strong Hall condition* if it satisfies

$$|\Gamma(A)| > |A| \text{ for all } \emptyset \neq A \subsetneq X. \tag{SHC}$$

Definition 4.6 (Sparse cover). We call a subset A of a Rees matrix semigroup $S = \mathcal{M}^0[G; I, \Lambda; P]$ a *sparse cover* of S if $|A| = \max\{|I|, |\Lambda|\}$ and A intersects every non-zero \mathcal{R} -class and every non-zero \mathcal{L} -class of S .

If S is a finite idempotent generated completely 0-simple semigroup, then every generating set of S of minimum cardinality is a sparse cover, but in general the converse does not hold. We now state Theorem 2.10 from [66], which ties these ideas together.

Theorem 4.7. *Let $S = \mathcal{M}^0[G; I, \Lambda; P]$ be a finite idempotent generated completely 0-simple semigroup with $|I| = |\Lambda|$. Then the following are equivalent:*

- (i) $\text{rank}(S) = \text{idrank}(S)$;
- (ii) *the bipartite graph $\Delta(S)$ satisfies the strong Hall condition;*
- (iii) *every sparse cover of S generates S .*

5. GENERATING SETS OF REGULAR *-SEMIGROUPS

The partition, Brauer and Jones monoids (see Section 7 for the definitions) belong to the variety, introduced by Nordahl and Scheiblich [121], of *regular *-semigroups*. More generally, every proper two-sided ideal of each of the above-mentioned diagram semigroups is a regular *-semigroup that happens to be idempotent generated (see below). So, for all these examples, the problems of determining the rank and idempotent rank, and of describing minimal (idempotent) generating sets, may all be considered.

In this section we present some general results about small generating sets of regular *-semigroups, and then see how they can be applied to answer the above questions for the two-sided ideals of all of the examples mentioned above. In particular, in each case, by applying Theorem 4.7(iii), we will be able to completely describe the generating sets of minimal cardinality.

Definition 5.1. A semigroup S is called a *regular $*$ -semigroup* if there is a unary operation $*$: $S \rightarrow S$: $s \mapsto s^*$ satisfying the following conditions, for all $a, b \in S$:

$$(a^*)^* = a, \quad (ab)^* = b^*a^*, \quad aa^*a = a.$$

(We note that regular $*$ -semigroups are occasionally called *regular involution semigroups* [120] or *$*$ -regular semigroups* [123], even though the latter also refers to a completely different class of (involution) semigroups [15, 32]. We will use the original *regular $*$ -semigroup* terminology [121].)

(Note that $a^*aa^* = a^*$ follows from these axioms.) Clearly every regular $*$ -semigroup is regular, and aa^* and a^*a are idempotents \mathcal{R} - and \mathcal{L} -related, respectively, to a . But not all regular semigroups are regular $*$ -semigroups; for instance, regular $*$ -semigroups have square \mathcal{D} -classes (the number of \mathcal{R} - and \mathcal{L} -classes contained in any given \mathcal{D} -class must be equal), and not every regular semigroup has this property (consider the full transformation monoid, for example). On the other hand, every inverse semigroup (and hence every group) is easily seen to be a regular $*$ -semigroup, with $*$ taken as the inverse operation.

Many naturally arising semigroups are regular $*$ -semigroups, including many semigroups whose elements admit a diagrammatic representation, where the $*$ operation corresponds to taking the vertical mirror image of the diagram representing the element; this is the case for the Brauer, Jones and partition monoids (see below).

There is a special type of idempotent in regular $*$ -semigroups, the so-called *projections*; these play an important role in understanding the structure and generating sets of the semigroup.

Definition 5.2. An idempotent p in a regular $*$ -semigroup S is called a *projection* if $p^* = p$. If $A \subseteq S$, we write $E(A)$ (respectively, $P(A)$) for the set of all idempotents (respectively, projections) of S contained in A .

5.1. Idempotent generated regular $*$ -semigroups. Of course, not every regular $*$ -semigroup is idempotent generated; consider groups or inverse semigroups, for instance. However, some important, naturally arising semigroups of diagrams turn out to be idempotent generated; we shall see several examples below.

Our next aim is to show how generating sets of idempotents may be replaced by small generating sets of idempotents consisting only of projections. We begin with some basic observations about the behaviour of projections. These results are well known and we include proofs only for the sake of completeness.

Lemma 5.3. *Let S be a regular $*$ -semigroup. Then:*

- (i) $P(S) = \{aa^* : a \in S\} = \{a^*a : a \in S\}$;
- (ii) $E(S) = P(S)^2$. *In particular, the subsemigroup generated by the idempotents coincides with the subsemigroup generated by the projections, and S is idempotent generated if and only if it is generated by its projections;*
- (iii) *Every \mathcal{R} -class of S contains precisely one projection, as does every \mathcal{L} -class.*

Proof. (i) If $p \in P(S)$, then $p = pp^* = p^*p$. Conversely, it is easy to check that $aa^*, a^*a \in P(S)$ for any $a \in S$.

(ii) If $e \in E(S)$, then $e = ee^*e = e(ee^*)^*e = (ee^*)(e^*e)$. Conversely, if $p, q \in P(S)$, then $pq = pq(pq)^*pq = pqq^*p^*pq = pqqppq = (pq)^2$.

(iii) We prove the result for \mathcal{R} -classes; the result for \mathcal{L} -classes is dual. Let R be an \mathcal{R} -class. For any $a \in R$, $aa^* \in R$ is a projection. So each \mathcal{R} -class contains at least one projection. Now let $p, q \in R$ be projections. Then $p = qp$ and $q = pq$ (as $p\mathcal{R}q$), so $p = p^* = (qp)^* = p^*q^* = pq = q$, as required. \square

Theorem 5.4. *Let S be a finite idempotent generated completely 0-simple regular $*$ -semigroup with set of non-zero \mathcal{R} -classes and \mathcal{L} -classes indexed by I and Λ respectively. Then:*

- (i) $\text{rank}(S) = \text{idrank}(S) = |I| = |\Lambda|$;
- (ii) *a subset X of S is a minimal generating set for S if and only if X intersects each non-zero \mathcal{R} - and \mathcal{L} -class of S precisely once.*

Proof. By Lemma 5.3, since S is idempotent generated it is generated by its set P of non-zero projections and, since this set intersects each \mathcal{R} -class and \mathcal{L} -class exactly once, we conclude that $|P| = |I| = |\Lambda|$. Clearly (thinking of S as a Rees matrix semigroup), every generating set must intersect every non-zero \mathcal{R} - and \mathcal{L} -class, giving:

$$|I| = |\Lambda| \leq \text{rank}(S) \leq \text{idrank}(S) \leq |P| = |I| = |\Lambda|.$$

Part (ii) is then an immediate consequence of Theorem 4.7. \square

Definition 5.5 (Projection graph). Let S be a finite idempotent generated completely 0-simple regular $*$ -semigroup. Let $\Gamma(S)$ be the digraph with vertex set $P(S) \setminus \{0\}$ and edges $p \rightarrow q$ if and only if $pq \neq 0$. We call $\Gamma(S)$ the *projection graph* of S . If T is a finite regular $*$ -semigroup generated by the idempotents in a maximal \mathcal{J} -class J , we will often write $\Gamma(T) = \Gamma(J^*)$.

For some examples, see Figures 4, 11 and 20, below. Note that the edge relation on $\Gamma(S)$ is symmetric and reflexive (symmetry holds because $pq = 0 \Rightarrow qp = q^*p^* = (pq)^* = 0^* = 0$, for all $p, q \in P(S) \setminus \{0\}$). The relationship between the projection graph $\Gamma(S)$ and the Graham-Houghton graph $\Delta(S)$ will be explained in Section 6. By Lemma 5.3, the edges of $\Gamma(S)$ are in one-one correspondence with the nonzero idempotents of S .

Definition 5.6 (Balanced subgraph). We say a subgraph H of a digraph G is *balanced* if $V(H) = V(G)$ and the in- and out-degree of each vertex of H is equal to 1.

So a balanced subgraph partitions the digraph into disjoint directed cycles, which may contain one, two, or more vertices. Note that this generalises the notion of disjoint cycle decompositions of permutations. For example, if we begin with the complete directed graph KD_n (the digraph with n vertices, a loop at each vertex, and one arc in each direction between every pair of distinct vertices), then there is a natural correspondence between balanced subgraphs of KD_n and elements of the symmetric group \mathcal{S}_n , given by the obvious translation between balanced subgraphs and the elements of \mathcal{S}_n written as products of disjoint cycles. Thus, balanced subgraphs may be thought of as *permutation subgraphs*: subgraphs that involve all the vertices of the graph, and whose edges induce a permutation of the vertex set of the digraph. The set of all permutations arising from the balanced subgraphs of a directed graph is in general not a group.

Theorem 5.7. *Let S be a finite idempotent generated completely 0-simple regular $*$ -semigroup. Let G be a subgraph of $\Gamma(S)$, and let $X_G = \{pq : p \rightarrow q \text{ is an edge of } G\}$ be the corresponding set of idempotents of S . Then X_G is a minimal generating set for S if and only if G is balanced.*

Proof. Let $p \in P(S) \setminus \{0\}$. It is easy to check that $|X_G \cap R_p|$ is equal to the out-degree of p in G (here, R_p denotes the \mathcal{R} -class containing p). Together with the dual statement, and Theorem 5.4 (ii), this proves the result. \square

Theorem 5.8. *Let S be a finite regular semigroup, let J be a \mathcal{J} -class of S , X a subset of J , and $T = \langle X \rangle$, the subsemigroup of S generated by X . If T is regular, then*

$$J \cap T = J_1 \cup \dots \cup J_l,$$

where J_1, \dots, J_l are the maximal \mathcal{J} -classes of T .

Proof. Throughout the proof, we write \mathcal{J}^S and \mathcal{J}^T for Green's \mathcal{J} -relations on S and T , and similarly for the other Green's relations. We also write $J^T(x)$ for the \mathcal{J}^T -class in T of $x \in T$. Now let K be a maximal \mathcal{J}^T -class of T . Since $\langle X \rangle = T$ it follows that $K \leq_{\mathcal{J}^T} J^T(x)$ for some $x \in X$. Since K is maximal, we conclude that $K = J^T(x) \subseteq J$. Thus, every maximal \mathcal{J}^T -class of T is a subset of J .

Conversely, suppose K is a \mathcal{J}^T -class of T with $K \subseteq J$. The proof will be complete if we can show that K is maximal (in the ordering on \mathcal{J}^T -classes). Now, $K \leq_{\mathcal{J}^T} M$ for some maximal \mathcal{J}^T -class M of T . Let $a \in K$ and $b \in M$ be arbitrary. Since $K \leq_{\mathcal{J}^T} M$ (and since T is regular), there exist $u, v \in T$ such that $a = ubv$. Since $K, M \subseteq J$, we have $a \mathcal{J}^S b$, so $b \leq_{\mathcal{J}^S} ub \leq_{\mathcal{J}^S} ubv = a \leq_{\mathcal{J}^S} b$. Thus, all these elements are \mathcal{J}^S -related. In particular, $b \mathcal{J}^S ub$ and $ub \mathcal{J}^S ubv$. Since S is finite, stability (see [126, Definition A.2.1 and Theorem A.2.4]) gives $b \mathcal{L}^S ub$ and $ub \mathcal{R}^S ubv = a$. Since T is regular, it follows that $b \mathcal{L}^T ub \mathcal{R}^T a$, whence $b \mathcal{D}^T a$ and $b \mathcal{J}^T a$. Thus, $K = J^T(a) = J^T(b) = M$. \square

The above result fails if one lifts the assumption that the subsemigroup T is regular. One can easily construct a counterexample where T is a 4-element subsemigroup of the 5-element Brandt semigroup (see [86, Section 5.1] for more on Brandt semigroups).

Lemma 5.9. *Let S be a finite semigroup generated by the elements in its maximal \mathcal{J} -classes J_1, \dots, J_m . Then*

$$\text{rank}(S) = \sum_{i=1}^m \text{rank}(J_i^*).$$

If, in addition, S is idempotent generated, then so are all of J_1^, \dots, J_m^* and*

$$\text{idrank}(S) = \sum_{i=1}^m \text{idrank}(J_i^*).$$

Proof. This is an easy generalization of Lemma 3.2 above. \square

The following theorem, which is the main result of this section, says that in general, generating sets consisting of projections all taken from the same \mathcal{J} -class of a finite regular $*$ -semigroup always constitute a minimal generating set for the semigroup they generate. As we shall see, all of the examples of subsemigroups of \mathcal{P}_n discussed above, and their ideals, may be obtained as subsemigroups of the partition monoid generated by projections of fixed rank, and this allows us to apply the following result to determine the ranks and idempotent ranks in all cases, and ultimately to describe all the minimal generating sets.

Theorem 5.10. *Let S be a finite regular $*$ -semigroup, let J be a \mathcal{J} -class of S and let $X \subseteq J$ be a set of projections. Then the subsemigroup $T = \langle X \rangle$ of S generated by X satisfies*

$$\text{rank}(T) = \text{idrank}(T) = |X|.$$

Furthermore, a subset Y of T is a minimal generating set for T if and only if Y forms a transversal of the set of \mathcal{R} -classes, and set of \mathcal{L} -classes, contained in the maximal \mathcal{J} -classes of T .

Proof. First observe that T is an idempotent generated regular $*$ -semigroup. Indeed, to see that T is closed under the $*$ operation let $t \in T$ be arbitrary. Then $t = p_1 \cdots p_k$ for some $p_1, \dots, p_k \in X$, and so $t^* = p_k^* \cdots p_1^* = p_k \cdots p_1 \in T$.

Next, we claim that T is generated by the elements in its maximal \mathcal{J} -classes. Let J_1, \dots, J_l be the maximal \mathcal{J} -classes of T . It follows from Theorem 5.8 that $J_1 \cup \cdots \cup J_l = J \cap T \supseteq X$. Therefore, T is an idempotent generated regular $*$ -semigroup generated by

the elements in its maximal \mathcal{J} -classes $J_1 \cup \dots \cup J_l$. By Lemma 5.9,

$$\text{rank}(T) = \sum_{i=1}^l \text{rank}(J_i^*) \quad \text{and} \quad \text{idrank}(T) = \sum_{i=1}^l \text{idrank}(J_i^*).$$

For each i , Theorem 5.4 gives

$$\text{rank}(J_i^*) = \text{idrank}(J_i^*) = |X \cap J_i|,$$

since X intersects each \mathcal{R} - and \mathcal{L} -class of J_i exactly once. It follows that $\text{rank}(T) = \text{idrank}(T) = |X|$.

The last clause follows from Theorem 5.4. \square

Note that in general, in the above theorem, the semigroup T will have more than one maximal \mathcal{J} -class.

In each of the examples we will study, the \mathcal{J} -classes will form a chain, so the next general result will be of use.

Proposition 5.11. *Suppose S is a finite idempotent generated regular $*$ -semigroup and that the \mathcal{J} -classes of S form a chain $J_0 < J_1 < \dots < J_k$. For $0 \leq r \leq k$, let $I_r = J_0 \cup J_1 \cup \dots \cup J_r$. Then the ideals of S are precisely the sets I_0, I_1, \dots, I_k .*

Suppose further that $P(J_s) \subseteq \langle J_{s+1} \rangle$ for all $0 \leq s \leq k-1$. Then for each $0 \leq r \leq k$:

- (i) $I_r = \langle J_r \rangle$;
- (ii) I_r is idempotent generated;
- (iii) $\text{rank}(I_r) = \text{idrank}(I_r) = \rho_r$, where ρ_r is the number of \mathcal{R} -classes (which equals the number of \mathcal{L} -classes) in J_r .

Proof. The statement concerning the ideals of S is easily checked (and is true for any semigroup S whose \mathcal{J} -classes form a finite chain).

(i) Let $0 \leq s \leq k-1$. We first show that

$$J_s \subseteq \langle J_{s+1} \rangle. \tag{5.12}$$

First note that if $e \in E(J_s)$, then, as in the proof of Lemma 5.3, $e = (ee^*)(e^*e)$ where $ee^*, e^*e \in P(J_s)$. So $E(J_s) \subseteq \langle P(J_s) \rangle \subseteq \langle J_{s+1} \rangle$. By Corollary 3.5, every element of J_s is a product of idempotents from J_s , and (5.12) follows. Next note that $I_0 = J_0 = \langle J_0 \rangle$. If $1 \leq r \leq k$, then by (5.12) and an induction hypothesis, $I_r = I_{r-1} \cup J_r = \langle J_{r-1} \rangle \cup J_r \subseteq \langle J_r \rangle$.

(ii) Since S is idempotent generated, it follows from Corollary 3.5 that J_r^* is an idempotent generated completely 0-simple semigroup. Lemma 3.2 then implies that $I_r = \langle J_r \rangle$ is idempotent generated.

(iii) Since I_r is a regular $*$ -semigroup, Theorem 5.4 gives $\text{rank}(J_r^*) = \text{idrank}(J_r^*) = \rho_r$. By Lemma 3.2, $\text{rank}(I_r) = \text{rank}(J_r^*)$ and $\text{idrank}(I_r) = \text{idrank}(J_r^*)$. \square

Remark 5.13. In the notation of the previous proposition, it is clear that the number of minimal generating sets of the ideal I_r is equal to $\rho_r! \times h_r^{\rho_r}$, where h_r is the (common) size of the \mathcal{H} -classes of S contained in J_r . In general, a formula for the number of minimal idempotent generating sets is harder to come by. For example, to the authors' knowledge, such a general formula is unknown even in the case of the proper ideals of the singular part $S = \text{Sing}_n$ of the full transformation monoid \mathcal{T}_n .

6. NON-MINIMAL GENERATING SETS OF IDEMPOTENTS

Theorem 5.7 gives a necessary and sufficient condition, in terms of the projection graph $\Gamma(S)$, for a set of idempotents to be a minimal idempotent generating set for a finite idempotent generated completely 0-simple regular $*$ -semigroup S . Here we consider idempotent generating sets in general, not just those of minimal size. We begin by giving some background on Graham-Houghton graphs taken from [126, Section 4.13].

Definition 6.1. Let $S = \mathcal{M}^0[G; I, \Lambda; P]$ be a completely 0-simple semigroup represented as a Rees matrix semigroup over a group G . Recall from Definition 4.3 that the *Graham-Houghton graph*, denoted $\Delta(S)$, is the directed bipartite graph with vertex set $I \cup \Lambda$ and an arc $\lambda \rightarrow i$ if and only if $p_{\lambda i} \neq 0$. Now let F be a set of non-zero idempotents of S . We denote by $\Delta(S, F)$ the directed bipartite graph obtained from $\Delta(S)$ by adding arcs $i \rightarrow \lambda$ for every pair (i, λ) such that $H_{i, \lambda} \cap F \neq \emptyset$. (Here, $H_{i, \lambda} = R_i \cap L_\lambda$ denotes the \mathcal{H} -class that is the intersection of the \mathcal{R} - and \mathcal{L} -classes indexed by $i \in I$ and $\lambda \in \Lambda$, respectively, and we note that such an \mathcal{H} -class contains an idempotent if and only if $p_{\lambda i} \neq 0$, in which case this idempotent is $(i, p_{\lambda i}^{-1}, \lambda)$.)

The various parts of the next result follow from [64, Theorem 2 and subsequent discussion]; we include a proof for convenience.

Proposition 6.2. *Let $S = \mathcal{M}^0[G; I, \Lambda; P]$ be a finite idempotent generated completely 0-simple semigroup, and let F be a set of non-zero idempotents of S . Then the following are equivalent:*

- (i) $S = \langle F \rangle$;
- (ii) For all $(i, \lambda) \in I \times \Lambda$ there is a directed path in $\Delta(S, F)$ from i to λ ;
- (iii) The digraph $\Delta(S, F)$ is strongly connected.

Proof. First assume (i). Let $i \in I$ and $\lambda \in \Lambda$ be arbitrary, and consider a product $(i, 1, \lambda) = (i_1, p_{\lambda_1, i_1}^{-1}, \lambda_1) \cdots (i_k, p_{\lambda_k, i_k}^{-1}, \lambda_k)$, where each $(i_m, p_{\lambda_m, i_m}^{-1}, \lambda_m) \in F$ (and where 1 denotes the identity of G). First, we must have $i_1 = i$ and $\lambda_k = \lambda$. For the term $(i_m, p_{\lambda_m, i_m}^{-1}, \lambda_m)$ to belong to F , the graph $\Delta(S, F)$ must contain the edge $i_m \rightarrow \lambda_m$. And for the product to be non-zero, each $p_{\lambda_m, i_{m+1}}$ must be non-zero, so $\Delta(S, F)$ contains the edge $\lambda_m \rightarrow i_{m+1}$. So $\Delta(S, F)$ contains the path $i = i_1 \rightarrow \lambda_1 \rightarrow i_2 \rightarrow \cdots \rightarrow \lambda_k = \lambda$, giving (ii).

Next assume (ii). To prove (iii), it remains to show that for any $i, j \in I$ and $\lambda, \mu \in \Lambda$, $\Delta(S, F)$ contains a path from i to j , and from λ to μ . We just do the first of these, as the other is similar. By definition, there exists some $\kappa \in \Lambda$ such that $p_{\kappa, j} \neq 0$, so $\Delta(S, F)$ has the edge $\kappa \rightarrow j$. But, by assumption, $\Delta(S, F)$ also contains a path from i to κ . Appending the edge $\kappa \rightarrow j$ to this path yields a path from i to j .

Finally, assume (iii). Let $e \in E(S) \setminus \{0\}$. So $e = (i, p_{\lambda, i}^{-1}, \lambda)$ for some $i \in I$ and $\lambda \in \Lambda$. By assumption, there is a path $i = i_1 \rightarrow \lambda_1 \rightarrow i_2 \rightarrow \cdots \rightarrow \lambda_k = \lambda$ in $\Delta(S, F)$. It follows that the product $(i_1, p_{\lambda_1, i_1}^{-1}, \lambda_1) \cdots (i_k, p_{\lambda_k, i_k}^{-1}, \lambda_k)$ is non-zero and is \mathcal{H} -related to $e = (i, p_{\lambda, i}^{-1}, \lambda)$; note that each term in this product belongs to F . Since S is finite, and since $H_{i, \lambda}$ is a group (as it contains an idempotent), some positive power of this product is equal to e . In particular, $e \in \langle F \rangle$. Since this is true of an arbitrary non-zero idempotent e , and since S is idempotent generated, it follows that $S = \langle F \rangle$, giving (i). \square

In the special case that S is also a regular $*$ -semigroup, the digraph $\Delta(S)$ has an especially nice form, and this allows us to re-express Proposition 6.2 in terms of the digraph $\Gamma(S)$, defined in Definition 5.5, as follows.

Let $S = \mathcal{M}^0[G; I, \Lambda; P]$ be an idempotent generated completely 0-simple regular $*$ -semigroup. Then we may set $\Lambda = I' = \{i' : i \in I\}$, with the \mathcal{R} - and \mathcal{L} -classes indexed in such a way that the projections lie on the main diagonal \mathcal{H} -classes, $H_{i, i'}$ ($i \in I$). From this it then follows that the idempotents are distributed in the non-zero \mathcal{D} -class of S with diagonal symmetry, that is $H_{i, j'}$ contains an idempotent if and only if $H_{j, i'}$ does.

Using these observations it is easy to see that the graph $\Gamma(S)$ is isomorphic to the quotient digraph of $\Delta(S)$ obtained by identifying the pairs of vertices $\{i, i'\}$ for all $i \in I$. More explicitly, $\Gamma(S)$ is isomorphic to the digraph with vertex set $\{\{i, i'\} : i \in I\}$ where there is an arc $\{i, i'\} \rightarrow \{j, j'\}$ if and only if $H_{i, j'}$ is a group (equivalently, $H_{j, i'}$ is a group).

The digraph $\Gamma(S)$ is clearly symmetric, and has loops at every vertex. No arcs are identified when passing from $\Delta(S)$ to $\Gamma(S)$, and the arcs of $\Delta(S)$ (and thus also in $\Gamma(S)$) are in natural bijective correspondence with the non-zero idempotents of S .

Now we would like to re-express Proposition 6.2 in terms of the digraph $\Gamma(S)$.

Definition 6.3. Let F be a set of non-zero idempotents from a finite idempotent generated completely 0-simple regular $*$ -semigroup $S = \mathcal{M}^0[G; I, \Lambda; P]$. We use $\Gamma(S, F)$ to denote the 2-coloured digraph obtained by first colouring all the edges blue in the projection graph $\Gamma(S)$, and then adding the additional red arcs $i \rightarrow j$ for each idempotent $f \in F \cap H_{i,j}$.

Definition 6.4. Let G be a digraph with directed edges coloured red or blue. Call a directed path p in G an *RBR-alternating path* if the first and last arcs are red, and, as we traverse the path, the arcs alternate in colour between red and blue. An RBR-alternating circuit is an RBR-alternating path whose initial and terminal vertices coincide. In particular, a red loop at a vertex is an example of an RBR-alternating circuit.

Since S is generated by the projections, F will generate S if and only if every projection may be expressed as a product of elements from F . Combining this with Proposition 6.2 we obtain the following result.

Theorem 6.5. *Let $S = \mathcal{M}^0[G; I, \Lambda; P]$ be a finite idempotent generated completely 0-simple regular $*$ -semigroup, and let F be a set of non-zero idempotents from S . Then $S = \langle F \rangle$ if and only if every vertex in $\Gamma(S, F)$ is the base point of some RBR-alternating circuit.*

Remark 6.6. To directly see the significance of RBR-alternating paths and circuits, consider a finite idempotent generated completely 0-simple regular $*$ -semigroup S with non-zero projections p_1, \dots, p_k . Let F be a set of non-zero idempotents from S , and let

$$p_{i_1} \rightarrow p_{i_2} \rightarrow p_{i_3} \rightarrow \cdots \rightarrow p_{i_r}$$

be an RBR-alternating path in the graph $\Gamma(S, F)$. The red edges mean that the (non-zero) idempotents $p_{i_1}p_{i_2}, p_{i_3}p_{i_4}, \dots, p_{i_{r-1}}p_{i_r}$ all belong to F , so the product $s = p_{i_1}p_{i_2} \cdots p_{i_r}$ belongs to $\langle F \rangle$. The blue edges mean that the products $p_{i_2}p_{i_3}, \dots, p_{i_{r-2}}p_{i_{r-1}}$ are all non-zero, and it follows (thinking of S as a Rees matrix semigroup) that s is non-zero too. In the case that $i_r = i_1$ (so the path is a circuit), $s \mathcal{H} p_{i_1}$ and so, since S is finite, some power $s^t \in \langle F \rangle$ is equal to p_{i_1} . See Example 7.17.

Remark 6.7. The condition that every vertex is the base of an RBR-circuit is not equivalent to saying that each vertex is simply *contained* in some RBR-alternating circuit. For example, consider the graph in Figure 1. In this example $1 \rightarrow 2 \rightarrow 3 \rightarrow 1$ is an RBR-circuit containing each vertex, but vertices 2 and 3 are clearly not the base of any RBR-circuit.

Remark 6.8. We have obtained a necessary and sufficient condition for $S = \mathcal{M}^0[G; I, \Lambda; P]$ to be generated by a given set F of non-zero idempotents from S in terms of the graph $\Gamma(S, F)$. It would be nice to give a necessary and sufficient condition just in terms of the subgraph induced by the edges corresponding to the elements from F ; that is, the red edges. One might be tempted to conjecture that $S = \langle F \rangle$ if and only if, in $\Gamma(S, F)$, either:

- (i) each vertex has at least one red edge coming in to it and at least one going out of it, or
- (ii) each vertex is contained in a red circuit.

By Theorem 6.5, we see immediately that condition (i) is necessary, while condition (ii) is sufficient. However, we will see that (i) need not be sufficient, and (ii) need not be necessary. See Examples 7.16 and 7.17, Remarks 8.13 and 9.12, and also Theorem 9.10.




FIGURE 1. A graph showing that the condition that every vertex is the base of an RBR-circuit is not equivalent to saying that each vertex is contained in some RBR-alternating circuit. In all diagrams of graphs with red and blue edges, the red edges are drawn to be thicker than blue edges (for the convenience of readers with black and white copies of the article).

7. THE PARTITION MONOID

In the following sections, we will apply the general results of the previous sections to calculate the ranks and idempotent ranks for the proper two-sided ideals of the partition monoid, and several of its submonoids. In each case, we will also describe the (minimal) idempotent generating sets for the largest proper two-sided ideal, and also enumerate the minimal such generating sets for the partition and Jones monoids; a solution is not currently available for the latter problem with respect to the Brauer monoid (see Remark 8.9).

Let n be a positive integer, which we fix throughout this section, and write $[n]$ for the finite set $\{1, \dots, n\}$. If $1 \leq r \leq s \leq n$, we write $[r, s] = \{r, r + 1, \dots, s\}$. We also write $[n]' = \{1', \dots, n'\}$ for a set in one-one correspondence with $[n]$. An n -partition (or simply a partition if n is understood from context) is an equivalence relation on $[n] \cup [n]'$. We regard the elements of \mathcal{P}_n formally as subsets of $([n] \cup [n]') \times ([n] \cup [n]')$ satisfying reflexivity, symmetry and transitivity; so we will write $(x, y) \in \alpha$ to mean that x and y are α -related elements of $[n] \cup [n]'$. The set \mathcal{P}_n of all n -partitions forms a monoid of size $|\mathcal{P}_n| = B_{2n}$ (the $2n$ th Bell number), known as the *partition monoid*, under an associative operation we will describe shortly.

A partition $\alpha \in \mathcal{P}_n$ may be represented by a graph on the vertex set $[n] \cup [n]'$ as follows. We arrange vertices $1, \dots, n$ in a row (increasing from left to right) and vertices $1', \dots, n'$ in a parallel row directly below. We then add edges in such a way that two vertices x, y are connected by a path if and only if $(x, y) \in \alpha$. For example, the partition from \mathcal{P}_6 with equivalence classes $\{1\}, \{2, 3', 4'\}, \{3, 4\}, \{5, 6, 1', 5', 6'\}, \{2'\}$ is represented by the graph . Of course, such a graphical representation is not unique, but we will identify two graphs on the vertex set $[n] \cup [n]'$ if they have the same connected components. In the same way, we will not distinguish between a partition and a graph that represents it.

In order to describe the product alluded to above, let $\alpha, \beta \in \mathcal{P}_n$. We first stack (graphs representing) α and β so that vertices $1', \dots, n'$ of α are identified with vertices $1, \dots, n$ of β . The connected components of this graph are then constructed, and we finally delete the middle row of vertices as well as any connected components that are contained within the middle row; the resulting graph is the product $\alpha\beta$. As an example, consider the two partitions $\alpha =$ and $\beta =$ from \mathcal{P}_6 . The product $\alpha\beta =$ is found by moving through the following intermediate stages:

$$\begin{array}{c} \alpha = \\ \beta = \end{array} \begin{array}{c} \text{graph} \\ \text{graph} \end{array} \rightarrow \begin{array}{c} \text{graph} \\ \text{graph} \end{array} \rightarrow \begin{array}{c} \text{graph} \\ \text{graph} \end{array} \rightarrow \begin{array}{c} \text{graph} \\ \text{graph} \end{array} = \alpha\beta.$$

We now introduce some notation and terminology that we will use throughout our study. Let $\alpha \in \mathcal{P}_n$. The equivalence classes of α are called its *blocks*. A block of α is called a

transversal block if it has nonempty intersection with both $[n]$ and $[n]'$, and a *nontransversal block* otherwise. The *rank* of α , denoted $\text{rank}(\alpha)$, is equal to the number of transversal blocks of α . For $x \in [n] \cup [n]'$, let $[x]_\alpha$ denote the block of α containing x . We define the *domain* and *codomain* of α to be the sets

$$\begin{aligned}\text{dom}(\alpha) &= \{x \in [n] : [x]_\alpha \cap [n]' \neq \emptyset\}, \\ \text{codom}(\alpha) &= \{x \in [n] : [x']_\alpha \cap [n] \neq \emptyset\}.\end{aligned}$$

We also define the *kernel* and *cokernel* of α to be the equivalences

$$\begin{aligned}\ker(\alpha) &= \{(x, y) \in [n] \times [n] : [x]_\alpha = [y]_\alpha\}, \\ \text{coker}(\alpha) &= \{(x, y) \in [n] \times [n] : [x']_\alpha = [y']_\alpha\}.\end{aligned}$$

To illustrate these ideas, let $\alpha = \begin{array}{c} \bullet \quad \bullet \quad \bullet \quad \bullet \\ \diagdown \quad \diagup \quad \diagdown \quad \diagup \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \in \mathcal{P}_6$. Then $\text{rank}(\alpha) = 2$, $\text{dom}(\alpha) = \{2, 5, 6\}$, $\text{codom}(\alpha) = \{1, 3, 4, 5, 6\}$, and α has kernel-classes $\{1\}$, $\{2\}$, $\{3, 4\}$, $\{5, 6\}$ and cokernel-classes $\{1, 5, 6\}$, $\{2\}$, $\{3, 4\}$.

It is immediate from the definitions that

$$\begin{aligned}\text{dom}(\alpha\beta) &\subseteq \text{dom}(\alpha), & \ker(\alpha\beta) &\supseteq \ker(\alpha), \\ \text{codom}(\alpha\beta) &\subseteq \text{codom}(\beta), & \text{coker}(\alpha\beta) &\supseteq \text{coker}(\beta)\end{aligned}$$

for all $\alpha, \beta \in \mathcal{P}_n$. Let Δ denote the trivial equivalence relation (that is, the equality relation) on $[n]$. It is also clear that

$$\begin{aligned}\text{dom}(\beta) = [n] &\Rightarrow \text{dom}(\alpha\beta) = \text{dom}(\alpha), \\ \text{codom}(\alpha) = [n] &\Rightarrow \text{codom}(\alpha\beta) = \text{codom}(\beta), \\ \ker(\beta) = \Delta &\Rightarrow \ker(\alpha\beta) = \ker(\alpha), \\ \text{coker}(\alpha) = \Delta &\Rightarrow \text{coker}(\alpha\beta) = \text{coker}(\beta)\end{aligned}$$

for all $\alpha, \beta \in \mathcal{P}_n$. In particular, the sets

$$\begin{aligned}\{\alpha \in \mathcal{P}_n : \text{dom}(\alpha) = [n]\}, & & \{\alpha \in \mathcal{P}_n : \ker(\alpha) = \Delta\}, \\ \{\alpha \in \mathcal{P}_n : \text{codom}(\alpha) = [n]\}, & & \{\alpha \in \mathcal{P}_n : \text{coker}(\alpha) = \Delta\}\end{aligned}$$

are all submonoids of \mathcal{P}_n . The intersection of these four submonoids is (isomorphic to) the symmetric group \mathcal{S}_n , which is easily seen to be the group of units of \mathcal{P}_n .

If $x \in [n]$, we write $x'' = x$. For $\alpha \in \mathcal{P}_n$, we define $\alpha^* = \{(x', y') : (x, y) \in \alpha\}$. Diagrammatically, α^* is obtained by reflecting (a graph representing) α in a horizontal axis. For example, if $\alpha = \begin{array}{c} \bullet \quad \bullet \quad \bullet \quad \bullet \\ \diagdown \quad \diagup \quad \diagdown \quad \diagup \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \in \mathcal{P}_6$, then $\alpha^* = \begin{array}{c} \bullet \quad \bullet \quad \bullet \quad \bullet \\ \diagup \quad \diagdown \quad \diagup \quad \diagdown \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$. The map $\mathcal{P}_n \rightarrow \mathcal{P}_n : \alpha \mapsto \alpha^*$ illustrates the regular $*$ -semigroup structure of \mathcal{P}_n ; for all $\alpha, \beta \in \mathcal{P}_n$, we have

$$(\alpha^*)^* = \alpha, \quad (\alpha\beta)^* = \beta^*\alpha^*, \quad \alpha\alpha^*\alpha = \alpha.$$

We also have $\text{codom}(\alpha) = \text{dom}(\alpha^*)$ and $\text{coker}(\alpha) = \ker(\alpha^*)$ and other such identities.

We say a partition $\alpha \in \mathcal{P}_n$ is *planar* if it has a graphical representation without any crossings. The set of all planar partitions forms a submonoid of \mathcal{P}_n , and we denote this submonoid by \mathcal{PP}_n . The Brauer monoid \mathcal{B}_n is the submonoid of \mathcal{P}_n consisting of all partitions whose blocks all have size 2. The Jones monoid \mathcal{J}_n is the intersection of \mathcal{PP}_n with \mathcal{B}_n . We will concentrate on the partition monoid itself in this section, and will return to the three submonoids in subsequent sections. The next result was first proved (using different language) in [136]; see also [51]. It also follows from some of the above identities.

Theorem 7.1 (Wilcox [136, Theorem 17]). *For each $\alpha, \beta \in \mathcal{P}_n$, we have:*

- (i) $\alpha\mathcal{R}\beta$ if and only if $\text{dom}(\alpha) = \text{dom}(\beta)$ and $\ker(\alpha) = \ker(\beta)$;
- (ii) $\alpha\mathcal{L}\beta$ if and only if $\text{codom}(\alpha) = \text{codom}(\beta)$ and $\text{coker}(\alpha) = \text{coker}(\beta)$;
- (iii) $\alpha\mathcal{J}\beta$ if and only if $\text{rank}(\alpha) = \text{rank}(\beta)$.

We will also require the following result from [37]; see also [44, 46]. For $1 \leq i \leq n$, let $\pi_i \in \mathcal{P}_n$ be the projection with domain $[n] \setminus \{i\}$ and kernel Δ . For $1 \leq i < j \leq n$, let \mathcal{E}_{ij} be the equivalence on $[n]$ whose only non-trivial equivalence class is $\{i, j\}$, and let $\pi_{ij} \in \mathcal{P}_n$ be the projection with domain $[n]$ and kernel \mathcal{E}_{ij} . See Figure 2 for an illustration.




FIGURE 2. The projections $\pi_i, \pi_{ij} \in \mathcal{P}_n$.

Theorem 7.2. *The singular part $\mathcal{P}_n \setminus \mathcal{S}_n$ of the partition monoid \mathcal{P}_n is idempotent generated. The set $\{\pi_i : 1 \leq i \leq n\} \cup \{\pi_{ij} : 1 \leq i < j \leq n\}$ is a minimal idempotent generating set.*

Note that this is precisely the set of all projections of rank $n - 1$. Defining relations for this generating set were given in [37], but we will not need them here.

If $A \subseteq [n]$, we write $A' = \{a' : a \in A\} \subseteq [n]'$. As in [46], if $\alpha \in \mathcal{P}_n$, we will write

$$\alpha = \left(\begin{array}{c|c} A_i & C_j \\ \hline B_i & D_k \end{array} \right)_{i \in I, j \in J, k \in K}$$

to indicate that α has transversal blocks $A_i \cup B'_i$ ($i \in I$), and nontransversal blocks C_j ($j \in J$) and D'_k ($k \in K$). Note that one or more (but not all) of I, J, K may be empty, and that $|I| = \text{rank}(\alpha)$. We will often use variations of this notation but it should always be clear what is meant. The proof of the following lemma from [46] is straightforward.

Lemma 7.3. *A partition is a projection if and only if it is of the form*

$$\left(\begin{array}{c|c} A_i & C_j \\ \hline A_i & C_j \end{array} \right)_{i \in I, j \in J}.$$

For $0 \leq r \leq n$, let

$$J_r(\mathcal{P}_n) = \{\alpha \in \mathcal{P}_n : \text{rank}(\alpha) = r\}.$$

By Theorem 7.1, these sets are precisely the \mathcal{J} -classes of \mathcal{P}_n , and they form a chain:

$$J_0(\mathcal{P}_n) < J_1(\mathcal{P}_n) < \cdots < J_{n-1}(\mathcal{P}_n) < J_n(\mathcal{P}_n).$$

It follows from Proposition 5.11 that the ideals of \mathcal{P}_n are precisely the sets

$$I_r(\mathcal{P}_n) = J_0(\mathcal{P}_n) \cup J_1(\mathcal{P}_n) \cup \cdots \cup J_r(\mathcal{P}_n) = \{\alpha \in \mathcal{P}_n : \text{rank}(\alpha) \leq r\}.$$

Note that $I_n(\mathcal{P}_n) = \mathcal{P}_n$, $J_n(\mathcal{P}_n) = \mathcal{S}_n$, $I_{n-1}(\mathcal{P}_n) = \mathcal{P}_n \setminus \mathcal{S}_n$, and that the maximal subgroups contained in $J_r(\mathcal{P}_n)$ are all isomorphic to \mathcal{S}_r . In what follows, we will apply the general results of Sections 4 and 5 to the (finite idempotent generated regular $*$ -) semigroup $S = I_{n-1}(\mathcal{P}_n) = \mathcal{P}_n \setminus \mathcal{S}_n$.

7.1. Rank and idempotent rank of ideals in \mathcal{P}_n . The key lemma that allows us to reduce the problem to the consideration of principal factors is the following, which shows how elements of large rank may be used to generate elements of smaller rank.

Lemma 7.4. *If $0 \leq r \leq n - 2$, then $J_r(\mathcal{P}_n) \subseteq \langle J_{r+1}(\mathcal{P}_n) \rangle$.*

Proof. Let $\alpha \in J_r(\mathcal{P}_n)$ be a projection where $0 \leq r \leq n - 2$. By (the proof of) Proposition 5.11, it is enough to show that $\alpha \in \langle J_{r+1}(\mathcal{P}_n) \rangle$. Write

$$\alpha = \left(\begin{array}{c|c} A_i & C_j \\ \hline A_i & C_j \end{array} \right)_{i \in I, j \in J},$$

where $|I| = r$ and $|J| = k$. Without loss of generality, we may suppose that $I = [r]$ and $J = [k]$. There are two cases to consider.

Case 1: First suppose $|J| = k \geq 1$. In this case $\alpha = \beta\gamma$ where

$$\beta = \left(\begin{array}{c|c|c} A_i & C_1 & C_j \\ i & r+1 & x \end{array} \right)_{i \in I, j \in [2, k], x \in [r+2, n]},$$

$$\gamma = \left(\begin{array}{c|c|c} i & n & x \\ A_i & C_1 & C_j \end{array} \right)_{i \in I, j \in [2, k], x \in [r+1, n-1]}$$

both belong to $J_{r+1}(\mathcal{P}_n)$.

Case 2: Next suppose $|J| = k = 0$. Without loss of generality, we may suppose that $|A_r| \geq 2$. Consider a non-trivial decomposition $A_r = A'_r \cup A''_r$ where $A'_r \cap A''_r = \emptyset$. Then $\alpha = \beta\gamma$ where

$$\beta = \left(\begin{array}{c|c|c|c} A_i & A'_r & A''_r & \emptyset \\ i & r & r+1, r+2 & x \end{array} \right)_{i \in [r-1], x \in [r+3, n]},$$

$$\gamma = \left(\begin{array}{c|c|c|c} i & r, r+1 & r+2 & x \\ A_i & A'_r & A''_r & \emptyset \end{array} \right)_{i \in [r-1], x \in [r+3, n]}$$

both belong to $J_{r+1}(\mathcal{P}_n)$. □

Theorem 7.5. *For $0 \leq r \leq n-1$, the ideal $I_r(\mathcal{P}_n)$ is idempotent generated, and*

$$\text{rank}(I_r(\mathcal{P}_n)) = \text{idrank}(I_r(\mathcal{P}_n)) = \sum_{j=r}^n S(n, j) \binom{j}{r} = \sum_{j=r}^n \binom{n}{j} S(j, r) B_{n-j}$$

where $S(j, r)$ denotes the Stirling number of the second kind, and B_k denotes the k th Bell number. Moreover, a subset $A \subseteq I_r(\mathcal{P}_n)$ of this cardinality is a generating set for $I_r(\mathcal{P}_n)$ if and only if the following three conditions hold:

- (1) $\text{rank}(\alpha) = r$ for all $\alpha \in A$;
- (2) for all $\alpha, \beta \in A$ with $\alpha \neq \beta$, either $\ker(\alpha) \neq \ker(\beta)$ or $\text{dom}(\alpha) \neq \text{dom}(\beta)$;
- (3) for all $\alpha, \beta \in A$ with $\alpha \neq \beta$, either $\text{coker}(\alpha) \neq \text{coker}(\beta)$ or $\text{codom}(\alpha) \neq \text{codom}(\beta)$.

Proof. It follows from Lemma 7.4 and Proposition 5.11 that $I_r(\mathcal{P}_n)$ is idempotent generated and $\text{rank}(I_r(\mathcal{P}_n)) = \text{idrank}(I_r(\mathcal{P}_n)) = \rho_{nr}$, where ρ_{nr} is the number of \mathcal{R} -classes in $J_r(\mathcal{P}_n)$. To specify an \mathcal{R} -class in $J_r(\mathcal{P}_n)$, we first choose j kernel classes for some $j \in [r, n]$. From these, we then choose r classes to be part of the transversal blocks. These choices may be made in $S(n, j)$ and $\binom{j}{r}$ ways, respectively. Multiplying these and summing over all $j \in [r, n]$ gives

$$\rho_{nr} = \sum_{j=r}^n S(n, j) \binom{j}{r}.$$

(Counting the \mathcal{R} -classes in another way shows that $\rho_{nr} = \sum_{j=r}^n \binom{n}{j} S(j, r) B_{n-j}$: here we first choose the domain, say of size j , in $\binom{n}{j}$ ways; we then choose the r kernel-classes contained in the domain in $S(j, r)$ ways; we choose the remaining kernel-classes in B_{n-j} ways; finally, we multiply these values, then sum over all $j \in [r, n]$.) Finally, a subset $A \subseteq I_r(\mathcal{P}_n)$ with $|A| = \rho_{nr}$ generates $I_r(\mathcal{P}_n) = \langle J_r(\mathcal{P}_n) \rangle$ if and only if it generates the principal factor $J_r(\mathcal{P}_n)^*$ which, by Theorem 5.4, occurs if and only if A is a transversal of the \mathcal{R} - and \mathcal{L} -classes of $J_r(\mathcal{P}_n)$. By Theorem 7.1, this is equivalent to saying that conditions (1), (2) and (3) hold. □

Remark 7.6. As expected, this theorem agrees with Theorem 7.2 in the particular case $r = n - 1$. Note also that $\text{rank}(I_0(\mathcal{P}_n)) = \text{idrank}(I_0(\mathcal{P}_n)) = B_n$, and that the identity $\rho_{n0} + \rho_{n1} = \rho_{n+1,0}$ is a consequence of the well-known recurrence $B_{n+1} = \sum_{i=0}^n \binom{n}{i} B_i$. See Table 3 for some computed values of $\text{rank}(I_r(\mathcal{P}_n)) = \text{idrank}(I_r(\mathcal{P}_n))$. These numbers do not appear on [1].

Remark 7.7. We note that the formula given in the proof of Theorem 7.5 for ρ_{nr} , the number of \mathcal{R} -classes (and \mathcal{L} -classes) contained in $J_r(\mathcal{P}_n)$, is valid for $r = n$, giving a value of $\rho_{nn} = 1$ (but not agreeing with the value $\text{rank}(I_n(\mathcal{P}_n)) = \text{rank}(\mathcal{P}_n) = 4$, as calculated in [37]). Since also any \mathcal{H} -class contained in $J_r(\mathcal{P}_n)$ has size $r!$, it follows that $|J_r(\mathcal{P}_n)| = \rho_{nr}^2 r!$. This yields a formula for the cardinality of the ideals of \mathcal{P}_n :

$$|I_r(\mathcal{P}_n)| = \sum_{i=0}^r \rho_{ni}^2 i! \quad \text{for any } 0 \leq r \leq n.$$

$n \setminus r$	0	1	2	3	4	5	6	7	8	9
1	1									
2	2	3								
3	5	10	6							
4	15	37	31	10						
5	52	151	160	75	15					
6	203	674	856	520	155	21				
7	877	3263	4802	3556	1400	287	28			
8	4140	17007	28337	24626	11991	3290	490	36		
9	21147	94828	175896	174805	101031	34671	6972	786	45	
10	115975	562595	1146931	1279240	853315	350889	88977	13620	1200	55

TABLE 3. Values of $\text{rank}(I_r(\mathcal{P}_n)) = \text{idrank}(I_r(\mathcal{P}_n))$.

7.2. Minimal idempotent generating sets of $\mathcal{P}_n \setminus \mathcal{S}_n$. Theorem 5.7 above gives a correspondence between minimal idempotent generating sets of

$$\mathcal{P}_n \setminus \mathcal{S}_n = I_{n-1}(\mathcal{P}_n) = \langle J_{n-1}(\mathcal{P}_n) \rangle$$

and balanced subgraphs of the projection graph $\Gamma(\mathcal{P}_n \setminus \mathcal{S}_n) = \Gamma(J_{n-1}(\mathcal{P}_n)^*)$, in the sense of Definitions 5.5 and 5.6, which, for simplicity, we will denote by Γ_n . We will also write \mathcal{G}_n for the set of all balanced subgraphs of Γ_n . Parts of the next lemma were also used in [41].

Lemma 7.8. *The set of idempotents of $J_{n-1}(\mathcal{P}_n)$ is*

$$\{\pi_i : 1 \leq i \leq n\} \cup \{\pi_{ij}, \lambda_{ij}, \lambda_{ji}, \rho_{ij}, \rho_{ji} : 1 \leq i < j \leq n\},$$

where these partitions are illustrated in Figures 2 and 3. The set of projections of $J_{n-1}(\mathcal{P}_n)$ is

$$\{\pi_i : 1 \leq i \leq n\} \cup \{\pi_{ij} : 1 \leq i < j \leq n\}.$$

In the principal factor $J_{n-1}(\mathcal{P}_n)^*$, the only nonzero products of pairs of projections are

$$\pi_{ij}^2 = \pi_{ij}, \quad \pi_i^2 = \pi_i, \quad \pi_{ij}\pi_j = \lambda_{ij}, \quad \pi_{ij}\pi_i = \lambda_{ji}, \quad \pi_i\pi_{ij} = \rho_{ij}, \quad \pi_j\pi_{ij} = \rho_{ji}.$$

Proof. The statement about projections follows quickly from Lemma 7.3. By Lemma 5.3, any idempotent is the product of two projections, and it is easy to check that the products of the stated projections give only the stated idempotents as well as lower rank ones. \square

Remark 7.9. So there are $n + 5\binom{n}{2} = (5n^2 - 3n)/2$ idempotents in $J_{n-1}(\mathcal{P}_n)$. The idempotents in an arbitrary \mathcal{J} -class $J_r(\mathcal{P}_n)$ are enumerated in [27].





FIGURE 3. The non-projection idempotents of $J_{n-1}(\mathcal{P}_n)$; see also Figure 2.

In light of Lemma 7.8, we see that the projection graph Γ_n is obtained from the complete graph on vertex set $[n]$ by replacing each edge



For convenience, we have labeled the vertices i and ij instead of π_i and π_{ij} . The graph Γ_5 is pictured in Figure 4, with the same labelling convention, and with loops omitted.




FIGURE 4. The graph $\Gamma_5 = \Gamma(\mathcal{P}_5 \setminus \mathcal{S}_5)$ with loops omitted.

The edges of Γ_n correspond to the idempotents of $J_{n-1}(\mathcal{P}_n)$ via:

$$\begin{aligned} i \rightarrow i &\equiv \pi_i, & ij \rightarrow ij &\equiv \pi_{ij}, \\ ij \rightarrow j &\equiv \lambda_{ij}, & ij \rightarrow i &\equiv \lambda_{ji}, \\ i \rightarrow ij &\equiv \rho_{ij}, & j \rightarrow ij &\equiv \rho_{ji}. \end{aligned}$$

As mentioned above, Theorem 5.7 shows that the minimal idempotent generating sets of $\mathcal{P}_n \setminus \mathcal{S}_n = I_{n-1}(\mathcal{P}_n)$ correspond to the balanced subgraphs of Γ_n . The generating set from Theorem 7.2, which consists of all projections from $J_{n-1}(\mathcal{P}_n)$, corresponds to the balanced subgraph whose edges are the $n + \binom{n}{2} = \binom{n+1}{2}$ loops of Γ_n .

We now turn to the task of enumerating \mathcal{G}_n , the set of all balanced subgraphs of Γ_n . In what follows, for convenience, we will use symmetric notation for the vertices ij , and use ij and ji interchangeably without intending to imply $i < j$ or $i > j$.

Let $G \in \mathcal{G}_n$. The in-degree/out-degree condition is equivalent to saying that G is a disjoint union of circuits. By inspecting Γ_n , we see that the circuits of G must be of one of the following four types:

- (1) $i_1 \rightarrow i_1 i_2 \rightarrow i_2 \rightarrow i_2 i_3 \rightarrow i_3 \rightarrow \cdots \rightarrow i_k \rightarrow i_k i_1 \rightarrow i_1$ where $k \geq 3$ and i_1, i_2, \dots, i_k are distinct,
- (2) $ij \rightarrow ij$,
- (3) $i \rightarrow i$,
- (4) $i \rightarrow ij \rightarrow i$.




FIGURE 5. Two balanced subgraphs in Γ_3 , both with $\mu_G = (1, 2, 3)$.

Note that if n is large, then most connected components of G will be loops of type (2). Also note that G is completely determined by its circuits of type (1), (3) and (4). Our goal is to show that G determines (and is determined by) a pair (π_G, τ_G) where π_G is a permutation of a subset $A_G \subseteq [n]$ that has no fixed points and no 2-cycles, and τ_G is a function $[n] \setminus A_G \rightarrow [n]$ that has no 2-cycles. Here we say that a function $\phi : Y \rightarrow X$ with $Y \subseteq X$ has a 2-cycle if there exists $x, y \in Y$ with $x \neq y$, $x\phi = y$ and $y\phi = x$. With this goal in mind, we define

$$A_G = \{i \in [n] : \text{vertex } i \text{ is contained in a circuit of type (1)}\}.$$

Note that $|A_G| \in \{0\} \cup [3, n]$. We define $\pi_G : A_G \rightarrow A_G$ to be the permutation whose cycle decomposition includes a cycle (i_1, i_2, \dots, i_k) corresponding to each circuit of G of type (1). Note that π_G has no 2-cycles or fixed points. Note also that if $A_G = \emptyset$, then π_G is the unique function $\emptyset \rightarrow \emptyset$. We also define $\tau_G : [n] \setminus A_G \rightarrow [n]$ by

$$i\tau_G = \begin{cases} i & \text{if } G \text{ contains the loop } i \rightarrow i \\ j & \text{if } G \text{ contains the circuit } i \rightarrow ij \rightarrow i. \end{cases}$$

Note that τ_G contains no 2-cycles, but might have fixed points. So G uniquely determines the pair (π_G, τ_G) . Conversely, a pair (π, τ) for which

- (B1) π is a permutation, of some subset $A \subseteq [n]$, with no fixed points and 2-cycles, and
- (B2) $\tau : [n] \setminus A \rightarrow [n]$ has no 2-cycles

determines a balanced subgraph of Γ_n in such a way that gives a bijective correspondence between $G \in \mathcal{G}_n$ and pairs (π, τ) satisfying (B1) and (B2). So it suffices to enumerate such pairs.

Remark 7.10. The functions π_G and τ_G could be combined to give a transformation $\mu_G : [n] \rightarrow [n]$ defined by

$$i\mu_G = \begin{cases} i\pi_G & \text{if } i \in A_G \\ i\tau_G & \text{if } i \in [n] \setminus A_G. \end{cases}$$

However, G is not uniquely determined by μ_G . For example, in Γ_3 , the two balanced subgraphs shown in Figure 5 both have $\mu_G = (1, 2, 3)$.

For $0 \leq k \leq n$, let

$$A_k = \{\pi \in \mathcal{S}_k : \pi \text{ has no fixed points or 2-cycles}\}$$

$$B_{nk} = \{\tau : [k] \rightarrow [n] : \tau \text{ has no 2-cycles}\},$$

and put

$$a_k = |A_k| \quad \text{and} \quad b_{nk} = |B_{nk}|.$$

Note that for any subset $X \subseteq [n]$ with $|X| = k$, the set

$$\{\tau : X \rightarrow [n] : \tau \text{ has no 2-cycles}\}$$

has cardinality b_{nk} . If $A \subseteq [n]$ with $|A| = k$, the number of pairs (π, τ) satisfying conditions (B1) and (B2) above is equal to $a_k b_{n, n-k}$. It follows that

$$|\mathcal{G}_n| = \sum_{k=0}^n \binom{n}{k} a_k b_{n, n-k},$$

so it remains to evaluate the sequences a_k and b_{nk} . The sequence a_k is well-known; it is A038205 on [1], but we prove the next result for completeness.

Lemma 7.11. *The sequence a_k satisfies the recurrence*

$$a_0 = 1, \quad a_1 = a_2 = 0, \quad a_{k+1} = ka_k + k(k-1)a_{k-2} \quad \text{for } k \geq 2.$$

Proof. The values for a_0, a_1, a_2 are clear. Now consider an element $\pi \in A_{k+1}$ where $k \geq 2$. There are two possibilities: either (i) $k+1$ is in an l -cycle of π where $l \geq 4$, or (ii) $k+1$ is in a 3-cycle of π . It is easy to see that there are ka_k elements of type (i), and $k(k-1)a_{k-2}$ of type (ii). \square

The first few values of a_k are given in Table 4.

k	0	1	2	3	4	5	6	7	8	9	10
a_k	1	0	0	2	6	24	160	1140	8988	80864	809856

TABLE 4. The sequence a_k .

Lemma 7.12. *For any $0 \leq k \leq n$, we have*

$$b_{nk} = \sum_{i=0}^{\lfloor \frac{k}{2} \rfloor} (-1)^i \binom{k}{2i} (2i-1)!! n^{k-2i},$$

where $(2i-1)!! = (2i-1)(2i-3)\cdots 3 \cdot 1$ and we interpret $(-1)!! = 1$.

Proof. For $1 \leq r < s \leq k$, let

$$C_{nk}(r, s) = \{\tau : [k] \rightarrow [n] : (r, s) \text{ is a 2-cycle of } \tau\}.$$

Then, since there are n^k functions $[k] \rightarrow [n]$,

$$b_{nk} = n^k - \left| \bigcup_{1 \leq r < s \leq k} C_{nk}(r, s) \right|. \quad (7.13)$$

Now, if $1 \leq i \leq \lfloor \frac{k}{2} \rfloor$ and $(r_1, s_1), \dots, (r_i, s_i)$ are disjoint cycles, then

$$|C_{nk}(r_1, s_1) \cap \cdots \cap C_{nk}(r_i, s_i)| = n^{k-2i},$$

and there are $\binom{k}{2i} (2i-1)!!$ ways to choose i disjoint cycles from $[k]$. So the inclusion-exclusion formula gives

$$\left| \bigcup_{1 \leq r < s \leq k} C_{nk}(r, s) \right| = \sum_{i=1}^{\lfloor \frac{k}{2} \rfloor} (-1)^{i+1} \binom{k}{2i} (2i-1)!! n^{k-2i}.$$

The result now follows from (7.13), since n^k is the $i = 0$ term of the sum in the statement of the lemma. \square

The numbers b_{nk} do not appear in [1]. The first few values are included in Table 5. We have proved the following.

$n \setminus k$	0	1	2	3	4	5	6	7	8	9	10
0	1										
1	1	1									
2	1	2	3								
3	1	3	8	18							
4	1	4	15	52	163						
5	1	5	24	110	478	1950					
6	1	6	35	198	1083	5706	28821				
7	1	7	48	322	2110	13482	83824	505876			
8	1	8	63	488	3715	27768	203569	1461944	10270569		
9	1	9	80	702	6078	51894	436656	3618540	29510268	236644092	
10	1	10	99	970	9403	90150	854485	8003950	74058105	676549450	6098971555

TABLE 5. The numbers b_{nk} .

Theorem 7.14. *The number of minimal idempotent generating sets for the singular part $\mathcal{P}_n \setminus \mathcal{S}_n$ of the partition monoid \mathcal{P}_n is equal to*

$$|\mathcal{G}_n| = \sum_{k=0}^n \binom{n}{k} a_k b_{n-k},$$

where formulae for the numbers a_k and b_{nk} are given in Lemmas 7.11 and 7.12.

The first few values of $|\mathcal{G}_n|$ are given in Table 6; this sequence also does not appear in [1].

n	0	1	2	3	4	5	6	7	8	9	10
$ \mathcal{G}_n $	1	1	3	20	201	2604	40915	754368	15960945	381141008	10139372451

TABLE 6. The numbers $|\mathcal{G}_n|$, which give the number of minimal idempotent generating sets for $\mathcal{P}_n \setminus \mathcal{S}_n$.

7.3. Arbitrary idempotent generating sets for $\mathcal{P}_n \setminus \mathcal{S}_n$. Given a set F consisting of idempotents from $J_{n-1}(\mathcal{P}_n)$, we would like to know whether F is a generating set of $\mathcal{P}_n \setminus \mathcal{S}_n$. For a subset F of

$$\{\alpha \in E(\mathcal{P}_n) : \text{rank}(\alpha) = n - 1\} = \{\pi_i : 1 \leq i \leq n\} \cup \{\pi_{ij}, \lambda_{ij}, \lambda_{ji}, \rho_{ij}, \rho_{ji} : 1 \leq i < j \leq n\},$$

let $\Gamma_n(F)$ be the two-coloured digraph obtained by colouring each edge of Γ_n blue, and then adding red edges corresponding to the idempotents from F :

- $i \rightarrow i$ if $\pi_i \in F$,
- $ij \rightarrow j$ if $\lambda_{ij} \in F$,
- $i \rightarrow ij$ if $\rho_{ij} \in F$,
- $ij \rightarrow ij$ if $\pi_{ij} \in F$,
- $ij \rightarrow i$ if $\lambda_{ji} \in F$,
- $j \rightarrow ij$ if $\rho_{ji} \in F$.

(As above, we will denote the vertices of Γ_n by i and ij rather than π_i and π_{ij} .) Applying the general result Theorem 6.5 we obtain the following.

Theorem 7.15. *For $F \subseteq \{\alpha \in E(\mathcal{P}_n) : \text{rank}(\alpha) = n - 1\}$, the following are equivalent:*

- (i) $\mathcal{P}_n \setminus \mathcal{S}_n = \langle F \rangle$;
- (ii) each vertex of $\Gamma_n(F)$ is the base point of an RBR-alternating circuit.

We currently do not know of any simpler necessary and sufficient condition for such a subset F of idempotents to be a generating set. It would be desirable to give such a condition in terms of only the red edges of the graph $\Gamma_n(F)$. As mentioned in Remark 6.8, one might be tempted to conjecture that $\langle F \rangle = \mathcal{P}_n \setminus \mathcal{S}_n$ if and only if, in $\Gamma_n(F)$, either:




FIGURE 6. The digraph $\Gamma_2(F)$ where $F = \{\pi_1, \pi_2, \lambda_{12}, \rho_{12}\}$ in the partition monoid \mathcal{P}_2 .

- (i) each vertex has at least one red edge coming in to it and at least one going out of it, or
- (ii) each vertex is contained in a red circuit.

As we also mentioned in Remark 6.8, Theorem 6.5 tells us that condition (i) is necessary, while condition (ii) is sufficient. But the following two examples show that neither condition is necessary *and* sufficient.

Example 7.16. Consider the set of idempotents $F = \{\pi_1, \pi_2, \lambda_{12}, \rho_{12}\}$ in the partition monoid \mathcal{P}_2 . The digraph $\Gamma_2(F)$ is illustrated in Figure 6. Clearly, there is no *RBR*-alternating circuit based at the vertex $\{12\}$. It follows that $\pi_{12} \notin \langle F \rangle$, and so F does not generate $\mathcal{P}_2 \setminus \mathcal{S}_2$. This example shows that (i) is not a sufficient condition.

Example 7.17. Consider the set of idempotents $F = \{\pi_2, \pi_3, \pi_{12}, \pi_{23}, \pi_{13}, \lambda_{31}, \rho_{12}\}$ in the partition monoid \mathcal{P}_3 . The digraph $\Gamma_3(F)$ is illustrated in Figure 7. Every vertex has a red loop (the simplest kind of *RBR*-circuit) with the exception of vertex 1. Vertex 1 is the basepoint of the *RBR*-alternating circuit:

$$1 \rightarrow 12 \rightarrow 2 \rightarrow 2 \rightarrow 23 \rightarrow 23 \rightarrow 3 \rightarrow 3 \rightarrow 13 \rightarrow 1.$$

This path corresponds to the product of idempotents $\rho_{12}\pi_2\pi_{23}\pi_3\lambda_{31}$, which is \mathcal{H} -related to the projection π_1 ; in fact, $\pi_1 = (\rho_{12}\pi_2\pi_{23}\pi_3\lambda_{31})^2$. It follows from Theorem 7.15 that $\mathcal{P}_3 \setminus \mathcal{S}_3 = \langle F \rangle$. This example shows that (ii) is not a necessary condition. It is also easy to check that for any $f \in F$, the graph $\Gamma_n(F \setminus \{f\})$ does not satisfy condition (i). It follows that F is an irreducible generating set, even though it is not of the minimal size $\binom{4}{2} = 6$. This contrasts to the situation for the singular part $\mathcal{T}_n \setminus \mathcal{S}_n$ of the full transformation semigroup \mathcal{T}_n , where every idempotent generating set contains an idempotent generating set of minimal size [23]. (The previous statement does not hold if “idempotent generating set” is replaced by “generating set”.)

We leave it as an open problem to determine a necessary and sufficient condition for F to be a generating set, stated in terms of the structure of the subgraph of $\Gamma_n(F)$ determined by F (that is, the subgraph determined by the red edges).

8. THE BRAUER MONOID

Recall that the Brauer monoid \mathcal{B}_n is the subsemigroup of \mathcal{P}_n consisting of all partitions whose blocks have cardinality 2. See Figure 8 for an example. Note that an element

$$\left(\begin{array}{c|c} A_i & C_j \\ \hline B_i & D_k \end{array} \right)_{i \in I, j \in J, k \in K}$$

of \mathcal{B}_n satisfies $|A_i| = |B_i| = 1$ and $|C_j| = |D_k| = 2$ for all i, j, k . In addition we must have $|J| = |K|$, and $|I| = n - 2|J|$, which means that the ranks of elements of \mathcal{B}_n are restricted to natural numbers of the form $n - 2k$ where k is a natural number. Note also that $|\mathcal{B}_n| = (2n - 1)!! = (2n - 1)(2n - 3) \cdots 3 \cdot 1$.

Since the symmetric group \mathcal{S}_n is clearly contained in \mathcal{B}_n , it follows that \mathcal{S}_n is the group of units of \mathcal{B}_n . And, since \mathcal{B}_n is closed under the $*$ operation, it is regular, so Green’s




FIGURE 7. The digraph $\Gamma_3(F)$ where $F = \{\pi_2, \pi_3, \pi_{12}, \pi_{23}, \pi_{13}, \lambda_{31}, \rho_{12}\}$ in the partition monoid \mathcal{P}_3 .




FIGURE 8. An element of the Brauer monoid \mathcal{B}_7 .

relations on \mathcal{B}_n are still given by the formulae in Theorem 7.1. In fact, since the domain (respectively, codomain) of an element of \mathcal{B}_n is determined by its kernel (respectively, cokernel), we have the following. See also [114, Theorem 7] where an equivalent characterization is given.

Theorem 8.1. *For each $\alpha, \beta \in \mathcal{B}_n$, we have:*

- (i) $\alpha \mathcal{R} \beta$ if and only if $\ker(\alpha) = \ker(\beta)$;
- (ii) $\alpha \mathcal{L} \beta$ if and only if $\text{coker}(\alpha) = \text{coker}(\beta)$;
- (iii) $\alpha \mathcal{J} \beta$ if and only if $\text{rank}(\alpha) = \text{rank}(\beta)$.

For $1 \leq i < j \leq n$, we define the partition

$$\tau_{ij} = \left(\begin{array}{c|c} x & i, j \\ x & i, j \end{array} \right)_{x \in [n] \setminus \{i, j\}}$$

See Figure 9 for an illustration. The following result was proved in [100].




FIGURE 9. The projection $\tau_{ij} \in \mathcal{B}_n$.

Theorem 8.2. *The singular part $\mathcal{B}_n \setminus \mathcal{S}_n$ of the Brauer monoid \mathcal{B}_n is idempotent generated. The set $\{\tau_{ij} : 1 \leq i < j \leq n\}$ is a minimal idempotent generating set.*

In [100], the authors go on to give a presentation for $\mathcal{B}_n \setminus \mathcal{S}_n$ with respect to the above generating set of projections, but we will not require this presentation here.

As before, for $r = n - 2k$ with $k \geq 0$ and $0 \leq r \leq n$, we write

$$J_r(\mathcal{B}_n) = \{\alpha \in \mathcal{B}_n : \text{rank}(\alpha) = r\} = J_r(\mathcal{P}_n) \cap \mathcal{B}_n.$$

By Theorem 7.1, these are precisely the \mathcal{J} -classes of \mathcal{B}_n , and they form a chain:

$$J_m(\mathcal{B}_n) < J_{m+2}(\mathcal{B}_n) < \cdots < J_{n-2}(\mathcal{B}_n) < J_n(\mathcal{B}_n),$$

where m denotes 0 if n is even, or 1 otherwise. By Proposition 5.11, it follows that the ideals of \mathcal{B}_n are precisely the sets

$$I_r(\mathcal{B}_n) = J_m(\mathcal{B}_n) \cup J_{m+2}(\mathcal{B}_n) \cup \cdots \cup J_r(\mathcal{B}_n) = \{\alpha \in \mathcal{B}_n : \text{rank}(\alpha) \leq r\} = I_r(\mathcal{P}_n) \cap \mathcal{B}_n.$$

Note that $I_n(\mathcal{B}_n) = \mathcal{B}_n$, $J_n(\mathcal{B}_n) = \mathcal{S}_n$, $I_{n-2}(\mathcal{B}_n) = \mathcal{B}_n \setminus \mathcal{S}_n$, and that the maximal subgroups contained in $J_r(\mathcal{B}_n)$ are all isomorphic to \mathcal{S}_r . In what follows, we will apply the general results of Sections 4 and 5 to the (finite idempotent generated regular $*$ -) semigroup $S = I_{n-2}(\mathcal{B}_n) = \mathcal{B}_n \setminus \mathcal{S}_n$.

8.1. Rank and idempotent rank of ideals of \mathcal{B}_n . Again, the key step is to show that elements of small rank in \mathcal{B}_n may be expressed as a product of higher rank elements.

Lemma 8.3. *If $0 \leq r \leq n - 4$, then $J_r(\mathcal{B}_n) \subseteq \langle J_{r+2}(\mathcal{B}_n) \rangle$.*

Proof. Let $\alpha \in J_r(\mathcal{B}_n)$ be a projection. By Proposition 5.11, it suffices to show that $\alpha \in \langle J_{r+2}(\mathcal{B}_n) \rangle$. Write

$$\alpha = \left(\begin{array}{c|c} A_i & C_j \\ \hline A_i & C_j \end{array} \right)_{i \in I, j \in J}.$$

Without loss of generality, we may assume that $J = [k]$, where $r = n - 2k$. For $j \in J$, let $C_j = \{a_j, b_j\}$, where $a_j < b_j$. So $\alpha = \tau_{a_1 b_1} \cdots \tau_{a_k b_k}$. Then $\alpha = \beta\gamma$, where $\beta = \tau_{a_1 b_1} \cdots \tau_{a_{k-1} b_{k-1}}$ and $\gamma = \tau_{a_2 b_2} \cdots \tau_{a_k b_k}$ both belong to $J_{r+2}(\mathcal{B}_n)$. \square

Theorem 8.4. *For $0 \leq r = n - 2k \leq n - 2$, the ideal $I_r(\mathcal{B}_n)$ is idempotent generated, and*

$$\text{rank}(I_r(\mathcal{B}_n)) = \text{idrank}(I_r(\mathcal{B}_n)) = \binom{n}{2k} (2k - 1)!! = \frac{n!}{2^k k! r!}.$$

Moreover, a subset $A \subseteq I_r(\mathcal{B}_n)$ of this cardinality is a generating set for $I_r(\mathcal{B}_n)$ if and only if the following three conditions hold:

- (1) $\text{rank}(\alpha) = r$ for all $\alpha \in A$;
- (2) for all $\alpha, \beta \in A$ with $\alpha \neq \beta$, $\ker(\alpha) \neq \ker(\beta)$;
- (3) for all $\alpha, \beta \in A$ with $\alpha \neq \beta$, $\text{coker}(\alpha) \neq \text{coker}(\beta)$.

Proof. It follows from Lemma 8.3 and Proposition 5.11 that $I_r(\mathcal{B}_n)$ is idempotent generated, and that $\text{rank}(I_r(\mathcal{B}_n)) = \text{idrank}(I_r(\mathcal{B}_n)) = \rho_{nr}$, where ρ_{nr} is the number of \mathcal{R} -classes in $J_r(\mathcal{B}_n)$. It follows from Theorem 8.1 that ρ_{nr} is equal to the number of equivalence relations on $[n]$ that have exactly k non-trivial blocks, each of size 2. There are $\binom{n}{2k}$ ways to choose the elements belonging to the non-trivial blocks, and then $(2k - 1)!!$ ways to choose the blocks of size 2; multiplying these together gives the required formula. The final clause follows by applying Theorem 5.4 in a virtually identical way to the argument in the final paragraph of the proof of Theorem 7.5. \square

Remark 8.5. Note that the $r = n - 2$ case agrees with Theorem 8.2. Note also that $\text{rank}(I_0(\mathcal{B}_n)) = (n - 1)!!$ if n is even, while $\text{rank}(I_1(\mathcal{B}_n)) = n!!$ if n is odd. Some calculated values of $\text{rank}(I_r(\mathcal{B}_n)) = \text{idrank}(I_r(\mathcal{B}_n))$ are given in Table 7.

Remark 8.6. As in Remark 7.7, we may derive formulae for the sizes of the \mathcal{J} -classes and ideals of \mathcal{B}_n :

$$|J_r(\mathcal{B}_n)| = \rho_{nr}^2 r! \quad \text{and} \quad |I_r(\mathcal{B}_n)| = \sum_{i=0}^r \rho_{ni}^2 i! \quad \text{for any } 0 \leq r \leq n,$$

where the numbers $\rho_{nr} = \text{rank}(I_r(\mathcal{B}_n))$ are as in Theorem 8.4, and for convenience we define $\rho_{nn} = 1$ and $\rho_{ni} = 0$ if $i \not\equiv n \pmod{2}$.

$n \setminus r$	0	1	2	3	4	5	6	7	8
2	1								
3		3							
4	3		6						
5		15		10					
6	15		45		15				
7		105		105		21			
8	105		420		210		28		
9		945		1260		378		36	
10	945		4725		3150		630		45

TABLE 7. Values of $\text{rank}(I_r(\mathcal{B}_n)) = \text{idrank}(I_r(\mathcal{B}_n))$.

8.2. Minimal idempotent generating sets of $\mathcal{B}_n \setminus \mathcal{S}_n$. As in Section 7.2, an enumeration of the minimal idempotent generating sets of $\mathcal{B}_n \setminus \mathcal{S}_n$ amounts to an enumeration of the balanced subgraphs of the projection graph $\Gamma(\mathcal{B}_n \setminus \mathcal{S}_n) = \Gamma(J_{n-2}(\mathcal{B}_n)^*)$, which we will denote by Λ_n .

For distinct $i, j, k \in [n]$, we define

$$\sigma_{ijk} = \left(x \mid \begin{array}{c|c} k & i, j \\ \hline x & i \\ \hline & j, k \end{array} \right)_{x \in [n] \setminus \{i, j, k\}}.$$

See Figure 10 for an illustration where, for convenience, we have only pictured the restriction of σ_{ijk} to $\{i, j, k\}$. It is easy to check that these partitions are idempotents.




FIGURE 10. Simplified illustrations of the idempotents $\sigma_{ijk} \in \mathcal{B}_n$ for all possible orderings of i, j, k . See text for more details.

The next lemma is verified in similar fashion to Lemma 7.8, and its proof is omitted. For simplicity, we will use symmetric notation and allow ourselves to write $\tau_{ji} = \tau_{ij}$ for all $1 \leq i < j \leq n$.

Lemma 8.7. *The set of idempotents of $J_{n-2}(\mathcal{B}_n)$ is*

$$\{\tau_{ij} : 1 \leq i < j \leq n\} \cup \{\sigma_{ijk} : i, j, k \in [n] \text{ distinct}\}.$$

The set of projections of $J_{n-2}(\mathcal{B}_n)$ is

$$\{\tau_{ij} : 1 \leq i < j \leq n\}.$$

In the principal factor $J_{n-2}(\mathcal{B}_n)^$, the only nonzero products of pairs of projections are, using symmetric notation for the projections,*

$$\tau_{ij}^2 = \tau_{ij}, \quad \tau_{ij}\tau_{jk} = \sigma_{ijk}.$$

It follows that the graph Λ_n has vertex set $\{\tau_{ij} : 1 \leq i < j \leq n\}$ with edges $\tau_{ij} \rightarrow \tau_{kl}$ if and only if $\{i, j\} \cap \{k, l\} \neq \emptyset$. The graphs Λ_4 and Λ_5 are pictured in Figure 11, where we have simplified matters by labelling the vertices ij instead of τ_{ij} , omitting loops, and displaying pairs of directed edges $ij \rightleftharpoons kl$ as single undirected edges $ij - kl$.




FIGURE 11. Simplified illustrations of the graphs $\Lambda_4 = \Gamma(\mathcal{B}_4 \setminus \mathcal{S}_4)$ and $\Lambda_5 = \Gamma(\mathcal{B}_5 \setminus \mathcal{S}_5)$; see text for further details.

Recall that the *Johnson graph* $J(n, k)$ is the graph with vertex set $\{A \subseteq [n] : |A| = k\}$, and with edges $A - B$ if and only if $|A \cap B| = k - 1$. In particular $J(n, 2)$ has vertex set $\{A \subseteq [n] : |A| = 2\}$ and edges $A - B$ if and only if A and B overlap in precisely one element. More background on Johnson graphs in the context of algebraic graph theory may be found in [56, Chapter 1.6]. Note that the underlying undirected loop-free graph of $\Lambda_n = \Gamma(\mathcal{B}_n \setminus \mathcal{S}_n)$ is isomorphic to $J(n, 2)$. So, in fact, Figure 11 pictures the Johnson graphs $J(4, 2)$ and $J(5, 2)$.

Factorizations of Johnson graphs have been considered, for instance, in [16]. Recall that a 1-factor of a graph is a collection of edges that spans the graph, while a 2-factor is a collection of cycles that spans all vertices of the graph. We define a $(0, 1, 2)$ -factor of a graph as a decomposition of the graph into vertices, edges, and cycles, such that each vertex is contained in precisely one of these pieces. An oriented $(0, 1, 2)$ -factor is a $(0, 1, 2)$ -factor such that all the cycles are assigned an orientation (clockwise or anticlockwise).

Note that there is a one-one correspondence between balanced subgraphs of Λ_n and oriented $(0, 1, 2)$ -factors of $J(n, 2)$. Indeed, consider a balanced subgraph H of Λ_n . The connected components of H come in three forms:

- (i) a component of size 1: $\{i, j\} \rightleftharpoons$;
- (ii) a component of size 2: $\{i, j\} \rightleftharpoons \{j, k\}$; or

(iii) a component of size $k \geq 3$: $\{i_1, i_2\} \rightarrow \{i_2, i_3\} \rightarrow \cdots \rightarrow \{i_{k-1}, i_k\} \rightarrow \{i_1, i_2\}$.

These three types of components give rise (respectively) to the vertices, edges and oriented cycles in a corresponding $(0, 1, 2)$ -factor of $J(n, 2)$. Combined with Theorem 5.7, which tells us that the minimal idempotent generating sets of $\mathcal{B}_n \setminus \mathcal{S}_n$ are in one-one correspondence with the balanced subgraphs of $\Lambda_n = \Gamma(\mathcal{B}_n \setminus \mathcal{S}_n)$, this proves the following.

Proposition 8.8. *There is a one-one correspondence between the minimal idempotent generating sets of $\mathcal{B}_n \setminus \mathcal{S}_n$ and the oriented $(0, 1, 2)$ -factors of the Johnson graph $J(n, 2)$.*

Remark 8.9. We do not know of any formula or recurrence relation for the number d_n of minimal idempotent generating sets of $\mathcal{B}_n \setminus \mathcal{S}_n$. However, the sequence d_n grows rapidly, as we now explain. By Proposition 8.8, d_n is equal to the number of $(0, 1, 2)$ -factors of the Johnson graph $J(n, 2)$. Consider the graph $J(n+1, 2)$. We decompose the vertex set of $J(n+1, 2)$ into two subsets:

$$V_1 = \{\{i, n+1\} : 1 \leq i \leq n\} \quad \text{and} \quad V_2 = \{\{i, j\} : 1 \leq i < j \leq n\}.$$

Let G_1 and G_2 be the induced subgraphs of Λ_{n+1} with these vertex sets (respectively). Any oriented $(0, 1, 2)$ -factor of G_1 and any oriented $(0, 1, 2)$ -factor of G_2 may be pieced together to give an oriented $(0, 1, 2)$ -factor of $J(n+1, 2)$. But G_1 is the complete graph on vertex set V_1 , and G_2 is in fact the Johnson graph $J(n, 2)$. Since there are $n!$ oriented $(0, 1, 2)$ -factors of G_1 (corresponding to permutations of the vertex set), and d_n oriented $(0, 1, 2)$ -factors of G_2 , it follows that $d_{n+1} \geq n! \times d_n$. This then gives

$$d_n \geq \prod_{i=1}^{n-1} i! = \prod_{i=1}^{n-1} i^{n-i}.$$

Remark 8.10. The first few values of d_n were calculated by James Mitchell using the Semigroups package in GAP; see [45, 116]. These values are given in Table 8, along with the sequence $c_n = \prod_{i=1}^{n-1} i!$. The calculation of $d_3 = 6$ is trivial, the $(0, 1, 2)$ -factors of $J(3, 2)$ being in one-one correspondence with the permutations of a three element set. To see that $d_4 = 265$, consider the graph $J(4, 2)$, depicted as an octahedron in Figure 11. Labelling the vertices a, b, c, A, B, C in such a way that vertex x is opposite vertex X for each $x \in \{a, b, c\}$, we see that the $(0, 1, 2)$ -factors of $J(4, 2)$ are in one-one correspondence with the permutations of $\{a, b, c, A, B, C\}$ such that no lower-case letter is mapped to its corresponding upper-case letter, and vice versa. These permutations are obviously in one-one correspondence with the fixed point free permutations of the set $[6] = \{1, 2, 3, 4, 5, 6\}$, of which there are 265. In general, for $n \geq 4$, d_n is bounded above by the number e_n of fixed point free permutations of a set of size $\binom{n}{2}$; this sequence is also given in Table 8. The sequences c_n , d_n and e_n are A000178, A244493 and (a subsequence of) A000166, respectively, on [1]; the sequence d_n was added to [1] after this article was written.

n	2	3	4	5	6	7
c_n	1	2	12	288	34560	24883200
d_n	1	6	265	126140	855966411	?
e_n	0	2	265	1334961	481066515734	895014631192902121
d_n/c_n	1	3	≈ 22	≈ 438	≈ 24768	?
e_n/d_n	0	1/3	1	≈ 11	≈ 562	?

TABLE 8. The sequences c_n, d_n, e_n . For $n \geq 2$, d_n is equal to the number of minimal idempotent generating sets of $\mathcal{B}_n \setminus \mathcal{S}_n$.

The Brauer monoid and Pfaffian orientations. We have established a correspondence between minimal idempotent generating sets of the singular part of the Brauer monoid, and certain factorizations of the Johnson graph $J(n, 2)$. As we mentioned above, we do not know of a formula or recurrence relation that gives the number of such factorizations. The following result shows that one cannot find such a formula by trying to compute a Pfaffian orientation for the corresponding Graham–Houghton graph.

A subgraph H of a graph G is called *central* if $G \setminus H$ has a perfect matching (here, \setminus stands for deletion, where we remove the vertices from H as well as any edges involving one or more vertices from H). An even circuit C in a directed graph D is called *oddly oriented* if for either choice of direction of traversal around C , the number of edges of C directed in the direction of traversal is odd. This is independent of the initial choice of direction of traversal, since C is even. An orientation D of the edges of a graph G is *Pfaffian* if every even central circuit of G is oddly oriented in D . We say that a graph G is Pfaffian if it has a Pfaffian orientation. The significance of Pfaffian orientations comes from the fact that if a bipartite graph G has one, then the number of perfect matchings of G can be computed in polynomial time. More on Pfaffian orientations may be found in [99, 128, 132, 133].

These ideas are relevant to us, since the number of distinct minimal idempotent generating sets is precisely the number of perfect matchings of the (bipartite) Graham–Houghton graph. So, if the Graham–Houghton graph in question did have a Pfaffian orientation this would mean that the number of distinct minimal idempotent generating sets could be computed in polynomial time. The following result shows that we cannot use the theory of Pfaffian orientations as an approach to computing the number of minimal generating sets for $\mathcal{B}_n \setminus \mathcal{S}_n$.

Proposition 8.11. *Let Δ_n be the unlabeled undirected Graham–Houghton graph of the \mathcal{J} -class $J_{n-2}(\mathcal{B}_n)$. If $n \geq 3$, then Δ_n does not admit a Pfaffian orientation.*

Proof. When $n = 3$, Δ_3 is isomorphic to the complete bipartite graph $K_{3,3}$, which is known not to admit a Pfaffian orientation; see [132, Theorem 4.1]. Now we shall apply results from [132] to show that Δ_n does not admit a Pfaffian orientation for any $n \geq 4$.

A digraph D is called *even* if whenever the arcs of D are assigned weights 0 or 1, D contains a cycle of even total weight. Now let $G = A \sqcup B$ be a bipartite graph and M a matching of G . Let $D = D(G, M)$ be the digraph obtained from G by (i) orienting arcs from A to B , and (ii) contracting every edge of M . Little [99] showed that a bipartite graph G is Pfaffian if and only if $D(G, M)$ is *not* even.

Now, carrying out this process with the bipartite graph Δ_n together with the natural choice of matching given by the set of projections, we obtain a directed version of the Johnson graph $J(n, 2)$; the graph obtained is essentially the graph $\Lambda_n = \Gamma(\mathcal{B}_n \setminus \mathcal{S}_n)$, defined above, but with the loops at each vertex removed. We claim that the resulting digraph, which we will denote by Λ'_n , is even. Once established, from the results above, this will complete the proof that Δ_n is not Pfaffian. To see that Λ'_n is even, consider an assignment of labels $\{0, 1\}$ to the arcs of this digraph. Since Λ'_n embeds triangles it follows that at least one of the four configurations displayed in Figure 12 must arise. By inspection it is now clear that in each case Λ'_n must contain a cycle of even total weight. \square

8.3. Arbitrary idempotent generating sets for $\mathcal{B}_n \setminus \mathcal{S}_n$.

For a subset F of $\{\alpha \in E(\mathcal{B}_n) : \text{rank}(\alpha) = n - 2\} = \{\tau_{ij} : 1 \leq i < j \leq n\} \cup \{\sigma_{ijk} : i, j, k \in [n] \text{ distinct}\}$, let $\Lambda_n(F)$ be the two-coloured digraph obtained by colouring each edge of Λ_n blue, and then adding red edges corresponding to the idempotents from F :

- $ij \rightarrow ij$ if $\tau_{ij} \in F$,
- $ij \rightarrow jk$ if $\sigma_{ijk} \in F$.




FIGURE 12. Four configurations, one of which must arise in Λ'_n after assigning labels $\{0, 1\}$ to the arcs of the digraph Λ'_n . Here dotted arcs are arcs labeled by 0, and the solid represent those labeled by 1.

(As above, we will denote the vertices of Λ_n by ij rather than τ_{ij} .)

Theorem 8.12. *For $F \subseteq \{\alpha \in E(\mathcal{B}_n) : \text{rank}(\alpha) = n - 2\}$, the following are equivalent:*

- (i) $\mathcal{B}_n \setminus \mathcal{S}_n = \langle F \rangle$;
- (ii) *each vertex of $\Lambda_n(F)$ is the base point of an RBR-alternating circuit.*

The authors are not aware of any formula for the number of subsets

$$F \subseteq \{\alpha \in E(\mathcal{B}_n) : \text{rank}(\alpha) = n - 2\}$$

that generate $\mathcal{B}_n \setminus \mathcal{S}_n$.

Remark 8.13. Neither condition (i) nor condition (ii) of Remark 6.8 gives a necessary and sufficient condition for F to be a generating set in the case of $\mathcal{B}_n \setminus \mathcal{S}_n$.

Condition (i) of Remark 6.8, though necessary, is not sufficient. For example, consider the graph $\Lambda_4(F)$ where $\Lambda_4 = \Gamma(\mathcal{B}_4 \setminus \mathcal{S}_4)$ and

$$F = \{\tau_{14}, \tau_{23}, \sigma_{214}, \sigma_{231}, \sigma_{234}, \sigma_{241}, \sigma_{314}, \sigma_{321}, \sigma_{324}, \sigma_{341}\}.$$

This graph is illustrated in Figure 13 (with blue edges simplified as in Figure 11). In this example, each vertex of $\Lambda_4(F)$ has at least one red edge coming in to it and at least one going out of it. However, the vertex 12 is not the base point of any RBR-alternating circuit, and therefore F is not a generating set for $\mathcal{B}_4 \setminus \mathcal{S}_4$.

Condition (ii) of Remark 6.8, though sufficient, is not necessary. For example, consider the graph $\Lambda_3(F)$, where $F = \{\tau_{13}, \tau_{23}, \sigma_{213}, \sigma_{321}\}$, which is illustrated in Figure 14. It is easy to check that each vertex is the base point of an RBR-alternating circuit, so that $\mathcal{B}_3 \setminus \mathcal{S}_3 = \langle F \rangle$, even though vertex 12 is not contained in a red circuit.

9. THE JONES MONOID

Recall that the Jones monoid \mathcal{J}_n is the subsemigroup of \mathcal{B}_n consisting of all partitions whose blocks have cardinality 2 and may be drawn in a planar fashion. By stretching the diagram of an element of \mathcal{J}_n so that the vertices appear in a single straight line $1, 2, \dots, n, n', \dots, 2', 1'$, it is straightforward to verify that the elements of \mathcal{J}_n are in one-to-one correspondence with the proper bracketings with n pairs of brackets; see Figure 15 for an example. It follows that $|\mathcal{J}_n| = C_n$, where $C_n = \frac{1}{n+1} \binom{2n}{n}$ is the n th Catalan number. The only planar permutation is the identity element of \mathcal{P}_n , denoted 1, so the group of units of \mathcal{J}_n is equal to $\{1\}$. But $\mathcal{J}_n \subseteq \mathcal{B}_n$ is again closed under the $*$ operation, so Green's relations are still described by Theorem 8.1. Note, however, that the planarity condition implies that \mathcal{J}_n is \mathcal{H} -trivial.

For $1 \leq i \leq n - 1$, write $\tau_i = \tau_{i,i+1}$. (The partitions τ_{ij} were defined in the previous section.) See Figure 16 for an illustration.

The following result is well known; see for example [96], where presentations are also discussed.

Theorem 9.1. *The singular part $\mathcal{J}_n \setminus \{1\}$ of the Jones monoid \mathcal{J}_n is idempotent generated. The set $\{\tau_i : 1 \leq i \leq n - 1\}$ is a minimal idempotent generating set.*




FIGURE 13. The graph $\Lambda_4(F)$, where

$$F = \{\tau_{14}, \tau_{23}, \sigma_{214}, \sigma_{231}, \sigma_{234}, \sigma_{241}, \sigma_{314}, \sigma_{321}, \sigma_{324}, \sigma_{341}\}.$$




FIGURE 14. The graph $\Lambda_3(F)$, where $F = \{\tau_{13}, \tau_{23}, \sigma_{213}, \sigma_{321}\}$.





FIGURE 15. An element of the Jones monoid \mathcal{J}_7 (above) along with its corresponding Catalan bracketing diagram (below).

Remark 9.2. Because the singular part of the Jones monoid \mathcal{J}_n is simply $\mathcal{J}_n \setminus \{1\}$, we could state many of the results of this section in terms of *monoid* generating sets of \mathcal{J}_n ,

FIGURE 16. The projection $\tau_i \in \mathcal{J}_n$.

rather than semigroup generating sets of $\mathcal{J}_n \setminus \{1\}$. However, for reasons of consistency, we will not do this.

For $0 \leq r = n - 2k \leq n$, let

$$J_r(\mathcal{J}_n) = \{\alpha \in \mathcal{J}_n : \text{rank}(\alpha) = r\} \quad \text{and} \quad I_r(\mathcal{J}_n) = \{\alpha \in \mathcal{J}_n : \text{rank}(\alpha) \leq r\}.$$

As before, these are precisely the \mathcal{J} -classes and ideals of \mathcal{J}_n , and the \mathcal{J} -classes form a chain; but this time all maximal subgroups are trivial.

9.1. Rank and idempotent rank of ideals of \mathcal{J}_n . In the proof of Lemma 8.3, it was shown that any projection $\alpha \in J_r(\mathcal{B}_n)$ with $r = n - 2k$ and $k \geq 2$ was the product of two partitions $\beta, \gamma \in J_{r+2}(\mathcal{B}_n)$, and we wish to establish the corresponding result for \mathcal{J}_n . However, in the proof of Lemma 8.3, even if α is planar, the partitions β, γ constructed in the proof need not be planar themselves. For example, with $\alpha = \tau_{14}\tau_{23}$, the proof of Lemma 8.3 gives $\beta = \tau_{14}$ and $\gamma = \tau_{23}$, with γ being planar but not β . So we must work a bit harder to prove the next result.

Lemma 9.3. *If $0 \leq r \leq n - 4$, then $J_r(\mathcal{J}_n) \subseteq \langle J_{r+2}(\mathcal{J}_n) \rangle$.*

Proof. As usual, it suffices to show that any projection from $J_r(\mathcal{J}_n)$ belongs to $\langle J_{r+2}(\mathcal{J}_n) \rangle$, so suppose $\alpha \in J_r(\mathcal{J}_n)$ is a projection. We consider two cases. For the duration of this proof, it will be convenient to introduce some terminology. If $1 \leq i < j \leq n$ and $1 \leq r < s \leq n$, we will say that $\{i, j\}$ is *surrounded by* $\{r, s\}$ if $r < i < j < s$.

Case 1: Suppose first that α has two blocks of the form $\{i, i + 1\}$ and $\{j, j + 1\}$ that are not surrounded by any other blocks of α ; see Figure 17. In this case, $\alpha = \beta\gamma$ where $\beta, \gamma \in J_{r+2}(\mathcal{J}_n)$ are also illustrated in Figure 17. In the diagram, it is understood that the shaded parts of β, γ are the same as the corresponding shaded parts of α . (Since α is an idempotent, these shaded parts are all idempotents in isomorphic copies of \mathcal{J}_{i-1} , \mathcal{J}_{j-i-2} and \mathcal{J}_{n-j-1} .)

Case 2: If we are not in Case 1, then α must have at least one block of the form $\{i, j\}$ where $j \geq i + 3$, and $\{i, j\}$ is not surrounded by any other block of α ; see Figure 18. Due to planarity, this implies that vertices $i + 1, i + 2, \dots, j - 1$ are all involved in non-transversal blocks. In this case, $\alpha = \beta\gamma$ where $\beta, \gamma \in J_{r+2}(\mathcal{J}_n)$ are also illustrated in Figure 18. \square

Remark 9.4. In Case 1 of the above proof, β and γ were both projections from $J_{r+2}(\mathcal{J}_n)$. In Case 2, they were not (and could not be; consider the above example of $\alpha = \tau_{14}\tau_{23}$). However, they are both idempotents. Indeed, consider β for example. Because α is a projection, the only part of β we do not automatically know is idempotent is the portion contained between points $i + 1, \dots, j$. But this is essentially a (planar) rank 1 Brauer diagram, and all of these are idempotents. Similarly we see that γ is an idempotent. Since β and γ are both idempotents, it follows that any projection from $J_r(\mathcal{J}_n)$ can be expressed as the product of at most four projections from $J_{r+2}(\mathcal{J}_n)$ if $0 \leq r \leq n - 4$. In general, this bound (of four projections) is sharp, as the above example of $\alpha = \tau_{14}\tau_{23} \in \mathcal{J}_4$ demonstrates.




FIGURE 17. The partitions $\alpha \in J_r(\mathcal{J}_n)$ and $\beta, \gamma \in J_{r+2}(\mathcal{J}_n)$ from Case 1 of the proof of Lemma 9.3.




FIGURE 18. The partitions $\alpha \in J_r(\mathcal{J}_n)$ and $\beta, \gamma \in J_{r+2}(\mathcal{J}_n)$ from Case 2 of the proof of Lemma 9.3.

Theorem 9.5. For $0 \leq r = n - 2k \leq n - 2$, the ideal $I_r(\mathcal{J}_n)$ is idempotent generated, and

$$\text{rank}(I_r(\mathcal{J}_n)) = \text{idrank}(I_r(\mathcal{J}_n)) = \frac{r+1}{n+1} \binom{n+1}{k}.$$

Moreover, a subset $A \subseteq I_r(\mathcal{J}_n)$ of this cardinality is a generating set for $I_r(\mathcal{J}_n)$ if and only if the following three conditions hold:

- (1) $\text{rank}(\alpha) = r$ for all $\alpha \in A$;
- (2) for all $\alpha, \beta \in A$ with $\alpha \neq \beta$, $\ker(\alpha) \neq \ker(\beta)$;
- (3) for all $\alpha, \beta \in A$ with $\alpha \neq \beta$, $\text{coker}(\alpha) \neq \text{coker}(\beta)$.

Proof. Fix some $0 \leq r = n - 2k \leq n$, and denote by ρ_{nr} the number of \mathcal{R} -classes in $J_r(\mathcal{J}_n)$. As usual, it suffices to show that $\rho_{nr} = \frac{r+1}{n+1} \binom{n+1}{k}$. As noted above, an element $\alpha \in \mathcal{J}_n$ is completely determined by a string of length $2n$, consisting of a proper bracketing of n pairs of brackets. We denote by $L(\alpha)$ the substring consisting of the first n symbols in this string. If $\alpha \in J_r(\mathcal{J}_n)$, then this string will have k right brackets and $k+r$ left brackets, r of which are unmatched. These unmatched left brackets of $L(\alpha)$ correspond to the elements of $\text{dom}(\alpha)$, while the pairs of matched brackets in $L(\alpha)$ correspond to the non-trivial $\ker(\alpha)$ -classes. Hence, by Theorem 8.1, it follows that ρ_{nr} is equal to the number of all such strings of n brackets (with the specified number of unmatched left brackets), and it is well known that there are $\frac{r+1}{n+1} \binom{n+1}{k}$ such strings; see for example [10, p. 303]. \square

Remark 9.6. As in Remarks 7.7 and 8.6, but noting that \mathcal{H} -classes of \mathcal{J}_n are singletons, we obtain

$$|J_r(\mathcal{J}_n)| = \rho_{nr}^2 \quad \text{and} \quad |I_r(\mathcal{J}_n)| = \sum_{i=0}^r \rho_{ni}^2 \quad \text{for any } 0 \leq r \leq n,$$

where the numbers ρ_{nr} are as in (the proof of) Theorem 9.5, and we interpret $\rho_{ni} = 0$ if $i \not\equiv n \pmod{2}$.

Remark 9.7. The sequence ρ_{nr} is A053121 on [1], and appears in a variety of contexts, most notably as the Catalan triangle (concerning U/D walks in the first quadrant). Of particular present interest is that these numbers give the dimensions of the irreducible representations of the *Temperley–Lieb algebra* of degree n in the semisimple case. The dimensions of the irreducible representations of the *Brauer* and *partition algebras* may also be obtained from our results on (idempotent) ranks of ideals of the corresponding diagram monoids. See Section 10 below for full details of this.

$n \setminus r$	0	1	2	3	4	5	6	7	8	9	10
0	1										
1		1									
2	1		1								
3		2		1							
4	2		3		1						
5		5		4		1					
6	5		9		5		1				
7		14		14		6		1			
8	14		28		20		7		1		
9		42		48		27		8		1	
10	42		90		75		35		9		1

TABLE 9. Values of ρ_{nr} from the proof of Theorem 9.5. For $0 \leq r \leq n-2$, $\rho_{nr} = \text{rank}(I_r(\mathcal{J}_n)) = \text{idrank}(I_r(\mathcal{J}_n))$.

9.2. Minimal idempotent generating sets of $\mathcal{J}_n \setminus \{1\}$. Again, an enumeration of the minimal idempotent generating sets of $\mathcal{J}_n \setminus \{1\}$ amounts to an enumeration of the balanced subgraphs of the projection graph $\Gamma(\mathcal{J}_n \setminus \{1\}) = \Gamma(J_{n-2}(\mathcal{J}_n)^*)$, which we will denote by Ξ_n .

For $1 \leq i \leq n-2$, put $\lambda_i = \sigma_{i,i+1,i+2}$ and $\rho_i = \sigma_{i+2,i+1,i}$. (The idempotents σ_{ijk} were defined in the previous section.) See Figure 19 for an illustration.

Again, the next lemma is easily verified, and its proof is omitted.


 FIGURE 19. The idempotents λ_i (left) and ρ_i (right) from \mathcal{J}_n .

Lemma 9.8. *The set of idempotents of $J_{n-2}(\mathcal{J}_n)$ is*

$$\{\tau_i : 1 \leq i \leq n-1\} \cup \{\lambda_i, \rho_i : 1 \leq i \leq n-2\}.$$


The set of projections of $J_{n-2}(\mathcal{J}_n)$ is

$$\{\tau_i : 1 \leq i \leq n-1\}.$$

In the principal factor $J_{n-2}(\mathcal{J}_n)^$, the only nonzero products of pairs of projections are*

$$\tau_i^2 = \tau_i, \quad \tau_i \tau_{i+1} = \lambda_i, \quad \tau_{i+1} \tau_i = \rho_i.$$

It follows that the graph Ξ_n has vertex set $\{\tau_i : 1 \leq i \leq n-1\}$ with edges $\tau_i \rightarrow \tau_j$ if and only if $|i-j| \leq 1$. The graph Ξ_5 is pictured in Figure 20, with vertices labeled i instead of τ_i .


 FIGURE 20. The graph $\Xi_5 = \Gamma(\mathcal{J}_5 \setminus \{1\})$.

Theorem 9.9. *The number of minimal idempotent generating sets for the singular part $\mathcal{J}_n \setminus \{1\}$ of the Jones monoid \mathcal{J}_n is equal to F_n , the n th Fibonacci number, where $F_1 = F_2 = 1$ and $F_n = F_{n-1} + F_{n-2}$ for $n \geq 3$.*

Proof. Let x_n be the number of balanced subgraphs of Ξ_n . It suffices to show that the numbers x_n satisfy the Fibonacci recurrence. It is clear that $x_1 = x_2 = 1$. (Note that $\mathcal{J}_1 \setminus \{1\} = \emptyset$, so there is only one generating set, \emptyset .) Next, suppose $n \geq 3$. A balanced subgraph of Ξ_n must contain either the loop at τ_{n-1} and a balanced subgraph of Ξ_{n-1} , or else the edges $\tau_{n-2} \rightleftarrows \tau_{n-1}$ and a balanced subgraph of Ξ_{n-2} . So $x_n = x_{n-1} + x_{n-2}$, as required. \square

Although these numbers are well-known (see A000045 on [1]), we include the first few values, for completeness, in Table 10.

n	1	2	3	4	5	6	7	8	9	10
F_n	1	1	2	3	5	8	13	21	34	55

TABLE 10. The Fibonacci sequence F_n . For $n \geq 1$, F_n is equal to the number of minimal idempotent generating sets of $\mathcal{J}_n \setminus \{1\}$.

9.3. Arbitrary idempotent generating sets for $\mathcal{J}_n \setminus \{1\}$. For a subset F of

$$\{\alpha \in E(\mathcal{J}_n) : \text{rank}(\alpha) = n - 2\} = \{\tau_1, \dots, \tau_{n-1}\} \cup \{\lambda_1, \dots, \lambda_{n-2}\} \cup \{\rho_1, \dots, \rho_{n-2}\},$$

let $\Xi_n(F)$ be the two-coloured digraph obtained by colouring each edge of Ξ_n blue, and then adding red edges corresponding to the idempotents from F :

- $i \rightarrow i$ if $\tau_i \in F$,
- $i \rightarrow i + 1$ if $\lambda_i \in F$,
- $i + 1 \rightarrow i$ if $\rho_i \in F$.

(As above, we will denote the vertices of Ξ_n by $1, \dots, n - 1$ rather than $\tau_1, \dots, \tau_{n-1}$.)

Theorem 9.10. *For $F \subseteq \{\alpha \in E(\mathcal{J}_n) : \text{rank}(\alpha) = n - 2\}$, the following are equivalent:*

- (i) $\mathcal{J}_n \setminus \{1\} = \langle F \rangle$;
- (ii) each vertex of $\Xi_n(F)$ is the base point of an RBR-alternating circuit;
- (iii) each vertex of $\Xi_n(F)$ is contained in a red circuit.

The number f_n of such subsets is given by the recurrence

$$f_2 = 1, \quad f_3 = 7, \quad f_n = 5f_{n-1} + 6f_{n-2} \quad \text{for } n \geq 4.$$

Proof. By Lemma 9.3, F generates $\mathcal{J}_n \setminus \{1\}$ if and only if it generates the principle factor $J_{n-2}(\mathcal{J}_n)^*$. So the equivalence of (i) and (ii) follows from Theorem 6.5. Any red circuit beginning and ending at vertex i clearly gives rise to an RBR-circuit based at i , since $\Xi_n(F)$ has a blue loop at each vertex, so (iii) implies (ii).

Now assume (ii), and suppose $1 \leq i \leq n - 1$. We must show that i is contained in a red circuit. Consider first the case in which $i = 1$. If $1 \rightarrow 1$ is an edge, then 1 is contained in a red circuit. If not, then, since there is an RBR-alternating circuit based at 1, we see that $\Xi_n(F)$ contains the edges $1 \rightleftarrows 2$, so 1 is contained in a red circuit. By symmetry, $n - 1$ is contained in a red circuit.

Finally, suppose $1 < i < n - 1$. If $\Xi_n(F)$ contains the loop $i \rightarrow i$ or the edges $i \rightleftarrows i + 1$ or $i + 1 \rightleftarrows i$, then i is contained in a red circuit. In order to obtain a contradiction, suppose this is not the case. By symmetry, we may suppose that $\Xi_n(F)$ contains the edges $i - 1 \rightarrow i \rightarrow i + 1$, but no other red edges at i . Now consider an RBR-alternating path beginning at i :

$$i = j_0 \rightarrow j_1 \rightarrow j_2 \rightarrow \cdots \rightarrow j_{2k} \rightarrow j_{2k+1}.$$

A simple induction shows that $j_{2r} \geq i$ and $j_{2r+1} \geq i + 1$ for each $0 \leq r \leq k$. In particular, $j_{2k+1} \geq i + 1$, so it follows that $\Xi_n(F)$ contains no RBR-alternating circuit based at vertex i , a contradiction. This completes the proof that (ii) implies (iii).

Now let \mathcal{X}_n denote the set of all graphs $\Xi_n(F)$ satisfying property (iii). We must show that the cardinalities $f_n = |\mathcal{X}_n|$ satisfy the given recurrence. It is easy to check that $f_2 = 1$ and $f_3 = 7$, so suppose now that $n \geq 4$. Let $G \in \mathcal{X}_n$. Since the vertex $n - 1$ is contained in a red circuit, there are three cases we must consider:

- (1) G does not contain both of the edges $n - 1 \rightleftarrows n - 2$, or
- (2) G contains both of the edges $n - 2 \rightleftarrows n - 1$ and the loop $n - 1 \rightarrow n - 1$, or
- (3) G contains both of the edges $n - 2 \rightleftarrows n - 1$ but not the loop $n - 1 \rightarrow n - 1$.

If (1) holds, then G contains the loop $n - 1 \rightarrow n - 1$, and the subgraph of G induced by the vertices $1, \dots, n - 2$ belongs to \mathcal{X}_{n-1} . Since there are three ways to choose at most one of the edges $n - 1 \rightleftarrows n - 2$, there are $3f_{n-1}$ graphs G satisfying (1).

Now suppose (2) holds. Let H be the subgraph of G induced by the vertices $1, \dots, n - 2$. Note that G is completely determined by H . There are two subcases:

- (2.1) H belongs to \mathcal{X}_{n-1} , or
- (2.2) H does not belong to \mathcal{X}_{n-1} .




FIGURE 21. The digraph $\Xi_4(F)$, where $F = \{\tau_1, \tau_3, \lambda_1, \lambda_2\}$.

There are obviously f_{n-1} graphs G satisfying (2.1). Now suppose (2.2) holds. Now, each vertex $1, \dots, n-3$ belongs to a red circuit of G . Since such a circuit does not need to include the edges $n-2 \rightleftarrows n-1$, we see that, in fact, each vertex $1, \dots, n-3$ belongs to a red circuit of H . Since we assumed that $H \notin \mathcal{X}_{n-1}$, it follows that H does not contain the loop $n-2 \rightarrow n-2$ and contains at most one of the edges $n-3 \rightleftarrows n-2$. It also follows that the subgraph of H induced by the vertices $1, \dots, n-3$ belongs to \mathcal{X}_{n-2} . So there are $3f_{n-2}$ graphs G satisfying (2.2). Thus, there are $f_{n-1} + 3f_{n-2}$ graphs G satisfying (2). The same argument shows that there are the same number of graphs G satisfying (3). Putting this together, $f_n = 3f_{n-1} + 2(f_{n-1} + 3f_{n-2}) = 5f_{n-1} + 6f_{n-2}$. This completes the proof. \square

Remark 9.11. Solving the above recurrence gives

$$f_n = \frac{2}{63} \cdot 6^n + \frac{1}{7} \cdot (-1)^{n+1}.$$

This is sequence A108983 on [1]. The first few values of the sequence f_n are given in Table 11.

n	2	3	4	5	6	7	8	9	10
f_n	1	7	41	247	1481	8887	53321	319927	1919561

TABLE 11. The sequence f_n , which gives the number of generating sets for $\mathcal{J}_n \setminus \{1\}$ consisting of idempotents belonging to $J_{n-2}(\mathcal{J}_n)$.

Remark 9.12. Theorem 9.10 shows that condition (i) from Remark 6.8 is necessary and sufficient in the case of $\mathcal{J}_n \setminus \{1\}$. While condition (ii) from Remark 6.8 is, as ever, necessary, it is not sufficient in general. For example, consider the set of idempotents $F = \{\tau_1, \tau_3, \lambda_1, \lambda_2\}$ from \mathcal{J}_4 . The digraph $\Xi_4(F)$ is displayed in Figure 21. As in Example 7.17, there is no RBR-alternating circuit at vertex 2, so $\mathcal{J}_n \setminus \{1\} \neq \langle F \rangle$.

9.4. The monoid of planar partitions. Recall that the monoid of planar partitions is denoted \mathcal{PP}_n . It is well known that \mathcal{PP}_n is isomorphic to the Jones monoid \mathcal{J}_{2n} ; see for example [79]. This isomorphism is easiest to describe diagrammatically, and we do so in Figure 22. Because of this isomorphism, we will not state the results concerning the ideals

$$I_r(\mathcal{PP}_n) = \{\alpha \in \mathcal{PP}_n : \text{rank}(\alpha) \leq r\}$$

and minimal idempotent generating sets, as these can be deduced easily from the results for \mathcal{J}_n in the preceding sections.

10. RANKS OF IDEALS AND DIMENSIONS OF CELL MODULES AND IRREDUCIBLE REPRESENTATIONS

As mentioned in the introduction, further motivation for the rank and idempotent rank formulae obtained above comes from the connection between these numbers and the dimensions of the irreducible representations of the corresponding algebras (realised as twisted semigroup algebras): namely, the partition, Brauer and Temperley–Lieb algebras.




FIGURE 22. A planar partition from \mathcal{PP}_8 (black) and its image (gray) under the isomorphism $\mathcal{PP}_8 \rightarrow \mathcal{J}_{16}$.

Specifically, the rank and idempotent rank formulae we obtained above can be used to recover formulae for dimensions of cell modules which, in the semisimple case, correspond to dimensions of irreducible representations. This fact was brought to the attention of the authors by Arun Ram (at the Workshop on Diagram Algebras, Stuttgart, 2014) who pointed out that the set of ranks (and idempotent ranks) of the two-sided ideals of the Jones monoid give precisely the dimensions of the irreducible representations of the Temperley–Lieb algebras. Analogous statements hold for the partition and Brauer algebras, and the purpose of this section is to explain this connection. We note that the formulae for the dimensions of the cell modules that we derive here are not new (we provide the relevant references below). Our aim here is to give a new way of deriving these formulae, and at the same time explain the relationship between the problem of computing such numbers and that of computing ranks (and idempotent ranks) of ideals of diagram monoids.

In each case the main idea is the same: the ranks of ideals of the semigroup are given by numbers of \mathcal{R} -classes in \mathcal{J} -classes of the semigroup. By [136], it follows that these numbers in turn arise in the construction of the cell modules of the corresponding twisted semigroup algebras when they are realised as cellular algebras. In the case that the algebras are semisimple, the cell modules give a complete set of irreducible representations of the algebra. The questions of determining the dimensions of these representations can then easily be reduced to the question of determining the ranks of the ideals of the semigroups, which are given by the formulae in Theorems 7.5, 8.4 and 9.5 above (depending on the semigroup–algebra pair under consideration).

We shall now explain this in more detail for the partition monoid and partition algebra. The arguments for the other pairs (the Brauer semigroup and Brauer algebra, and the Jones monoid and Temperley–Lieb algebra) are analogous, and for these we shall just state the corresponding results and give relevant references.

Dimensions of irreducible representations of partition algebras. In this subsection we follow [136, Section 7] to define the partition algebra and explain its cellular structure. Let R be a commutative ring with identity, let $\delta \in R$ be fixed, and let ψ be the mapping

$$\psi : \mathcal{P}_n \times \mathcal{P}_n \rightarrow R, \quad (\alpha, \beta) \mapsto \delta^{m(\alpha, \beta)},$$

called a *twisting* from \mathcal{P}_n to R , where $m(\alpha, \beta)$ denotes the number of connected components removed from the middle row when constructing the composition $\alpha\beta$ in \mathcal{P}_n . The resulting twisted semigroup algebra $R^\psi[\mathcal{P}_n]$ is called the *partition algebra*; see [89, 101, 102]. The algebra $R^\psi[\mathcal{P}_n]$ has \mathcal{P}_n as its basis, and its multiplication \star is defined on basis elements (and then extended linearly) by $\alpha \star \beta = \psi(\alpha, \beta)(\alpha\beta)$.

Recall from Section 7 above that there is a natural anti-involution $*$ on \mathcal{P}_n that reflects graphs representing elements in the horizontal axis. By linearity, this extends to an R -linear anti-involution (also denoted $*$) on the partition algebra $R^\psi[\mathcal{P}_n]$. Also recall from Section 7 that $\text{rank}(\alpha)$ is the number of transversal blocks of $\alpha \in \mathcal{P}_n$, that the \mathcal{J} -classes of \mathcal{P}_n are the sets

$$J_r(\mathcal{P}_n) = \{\alpha \in \mathcal{P}_n : \text{rank}(\alpha) = r\}$$

where $0 \leq r \leq n$, that they form a chain:

$$J_0(\mathcal{P}_n) < J_1(\mathcal{P}_n) < \cdots < J_{n-1}(\mathcal{P}_n) < J_n(\mathcal{P}_n),$$

and that the ideals of \mathcal{P}_n are precisely the sets

$$I_r(\mathcal{P}_n) = J_0(\mathcal{P}_n) \cup J_1(\mathcal{P}_n) \cup \cdots \cup J_r(\mathcal{P}_n) = \{\alpha \in \mathcal{P}_n : \text{rank}(\alpha) \leq r\}.$$

For each \mathcal{J} -class $J = J_r(\mathcal{P}_n)$, we may choose and fix a maximal subgroup G_J of J that is fixed setwise by $*$ (such a maximal subgroup contains a projection) and in this way the $*$ operation restricted to $G_J \cong \mathcal{S}_r$ corresponds to inversion in \mathcal{S}_r . Since the group algebra $R[\mathcal{S}_r]$ is cellular with the anti-involution induced by inversion (see for example [111]), it follows that $R[G_J]$ is cellular with respect to $*$, and thus (by [136, Corollary 7]) the partition algebra $R^\psi[\mathcal{P}_n]$ is cellular with anti-involution $*$. Viewing the partition algebra $R^\psi[\mathcal{P}_n]$ in this way, as a cellular algebra, gives information about its representation theory, by appealing to the general theory of cellular algebras in the following way.

Let A be a cellular algebra with cell datum $(\Lambda, M, \mathcal{C}, *)$ (see [62]). Here Λ is a partially ordered set, the algebra A has an R -basis

$$\mathcal{C} = \{C_{\mathfrak{s}, \mathfrak{t}}^\lambda : \lambda \in \Lambda, \mathfrak{s}, \mathfrak{t} \in M(\lambda)\},$$

and an anti-involution $*$: $A \rightarrow A$ is given by $(C_{\mathfrak{s}, \mathfrak{t}}^\lambda)^* = C_{\mathfrak{t}, \mathfrak{s}}^\lambda$. Moreover, for all $\lambda \in \Lambda$, $\mathfrak{s}, \mathfrak{t} \in M(\lambda)$ and $a \in A$

$$aC_{\mathfrak{s}, \mathfrak{t}}^\lambda \equiv \sum_{\mathfrak{s}' \in M(\lambda)} r_a(\mathfrak{s}', \mathfrak{s}) C_{\mathfrak{s}', \mathfrak{t}}^\lambda \pmod{A(< \lambda)}$$

where each $r_a(\mathfrak{s}', \mathfrak{s}) \in R$ is independent of \mathfrak{t} , and where $A(< \lambda)$ is the R -submodule of A generated by $\{C_{\mathfrak{u}, \mathfrak{v}}^\mu : \mu < \lambda, \mathfrak{u}, \mathfrak{v} \in M(\mu)\}$. It follows that for all $\lambda \in \Lambda$ and all $\mathfrak{s}_1, \mathfrak{s}_2, \mathfrak{t}_1, \mathfrak{t}_2 \in M(\lambda)$ we have

$$C_{\mathfrak{s}_1, \mathfrak{t}_1}^\lambda C_{\mathfrak{s}_2, \mathfrak{t}_2}^\lambda \equiv \phi(\mathfrak{t}_1, \mathfrak{s}_2) C_{\mathfrak{s}_1, \mathfrak{t}_2}^\lambda \pmod{A(< \lambda)}$$

for some $\phi(\mathfrak{t}_1, \mathfrak{s}_2) \in R$ that depends only on \mathfrak{t}_1 and \mathfrak{s}_2 .

For each $\lambda \in \Lambda$, let $W(\lambda)$ denote the left A -module with R -basis $\{C_{\mathfrak{s}} : \mathfrak{s} \in M(\lambda)\}$ and A -action given by

$$aC_{\mathfrak{s}} = \sum_{\mathfrak{s}' \in M(\lambda)} r_a(\mathfrak{s}', \mathfrak{s}) C_{\mathfrak{s}'}$$

for each $a \in A$. The $W(\lambda)$ are called *cell modules*. For $\lambda \in \Lambda$ define the bilinear form

$$\phi_\lambda : W(\lambda) \times W(\lambda) \rightarrow R, \quad \phi_\lambda(C_{\mathfrak{s}}, C_{\mathfrak{t}}) = \phi(\mathfrak{s}, \mathfrak{t}),$$

for $\mathfrak{s}, \mathfrak{t} \in M(\lambda)$. The radical of $\lambda \in \Lambda$ is then the A -submodule

$$\text{rad}(\lambda) = \{x \in W(\lambda) : \phi_\lambda(x, y) = 0 \ \forall y \in W(\lambda)\}$$

of $W(\lambda)$. Let us now quote two important results from [62].

Theorem 10.1 (Graham and Lehrer [62, Theorem 3.4(i)]). *Let R be a field and let $(\Lambda, M, \mathcal{C}, *)$ be a cell datum for the R -algebra A . For $\lambda \in \Lambda$ let $L_\lambda = W(\lambda)/\text{rad}(\lambda)$. Then*

$$\{L_\lambda : \lambda \in \Lambda, \phi_\lambda \neq 0\}$$

is a complete set of (representatives of equivalence classes of) absolutely irreducible A -modules.

Theorem 10.2 (Graham and Lehrer [62, Theorem 3.8]). *Let A be an R -algebra (R a field) with cell datum $(\Lambda, M, \mathcal{C}, *)$. Then the following are equivalent.*

- (i) *The algebra A is semisimple.*
- (ii) *The nonzero cell representations $W(\lambda)$ are irreducible and pairwise inequivalent.*
- (iii) *The form ϕ_λ is nondegenerate (i.e. $\text{rad}(\lambda) = 0$) for each $\lambda \in \Lambda$.*

It follows from these results that the irreducible left A -modules are parameterised by the set

$$\Lambda_0 = \{\lambda \in \Lambda : \phi_\lambda \neq 0\},$$

and the dimensions of the irreducible left A -modules are given by

$$\dim_R(L_\lambda) = |M(\lambda)| - \dim_R(\text{rad}(\lambda)).$$

In particular, in the case of semisimple cellular algebras, these dimensions are given simply by $|M(\lambda)|$ with $\lambda \in \Lambda$.

Returning our attention to the partition algebra $R^\psi[\mathcal{P}_n]$, applying [136, Theorem 5 and Corollary 7], Wilcox shows that the partition algebra $R^\psi[\mathcal{P}_n]$ is cellular with cell datum $(\Lambda, M, \mathcal{C}, *)$ where

$$\Lambda = \{(J, \lambda) : J \in \mathbb{J} \text{ and } \lambda \in \Lambda_J\},$$

with a partial order defined on it (which we shall not need here), the symbol \mathbb{J} denotes the set of \mathcal{J} -classes of the semigroup \mathcal{P}_n , and Λ_J comes from the cell datum $(\Lambda_J, M_J, \mathcal{C}_J, *)$ for the (non-twisted) group algebra $R[G_J]$ where G_J is the a maximal subgroup of D fixed set-wise by $*$ defined above. Moreover, for $(J, \lambda) \in \Lambda$ we have


$$M(J, \lambda) = \mathbb{L}_J \times M_J(\lambda),$$

where \mathbb{L}_J is the set of \mathcal{L} -classes of the \mathcal{J} -class J , and so in particular

$$|M(J, \lambda)| = |\mathbb{L}_J| \cdot |M_J(\lambda)|. \quad (10.3)$$

Finally, let us now consider the case that R is the field \mathbb{C} of complex numbers and consider the complex partition algebra $\mathbb{C}^\psi[\mathcal{P}_n]$. For fixed n , semisimplicity of the algebra $\mathbb{C}^\psi[\mathcal{P}_n]$ depends on ψ . This is explored in more detail in [79]. Specifically it is observed that, with δ in the definition of ψ , for all but a finite number of $\delta \in \mathbb{C}$, the algebra $\mathbb{C}^\psi[\mathcal{P}_n]$ is semisimple. In the cases that $\mathbb{C}^\psi[\mathcal{P}_n]$ is semisimple, a method for computing dimensions of irreducible $\mathbb{C}^\psi[\mathcal{P}_n]$ -modules in terms of counting paths in a certain graph \hat{A} , whose vertices are labelled by integer partitions, is given in [79, Theorem 2.24]. Here our aim is to state a result analogous to [79, Theorem 2.24(b)] but where we express the dimensions of the irreducible $\mathbb{C}^\psi[\mathcal{P}_n]$ -modules in terms of the rank formula we obtained in Theorem 7.5.

In order to do this we need to recall some basic notions about integer partitions and the representation theory of the symmetric group. Any integer partition λ can be identified with a sequence $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_k)$. Integer partitions are represented using Young diagrams; for example, the diagram corresponding to $\lambda = (543311)$ is



The *hook length* of the box b of λ is

$$h(b) = (\lambda_i - j) + (\lambda'_j - i) + 1 \text{ if } b \text{ is in position } (i, j) \text{ of } \lambda.$$

Here, $\lambda' = (\lambda'_1 \geq \lambda'_2 \geq \dots \geq \lambda'_{\lambda_1})$ denotes the partition obtained by reflecting (the Young diagram corresponding to) λ in the leading diagonal. In other words, $h(b)$ is the number of boxes to the right of b plus the number below b , plus one (to include the box b itself in the count). Write $\lambda \vdash r$ and $|\lambda| = r$ if λ is a partition with r boxes (so $\lambda_1 + \lambda_2 + \dots + \lambda_k = r$).

Let $0 \leq r \leq n - 1$ and consider the \mathcal{J} -class $J = J_r(\mathcal{P}_n)$ of the partition monoid \mathcal{P}_n . The maximal subgroups of this \mathcal{J} -class are isomorphic to the symmetric group \mathcal{S}_r . As noted above, the algebra $\mathbb{C}[\mathcal{S}_r]$ is known to be cellular with cell datum $(\Lambda_J, M_J, \mathcal{C}_J, *)$ where Λ_J is the set $\hat{\mathcal{S}}_r = \{\lambda : \lambda \vdash r\}$ of all partitions with r boxes, carrying a natural partial order (which will not be needed here). For a partition λ , let $\text{Std}(\lambda)$ denote the set of standard λ -tableaux (that is, all ways of filling the boxes of λ with the symbols 1 up

to r so that both rows and columns are strictly increasing). Then for $\lambda \in \Lambda_J$, we have $M_J(\lambda) = \text{Std}(\lambda)$.

So the irreducible $\mathbb{C}[\mathcal{S}_r]$ -modules, denoted S_r^λ , are indexed by the elements of $\Lambda_J = \hat{\mathcal{S}}_r = \{\lambda : \lambda \vdash r\}$. For a given $\lambda \in \Lambda_J$ we have $M_J(\lambda) = \text{Std}(\lambda)$ and the dimension of the corresponding irreducible module S_r^λ is given by $\dim(S_r^\lambda) = |M_J(\lambda)| = |\text{Std}(\lambda)|$. This number is well known to be given by

$$|M_J(\lambda)| = |\text{Std}(\lambda)| = \frac{r!}{\prod_{b \in \lambda} h(b)},$$

where $h(b)$ denotes the hook-length of the box b of λ , as defined above. In the last expression, we write $b \in \lambda$ to indicate that b is a box of (the Young diagram representing) λ . Now combining these observations, Equation 10.3 and Theorem 7.5 we obtain the following.

Proposition 10.4. *If $\mathbb{C}^\psi[\mathcal{P}_n]$ is semisimple, then the irreducible $\mathbb{C}^\psi[\mathcal{P}_n]$ -modules A_n^μ are indexed by elements of the set*

$$\hat{A}_n = \{\text{partitions } \mu : 0 \leq |\mu| \leq n\}.$$

Moreover, for $|\mu| < n$, we have

$$\dim(A_n^\mu) = \text{rank}(I_{|\mu|}(\mathcal{P}_n)) \cdot \left(\frac{|\mu|!}{\prod_{b \in \mu} h(b)} \right) = \left(\sum_{j=|\mu|}^n S(n, j) \binom{j}{|\mu|} \right) \cdot \frac{|\mu|!}{\prod_{b \in \mu} h(b)}.$$

We note that in the cases $|\mu| = n$, the right hand side of the formula in the above proposition still holds, but the expression with the term $\text{rank}(I_{|\mu|}(\mathcal{P}_n))$ is no longer correct (since the partition monoid $\mathcal{P}_n = I_n(\mathcal{P}_n)$ itself does not have rank 1). It is interesting to compare this statement with [79, Theorem 2.24(b)], where the dimensions are given by counting paths in a certain graph \hat{A} , whose vertices are labelled by partitions (so called, Bratteli diagrams).

Dimensions of irreducible representations of Brauer algebras. Recall from Section 8 above that the Brauer monoid \mathcal{B}_n is the subsemigroup of \mathcal{P}_n consisting of all partitions whose blocks have cardinality 2, that for $r = n - 2k$ with $k \geq 0$ and $0 \leq r \leq n$, we write

$$J_r(\mathcal{B}_n) = \{\alpha \in \mathcal{B}_n : \text{rank}(\alpha) = r\} = J_r(\mathcal{P}_n) \cap \mathcal{B}_n,$$

that these are precisely the \mathcal{J} -classes of \mathcal{B}_n , and that they form a chain:

$$J_m(\mathcal{B}_n) < J_{m+2}(\mathcal{B}_n) < \cdots < J_{n-2}(\mathcal{B}_n) < J_n(\mathcal{B}_n),$$

where m denotes 0 if n is even, or 1 otherwise. As in the partition monoid, the maximal subgroups of the \mathcal{J} -class $J_r(\mathcal{B}_n)$ are isomorphic to the symmetric group \mathcal{S}_r . The ideals of \mathcal{B}_n are precisely the sets

$$I_r(\mathcal{B}_n) = J_m(\mathcal{B}_n) \cup J_{m+2}(\mathcal{B}_n) \cup \cdots \cup J_r(\mathcal{B}_n) = \{\alpha \in \mathcal{B}_n : \text{rank}(\alpha) \leq r\}.$$

The twisted semigroup algebra $R^\psi[\mathcal{B}_n]$ is called the Brauer algebra. (For simplicity, we write ψ for the restriction of the twisting ψ to \mathcal{B}_n .) This algebra has been studied extensively in the literature; see for example [8, 81, 134]. Semisimplicity of Brauer algebras is considered in [129, 130, 134]. Cellularity of the Brauer algebra $R^\psi[\mathcal{B}_n]$ may be proved, as for the partition algebra, by appealing to cellularity of symmetric group algebras; see [136, Section 8]. Following the same argument used for the partition algebra above, and applying Theorem 8.4, we obtain the following.

Proposition 10.5. *If $\mathbb{C}^\psi[\mathcal{B}_n]$ is semisimple, then the irreducible $\mathbb{C}^\psi[\mathcal{B}_n]$ -modules, B_n^μ are indexed by elements of the set*

$$\hat{B}_n = \{\text{partitions } \mu : 0 \leq |\mu| = r = n - 2k \leq n\}.$$

Moreover, for $0 \leq |\mu| = r = n - 2k \leq n - 2$ we have

$$\dim(B_n^\mu) = \text{rank}(I_{|\mu|}(\mathcal{P}_n)) \cdot \left(\frac{|\mu|!}{\prod_{b \in \mu} h(b)} \right) = \frac{n!}{2^k k! r!} \left(\frac{|\mu|!}{\prod_{b \in \mu} h(b)} \right) = \frac{n!}{2^k k! \prod_{b \in \mu} h(b)}.$$

We have not been able to find this formula as stated in the above proposition anywhere in the literature. However, a similar (and equivalent) formula may be found in [122, Equation 3.4]. Also, in a similar way as for the partition algebra above, there is an alternative approach to computing these dimensions using Bratteli diagrams; see [134, Theorem 3.2].

Dimensions of irreducible representations of Temperley–Lieb algebras. Recall from Section 9 that the Jones monoid \mathcal{J}_n is the subsemigroup of \mathcal{B}_n consisting of all partitions whose blocks have cardinality 2 and may be drawn in a planar fashion, that for $0 \leq r = n - 2k \leq n$,

$$J_r(\mathcal{J}_n) = \{\alpha \in \mathcal{J}_n : \text{rank}(\alpha) = r\} \quad \text{and} \quad I_r(\mathcal{J}_n) = \{\alpha \in \mathcal{J}_n : \text{rank}(\alpha) \leq r\}$$

are precisely the \mathcal{J} -classes and ideals of \mathcal{J}_n , and that the \mathcal{J} -classes form a chain. In contrast to the partition and Brauer monoids, the maximal subgroups of the Jones monoid are all trivial.

The twisted semigroup algebra $R^\psi[\mathcal{J}_n]$ is called the Temperley–Lieb algebra; see [60, 89, 131]. In [135, Section 5] a simple criterion for semisimplicity of Temperley–Lieb algebras is given. Cellularity of the Temperley–Lieb algebras may be proved as for the partition and Brauer algebras above, but this time appealing to the fact that the maximal subgroups are trivial, and thus the corresponding group algebras are trivially cellular; see [136, Section 8].

Following the same approach as used for the partition and Brauer algebras above, and applying Theorem 9.5, we obtain the following.

Proposition 10.6. *If $\mathbb{C}^\psi[\mathcal{J}_n]$ is semisimple, then the irreducible $\mathbb{C}^\psi[\mathcal{J}_n]$ -modules, J_n^r are indexed by elements of the set*

$$\hat{J}_n = \{r \in \mathbb{Z}_{\geq 0} : 0 \leq r = n - 2k \leq n\}.$$

Moreover, for $0 \leq r = n - 2k \leq n - 2$, we have

$$\dim(J_n^r) = \text{rank}(I_r(\mathcal{J}_n)) = \frac{r+1}{n+1} \binom{n+1}{k}.$$

This agrees with known results from the literature; see the recent survey article [127] for details.

Acknowledgements. We thank the referees for their careful reading of the article, for several helpful suggestions that improved readability, and for drawing our attention to a number of interesting references.

REFERENCES

- [1] The on-line encyclopedia of integer sequences. Published electronically at <http://oeis.org/>, 2015.
- [2] K. Auinger. Krohn–Rhodes complexity of Brauer type semigroups. *Port. Math.*, 69(4):341–360, 2012.
- [3] K. Auinger. Pseudovarieties generated by Brauer type monoids. *Forum Math.*, 26(1):1–24, 2014.
- [4] K. Auinger, Y. Chen, X. Hu, Y. Luo, and M. Volkov. The finite basis problem for Kauffman monoids. *Algebra Universalis*, to appear.
- [5] K. Auinger, I. Dolinka, and M. V. Volkov. Equational theories of semigroups with involution. *J. Algebra*, 369:203–225, 2012.
- [6] G. Ayik, H. Ayik, L. Bugay, and O. Kelekci. Generating sets of finite singular transformation semigroups. *Semigroup Forum*, 86(1):59–66, 2013.
- [7] H. Barcelo and A. Ram. Combinatorial representation theory. In *New perspectives in algebraic combinatorics (Berkeley, CA, 1996–97)*, volume 38 of *Math. Sci. Res. Inst. Publ.*, pages 23–90. Cambridge Univ. Press, Cambridge, 1999.

- [8] R. Brauer. On algebras which are connected with the semisimple continuous groups. *Ann. of Math.* (2), 38(4):857–872, 1937.
- [9] M. Brittenham, S. W. Margolis, and J. Meakin. Subgroups of the free idempotent generated semigroups need not be free. *J. Algebra*, 321(10):3026–3042, 2009.
- [10] R. A. Brualdi. *Introductory combinatorics*. Pearson Prentice Hall, Upper Saddle River, NJ, fifth edition, 2010.
- [11] K. Byleen. Embedding any countable semigroup in a 2-generated bisimple monoid. *Glasgow Math. J.*, 25(2):153–161, 1984.
- [12] A. H. Clifford. Matrix representations of completely simple semigroups. *Amer. J. Math.*, 64:327–342, 1942.
- [13] A. H. Clifford. Basic representations of completely simple semigroups. *Amer. J. Math.*, 82:430–434, 1960.
- [14] A. H. Clifford and G. B. Preston. *The algebraic theory of semigroups. Vol. I*. Mathematical Surveys, No. 7. American Mathematical Society, Providence, R.I., 1961.
- [15] S. Crvenković and I. Dolinka. Congruences on $*$ -regular semigroups. *Period. Math. Hungar.*, 45(1-2):1–13, 2002.
- [16] M. C. Cuaresma, M. Giudici, and C. E. Praeger. Homogeneous factorisations of Johnson graphs. *Des. Codes Cryptogr.*, 46(3):303–327, 2008.
- [17] Y. Dandan, I. Dolinka, and V. Gould. Free idempotent generated semigroups and endomorphism monoids of free G -acts. *J. Algebra*, 429:133–176, 2015.
- [18] Y. Dandan and V. Gould. Free idempotent generated semigroups over bands and biorordered sets with trivial products. *Internat. J. Algebra Comput.*, 26(3):473–507, 2016.
- [19] R. J. H. Dawlings. Products of idempotents in the semigroup of singular endomorphisms of a finite-dimensional vector space. *Proc. Roy. Soc. Edinburgh Sect. A*, 91(1-2):123–133, 1981/82.
- [20] R. J. H. Dawlings. Sets of idempotents that generate the semigroup of singular endomorphisms of a finite-dimensional vector space. *Proc. Edinburgh Math. Soc. (2)*, 25(2):133–139, 1982.
- [21] I. Dolinka. A note on maximal subgroups of free idempotent generated semigroups over bands. *Period. Math. Hungar.*, 65(1):97–105, 2012.
- [22] I. Dolinka and J. East. Variants of finite full transformation semigroups. *Internat. J. Algebra Comput.*, 25(8):1187–1222, 2015.
- [23] I. Dolinka and J. East. Idempotent generation in the endomorphism monoid of a uniform partition. *Comm. Algebra*, 44(12):5179–5198, 2016.
- [24] I. Dolinka and J. East. Semigroups of rectangular matrices under a sandwich operation. (preprint 2015).
- [25] I. Dolinka and J. East. The idempotent generated subsemigroup of the Kauffman monoid. (preprint 2016).
- [26] I. Dolinka and J. East. Twisted Brauer monoids. (preprint 2016).
- [27] I. Dolinka, J. East, A. Evangelou, D. FitzGerald, N. Ham, J. Hyde, and N. Loughlin. Enumeration of idempotents in diagram semigroups and algebras. *J. Combin. Theory Ser. A*, 131:119–152, 2015.
- [28] I. Dolinka, J. East, A. Evangelou, D. FitzGerald, N. Ham, J. Hyde, and N. Loughlin. Idempotent statistics of the Motzkin and Jones monoids. (preprint 2015).
- [29] I. Dolinka, J. East, and R. D. Gray. Motzkin monoids and partial brauer monoids. (preprint 2016).
- [30] I. Dolinka, J. East, and J. Mitchell. Idempotent rank in the endomorphism monoid of a nonuniform partition. *Bull. Aust. Math. Soc.*, 93(1):73–91, 2016.
- [31] I. Dolinka and R. D. Gray. Maximal subgroups of free idempotent generated semigroups over the full linear monoid. *Trans. Amer. Math. Soc.*, 366(1):419–455, 2014.
- [32] M. P. Drazin. Regular semigroups with involution. In *Proc. Symp. on Regular Semigroups (DeKalb, 1979)*, Northern Illinois University, DeKalb, pages 29–46. 1979.
- [33] D. Easdown. Biorordered sets come from semigroups. *J. Algebra*, 96(2):581–591, 1985.
- [34] D. Easdown, M. V. Sapir, and M. V. Volkov. Periodic elements of the free idempotent generated semigroup on a biorordered set. *Internat. J. Algebra Comput.*, 20(2):189–194, 2010.
- [35] J. East. Cellular algebras and inverse semigroups. *J. Algebra*, 296(2):505–519, 2006.
- [36] J. East. Generators and relations for partition monoids and algebras. *J. Algebra*, 339:1–26, 2011.
- [37] J. East. On the singular part of the partition monoid. *Internat. J. Algebra Comput.*, 21(1-2):147–178, 2011.
- [38] J. East. Defining relations for idempotent generators in finite full transformation semigroups. *Semigroup Forum*, 86(3):451–485, 2013.
- [39] J. East. Defining relations for idempotent generators in finite partial transformation semigroups. *Semigroup Forum*, 89(1):72–76, 2014.
- [40] J. East. Infinite partition monoids. *Internat. J. Algebra Comput.*, 24(4):429–460, 2014.

- [41] J. East. Partition monoids and embeddings in 2-generator regular $*$ -semigroups. *Period. Math. Hungar.*, 69(2):211–221, 2014.
- [42] J. East. A symmetrical presentation for the singular part of the symmetric inverse monoid. *Algebra Universalis*, 74(3-4):207–228, 2015.
- [43] J. East. Presentations for rook partition monoids and algebras and their singular ideals. (preprint 2016).
- [44] J. East. Presentations for (singular) partition monoids: a new approach. (preprint 2016).
- [45] J. East, A. Egri-Nagy, J. Mitchell, and Y. Péresse. Computing finite semigroups. (preprint 2015).
- [46] J. East and D. G. FitzGerald. The semigroup generated by the idempotents of a partition monoid. *J. Algebra*, 372:108–133, 2012.
- [47] J. Enyang. A seminormal form for partition algebras. *J. Combin. Theory Ser. A*, 120(7):1737–1785, 2013.
- [48] K. Erdmann and R. M. Green. On representations of affine Temperley-Lieb algebras. II. *Pacific J. Math.*, 191(2):243–273, 1999.
- [49] J. A. Erdos. On products of idempotent matrices. *Glasgow Math. J.*, 8:118–122, 1967.
- [50] D. G. Fitz-Gerald. On inverses of products of idempotents in regular semigroups. *J. Austral. Math. Soc.*, 13:335–337, 1972.
- [51] D. G. FitzGerald and K. W. Lau. On the partition monoid and some related semigroups. *Bull. Aust. Math. Soc.*, 83(2):273–288, 2011.
- [52] J. Fountain and A. Lewin. Products of idempotent endomorphisms of an independence algebra of finite rank. *Proc. Edinburgh Math. Soc. (2)*, 35(3):493–500, 1992.
- [53] J. Fountain and A. Lewin. Products of idempotent endomorphisms of an independence algebra of infinite rank. *Math. Proc. Cambridge Philos. Soc.*, 114(2):303–319, 1993.
- [54] O. Ganyushkin, V. Mazorchuk, and B. Steinberg. On the irreducible representations of a finite semigroup. *Proc. Amer. Math. Soc.*, 137(11):3585–3592, 2009.
- [55] G. U. Garba. Idempotents in partial transformation semigroups. *Proc. Roy. Soc. Edinburgh Sect. A*, 116(3-4):359–366, 1990.
- [56] C. Godsil and G. Royle. *Algebraic graph theory*, volume 207 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2001.
- [57] G. M. S. Gomes and J. M. Howie. On the ranks of certain finite semigroups of transformations. *Math. Proc. Cambridge Philos. Soc.*, 101(3):395–403, 1987.
- [58] G. M. S. Gomes and J. M. Howie. On the ranks of certain semigroups of order-preserving transformations. *Semigroup Forum*, 45(3):272–282, 1992.
- [59] G. M. S. Gomes and N. Ruškuc. John Macintosh Howie: work and legacy. *Semigroup Forum*, 89(1):2–19, 2014.
- [60] F. M. Goodman and H. Wenzl. The Temperley-Lieb algebra at roots of unity. *Pacific J. Math.*, 161(2):307–334, 1993.
- [61] V. Gould and D. Yang. Every group is a maximal subgroup of a naturally occurring free idempotent generated semigroup. *Semigroup Forum*, 89(1):125–134, 2014.
- [62] J. J. Graham and G. I. Lehrer. Cellular algebras. *Invent. Math.*, 123(1):1–34, 1996.
- [63] J. J. Graham and G. I. Lehrer. The representation theory of affine Temperley-Lieb algebras. *Enseign. Math. (2)*, 44(3-4):173–218, 1998.
- [64] R. L. Graham. On finite 0-simple semigroups and graph theory. *Math. Systems Theory*, 2:325–339, 1968.
- [65] R. Gray. Idempotent rank in endomorphism monoids of finite independence algebras. *Proc. Roy. Soc. Edinburgh Sect. A*, 137(2):303–331, 2007.
- [66] R. Gray. Hall’s condition and idempotent rank of ideals of endomorphism monoids. *Proc. Edinb. Math. Soc. (2)*, 51(1):57–72, 2008.
- [67] R. Gray and N. Ruškuc. Generating sets of completely 0-simple semigroups. *Comm. Algebra*, 33(12):4657–4678, 2005.
- [68] R. Gray and N. Ruškuc. Maximal subgroups of free idempotent-generated semigroups over the full transformation monoid. *Proc. Lond. Math. Soc. (3)*, 104(5):997–1018, 2012.
- [69] R. Gray and N. Ruskuc. On maximal subgroups of free idempotent generated semigroups. *Israel J. Math.*, 189:147–176, 2012.
- [70] R. D. Gray. The minimal number of generators of a finite semigroup. *Semigroup Forum*, 89(1):135–154, 2014.
- [71] U. Grimm and S. O. Warnaar. Solvable RSOS models based on the dilute BWM algebra. *Nuclear Phys. B*, 435(3):482–504, 1995.
- [72] C. Grood. Brauer algebras and centralizer algebras for $SO(2n, \mathbb{C})$. *J. Algebra*, 222(2):678–707, 1999.
- [73] N. Guay and S. Wilcox. Almost cellular algebras. *J. Pure Appl. Algebra*, 219(9):4105–4116, 2015.

- [74] X. Guo and C. Xi. Cellularity of twisted semigroup algebras. *J. Pure Appl. Algebra*, 213(1):71–86, 2009.
- [75] P. Hall. *The collected works of Philip Hall*. Oxford Science Publications. The Clarendon Press Oxford University Press, New York, 1988. Compiled and with a preface by K. W. Gruenberg and J. E. Roseblade, With an obituary by Roseblade.
- [76] T. Halverson. Characters of the partition algebras. *J. Algebra*, 238(2):502–533, 2001.
- [77] T. Halverson and E. delMas. Representations of the Rook-Brauer algebra. *Comm. Algebra*, 42(1):423–443, 2014.
- [78] T. Halverson and T. Lewandowski. RSK insertion for set partitions and diagram algebras. *Electron. J. Combin.*, 11(2):Research Paper 24, 24 pp. (electronic), 2004/06.
- [79] T. Halverson and A. Ram. Partition algebras. *European J. Combin.*, 26(6):869–921, 2005.
- [80] T. Halverson and N. Thiem. q -partition algebra combinatorics. *J. Combin. Theory Ser. A*, 117(5):507–527, 2010.
- [81] P. Hanlon and D. Wales. A tower construction for the radical in Brauer’s centralizer algebras. *J. Algebra*, 164(3):773–830, 1994.
- [82] R. Hartmann, A. Henke, S. Koenig, and R. Paget. Cohomological stratification of diagram algebras. *Math. Ann.*, 347(4):765–804, 2010.
- [83] C. H. Houghton. Completely 0-simple semigroups and their associated graphs and groups. *Semigroup Forum*, 14(1):41–67, 1977.
- [84] J. M. Howie. The subsemigroup generated by the idempotents of a full transformation semigroup. *J. London Math. Soc.*, 41:707–716, 1966.
- [85] J. M. Howie. Idempotent generators in finite full transformation semigroups. *Proc. Roy. Soc. Edinburgh Sect. A*, 81(3-4):317–323, 1978.
- [86] J. M. Howie. *Fundamentals of semigroup theory*. Academic Press [Harcourt Brace Jovanovich Publishers], London, 1995. L.M.S. Monographs, No. 7.
- [87] J. M. Howie and R. B. McFadden. Idempotent rank in finite full transformation semigroups. *Proc. Roy. Soc. Edinburgh Sect. A*, 114(3-4):161–167, 1990.
- [88] J. T. Hyde, N. J. Loughlin, M. Quick, N. Ruskuc, and A. R. Wallis. On the growth of generating sets for direct powers of semigroups. *Semigroup Forum*, 84:116–130, 2012.
- [89] V. F. R. Jones. The Potts model and the symmetric group. In *Subfactors (Kyuzeso, 1993)*, pages 259–267. World Sci. Publ., River Edge, NJ, 1994.
- [90] V. F. R. Jones. A quotient of the affine Hecke algebra in the Brauer algebra. *Enseign. Math. (2)*, 40(3-4):313–344, 1994.
- [91] S. König and C. Xi. On the structure of cellular algebras. In *Algebras and modules, II (Geiranger, 1996)*, volume 24 of *CMS Conf. Proc.*, pages 365–386. Amer. Math. Soc., Providence, RI, 1998.
- [92] S. König and C. Xi. When is a cellular algebra quasi-hereditary? *Math. Ann.*, 315(2):281–293, 1999.
- [93] S. König and C. Xi. A characteristic free approach to Brauer algebras. *Trans. Amer. Math. Soc.*, 353(4):1489–1505, 2001.
- [94] G. Kudryavtseva and V. Mazorchuk. On presentations of Brauer-type monoids. *Cent. Eur. J. Math.*, 4(3):413–434 (electronic), 2006.
- [95] T. J. Laffey. Products of idempotent matrices. *Linear and Multilinear Algebra*, 14(4):309–314, 1983.
- [96] K. W. Lau and D. G. FitzGerald. Ideal structure of the Kauffman and related monoids. *Comm. Algebra*, 34(7):2617–2629, 2006.
- [97] I. Levi and S. Seif. Combinatorial techniques for determining rank and idempotent rank of certain finite semigroups. *Proc. Edinb. Math. Soc. (2)*, 45(3):617–630, 2002.
- [98] I. Levi and S. Seif. Counting techniques to label constant weight Gray codes with links to minimal generating sets of semigroups. *J. Algebra*, 266(1):220–238, 2003.
- [99] C. H. C. Little. A characterization of convertible $(0,1)$ -matrices. *J. Combinatorial Theory Ser. B*, 18:187–208, 1975.
- [100] V. Maltcev and V. Mazorchuk. Presentation of the singular part of the Brauer monoid. *Math. Bohem.*, 132(3):297–323, 2007.
- [101] P. Martin. *Potts models and related problems in statistical mechanics*, volume 5 of *Series on Advances in Statistical Mechanics*. World Scientific Publishing Co. Inc., Teaneck, NJ, 1991.
- [102] P. Martin. Temperley-Lieb algebras for nonplanar statistical mechanics—the partition algebra construction. *J. Knot Theory Ramifications*, 3(1):51–82, 1994.
- [103] P. Martin. The structure of the partition algebras. *J. Algebra*, 183(2):319–358, 1996.
- [104] P. Martin and V. Mazorchuk. Partitioned binary relations. *Math. Scand.*, 113(1):30–52, 2013.
- [105] P. Martin and V. Mazorchuk. On the representation theory of partial Brauer algebras. *Q. J. Math.*, 65(1):225–247, 2014.
- [106] P. Martin and D. Woodcock. On central idempotents in the partition algebra. *J. Algebra*, 217(1):156–169, 1999.

- [107] P. P. Martin. All direct sum representations of the Temperley-Lieb algebra. *J. Phys. A*, 20(8):L539–L542, 1987.
- [108] P. P. Martin. Representations of graph Temperley-Lieb algebras. *Publ. Res. Inst. Math. Sci.*, 26(3):485–503, 1990.
- [109] P. P. Martin. Temperley-Lieb algebras and the long distance properties of statistical mechanical models. *J. Phys. A*, 23(1):7–30, 1990.
- [110] P. P. Martin and A. Elgamal. Ramified partition algebras. *Math. Z.*, 246(3):473–500, 2004.
- [111] A. Mathas. *Iwahori-Hecke algebras and Schur algebras of the symmetric group*, volume 15 of *University Lecture Series*. American Mathematical Society, Providence, RI, 1999.
- [112] R. May. Cell algebras. *J. Algebra*, 425:107–132, 2015.
- [113] R. D. May. Cell algebra structures on monoid and twisted monoid algebras. (preprint 2015).
- [114] V. Mazorchuk. On the structure of Brauer semigroup and its partial analogue. *Problems in Algebra*, 13:29–45, 1998. Available at <http://www2.math.uu.se/mazor/papers.html>.
- [115] V. Mazorchuk. Endomorphisms of \mathfrak{B}_n , $\mathcal{P}\mathfrak{B}_n$, and \mathfrak{C}_n . *Comm. Algebra*, 30(7):3489–3513, 2002.
- [116] J. D. Mitchell. The Semigroups package for GAP, Version 2.8.0. <http://tinyurl.com/semigroups>, 2016.
- [117] W. D. Munn. On semigroup algebras. *Proc. Cambridge Philos. Soc.*, 51:1–15, 1955.
- [118] W. D. Munn. Matrix representations of semigroups. *Proc. Cambridge Philos. Soc.*, 53:5–12, 1957.
- [119] K. S. S. Nambooripad. Structure of regular semigroups. I. *Mem. Amer. Math. Soc.*, 22(224):vii+119, 1979.
- [120] K. S. S. Nambooripad and F. J. C. M. Pastijn. Regular involution semigroups. In *Semigroups (Szeged, 1981)*, volume 39 of *Colloq. Math. Soc. János Bolyai*, pages 199–249. North-Holland, Amsterdam, 1985.
- [121] T. E. Nordahl and H. E. Scheiblich. Regular $*$ -semigroups. *Semigroup Forum*, 16(3):369–377, 1978.
- [122] F. Pan. Irreducible representations of Brauer algebras. *J. Phys. A*, 28(11):3139–3156, 1995.
- [123] L. Polák. A solution of the word problem for free $*$ -regular semigroups. *J. Pure Appl. Algebra*, 157(1):107–114, 2001.
- [124] I. S. Ponizovskii. On matrix representations of associative systems. *Mat. Sb. N.S.*, 38(80):241–260, 1956.
- [125] M. S. Putcha. Products of idempotents in algebraic monoids. *J. Aust. Math. Soc.*, 80(2):193–203, 2006.
- [126] J. Rhodes and B. Steinberg. *The q -theory of finite semigroups*. Springer Monographs in Mathematics. Springer, New York, 2009.
- [127] D. Ridout and Y. Saint-Aubin. Standard modules, induction and the structure of the Temperley-Lieb algebra. *Adv. Theor. Math. Phys.*, 18(5):957–1041, 2014.
- [128] N. Robertson, P. D. Seymour, and R. Thomas. Permanents, Pfaffian orientations, and even directed circuits. *Ann. of Math. (2)*, 150(3):929–975, 1999.
- [129] H. Rui. A criterion on the semisimple Brauer algebras. *J. Combin. Theory Ser. A*, 111(1):78–88, 2005.
- [130] H. Rui and M. Si. A criterion on the semisimple Brauer algebras. II. *J. Combin. Theory Ser. A*, 113(6):1199–1203, 2006.
- [131] H. N. V. Temperley and E. H. Lieb. Relations between the “percolation” and “colouring” problem and other graph-theoretical problems associated with regular planar lattices: some exact results for the “percolation” problem. *Proc. Roy. Soc. London Ser. A*, 322(1549):251–280, 1971.
- [132] R. Thomas. A survey of Pfaffian orientations of graphs. In *International Congress of Mathematicians. Vol. III*, pages 963–984. Eur. Math. Soc., Zürich, 2006.
- [133] V. V. Vazirani and M. Yannakakis. Pfaffian orientations, 0-1 permanents, and even cycles in directed graphs. *Discrete Appl. Math.*, 25(1-2):179–190, 1989. *Combinatorics and complexity* (Chicago, IL, 1987).
- [134] H. Wenzl. On the structure of Brauer’s centralizer algebras. *Ann. of Math. (2)*, 128(1):173–193, 1988.
- [135] B. W. Westbury. The representation theory of the Temperley-Lieb algebras. *Math. Z.*, 219(4):539–565, 1995.
- [136] S. Wilcox. Cellularity of diagram algebras as twisted semigroup algebras. *J. Algebra*, 309(1):10–31, 2007.
- [137] E. M. Wright. The number of irreducible tournaments. *Glasgow Math. J.*, 11:97–101, 1970.
- [138] C. Xi. Partition algebras are cellular. *Compositio Math.*, 119(1):99–109, 1999.