

Association of flavonoid-rich foods and flavonoids with risk of all-cause mortality

Kerry L Ivey PhD^{1, 2, 6}, Majken K Jensen PhD^{1,3}, Jonathan M Hodgson PhD⁴,

A Heather Eliassen ScD^{2,3}, Aedín Cassidy PhD⁵, Eric B Rimm ScD^{1,2,3}.

Author affiliations: *Harvard T. H. Chan School of Public Health Departments of Nutrition¹, and Epidemiology²; Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School³; University of Western Australia School of Medicine and Pharmacology⁴; Department of Nutrition & Preventive Medicine, Norwich Medical School, University of East Anglia, Norwich, UK⁵. South Australian Health and Medical Research Institute, Infection and Immunity Theme, School of Medicine, Flinders University, Adelaide, Australia⁶.*

Address for correspondence: *Kerry L Ivey: Department of Nutrition, Harvard T. H. Chan School of Public Health, 677 Huntington Ave, 02115, Boston, MA, USA;*

Email: kivey@hsph.harvard.edu.

Running title: *Flavonoid-rich foods and mortality*

Key words: *Flavonoids; mortality; cancer; cardiovascular disease; red wine; tea*

1 ABSTRACT

2 Flavonoids are bioactive compounds found in foods such as tea, red wine, fruits, and vegetables.
3 Higher intakes of specific flavonoids, and flavonoid rich foods, have been linked to reduced
4 mortality from specific vascular diseases and cancers. However, the importance of flavonoid-rich
5 foods, and flavonoids, in preventing all-cause mortality remains uncertain. As such, we
6 examined the association of intake of flavonoid-rich foods and flavonoids with subsequent
7 mortality among 93,145 young and middle-aged women in the Nurses' Health Study II. During
8 1,838,946 person-years of follow-up, 1,808 participants died. When compared to non-consumers,
9 frequent consumers of red wine, tea, peppers, blueberries, and strawberries were at reduced risk
10 of all-cause mortality ($P<0.05$), with the strongest associations observed for red wine and tea;
11 multivariable-adjusted hazard ratios (95%CI): 0.60 (0.49-0.74), and 0.73 (0.65-0.83),
12 respectively. Conversely, frequent grapefruit consumers were at increased risk of all-cause
13 mortality, compared to their non-grapefruit consuming counterparts ($P<0.05$). When compared to
14 those in the lowest consumption quintile, participants in the highest quintile of total-flavonoid
15 intake were at reduced risk of all-cause mortality in the age-adjusted model; 0.81 (0.71-0.93).
16 However, this association was attenuated following multivariable-adjustment; 0.92 (0.80-1.06).
17 Similar results were observed for consumption of flavan-3-ols, proanthocyanins and
18 anthocyanins. Flavonols, flavanones and flavones were not associated with all-cause mortality in
19 any model. Despite null associations at the compound level and select foods, higher consumption
20 of red wine, tea, peppers, blueberries and strawberries, was associated with reduced risk of total
21 and cause-specific mortality. These findings support the rationale for making food-based dietary
22 recommendations.

23 INTRODUCTION

24 Flavonoids represent a structurally diverse group of polyphenolic compounds which are
25 synthesized during plant metabolism¹ and are present in many commonly consumed foods².

26 Particular fruits and vegetables, such as blueberries, apples, spinach and onions are considered
27 rich sources of flavonoids. So too are beverages, such as tea and red wine³.

28 Meta-analyses have shown that consumption of flavonoid-rich foods are associated with reduced
29 risk of cause-specific mortalities such as those attributable to cancer, diabetes, and
30 cardiovascular disease⁴⁻⁷. Much of the beneficial effects of these foods have been attributed to
31 their high content of biologically active flavonoids, which have been shown to improve nitric
32 oxide homeostasis and endothelial function, and reduce platelet aggregation and oxidative
33 stress⁸⁻¹³. Flavonoids are also thought to play roles in inactivating carcinogens, inducing
34 antiproliferation, cell cycle arrest and apoptosis, and inhibiting angiogenesis¹⁴.

35 Despite this growing evidence for reduced risk of specific cancer and cardiovascular disease
36 mortalities associated with consumption of flavonoid rich foods, the relationship of flavonoid-
37 rich food and flavonoid compound intake with risk of all-cause mortality is less clear. Following
38 the advent of the comprehensive United States Department of Agriculture (USDA) flavonoid-
39 class food content database in 2007¹⁵, the few studies exploring the relation of flavonoid intake
40 with all-cause mortality have yielded inconsistent patterns of association, likely due to
41 insufficient sample size, limited variation in intake, or incomplete food composition table¹⁶⁻¹⁸.

42 The varying levels of flavonoid-class intake and different patterns of flavonoid-class intake
43 across countries¹⁹ have also likely contributed to the lack of clarity regarding the relationship
44 between flavonoid-compound intake and risk of all-cause mortality in the population.

45 Furthermore, the role that the whole-food, as distinct from the isolated food-constituent, plays in

46 the relationship with all-cause mortality is yet to be elucidated. We have recently shown that a
47 higher intake of total-flavonoids, as well as individual flavonoid classes, was associated with
48 lower risk of cardiovascular, cancer and all-cause mortality in elderly, postmenopausal women¹⁶.
49 Therefore, this study sought to explore the relationship between the consumption of flavonoid
50 rich foods, and flavonoid-compounds, and the risk of all-cause mortality in a cohort of young
51 and middle-aged US women.

52 PARTICIPANTS AND METHODS

53 *Participants*

54 In 1989, 116 430 women aged between 25 and 42 years, were enrolled into the Nurses' Health
55 Study II. Baseline for this analysis was 1991, where 93 145 participants had complete dietary
56 intake data and were free of previous myocardial infarction, angina, stroke and cancer. The
57 institutional review board at Brigham and Women's Hospital reviewed and approved this study,
58 and participants provided implied consent by virtue of questionnaire return.

59 *Mortality ascertainment*

60 Mortality incidence were identified through reports from family members and postal authorities,
61 allowing ascertainment of an estimated 98% or more of all deaths²⁰. Further mortality cases
62 were identified through the National Death Index. Using data from death certificates and medical
63 records, a physician blinded to exposure intake classified causes of mortality according to the
64 eighth and ninth revisions of the International Classification of Diseases (ICD)^{21, 22}.

65 Deaths attributable to cardiovascular disease were defined using the ICD8 codes 390-458 (ICD9
66 390-459) and cancer mortalities were those with ICD8 codes ranging from 140 to 207 (ICD9
67 140-208). The other-cause mortality variable refers to all mortalities not attributable to CVD or
68 cancer based on ICD8 codes.

69 *Dietary intake assessment*

70 At baseline (1991) and every subsequent 4 years until 2007, participants completed a semi-
71 quantitative food-frequency questionnaire (FFQ). From this, habitual daily intake, in mg/d, of
72 total-flavonoids and flavonoid-classes was estimated using previously described methods²³.

73 Flavonoid-classes in this analysis include: *i*) flavonols; *ii*) flavan-3-ols (including catechins and
74 epicatechins, and excluding proanthocyanins); *iii*) proanthocyanins; *iv*) flavones; *v*) flavanones;
75 and *vi*) anthocyanins. Frequency of consumption of flavonoid-rich foods were recorded as
76 number of servings per day, week, or month²⁴.

77 As an indicator of adherence to a healthy dietary pattern, the Alternative Healthy Eating Index
78 score was calculated using methods previously described²⁵.

79 In order to reflect long-term dietary intake, and to minimize effects of within-person variation,
80 flavonoid exposure was considered the cumulative average of flavonoid intake, updated with
81 every 4-year FFQ return. To account for potential alterations in dietary patterns following a
82 major illness diagnosis, the primary flavonoid exposure was computed by suspending dietary
83 intake updates following reported diagnoses of stroke, heart disease, angina, or cancer, although
84 follow-up continued until death or the end of the study period at 2009.

85 *Risk factor assessment*

86 At baseline, and every two years thereafter, participants completed questionnaires on lifestyle,
87 medical conditions, medications and family medical history.

88 *Statistical Analysis*

89 Analyses for habitual consumption of flavonoid-rich foods based on categories of consumption
90 from the FFQ; ranging from non-consumers to frequent consumers, as defined as consuming the
91 food more than once per week. Exposure of total-flavonoid or flavonoid-class consumption was
92 divided into quintiles. Hazard ratios (HR) and 95% confidence intervals for risk of all-cause, and
93 cause-specific, mortalities were estimated using age-adjusted and multivariable-adjusted Cox
94 proportional-hazards models. P values for trend were calculated with the use of the Wald test of

95 a score variable based on the median consumption level for each quintile of flavonoid
96 consumption.

97 The multivariable-adjusted model included age, body mass index (BMI), smoking status,
98 menopausal status, family history of diabetes/cancer/myocardial infarction, multivitamin
99 supplement use, aspirin use, race, type 2 diabetes, hypercholesterolemia, hypertension, physical
100 activity, alcohol consumption, and caloric intake. The multivariable plus diet-adjusted model
101 incorporated the multivariable-adjusted model plus the Alternative Healthy Eating Index (minus
102 alcohol) score²⁶.

103 For sensitivity analyses, baseline flavonoid intake and unrestricted cumulative average flavonoid
104 intake, where updates continued until death or end of study irrespective of chronic disease
105 diagnosis, were also computed. To address the concern that occult chronic diseases in the years
106 that preceded diagnosis may have influenced dietary intake, we excluded the first 2 years of
107 follow-up data and added a 2-year lag period between flavonoid-intake assessment and each
108 follow-up period.

109 We conducted several additional sensitivity analyses to assess the robustness of the results. To
110 minimize the influence of smoking or an extremely low or high body-mass index on the results,
111 we excluded participants who had ever smoked or who had a BMI of less than 18.5 or more than
112 40 kg/m². We also excluded participants who had diabetes at baseline, and we suspended
113 updating of dietary variables after a diagnosis of diabetes during study follow-up.

114 Analyses were performed with the SAS statistical package (version 9.3, SAS Institute).

115 Statistical tests were two-sided, and P values of less than 0.05 were considered to indicate
116 statistical significance.

117 RESULTS

118 *Cohort characteristics*

119 At baseline, the mean age of participants was 36.1 (± 4.7) years, with a mean BMI of 24.6 (± 5.3)
120 kg/m². Over the 18-year (1 838 904 person-year) follow-up, there were 1 894 deaths. Cancer was
121 the leading cause of mortality in this cohort, accounting for 47% (n=887) of all deaths.

122 Cardiovascular disease contributed 10% (n=189) to all follow-up mortalities, and the remaining
123 818 (43%) mortalities comprised the other cause mortality group. **The majority of other-cause**
124 **mortalities were due to infections (n=182, ICD8 000-136), diseases of other endocrine glands**
125 **(n=192, ICD8 250-258), and diseases of the nervous system (n=315, ICD8 320-358).**

126 Mean daily total-flavonoid consumption was 379 (± 374) mg. **Proanthocyanins** contribute 57%,
127 **and flavan-3-ols** 28%, to total-flavonoid intake (**Table 1**).

128 Participants were similar in terms of baseline risk factors across all levels of total-flavonoid
129 consumption. However, high flavonoid consumers were more physically active and were less
130 likely to be current smokers at baseline (**Table 2**).

131 *Flavonoid-rich foods and risk of all-cause, and cause specific, mortality*

132 We explored potential whole-food contributors by analyzing foods rich in the flavonoid-classes
133 (**Figure 1**). Frequent consumption of blueberries, strawberries, apples, peppers, red-wine and tea
134 were all significantly inversely associated with risk of all-cause mortality in age-adjusted,
135 multivariable-adjusted and multivariable-plus-diet- adjusted models. When compared to non-
136 consumers, frequent tea and red wine consumption showed the greatest magnitude of reduction
137 in risk. Conversely, when compared to infrequent consumers, risk of all-cause mortality was
138 greater in participants with frequent grapefruit consumption. **Continuing to update of intake**

139 **irrespective of chronic disease diagnosis did not substantially impact results, and results were**
140 similar in all sensitivity analyses.

141 In cause-specific age and multivariable-adjusted analyses (**Table 3**), when compared to non-
142 consumers, the benefit of frequent blueberry and strawberry consumption was restricted to
143 cancer mortality, and the benefit of peppers restricted to mortalities from other-causes.

144 Although not associated with all-cause mortality, frequent orange-fruit consumers were at
145 reduced risk of cancer and other-cause mortalities, respectively. Congruent with the all-cause
146 mortality results, we observed that frequent grapefruit consumers were at increased risk of
147 mortalities from other causes.

148 Both red wine and tea showed the greatest magnitude of benefit in the all-cause mortality
149 analyses. Specifically, in the multivariable-adjusted model, when compared to the non-
150 consumers, the relative risk (95% CI) of all-cause mortality for frequent consumers of red wine
151 and tea (more than once per week) was 0.60 (0.49, 0.74), and 0.73 (0.65, 0.83), respectively.

152 When looking at cause-specific mortalities, frequent consumption of red wine and tea was
153 associated with reduced risk of both cancer and other-cause mortalities, in both age-adjusted and
154 multivariable-adjusted models. Results were not significantly altered in sensitivity analyses.

155 ***Flavonoid compounds and risk of all-cause, and cause specific, mortality***

156 In age-adjusted models, participants in the highest quintile of total-flavonoid consumption were
157 19% (7-29%) less likely to have died in the 18-year follow-up period, when compared to those in
158 the lowest quintile (**Table 4**). Similar beneficial associations were observed with increased
159 consumption of flavan-3-ols, flavonols, flavones, proanthocyanins and anthocyanins, however,
160 relationships were attenuated and no longer statistically significant following multivariable-

161 adjustment. Despite multivariable-adjustment substantially attenuating the relationships for
162 proanthocyanins and anthocyanidins, no one factor in the multivariable-adjusted model was
163 responsible for attenuation of the findings. Results were not significantly altered in sensitivity
164 analyses.

165 We then examined the relation of individual flavonoid classes with cause-specific mortality. In
166 age-adjusted models, when compared to the lowest quintile, participants in the highest quintile of
167 anthocyanin intake were at lower risk of mortality from cancer, cardiovascular disease and other-
168 causes. This beneficial association remained for cancer mortalities following multivariable-
169 adjustment (**Table 5**). Flavan-3-ols and proanthocyanins followed a similar inverse pattern,
170 whereas flavonols, flavanones and flavones showed no association in either the unadjusted or
171 multivariable-adjusted models, with any of the mortality types.

172 DISCUSSION

173 This prospective cohort study of middle-aged US women found that participants with higher
174 intakes of specific flavonoid-rich foods, namely blueberries, strawberries, peppers, red wine and
175 tea, were associated with reduced risk of all-cause mortality. When exploring contributors to
176 these relationships, the association with all-cause mortality appeared to be largely driven by
177 mortalities from cancer, as well as other causes. These beneficial relations did not extend to the
178 other flavonoid-rich foods, or intakes of flavonoid compounds.

179 Despite null associations at a compound level, numerous significant associations with all-cause
180 mortality were observed for many flavonoid-rich foods. Our finding of a null association of total-
181 flavonoid intake with risk of all-cause mortality in US women is congruent with the Iowa
182 Womens' Health Study¹⁷. However, in our previous analysis in Australian women¹⁶ we observed
183 a strong relation between increased total-flavonoid intake and reduced risk of all-cause mortality.
184 This incongruence in findings between studies, **and the differences we observed in associations**
185 **with compounds and whole foods**, is likely explained by the complexity of flavonoid intake
186 assessment and regional differences in the compositional variation in the whole food sources of
187 dietary flavonoids¹⁹, which in turn shapes the pattern of over 4,000 different flavonoid
188 compounds consumed on a daily basis²⁷.

189 When looking at whole-food associations, we observed that increased consumption of
190 blueberries, strawberries, peppers, red wine and tea was associated with reduced risk of all-cause
191 mortality. These associations remained after adjusting for dietary pattern, suggesting that the
192 relations are not explained by their contribution to a healthy dietary pattern. Furthermore, our
193 results are supported by clinical trial data showing effects of these foods in improving
194 endothelial function, nitric oxide status, blood pressure and platelet function, and by reducing

195 oxidative stress and inflammation²⁹⁻³³. The strongest beneficial relation with all-cause mortality
196 was observed with the frequency of red wine consumption, which remained even after adjusting
197 for total alcohol consumption, which has been shown to be a strong predictor of all-cause
198 mortality³⁴. When looking at cause-specific mortalities, the strongest associations for with red
199 wine were observed with reduced risk of cancer and other-cause mortalities, in both unadjusted
200 and multivariate-adjusted models. **The lack of beneficial association with cardiovascular disease
201 may be due to the cohort characteristics itself, namely the low cardiovascular disease mortality
202 rate in this middle aged female population.**

203 In contrast to the beneficial whole foods listed above, which are rich sources of flavan-3-ols,
204 proanthocyanins and anthocyanins, the foods rich in flavanones showed markedly different
205 results. Oranges showed no association with all-cause mortality, and grapefruit had a small
206 positive association with all-cause mortality. **This inverse association may be due to the
207 contribution of sugar-rich juices to total grapefruit intake. Furthermore, this detrimental
208 association** may also be explained by be due to the findings that grapefruit components have
209 clinically significant interactions with drugs, which appear to be independent of their flavonoid
210 content³⁵. However, this hypotheses were unable to be explored in this cohort.

211 **We observed that flavonoid-rich whole foods, and not flavonoid subclasses, showed the strongest
212 associations with risk of all-cause mortality. Although not reaching statistical significance, many
213 of the flavonoid subclasses followed similar trends to that of their predominant whole food
214 constituents. For example, the positive and null associations of flavanone-rich grapefruit and
215 oranges, respectively, were reflected in a non-significant trend in the multivariate-adjusted model
216 whereby high flavanone consumers tended to have higher mortality rates. Conversely, the
217 beneficial associations of anthocyanin-rich blueberries and strawberries was reflected in a no-**

218 significant observed trend high anthocyanin consumers tended to have lower mortality rates. The
219 role of the whole-food in influencing relationships has not yet been fully elucidated, and results
220 from the literature are conflicting. In understanding the strength of association differences at a
221 whole food level as opposed to a compound level, it is important to note that flavonoid intake
222 estimates are derived from intake data for many different individual food items, the majority of
223 which were not included in our study, which only looked at foods which contribute substantially
224 to flavonoid-class intake. The importance of whole foods, as opposed to isolated nutrients, are
225 becoming increasingly recognized for public health guidelines and dietary recommendations ²⁸.

226 Although results were not substantially altered by conducting sensitivity analyses, such as the lag
227 analyses, it is important to note that causality of observed relationships cannot be established due
228 to the observational nature of the study. Also, despite the inclusion of dietary and lifestyle factors
229 into statistical models, residual or unmeasured confounders cannot be ruled out. Identification of
230 causality is further limited by the complexity associated with assessing food composition and
231 dietary intake including for flavonoids³⁶, which further highlights the importance of conducting
232 both nutrient-based and whole-food based analyses.

233 In summary, in this prospective cohort study of female US nurses, we found a beneficial
234 relationship between the dietary intake of select whole-food sources of flavonoids and risk of
235 mortality. Specifically, frequent consumption of blueberries, strawberries, peppers, red wine and,
236 was associated with reduced risk of all-cause mortality. These beneficial associations did not
237 extend to total-flavonoids or flavonoid subclasses, and when considering the literature as a
238 whole, future prospective association studies are warranted.

239

240 **ACKNOWLEDGEMENTS**

241 We would like to thank the participants and staff of the Nurses' Health Study II for their valuable
242 contributions as well as the following state cancer registries for their help: AL, AZ, AR, CA, CO,
243 CT, DE, FL, GA, ID, IL, IN, IA, KY, LA, ME, MD, MA, MI, NE, NH, NJ, NY, NC, ND, OH,
244 OK, OR, PA, RI, SC, TN, TX, VA, WA, WY. The authors assume full responsibility for
245 analyses and interpretation of these data.

246 **AUTHOR CONTRIBUTIONS**

247 All authors were responsible for either drafting of the work or revising it critically for important
248 intellectual content, and provided final approval of the version to be published. KLI, AHE, EBR
249 made substantial contributions to conception or design of the work, or the acquisition, analysis,
250 or interpretation of data for the work.

251 **CONFLICT OF INTEREST STATEMENT**

252 EBR, KLI and AC received funds from the US Highbush Blueberry Research Council, for
253 activities unrelated to this research paper. No other disclosures were reported.

254 **SOURCES OF SUPPORT**

255 Components of the Nurses' Health Study 2 were supported by the following National Institute of
256 Health Grants: UM1 CA176726 and R01 CA67262. The Salary of Kerry L Ivey was supported
257 by a National Health and Medical Research Council Early Career Fellowship.

REFERENCES

1. Robards, K.; Prenzler, P. D.; Tucker, G.; Swatsitang, P.; Glover, W., Phenolic compounds and their role in oxidative processes in fruits. *Food Chemistry* **1999**, *66* (4), 401-436.
2. Beecher, G. R., Overview of dietary flavonoids: nomenclature, occurrence and intake. *The Journal of Nutrition* **2003**, *133* (10), 3248S-3254S.
3. Bhagwat, S., Haytowitz, D.B. Holden, J.M. , USDA database for the flavonoid content of selected foods, release 3.1. *U.S. Department of Agriculture, Agricultural Research Service. Nutrient Data Laboratory* **2014**.
4. Sun, C.-L.; Yuan, J.-M.; Koh, W.-P.; Mimi, C. Y., Green tea, black tea and breast cancer risk: a meta-analysis of epidemiological studies. *Carcinogenesis* **2006**, *27* (7), 1310-1315.
5. Tang, N.; Wu, Y.; Zhou, B.; Wang, B.; Yu, R., Green tea, black tea consumption and risk of lung cancer: a meta-analysis. *Lung Cancer* **2009**, *65* (3), 274-283.
6. Arab, L.; Liu, W.; Elashoff, D., Green and black tea consumption and risk of stroke. *Stroke* **2009**, *40* (5), 1786-1792.
7. Wang, Z.-M.; Zhou, B.; Wang, Y.-S.; Gong, Q.-Y.; Wang, Q.-M.; Yan, J.-J.; Gao, W.; Wang, L.-S., Black and green tea consumption and the risk of coronary artery disease: a meta-analysis. *American Journal of Clinical Nutrition* **2011**, *93* (3), 506-515.
8. Ishige, K.; Schubert, D.; Sagara, Y., Flavonoids protect neuronal cells from oxidative stress by three distinct mechanisms. *Free Radical Biology and Medicine* **2001**, *30* (4), 433-446.
9. Pignatelli, P.; Pulcinelli, F. M.; Celestini, A.; Lenti, L.; Ghiselli, A.; Gazzaniga, P. P.; Violi, F., The flavonoids quercetin and catechin synergistically inhibit platelet function by antagonizing the intracellular production of hydrogen peroxide. *The American Journal of Clinical Nutrition* **2000**, *72* (5), 1150-1155.
10. Loke, W. M.; Hodgson, J. M.; Proudfoot, J. M.; McKinley, A. J.; Puddey, I. B.; Croft, K. D., Pure dietary flavonoids quercetin and (-)-epicatechin augment nitric oxide products and reduce endothelin-1 acutely in healthy men. *American Journal of Clinical Nutrition* **2008**, *88* (4), 1018-1025.
11. Bondonno, C. P.; Yang, X.; Croft, K. D.; Considine, M. J.; Ward, N. C.; Rich, L.; Puddey, I. B.; Swinny, E.; Mubarak, A.; Hodgson, J. M., Flavonoid-rich apples and nitrate-rich spinach augment nitric oxide status and improve endothelial function in healthy men and women: a randomized controlled trial. *Free Radical Biology and Medicine* **2012**, *52* (1), 95-102.
12. Hodgson, J. M.; Puddey, I. B.; Burke, V.; Watts, G. F.; Beilin, L. J., Regular ingestion of black tea improves brachial artery vasodilator function. *Clinical Science* **2002**, *102* (2), 195-201.
13. Hodgson, J. M.; Burke, V.; Puddey, I. B., Acute effects of tea on fasting and postprandial vascular function and blood pressure in humans. *Journal of Hypertension* **2005**, *23* (1), 47-54.
14. Chahar, M. K.; Sharma, N.; Dobhal, M. P.; Joshi, Y. C., Flavonoids: A versatile source of anticancer drugs. In *Pharmacogn Rev*, 2011; Vol. 5, pp 1-12.

15. US Department of Agriculture, USDA database for the flavonoid content of selected foods; release 2.1. Maryland, 2007.
16. Ivey KL; Hodgson JM; Croft KD; Lewis JR; Prince RL, Flavonoid intake and all-cause mortality. *American Journal of Clinical Nutrition* **2015** 101 (5), 1012-1020.
17. Mink, P. J.; Scrafford, C. G.; Barraj, L. M.; Harnack, L.; Hong, C.-P.; Nettleton, J. A.; Jacobs, D. R., Flavonoid intake and cardiovascular disease mortality: a prospective study in postmenopausal women. *American Journal of Clinical Nutrition* **2007**, 85 (3), 895-909.
18. Zamora-Ros, R.; Jiménez, C.; Cleries, R.; Agudo, A.; Sánchez, M.-J.; Sánchez-Cantalejo, E.; Molina-Montes, E.; Navarro, C.; Chirlaque, M.-D.; Huerta, J. M., Dietary flavonoid and lignan intake and mortality in a Spanish cohort. *Epidemiology* **2013**, 24 (5), 726-733.
19. Zamora-Ros, R.; Knaze, V.; Luján-Barroso, L.; Romieu, I.; Scalbert, A.; Slimani, N.; Hjartåker, A.; Engeset, D.; Skeie, G.; Overvad, K., Differences in dietary intakes, food sources and determinants of total flavonoids between Mediterranean and non-Mediterranean countries participating in the European Prospective Investigation into Cancer and Nutrition (EPIC) study. *British Journal of Nutrition* **2013**, 109 (08), 1498-1507.
20. Rich-Edwards, J. W.; Corsano, K. A.; Stampfer, M. J., Test of the national death index and Equifax nationwide death search. *American Journal of Epidemiology* **1994**, 140 (11), 1016-1019.
21. Slee, V. N., The International Classification of Diseases: Ninth Revision (ICD-9). *Annals of Internal Medicine* **1978**, 88 (3), 424-426.
22. Moriyama, I., The eighth revision of the International Classification of Diseases. *American Journal of Public Health and the Nations Health* **1966**, 56 (8), 1277-1280.
23. Willett, W. C.; Sampson, L.; STAMPFER, M. J.; Rosner, B.; Bain, C.; Witschi, J.; Hennekens, C. H.; Speizer, F. E., Reproducibility and validity of a semiquantitative food frequency questionnaire. *American Journal of Epidemiology* **1985**, 122 (1), 51-65.
24. Cassidy, A.; O'Reilly, É. J.; Kay, C.; Sampson, L.; Franz, M.; Forman, J.; Curhan, G.; Rimm, E. B., Habitual intake of flavonoid subclasses and incident hypertension in adults. *The American Journal of Clinical Nutrition* **2011**, 93 (2), 338-347.
25. McCullough, M. L.; Willett, W. C., Evaluating adherence to recommended diets in adults: the Alternate Healthy Eating Index. *Public health nutrition* **2007**, 9 (1a), 152-157.
26. Chiuve, S. E.; Fung, T. T.; Rimm, E. B.; Hu, F. B.; McCullough, M. L.; Wang, M.; Stampfer, M. J.; Willett, W. C., Alternative dietary indices both strongly predict risk of chronic disease. *The Journal of Nutrition* **2012**, jn. 111.157222.
27. Cook, N.; Samman, S., Flavonoids: chemistry, metabolism, cardioprotective effects, and dietary sources. *Journal of Nutritional Biochemistry* **1996**, 7 (2), 66-76.
28. Mozaffarian, D.; Ludwig, D. S., Dietary Guidelines in the 21st Century—a Time for Food. *Journal of the American Medical Association* **2017**, 304 (6), 681-682.

29. Riso, P.; Klimis-Zacas, D.; Del Bo, C.; Martini, D.; Campolo, J.; Vendrame, S.; Møller, P.; Loft, S.; De Maria, R.; Porrini, M., Effect of a wild blueberry (*Vaccinium angustifolium*) drink intervention on markers of oxidative stress, inflammation and endothelial function in humans with cardiovascular risk factors. *European Journal of Nutrition* **2013**, *52* (3), 949-961.
30. Alvarez-Suarez, J. M.; Giampieri, F.; Tulipani, S.; Casoli, T.; Di Stefano, G.; González-Paramás, A. M.; Santos-Buelga, C.; Busco, F.; Quiles, J. L.; Cordero, M. D., One-month strawberry-rich anthocyanin supplementation ameliorates cardiovascular risk, oxidative stress markers and platelet activation in humans. *The Journal of Nutritional Biochemistry* **2014**, *25* (3), 289-294.
31. Diebolt, M.; Bucher, B.; Andriantsitohaina, R., Wine polyphenols decrease blood pressure, improve NO vasodilatation, and induce gene expression. *Hypertension* **2001**, *38* (2), 159-165.
32. Ras, R. T.; Zock, P. L.; Draijer, R., Tea consumption enhances endothelial-dependent vasodilation; a meta-analysis. *PLoS One* **2011**, *6* (3).
33. Spiller, F.; Alves, M. K.; Vieira, S. M.; Carvalho, T. A.; Leite, C. E.; Lunardelli, A.; Poloni, J. A.; Cunha, F. Q.; Oliveira, J. R., Anti-inflammatory effects of red pepper (*Capsicum baccatum*) on carrageenan- and antigen-induced inflammation. *Journal of Pharmacy and Pharmacology* **2008**, *60* (4), 473-478.
34. Di Castelnuovo, A.; Costanzo, S.; Bagnardi, V.; Donati, M. B.; Iacoviello, L.; de Gaetano, G., Alcohol dosing and total mortality in men and women: an updated meta-analysis of 34 prospective studies. *Archives of Internal Medicine* **2006**, *166* (22), 2437-2445.
35. Huang, S. M.; Hall, S. D.; Watkins, P.; Love, L. A.; Serabjit-Singh, C.; Betz, J. M.; Hoffman, F. A.; Honig, P.; Coates, P. M.; Bull, J., Drug interactions with herbal products and grapefruit juice: a conference report. *Clinical Pharmacology & Therapeutics* **2004**, *75* (1), 1-12.
36. Ivey, K. L.; Croft, K.; Prince, R. L.; Hodgson, J. M., Comparison of flavonoid intake assessment methods. *Food and Function* **2016**.

TABLES AND FIGURES

Table 1: Baseline total-flavonoid and flavonoid-class consumption, and frequency of consumption of major whole food contributors ¹

	Level of consumption (mg/d)	Major whole food contributors	Frequency of consumption ³		
			Never	Rarely	Frequent
Total-flavonoids	379 ± 374	Tea Apples Oranges ²	33.0 11.0 9.75	23.8 49.6 32.2	43.2 39.45 58.13
Flavonoid-classes					
Flavonols	19 ± 13	Tea Onions Apples		16.3 58.0	25.8
Flavan-3-ols	61 ± 82	Tea Apples Blueberries		65.9 31.9	2.25
Proanthcyanins	257 ± 278	Tea Apples Strawberries		22.0 68.1	9.9
Flavones	2 ± 1	Oranges ² Red wine Peppers		80.6 31.4	16.8 15.7
Flavanones	33 ± 33	Oranges ² Grapefruit ² Red wine		51.7 33.4	14.9
Anthocyanins	11 ± 14	Blueberries Strawberries Apples			

¹ Results are energy-adjusted mean ± SD. n = 93,145. ² Includes both fresh fruit and juice products.

³ Results presented as %. Rarely consumed defined as > never, and < once per day. Frequently consumed defined as ≥ once per day.

Table 2: Baseline characteristics of the cohort stratified by quintiles of total-flavonoid consumption ¹

	Quintile 1 < 150 mg/d	Quintile 3 222 - < 329 mg/d	Quintile 5 ≥ 587 mg/d
Number	18 617	18 612	18 651
Demographic variables			
Age (years) ²	36.0 ± 4.7	36.0 ± 4.7	36.5 ± 4.6
Body mass index (kg/m ²)	25.0 ± 5.7	24.4 ± 5.1	24.7 ± 5.3
Caucasian (%)	93.1	93.6	94.7
Current smoker (%)	18.1	9.9	11.3
Postmenopausal (%)	3.2	3.1	3.9
Physical activity (MET-hrs/wk) ³	16.8 ± 24.2	22.6 ± 28.1	21.6 ± 28.8
Prevalent disease			
Type 2 diabetes (%)	1.0	1.0	1.0
Hypercholesterolemia (%)	11.3	10.1	10.9
Hypertension (%)	6.4	6.0	7.0
Family history of disease			
Diabetes (%)	17.2	16.0	17.4
Myocardial infarction (%)	22.7	20.7	22.2
Cancer (%)	22.7	22.9	22.2
Dietary intake and medications			
Current Aspirin use (%)	11.7	10.7	12.0
Current multivitamin use (%)	38.6	47.1	42.7
Calorie intake (Kcal/d)	1 703.2 ± 540.8	1 851.4 ± 548.5	1 735.4 ± 565.7
Alcohol intake (g/d)	3.1 ± 6.4	3.3 ± 5.9	2.6 ± 5.6
AHEI (score) ⁴	40.8 ± 10.1	45.5 ± 10.4	44.9 ± 10.5

¹Results are mean ± SD or percentage where appropriate. Values standardized to the age distribution of the study population. Flavonoid consumption is standardized to total-energy intake. n = 93,145; ² Value is not age adjusted;

³ Met: metabolic equivalent; ⁴ AHEI: Alternative Healthy Eating Index (excluding alcohol) score.

Table 3: Multivariable-adjusted risk of mortality subtypes by flavonoid-rich foods, comparing non-consumers (referent group) to frequent consumers (more than once per week).¹

	Cancer mortality ³ ≥ once per week	CVD mortality ⁴ ≥ once per week	Other-cause ⁴ ≥ once per week
Orange fruit consumption			
Age-adjusted	0.75 (0.60-0.95)	0.66 (0.41-1.07)	0.79 (0.64-0.97)
Multivariable-adjusted ²	0.76 (0.60-0.98)	0.89 (0.54-1.48)	0.98 (0.78-1.24)
Orange juice consumption			
Age-adjusted	0.90 (0.72-1.13)	0.76 (0.48-1.21)	0.83 (0.68-1.03)
Multivariable-adjusted ²	0.96 (0.76-1.21)	0.89 (0.55-1.43)	0.97 (0.78-1.21)
Grapefruit consumption			
Age-adjusted	1.01 (0.83-1.22)	0.72 (0.48-1.08)	1.19 (1.00-1.42)
Multivariable-adjusted ²	1.07 (0.88-1.31)	0.92 (0.60-1.42)	1.47 (1.22-1.77)
Apple consumption			
Age-adjusted	0.63 (0.47-0.84)	0.74 (0.37-1.48)	0.60 (0.45-0.79)
Multivariable-adjusted ²	0.68 (0.50-0.93)	1.19 (0.58-2.45)	0.84 (0.63-1.14)
Blueberry consumption			
Age-adjusted	0.67 (0.50-0.89)	0.41 (0.19-0.89)	0.77 (0.60-1.01)
Multivariable-adjusted ²	0.64 (0.47-0.87)	0.64 (0.29-1.41)	1.00 (0.75-1.32)
Strawberry consumption			
Age-adjusted	0.69 (0.54-0.89)	0.49 (0.29-0.82)	0.65 (0.51-0.81)
Multivariable-adjusted ²	0.73 (0.56-0.95)	0.72 (0.41-1.24)	0.86 (0.67-1.10)
Onion consumption			
Age-adjusted	0.84 (0.65-1.08)	0.92 (0.53-1.62)	1.00 (0.78-1.28)
Multivariable-adjusted ²	0.83 (0.64-1.09)	0.95 (0.53-1.70)	1.05 (0.81-1.35)
Pepper consumption			
Age-adjusted	0.78 (0.63-0.96)	0.74 (0.47-1.17)	0.58 (0.47-0.71)
Multivariable-adjusted ²	0.80 (0.64-1.01)	1.04 (0.64-1.71)	0.67 (0.54-0.84)
Red wine consumption			
Age-adjusted	0.60 (0.46-0.78)	0.43 (0.22-0.82)	0.57 (0.44-0.74)
Multivariable-adjusted ²	0.53 (0.39-0.72)	0.74 (0.35-1.58)	0.65 (0.48-0.89)
Tea consumption			
Age-adjusted	0.67 (0.56-0.81)	0.62 (0.43-0.90)	0.71 (0.60-0.85)
Multivariable-adjusted ²	0.68 (0.56-0.82)	0.70 (0.48-1.02)	0.79 (0.66-0.95)

¹ Results are HR (95% CI) and n(%) where appropriate. n = 93,145. ² Multivariable adjusted model includes: age, body mass index, smoking status, menopausal status, family history of diabetes, cancer and myocardial infarction, multivitamin supplement use, Aspirin use, race, type 2 diabetes, hypercholesterolemia, hypertension, physical activity, caloric intake, alcohol consumption and the Alternative Health Eating Index (minus alcohol) score. ³ Total number of cancer mortalities: 887. ⁴ Total number of cardiovascular disease (CVD) mortalities: 189. ⁵ Total number of mortalities from other causes: 818.

Table 4: Association of quintiles of total-flavonoid and flavonoid-class consumption with risk of all-cause mortality ¹

	Quintile 1	Quintile 2	Quintile 3	Quintile 4	Quintile 5	P for trend
Total-flavonoid intake (mg/d)	< 138	138 - < 207	207 - < 308	308 - < 518	≥ 518	
Person-years	355 151	368 485	370 364	372 609	372 295	
Deaths (n)	433	390	323	351	397	
Age-adjusted	1.00 (referent)	0.85 (0.74-0.97)	0.69 (0.60-0.80)	0.73 (0.63-0.84)	0.81 (0.71-0.93)	0.10
Multivariable-adjusted ²	1.00 (referent)	0.96 (0.84-1.11)	0.81 (0.70-0.94)	0.86 (0.74-0.99)	0.92 (0.80-1.06)	0.59
Flavonol intake (mg/d)	< 9	9 - < 13	13 - < 17	17 - < 26	≥ 26	
Person-years	361 301	368 810	370 972	372 134	365 687	
Deaths (n)	398	347	345	356	448	
Age-adjusted	1.00 (referent)	0.83 (0.71-0.95)	0.82 (0.69-0.92)	0.80 (0.69-0.92)	1.00 (0.87-1.14)	0.030
Multivariable-adjusted ²	1.00 (referent)	0.90 (0.78-1.05)	0.89 (0.77-1.04)	0.88 (0.76-1.03)	1.08 (0.94-1.25)	0.006
Flavan-3-ol intake (mg/d)	< 12	12 - < 19	19 - < 39	39 - < 86	≥ 86	
Person-years	352 706	366 094	372 429	373 445	374 230	
Deaths (n)	428	399	335	335	397	
Age-adjusted	1.00 (referent)	0.88 (0.77-1.01)	0.72 (0.62-0.83)	0.71 (0.62-0.82)	0.82 (0.72-0.94)	0.30
Multivariable-adjusted ²	1.00 (referent)	0.99 (0.86-1.14)	0.81 (0.70-0.94)	0.79 (0.69-0.92)	0.90 (0.78-1.03)	0.30
Proanthocyanin intake (mg/d)	< 79	79 - < 126	126 - < 200	200 - < 356	≥ 356	
Person-years	352 476	367 583	372 126	373 157	373 562	
Deaths (n)	435	400	327	331	401	
Age-adjusted	1.00 (referent)	0.87 (0.76-1.00)	0.70 (0.61-0.81)	0.69 (0.60-0.80)	0.82 (0.72-0.94)	0.78
Multivariable-adjusted ²	1.00 (referent)	0.98 (0.85-1.13)	0.81 (0.70-0.94)	0.80 (0.69-0.93)	0.92 (0.80-1.05)	0.42
Flavone intake (mg/d)	< 0.7	0.7 - < 1.1	1.1 - < 1.5	1.5 - < 2.2	≥ 2.2	
Person-years	362 321	371 169	371 600	371 692	362 121	
Deaths (n)	404	381	369	335	405	
Age-adjusted	1.00 (referent)	0.90 (0.78-1.03)	0.84 (0.73-0.97)	0.75 (0.65-0.86)	0.89 (0.77-1.02)	0.007
Multivariable-adjusted ²	1.00 (referent)	1.00 (0.87-1.16)	1.00 (0.87-1.16)	0.92 (0.79-1.07)	1.11 (0.96-1.29)	0.96
Flavanone intake (mg/d)	< 9	9 - < 17	17 - < 30	30 - < 51	≥ 51	
Person-years	360 666	370 612	373 056	372 588	361 982	
Deaths (n)	373	378	389	380	374	
Age-adjusted	1.00 (referent)	0.98 (0.85-1.13)	0.98 (0.85-1.13)	0.94 (0.81-1.08)	0.92 (0.80-1.06)	0.75
Multivariable-adjusted ²	1.00 (referent)	1.07 (0.93-1.24)	1.11 (0.96-1.28)	1.11 (0.96-1.28)	1.11 (0.97-1.30)	0.015
Anthocyanin intake (mg/d)	< 3	3 - < 5	5 - < 9	9 - < 17	≥ 17	
Person-years	354 483	368 366	374 278	373 382	368 395	
Deaths (n)	456	407	331	336	364	
Age-adjusted	1.00 (referent)	0.85 (0.74-0.97)	0.67 (0.58-0.77)	0.67 (0.58-0.77)	0.71 (0.62-0.82)	<0.001
Multivariable-adjusted ²	1.00 (referent)	0.96 (0.84-1.10)	0.81 (0.70-0.94)	0.85 (0.73-0.99)	0.92 (0.79-1.08)	0.10

¹ Results are HR (95% CI) and n(%) where appropriate. n = 93,145. ² Multivariable adjusted model includes: age, body mass index, smoking status, menopausal status, family history of diabetes, cancer and myocardial infarction, multivitamin supplement use, Aspirin use, race, type 2 diabetes, hypercholesterolemia, hypertension, physical activity, caloric intake, alcohol consumption and the Alternative Health Eating Index (minus alcohol) score.

Table 5: Multivariable-adjusted risk of mortality subtypes by total flavonoid and flavonoid class, comparing participants in the lowest quintile of flavonoid intake (referent group) to those in the highest quintile.¹

	Cancer mortality ³ Quintile 5	CVD mortality ⁴ Quintile 5	Other-cause ⁵ Quintile 5
Total flavonoid intake			
Age-adjusted	0.80 (0.64-0.98)	0.66 (0.43-1.02)	0.86 (0.71-1.05)
Multivariable-adjusted ²	0.84 (0.67-1.04)	0.83 (0.53-1.29)	1.03 (0.84-1.26)
Flavonol intake			
Age-adjusted	0.98 (0.80-1.20)	0.76 (0.50-1.16)	1.08 (0.88-1.31)
Multivariable-adjusted ²	0.99 (0.80-1.24)	0.91 (0.58-1.41)	1.22 (0.99-1.51)
Flavan-3-ol intake			
Age-adjusted	0.84 (0.68-1.04)	0.64 (0.42-0.98)	0.85 (0.70-1.04)
Multivariable-adjusted ²	0.87 (0.70-1.08)	0.75 (0.49-1.16)	0.96 (0.78-1.17)
Proanthcyanin intake			
Age-adjusted	0.86 (0.70-1.06)	0.64 (0.41-0.99)	0.83 (0.68-1.00)
Multivariable-adjusted ²	0.90 (0.72-1.11)	0.77 (0.49-1.20)	0.97 (0.79-1.18)
Flavone intake			
Age-adjusted	0.89 (0.72-1.10)	0.74 (0.48-1.16)	0.91 (0.75-1.12)
Multivariable-adjusted ²	1.00 (0.80-1.06)	1.15 (0.72-1.83)	1.22 (0.99-1.51)
Flavanone intake			
Age-adjusted	0.91 (0.73-1.13)	0.81 (0.51-1.28)	0.95 (0.77-1.18)
Multivariable-adjusted ²	1.03 (0.83-1.29)	1.10 (0.69-1.76)	1.21 (0.97-1.50)
Anthocyanin intake			
Age-adjusted	0.74 (0.59-0.91)	0.48 (0.30-0.78)	0.74 (0.61-0.90)
Multivariable-adjusted ²	0.77 (0.61-0.98)	0.85 (0.50-1.43)	1.10 (0.88-1.37)

¹ Results are HR (95% CI) and n(%) where appropriate. n = 93,145). ² Multivariable adjusted model includes: age, body mass index, smoking status, menopausal status, family history of diabetes, cancer and myocardial infarction, multivitamin supplement use, Aspirin use, race, type 2 diabetes, hypercholesterolemia, hypertension, physical activity, caloric intake, alcohol consumption and the Alternative Health Eating Index (minus alcohol) score. ³ Total number of cancer mortalities: 887. ⁴ Total number of cardiovascular disease (CVD) mortalities: 189. ⁵ Total number of mortalities from other causes: 818.