

Burnout syndrome among psychiatric trainees in 22 countries: risk increased by long working hours, lack of supervision, and psychiatry not being first career choice

Jovanovic N¹, Podlesek A², Volpe U³, Barrett E⁴, Ferrari S⁵, Rojnic Kuzman M⁶, Wuyts P^{7,8}, Papp S⁹, Nawka A^{10,11}, Vaida A^{12,13}, Moscoso A¹⁴, Andlauer O^{15,16}, Tateno M¹⁷, Lydall G¹⁸, Wong V¹⁹, Rujevic J²⁰, Platz Clausen N²¹, Psaras R²², Delic A²³, Losevica M²⁴, Flegar S²⁵, Crépin P^{26,27}, Shmunk E²⁸, Kuvshinov I²⁹, Loibl-Weiß E³⁰, Beezhold J^{31,32}.

¹ Unit for Social and Community Psychiatry, WHO Collaborating Centre for Mental Health Service Development, Queen Mary University of London, London, UK

² Department of Psychology, University of Ljubljana, Aškerčeva 2, SI-1000 Ljubljana, Slovenia

³ Department of Psychiatry, University of Naples SUN, Largo Madonna delle Grazie, 80138 Naples, Italy

⁴ Department of Liaison Child and Adolescent Psychiatry, Our Lady's Hospital Crumlin, Dublin, Ireland

⁵ Department of Diagnostic-Clinical Medicine and Public Health University of Modena & Reggio Emilia, Modena, Italy.

⁶ Department of Psychiatry, Zagreb University Hospital Centre, Kispaticeva 12, 10 000 Zagreb, Croatia

⁷ Private psychiatric practice, 140 Avenue Victor Hugo, 75116 Paris

⁸ UPC KU Leuven, Leuven, Belgium

⁹ Department of Psychiatry and Psychoterapy, Semmelweis University, Budapest, Hungary

¹⁰ Department of Psychiatry, First Faculty of Medicine, Charles University in Prague;

¹¹ Institute of Neuropsychiatric Care, Prague

¹² Clinical Hospital of Psychiatry 2, Tirgu Mures, Romania

¹³ Specialist Psychotherapies Services, BSMHFT, Birmingham, UK

¹⁴ Hospital de D. Estefânia, Lisbon, Portugal

¹⁵ Laboratoire de Neurosciences de Besancon, Université de France-Comté, France

¹⁶ Newham centre for mental health, East London NHS Foundation Trust, UK

¹⁷ Department of Neuropsychiatry, Sapporo Medical University, Japan

¹⁸ HSSD Guernsey, UK

¹⁹ Private psychiatric practice, 122A, New Henry House, 10 Ice House Street, Central, Hong Kong

²⁰ Department of child psychiatry, University Children's Hospital, Bohoričeva 20, 1525 Ljubljana, Slovenia

²¹ Department for affective disorders, Aarhus University Hospital, Denmark

²² Private Psychiatric Practice, Athens, Greece

²³ Asociation XY - Mental Health Project in Bosnia and Herzegovina, Sarajevo, Bosnia and Herzegovina

²⁴ Riga Centre of Psychiatry and Addiction Disorders, Outpatient Department, Tvaika 2, Riga, Latvia

²⁵ Private psychiatric practice, Cape Town, South Africa

²⁶ Union Sanitaire et Sociale Aude Pyrénées, Clinique Via Domitia, Narbonne, France

²⁷ Psychiatry Clinic of University of Tartu, Estonia

²⁸ Siberian State Medical University, Tomsk, Russia

²⁹ Republican Research and Practice Centre of Mental Health, Minsk, Belarus

³⁰ Landeskrankenhaus Baden-Mödling, Standort Baden, Austria

³¹ Norfolk and Suffolk NHS Foundation Trust

³² University of East Anglia, UK

Corresponding author:

Nikolina Jovanovic, MD, PhD

Unit for Social and Community Psychiatry,

WHO Collaborating Centre for Mental Health Services Development

Barts and The London School of Medicine and Dentistry

Queen Mary University of London

E-mail: n.jovanovic@qmul.ac.uk

ABSTRACT

Background

Postgraduate medical trainees experience high rates of burnout, but evidence regarding psychiatric trainees is missing. We aim to determine burnout rates among psychiatric trainees, and identify individual, educational and work-related factors associated with severe burnout.

Methods

In an online survey psychiatric trainees from 22 countries were asked to complete the Maslach Burnout Inventory (MBI-GS) and provide information on individual, educational and work-related parameters. Linear mixed models were used to predict the MBI-GS scores, and a generalized linear mixed model to predict severe burnout.

Results

This is the largest study on burnout and training conditions among psychiatric trainees to date. Complete data were obtained from 1980 out of 7625 approached trainees (26%; range 17.8 - 65.6%). Participants were 31.9 (SD 5.3) years old with 2.8 (SD 1.9) years of training. Severe burnout was found in 726 (36.7%) trainees. The risk was higher for trainees who were younger ($p<0.001$), without children ($p=0.010$), and had not opted for psychiatry as a first career choice ($p=0.043$). After adjustment for socio-demographic characteristics, years in training and country differences in burnout, severe burnout remained associated with long working hours ($p<0.001$), lack of supervision ($p<0.001$), and not having regular time to rest ($p=0.001$). Main findings were replicated in a sensitivity analysis with countries with response rate above 50%.

Conclusions

Besides previously described risk factors such as working hours and younger age, this is the first evidence of negative influence of lack of supervision and not opting for psychiatry as a first career choice on trainees' burnout.

Key words – burnout, psychiatry, training, risk factors, supervision

1. Introduction

Recruitment and retention in psychiatry have been two issues of major concern in recent decades. While negative perception of psychiatry may discourage medical graduates from entering the field (1,2), reports from doctors leaving psychiatry have indicated reasons such as high numbers of challenging patients, depressing work conditions, job stress and low morale among staff (3). With shortage of qualified psychiatrists, most countries today cannot afford to lose trainees for reasons such as poor training conditions and development of work-related burnout. However, very little is known regarding burnout rates, or indeed factors influencing burnout syndrome among psychiatric trainees.

Work-related burnout syndrome is characterised by a triad of emotional exhaustion, cynicism, and low sense of professional efficacy (4). It has been associated with impaired patient care (5), reduced learning capacity (6), stress-related health problems (7), and broken personal relationships (8). Medical doctors experience higher rates of burnout syndrome than the general population (9). Persistent imbalance between demands and resources seems to be a crucial contributor to development of burnout. Training years, more than any other stage in physicians' career, are characterised by this type of imbalance due to long and irregular work hours (10) and high levels of responsibility combined with lack of professional experience (11). Psychiatry itself adds several very specific stressors such as perceived stigma of this profession, demanding therapeutic relationships, personal threats from violent patients and patient suicide (12).

The reported burnout rates among medical trainees vary from 27% to 75%, depending on specialty, country and methods (13, 14). A small number of studies explored this issue among psychiatric trainees. A national multi-specialty study from Netherlands included 242 psychiatric trainees (15). Reported burnout rates of 23.2% were higher than for the whole sample (21%). This was the only extant study that analysed the role of psychiatric training conditions in this context. The authors reported that trainees who worked with chronically-ill patients were more emotionally exhausted than others. Similarly, an US study found that psychiatric trainees (n=11) reported higher levels of emotional exhaustion than trainees from other specialties (16). Martini and colleagues compared US trainees from eight different specialties and with a sample of 15 psychiatric trainees found that 40% met criteria for burnout, while the rate was 50% for the whole sample (17). Several other studies included rather small subsamples of psychiatric trainees that were merged with trainees from other specialties, without reporting burnout rates or risk factors specifically for psychiatric trainees (18-21). Studies in other disciplines indicated that occupational factors such as excessive workload, high degree of work-home interference, and perception of work as stressful, were strongly correlated with burnout syndrome (13, 14). Individual factors such as neuroticism and being unmarried were also found to be associated with burnout, but these associations were weak. Overall, there is a lack of evidence regarding burnout rates and factors influencing burnout among psychiatric trainees.

Thus, the aim of this study was to assess burnout rates among psychiatric trainees, and explore which individual, educational and work-related factors are associated with severe burnout.

2. Methods

The study was created by a group of European trainees involved as national representatives in the European Federation of Psychiatric Trainees and early career psychiatrists involved with the European Psychiatric Association. The driving force behind the study was a shared awareness of the impact of burnout on service provision, and of the challenges for retention and recruitment in psychiatry.

2.1. Study protocol

This is a cross-sectional, multicountry, online survey in which participants were sent e-mail invitations to participate in the study by entering data via an online questionnaire directly into an anonymised, encrypted, secure database. While planning the study, we explored available avenues to access trainees' e-mail contacts. In the majority of countries (N=15) all trainees were invited to participate in the study. If centralised database of trainees was not available, other options were considered. In 6 countries, we were able to invite all trainees from at least three training institutions. While choosing participating institutions, trainee organisations acted to balance geographical and educational characteristics to ensure the most representative sample within local limitations. Total number of trainees in each country and sampling method are listed in **Table 1**. An ethical approval was obtained in each country in line with best practice. The study was first piloted in Croatia and France in 2007/2008 (22), and subsequently implemented in other countries during 2009–2012. The only inclusion criterion was being a psychiatric trainee, defined as a fully qualified medical doctor enrolled in a nationally recognised specialist training program in psychiatry. All participants received two reminders within a one month timeframe. Taking part in the study was voluntary.

The main outcome was trainees' burnout syndrome measured using the Maslach Burnout Inventory MBI-GS (4). The instrument consists of 16 items divided into three subscales: *emotional exhaustion* (MBI-EX), *cynicism* (MBI-CY), and *professional efficacy* (MBI-PE). All items use a frequency rating scale ranging from 0 (never) to 6 (daily). The MBI-GS cut-off scores for severe burnout are mean sum scores ≥ 2.20 on the MBI-EX and ≥ 2.00 on the MBI-CY, or ≤ 3.66 on the MBI-PE (4, 23). To determine severe burnout cases as a dichotomous outcome, only the first two criteria were used as there is empirical evidence that exhaustion and cynicism constitute the core of burnout and are often used to distinguish clinically burned out from non-burned out subjects, whereas lack of professional efficacy seems to play a less prominent role (7).

We also collected data on three groups of predictors - individual (age, gender, relationship status, having children, psychiatry as first career choice), educational (completed years of training, postgraduate education other than training), and work-related (weekly working hours, daily rest and weekly clinical supervision) characteristics. Relationship status was dichotomized into being single (comprising being single, divorced, separated, and widowed), and being in a relationship (married or in a relationship).

2.2. Statistical analysis

Burnout syndrome was explored across three dimensions (continuous variables) and as severe burnout (dichotomous variable). Descriptive statistics were used to report burnout rates and trainees' general characteristics. Sensitivity analysis was performed to examine the possible impact of non-response on burnout rates. First, to gain some insight into relations between the studied variables, all predictors were compared between severe burnout and others using chi square and analyses of variance, depending on the type of data.

Secondly, due to inter-country variance (MBI-EX: 9.2%, MBI-PA: 16.1%, and MBI-CY: 5.0%), linear mixed models (provided by IBM SPSS Statistics 22 software) were used to predict MBI scores. A random-intercept model was used to account for the nesting (clustering) of residents within countries. Country was considered a level-2 variable, while the other variables were considered level-1 variables. A random-intercept model where all predictors were entered simultaneously in the regression equation as main effects only was compared to the null random-intercept model. Because the means of various predictors differed among the countries, we decided to centre all continuous predictors using group-mean centering (24). The maximum likelihood method was used for estimating parameters. As residuals were not normally distributed for any of the studied criteria, the bias-corrected and accelerated bootstrap approach with 10,000 generated samples stratified by country was used for deriving 95% CIs for the estimated regression parameters.

Next, a generalized linear mixed model was used to examine how different predictors were associated with severe burnout. Subjects were grouped within countries and the same predictors as in predicting the MBI scale scores were entered simultaneously in the model with random intercepts.

A binary logistic regression model (with binomial probability distribution and logit link function specified) was used with Satterthwaite's approximation for determining degrees of freedom and robust estimation of covariances. A 5% alpha error rate was used as the limit of statistical significance for each predictor.

3. Results

3.1. Subjects

Out of 7468 trainees who received an e-mail invitation to participate, 2698 (36.1%) partially responded and 1980 trainees (26%) fully completed the main outcome measure (MBI-GS). Thus, a total of 1980 trainees from 22 countries were included in the final analysis presented in this paper. Please see the **Flow diagram**. Trainees' individual, educational and work profile is given in **Table 2**. On average, participants had started psychiatry training at the age of 28.5. At the time of study they had completed 2.8 years of training and were 32 years of age. Half of the sample was engaged in some form of postgraduate education outside official psychiatry training curriculum (n=1092, 55.2%) such as having a master's degree (n=525), being a master's degree student (n=27), having a PhD (n=82), being a PhD student (n=117), or having completed/being in psychotherapy training (n=439). For the vast majority,

psychiatry was the first career choice (n=1588, 80.2%) and this decision was made mostly after medical school (n=947, 59.6%). Participants reported an average of 53.7 working hours per week divided between the workplace (mean 48.1, SD 14.3) and doing work assignments at home (mean 5.9, SD 7.1). The majority of the sample (n=1349, 68.2%) was working more than 48 hours per week on average. Regular weekly clinical supervision was available to 1666 (83.9%) trainees, ranging from one hour (n=722, 36.9%) to five hours per week (n=110, 5.6%). A total of 286 trainees (14.4%) reported not having any supervision for their clinical work. It was also found that majority were able to get rest breaks such as eleven hours of continuous rest a day (n=1037, 52.4%), a day off each week (n=1638, 82.7 %), and a 20 minutes rest break every six hours (n=1297, 65.5%).

3.2. Burnout syndrome

Table 1 shows mean scores on all three MBI-GS subscales for each country. When compared to standard values provided in the MBI-GS Manual (4), the overall level of burnout was moderate across all three dimensions. The number of trainees scoring above cut-off values on emotional exhaustion was 1167 (58.9%), on cynicism 892 (45.1%) and below cut-off on professional efficacy 414 (20.2%). The number of severe burnout cases was 726 (36.7%). The prevalence ranged from 15% in Bosnia and Herzegovina to 60% in South Africa. Internal consistency of the three subscales was good, Cronbach's alpha for MBI-EX was 0.89, for MBI-CY 0.81, and for MBI-PE 0.82.

3.3. Non-response bias

There are several ways to address non-response bias (25). Firstly, countries with response rate above 50% of all trainees in the country (i.e. Bosnia and Herzegovina, Croatia, Estonia, and Slovenia) were compared to the rest of the sample. No statistically significant difference was found in emotional exhaustion (mean 2.6, SD 1.4 vs. mean 2.5, SD 1.4, $F(1)=0.333$, $p=0.467$), cynicism (mean 1.9, SD 1.2 vs. mean 2.1, SD 1.4, $F(1)=2.43$, $p=0.119$) or rates of severe burnout (49/127 vs. 677/1763, $\chi^2(1)=0.002$, $p=0.96$). However, trainees in these countries scored significantly higher on professional efficacy (mean 4.7, SD 0.9 vs. mean 4.5, SD 1.1, $F(1)=8.118$, $p\leq 0.001$). Associations between predictors and severe burnout were all replicated with the exception of weekly working hours (mean 53.8, SD 16.1 vs. mean 50.6, SD 14.7, $F(1)=2.55$, $p=0.111$). Secondly, we assumed that partial responders are likely to resemble non-responders. When compared to full responders, partial respondents ($n=718$) did not differ in socio-demographic characteristics or the reported weekly working hours (see flow diagram). Thirdly, we searched for national data to compare basic socio-demographic characteristics of responders vs. all psychiatric trainees within the country. Data on trainees' age and gender were available from UK, Hong Kong, Croatia, Latvia, Estonia and Belgium. No difference in gender distribution was found. In terms of age, the UK sample in our study differed from the UK national sample. The proportion of younger trainees (less than 34 years) was higher in our study than in the national sample (575/811 vs. 1317/1984).

3.4. The role of individual, educational and work-related characteristics

To explore the role of these predictors on emotional exhaustion, cynicism and professional efficacy and at the same time control for the differences in burnout rates

between countries, a hierarchical linear model was created and results are shown in **Table 3**. The models were statistically significant (MBI-EX $\chi^2(10) = 133.9, p < 0.001$; MBI-CY $\chi^2(10) = 119.5, p < 0.001$; MBI-PE $\chi^2(10) = 65.7, p < 0.001$). A higher level of emotional exhaustion was statistically significantly associated with longer working hours, lack of clinical supervision and less than 11 hours of daily rest. Cynicism was statistically significantly higher in younger participants and in participants without children, male trainees, those who stated that psychiatry was not their first career choice, and those who had no clinical supervision. Professional efficacy was statistically significantly lower in those without postgraduate education.

To explore the influence of these predictors on severe burnout, we created a generalized linear model and results are shown in **Table 4**. The probability for the presence of severe burnout was positively related to younger age (for every one-year increase, the odds ratio of severe burnout decreased for 17%), not having children (the odds ratio of severe burnout was increased by 44% compared to the odds ratio in participants having children), and not choosing psychiatry as the first career choice (the odds ratio of severe burnout was 21% higher than in the participants who chose psychiatry as the first career choice). Not being enrolled in or not having completed postgraduate education was of borderline statistical significance. All work-related characteristics showed significant association even when controlled for country differences and individual and educational predictors. The odds ratio of severe burnout was higher with more weekly working hours (it increased by 9% with every additional hour), not having clinical supervision (it was 63% higher than in the group receiving supervision), and not having regular 11 hours rest in every 24 hours (the odds ratio of severe burnout was 38% higher for those who did not have regular rest compared to

those who did have it). The relationship between severe burnout and working hours remained positively significant even when only working hours at the workplace were taken into account (without hours spent doing assignments at home).

4. Discussion

Key findings:

- Besides national variations, 36.7% of psychiatric trainees met criteria for severe burnout syndrome.
- The risk was increased by several individual factors such as not opting for psychiatry as a first career choice, not having children and younger age. Educational characteristics such as years spent in training and other postgraduate education were not statistically significantly associated with severe burnout.
- Even after adjusting for these individual differences, country differences in burnout rates, and years in training, three work-related factors remained positively associated with severe burnout: long working hours, lack of clinical supervision, and not having regular time to rest.

Burnout syndrome rates among psychiatric trainees in most countries included in the study were substantial and double that in the general population (26). However, they seemed to be in line or lower than among other medical trainees (15) or fully qualified psychiatrists (9, 23). This rather optimistic finding might be beneficial for the image of psychiatry and potentially attract medical graduates considering this career. When analysing trainees' individual and educational characteristics, we found that years of

completed training were not protective from severe burnout. Postgraduate education other than core psychiatric training (such as a master's degree, PhD or psychotherapy training) was of borderline protective value. Notably, trainees with additional postgraduate education did report more feelings of professional efficacy which might be one of factors that is preventing them from developing burnout syndrome. For example, it might be expected that for many trainees psychotherapy training would entail personal therapy. Although we were not able to explore this specific point in our sample as data was not available, it might be worth exploring it in future research because personal therapy could be expected to be a key factor in supporting trainees and mitigating burnout. We have also found that having children, but not relationship status, decreased the risk for severe burnout. This finding was previously reported for medical trainees (27) and might indicate a protective role of parenthood, possibly through its contribution to healthier work-life balance.

Most importantly, even after adjusting for these individual differences, country differences in burnout rates, and years in training - three work-related factors remained positively associated with severe burnout. These were long working hours, lack of clinical supervision, and not having regular time to rest. The effect of working hour limitations has been studied extensively and it was found that trainees who reported working more than 80 hours had higher rates of burnout (69.2%) compared with 38.5% after the time restriction (10). In our sample long working hours increased the risk for high emotional exhaustion and severe burnout even when only hours at the workplace were taken into account (without hours spent on work assignments at home).

This is the first study to show the negative influence of lack of supervision, and the impact of working in a field where this is not the trainees' first career choice, on developing severe burnout. Supervision has been often considered as a unique learning experience for psychiatric trainees. It offers an effective platform for trainees to reflect on their practice, and supports development of competence and professional identity (28). In the light of the known importance of clinical supervision, it is surprising that almost 15% of trainees did not receive regular weekly supervision (or did not perceive it as such). Despite several reports which have highlighted problems over structure and content of supervision (29-31), the topic lacks systematic research and remains a largely under researched area. The impact of not choosing psychiatry as first career choice (in 19.8% trainees) on developing severe burnout could be related to the lack of interest and motivation to persevere with training, and endure its more demanding aspects. This finding needs to be discussed in the light of recruitment challenges in psychiatry (32). With current unfilled training posts, it remains likely that trainees who fail to attain a training post in a preferred discipline will be recruited into psychiatry training presenting both challenges and opportunities for trainees and training systems.

The study has both strengths and limitations. This is the largest study on training conditions and burnout syndrome among psychiatric trainees to date. In most countries included in the study no previous information on training conditions from the trainees' perspective or trainee burnout is available. We hope this report will serve as a starting point for discussions regarding training and working conditions in individual countries and indeed in national organisations and at local and international policy levels. Although similar in response rate to extant data from other studies utilising online

surveys, response rate varied across participating countries (range 17.8 - 65.6%; overall 26%). However, a sensitivity analysis revealed that the prevalence of severe burnout and associations with studied predictors did not statistically significantly differ between countries with response rates of more or less than 50% which strengthens the representativeness of our results. We note that further evaluation of non-response bias indicated that UK subsample in our study is younger than the UK national sample of trainees in the same year when data was collected. Perhaps given the younger age of UK participants, this may indicate a response bias in our finding that younger trainees are at higher risk for severe burnout. Although several other studies also found younger age as a risk factor (7), it remains unclear whether younger trainees are indeed more affected or maybe more likely to participate in similar research. In light of recent findings that burnout might be more strongly associated with generic than with job-related factors (33), it might be expected parameters such as global stress, economic stability or global life satisfaction play significant role in trainees' burnout. Lastly, countries included in the study differ in cultural, educational and organisational backgrounds. We were not able to explore in detail how different historic and cultural backgrounds, work ethos, health care settings, political settings and economic inequality for both trainees and patients might impact upon the prevalence of burnout rates, although we are aware that the impact might be both positive and negative. Extant research shows that burnout is a global phenomenon that can, to some extent, go beyond intercultural differences (34). We did explore further differences between training systems in countries included in the study and thus determined that duration of training and rotations were rather similar across countries, exceptions being Belarus and Russia with significantly shorter duration of training (1 and 2 years, respectively). Since data collection most countries did not reform their training curricula, while several

countries slightly modified duration of non-psychiatric rotations (Czech Republic, Denmark and Hungary). In two countries curriculum has recently been prolonged from 4 to 5 years (Croatia and Bosnia and Herzegovina).

5. Conclusions

Working conditions influence burnout rates, even if other influential factors (such as personal and educational characteristics) are controlled for, so interventions such as limited working hours and regular supervision could represent a way of reducing severe burnout. Since previous research identified that job stress and low morale are associated with worse patient care and low retainment, we contend that these interventions may improve patient care and may increase retention rates in psychiatry. Clearly, these results only represent a part of a complex picture which is not yet fully understood or detailed. The study was created using principles of model building methodology to provide basic understanding of associations between individual, educational and training characteristics, and severe burnout among psychiatric trainees. Future studies are needed to expand our knowledge in this area. Ideally, future studies could use individual domain data as continuous variables to further evaluate the relationship between severe burnout, supervision and career choice, and other potential contributing factors. They can also be expected to provide additional evidence regarding potentially resilience-promoting factors such as age, professional experience, psychotherapy training, generic factors, and work engagement. We hope this study provides a foundation for future work to benefit psychiatry, trainees, and ultimately psychiatric patients.

Aknowledgements

We would like to thank Professor Stefan Priebe, Dr Amit Malik, Professor Andrea Fiorillo, Dr Adriana Mihai, Dr Olga Kazakova, Dr João Gama Marques, Dr Cecile Hanon, Dr Ulrike Weisz, Dr Goran Racetovic, Dr Teelia Rolko, Dr Mia Soendergaard, Dr Marija Rusaka, members of the European Federation of Psychiatric Trainees (EFPT) and the European Psychiaytric Association – European Early Career Psychiatrists (EPA-EECP).

We would like to thank the Royal College of Psychiatrists in the UK, the Hong Kong College of Psychiatrists, and the Croatian Institute of Public Health for their help in aquiring socio-demographic data of psychiatric trainees.

Declaration of interests

We declare no competing interests.

Funding

None.

REFERENCES

1. Goldacre MJ, Fazel S, Smith F, Lambert T. Choice and rejection of psychiatry as a career: surveys of UK medical graduates from 1974 to 2009. *Br J Psychiatry* 2013; 202: 228-34.
2. Farooq K, Lydall GJ, Malik A, Ndeti DM; ISOSCCIP Group, Bhugra D. Why medical students choose psychiatry - a 20 country cross-sectional survey. *BMC Med Educ*. 2014 15;14:12.
3. Lambert TW, Turner G, Fazel S, Goldacre MJ. Reasons why some UK medical graduates who initially choose psychiatry do not pursue it as a long-term career. *Psychol Med* 2006; 36: 679-84.
4. Maslach C, Jackson SE, Leiter MP. *Maslach Burnout Inventory Manual*. Consulting Psychologists Press, Palo Alto, 1996.
5. Demerouti E, Bakker AB, Leiter M. Burnout and job performance: the moderating role of selection, optimization, and compensation strategies. *J Occup Health Psychol* 2014; 19: 96–107.
6. Girard DE, Hickam DH. Predictors of clinical performance among internal medicine residents. *J Gen Intern Med* 1991; 20: 150–4.
7. Maslach C, Schaufeli WB, Leiter MP. Job burnout. *Annu Rev Psychol* 2001; 52: 397–422.
8. Shanafelt TD, Sloan JA, Habermann TM. The well-being of physicians. *Am J Med* 2003; 114: 513–19
9. Shanafelt TD, Boone S, Tan L et al. Burnout and satisfaction with work-life balance among US physicians relative to the general US population. *Arch Intern Med* 2012; 172: 1377–85.

10. Martini S, Arfken CL, Balon R. Comparison of burnout among medical residents before and after the implementation of work hours limits. *Acad Psych* 2006; 30: 352–5.
11. Schaufeli WB. Burnout. In: Firth-Cozens J, Payne R (eds). *Stress in Health Professionals: Psychological and Organizational Causes and Interventions*. Wiley, Chichester, NY, 1999.
12. Rössler W. Stress, burnout, and job dissatisfaction in mental health workers. *Eur Arch Psychiatry Clin Neurosci*. 2012; 262: S65-9.
13. Prins JT, Gazendam-Donofrio SM, Tubben BJ, van der Heijden FM, van de Wiel HB, Hoekstra-Weebers JE. Burnout in medical residents: a review. *Med Educ* 2007; 41: 788–800.
14. Thomas NK. Resident burnout. *JAMA*. 2004; 292: 288–9.
15. Prins JT, Hoekstra-Weebers JE, Gazendam-Donofrio SM et al. Burnout and engagement among resident doctors in the Netherlands: a national study. *Med Educ* 2010; 44: 236–47.
16. Afzal KI, Khan FM, Mulla Z, Akins R, Ledger E, Giordano FL. Primary language and cultural background as factors in resident burnout in medical specialties: a study in a bilingual US city. *South Med J* 2010; 103: 607–5.
17. Martini S, Arfken CL, Churchill A, Balon R. Burnout comparison among residents in different medical specialties. *Acad Psychiatry* 2004; 28: 240-2.
18. Dyrbye LN, West CP, Satele D, Boone S, Tan L, Sloan J, Shanafelt TD. Burnout among U.S. medical students, residents, and early career physicians relative to the general U.S. population. *Acad Med* 2014; 89: 443–51.

19. Msaouel P, Keramaris NC, Tasoulis A et al. Burnout and training satisfaction of medical residents in Greece: will the European Work Time Directive make a difference? *Hum Resour Health* 2010; 8: 16.
20. Zis P, Anagnostopoulos F, Sykioti P. Burnout in medical residents: a study based on the job demands-resources model. *ScientificWorldJournal* 2014; 2014: 673279.
21. Prins JT, Hoekstra-Weebers JE, van de Wiel HB, Gazendam-Donofrio SM, Sprangers F, Jaspers FC, van der Heijden FM. Burnout among Dutch medical residents. *Int J Behav Med* 2007; 14: 119-25.
22. Jovanovic N, Beezhold J, Andlauer O, Rojnic Kuzman M, Hanon C, Schulze B. Burnout among psychiatry residents – The International Psychiatry Resident / Trainee Burnout Syndrome Study (BoSS). *Die Psychiatrie* 2009; 6: 1–5.
23. Lasalvia A, Bonetto C, Bertani M et al. Influence of perceived organisational factors on job burnout: survey of community mental health staff. *Br J Psychiatry* 2009; 195: 537–44.
24. Enders CK, Tofighi D. Centering predictor variables in cross-sectional multilevel models: A new look at an old issue. *Psychol Methods* 2007; 12: 121–138.
25. Johnson TP, Wislar JS. Response rates and nonresponse errors in surveys. *JAMA*. 2012; 307: 1805-6.
26. Lindblom KM, Linton SJ, Fedeli C, Bryngelsson IL. Burnout in the working population: relations to psychosocial work factors. *Int J Behav Med* 2006; 13: 51–9.
27. Collier VU, McCue JD, Markus A, Smith L. Stress in medical residency: status quo after a decade of reform? *Ann Intern Med*. 2002; 136: 384-390.
28. Falender CA, Shafranske EP. *Clinical supervision: A competency-based approach*. American Psychological Association, Washington, DC, 2004.

29. Sembhi S, Livingston G. What trainees and trainers think about supervision. *Psych Bull.* 2000; 24: 376-9.
30. Julyan TE. Educational supervision and the impact of workplace-based assessments: a survey of psychiatry trainees and their supervisors. *BMC Med Educ.* 2009; 29; 9:51.
31. Viswanath B, Chandra PS, Chaturvedi SK. Supervision as-usual--is it enough? *Asian J Psychiatr.* 2013; 6: 488-93.
32. Katschnig H. Are psychiatrists an endangered species? Observations on internal and external challenges to the profession. *World Psychiatry* 2010; 9: 21–8.
33. Bianchi R, Boffy C, Laurent E. Burnout is More Strongly Associated with Generic Factors Than with Job-related Factors. *Eur Psychiatry.* 2015;30:210.
34. Schaufeli WB, Leiter MP, Maslach C. Burnout: 35 years of research and practice, *Career Development International* 2009; 14: 204–20.

Flow diagram




Table 1 – Sampling description with response rates, age, gender, weekly working hours and burnout syndrome rates in participants included in the study (n=1980)

Country	Time of study	N of trainees at the time of study	Sampling method ¹	Sample ²	Response rate ³	Age	Gender (M)	Weekly working hours	MBI-EX ⁴	MBI-CY ⁴	MBI-PE ⁴	Severe burnout cases ⁵
	-	-	-	N (%)	%	M(SD)	N (%)	M (SD)	M (SD)	M (SD)	M (SD)	N (%)
Austria	2010	302	A	81 (4.1)	26.8	35.2 (6.1)	29(35.8)	62.4 (17.5)	2.5(11)	1.6 (1.3)	4.9 (0.9)	28 (35)
Belarus	2009	37	A	14 (0.7)	37.8	25.1 (1.1)	2 (14.3)	38.9 (23.6)	2.5 (1.5)	2.5 (1.6)	4.4 (1.0)	5 (36)
Belgium ⁶	2012	131	D	28 (1.4)	21.8	28.4 (3.4)	4 (14.3)	63.6 (10.1)	2.8 (1.3)	2.1 (1.3)	4.3 (0.8)	10 (36)
Bosnia and Herzegovina	2009	40	A	20 (1)	50.0	33.6 (3.4)	6 (30.0)	55.5 (18.1)	1.9 (1.1)	1.8 (1.1)	5.1 (0.9)	3 (15)
Croatia	2008	106	A	56 (2.8)	52.8	31.9 (3.6)	17 (30.4)	59.0 (13.5)	2.7 (1.6)	2.2 (1.3)	5.0 (0.8)	28 (50)
Czech Republic	2009	114	A	43 (2.2)	37.7	29.9 (2.7)	18 (41.9)	55.3 (15.3)	2.9 (1.4)	2.5 (1.5)	4.8 (0.8)	19 (44)
Denmark	2010	155	A	64 (3.2)	41.3	39.8 (6.9)	15 (23.4)	42.7 (13.5)	2.6 (1.3)	1.8 (1.1)	5.3 (0.7)	20 (31)
Estonia	2010	43	A	28 (1.4)	65.1	29.4 (4.1)	6 (21.4)	50.9 (19.9)	2.6 (1.3)	2.4 (1.3)	4.8 (0.9)	11 (39)
France	2008	1131	D	134 (6.8)	35.9	27.5 (2.2)	42 (31.3)	55.5 (13.3)	2.1 (1.3)	1.4 (1.0)	3.8 (1.1)	31 (23)
Greece	2009	330	A	58 (2.9)	17.8	34.2 (2.9)	36 (62.1)	66.4 (21.6)	2.2 (1.4)	1.8 (1.5)	5.1 (0.8)	14 (24)
Hong Kong	2010	142	A	44 (2.2)	30.9	27.8 (2.5)	11 (25.0)	57.2 (7.8)	4.0 (1.3)	2.8 (1.5)	4.1 (0.8)	29 (66)
Hungary	2009	150	F	37 (1.9)	45.7	29.8 (5.2)	8 (21.6)	61.4 (10.9)	3.0 (1.5)	2.2 (1.6)	4.3 (1.3)	18 (49)
Ireland	2010	380	A	101 (5.1)	26.6	34.1 (5.1)	53 (52.5)	54.1 (11.7)	2.7 (1.4)	2.0 (1.3)	4.8 (1.2)	39 (39)
Italy	2009	730	D	118 (6)	65.6	30.4 (3.8)	27 (22.9)	50.5 (12.2)	2.6 (1.3)	1.6 (1.2)	4.6 (0.9)	26 (22)
Japan	2010	1,922	D	95 (4.8)	41.5	31.8 (4.8)	64 (67.4)	72.3 (27.1)	2.9 (1.4)	2.3 (1.4)	3.7 (1.1)	40 (42)
Latvia	2010	20	A	8 (0.4)	40.0	30.0 (6.2)	0 (0.0)	58.8 (22.0)	2.3 (1.6)	2.4 (2.0)	3.4 (1.6)	3 (38)
Portugal	2010	181	A	66 (3.3)	36.5	29.3 (2.7)	22 (33.3)	47.9 (13.0)	2.5 (1.4)	2.0 (1.2)	4.4 (0.9)	27 (41)
Romania	2010	726	A	106 (5.4)	18.8	29.4 (3.1)	23 (21.7)	56.9 (16.1)	2.4 (1.3)	2.2 (1.3)	4.9 (0.8)	42 (40)
Russia	2011	N/A	D	25 (1.3)	31.6	24.4 (0.9)	11 (44.0)	42.2 (10.5)	1.9 (1.4)	1.4 (1.5)	4.7 (0.9)	5 (20)
Slovenia	2009	45	A	23 (1.2)	51.1	39.8 (6.9)	2 (8.7)	49.2 (11.2)	2.5 (1.4)	2.0 (1.1)	4.0 (0.9)	7 (30)

South Africa	2010	47	D	20 (1)	42.5	31.8 (5.9)	9 (45.0)	65.6 (10.5)	4.1 (1.4)	2.7 (1.4)	4.5 (0.8)	12 (60)
UK	2009	3964	A	811 (41)	20.5	33 (5.2) ^a	398 (49.1)	51.0 (13.5)	2.6 (1.5)	2.1 (1.5)	4.5 (1.1)	309 (38)
Total	-	7190	-	1980 (100)	26.0	31.9 (5.3)	803 (40.6)	54.2 (16.2)	2.6 (1.4)	2.0 (1.4)	4.5 (1.1)	726 (37)

¹Hierarchical Sample-Type Options: A – all trainees in the country, B – statistically rigorous randomised sample of all trainees in the country that controls for possible sources of bias, C – randomised sample consisting of a significant proportion of all trainees in the country but without clear controls for possible sources of bias, D – all trainees from more than one training institution in the country, E – trainees from more than one training institution in the country, F – all trainees from the largest training institution in the country. ²Number of participants who fully completed the MBI-GS and were included in the final analysis presented in this paper. ³Percentage of trainees who fully completed the MBI-GS out of all approached trainees in each country. ⁴MBI-GS subscales were abbreviated as follows: Emotional exhaustion (MBI-EX), Cynicism (MBI-CY) and Professional efficacy (MBI-PE). ⁵Severe burnout was determined as the result of the following logical operation: (MBI-EX > 2.20) and (MBI-CY > 2.00). ⁶The sample was taken from Flanders only. ^aTwo participants did not provide data on their age.

Table 2 – Individual, educational and work-related predictors among all study participants (n=1980) and severe burnout cases (n=726, 36.7%)

PREDICTORS	ALL PARTICIPANTS n=1980	BURNOUT SYNDROME	
		Severe burnout cases ¹ n=726 (36.7%)	Not meeting criteria for severe burnout n=1254 (63.3%)
INDIVIDUAL			
AGE, yrs	31.9 (SD 5.3)	31.3 (SD 4.7)	32.3 (SD 5.6)
GENDER			
female	1177 (59.4%)	428 (58.9%)	749 (59.7%)
male	803 (40.6%)	298 (41.0%)	505 (40.3%)
RELATIONSHIP STATUS			
single	559 (28.2%)	219 (27.1%)	340 (25.2%)
in a relationship	1421 (71.8%)	50.07 (27.7%)	914 (26.9%)
HAVING CHILDREN			
yes	648 (33.2%)	196 (26.9%)	452 (36.0%)
no	1332 (67.3%)	530 (73.0%)	802 (63.9%)
PSYCHIATRY AS FIRST CAREER CHOICE			
yes	1588 (80.2%)	577 (79.5%)	1011 (80.6%)
no	392 (19.8%)	149 (20.5%)	243 (19.4%)
EDUCATIONAL			
COMPLETED TRAINING, yrs	2.8 (1.9)	2.7 (1.9)	2.8 (1.9)
POSTGRADUATE EDUCATION²			
yes	1092 (55.2%)	381 (52.5%)	711 (56.7%)
no	888 (44.8%)	345 (47.5%)	543 (43.3%)
WORK-RELATED			
TOTAL WEEKLY WORKING HOURS ²	53.7 (SD 15.6) Median 53	55.6 (SD 15.7) Median 54	53.3 (SD 16.5) Median 52
≤ 48 per week	629 (31.8 %)	199 (27.4%)	430 (34.3%)
> 48 per week	1349 (68.2%)	526 (72.6%)	823 (65.7%)
CLINICAL SUPERVISION			
yes	1662 (83.9%)	581 (80.0%)	1081 (86.2%)
no	286 (14.4%)	132 (18.2%)	154 (12.3%)
MIN. 11 HOURS REST IN EVERY 24 HOURS			
yes	1037 (52.4%)	332 (45.7%)	705 (56.2%)
no	943 (47.6%)	394 (54.3%)	549 (43.8%)

Data reported as mean (SD) for continuous and N (%) for categorical variables. ¹Severe burnout was a dichotomous outcome, defined as mean sum scores ≥ 2.20 on the *emotional exhaustion* (MBI-EX) and ≥ 2.00 on the *cynicism* (MBI-CY). ²Total weekly working hours are sum of time spent doing work assignments at one's workplace (mean 48.1, SD 14.3) and at home (mean 5.9, SD 7.1). Two participants did not provide data on their weekly working hours.

Table 3 - Linear mixed models for predicting scores on the three Maslach Burnout Inventory (MBI-GS)¹ scales among all study participants (n=1978)²

Parameters ³	Emotional Exhaustion (MBI-EX) ¹		Cynism (MBI-CY) ¹		Professional efficacy (MBI-PE) ¹	
	Coefficient [95% CI] ⁴	p	Coefficient [95% CI] ⁴	p	Coefficient [95% CI] ⁴	p
Intercept	2.31 [2.14, 2.48]	< 0.001	1.60 [1.43, 1.77]	< 0.001	4.69 [4.56, 4.82]	< 0.001
Gender (males)	-0.11 [-0.23, 0.01]	0.094	0.14 [0.01, 0.28]	0.030	0.05 [-0.05, 0.14]	0.343
Age	-0.05 [-0.12, 0.02]	0.119	-0.13 [-0.19, -0.06]	<0.001	0.04 [-0.01, 0.09]	0.095
Not in a relationship	0.13 [-0.01, 0.27]	0.076	0.08 [-0.06, 0.22]	0.265	-0.05 [-0.17, 0.06]	0.309
Not having children	0.11 [-0.05, 0.27]	0.165	0.24 [0.10, 0.39]	0.002	-0.07 [-0.19, 0.05]	0.229
Completed years of training	0.00 [-0.06, 0.07]	0.901	0.06 [-0.01, 0.12]	0.094	-0.03 [-0.08, 0.01]	0.171
Psychiatry was not first career choice	0.07 [-0.10, 0.24]	0.406	0.22 [0.05, 0.38]	0.007	-0.10 [-0.22, 0.02]	0.093
No postgraduate education	0.09 [-0.05, 0.22]	0.196	0.12 [-0.01, 0.25]	0.081	-0.15 [-0.24, -0.06]	0.003
Working hours	0.11 [0.05, 0.18]	<0.001	-0.00 [-0.07, 0.07]	0.984	0.02 [-0.03, 0.07]	0.467
Not enough daily rest	0.33 [0.20, 0.46]	<0.001	0.12 [-0.00, 0.24]	0.058	-0.01 [-0.10, 0.08]	0.830

No clinical supervision	0.34 [0.16, 0.52]	<0.001	0.34 [0.17, 0.52]	<0.001	-0.11 [-0.23, -0.00]	0.064
-------------------------	-------------------	--------	-------------------	--------	----------------------	-------

¹Maslach Burnout Inventory (MBI-GS) consists of 16 items divided into three subscales: *emotional exhaustion* (MBI-EX), *cynicism* (MBI-CY), and *professional efficacy* (MBI-PE). In our analysis, the three outcome variables were treated as continuous and ranged from 0 to 6.

²As two participants had some missing data, the analyses were performed on 1978 cases.

³Predictor variables were dichotomous (gender, not in a relationship, not having children, psychiatry was not first career choice, no postgraduate education, not enough daily rest, and no clinical supervision) or continuous (age, completed years of training, and working hours). All predictors were entered simultaneously in the model. Please note that for dichotomous predictors the value of the coefficient shows the difference in the outcome variable between the tested and the reference group (e. g., Not in a relationship represented the tested group compared to the reference group that was composed of participants who were in a relationship) when all other variables are held constant. In continuous variables, the value of the coefficient shows the amount of change in the outcome variable for every one-year increase in age and completed years of training, or for every one-hour increase in working hours when all other variables are held constant.

⁴95% CI for parameter estimates were obtained with the bias-corrected and accelerated bootstrap.

Table 4 - Generalized linear mixed model for predicting severe burnout¹ among all study participants (n=1978)²

Parameters ³	Coefficient [95% CI]	p	OR [95% CI]
Intercept	-1.18 [-1.51, -0.85]	< 0.001	0.31 [0.22, 0.43]
Gender (males)	0.06 [-0.13, 0.24]	0.561	1.06 [0.88, 1.27]
Age	-0.18 [-0.28, -0.09]	< 0.001	0.83 [0.76, 0.91]
Not in a relationship	0.01 [-0.16, 0.17]	0.935	1.01 [0.86, 1.18]
Not having children	0.37 [0.09, 0.65]	0.010	1.44 [1.09, 1.91]
Completed years of training	0.05 [-0.07, 0.17]	0.439	1.05 [0.93, 1.19]
Psychiatry was not first career choice	0.19 [0.01, 0.38]	0.043	1.21 [1.01, 1.46]
No postgraduate education	0.14 [-0.00, 0.29]	0.056	1.16 [1.00, 1.34]
Working hours	0.09 [0.04, 0.17]	<0.001	1.09 [1.04, 1.15]
Not enough daily rest	0.33 [0.13, 0.52]	0.001	1.38 [1.14, 1.69]
No clinical supervision	0.49 [0.32, 0.66]	<0.001	1.63 [1.37, 1.93]

¹Severe burnout was a dichotomous outcome, defined as mean sum scores ≥ 2.20 on the *emotional exhaustion* (MBI-EX) and ≥ 2.00 on the *cynicism* (MBI-CY). Please also see notes 2 and 3 to Table 3.