

Heightened dynamics of the oxidized Y48H variant of human cytochrome c increases its peroxidatic activity

Oliver M. Deacon, Andreas Ioannis Karsisiotis, Tadeo Moreno Chicano, Michael Alexander Hough, Colin J. Macdonald, Tharin Blumenschein, Michael T. Wilson, Geoffrey R. Moore, and Jonathan A.R. Worrall

Biochemistry, **Just Accepted Manuscript** • DOI: 10.1021/acs.biochem.7b00890 • Publication Date (Web): 30 Oct 2017

Downloaded from <http://pubs.acs.org> on November 2, 2017

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.



1
2
3 **Heightened dynamics of the oxidized Y48H variant of human cytochrome *c* increases its**
4 **peroxidatic activity**
5
6
7

8 Oliver M. Deacon¹, Andreas Ioannis Karsisiotis¹, Tadeo Moreno-Chicano¹, Michael A.
9 Hough¹, Colin Macdonald², Tharin M.A. Blumenschein², Michael T. Wilson¹, Geoffrey R.
10 Moore^{2*}, Jonathan A.R. Worrall^{1*}
11
12

13
14 ¹School of Biological Sciences, University of Essex, Wivenhoe Park, Colchester, CO4 3SQ,
15 U.K. ²School of Chemistry, University of East Anglia, Norwich Research Park, NR4 7TJ,
16 U.K.
17
18
19

20
21 ***To whom correspondence should be addressed:**
22

23 Geoffrey R. Moore: g.moore@uea.ac.uk; Tel: +44 1603 592003
24

25 Jonathan A.R. Worrall: jworrall@essex.ac.uk; Tel: +44 1206 872095
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Proteins performing multiple biochemical functions are called ‘moonlighting proteins’ or extreme multifunctional (EMF) proteins. Mitochondrial cytochrome *c* is an EMF protein that binds multiple partner proteins to act as a signalling molecule, transfers electrons in the respiratory chain, and acts as a peroxidase in apoptosis. Mutations in the cytochrome *c* gene lead to the disease thrombocytopenia, which is accompanied by enhanced apoptotic activity. The Y48H variant arises from one such mutation and is found in the 40-57 Ω -loop, the lowest unfolding free energy sub-structure of the cytochrome *c* fold. A 1.36 Å resolution X-ray structure of the Y48H variant reveals minimal structural changes compared to the wild type structure, with the axial Met80 ligand coordinated to the heme iron. Despite this, the intrinsic peroxidase activity is enhanced implying that a pentacoordinate heme state is more prevalent in the Y48H variant, corroborated through determination of a Met80 ‘off-rate’ of $> 125 \text{ s}^{-1}$ compared to $\sim 6 \text{ s}^{-1}$ for the wild type protein. Heteronuclear NMR measurements with the oxidised Y48H variant reveal heightened dynamics in the 40-57 Ω -loop and the Met80 containing 71-85 Ω -loop relative to the wild type protein, illustrating communication between these sub-structures. Placed into context with the G41S cytochrome *c* variant, also implicated in thrombocytopenia, a dynamic picture associated with this disease relative to cytochrome *c* is emerging whereby increasing dynamics in sub-structures of the cytochrome *c* fold serve to facilitate an increased population of the peroxidatic pentacoordinate heme state of the order, wild type $<$ G41S $<$ Y48H.

INTRODUCTION

The discovery that many proteins have more than one biological function was an important advance in understanding how complex organisms can have relatively few genes to carry out the many thousands of biochemical functions that occur within it^{3, 4}. Jeffery listed a variety of mechanisms by which the function of a protein can switch between its different activities – and termed this ability ‘moonlighting’⁶. In addition to genetic origins governing expression, these mechanisms included: differential localization within the cell, intracellular versus extracellular location, oligomerization, variable ligand/substrate concentrations, chemical regulation of activity, different binding sites for different partner molecules, and being part of different multi-protein complexes⁶. Advances in the field have led to the recent suggestion that the phrase ‘extreme multifunctional proteins’ (EMF) be employed instead of moonlighting proteins⁷. The rebranding is to create focus on the functional aspects of the phenomenon, rather than the mechanistic, noting that what makes moonlighting proteins worth studying is that they perform diverse functions and so that is what their definition should emphasize⁸. However, from a chemical perspective it is important to understand how moonlighting or EMF proteins operate. Energy landscape theory⁹ suggests that one way a protein could moonlight is if the ground-state structure, which is the prevalent form under physiological conditions, is in equilibrium with one or more higher-energy forms of different structure present in small amounts. Such excited states of proteins have been identified in numerous systems in recent years, particularly through studies of protein folding, but also in ligand binding and enzyme catalysis^{10, 11}, and thus it is reasonable to think that this offers a route to enable proteins to moonlight.

Cytochrome *c* (Cc) is well known for its role in mitochondrial electron-transfer as an electron carrier between the cytochromes *bc₁* and *aa₃* complexes¹² but it also plays two important roles in apoptosis^{13, 14}. One role is as a peroxidase to oxidise lipids as part of the process of creating a pore in the outer mitochondrial membrane¹⁵, and the second set of roles is as a component of various multiprotein complexes, including: the cytoplasmic apoptosome multi-protein complex¹⁶, heat shock protein 27¹⁷ and inositol triphosphate receptors in the endoplasmic reticulum¹⁸. Cc also functions as an inhibitor of a histone chaperone activity in the nucleus following DNA damage by binding to a nucleosome assembly protein¹⁹ and as an extracellular ligand to serum leucine-rich alpha-2-glycoprotein-1²⁰, an interaction that helps lymphocytes to survive from the toxicity resulting from Cc being released from cells undergoing apoptosis. From the structural information available, it appears that Cc in its role as a histone chaperone inhibitor¹⁹ and as a component of the apoptosome²¹ binds to the

1
2
3 relevant partner proteins at sites overlapping its binding sites for the cytochromes *bc₁* and *aa₃*
4 complexes^{22, 23}. Thus, in these moonlighting roles Cc appears to fit in well with the possible
5 mechanisms given by Jeffery⁶. However, to turn Cc from an electron carrier into a
6 peroxidase requires a major change in the way the heme group interacts with the polypeptide
7 chain. The electron-transfer form of Cc, which we can call the ground state conformation,
8 possesses a hexacoordinate heme iron with the side chains of His18 and Met80 acting as the
9 axial heme iron ligands. To act as a peroxidase, Cc needs a pentacoordinate heme to allow
10 the Fe(III) to react directly with H₂O₂, and it is this pentacoordinate state that we have
11 suggested is the poorly populated excited state in equilibrium with the ground state¹.

12
13
14
15
16
17
18 Mutations in the human Cc (hCc) gene (*CYCS*) give rise to an inherited autosomal
19 dominant disease, thrombocytopenia 4 (THC4; OMIM 612004), which is accompanied by
20 enhanced mitochondrial apoptotic activity²⁴. To date three mutations have been identified in
21 the *CYCS* gene that give rise to the G41S, Y48H and A51V protein variants²⁴⁻²⁶. We, and
22 others, have shown that the oxidized form of the hCc G41S variant has a greater intrinsic
23 peroxidase activity than the wild-type (WT) protein^{27, 28}. Furthermore, using NMR and azide
24 binding kinetics we have shown that an equilibrium exists between the hexacoordinate
25 ground state and a pentacoordinate excited state, with more of the G41S variant in the excited
26 state than for the WT protein¹. Here we report on studies of the second pro-apoptotic variant
27 of hCc, Y48H, and show for the ferric form even more of this variant is in the excited
28 pentacoordinate state than the G41S variant, and that this population difference is correlated
29 with their relative stabilities and peroxidase activities.

30 31 32 33 34 35 36 37 38 39 40 **MATERIALS AND METHODS**

41 *Site-directed mutagenesis and purification of the Y48H human cyt c variant*

42
43 The Y48H variant of hCc was constructed using the Quikchange mutagenesis (Stratagene)
44 protocol. The forward and reverse mutagenic primers were as follows: 5'-GCC CCT GGA
45 TAC TCT CAT ACA GCC GCC AAT AAG AAC-3' and 5'-GTT CTT ATT GGC GGC
46 TGT ATG AGA GTA TCC AGG GGC-3', respectively. A PCR mix consisting of primers
47 (75 ng/μl), the wild-type template (15 ng/μl), 10 mM dNTPs (Fermentas), Pfu Turbo
48 polymerase (Stratagene), 10 X Pfu Buffer (Agilent) and deionised water to give a final
49 volume of 50 μl was prepared and subjected to the following PCR cycle; 95 °C – 3 min,
50 (95 °C – 1 min, 65 °C – 1 min, 72 °C – 6 min) x18, 72 °C – 10 min. Over-expression of the
51 Y48H variant together with the wild-type and G41S variant in *Escherichia coli* were carried
52
53
54
55
56
57
58
59
60

1
2
3 out in BL21(DE3) RIL (Invitrogen) cells with isolation and purification carried out as
4 reported previously²⁷.
5
6

7 8 *Protein preparation*

9
10 Oxidized Cc proteins were prepared by the addition of excess $K_3[Fe(CN)_6]$, followed by
11 removal of $K_3[Fe(CN)_6]$ and $K_4[Fe(CN)_6]$ and exchange into a desired buffer using a PD-10
12 column (GE Healthcare). Protein concentration was determined with a Cary 60
13 spectrophotometer (Agilent) and a molar extinction coefficient of $\epsilon = 106 \text{ mM}^{-1} \text{ cm}^{-1}$ at 409
14 nm.
15
16
17
18

19 20 *Chemical denaturation and alkaline pH titrations*

21 Far-UV CD spectra at 15 °C for oxidized Cc proteins (20 μM) in 10 mM potassium
22 phosphate, 50 mM potassium fluoride pH 6.5 were recorded in the wavelength range 250-190
23 nm with an Applied Photophysics Chirascan CD spectrophotometer (Leatherhead, UK).
24 Oxidized Cc stabilities were determined by titrating a 6 M stock solution of guanidine
25 hydrochloride (GuHCl) (Fluka) to a 20 μM protein sample and the changes in molar
26 ellipticity at 222 nm ($\theta_{222\text{nm}}$) monitored. All titrations were carried out in triplicate. The
27 fraction unfolded (F_u) at any given [GuHCl] was determined²⁹ and the free energy of
28 unfolding (ΔG_{unf}) and dependence of ΔG_{unf} on denaturant concentration (m value) calculated
29 using an equation for a two-state unfolding process². The pH dependence of the 695 nm band
30 in the UV-visible spectrum was monitored at room temperature by determining its
31 absorbance at various values of pH for a solution of 100 μM oxidized Cc in a quartz cuvette
32 (Hellma) with a small aliquot of $K_3[Fe(CN)_6]$ present to maintain an oxidising environment.
33 The pH of the buffer (20 mM sodium phosphate pH 6.0) was adjusted with microliter
34 aliquots of 1 M NaOH and measured after each addition using a semi-micro glass pH
35 electrode. pH titrations were repeated up to three times and with different batches of proteins.
36
37 Data were fitted to a one-proton ionization equilibrium equation to yield an apparent pK_a .
38
39
40
41
42
43
44
45
46
47
48
49

50 51 *Preparation of liposomes and peroxidase assays*

52 The phospholipids 1,1',2,2'-tetraoleoyl cardiolipin (TOCL) and 1,2-dioleoyl-*sn*-glycero-3-
53 phosphocholine (DOPC) (AVANTI Polar Lipids, USA) were mixed in a 1:1 ratio and
54 vortexed with the appropriate amount of buffer (20 mM sodium phosphate pH 6.5) and
55 sonicated for 5 min in a high power sonicating water bath filled with ice-cold water to give
56
57
58
59
60

1
2
3 the desired stock concentration as previously described ²⁷. Peroxidase assays of the ferriCc
4 proteins were carried out in the absence and presence of liposomes using H₂O₂ (Sigma) and
5 2,2-Azinobis(3-ethylbenthiiazoline-6-sulfonic acid) (ABTS) (Sigma). The oxidation of ABTS
6 was monitored at 730 nm on a Hewlett-Packard 8453 diode-array spectrophotometer
7 scanning between 190 and 1100 nm and thermostatted at 20 °C. The reaction was initiated by
8 the addition of 1 mM H₂O₂ (after 40 s) to a series of cuvettes containing 5 μM ferriCc and
9 200 μM ABTS with or without 25 μM TOCL/DOPC from a 2.5 mM stock. Using the
10 wavelength pair 475-730 nm the slope of each trace (post-lag phase) was determined to
11 obtain a rate in AU/s. The rate of ferriCc turnover is reported in s⁻¹ obtained by dividing AU/s
12 by the product of the concertation of oxidized ABTS ($\epsilon = 14 \text{ mM}^{-1} \text{ cm}^{-1}$) and the total protein
13 concentration of 5 μM. All assays were carried out in triplicate with errors reported as the
14 standard error.
15
16
17
18
19
20
21
22
23

24 25 *Stopped-flow kinetics*

26 An Applied Photophysics (Leatherhead, UK) SX20 stopped-flow spectrophotometer
27 thermostatted at 25 °C and equipped with both photomultiplier and diode array detection
28 systems was used to monitor the kinetics of azide (N₃⁻) binding to the oxidized Y48H hCc
29 variant. A stock solution of 2 M sodium azide (Sigma) were prepared in 50 mM MES pH 7.0
30 and diluted to the desired [N₃⁻] with the same buffer containing 2 M NaCl to maintain the
31 ionic strength. Reaction time-courses were taken at 420 nm with [N₃⁻] varying between 0.08
32 and 2 M before mixing and 10 μM protein (before mixing). All transients were fitted to a
33 single exponential function yielding both pseudo first-order rate constants and amplitudes.
34
35
36
37
38
39
40
41

42 *Crystallisation and X-ray structure determination*

43 Crystals were grown using the sitting-drop vapour diffusion method at 18 °C. A 0.2 μl
44 solution containing 56 mg/ml of oxidized Y48H Cc variant in 20 mM sodium phosphate pH
45 6.5 was mixed with an equal volume of reservoir solution (33 % PEG6000, 100 mM Tris/HCl
46 pH 7.5) using a Crystal Gryphon robot (Art Robbins Instruments). Crystals were transferred
47 to a cryoprotectant solution containing the reservoir solution and 15 % glycerol, before flash
48 cooling by plunging into liquid nitrogen. Crystallographic data were measured at the Swiss
49 Light Source (SLS, Villigen, Switzerland) at beamline X10SA using a Pilatus 6M-F detector
50 (Dectris) and an X-ray wavelength of 1.0 Å. Data were indexed using XDS ³⁰ and scaled and
51 merged using Aimless ³¹ in the CCP4 suite with the CCP4i2 interface. The structure was
52
53
54
55
56
57
58
59
60

solved by molecular replacement in Phaser³² using the WT hCc structure²⁷ (pdb code 3zcf) as the search model. The structure was refined with Refmac5³³, with model building between refinement cycles in Coot³⁴. Riding hydrogen atoms were added when refinement of the protein atoms had converged. Structures were validated using the Molprobity server³⁵ the JCSG Quality Control Server and tools within Coot³⁴. Structural superpositions were carried out using GESAMT in CCP4i2³⁶. Coordinates and structure factors were deposited in the RCSB Protein Data Bank with entry 5O10. A summary of data, refinement statistics and the quality indicators for the structure is given in Table 1.

Table 1: Crystallographic data processing and refinement statistics for the Y48H hCc variant. Values in parenthesis refer to the outermost resolution shell 1.39-1.36 Å.

Parameter	hCc Y48H
Space group	P2 ₁
Unit cell	a = 56.9, b = 36.5, c = 60.7 Å, β = 116.7°
Resolution (Å)	1.36
R _{merge}	0.072 (1.115)
Unique reflections	45812 (2124)
Mn (I/SD)	13.3 (1.4)
CC _{1/2}	0.998 (0.512)
Completeness (%)	95.3 (90.2)
Redundancy	6.2 (5.4)
R _{cryst}	0.144
R _{free}	0.175
RMS dev. bond lengths (Å)	0.015
RMS dev. bond angles (°)	2.18
Ramachandran favoured (%)	97.0
Accumulated dose (MGy)	2.54
PDB accession code	5O10

NMR spectroscopy

All NMR data were collected with a Bruker 500 MHz (H/D exchange) or 800 MHz (1D, ¹⁵N-relaxation and sequence specific assignments) spectrometer, both equipped with 5 mm HCN inverse triple resonance z-axis gradient probes. Samples for 1D ¹H NMR spectroscopy were oxidized and buffer exchanged into 20 mM sodium phosphate pH 6.5 and freeze dried.

1
2
3 Samples were redissolved in 500 μl of 99.9 % D_2O (sigma) and allowed to deuterate at room
4 temperature for ~ 72 h. 4,4-dimethyl-4-silapentane-1-sulfonic acid (DSS; Sigma) (200 μM
5 final concentration) and $\text{K}_3[\text{Fe}(\text{CN})_6]$ (100 μM final concentration to maintain an oxidising
6 environment) were added to each sample and 1D spectra were recorded at temperatures
7 ranging from 288 to 313 K using presaturation to suppress the solvent signal and with spectral
8 widths of 20 and 80 ppm. Spectra were processed in Topspin (Bruker Biospin).
9

10
11
12
13 Single (^{15}N) and doubly (^{13}C , ^{15}N) labelled samples of the Y48H hCc variant were
14 prepared using procedures previously described³⁷. NMR samples of the oxidized Y48H
15 variant were prepared in 20 mM sodium phosphate pH 6.5 and contained 1 mM protein with
16 10 % D_2O (Sigma) for the lock. A set of 2D [^1H , ^{15}N]-HSQC and 3D HNCACB and
17 CBCA(CO)NH spectra were recorded at 288 K and processed in Topspin (Bruker Biospin).
18 Sequence specific assignments were obtained using the CCPNmr Analysis³⁸ and Sparky³⁹
19 programs.
20

21
22
23
24
25 H/D exchange NMR data was acquired at 288 K. Freeze-dried samples of the
26 oxidized Y48H variant (1 mM) were re-dissolved in 20 mM sodium phosphate pH 6.5, 50
27 mM NaCl pre-made in 99.9 % D_2O (Sigma). Spectra were recorded as matrices of 1024 x 64
28 complex data points with 4 scans and spectral widths of 15 (^1H) and 42 (^{15}N) ppm centred at
29 4.7 and 118.6 ppm, respectively. The acquisition of consecutive [^1H , ^{15}N]-HSQC spectra
30 started ~ 5 min after sample resuspension. Acquisition proceeded with consecutive HSQC
31 spectra of ~ 5 min acquisition time for a period of 134 h, totalling 191 experiments. The first
32 100 experiments were collected with a 10 s delay between each to ensure maximum coverage
33 of the fast-exchanging residues, with the following 74 experiments at 45 min intervals,
34 followed by 17 experiments at 4 h intervals. CCPNmr Analysis³⁸ was used to fit peak
35 volumes as a function of time to a single exponential decay function to yield decay rates from
36 which protection factors (PFs) were calculated^{40, 41} (Table S1). The free energies of
37 exchange (ΔG_{ex}) were calculated from $\Delta G_{\text{ex}} = -RT\ln(\text{PF})$ where R is the gas constant and T is
38 the temperature in Kelvin⁴¹ (Table S1).
39

40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
 ^{15}N T_1 , T_2 and [^1H]- ^{15}N NOE data were collected at 288 K with spectra acquired as
matrices of 1024 x 256 complex data points and 8 scans for T_1 and T_2 data and 64 scans for
[^1H]- ^{15}N NOE data. Spectral widths of 15 (^1H) and 39.3 (^{15}N) ppm centred at 4.7 and 118.6
ppm, respectively, were used. T_1 experiments were acquired with relaxation delays of 20, 50,
80, 200, 500, 750, 2000 and 3500 ms with duplicates at 20, 200 and 500 ms. T_2 experiments
were acquired with relaxation delays of 16.96, 33.92, 50.88, 67.84, 101.76, 135.68, 203.52

1
2
3 and 254.40 ms with duplicates at 16.96, 50.88 and 101.76 ms. Recycle delays for both
4 experiments were 5 s. The repeated relaxation delays were used for the determination of peak
5 height uncertainties⁴². Both the T₁ and T₂ series of experiments were acquired as a pseudo-
6 3D experiment and processed as individual planes 2D planes using NMRPipe⁴³. Residues
7 with peak overlap or weak intensity were excluded from the analysis. T₁ and T₂ values were
8 extracted through the fitting of peak heights as a function of time to a single exponential
9 function in CCPNmr Analysis³⁸. These were subsequently converted to R₁ and R₂ rate
10 constants (R₁ = 1/T₁ and R₂ = 1/T₂). The fit errors for T₁ and T₂ were also converted into R₁
11 and R₂ errors through the equality of the T₁/T₂ and R₂/R₁ errors. Heteronuclear [¹H]-¹⁵N
12 NOEs were measured with a proton saturation period of 5 s. Interleaved saturated and
13 unsaturated experiments were acquired in triplicate to determine the experimental errors as
14 the standard deviation of the average NOE value. NOE values were calculated as the ratio of
15 peak heights with and without proton saturation.
16
17
18
19
20
21
22
23
24
25

26 RESULTS

27
28 *X-ray structure of the pro-apoptotic Y48H hCc variant* The X-ray structure of the Y48H
29 variant of hCc was determined to 1.36 Å resolution. A space group of P2₁ was determined
30 with two hCc molecules found in the crystallographic asymmetric unit, contrasting with the
31 reported X-ray structures of the WT protein²⁷ and G41S variant⁴⁴, both of which had a P₁
32 space group and four hCc molecules in the asymmetric unit. The diffraction-weighted dose⁴⁵
33 incurred during collection of the high-resolution X-ray dataset for the Y48H variant was 2.54
34 MGy, a dose greatly exceeding that required to fully reduce ferric heme proteins to the
35 ferrous form (typically 10s-100s kGy)^{46, 47}. Hence, while the crystals were obtained from
36 solutions of the ferric protein, in light of the extensive previous studies, measuring the doses
37 typically required for ferric to ferrous heme protein reduction, we consider the structure to
38 likely be predominantly that of the ferrous state. It is important to note that earlier X-ray
39 studies of tuna Cc⁴⁸ and *Saccharomyces cerevisiae* iso-1 Cc⁴⁹ with much lower X-ray doses
40 than the present study show that the conformation of the protein is largely independent of the
41 heme oxidation state. Electron density in the heme region is consistent with a hexacoordinate
42 species, with the Nε2 atom of His18 and the Sδ atom of Met80 coordinating the heme iron.
43 The overall fold of each molecule in the asymmetric unit is identical to that of the WT protein
44 and the G41S variant, with an overall RMS deviation in Cα positions of 0.18 Å between the
45 three X-ray structures (Figure 1A). It is noted that the Cα deviation between residues 37-61
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

increases to 0.27 Å. Replacing Tyr48 with a His residue alters a H-bonding arrangement involving the heme propionates. In the WT protein structure the OH group of the Tyr48 side chain forms a H-bond (2.57 Å) to the O1A atom of heme propionate-7²⁷. Such an interaction is lost in the Y48H variant and the Nδ1 atom of the His48 side chain is favourably orientated to form a medium length polar interaction with the O2D atom of the heme propionate-6 (3.19 Å) (Figure 1B). The number of H-bond interactions with the OD2 atom of heme propionate-6




Figure 1: X-ray crystal structure of the Y48H variant of hCc. A) Ribbon representations of a C α superposition (molecule A) of the Y48H variant (green; pdb 5O10) with the G41S variant (red; pdb 3nvw) and WT protein (blue; pdb 3zcf). The heme groups are shown in sticks. B) A cartoon and stick representation of the region between residues 37 and 49 of the WT protein (blue), the Y48H (green) and G41S (red) variants. The coordinate bonds (black dashes) of the axial His18 and Met80 ligands to the heme iron in the Y48H variant are indicated and H-bonding is represented as red dashes. The side chain position of His48, Asn52 and Tyr67 in the Y48H variant are shown together with the side chain position of Arg38 in all three structures. Water molecules are shown in red spheres for the WT and Y48H structures and in grey for the G41S variant.

therefore increases to two, with the second H-bond interaction involving the backbone amide of Thr49. A second medium length polar interaction between the Nδ1 atom of His48 and the OD1 atom of Asn52 (3.19 Å) is also noted leading to His48 communicating with the Met80 ligand through a H-bond network that extends from Asn52, involving a H₂O molecule and Tyr67 (Figure 1B). Interestingly, the Arg38 side chain in the Y48H structure is positioned as in the WT structure, with its ε-amido group coupled to heme propionate-7 via a H₂O molecule (Figure 1B). In the G41S structure the Arg38 side chain moves away from the heme propionate-7 leading to the presence of two additional H₂O molecules (Figure 1B). These waters enable the ε-amido group of Arg38 to remain coupled to the heme propionate-7 via a forked hydrogen bond network (Figure 1B).

The Y48H variant has a lower global stability than the G41S variant but a similar alkaline isomerisation pK

The Y48H variant of hCc displayed optical absorbance properties in the ferric and ferrous




Figure 2: Global stability and pK_{695} measurement of the oxidised Y48H variant of hCc compared under identical experimental conditions with the WT protein and G41S variant. A) Plots of fraction unfolded as a function of GuHCl at pH 6.5 and 15 °C. Solid-lines are fits to the data using a two-state equilibrium unfolding equation², with the determined ΔG_{unf} values reported in Table 2. B) The pH dependence of the 695 nm absorbance band. Solid-lines are representative fits to the data using a one-proton equilibrium equation with pK_{695} values reported in Table 2.

oxidation states and a far-UV CD spectrum identical to that of the WT protein. Far-UV CD spectroscopy is a convenient method to determine the global stabilization of a protein through chemical denaturation studies using GuHCl as illustrated in Figure 2A. It is apparent from Figure 2A that there is a progressive loss in the cooperativity of unfolding (WT > G41S > Y48H) with the Y48H variant having the lowest global stability (ΔG_{unf} , Table 2). The m -value (Table 2), which is a measure of the change in hydrophobic surface area that has become desolvated upon denaturation, decreases in the order WT > G41S > Y48H, suggesting that the variants have an increasingly more compact denatured state relative to the WT protein⁵⁰. Interestingly the K54H variant of *S. cerevisiae* iso-1 Cc stabilizes the denatured state through a His-heme loop leading to a decreased m -value relative to WT protein⁵¹. It may be that a similar effect occurs for the Y48H variant. Alternatively, the Y48H variant may simply be partially unfolded in buffer, which would also lower the m -value⁵², with the NMR data (Fig. 6A) providing some support for this interpretation (*vide infra*).

Table 2: Summary of thermodynamic and kinetic parameters for the Y48H human ferricyt *c* variant and comparison with the WT protein and the G41S variant.

Protein	GuHCl denaturation				Azide binding		
	ΔG_{unf} (kcal mol ⁻¹)	<i>m</i> (kcal mol ⁻¹ M ⁻¹)	<i>C_m</i> (M)	<i>pK</i> ₆₉₅	<i>K</i> _{app} (M)	<i>k</i> ₁ (s ⁻¹)	<i>k</i> ₂ (s ⁻¹)
Y48H	5.09 ± 0.16	2.57 ± 0.1	1.98 ± 0.05	8.4 ± 0.1	0.07 ± 0.01	125	12.1 ± 1.1
WT	10.65 ± 0.55	4.10 ± 0.25	2.60 ± 0.05	9.3 ± 0.2	0.31 ± 0.03 ^a	5.77 ± 1.5 ^a	3.5 ± 0.44 ^a
G41S	6.75 ± 0.10	3.0 ± 0.1	2.25 ± 0.05	8.5 ± 0.2	0.11 ± 0.01 ^a	45 ± 18 ^a	8.9 ± 0.3 ^a

^aData taken from reference ¹.

Figure 2B shows the variation in absorbance of the 695 nm band for the oxidized Y48H variant as a function of pH compared to data for WT and the G41S variant. Although the latter *pK*₆₉₅ values for the WT and G41S variant have been previously reported ²⁷, they have been re-determined in the present study so that a true comparison under identical conditions can be made (*i.e.* in the absence of NaCl that is known to affect the *pK*₆₉₅ value ⁵³). The bleaching of the 695 nm band is a consequence of a deprotonation event in which the axial Met80 heme ligand becomes dissociated. The apparent *pK*₆₉₅ value for the Y48H variant, which was obtained from these data, is near identical to that of the G41S variant but significantly lower than that of the WT protein (Table 2). Thus, whereas the global stability for the Y48H variant is clearly decreased, the Met80-Fe(III) bond lability as determined through *pK*₆₉₅ values is indistinguishable from the G41S pro-apoptotic variant.

Exogenous ligand binding reveals the pentacoordinate form of the oxidized Y48H variant is more populated than for either the G41S variant or WT hCc

To probe further variances in Met80-Fe(III) bond lability, the kinetics of N_3^- binding was




Figure 3: Azide binding to the oxidized Y48H variant of hCc variant compared with data for the WT protein and G41S variant ¹ at pH 7.0 and 25 °C. A) Normalized amplitudes from the reaction time-courses obtained using stopped-flow absorption spectroscopy at 420 nm plotted against azide concentration. Data are fitted to a hyperbolic equation to yield K_{app} values reported in Table 2. B) The rate constants (k_{obs}) determined for azide binding with solid lines representative of fits to Equation 2, to give k_1 and k_2 reported in Table 2. C) S_N1 mechanism of azide binding to oxidized Cc.

studied using stopped-flow spectrometry. Upon mixing N_3^- with the oxidized Y48H hCc an optical transition occurred caused by the dissociation of Met80 from the heme iron and the binding of N_3^- in its place. The reaction time-course of this transition conformed to a simple exponential, as reported previously for the WT protein and the G41S variant under the same experimental conditions ¹. Over a broad range of $[N_3^-]$ the normalized amplitudes of such reaction time-courses follow a simple hyperbolic binding isotherm (Figure 3A) enabling an apparent equilibrium constant (K_{app}) for N_3^- binding to be determined. From Table 2 it is apparent that the Y48H variant binds N_3^- with a higher affinity than either the G41S variant or the WT protein. As previously reported the relationship between k_{obs} and $[N_3^-]$ under the conditions employed is almost linear ¹, and is also the case here for the Y48H variant (Figure 3B). Assuming N_3^- binding to oxidized Cc is an S_N1 mechanism, in which the hexacoordinate heme form is in equilibrium with a pentacoordinate form, the latter being the form that binds N_3^- (Figure 3C), then, as outlined previously ¹, Equations 1 and 2 may be

derived that describe the K_{app} and the dependence of k_{obs} for N_3^- binding as a function of $[N_3^-]$.

$$K_{app} = K_D \left(\frac{K+1}{K} \right) \quad (1)$$

$$k_{obs} = \frac{(k_1 - k_{-2})[N_3^-]}{\left(\frac{k_1 + k_{-1}}{k'} \right) + [N_3^-]} + k_{-2} \quad (2)$$

In these equations, $K = k_1/k_{-1}$, where k_1 and k_{-1} are the rate constants for Met80 dissociation and association respectively and $K_D = k_{-2}/k_2$, with k_2 the pseudo-first order rate constant for N_3^- binding *i.e.* $k_2 = k'[N_3^-]$, where k' is the second order rate constant and k_{obs} is $\sim k_1$ at high $[N_3^-]$. Eq. 2 predicts that plots of k_{obs} vs $[N_3^-]$ (Figure 3B) follow hyperbolae that intercept the k_{obs} axis at the value of k_{-2} (N_3^- dissociation rate constant) and plateau at k_1 , the Met80 ligand dissociation rate constant. Fits of the normalized amplitude data are satisfactory and yield the values of K_{app} given in Table 2. Fitting the Y48H kinetic data to Eq. 2 presented a problem because it was not possible to explore a sufficiently high $[N_3^-]$ where the curve is expected to reach a plateau region and thus it was difficult to discriminate between linear and hyperbolic curves. Given this we have estimated a minimum value of k_1 that provides a fit (Figure 3B), with a value (125 s^{-1}) quoted without error in Table 2, and a reliable k_{-2} value (Table 2). Nevertheless, from this estimation of k_1 it is apparent that the Y48H variant has a Met80 off-rate significantly higher than the WT protein and the G41S variant. Furthermore, Eq. 1 shows that when $K < 1$, as here since the concentration of the pentacoordinate species is low, then K_{app} is $\sim K_D/K$. With the assumption that N_3^- binding to the pentacoordinate species is not directly affected by a distant mutation in the protein then K_D will be similar for both the WT protein and the Y48H variant, so that the ratio of K_{app} values ($^{WT}K_{app}/^{Y48H}K_{app}$) reflects the ratio of K values ($^{Y48H}K/^{WT}K$). On this basis $^{Y48H}K > ^{G41S}K > ^{WT}K$ and thus the pentacoordinate form of Y48H variant is more populated than in either the G41S variant or the WT protein.

The population of the peroxidase active pentacoordinate form of the Y48H variant is not increased upon binding CL

The peroxidase activity of the oxidized Y48H hCc variant was compared to that of the WT protein and the G41S variant under the same experimental conditions. From the reaction time-courses given in Figure 4A it is apparent that the WT protein displays a considerable lag-phase (~ 160 s following addition of H_2O_2) in the absence of CL-containing liposomes.




Figure 4: Peroxidase activity at pH 6.5. A) Illustrative reaction time-courses for the oxidation of ABTS in the presence of H_2O_2 with (+ CL) and without CL-containing liposomes. The vertical dashed line indicates the time point in each assay where H_2O_2 was added. B) The maximum peroxidation rates of ABTS with the various forms of ferriCc with (+ CL) and without CL-containing liposomes. The error bars are the standard error from measurements carried out in triplicate.

The lag phase is essentially absent for both the G41S and Y48H variants (Figure 4A). Peroxidase activity for the G41S variant is greater than that of the WT protein but less than that of the Y48H variant (Figure 4B). The progressive increase in peroxidase activity observed, correlates with the increase in less cooperative global unfolding (Figure 2A). In the presence of CL-containing liposomes the lag phase is now absent for the WT protein (Figure 4A) with a significant boost in peroxidase activity evident (Figure 4B). No change in peroxidase activity is observed for the Y48H variant in the presence of CL-containing liposomes, in contrast to the G41S variant where peroxidase activity is increased to a similar level as the Y48H variant (Figure 4B). The secondary increase would indicate that the binding of CL to this variant pushes the equilibrium between the hexa- and pentacoordinate heme forms further towards the latter (Figure 3C). A finding that further confirms the interpretation from the N_3^- binding kinetics, that for the Y48H variant the pentacoordinate form of the oxidized protein is energetically more favoured than it is for the G41S variant and the WT protein.

¹H NMR spectroscopy reveals an increase in heme pocket dynamics for the Y48H variant

The ¹H NMR spectrum of oxidized Cc has many resonances that are both downfield and upfield shifted due to their nuclei experiencing a hyperfine interaction from the unpaired




Figure 5: Paramagnetic ¹H NMR spectra in D₂O and pH 6.5 of oxidised hCc. A) WT Cc spectrum with several well resolved hyperfine shifted resonances arising from heme substituents and protein labelled with their assignment. B and C) Temperature dependence in the downfield and upfield regions of the spectra for the various hCc proteins. Change in chemical shift with increasing temperature for the 8- and 3-methyl heme substituents are indicated with dashed lines, along with those of the Met 80 CεH₃, the Met80 CγH and the His18 C₂H.

downfield shifted hyperfine resonances of known assignment labelled^{56, 57}. For both the WT

electron on the low-spin heme iron⁵⁴.

These paramagnetic signals can inform on the

electronic structure of the heme and also provide insight into heme-ligand dynamics and non-native heme-ligand states⁵⁵. The ¹H

NMR spectrum of the oxidized WT hCc is shown in Figure 5A with some of the upfield and

1
2
3 protein and G41S variant the pattern of chemical shifts for the heme-methyl signals is $8 > 3 >$
4 $5 > 1$ as expected⁵⁸. On increasing the temperature (from 283 to 313K) the hyperfine
5 chemical shift for the 8- and 3-methyl groups of the WT protein and G41S variant display a
6 Curie-type temperature dependence (Figure 5B), whereas the 5-methyl group displays an
7 anti-Curie dependence. Similar temperature dependence is observed for the upfield
8 hyperfine-shifted resonances assigned to protons of the axial His18 and Met80 heme-iron
9 ligands (Figure 5C). For the Y48H variant the spectrum is notably different in both the
10 upfield and downfield regions (Figure 5B and C). At 283 K there is a splitting of the
11 hyperfine peak found at ~ 36 ppm (Figure 5B). It is known that the 3-methyl group in certain
12 Cc's including horse heart, displays excess line broadening compared with the other heme
13 methyl peaks at room temperature and splits into two resonances at lower temperature^{59, 60}. A
14 view consistent with exchange broadening due to rapid interconversion between
15 heterogeneous protein environments around the 3-methyl group⁶⁰. The splitting into two
16 peaks at 283 K observed for the peak at ~ 36 ppm in the Y48H variant is therefore consistent
17 with it being assigned to the 3-methyl group (Figure 5B). Raising the temperature causes the
18 interconversion rate to increase and the two components become dynamically averaged
19 leading to a single broadened peak. (Figure 5B). Burns and La Mar⁶⁰ showed at 298 K and
20 pH 5.8 that the rate of the exchange process in oxidized horse heart Cc was $\sim 10^3$ s⁻¹ and a
21 similar rate for Y48H Cc at 283 K and pH 6.5 can be determined from the chemical shift
22 difference between the heme methyl 3 resonances in the two environments (Fig. 5B). A rate
23 considerably faster than the rate (k_1) we determine for the Met80-Fe(III) bond dissociation
24 (Table 2), showing that the unusual dynamic process affecting heme 3-methyl is not
25 associated with the formation of the pentacoordinate state. Furthermore, the Y48H variant
26 alters the pattern of heme methyl chemical shifts which now become $3 > 8 > 5 > 1$. Such an
27 order of heme methyl shifts has not been previously observed in mitochondrial or bacterial
28 oxidized Cc, where the order is either $8 > 3 > 5 > 1$ or $5 > 1 > 8 > 3$, noting that 8 is always > 3
29 and 5 is always > 1 in $S = \frac{1}{2}$ oxidized Cc⁶¹. These patterns are governed by the distribution
30 of the unpaired electron, which reflects the orientation of heme axial ligands and is largely
31 dependent on two conformers of the Met80 side chain related through inversion of the Met
32 thioether sulfur⁵⁸. In this context, we note that the upfield region of the spectrum where
33 hyperfine signals for protons of the His18 and Met80 sidechains are observed in the WT and
34 G41S spectrum is distinctly different in the Y48H variant. Thus, although the conformation
35 of the axial ligands is the same in the three X-ray structures the orientation of at least Met80
36 for the Y48H variant in solution is different. Bren *et al.* have demonstrated that fluxional
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 solution behaviour of the Met80 ligand is commonplace in Cc's ⁶¹. We suggest that the
4 altered fluxional behaviour causes a change in the magnetic susceptibility axes which
5 perturbs the relative chemical shifts of the 3- and 8-methyl resonances ⁶¹.
6
7

8 Finally, we note the presence of non-native hyperfine signals in both the G41S and
9 Y48H variants in the downfield region between 18 and 26 ppm at 303 K and above (Figure
10 5B). A similar set of non-native resonances has been observed for oxidized horse Cc at
11 temperatures greater than 315 K ⁶², along with the WT resonances, and been assigned to the
12 alkaline conformer with Lys and His coordination of the heme Fe(III). For the G41S and
13 Y48H variants we suggest that as the equilibrium between the hexa- and pentacoordinate
14 heme form is pushed further towards the latter species with increasing temperature these non-
15 native species similarly arise from a fraction of the protein that populates the alkaline
16 conformation. The fact that these resonances are not observed for the WT protein under
17 similar conditions is a further indication that these pro-apoptotic variants have a more labile
18 Met80-Fe(III) ligand. Note that the N₃⁻ binding study of the G41S variant ¹ showed that
19 under the conditions employed for these measurements the alkaline conformer was not
20 significantly populated. With the higher protein concentrations and longer experimental times
21 needed for the NMR experiments it is clear that at elevated temperatures the alkaline
22 conformer becomes populated.
23
24
25
26
27
28
29
30
31
32
33

34 *The Y48H variant has increased main-chain mobility relative to the G41S variant*

35 To explore in more detail the dynamics of the oxidized Y48H variant, ¹⁵N backbone amide
36 H/D exchange and ¹⁵N relaxation studies using NMR spectroscopy were carried out. The
37 [¹H,¹⁵N]-HSQC spectrum of the Y48H variant is shown in Figure 6A overlaid with the WT
38 protein spectrum under identical conditions ¹. Clear differences between the two spectra exist
39 (Figure 6A). Most notable is the significant number of backbone amide resonances absent in
40 the Y48H spectrum compared with the WT spectrum, with a total of 64 backbone amide
41 resonances out of a possible 99 non-proline residues being assigned (Table S1). The missing
42 resonances are found predominately in the 40-57 Ω-loop encompassing helix-II and the 71-
43 85 Ω-loop. At a lower magnetic field strength (500 MHz) no new peaks in the Y48H
44 [¹H,¹⁵N]-HSQC spectrum arise, which coupled with the increased line widths of the
45 backbone amide resonances at 800 MHz is indicative of the Y48H variant having
46 significantly greater conformational heterogeneity compared to the WT protein ^{1, 37}.
47
48
49
50
51
52
53
54
55

56 H/D-exchange behaviour of the oxidized Y48H hCc variant was monitored and
57
58
59
60

backbone amide PFs, defined as the ratio between the observed experimental and intrinsic




Figure 6: Backbone amide assignments and amide proton H/D exchange analysis of the oxidized Y48H hCc variant. A) An overlay of the 800 MHz $[^1\text{H}, ^{15}\text{N}]$ -HSQC spectrum of the Y48H variant (blue) with the WT protein spectrum (orange) at 288 K. Assignments for most of the assigned backbone amide resonances in the Y48H spectrum are indicated in blue. B) PFs for the Y48H variant plotted against residue number. For the PFs that could be calculated 3 residues underwent fast-exchange ($\log\text{PF} \leq 1$), 13 medium ($1 < \log\text{PF} \leq 2$), 10 slow ($2 < \log\text{PF}$) and 3 extremely slow-exchange (residual proton occupancy $> 80\%$ after ~ 134 h corresponding to exchange rates $< 0.003 \text{ h}^{-1}$, shown in red). Secondary structure (α -helix) is indicated by the bottom red squares with the 5 α -helices numbered I-V, while red squares above represent Pro residues. Unassigned and fast-exchanging residues are indicated in dark blue and yellow squares, respectively. Residues for which PFs could not be calculated due to the slowness of the exchange are indicated with black squares (3 residues).

exchange rates ($k_{\text{ex}}/k_{\text{in}}$) have been determined⁴⁰ (Table S1). Exchange rates are influenced by H-bonding, particularly in secondary structure and solvent accessibility⁶³. Out of the 64 assigned backbone amides 32 are fully exchanged within the first 10 min, leaving only 32 residues for which H/D exchange behaviour could be analysed. From these, PFs were calculated for 29 residues with 3 residues having exchange that is too slow to calculate within

the experimental timeframe (Figure 6B). The rapid H/D exchange of many backbone amide resonances and the absence of resonances for 28 backbone amides can be seen from the sequence dependence of the PFs plotted in Figure 6B for the Y48H variant. Only the interlocking helices-I and V along with portions of helix-III and the 32-36 turn maintain various levels of protection, with the remaining segments of the protein likely to be undergoing extensive dynamic fluctuations (Figure 6B).

^{15}N relaxation parameters, T_1 , T_2 and NOE were obtained for 55 residues, with plots of these parameters versus residue number shown in Figure 7 for the Y48H variant. The




Figure 7: ^{15}N relaxation parameters for the oxidized Y48H hCc variant acquired at 800 MHz and 288 K. A) T_1 and B) T_2 times plotted against residue number. C) R_2/R_1 ratio (black open circles) and $[^1\text{H}]-^{15}\text{N}$ NOE (blue circles) plotted against residue number.

average relaxation rates were $R_1 = 1.08 \pm 0.03 \text{ s}^{-1}$ and $R_2 = 14.99 \pm 0.72 \text{ s}^{-1}$ with an average NOE = 0.84 ± 0.06 . These values are similar to the WT protein ($R_1 = 0.99$, $R_2 = 14.23$, NOE = 0.84) and the G41S variant ($R_1 = 1.07$, $R_2 = 14.48$, NOE = 0.84)¹. The T_1 values, reporting on ns-ps motions are relatively constant, with the notable exceptions of His18 and Ile75 (Figure 7A). In contrast, the T_2 plot reveals a significant number of residues that deviate below the average (Figure 7B), probably reflecting contributions to relaxation from slower conformational exchange equilibria. A useful indicator to identify residues experiencing chemical exchange is a plot of the R_2/R_1 ratio (Figure 7C). Despite the smaller size of the relaxation data set for the Y48H variant (55 residues) compared to the G41S variant (92 residues)¹, a similar pattern of residues experiencing slow conformational exchange emerges (*i.e.* 14-36, 40-57 and 71-85 Ω -loops) and is also in keeping with the regions experiencing rapid H/D-exchange (Figure 6B).

DISCUSSION

1
2
3 The past two decades of Cc research has placed this essential redox protein firmly into the
4 category of a moonlighting or EMF protein ⁶⁴. For Cc to have a peroxidatic moonlighting
5 function, then access to an alternative structural state or conformer is required. Such a form
6 could come about because Cc binding of CL results in a conformational change to create the
7 peroxidatic state or because the ground-state structure is in equilibrium with one or more
8 minor higher-energy forms of different structure and CL binding displaces the equilibrium by
9 binding preferentially to a higher-energy state. We have shown for the pro-apoptotic G41S
10 variant of hCc enhanced conformational dynamics of the 40-57 Ω -loop exists and is directly
11 linked to the dissociation of the Met80-Fe(III) ligand in the 71-85 Ω -loop. Cross-talk
12 between Ω -loop results in an increased population of the peroxidatic pentacoordinate state ¹,
13 which under healthy conditions aids in controlling levels of reactive oxygen species (ROS),
14 but during the early phase of mitochondrial apoptosis results in Cc peroxidising the
15 membrane lipid CL. Furthermore, once released from the mitochondrion Cc has also been
16 implicated in the peroxidation of the plasma membrane lipid phosphatidylserine ⁶⁵. The
17 second thrombocytopenia 4 associated mutation of hCc to be discovered was the Y48H
18 variant ²⁵ and is also located within the 40-57 Ω -loop. In Cc evolution, this is the least
19 conserved Ω -loop and has the lowest free energy of the five-cooperative folding/unfolding
20 units (foldons) assigned in Cc ^{66, 67}. Perhaps not surprisingly therefore, the Y48H variant
21 displays enhanced mobility and dynamics. However, it is apparent that the dynamics, both
22 local and global, are further enhanced in the Y48H variant compared to the G41S variant.
23 Such behaviour is revealed by ~ 35 % of backbone amide resonances for the Y48H variant
24 remaining unassigned due to their absence in the ¹H-¹⁵N HSQC spectrum compared to 2 and
25 6 % for the WT protein and G41S variant, respectively, under identical conditions. Except for
26 Ala43 the whole of the 40-57 Ω -loop is unassigned, along with the latter half (residues 79-85)
27 of the 71-85 Ω -loop, the second least stable foldon ⁶⁷, housing the Met80-Fe(III) ligand.
28 These missing resonances in the Y48H variant are a likely consequence of a decreased T₂
29 resulting from exchange between different conformations. Although we cannot demonstrate
30 unequivocally from direct NMR measurements, because of the missing resonances, that the
31 Y48H variant has enhanced dynamics on the μ s-ms timescale (*i.e.* compute R_{ex} terms), it is
32 apparent for the resonances which allow for an amide H/D exchange PF to be calculated (*i.e.*
33 those from the N- and C-terminal helices, the end of the 14-36 Ω -loop and some of helix 3)
34 that these are generally smaller for the Y48H variant than for the other proteins ¹.
35 Visualization of the spread in increasing dynamics can be appreciated from mapping amide
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

H/D exchange data onto the respective Cc X-ray structures (Figure 8A). From this it is apparent that helices I, III and V for all proteins maintain similar H/D profiles with the mutations clearly resulting in more rapid exchange and thus enhanced dynamics (WT < G41S < Y48H) in the 40-57 and 71-85 Ω -loops, as well as helix II (Figure 8A).

Englander and colleagues have elegantly determined how Cc folds through uncovering its foldon substructure^{66, 67}. Of the five foldons, the two with the lowest stability are the 40-57 and 71-85 Ω loops^{5, 68}. The correlation between the conformational dynamics linking the G41S and Y48H mutations with the displacement of the Met80 ligand and the energetics of




Figure 8: Dynamics in the hCc variants attributed to causing thrombocytopenia 4. A) PF mapping onto the respective X-ray structures of hCc, with PFs colour coded as indicated. The data for the ferric WT protein and ferric G41S variant are taken from¹ with the data for the Y48H variant taken from the LogP plot in Figure 6B. The heme and axial ligands are shown in stick and helices labelled I to V. B) Energy diagrams illustrating the calculated differences in free energy change ($\Delta\Delta G$) between the pentacoordinate forms in the various species derived from $\Delta G = -RT\ln(\frac{K_{app}^{WT}}{K_{app}^{variant}})$ from N_3^- binding kinetics. The ΔG for the WT protein is based on Hu *et al.*⁵.

the foldon substructure is striking. Making the assumption that the pentacoordinate state of oxidized Cc is equivalent to the partially unfolded form, with the two lowest energy foldons unfolded, places the pentacoordinate state at 6 kcal/mol above the ground state structure, based on the analyses of Hu *et al.*⁵ and assuming the energetics of horse Cc are the same as those of hCc. From $\Delta G = -RT\ln(\frac{K_{app}^{WT}}{K_{app}^{Y48H}})$ (determined from the kinetics of N_3^- binding) the relative $\Delta\Delta G$ of the pentacoordinate state of the various Cc forms may be calculated as illustrated by the energy level diagrams in Figure 8B, since, as noted earlier, the ratio $\frac{K_{app}^{WT}}{K_{app}^{Y48H}}$ reflects the ratio of equilibrium constants for dissociation and re-association

1
2
3 of the Met80 ligand ($^{Y48H}K/^{WT}K$), with the assumption that the free energy of the ground state
4 is the same for both the variant and the WT protein. Thus, as illustrated in Figure 8B the
5 pentacoordinate form is 0.9 kcal/mol and 0.6 kcal/mol lower in energy than the WT protein
6 for the Y48H and G41S variants, respectively, than for the WT protein. Each kcal/mol
7 change denotes a \sim 5-fold change in the equilibrium constant governing the distribution
8 between the hexa- and pentacoordinated forms. If the minor component in such an
9 equilibrium, here the pentacoordinated form, plays a significant mechanistic role, this mean
10 that for a kcal/mol change in stability the concentration of this minor component increases \sim
11 5-fold. Our proposal that the G41S and Y48H variants cause small changes in the oxidized
12 hCc energy landscape should be considered alongside the recent finding that the Fe(III)-
13 S(Met) bond of Cc is weak and requires constraints imposed by the protein structure, such as
14 extensive H-bonding of neighbouring residues, to be maintained under physiological
15 conditions^{69, 70}. Stabilisation of a weak bond by a protein structure is an example of entasis,
16 originally proposed to account for some of the apparently unusual properties of
17 metalloproteins by suggesting that the metal centre ‘constituted an energetically poised
18 domain’⁷¹, and Mara *et al.*⁷⁰ have determined that the protein entatic contribution to the
19 Fe(III)-S(Met) bond is \sim 4 kcal/mol. It is readily appreciated therefore that small changes in
20 the energy landscape can have mechanistic effects and contribute to this moonlighting
21 function of Cc.
22
23
24
25
26
27
28
29
30
31
32
33

34
35 From the peroxidase assays it is particularly noteworthy that the turnover of H₂O₂ by
36 the Y48H variant is not boosted in the presence of CL-liposomes, as is the case for the WT
37 protein and the G41S variant (Figure 4B). An absence of a CL boosting effect for G41
38 variants has been reported by others²⁸ and thus contrasts with both our previous observations
39 using an Amplex red assay²⁷ and in the present study using ABTS. Our findings that there is
40 a boost in peroxidase activity for the G41S variant in the presence of CL-liposomes are
41 wholly consistent with our dynamic appraisal of these two variants. The Y48H variant is
42 more dynamic than G41S variant with the equilibrium between the hexa- and pentacoordinate
43 species no longer rate-limiting in this peroxidase assay for the Y48H variant as the
44 concentration of the reactive pentacoordinate form is further increased (Figure 8B).
45
46
47
48
49
50

51 Access of small molecules such as H₂O₂ to the heme has been suggested to occur
52 through a gating motion of Arg38 based on its side chain repositioning in the G41S variant
53 creating a ‘channel’ to enable H₂O molecules to access the heme^{27, 44}. A subsequent X-ray
54 structure of the K72A variant of *Saccharomyces cerevisiae* Cc revealed the Met80 ligand to
55 be dissociated from the heme iron and the Arg38 side chain positioned in an ‘out’ position as
56
57
58
59
60

1
2
3 observed in the G41S structure ⁷². A network of H-bonded H₂O molecules connects the
4 Met80 side of the heme to the bulk solvent in the K72A structure ⁷². Repositioning of Arg38
5 thus causes perturbations to the environment of the heme propionate substituent at position 7
6 and may be considered to influence the dynamics of the 40-57 Ω-loop, which is linked to the
7 dissociation of the Met80 ligand ^{1, 27}. However, we note that in the X-ray structure of the
8 Y48H variant the Arg38 side chain is in the same position as in the WT protein (Figure 1B).
9 Comparison of the G41S and Y48H X-ray structures reveals that the introduction of a Ser or
10 His at positions 41 and 48, respectively, creates the opportunity for both side chains to form a
11 H-bond to the side chain of Asn52 (Figure 1B). The latter communicates with the Met80
12 ligand through further H-bond interactions (Figure 1B). Thus, we propose that there are two
13 possible conformational routes by which dynamical changes at the base of the heme could be
14 communicated to the Met80 ligand. Indeed, the literature contains many reports showing that
15 the 40-57 Ω-loop and heme crevice have an important conformational linkage, though this
16 was not something always recognised at the time. For example, Robinson *et al.* ⁷³, based on
17 incomplete NMR data showed that residues in the 40-57 Ω-loop rapidly interconverted
18 between different local conformational states under native conditions. Our suggested role of
19 Asn52 is consistent with observations of Schejter *et al.* ⁷⁴, who present data on CN⁻ binding
20 to a N52I variant of *S. cerevisiae* Cc, that is indicative of a reduced population of the
21 pentacoordinate form. A linkage has also been discussed in connection with the alkaline
22 isomerisation of oxidized Cc ⁷⁵ and is supported by the NMR structure of a triple variant of *S.*
23 *cerevisiae* ferric Cc engineered to stabilize the alkaline conformer ⁷⁶. The triple variant
24 structure showed that the 71-85 Ω loop with the displaced Met80 ligand and the 40-57 Ω-
25 loop were the most perturbed feature compared with the WT structure. ⁷⁶

26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42 Recent studies have highlighted that Cc function can be governed by phosphorylation,
43 where it regulates the electron-transport chain flux preventing hyperpolarization of the
44 mitochondrial membrane potential, a known cause of ROS and trigger of apoptosis ^{77, 78}.
45 Three phosphorylation sites on isolates of bovine Cc have been identified from different
46 tissues and include Thr28, Tyr48 and Tyr97 ⁷⁹⁻⁸¹. Considering the reported pro-apoptotic
47 behavior of the Y48H variant ²⁵, it is interesting to note that one of the phosphorylation sites
48 is Tyr48, which when phosphorylated abolishes the capability to trigger apoptosis at the
49 apoptosome level ⁸²⁻⁸⁴. A recent study has characterized a Tyr48 phosphomimetic (designated
50 Tyr48p) using a non-native amino acid to assess the effect of phosphorylation at this position
51 on the hCc structure ⁸⁴. Structural studies were conducted with the ferrous form, which is the
52
53
54
55
56
57
58
59
60

1
2
3 heme oxidation state that donates an electron to Cc oxidase ¹², and not the state that gives rise
4 to the peroxidatic pentacoordinate form, central to our work. The WT reduced Cc has a more
5 rigid structure than its oxidized counterpart, and an equilibrium between the Met80-on and
6 Met80-off has not been detected under non-apoptosis physiological conditions ⁸⁵. Despite this,
7 the dynamics on the μ s-ms timescale for the reduced Tyr48p Cc showed increased mobility
8 compared to the WT protein, particularly, in the 40-57 Ω -loop, but with an overall increase
9 significantly lower than the G41S and Y48H variants in their oxidized forms. Furthermore,
10 an increase in mobility for the Tyr48p reduced Cc was also observed for residues 75-78 in the
11 71-85 Ω -loop again highlighting the link between the 40-57 Ω -loop and the Met80 ligand. In
12 the ferric state the Tyr48p exhibits a 3-fold increase in peroxidase activity relative to the WT
13 protein ⁸⁴, in contrasts to the Y48H variant where a 7-fold increase in activity is observed
14 (Figure 4B). In the presence of CL-liposomes the Tyr48p displays a 7-fold boost in
15 peroxidase activity relative to the WT protein and thus on a par with the Y48H variant in the
16 absence or presence of CL (Figure 4B). Therefore, these data suggest that the equilibrium
17 constant between the hexacoordinate and pentacoordinate heme forms (Figure 3C) favours
18 the hexacoordinate form in Tyr48p relative to the Y48H variant, enabling CL to further
19 regulate the activity of the phosphorylated protein which may have consequences for pro- and
20 anti-apoptotic behaviour insofar as we have shown that the pentacoordinate heme form is
21 critical for this.
22
23
24
25
26
27
28
29
30
31
32
33
34

35 In conclusion, we have shown that the naturally occurring Y48H variant of hCc in its
36 oxidized heme state is more peroxidatic than either the WT protein or the G41S variant that is
37 also implicated in thrombocytopenia. The dynamic fluctuations in the 40-57 Ω -loop resulting
38 in cross-talk with the 71-85 Ω -loop containing the Met80 ligand are further amplified in the
39 Y48H variant. Consequently, the Y48H protein has intrinsic peroxidase activity that is no
40 longer enhanced in the presence of CL because of the increased population of an excited
41 pentacoordinate heme state structure lying close in energy to the ground state Met80
42 coordinated structure. These finding underpins our proposal that the switch in function of the
43 moonlighting Cc from electron-transfer to peroxidase is, at least partly, a result of its pre-
44 existing conformational equilibrium involving an excited pentacoordinate state. A third
45 naturally occurring variant, A51V, also implicated in thrombocytopenia has not yet been
46 characterized. Owing to the location of the variant on the 40-57 Ω -loop it may be predicted to
47 have similar traits to the G41S and Y48H variants and it will therefore be interesting to
48
49
50
51
52
53
54
55
56
57
58
59
60

discover where this variant is located within the dynamic window (WT < G41S < Y48H) that has emerged from the studies so far.

ACKNOWLEDGEMENTS

Our work was supported by a Leverhulme Trust project grant (RPG-2013-164) to J.A.R.W. and a Leverhulme Trust emeritus fellowship (EM-2014-088) to G.R.M. Access to the Swiss Light Source was via Long-Term award 20160704 to MAH.

SUPPORTING INFORMATION

A supporting information Table is available, Table S1 that reports on the chemical shift assignments and the H/D exchange rates (k_{ex}), calculated protection factors (log P), free energy of exchange (ΔG_{ex}) and proton occupancy values (Occ) for oxidized Y48H hCc (pH 6.5 and 288 K).

REFERENCES

- [1] Karsisiotis, A. I., Deacon, O. M., Wilson, M. T., Macdonald, C., Blumenschein, T. M., Moore, G. R., and Worrall, J. A. (2016) Increased dynamics in the 40-57 Ω -loop of the G41S variant of human cytochrome c promote its pro-apoptotic conformation, *Sci Rep* 6, 30447.
- [2] Santoro, M. M., and Bolen, D. W. (1992) A test of the linear extrapolation of unfolding free energy changes over an extended denaturant concentration range, *Biochemistry* 31, 4901-4907.
- [3] Henderson, B., and Martin, A. C. (2014) Protein moonlighting: a new factor in biology and medicine, *Biochem Soc Trans* 42, 1671-1678.
- [4] Jeffery, C. J. (2014) An introduction to protein moonlighting, *Biochem Soc Trans* 42, 1679-1683.
- [5] Hu, W., Kan, Z. Y., Mayne, L., and Englander, S. W. (2016) Cytochrome c folds through foldon-dependent native-like intermediates in an ordered pathway, *Proc Natl Acad Sci U S A* 113, 3809-3814.
- [6] Jeffery, C. J. (1999) Moonlighting proteins, *Trends Biochem Sci* 24, 8-11.
- [7] Chapple, C. E., Robisson, B., Spinelli, L., Guien, C., Becker, E., and Brun, C. (2015) Extreme multifunctional proteins identified from a human protein interaction network, *Nat Commun* 6, 7412.
- [8] Chapple, C. E., and Brun, C. (2015) Redefining protein moonlighting, *Oncotarget* 6, 16812-16813.
- [9] Bryngelson, J. D., Onuchic, J. N., Socci, N. D., and Wolynes, P. G. (1995) Funnels, pathways, and the energy landscape of protein folding: a synthesis, *Proteins* 21, 167-195.
- [10] Henzler-Wildman, K., and Kern, D. (2007) Dynamic personalities of proteins, *Nature* 450, 964-972.
- [11] Baldwin, A. J., and Kay, L. E. (2009) NMR spectroscopy brings invisible protein states into focus, *Nat Chem Biol* 5, 808-814.

- 1
2
3 [12] Ferguson-Miller, S., and Babcock, G. T. (1996) Heme/Copper Terminal Oxidases, *Chem*
4 *Rev* 96, 2889-2908.
- 5 [13] Ow, Y. P., Green, D. R., Hao, Z., and Mak, T. W. (2008) Cytochrome c: functions
6 beyond respiration, *Nat Rev Mol Cell Biol* 9, 532-542.
- 7 [14] Huttemann, M., Pecina, P., Rainbolt, M., Sanderson, T. H., Kagan, V. E., Samavati, L.,
8 Doan, J. W., and Lee, I. (2011) The multiple functions of cytochrome c and their
9 regulation in life and death decisions of the mammalian cell: From respiration to
10 apoptosis, *Mitochondrion* 11, 369-381.
- 11 [15] Kagan, V. E., Tyurin, V. A., Jiang, J., Tyurina, Y. Y., Ritov, V. B., Amoscato, A. A.,
12 Osipov, A. N., Belikova, N. A., Kapralov, A. A., Kini, V., Vlasova, II, Zhao, Q., Zou,
13 M., Di, P., Svistunenko, D. A., Kurnikov, I. V., and Borisenko, G. G. (2005)
14 Cytochrome c acts as a cardiolipin oxygenase required for release of proapoptotic
15 factors, *Nat Chem Biol* 1, 223-232.
- 16 [16] Wang, X. (2001) The expanding role of mitochondria in apoptosis, *Genes Dev* 15, 2922-
17 2933.
- 18 [17] Bruey, J. M., Ducasse, C., Bonniaud, P., Ravagnan, L., Susin, S. A., Diaz-Latoud, C.,
19 Gurbuxani, S., Arrigo, A. P., Kroemer, G., Solary, E., and Garrido, C. (2000) Hsp27
20 negatively regulates cell death by interacting with cytochrome c, *Nat Cell Biol* 2, 645-
21 652.
- 22 [18] Boehning, D., Patterson, R. L., Sedaghat, L., Glebova, N. O., Kurosaki, T., and Snyder,
23 S. H. (2003) Cytochrome c binds to inositol (1,4,5) trisphosphate receptors,
24 amplifying calcium-dependent apoptosis, *Nat Cell Biol* 5, 1051-1061.
- 25 [19] Gonzalez-Arzola, K., Diaz-Moreno, I., Cano-Gonzalez, A., Diaz-Quintana, A.,
26 Velazquez-Campoy, A., Moreno-Beltran, B., Lopez-Rivas, A., and De la Rosa, M. A.
27 (2015) Structural basis for inhibition of the histone chaperone activity of SET/TAF-
28 Ibeta by cytochrome c, *Proc Natl Acad Sci U S A* 112, 9908-9913.
- 29 [20] Cummings, C., Walder, J., Treeful, A., and Jemmerson, R. (2006) Serum leucine-rich
30 alpha-2-glycoprotein-1 binds cytochrome c and inhibits antibody detection of this
31 apoptotic marker in enzyme-linked immunosorbent assay, *Apoptosis* 11, 1121-1129.
- 32 [21] Yu, T., Wang, X., Purring-Koch, C., Wei, Y., and McLendon, G. L. (2001) A mutational
33 epitope for cytochrome C binding to the apoptosis protease activation factor-1, *J Biol*
34 *Chem* 276, 13034-13038.
- 35 [22] Speck, S. H., Ferguson-Miller, S., Osheroff, N., and Margoliash, E. (1979) Definition of
36 cytochrome c binding domains by chemical modification: kinetics of reaction with
37 beef mitochondrial reductase and functional organization of the respiratory chain,
38 *Proc Natl Acad Sci U S A* 76, 155-159.
- 39 [23] Rieder, R., and Bosshard, H. R. (1980) Comparison of the binding sites on cytochrome c
40 for cytochrome c oxidase, cytochrome bc1, and cytochrome c1. Differential
41 acetylation of lysyl residues in free and complexed cytochrome c, *J Biol Chem* 255,
42 4732-4739.
- 43 [24] Morison, I. M., Cramer Borde, E. M., Cheesman, E. J., Cheong, P. L., Holyoake, A. J.,
44 Fichelson, S., Weeks, R. J., Lo, A., Davies, S. M., Wilbanks, S. M., Fagerlund, R. D.,
45 Ludgate, M. W., da Silva Tatley, F. M., Coker, M. S., Bockett, N. A., Hughes, G.,
46 Pippig, D. A., Smith, M. P., Capron, C., and Ledgerwood, E. C. (2008) A mutation of
47 human cytochrome c enhances the intrinsic apoptotic pathway but causes only
48 thrombocytopenia, *Nat Genet* 40, 387-389.
- 49 [25] De Rocco, D., Cerqua, C., Goffrini, P., Russo, G., Pastore, A., Meloni, F., Nicchia, E.,
50 Moraes, C. T., Pecci, A., Salviati, L., and Savoia, A. (2014) Mutations of cytochrome
51 c identified in patients with thrombocytopenia THC4 affect both apoptosis and
52 cellular bioenergetics, *Biochim Biophys Acta* 1842, 269-274.
- 53
54
55
56
57
58
59
60

- 1
2
3 [26] Johnson, B., Lowe, G. C., Futterer, J., Lordkipanidze, M., MacDonald, D., Simpson, M.
4 A., Sanchez-Guiu, I., Drake, S., Bem, D., Leo, V., Fletcher, S. J., Dawood, B., Rivera,
5 J., Allsup, D., Biss, T., Bolton-Maggs, P. H., Collins, P., Curry, N., Grimley, C.,
6 James, B., Makris, M., Motwani, J., Pavord, S., Talks, K., Thachil, J., Wilde, J.,
7 Williams, M., Harrison, P., Gissen, P., Mundell, S., Mumford, A., Daly, M. E.,
8 Watson, S. P., and Morgan, N. V. (2016) Whole exome sequencing identifies genetic
9 variants in inherited thrombocytopenia with secondary qualitative function defects,
10 *Haematologica* 101, 1170-1179.
11
12 [27] Rajagopal, B. S., Edzuma, A. N., Hough, M. A., Blundell, K. L., Kagan, V. E.,
13 Kapralov, A. A., Fraser, L. A., Butt, J. N., Silkstone, G. G., Wilson, M. T.,
14 Svistunenko, D. A., and Worrall, J. A. (2013) The hydrogen-peroxide-induced radical
15 behaviour in human cytochrome c-phospholipid complexes: implications for the
16 enhanced pro-apoptotic activity of the G41S mutant, *Biochem J* 456, 441-452.
17
18 [28] Josephs, T. M., Morison, I. M., Day, C. L., Wilbanks, S. M., and Ledgerwood, E. C.
19 (2014) Enhancing the peroxidase activity of cytochrome c by mutation of residue 41:
20 implications for the peroxidase mechanism and cytochrome c release, *Biochem J* 458,
21 259-265.
22
23 [29] Mason, J. M., Bendall, D. S., Howe, C. J., and Worrall, J. A. (2012) The role of a
24 disulfide bridge in the stability and folding kinetics of Arabidopsis thaliana
25 cytochrome c(6A), *Biochim Biophys Acta* 1824, 311-318.
26
27 [30] Kabsch, W. (2010) XDS, *Acta Cryst D*, 66, 125-132.
28
29 [31] Evans, P. R., and Murshudov, G. N. (2013) How good are my data and what is the
30 resolution?, *Acta Cryst D* 69, 1204-1214.
31
32 [32] McCoy, A. J., Grosse-Kunstleve, R. W., Adams, P. D., Winn, M. D., Storoni, L. C., and
33 Read, R. J. (2007) Phaser crystallographic software, *J Appl Crystallogr* 40, 658-674.
34
35 [33] Murshudov, G. N., Vagin, A. A., and Dodson, E. J. (1997) Refinement of
36 macromolecular structures by the maximum-likelihood method, *Acta Cryst D*, 53,
37 240-255.
38
39 [34] Emsley, P., and Cowtan, K. (2004) Coot: model-building tools for molecular graphics,
40 *Acta Cryst D* 60, 2126-2132.
41
42 [35] Davis, I. W., Leaver-Fay, A., Chen, V. B., Block, J. N., Kapral, G. J., Wang, X., Murray,
43 L. W., Arendall, W. B., 3rd, Snoeyink, J., Richardson, J. S., and Richardson, D. C.
44 (2007) MolProbity: all-atom contacts and structure validation for proteins and nucleic
45 acids, *Nucleic Acids Res* 35, W375-383.
46
47 [36] Krissinel, E. (2012) Enhanced fold recognition using efficient short fragment clustering,
48 *J Mol Biochem* 1, 76-85.
49
50 [37] Karsisiotis, A. I., Deacon, O. M., Rajagopal, B. S., Macdonald, C., Blumenschein, T. M.,
51 Moore, G. R., and Worrall, J. A. (2015) Backbone resonance assignments of ferric
52 human cytochrome c and the pro-apoptotic G41S mutant in the ferric and ferrous
53 states, *Biomol NMR Assign* 9, 415-419.
54
55 [38] Vranken, W. F., Boucher, W., Stevens, T. J., Fogh, R. H., Pajon, A., Llinas, M., Ulrich,
56 E. L., Markley, J. L., Ionides, J., and Laue, E. D. (2005) The CCPN data model for
57 NMR spectroscopy: development of a software pipeline, *Proteins* 59, 687-696.
58
59 [39] Goddard, T. D., and Kneller, D. G. (2008) SPARKY 3, *University of California, San*
60 *Francisco*.
61
62 [40] Bai, Y., Milne, J. S., Mayne, L., and Englander, S. W. (1993) Primary structure effects
63 on peptide group hydrogen exchange, *Proteins* 17, 75-86.
64
65 [41] Hoshino, M., Katou, H., Yamaguchi, K.-I., and Goto, Y. (2007) Dimethylsulfoxide-
66 quenched hydrogen/deuterium exchange method to study amyloid fibril structure,
67 *Biochim Biophys Acta* 1768, 1886-1899.

- 1
2
3 [42] Palmer, A. G., Rance, M., and Wright, P. E. (1991) Intramolecular Motions of a Zinc
4 Finger DNA-Binding Domain from Xfin Characterized by Proton-Detected Natural
5 Abundance C-12 Heteronuclear Nmr-Spectroscopy, *J Am Chem Soc* 113, 4371-4380.
- 6 [43] Delaglio, F., Grzesiek, S., Vuister, G. W., Zhu, G., Pfeifer, J., and Bax, A. (1995)
7 NMRPipe: a multidimensional spectral processing system based on UNIX pipes, *J*
8 *Biomol NMR* 6, 277-293.
- 9
10 [44] Liptak, M. D., Fagerlund, R. D., Ledgerwood, E. C., Wilbanks, S. M., and Bren, K. L.
11 (2011) The Proapoptotic G41S Mutation to Human Cytochrome c Alters the Heme
12 Electronic Structure and Increases the Electron Self-Exchange Rate, *J Am Chem Soc*
13 133, 1153-1155.
- 14 [45] Zeldin, O. B., Gerstel, M., and Garman, E. F. (2013) RADDPOSE-3D: time- and space-
15 resolved modelling of dose in macromolecular crystallography, *J Appl Crystallogr* 46,
16 1225-1230.
- 17 [46] Beitlich, T., Kuhnel, K., Schulze-Briese, C., Shoeman, R. L., and Schlichting, I. (2007)
18 Cryoradiolytic reduction of crystalline heme proteins: analysis by UV-Vis
19 spectroscopy and X-ray crystallography, *J Synchro Rad* 14, 11-23.
- 20 [47] Kekilli, D., Moreno-Chicano, T., Chaplin, A. K., Horrell, S., Dworkowski, F. S. N.,
21 Worrall, J. A. R., Strange, R. W., and Hough, M. A. (2017) Photoreduction and
22 validation of haem-ligand intermediate states in protein crystals by in situ single-
23 crystal spectroscopy and diffraction, *IUCrJ* 4, 263-270.
- 24 [48] Takano, T., and Dickerson, R. E. (1981) Conformation change of cytochrome c. II.
25 Ferricytochrome c refinement at 1.8 Å and comparison with the ferrocyclochrome
26 structure, *J Mol Biol* 153, 95-115.
- 27 [49] Brayer, G. D., and Murphy, M. P. (1996) Structural studies of eukaryotic cytochromes c,
28 In *Cytochrome c: A multidisciplinary approach* (Scott, R. A., and Mauk, A. G., Eds.) 1
29 ed., pp 103-166, University Science Books, Sausalito, California.
- 30 [50] Shortle, D. (1995) Staphylococcal nuclease: a showcase of m-value effects, *Advances in*
31 *protein chemistry* 46, 217-247.
- 32 [51] Hammack, B. N., Smith, C. R., and Bowler, B. E. (2001) Denatured state
33 thermodynamics: residual structure, chain stiffness and scaling factors, *J Mol Biol*
34 311, 1091-1104.
- 35 [52] Godbole, S., Dong, A., Garbin, K., and Bowler, B. E. (1997) A lysine 73-->histidine
36 variant of yeast iso-1-cytochrome c: evidence for a native-like intermediate in the
37 unfolding pathway and implications for m value effects, *Biochemistry* 36, 119-126.
- 38 [53] Rosell, F. I., Ferrer, J. C., and Mauk, A. G. (1998) Proton-linked protein conformational
39 switching: Definition of the alkaline conformational transition of yeast iso-1-
40 ferricytochrome c, *J Am Chem Soc* 120, 11234-11245.
- 41 [54] Wuthrich, K. (1969) High-resolution proton nuclear magnetic resonance spectroscopy of
42 cytochrome, *Proc Natl Acad Sci U S A* 63, 1071-1078.
- 43 [55] Ubbink, M., Worrall, J. A., Canters, G. W., Groenen, E. J., and Huber, M. (2002)
44 Paramagnetic resonance of biological metal centers, *Annu Rev Biophys Biomol Struct*
45 31, 393-422.
- 46 [56] Santos, H., and Turner, D. L. (1987) Proton NMR studies of horse ferricytochrome c.
47 Completion of the assignment of the well resolved hyperfine shifted resonances,
48 *FEBS Lett* 226, 179-185.
- 49 [57] Feng, Y. Q., Roder, H., and Englander, S. W. (1990) Assignment of paramagnetically
50 shifted resonances in the ¹H NMR spectrum of horse ferricytochrome c, *Biophys J* 57,
51 15-22.
- 52
53
54
55
56
57
58
59
60

- 1
2
3 [58] Senn, H., and Wuthrich, K. (1985) Amino acid sequence, haem-iron co-ordination
4 geometry and functional properties of mitochondrial and bacterial c-type
5 cytochromes, *Q Rev Biophys* 18, 111-134.
- 6 [59] Burns, P. D., and Lamar, G. N. (1979) Detection of Localized Conformational
7 Flexibility in Horse Heart Cytochrome-C by Proton Nuclear Magnetic-Resonance, *J*
8 *Amer Chem Soc* 101, 5844-5846.
- 9 [60] Burns, P. D., and La Mar, G. N. (1981) Characterization of conformational
10 heterogeneity in the heme pocket of ferricytochrome c using high field proton nuclear
11 magnetic resonance spectroscopy, *J Biol Chem* 256, 4934-4939.
- 12 [61] Zhong, L., Wen, X., Rabinowitz, T. M., Russell, B. S., Karan, E. F., and Bren, K. L.
13 (2004) Heme axial methionine fluxionality in *Hydrogenobacter thermophilus*
14 cytochrome c552, *Proc Natl Acad Sci U S A* 101, 8637-8642.
- 15 [62] Angstrom, J., Moore, G. R., and Williams, R. J. P. (1982) The Magnetic-Susceptibility
16 of Ferricytochrome-C, *Biochim Biophys Acta* 703, 87-94.
- 17 [63] Englander, S. W., and Kallenbach, N. R. (1983) Hydrogen-Exchange and Structural
18 Dynamics of Proteins and Nucleic-Acids, *Q Revs Biophys* 16, 521-655.
- 19 [64] Hannibal, L., Tomasina, F., Capdevila, D. A., Demicheli, V., Tortora, V., Alvarez-Paggi,
20 D., Jemmerson, R., Murgida, D. H., and Radi, R. (2016) Alternative Conformations of
21 Cytochrome c: Structure, Function, and Detection, *Biochemistry* 55, 407-428.
- 22 [65] Bayir, H., Fadeel, B., Palladino, M. J., Witasp, E., Kurnikov, I. V., Tyurina, Y. Y.,
23 Tyurin, V. A., Amoscato, A. A., Jiang, J., Kochanek, P. M., DeKosky, S. T.,
24 Greenberger, J. S., Shvedova, A. A., and Kagan, V. E. (2006) Apoptotic interactions
25 of cytochrome c: redox flirting with anionic phospholipids within and outside of
26 mitochondria, *Biochim Biophys Acta* 1757, 648-659.
- 27 [66] Krishna, M. M., Lin, Y., Rumbley, J. N., and Englander, S. W. (2003) Cooperative
28 omega loops in cytochrome c: role in folding and function, *J Mol Biol* 331, 29-36.
- 29 [67] Krishna, M. M., Maity, H., Rumbley, J. N., Lin, Y., and Englander, S. W. (2006) Order
30 of steps in the cytochrome c folding pathway: evidence for a sequential stabilization
31 mechanism, *J Mol Biol* 359, 1410-1419.
- 32 [68] Maity, H., Maity, M., Krishna, M. M., Mayne, L., and Englander, S. W. (2005) Protein
33 folding: the stepwise assembly of foldon units, *Proc Natl Acad Sci U S A* 102, 4741-
34 4746.
- 35 [69] Kroll, T., Hadt, R. G., Wilson, S. A., Lundberg, M., Yan, J. J., Weng, T. C., Sokaras, D.,
36 Alonso-Mori, R., Casa, D., Upton, M. H., Hedman, B., Hodgson, K. O., and Solomon,
37 E. I. (2014) Resonant inelastic X-ray scattering on ferrous and ferric bis-imidazole
38 porphyrin and cytochrome c: nature and role of the axial methionine-Fe bond, *J Am*
39 *Chem Soc* 136, 18087-18099.
- 40 [70] Mara, M. W., Hadt, R. G., Reinhard, M. E., Kroll, T., Lim, H., Hartsock, R. W., Alonso-
41 Mori, R., Chollet, M., Glowina, J. M., Nelson, S., Sokaras, D., Kunnus, K., Hodgson,
42 K. O., Hedman, B., Bergmann, U., Gaffney, K. J., and Solomon, E. I. (2017)
43 Metalloprotein entatic control of ligand-metal bonds quantified by ultrafast x-ray
44 spectroscopy, *Science* 356, 1276-1280.
- 45 [71] Vallee, B. L., and Williams, R. J. (1968) Metalloenzymes: the entatic nature of their
46 active sites, *Proc Natl Acad Sci U S A* 59, 498-505.
- 47 [72] McClelland, L. J., Mou, T. C., Jeakins-Cooley, M. E., Sprang, S. R., and Bowler, B. E.
48 (2014) Structure of a mitochondrial cytochrome c conformer competent for
49 peroxidase activity, *Proc Natl Acad Sci U S A* 111, 6648-6653.
- 50 [73] Robinson, M. N., Boswell, A. P., Huang, Z. X., Eley, C. G., and Moore, G. R. (1983)
51 The conformation of eukaryotic cytochrome c around residues 39, 57, 59 and 74,
52 *Biochem J* 213, 687-700.
- 53
54
55
56
57
58
59
60

- 1
2
3 [74] Schejter, A., Koshy, T. I., Luntz, T. L., Sanishvili, R., Vig, I., and Margoliash, E. (1994)
4 Effects of mutating Asn-52 to isoleucine on the haem-linked properties of cytochrome
5 c, *Biochem J* 302 (Pt 1), 95-101.
- 6 [75] Maity, H., Rumbley, J. N., and Englander, S. W. (2006) Functional role of a protein
7 foldon--an Ω -loop foldon controls the alkaline transition in ferricytochrome c,
8 *Proteins* 63, 349-355.
- 9 [76] Assfalg, M., Bertini, I., Dolfi, A., Turano, P., Mauk, A. G., Rosell, F. I., and Gray, H. B.
10 (2003) Structural model for an alkaline form of ferricytochrome C, *J Am Chem Soc*
11 125, 2913-2922.
- 12 [77] Huttemann, M., Lee, I., Pecinova, A., Pecina, P., Przyklenk, K., and Doan, J. W. (2008)
13 Regulation of oxidative phosphorylation, the mitochondrial membrane potential, and
14 their role in human disease, *J Bioenerg Biomembr* 40, 445-456.
- 15 [78] Huttemann, M., Lee, I., Grossman, L. I., Doan, J. W., and Sanderson, T. H. (2012)
16 Phosphorylation of mammalian cytochrome c and cytochrome c oxidase in the
17 regulation of cell destiny: respiration, apoptosis, and human disease, *Adv Exp Med*
18 *Biol* 748, 237-264.
- 19 [79] Lee, I., Salomon, A. R., Yu, K., Doan, J. W., Grossman, L. I., and Huttemann, M. (2006)
20 New prospects for an old enzyme: mammalian cytochrome c is tyrosine-
21 phosphorylated in vivo, *Biochemistry* 45, 9121-9128.
- 22 [80] Yu, H., Lee, I., Salomon, A. R., Yu, K., and Huttemann, M. (2008) Mammalian liver
23 cytochrome c is tyrosine-48 phosphorylated in vivo, inhibiting mitochondrial
24 respiration, *Biochim Biophys Acta* 1777, 1066-1071.
- 25 [81] Mahapatra, G., Varughese, A., Ji, Q., Lee, I., Liu, J., Vaishnav, A., Sinkler, C.,
26 Kapralov, A. A., Moraes, C. T., Sanderson, T. H., Stemmler, T. L., Grossman, L. I.,
27 Kagan, V. E., Brunzelle, J. S., Salomon, A. R., Edwards, B. F., and Huttemann, M.
28 (2017) Phosphorylation of Cytochrome c Threonine 28 Regulates Electron Transport
29 Chain Activity in Kidney: IMPLICATIONS FOR AMP KINASE, *J Biol Chem* 292,
30 64-79.
- 31 [82] Pecina, P., Borisenko, G. G., Belikova, N. A., Tyurina, Y. Y., Pecinova, A., Lee, I.,
32 Samhan-Arias, A. K., Przyklenk, K., Kagan, V. E., and Huttemann, M. (2010)
33 Phosphomimetic substitution of cytochrome C tyrosine 48 decreases respiration and
34 binding to cardiolipin and abolishes ability to trigger downstream caspase activation,
35 *Biochemistry* 49, 6705-6714.
- 36 [83] Garcia-Heredia, J. M., Diaz-Moreno, I., Diaz-Quintana, A., Orzaez, M., Navarro, J. A.,
37 Hervas, M., and De la Rosa, M. A. (2012) Specific nitration of tyrosines 46 and 48
38 makes cytochrome c assemble a non-functional apoptosome, *FEBS Lett* 586, 154-158.
- 39 [84] Moreno-Beltran, B., Guerra-Castellano, A., Diaz-Quintana, A., Del Conte, R., Garcia-
40 Maurino, S. M., Diaz-Moreno, S., Gonzalez-Arzola, K., Santos-Ocana, C.,
41 Velazquez-Campoy, A., De la Rosa, M. A., Turano, P., and Diaz-Moreno, I. (2017)
42 Structural basis of mitochondrial dysfunction in response to cytochrome c
43 phosphorylation at tyrosine 48, *Proc Natl Acad Sci U S A* 114, 3041-3050.
- 44 [85] Moore, G. R., and Pettigrew, G. W. (1990) Cytochrome c: Evolutionary, Structural and
45 Physicochemical Aspects, *Springer-Verlag, London*.
- 46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **For Table of Contents Use Only**
4

5 **Heightened dynamics of the oxidized Y48H variant of human cytochrome *c* increases its**
6 **peroxidatic activity**
7
8

9
10 Oliver M. Deacon, Andreas Ioannis Karsisiotis, Tadeo Moreno-Chicano, Michael A. Hough,
11 Colin Macdonald, Tharin M.A. Blumenschein, Michael T. Wilson, Geoffrey R. Moore,
12 Jonathan A.R. Worrall
13
14

