

ACCEPTED MANUSCRIPT • OPEN ACCESS

Temperature and precipitation extremes under current, 1.5°C and 2.0°C global warming above pre-industrial levels over Botswana, and implications for climate change vulnerability

To cite this article before publication: Tiro Nkemelang *et al* 2018 *Environ. Res. Lett.* in press <https://doi.org/10.1088/1748-9326/aac2f8>

Manuscript version: Accepted Manuscript

Accepted Manuscript is “the version of the article accepted for publication including all changes made as a result of the peer review process, and which may also include the addition to the article by IOP Publishing of a header, an article ID, a cover sheet and/or an ‘Accepted Manuscript’ watermark, but excluding any other editing, typesetting or other changes made by IOP Publishing and/or its licensors”

This Accepted Manuscript is © 2018 The Author(s). Published by IOP Publishing Ltd.

As the Version of Record of this article is going to be / has been published on a gold open access basis under a CC BY 3.0 licence, this Accepted Manuscript is available for reuse under a CC BY 3.0 licence immediately.

Everyone is permitted to use all or part of the original content in this article, provided that they adhere to all the terms of the licence <https://creativecommons.org/licences/by/3.0>

Although reasonable endeavours have been taken to obtain all necessary permissions from third parties to include their copyrighted content within this article, their full citation and copyright line may not be present in this Accepted Manuscript version. Before using any content from this article, please refer to the Version of Record on IOPscience once published for full citation and copyright details, as permissions may be required. All third party content is fully copyright protected and is not published on a gold open access basis under a CC BY licence, unless that is specifically stated in the figure caption in the Version of Record.

View the [article online](#) for updates and enhancements.

1 Temperature and Precipitation Extremes under current, 1.5°C and
2 2.0°C Global Warming above Pre-Industrial Levels over Botswana,
3 and Implications for Climate Change Vulnerability

5 Tiro Nkemelang^{1,2}, Mark New^{2,3,4}, Modathir Zaroug^{5,2,6}

6 ¹Botswana Institute for Technology Research and Innovation, Botswana. (ORDIC ID: 0000-0002-9588-7382)

7 ²African Climate and Development Initiative, University of Cape Town, South Africa

8 ³Department of Environmental and Geographical Science, University of Cape Town, South Africa

9 ⁴School of International Development, University of East Anglia, UK

10 ⁵Nile Basin Initiative, Uganda

11 ⁶Climate System Analysis Group, University of Cape Town, South Africa

12 **Abstract**

14 Climate extremes are widely projected to become more severe as the global climate continues to warm
15 due to anthropogenic greenhouse gas emissions. These extremes often cause the most severe impacts on
16 society. Therefore, the extent to which the extremes might change at regional level as the global climate
17 warms from current levels to proposed policy targets of 1.5 and 2.0 °C above preindustrial levels need to
18 be understood to allow for better preparedness and informed policy formulation. This paper analysed
19 projected changes in temperature and precipitation extremes at 1.0, 1.5 and 2.0 °C warming over
20 Botswana, a country highly vulnerable to the impacts of climate change. Projected changes in temperature
21 extremes are significantly different from each other at the three levels of global warming, across three
22 main climatic zones in the country. Specifically, at 2.0 °C global warming relative to preindustrial, for the
23 ensemble median: (a) country average Warm Spell Duration Index (WSDI) increases by 80, 65, 62 days per
24 year across different climatic zones, approximately three (and two) times the change at 1.0 (1.5) °C; (b)
25 cold night (TN10P) and cold day (TX10P) frequencies decrease by 12 and 9 days per year across all regions,
26 respectively, while hot nights (TN90P) and hot days (TX90P) both increase by 8-9 days across all regions.
27 Projected changes in drought-related indices are also distinct at different warming levels. Specifically: (a)
28 projected mean annual precipitation decreases across the country by 5-12% at 2°C, 3-8% at 1.5 °C and 2-
29 7% at 1.0 °C; (b) dry spell length (ALTCDD) increases by 15-19 days across the three climatic zones at 2.0
30 °C, about three (and two) times as much as the increase at 1.0 (1.5) °C. Ensemble mean projections
31 indicate increases in heavy rainfall indices, but the inter-model spread is large, with no consistent direction
32 of change, and so changes are not statistically significant. The implications of these changes in extreme
33 temperature and precipitation for key socio-economic sectors are explored, and reveal progressively
34 severe impacts, and consequent adaptation challenges for Botswana as the global climate warms from its
35 present temperature of 1.0 °C above preindustrial levels to 1.5, and then 2.0 °C.

1. Introduction

Some of the harshest impacts of climate change are likely to occur through more extreme climate and weather events (Trenberth, 2012). Sub-Saharan Africa is among the most vulnerable of regions (Boko *et al.*, 2008; Niang *et al.*, 2014), with 34 of the top 50 most climate-vulnerable countries located in the continent (ND-GAIN, 2016). Therefore, climate change and extremes will continue to pose significant social and economic pressures on livelihoods within the continent under a warming global climate (King *et al.*, 2015; ASC, 2016). The frequency and intensity of extreme weather events is set to change, often increasing, across most regions thereby increasing vulnerability of those already exposed and sensitive especially rural communities (McElroy and Baker, 2012; World Bank, 2013). These extreme events, including heat waves (Ceccherini *et al.*, 2017), heavy precipitation events causing floods (Cook *et al.*, 2004; Reason, 2007; Manhique *et al.*, 2015; Moyo and Nangombe, 2015) as well as droughts (Richard *et al.*, 2001; Rouault and Richard, 2005) are already common drivers of vulnerability in Southern Africa.

The 2015 Paris climate change accord, drawing on growing evidence from successive IPCC reports, agreed to work to keeping Global Mean Surface Temperature (GMST) warming due to greenhouse gas and other emissions to below 2.0 °C, with the ambition to keep temperature change below 1.5 °C of preindustrial temperature (UNFCCC, 2015). Based on these two temperature warmings, various studies have been conducted on shifts in climates of different localities across the globe, assessing the extent to which an additional 0.5 °C temperature rise would influence both the mean and the climate extremes (e.g., James *et al.*, 2017; King *et al.*, 2017; Sanderson *et al.*, 2017; Wang *et al.*, 2017). With some studies proposing that the 1.5 °C warming level could be crossed as early as 2026 (Henley and King, 2017), the need for early action becomes even more critical. This therefore calls for more detailed analysis of the potential regional and local changes in extremes that might accompany these levels of global temperature increase, especially in the most vulnerable regions of the world.

Previous work on assessing observed and projected trends in climatic extremes have largely been based on a set of 27 temperature and precipitation indices, devised by the Expert Team on Climate Change Detection and Indices (ETCCDI) (Zhang *et al.*, 2011). Several studies have shown changes in these indices over the last several decades across the globe, based on observed data (e.g., Frich *et al.*, 2002; Alexander *et al.*, 2006; Manton, 2010) as well reanalysis datasets (e.g., Sillmann *et al.*, 2013a). These studies have shown a general trend towards more extreme weather under climate change. For future climate change, a number of studies have looked at the projected changes in both extremes of temperature (e.g., Sillmann *et al.*, 2013b; Lewis and King, 2017) and precipitation (Sillmann *et al.*, 2013b) globally. Most of these studies looked at these changes for a particular period in the 21st century following a particular Representative Concentration Pathway (RCP) (Fischer *et al.*, 2013; Lewis and King, 2017) or emissions scenario (Kharin *et al.*, 2007).

At regional scales, a number of studies have been carried out to assess both the historical and projected future trends in temperature and precipitation extremes (e.g., You *et al.*, 2011; Monier and Gao, 2015; Dosio, 2016; Razavi *et al.*, 2016). For Southern Africa, New *et al.* (2006) investigated the historical context of extremes from observations, while Shongwe *et al.* (2009) and Pinto *et al.* (2016) looked at the historical and future projection of precipitation extremes from reanalysis and downscaled climate model outputs.

1 There has so far been little research on the relative impact of global warming of 1.5 and 2.0 °C on climate
2 extremes at the national or sub-national scale.

3 Botswana, though highly exposed and vulnerable to the impacts of climate change, is under-researched
4 on physical climate change, especially on the evolution of climate extremes. A recent study ranked
5 Botswana among the top few countries in Africa in terms of national mean temperature and precipitation
6 changes as global temperature rises to 1.5 and 2.0 °C (Zaroug *et al.*, in review). This potential vulnerability
7 is reflected in policy and practice in Botswana: Hambira and Saarinen (2015) note the widespread
8 agreement among policy makers on the perceived worsening climate patterns in Botswana; Akinyemi
9 (2017) reports similar perceptions amongst farming communities in Eastern Botswana; in both these
10 cases, there is no quantitative research available to back up these views. This paper therefore aims to
11 quantify the projected changes in climate extremes that can be expected at the global temperature
12 targets agreed in Paris, relative to both pre-industrial and present-day global temperature levels. The
13 paper then discusses the implications of these projected changes for vulnerability of key sectors in
14 Botswana.

15 2. Data and Methodology

16 2.1. Study Area

17 Botswana is a landlocked, subtropical country in central Southern Africa (*Figure 1a*), sharing borders with
18 South Africa to the south, Namibia to the west, Zambia to the north and Zimbabwe to the north east. The
19 subsiding limb of the tropical Hadley circulation defines much of the climate in Botswana, with a semi-
20 permanent high pressure system (the Botswana high) persistent over the region (Driver and Reason,
21 2017). The south-western part of the country is hyper-arid, receiving the lowest rainfall, with aridity then
22 decreasing to North and East of the country (*Figure 1b*). Temperatures in Botswana are lowest during the
23 austral winter and highest during the austral summer. In winter, the occasional passing of westerly cold
24 frontal systems can cause minimum temperatures to fall to below freezing resulting in frosts over most
25 parts of the country (Andringa, 1984). Summer mean maximum temperatures range between 30.9 – 33.0
26 °C across the country, the western parts of the country being hotter (Moses, 2017). Occasional heat waves
27 are also experienced in austral summer with temperatures reaching highs of over 42 °C in some places
28 (Moses, 2017). Botswana's interannual climate variability is mainly driven by the El Nino Southern
29 Oscillation (ENSO), where El-Nino events are associated with dry conditions (Nicholson *et al.*, 2001) and
30 La Nina events are associated with wet years (Mason and Jury, 1997). Rain bearing weather systems
31 affecting the country include among others temperate tropical troughs (TTT) (Williams *et al.*, 2007) and
32 mesoscale convective systems(MCCs) (Blamey and Reason, 2012). These systems are mainly convection
33 driven and associated with the southward movement of the Intertropical Convergence Zone (ITCZ) in the
34 austral summer months (mainly November to March) (Bhalotra, 1987). Tropical depressions that form in
35 the Indian Ocean also occasionally penetrate the subcontinent from the east, bringing along heavy
36 precipitation (Reason and Keibel, 2004; Reason, 2007). Some heavy precipitation events could be a
37 combination of various weather systems such as westerly troughs (cold fronts), tropical lows and ridging
38 anticyclones creating conducive environment for such events (Crimp and Mason, 1999). Cut off lows are

also a common occurrence over the region, they tend to bring along heavy precipitation episodes that can cause flooding in some places in Southern Africa including Botswana (Singleton and Reason, 2007; Favre et al., 2013; Molekwa, 2013). Cut off lows also tend to be associated with extreme rainy days that fall outside of the normal rainy season (Favre et al., 2013). Dry spells within the rainy season are also a common occurrence in Botswana (Vossen, 1990), being more frequent during ENSO warm phases and over the south-western parts of the country (Usman and Reason, 2004).




Figure 1: (a)-Map of Southern Africa with Botswana highlighted (Mapchart.net). (b) Mean annual precipitation (1979-2013) in mm over Botswana derived from the WFDEI dataset and the 3 regions of homogeneous monthly rainfall in Botswana (Region 1, 2 and 3)

2.2. Data

The expert Team on Climate Change Detection and Indices (ETCCDI) temperature and precipitation extreme indices derived from the fifth version of Coupled Model Intercomparison Project (CMIP5) program participating models were analyzed. A total of 24 CMIP5 Global Climate Model (GCM) outputs developed by Sillmann et al. (2013a) (Table S1), were downloaded from KNMI Climate Explorer website database, <https://climexp.knmi.nl> (Trouet and Van Oldenborgh, 2013). A multi-model ensemble is used to account for uncertainty in the projections of the climate extremes, as both the ensemble mean, and associated spread, provide a more robust assessment of signal and noise than results from one or a few models (Tebaldi and Knutti, 2007; Knutti et al., 2010; IPCC, 2012).

The indices are available re-gridded to a common spatial resolution of $2.5^0 \times 2.5^0$ from their native model resolutions (Sillmann et al., 2013b). Historical simulations (1861 – 2005) combined with the high emissions scenario Representative Concentration Pathway that projects an 8.5 W/m^2 radiative forcing on the climate system by 2100 (RCP8.5) (Taylor et al., 2012) are chosen for analysis. This RCP is chosen as the forcing is sufficiently intense to guarantee all models reach a warming of $2.0 \text{ }^\circ\text{C}$ by the end of their simulations; further it is most representative of current forcing trends (Lewis and King, 2017). As described below, we analyze indices from individual models at the time they reach specific global temperature targets, so the forcing scenario is not particularly important. Additionally, Pendergrass et al.

(2015) and Shi *et al.* (2017) have shown that, in general, changes in climate extremes are indistinguishable across different RCPs when using multi-model ensembles; although Wang *et al.* (2017) shows that differences in regional aerosol emissions do produce differences in projected changes over high emission areas (not Southern Africa).

This study focuses on extreme indices that relate directly to climate change vulnerability in Botswana (*Table 1*) as determined from a review of vulnerability to climate in semi-arid countries of Southern Africa (Spear *et al.*, 2015). Rainfall deficit, rainfall variability and high temperature extremes have been found to be the main climate factors driving vulnerability (Batisani and Yarnal, 2010; Omari, 2010; Kgosikoma and Batisani, 2014; Masundire *et al.*, 2016). PRCPTOT relates to annual rainfall deficits, R99P, R95P, RX1DAY, RX5DAY, R20MM and R10MM, ALTCWD heavy rainfall extremes and ALTCD dry periods (irrespective of season). We note that rainfall extreme indices from GCMs may be biased compared to observations due to limitations in how models represent processes that drive precipitation extremes (Dai, 2006; Westra *et al.*, 2014); they tend to precipitate more frequently, and in smaller amounts, affecting both dry spell length, as well the absolute value of heavy rainfall. However, within their own climate, they do simulate precipitation extremes (O’Gorman, 2015), and so trends in these extremes can provide insights for the real world. For temperature-derived indices, WSDI is used to relate the potential impact of continuous high temperature instances (heat waves as defined by Moses, 2017). TN90P, TN10P, TX10P and TN10P are also included to help in defining the potential impact of individual hot and cold events (Klein Tank *et al.*, 2009).

Table 1: Temperature and precipitation climate extreme indices relevant to vulnerability assessment in Botswana. The indices are available from the KNMI Climate Explorer website [http://climexp.knmi.nl]

| Symbol | Name | Units |
|-----------------------|--|--------|
| Precipitation Indices | | |
| PRCPTOT | Annual Total Precipitation in Wet Days | mm/yr |
| ALTCD | Maximum Number of Consecutive Days Per Year with Less Than 1mm of Precipitation | days |
| ALTCWD | Maximum Number of Consecutive Days Per Year with At Least 1mm of Precipitation | days |
| RX1DAY | Annual Maximum 1-day Precipitation | mm/dy |
| RX5DAY | Annual Maximum 5-day Precipitation | mm/5dy |
| R99P | Annual Total Precipitation when Daily Precipitation Exceeds the 99 th Percentile of Wet Day Precipitation | mm/yr |
| R95P | Annual Total Precipitation when Daily Precipitation Exceeds the 95 th Percentile of Wet Day Precipitation | mm/yr |
| R20MM | Annual Count of Days with At Least 20mm of Precipitation | days |
| R10MM | Annual Count of Days with At Least 10mm of Precipitation | days |
| Temperature Indices | | |
| TX90P | Percentage of Days when Daily Maximum Temperature is Above the 90 th Percentile | % |
| TN90P | Percentage of Days when Daily Minimum Temperature is Above the 90 th Percentile | % |
| TX10P | Percentage of Days when Daily Minimum Temperature is Below the 10 th Percentile | % |
| TN10P | Percentage of Days when Daily Minimum Temperature is Below the 10 th Percentile | % |
| WSDI | Maximum Number of Consecutive Days Per Year when Daily Maximum Temperature is Above the 90 th Percentile | days |

2.3. Methodology

A 40 year period (1861-1900) base period for the preindustrial era was defined from which the changes in extreme indices are compared following Huang *et al.* (2017). The years at which each participating model reaches 1.0, 1.5 and 2.0 °C global warming above preindustrial levels is defined using a time sampling method initially used by Kaplan and New (2006) and applied by Zaroug *et al.* (in review). The 1.0 °C temperature warming above preindustrial is used to represent the current climate (warming to date, from preindustrial) given the GMST reached 1.1 °C in 2016 (WMO, 2017). A 31-year running mean is applied to the entire time-series for each model ensemble member. The climatology at a given GMST warming above pre-industrial is defined by the year the running mean reaches the GMST of interest and then stays consistently above the GMST. *Figure 2* shows the spread of the years the participating models reach 1.0, 1.5 and 2.0 °C warming above preindustrial levels.




Figure 2: Timing of when participating ensemble members reach GMST of 1.0, 1.5 and 2.0 °C warming above the 1861-1900 preindustrial baseline period following the RCP8.5 emissions scenario pathway

Monthly means of observed climate for 1979-2013 from the WATCH Forcing Data methodology applied to ERA-Interim data (WFDEI) (Weedon *et al.*, 2014) was used to cluster Botswana into 3 regions of homogeneous rainfall (*Figure 1b*). The WFDEI dataset was found to simulate precipitation well over southern Africa (Li *et al.*, 2013) and has been used in a number of studies in Sub Saharan Africa (Andersson *et al.*, 2016; Nkiaka *et al.*, 2017). Shapefiles of Botswana and the three regions were created using ArcMap and used to extract the gridded sub-sets of the indices over each region, and the whole country.

Area-weighted average climatological means of the indices at a given GMST warming above pre-industrial levels were calculated within the subsets and used to determine the change relative to pre-industrial levels. For all the 24 members, the change for each extreme index relative to preindustrial levels is calculated as;

$$\Delta I = I_n - I_0 \quad (1)$$

where $I_n, n \in \{1.0, 1.5 \text{ and } 2.0\}$ represents the area-averaged climatological mean calculated from the 31-year period surrounding the date of GMST, and I_0 is the area averaged climatological mean of the

index of interest for 1861-1900 preindustrial times. Box-and-whisker plots of the absolute changes for each climate extreme index are plotted spanning the 24-member ensemble for each for the 3 regions and over the country areal average.

The non-parametric Wilcoxon Paired Signed Rank test (WPSR) is applied to test if the distributions of ensembles of the indices at 1.0, 1.5 and 2.0 °C are statistically significantly different from preindustrial levels, and from each other. This test has been used in previous climate studies looking to determine the significance of changes in various climate indices/variables at different warming levels (Kharin *et al.*, 2013; Sillmann *et al.*, 2013b). To determine whether the models agree on the sign of change, a criterion that at least 75% of the members need to agree on the sign was adopted (Sillmann *et al.*, 2013b; Pinto *et al.*, 2016).

3. RESULTS

Results are presented in box-and-whisker plot format, representing the ensemble spread for the change in the indices relative to the preindustrial baseline period. Changes are presented first for precipitation extreme indices followed by temperature extreme indices. From the plots, the ensemble median, interquartile range (IQR) and the outliers are represented. Ensemble member values that exceed $1.5 \times IQR$ are considered to be outliers. The box-and-whisker plots are made for each of the 3 regions (*Figure 1b*) and the country average. Results obtained from testing model agreement on the sign of change are summarized in *Table S2* while *Table S3* summarizes the median and IQR changes in both of the precipitation and temperature indices relative to preindustrial levels. The inter-model ensemble spread, shown by the box-and-whisker plots together with the test for model agreement on change of sign depict the uncertainty in the projected changes of the indices. WPSR test results are presented using p-values.

3.1. Precipitation Extremes Changes

Changes in the precipitation indices at 1.0, 1.5 and 2.0 °C GMST warming above preindustrial levels indicate a progressive drying across Botswana, accompanied by an increase in heavy precipitation events, reduced wet spell events and increased dry spells (*Figure 3*). The country average ensemble median change for total annual precipitation, PRCPTOT at 1.0, 1.5 and 2.0 °C GMST warming above preindustrial levels indicate a reduction of -13 [range of -175; +75], -33 [-187; +46] and -63 [-192; +40] mm/yr (*Figure 3a*). Of the 3 regions, Region 1 representing the northern and wettest parts of the country has the largest median reduction, and range, across the 3 warming periods with reductions of 20.2, 55.1 and 75.9 mm/yr respectively (these are also the largest relative reductions, compared to preindustrial mean precipitation; see *Figure S1a*). This region, as opposed to the rest of the country tends to be the one that receives most of its rainfall when the ITCZ shifts south in austral summer. The stabilizing and equatorward shifting of the mean position of the ITCZ under climate change could therefore be the reason behind the reduction (Giannini *et al.*, 2008). Region 3 has the least reductions (10.0-43.0mm) in total annual precipitation, because of its already low annual precipitation totals (*Figure 1a*); relative reductions are much larger (*Figure S1a*). When testing for model agreement on change of sign, the change in PRCPTOT at 2.0 °C is the only one that shows consensus among models with >75% of the ensemble members projecting a

1 reduction across all regions. Taking the country average, 58.3% of the models project a decrease in
2 PRCPTOT at 1.0 °C, 62.5% at 1.5 °C and 83.3% at 2.0 °C GMST above preindustrial levels (*Table S2*). WPSR
3 test results show that the change in PRCPTOT between the three levels of warming is statistically
4 significant across all regions, except for 1.0 versus 1.5 for a couple of the regions. An additional 0.5 °C
5 increment in GMST from 1.5 and 2.0 °C therefore has significant impacts on the annual precipitation across
6 the country. We note that PRCPTOT is not strictly a climate extreme, but it is included in climate extreme
7 studies as total annual precipitation is a measure of interannual drought, and has implications on various
8 economic sectors especially in water stressed countries like Botswana.




Figure 3: Box and whisker plots showing the changes in extreme precipitation indices across an ensemble of 24 participating model members. The changes are at 1.0, 1.5 and 2.0 °C above preindustrial levels (1861-1900). P-values from the WRS test are shown: P_0 , P_1 and P_2 compares the ensemble spread of 1.0 to 1.5 °C, 1.5 to 2.0 °C and 1.0 to 2.0 °C warmer climate regimes respectively

The number of consecutive dry days (ALTCDD) (*Figure 3b*) show statistically significant increases across all regions for all warming levels. The ALTCDD median increases of 7.2, 13.5 and 18.7 days for the 1.0, 1.5 and 2.0 °C respectively are projected for Region 1, these being the largest of the changes. On average, Botswana is projected to experience median increases of 5.7, 10.4 and 17.8 days in ALTCDD under the respective warmings. There is general consensus on the sign of change of ALTCDD across ensemble members with more than 80% of members depicting an increase across the three warming periods over the entire country. The increases in ALTCDD imply longer dry seasons with late onsets and early cessation of rainfall, as noted by Pinto *et al.* (2016) and Sillmann *et al.* (2013b). Median changes in the number of consecutive wet days (ALTCWD) are generally small in magnitude (*Figure 3c*). Relative to preindustrial levels, median changes at 1.0 °C warming are for a decrease of 0.21 days in Region 1 while Region 2 and 3 show median increases of even smaller magnitude. On average, an additional 0.5 °C warming to 1.5 and 2.0 °C reduces ALTCWD by about half a day. The reductions in ALTCWD may be small in magnitude but are very significant given the short-lived and convective nature of rain bearing weather systems in Botswana. The shorter rainfall seasons described by increases in ALTCDD coupled with potentially shorter wet-spells could have serious implications on various economic sectors, with the agricultural sector likely to be particularly vulnerable.

For the heavy precipitation indices, the projected ensemble median changes in total accumulated precipitation from heavy (R95P) and very heavy (R99P) precipitation days are a small, but generally non-significant increase as the climate system warms further (*Figure 3d and 3e*). For R99P, the changes are mostly significant between the three warming levels in Regions 1 and 2, but not in Region 3. The ensemble spread generally disagrees on the sign of change for R95P, but there is more agreement for R99P, especially at the 1.5 and 2.0 °C warming levels.

Median changes in one-day and five-day maximum precipitation (RX1DAY and RX5DAY) show a general increase across all regions as the models progress to warmer climates. These changes are generally statistically significant over the wetter and semi-arid Regions 1 and 2, but not in the arid Region 3. (*Figure 3f and 3g*). These results slightly contradict the findings by Sillmann *et al.* (2013b) who concluded that there is a general decrease (though statistically insignificant) in RX5DAY in Southern Africa when looking at the changes for the period 2081-2100 relative to 1981-2000. Pinto *et al.* (2016)'s findings using downscaled projections over Southern Africa are consistent with our results, though they looked the changes at 2069-2098 relative to the 1976-2005 period. Changes in the frequency of very heavy (R20MM) rainfall events do not show any significant changes, while the frequency of moderately heavy rainfall events (R10MM) show statistically significant decreases across all regions (*Figure 3h and 3i*).

To investigate whether biases inherent in climate models especially in simulating accumulated precipitation may influence the results, box-and-whisker plots of percentage changes in the total annual precipitation (PRCPTOT), one day maximum precipitation (RX1day) and the five-day maximum accumulated precipitation (RX5DAY) relative to industrial levels were also analysed (*Figure S1*). Similar results to those obtained using absolute changes were found. An exception was that P_0 for PRCPTOT in Region 2 decreases from 0.117 when using absolute changes to 0.097 when using percentage changes (*Figure S1a*). The difference suggests that the change in PRCPTOT between the current climate and 1.5 °C

are statistically significantly different in Region 2. This could be because of the aridity in Region 2 meaning small percentage changes in total annual precipitation make a significant difference.

3.2. Temperature Changes

The changes in temperature indices agree strongly on the direction of change across all indices (*Figure 4* and *Table S2*). P-values for temperature derived indices obtained from the WPSR test are all very small ($\ll 0.10$) across all regions and warming levels above preindustrial levels. The p-values here imply that the changes in these indices are statistically significant across all warming levels. The inter-model spread is small for most temperature extremes therefore results are generally associated with less uncertainty.

Of the percentile based indices, the hot day and hot night extremes, TX90P (*Figure 4a*) and TN90P (*Figure 4b*) show the greatest changes, though the change is more pronounced for TN90P especially over Region 1. TX90P is projected to increase by 30% above preindustrial levels on average in Botswana when the climate system reaches 2.0 °C, an increment of 10% from 1.5 °C levels. For TN90P the average increase at 2.0 °C is even higher, at 35%. Decreases in cold day and night extremes, TX10P (*Figure 4c*) and TN10P (*Figure 4d*), occur over the entire country with minimum temperature based extremes showing the larger reductions. Hot nights and mild winters are therefore expected to become a common occurrence with a warming climate leading to a decrease in frost occurrences. Warm spells (heat waves; WSDI) increase are projected across all regions, by 80 days compared to preindustrial levels at 2.0 °C for Region 1, and by 65 and 62 days for Region 2 and 3 (*Figure 4e*). Even though all models show an increase in warm spells with increased warming, the ensemble spread also increases significantly at 2.0 °C (ranging between 26 and 96 days) compared to present (5 to 39 days) and 1.5 °C (13 and 65 days) indicating an increasing uncertainty as models warm further. Relating to findings by Moses (2017), the increases in TX90P and WSDI suggest a significant increase in heat wave events across the country. We note here the need to look at these indices seasonally as an opportunity for further investigation (Sillmann *et al.*, 2013b), and that the WSDI does not consider the intensity of heat waves (Dosio, 2016).




Figure 4: Box and whisker plots showing the changes in extreme temperature indices across an ensemble of 24 participating model members. The changes are at 1.0, 1.5 and 2.0 °C above preindustrial levels (1861-1900). P-values from the WRS test are shown: P_0 , P_1 and P_2 compares the ensemble spread of 1.0 to 1.5 °C, 1.5 to 2.0 °C and 1.0 to 2.0 °C warmer climate regimes respectively

3.3. Implications for Vulnerability to Climate Change in Botswana

The projected changes in both temperature and precipitation extremes under warming climates are likely to have significant negative impacts on many social and economic activities in Botswana, most especially in the agricultural sector, and those dependent on water. The majority of Botswana's population is highly reliant on rain-fed agriculture for livelihoods, so these changes in extremes are likely to produce severe impacts, especially on the most vulnerable, women (Omari, 2010). Many crops suffer sharp drops in yield after periods of cumulative heat stress (e.g., Schlenker *et al.*, 2010). The large increases in temperature extremes projected for Botswana as one moves from 1.5 to 2.0 °C, suggest potentially large, and growing impacts on crops that are currently farmed, such as maize (Barnabás *et al.*, 2008), significantly reducing yields. This is in agreement with previous studies showing that Botswana will be among the most impacted countries with regards to agriculture in Africa (Chipanshi *et al.*, 2004; Schlenker and Lobell, 2010). These impacts will be exacerbated, particularly for rain-fed agriculture, by the projected decreases in mean annual rainfall and longer dry spells causing plant water stress, more frequently crossing the tipping point between good and average harvests, and complete agricultural failure (Batisani and Yarnal, 2010). More

intense rainfall will potentially cause crop damage and lower soil moisture. These multiple impacts warn of a growing adaptation challenge as the climate warms from today, through 1.5 to 2.0 degrees, requiring more heat and drought-tolerant varieties to be developed or adopted. The options for expanding irrigated agriculture are small, apart from perhaps the North West of the country in the Okavango basin, but this would involve trade-offs with biodiversity conservation and ecotourism in this unique wetland system. Livestock production is also likely to be negatively impacted as increased dry spells will reduce pasture productivity (Setshwaelo, 2001; Mberego, 2017), compounded by inadequate adaptation strategies (Kgosikoma and Batisani, 2014).

Water resources are also likely to be heavily impacted by the reduced total precipitation and increased intensity and longer dry spells, and greater evaporation under more extreme temperatures. Water stress is already a challenge with various economic sectors competing for the scarce resource; drought in 2014-2016 led to water-supply failure in the capital city, Gaborone (Siderius *et al.*, 2017). For Region 3, in the south-western parts of the country where rainfall is already low (annual accumulations of less than 300mm), further reductions in rainfall imply increased pressure on the already stressed water sources (Batisani, 2011). The increasing RX5DAY implies a potential increase in very heavy rainfall events that could cause flooding and lead to economic losses (Tsheko, 2003). Though this is the case, some of these heavy rainfall events could come as a relief, replenishing stressed water resources. An example of such a case is the heavy downpours that came with post cyclone Dineo in February 2017, filling up the Gaborone dam that had run dry the previous year.

Climate sensitive diseases such as malaria will also be affected by the changes in climate extremes. As malaria epidemics thrive in wet and warmer climates, a general drying of the climate and shorter rainy seasons could lead to a reduction in the extent of the disease (Tanser *et al.*, 2003). Tanser *et al.* (2003) also note that though for drier climates climate change might lead to a reduction in malaria incidences, epidemics can rise during times of heavy precipitation in these generally dry climates (Huang *et al.*, 2017). Based on the reasoning above, epidemics of malaria cases could also become a challenge as the climate systems warms to 2.0 °C. Other implications for health include increased chances of heat related mortalities as heat waves become a common occurrence, malnutrition due to reduced food supply as the agricultural sector is negatively impacted, and direct mortalities and injuries from floods (McMichael *et al.*, 2006).

4. Conclusions

This study found that Botswana is projected to experience significant increases in all temperature and many rainfall extremes as GMST increases from 1.0 °C through 1.5 to 2.0 °C above preindustrial levels. The changes are particularly strong for temperature extremes; for each increment of global warming, temperature extremes are statistically different, indicating markedly different regional climates over Botswana at different levels of global warming. Similarly, there are projected statistically significant decreases in mean rainfall and increases in dry-spell length at each global temperature level. In contrast, although intense rainfall indices do show ensemble median increases, the spread in model results means that changes at each increment are not statistically distinguishable.

These changes in extremes present a growing adaptation challenge between 1.5 and 2.0 °C for key economic sectors in the country. Rain-fed agriculture is already marginal across much of the country, and the combined changes in heat extremes and decrease in moisture may well make current agricultural practices unviable at 1.5 and 2.0 °C warming. Botswana is already water-stressed; the projected decreases in mean annual rainfall, as well increased dry spell length, will escalate stress, leading to more frequent water shortages in today's urban and agricultural supply systems. Further work is needed to better quantify the impacts, and resultant costs of adaptation at these different levels of global mean warming. However, our results suggest that, for a climate-stressed country such as Botswana, even small increments in global mean temperature have serious societal consequences that will demand progressively more radical adaptation responses.

A key limitation in our study relates to the use of GCM data, which are known to not completely represent extremes, especially rainfall, due to their relatively coarse spatial resolution. Further work is needed to apply downscaling methods to the GCM data, to add more information at finer space scales, subject to suitable evaluation of the downscaled data, as suggested by Pinto *et al.* (2016). Another limitation to this study is that all extremes analysed are calculated on annual timescales; given that some of the indices such as ALTCD and WSDI are also significant when looked at within seasons, further work is needed to look at the changes at seasonal timescales (Sillmann *et al.*, 2013b; Sillmann *et al.*, 2017).

5. References

- Akinyemi, F.O. 2017. Climate change and variability in semiarid palapye, eastern Botswana: An assessment from smallholder farmers' perspective. *Weather, Climate, and Society*. 9(3). DOI: 10.1175/WCAS-D-16-0040.1.
- Alexander, L. V., Zhang, X., Peterson, T.C., Caesar, J., Gleason, B., Klein Tank, A.M.G., Haylock, M., Collins, D., et al. 2006. Global observed changes in daily climate extremes of temperature and precipitation. *Journal of Geophysical Research Atmospheres*. 111(5):1–22. DOI: 10.1029/2005JD006290.
- Andersson, J.C.M., Ali, A., Arheimer, B., Gustafsson, D. & Minoungou, B. 2016. Providing peak river flow statistics and forecasting in the Niger River basin. *Physics and Chemistry of the Earth*. DOI: 10.1016/j.pce.2017.02.010.
- Andringa, J. 1984. The climate of Botswana in histograms. *Botswana Notes and Records*. 16(1):117–125. Available: <http://journals.co.za/docserver/fulltext/botnotes/16/1/748.pdf?expires=1512601715&id=id&acname=guest&checksum=C606DC53AD270E341F433DF0D3E11203> [2017, December 07].
- ASC. 2016. *UK Climate Change Risk Assessment 2017 Synthesis Report: priorities for the next five years*. London: Adaptation Sub-Committee of the Committee on Climate Change. Available: <https://www.theccc.org.uk/wp-content/uploads/2016/07/UK-CCRA-2017-Synthesis-Report-Committee-on-Climate-Change.pdf>.
- Barnabás, B., Jäger, K. & Fehér, A. 2007. The effect of drought and heat stress on reproductive processes in cereals. *Plant, Cell & Environment*. 31(0):071030190532001–??? DOI: 10.1111/j.1365-

- 1 3040.2007.01727.x.
- 2 Batisani, N. 2011. Spatio-temporal ephemeral streamflow as influenced by climate variability in
- 3 Botswana. *Journal of Geographical Sciences*. 21(3):417–428. DOI: 10.1007/s11442-011-0854-5.
- 4 Batisani, N. & Yarnal, B. 2010. Rainfall variability and trends in semi-arid Botswana: Implications for
- 5 climate change adaptation policy. *Applied Geography*. 30(4):483–489. DOI:
- 6 10.1016/j.apgeog.2009.10.007.
- 7 Bhalotra, Y.P.R. 1987. *Climate of Botswana, part II: Elements of Climate*. Gaborone, Botswana.
- 8 Blamey, R.C. & Reason, C.J.C. 2012. Mesoscale convective complexes over Southern Africa. *Journal of*
- 9 *Climate*. 25(2):753–766. DOI: 10.1175/JCLI-D-10-05013.1.
- 10 Boko, M., Niang, I., Nyong, A., Vogel, C., Githeko, A., Medany, M., Osman-Elasha, B., Tabo, R., et al. 2008.
- 11 Africa. In *Climate change 2007: Impacts, adaptation and vulnerability. Contribution of Working Group II*
- 12 *to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. M.L. Parry, O.
- 13 Canziani, J. Palutikof, P.J. van de Linden, & C.E. Hanson, Eds. 433–467. DOI: 10.2134/jeq2008.0015br.
- 14 Ceccherini, G., Russo, S., Amezttoy, I., Francesco Marchese, A. & Carmona-Moreno, C. 2017. Heat waves
- 15 in Africa 1981-2015, observations and reanalysis. *Natural Hazards and Earth System Sciences*. 17(1):115–
- 16 125. DOI: 10.5194/nhess-17-115-2017.
- 17 Chipanshi, A.C., Chanda, R. & Totolo, O. 2003. Vulnerability Assessment of the Maize and Sorghum Crops
- 18 to Climate Change in Botswana. *Climatic Change*. 61(3):339–360. DOI:
- 19 10.1023/B:CLIM.0000004551.55871.eb.
- 20 Cook, C., Reason, C.J.C. & Hewitson, B.C. 2004. Wet and dry spells within particularly wet and dry
- 21 summers in the South African summer rainfall region. *Climate Research*. 26(1):17–31. DOI:
- 22 10.3354/cr026017.
- 23 Crimp, S.J. & Mason, S.J. 1999. The Extreme Precipitation Event of 11 to 16 February 1996 over South
- 24 Africa. *Meteorology and Atmospheric Physics*. 70(1–2):29–42. DOI: 10.1007/s007030050023.
- 25 Dai, A. 2006. Precipitation characteristics in eighteen coupled climate models. *Journal of Climate*.
- 26 19(18):4605–4630. DOI: 10.1175/JCLI3884.1.
- 27 Dosio, A. 2016. Projection of temperature and heat waves for Africa with an ensemble of CORDEX
- 28 Regional Climate Models. *Climate Dynamics*. 49(1):1–27. DOI: 10.1007/s00382-016-3355-5.
- 29 Driver, P. & Reason, C.J.C. 2017. Variability in the Botswana High and its relationships with rainfall and
- 30 temperature characteristics over southern Africa. *International Journal of Climatology*. (March, 1). DOI:
- 31 10.1002/joc.5022.
- 32 Favre, A., Hewitson, B., Lennard, C., Cerezo-Mota, R. & Tadross, M. 2013. Cut-off Lows in the South
- 33 Africa region and their contribution to precipitation. *Climate Dynamics*. 41(9–10):2331–2351. DOI:
- 34 10.1007/s00382-012-1579-6.
- 35 Fischer, E.M., Beyerle, U. & Knutti, R. 2013. Robust spatially aggregated projections of climate extremes.
- 36 *Nature Climate Change*. 3(12):1033–1038. DOI: 10.1038/nclimate2051.
- 37 Frich, P., Alexander, L. V., Della-Marta, P., Gleason, B., Haylock, M., Tank Klein, A.M.G. & Peterson, T.

- 1 2002. Observed coherent changes in climatic extremes during the second half of the twentieth century.
2 *Climate Research*. 19(3):193–212. DOI: 10.3354/cr019193.
- 3 Giannini, A., Held, I.M. & Sobel, A.H. 2008. A global perspective on African climate. *Climatic Change*.
4 90(4):359–383. DOI: 10.1007/s10584-008-9396-y.
- 5 Hambira, W.L. & Saarinen, J. 2015. Policy-makers' perceptions of the tourism –climate change nexus:
6 Policy needs and constraints in Botswana. *Development Southern Africa*. 32:350–362. DOI:
7 10.1080/0376835X.2015.1010716.
- 8 Henley, B.J. & King, A.D. 2017. Trajectories towards the 1.5°C Paris target: modulation by the
9 Interdecadal Pacific Oscillation. *Geophysical Research Letters*. 44(9):4256–4262. DOI:
10 10.1002/2017GL073480.
- 11 Huang, J., Yu, H., Dai, A., Wei, Y. & Kang, L. 2017. Drylands face potential threat under 2°C global
12 warming target. *Nature Climate Change*. 7(6):417–422. DOI: 10.1038/nclimate3275.
- 13 IPCC. 2012. *Managing the risks of extreme events and disasters to advance climate change adaptation*.
14 C.B. Field, V. Barros, T.F. Stocker, D. Qin, D.J. Dokken, K.L. Ebi, M.D. Mastrandrea, K.J. Mach, G.-K.
15 Plattner, S.K. Allen, M. Tignor, & P.M. Midgley, Eds. Cambridge, UK, and New York, NY, USA: Cambridge
16 University Press. DOI: 10.1596/978-0-8213-8845-7.
- 17 James, R., Washington, R., Schleussner, C.-F., Rogelj, J. & Conway, D. 2017. Characterizing half-a-degree
18 difference: a review of methods for identifying regional climate responses to global warming targets.
19 *Wiley Interdisciplinary Reviews: Climate Change*. 8(2):e457. DOI: 10.1002/wcc.457.
- 20 Kaplan, J.O. & New, M. 2006. Arctic climate change with a 2°C global warming: Timing, climate patterns
21 and vegetation change. *Climatic Change*. 79(3–4):213–241. DOI: 10.1007/s10584-006-9113-7.
- 22 Kgosi, O.E. & Batisani, N. 2014. Livestock population dynamics and pastoral communities'
23 adaptation to rainfall variability in communal lands of Kgalagadi South, Botswana. *Pastoralism*. 4:1–9.
24 DOI: 10.1186/s13570-014-0019-0.
- 25 Kharin, V. V., Zwiers, F.W., Zhang, X. & Hegerl, G.C. 2007. Changes in temperature and precipitation
26 extremes in the IPCC ensemble of global coupled model simulations. *Journal of Climate*. 20(8):1419–
27 1444. DOI: 10.1175/JCLI4066.1.
- 28 Kharin, V. V., Zwiers, F.W., Zhang, X. & Wehner, M. 2013. Changes in temperature and precipitation
29 extremes in the CMIP5 ensemble. *Climatic Change*. 119(2):345–357. DOI: 10.1007/s10584-013-0705-8.
- 30 King, A.D., Karoly, D.J., Henley, B.J., Karoly, D.J. & Henley, B.J. 2017. Australian climate extremes at 1.5°C
31 and 2°C of global warming. *Nature Climate Change*. 7(May):1–10. DOI: 10.1038/nclimate3296.
- 32 King, D., Schrag, D., Dadi, Z., Ye, Q. & Ghosh, A. 2015. *Climate Change: A Risk Assessment*. London,
33 Beijing, Delhi, Cambridge. DOI: 10.1016/j.ecocom.2014.11.002.
- 34 Knutti, R., Furrer, R., Tebaldi, C., Cermak, J. & Meehl, G.A. 2010. Challenges in combining projections
35 from multiple climate models. *Journal of Climate*. 23(10):2739–2758. DOI: 10.1175/2009JCLI3361.1.
- 36 Lewis, S.C. & King, A.D. 2017. Evolution of mean, variance and extremes in 21st century temperatures.
37 *Weather and Climate Extremes*. 15:1–10. DOI: 10.1016/j.wace.2016.11.002.

- 1 Li, L., Ngongondo, C.S., Xu, C. & Gong, L. 2013. Comparison of the global TRMM and WFD precipitation
2 datasets in driving a large-scale hydrological model in southern Africa. *Hydrology Research*. 44(5):770.
3 DOI: 10.2166/nh.2012.175.
- 4 Manhique, A.J., Reason, C.J.C., Silinto, B., Zucula, J., Raiva, I., Congolo, F. & Mavume, A.F. 2015. Extreme
5 rainfall and floods in southern Africa in January 2013 and associated circulation patterns. *Natural*
6 *Hazards*. 77(2):679–691. DOI: 10.1007/s11069-015-1616-y.
- 7 Manton, M.J. 2010. Trends in climate extremes affecting human settlements. *Current Opinion in*
8 *Environmental Sustainability*. 2(3):151–155. DOI: 10.1016/j.cosust.2010.05.001.
- 9 Mason, S.J. & Jury, M.R. 1997. Climatic variability and change over southern Africa: a reflection on
10 underlying processes. *Progress in Physical Geography*. 21(1):23–50. DOI:
11 10.1177/030913339702100103.
- 12 Masundire, H., Morchain, D., Raditloaneng, N., Hegga, S., Ziervogel, G., Molefe, C., Angula, M. & Omari,
13 K. 2016. *Vulnerability and Risk Assessment in Botswana's Bobirwa Sub-District: Fostering People-Centred*
14 *Adaptation to Climate Change*. Ottawa.
- 15 Mberego, S. 2017. Temporal patterns of precipitation and vegetation variability over Botswana during
16 extreme dry and wet rainfall seasons. *International Journal of Climatology*. 37:2947–2960. DOI:
17 10.1175/JAMC-D-14-0006.1.
- 18 McElroy, M. & Baker, D.J. 2012. *Climate Extremes: Recent Trends with Implications for National Security*.
19 Cambridge. Available:
20 http://environment.harvard.edu/sites/default/files/climate_extremes_report_2012-12-04.pdf.
- 21 McMichael, A.J., Woodruff, R.E. & Hales, S. 2006. Climate change and human health: Present and future
22 risks. *Lancet*. 367(9513):859–869. DOI: 10.1016/S0140-6736(06)68079-3.
- 23 Molekwa, S. 2013. Cut-off lows over South Africa and their contribution to the total rainfall of the
24 Eastern Cape Province. University of Pretoria. Available:
25 <https://repository.up.ac.za/handle/2263/36767>.
- 26 Monier, E. & Gao, X. 2015. Climate change impacts on extreme events in the United States: an
27 uncertainty analysis. *Climatic Change*. 131(1):67–81. DOI: 10.1007/s10584-013-1048-1.
- 28 Moses, O. 2017. Heat wave characteristics in the context of climate change over the past 50 years in
29 Botswana. *Botswana Notes and Records*. 49(0):13–25. Available:
30 <http://journals.ub.bw/index.php/bnr/article/view/973/589>.
- 31 Moyo, E.N. & Nangombe, S.S. 2015. Southern Africa's 2012-13 violent storms: Role of climate change.
32 *Procedia IUTAM*. 17(2013):69–78. DOI: 10.1016/j.piutam.2015.06.011.
- 33 ND-GAIN. 2016. *Rankings [ND-GAIN Country Index]*. Available: [https://gain.nd.edu/our-work/country-](https://gain.nd.edu/our-work/country-index/rankings/)
34 [index/rankings/](https://gain.nd.edu/our-work/country-index/rankings/) [2018, February 08].
- 35 New, M., Hewitson, B., Stephenson, D.B., Tsiga, A., Kruger, A., Manhique, A., Gomez, B., Coelho, C.A.S.,
36 et al. 2006. Evidence of trends in daily climate extremes over southern and west Africa. *Journal of*
37 *Geophysical Research Atmospheres*. 111(14):1–11. DOI: 10.1029/2005JD006289.

- 1 Niang, I., Ruppel, O.C., Abdrabo, M.A., Essel, A., Lennard, C., Padgham, J. & Urquhart, P. 2014. Africa.
2 *Climate Change 2014: Impacts, Adaptation and Vulnerability - Contributions of the Working Group II to*
3 *the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. 1199–1265. DOI:
4 10.1017/CBO9781107415386.002.
- 5 Nicholson, S.E., Leposo, D. & Grist, J. 2001. The Relationship between El Niño and Drought over
6 Botswana. *Journal of Climate*. 14(3):323–335. DOI: 10.1175/1520-
7 0442(2001)014<0323:TRBENO>2.0.CO;2.
- 8 Nkiaka, E., Nawaz, N. & Lovett, J. 2017. Evaluating Global Reanalysis Datasets as Input for Hydrological
9 Modelling in the Sudano-Sahel Region. *Hydrology*. 4(1):13. DOI: 10.3390/hydrology4010013.
- 10 O’Gorman, P.A. 2015. Precipitation Extremes Under Climate Change. *Current Climate Change Reports*.
11 1(2):49–59. DOI: 10.1007/s40641-015-0009-3.
- 12 Omari, K. 2010. Gender and Climate change: Botswana Case Study. 35(3):42–49. DOI: 10.1111/j.1759-
13 5436.2004.tb00133.x.
- 14 Pendergrass, A.G., Lehner, F., Sanderson, B.M. & Xu, Y. 2015. Does extreme precipitation intensity
15 depend on the emissions scenario? *Geophysical Research Letters*. 42(20):8767–8774. DOI:
16 10.1002/2015GL065854.
- 17 Pinto, I., Lennard, C., Tadross, M., Hewitson, B., Dosio, A., Nikulin, G., Panitz, H.J. & Shongwe, M.E. 2016.
18 Evaluation and projections of extreme precipitation over southern Africa from two CORDEX models.
19 *Climatic Change*. 135(3–4):655–668. DOI: 10.1007/s10584-015-1573-1.
- 20 Razavi, T., Switzman, H., Arain, A. & Coulibaly, P. 2016. Regional climate change trends and uncertainty
21 analysis using extreme indices: A case study of Hamilton, Canada. *Climate Risk Management*. 13:43–63.
22 DOI: 10.1016/j.crm.2016.06.002.
- 23 Reason, C.J.C. 2007. Tropical cyclone Dera, the unusual 2000/01 tropical cyclone season in the South
24 West Indian Ocean and associated rainfall anomalies over Southern Africa. *Meteorology and*
25 *Atmospheric Physics*. 97(1–4):181–188. DOI: 10.1007/s00703-006-0251-2.
- 26 Reason, C.J.C. & Keibel, A. 2004. Tropical Cyclone Eline and Its Unusual Penetration and Impacts over the
27 Southern African Mainland. *Weather and Forecasting*. 19(5):789–805. DOI: 10.1175/1520-
28 0434(2004)019<0789:TCEAIU>2.0.CO;2.
- 29 Richard, Y., Fauchereau, N., Pocard, I., Rouault, M. & Trzaska, S. 2001. 20th century droughts in
30 southern Africa: Spatial and temporal variability, teleconnections with oceanic and atmospheric
31 conditions. *International Journal of Climatology*. 21:873–885. DOI: 10.1002/joc.656.
- 32 Rouault, M. & Richard, Y. 2005. Intensity and spatial extent of droughts in southern Africa. *Geophysical*
33 *Research Letters*. 32(15):2–5. DOI: 10.1029/2005GL022436.
- 34 Sanderson, B.M., Xu, Y., Tebaldi, C., Wehner, M., O’Neill, B., Jahn, A., Pendergrass, A.G., Lehner, F., et al.
35 2017. Community Climate Simulations to assess avoided impacts in 1.5°C and 2.0°C futures. *Earth*
36 *System Dynamics Discussions*. 2017(April):1–33. DOI: 10.5194/esd-2017-42.
- 37 Schlenker, W. & Lobell, D.B. 2010. Robust negative impacts of climate change on African agriculture.
38 *Environmental Research Letters*. 5(1):14010. DOI: 10.1088/1748-9326/5/1/014010.

- 1 Setshwaelo, L. 2001. *A Study of Grasslands and Livestock Vulnerability and Adaptation to Climate Change in Botswana*. Gaborone: Government of Botswana.
- 2
- 3 Shi, C., Jiang, Z.-H., Chen, W.-L. & Li, L. 2017. Changes in temperature extremes over China under 1.5 °C and 2 °C global warming targets. *Advances in Climate Change Research*. (December):1–8. DOI: 10.1016/j.accre.2017.11.003.
- 4
- 5
- 6 Shongwe, M.E., Van Oldenborgh, G.J., Van Den Hurk, B.J.J.M., De Boer, B., Coelho, C.A.S. & Van Aalst, M.K. 2009. Projected changes in mean and extreme precipitation in Africa under global warming. Part I: Southern Africa. *Journal of Climate*. 22(13):3819–3837. DOI: 10.1175/2009JCL12317.1.
- 7
- 8
- 9
- 10 Siderius, C., Gannon, K.E., Ndiyoi, M., Opere, A., Batisani, N., Olago, D., Pardoe, J. & Conway, D. 2017. Hydrological Response and Complex Impact Pathways of the 2015/2016 El Niño in Eastern and Southern Africa. *Earth's Future*. 5:1–21. DOI: 10.1002/2017EF000680.
- 11
- 12
- 13 Sillmann, J., Kharin, V. V., Zhang, X., Zwiers, F.W. & Bronaugh, D. 2013. Climate extremes indices in the CMIP5 multimodel ensemble: Part 1. Model evaluation in the present climate. *Journal of Geophysical Research Atmospheres*. 118(4):1716–1733. DOI: 10.1002/jgrd.50203.
- 14
- 15
- 16 Sillmann, J., Kharin, V. V., Zwiers, F.W., Zhang, X. & Bronaugh, D. 2013. Climate extremes indices in the CMIP5 multimodel ensemble: Part 2. Future climate projections. *Journal of Geophysical Research Atmospheres*. 118(6):2473–2493. DOI: 10.1002/jgrd.50188.
- 17
- 18
- 19 Sillmann, J., Thorarinsdottir, T., Keenlyside, N., Schaller, N., Alexander, L. V., Hegerl, G., Seneviratne, S.I., Vautard, R., et al. 2017. Understanding, modeling and predicting weather and climate extremes: Challenges and opportunities. *Weather and Climate Extremes*. 18:65–74. DOI: 10.1016/j.wace.2017.10.003.
- 20
- 21
- 22 Singleton, A.T. & Reason, C.J.C. 2007. Variability in the characteristics of cut-off low pressure systems over subtropical southern Africa. *International Journal of Climatology*. 27(3):295–310. DOI: 10.1002/joc.1399.
- 23
- 24
- 25 Spear, D., Baudoin, M., Hegga, S., Zaroug, M., Okeyo, A. & Haimbili, E. 2015. *Vulnerability and Adaptation To Climate Change in the Semi-Arid Regions of Southern Africa*. Available: <http://www.assar.uct.ac.za>.
- 26
- 27
- 28 Tanser, F.C., Sharp, B. & Le Sueur, D. 2003. Potential effect of climate change on malaria transmission in Africa. *Lancet*. 362(9398):1792–1798. DOI: 10.1016/S0140-6736(03)14898-2.
- 29
- 30
- 31 Taylor, K.E., Stouffer, R.J. & Meehl, G.A. 2012. An overview of CMIP5 and the experiment design. *Bulletin of the American Meteorological Society*. 93(4):485–498. DOI: 10.1175/BAMS-D-11-00094.1.
- 32
- 33
- 34 Tebaldi, C. & Knutti, R. 2007. The use of the multi-model ensemble in probabilistic climate projections. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*. 365(1857):2053–2075. DOI: 10.1098/rsta.2007.2076.
- 35
- 36
- 37 Trenberth, K.E. 2012. Framing the way to relate climate extremes to climate change. *Climate Change*. 115:283–290. DOI: 10.1007/s10584-012-0441-5.
- 38
- 39
- 40
- 41 Trouet, V. & Van Oldenborgh, G.J. 2013. KNMI Climate Explorer: A Web-Based Research Tool for High-Resolution Paleoclimatology. *Tree-Ring Research*. 69(1):3–13. DOI: 10.3959/1536-1098-69.1.3.
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60

- 1 Tsheko, R. 2003. Rainfall reliability, drought and flood vulnerability in Botswana. *Water SA*. 29(4):389–392. DOI: 10.4314/wsa.v29i4.5043.
- 2
- 3 UNFCCC. 2015. *Decision 1/CP.21. The Paris Agreement*. DOI: FCCC/CP/2015/L.9/Rev.1.
- 4
- 5 Usman, M.T. & Reason, C.J.C. 2004. Dry spell frequencies and their variability over southern Africa. *Climate Research*. 26(3):199–211. DOI: 10.3354/cr026199.
- 6
- 7 Vossen, P. 1990. Rainfall and Agricultural Production in Botswana. *Afrika Focus*. 6(2):141–155.
- 8
- 9 Wang, Z., Lin, L., Zhang, X., Zhang, H., Liu, L. & Xu, Y. 2017. Scenario dependence of future changes in climate extremes under 1.5°C and 2°C global warming. *Scientific Reports*. 7:46432. DOI: 10.1038/srep46432.
- 10
- 11 Weedon, G.P., Balsamo, G., Bellouin, N., Gomes, S., Best, M.J. & Viterbo, P. 2014. The WFDEI meteorological forcing data set: WATCH Forcing Data methodology applied to ERA-Interim reanalysis data. *Water Resources Research*. 50(9):7505–7514. DOI: 10.1002/2014WR015638.
- 12
- 13 Westra, S., Fowler, H.J., Evans, J.P., Alexander, L. V., Berg, P., Johnson, F., Kendon, E.J., Lenderink, G., et al. 2014. Future changes to the intensity and frequency of short-duration extreme rainfall. *Reviews of Geophysics*. 52(3):522–555. DOI: 10.1002/2014RG000464.
- 14
- 15 Williams, C.J.R., Kniveton, D.R. & Layberry, R. 2007. Climatic and oceanic associations with daily rainfall extremes over southern Africa. *International Journal of Climatology*. 27:93–108. DOI: 10.1002/joc1376.
- 16
- 17 WMO. (in press). WMO statement on the status of the global climate in 2016. *World Meteorological Organization*. (1189):WMO-No. 1189. Available: http://www.wmo.int/pages/mediacentre/press_releases/documents/WMO_1108_EN_web_000.pdf.
- 18
- 19 WMO, Klein Tank, A.M.G., Zwiers, F.W. & Zhang, X. 2009. *Guidelines on Analysis of extremes in a changing climate in support of informed decisions for adaptation*. Geneva, Switzerland. Available: File Attachment.
- 20
- 21 World Bank. 2013. *Turn Down the Heat: Climate Extremes, Regional Impacts, and the Case for Resilience*. Washington D.C.
- 22
- 23 You, Q., Kang, S., Aguilar, E., Pepin, N., Flugel, W.A., Yan, Y., Xu, Y., Zhang, Y., et al. 2011. Changes in daily climate extremes in China and their connection to the large scale atmospheric circulation during 1961–2003. *Climate Dynamics*. 36(11–12):2399–2417. DOI: 10.1007/s00382-009-0735-0.
- 24
- 25 Zaroug, M., New, M. & Lennard, C. n.d. National Climate Change at 1.5 and 2.0°C Global Warming over Africa. *Environmental Research Letters*.
- 26
- 27 Zhang, X., Alexander, L., Hegerl, G.C., Jones, P., Tank, A.K., Peterson, T.C., Trewin, B. & Zwiers, F.W. 2011. Indices for monitoring changes in extremes based on daily temperature and precipitation data. *Wiley Interdisciplinary Reviews: Climate Change*. 2(6):851–870. DOI: 10.1002/wcc.147.
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35