

Association of General and Central Adiposity with Blood Pressure among Chinese adults: Results from the China National Stroke Prevention Project (CSPP)

Running title: Adiposity and blood pressure

Wenning Fu¹, Shiyi Cao¹, Bing Liu², Haibin Li³, Fujian Song⁴, Yong Gan¹,
Wenzhen Li¹, Qiaoyan Liu¹, Longde Wang⁵, Shijiao Yan¹, Wei Yue⁶, Feng Yan⁶,
Chao Wang¹, Hui Li¹, Xiaojun Wang¹, Zhihong Wang^{8*}, Zuxun Lu^{1*}

¹ School of Public Health, Tongji Medical College, Huazhong University of Science and Technology, China

² Center of Health Administration and Development Studies, Hubei University of Medicine, China

³ Department of Epidemiology and Health Statistics, School of Public Health Capital Medical University, China

⁴ Norwich Medical School, Faculty of Medicine and Health Science, University of East Anglia, Norwich, United Kingdom

⁵ The National Health and Family Planning Commission, Beijing, China

⁶ Department of Neurology, Tianjin Huanhu Hospital, Tianjin, China

⁷ Department of Neurosurgery, Xuanwu Hospital, Capital medical University, Beijing, China

⁸ Department of Neurosurgery, Shenzhen Second People's Hospital, Shenzhen University, Shenzhen, Guangdong, China

*Corresponding author

Address: No. 13 Hangkong Road, Wuhan 430030, China

E-mail: Zuxun Lu: zuxunlu@yahoo.com

Zhihong Wang: 247506385@qq.com

Telephone and fax numbers: +86-027-83693756

Conflicts of Interest and Source of Funding

We declare we have no conflict of interest. This study was supported by the Fundamental Research Funds for the Central Universities, Huazhong University of Science and Technology, China. (2016YXMS215).

Word count:4734

Number of table: 1

Number of figures: 4

Supplementary material: 4 figures

Abstract

Background: The American Heart Association recommended that waist circumference was a better predictor of blood pressure risk than body mass index in Asians. However, data are inconsistent and information in Chinese, the largest global population group, is limited.

Methods: Data was obtained from the Chinese National Stroke Prevention Project Survey of a nationally representative sample of middle-aged and older Chinese adults. A total of 135825 individuals not taking any antihypertensive drugs were included in this study. Multiple linear regression analyses were conducted to examine the association between blood pressure and parameters of general adiposity, including body mass index, height-adjusted weight, and parameters of central adiposity, including waist circumference, hip circumference, waist-hip ratio, and waist-height ratio. Results were shown as mean difference in blood pressure associated with one standard deviation higher level of adiposity.

Results: The overall means \pm standard deviation of BMI and waist circumference were 24.3 \pm 3.18 kg/m² and 84.0 \pm 8.88 cm, respectively. BMI seemed more strongly associated with SBP/DBP (4.22 mmHg/SD; 2.60 mmHg/SD) than central adiposity markers. In addition, there were sex differences. For men, waist circumference showed a stronger association with SBP/DBP than BMI (4.04 vs. 3.79, $P < 0.05$; 2.26 vs. 2.13, $P < 0.05$). For women, BMI was more closely related to SBP/DBP than central adiposity parameters, such as waist circumference (4.59 vs. 3.41, $P < 0.05$; 2.98 vs. 2.24, $P < 0.05$). Additionally, in both urban and rural areas, waist circumference was mostly associated with SBP/DBP

among men, whereas it was BMI among women.

Conclusions: Compared with central adiposity, blood pressure is more strongly associated with general adiposity in Chinese adults. Interestingly, there are significant sex differences in the relationship of blood pressure with general and central adiposity. Waist circumference is the strongest predictor for men but suboptimal for women, and BMI tend to a better predictor of blood pressure for women. In addition, our results for men are consistent with the recommendation of the American Heart Association in 2015 that waist circumference could be used for assessing the risk of blood pressure.

Keywords: general adiposity; central adiposity; blood pressure; China

Introduction

High blood pressure (BP) is a major cause of coronary heart disease and stroke, and contributes to morbidity and premature mortality in both developed and developing countries^[1-4]. The causes of high BP are complex and are correlated with numerous environmental and genetic factors. Among them, adiposity is a well-established modifiable risk factor^[5-7]. Previous studies have reported that body mass index (BMI), a general measure of adiposity, is a predictor of BP^[8-9]. Subsequently, a growing amount of evidence has indicated that central adiposity parameters, such as waist circumference (WC), hip circumference (HC), waist-hip ratio (WHR), and waist-to-height ratio (WHtR), are also related to BP^[10-11]. Several studies have investigated the association of different measures of central and general adiposity with BP, but findings have been inconsistent. Some studies tended to suggest that WC is more strongly associated with BP as compared to BMI^[8-9]. But others did not support this, showing that BMI was a better predictor of BP than WC^[12-13]. Moreover, the majority of existing studies were conducted in Western populations^[14-15]. China has the largest population in the world (with one-fifth of the world's population), but there has been no research based on a nationally representative adult sample in China. Additionally, great differences in genetic backgrounds and the criteria for evaluating obesity by BMI exist between Chinese and Western populations^[16]. A good understanding of associations between BP and different measures of general obesity or central adiposity in China is of particular importance, which may not only help to inform public health policies for the control of high BP and related cardiovascular diseases, but also provide

valuable evidence on the topic in the international cardiovascular diseases -related research field.

Although BMI has been used to measure adiposity in many countries including Asians, its sensitivity was low in the assessment of the risk of cardiovascular disease^[17]. In 2015, the American Heart Association (AHA) recommended that the WC should be used to assess the risk of cardiovascular diseases in Asians^[17]. Studies have shown that control of blood pressure could reduce the risk of cardiovascular disease^[18-19], and recent studies have reported that WC was associated with BP^[20-21]. However, these studies mainly focused on the relationship between WC and BP in western population and Chinese children. Research in a nationally representative sample of Chinese adults has not been available.

The present study aimed to evaluate the association between BP and different measures of general or central obesity based on data from a national population-based study of Chinese adults.

Subjects and Methods

Study population and sampling

The China National Stroke Prevention Project (CSPP) was a national campaign on stroke prevention and control launched in 2011 by the Chinese government. The CSPP survey was carried out in 30 provinces of China from September 2014 to May 2015, administered by the National Project Office of CSPP. The CSPP survey used a two-

stage stratified cluster sampling method. At the first stage, 200 project areas were determined in proportion to the local population size and numbers of total counties. At the second stage, an urban community and a rural village were selected as primary sampling units according to geographical location and suggestions from local hospitals. The cluster sampling method was used in every primary sampling unit, and all residents aged 40 years or older were surveyed during the primary screening. A total of 849,874 adults aged ≥ 40 years were registered at the local government office. Of these adults, 726,451 received the stroke screening, yielding a response rate of 85.48%. Questionnaire completion and the assessment of stroke risk factors were conducted in primary health care institutions. About 180000 subjects were selected randomly for further assessment of blood pressure and physical examination.

Written informed consent was obtained from all individuals, and the study protocol was approved by the Ethics Committee of the Xuanwu Hospital Institutional Review Board, Capital Medical University (Beijing, China).

Data collection

Data collection was conducted by trained medical staff using a structured questionnaire in community health centers. Information obtained included sociodemographic characteristics, lifestyle factors (including physical activity, smoking and alcohol consumption), and medical history. The senior investigators checked the completed questionnaires daily as a quality control measure. Data were entered into a database in a double-blind manner by two researchers using EpiData 3.0.

Blood pressure measurement

Subjects sat quietly in a chair with their back supported, both feet flat on the floor, and arms supported at the heart level during blood pressure measurement. A trained nurse used an automated Sphygmomanometer to measure blood pressure. Three measurements were taken in the morning between 08:00 and 09:00 am at 5-min intervals after a 20-min rest. We used the mean values of the three recorded measurements for the analysis. Considering that systolic blood pressure (SBP) is a better predictor of vascular mortality than diastolic blood pressure (DBP), SBP was used in the main analyses.^[22] Results for DBP were provided in a supplementary file.

Body adiposity measures

The anthropometric indices were measured using standard techniques and equipment with the subjects in light clothing after an overnight fast. Height was measured by a portable stadiometer to within 0.1cm without shoes. Weight was measured using a digital scale to within 0.1 kg without heavy outer garments. WC was measured at the midpoint between the iliac crest and the lower rib, and HC was measured at the widest circumference over the gluteal muscles. Both WC, and HC were measured to the nearest 0.1 cm.

Covariate variables

Physical activity was defined as ≥ 3 times per week for at least 30 minutes each time, or engaged in heavy physical work. Low level of physical activity refers to the action of rare exercise or light physical work lasting for more than 1 year, during which the

exercise is done for less than 30mins each time and less than 3 times a week. Smoking was defined as smoking ≥ 1 cigarette per day in the last 3 months. Drinking was defined as alcohol drinking at least once per week for more than half a year.

Statistical analysis

The present study assessed six adiposity variables either directly measured or derived, including two general and four central adiposity measurements. The two anthropometric markers of general adiposity are height-adjusted weight (HtaW) and BMI, and the four anthropometric markers of central adiposity are WC, HC, WHR, and WHtR. BMI was calculated as weight (in kilograms) divided by height (in meters) squared. WHR was calculated as WC divided by HC, and WHtR was WC divided by height.

To control measurement errors, individuals with implausible values of adiposity measures (e.g. BMI <15 or >50 kg/m²; n=699) and blood pressures (SBP <80 or >250 mm Hg; DBP <40 or >150 mm Hg; n=97) or missing data (n=758) were excluded. We also excluded participants currently taking antihypertensive drugs (42926). Finally, a total of 135825 participants remained in the analysis.

Categorical data were described as percentages and continuous data were described as means \pm standard deviation (SD). All continuous data was abnormally distributed even after logarithmic transformation using a Shapiro-Wilk test. Differences in the distribution of baseline characteristics between males and females were tested using a Student's t-test for continuous variables (age, height, weight, BMI, WC, HC, WHR, WHtR, SBP and DBP) and a chi-square test for categorical variables (smoking, drinking

status and physical activity). We used multiple linear regression to estimate the mean difference in SBP and DBP associated with a 1 SD higher level of adiposity. In the multiple linear regression analysis, covariates included age (year in continuous), smoking status (yes, no), alcohol consumption (yes, no) and physical activity (yes, no). All statistical analyses were performed using SPSS19.0 (SPSS Inc, Chicago, Ill), and Stata 13.0. A value of $P < 0.05$ was considered as statistically significant.

Results

Of the 135825 individuals remained in the final analysis, there were 62451 men (mean age, 58 ± 11 years) and 73374 women (mean age, 59 ± 10 years). Demographic, lifestyle, and anthropometric characteristics stratified by gender of the overall sample and final participants are shown in Table 1. The 135825 subjects had a mean BMI of 24.3 ± 3.18 kg/m² (men: 24.3 ± 3.01 kg/m², women: 24.2 ± 3.31 kg/m²), mean WC of 84.0 ± 8.88 cm (men: 85.8 ± 8.89 cm, women 82.4 ± 8.57 cm), mean HC of 95.3 ± 9.35 cm (men: 96.5 ± 9.27 cm, women: 94.2 ± 9.29 cm), mean WHR of 0.883 ± 0.061 (men: 0.891 ± 0.061 , women: 0.876 ± 0.061), and mean WHtR of 0.515 ± 0.056 (men: 0.509 ± 0.053 , women: 0.520 ± 0.057). Compared with women, men had significantly higher SBP, DBP, and were more likely to be smokers, drinkers, and physically inactive (all $p < 0.05$).

Associations between markers of general and central adiposity with SBP are shown in Figure 1. Compared to central adiposity measures, BMI was the most significant predictor of SBP after considering age, smoking, drinking, physical activity and

anthropometric variable. That is, one SD increase in BMI was corresponding to an increase of SBP by about 4.0 mmHg. The next strongest association after adjusting covariates was the HtaW of general adiposity, with each one SD increase associated with about a 3.8 mm Hg higher SBP. The associations for other measures of adiposity were less significant. Similar relationships were observed between different measures of obesity and DBP.

Figure 2 shows the relation of SBP with different anthropometric indices of adiposity separately for men and women. For males, WC was the most strongly related to SBP after adjusting for age, smoking, drinking, physical activity and anthropometric variable, in which 1 SD higher WC (8.89 cm) was associated with an increase in SBP by 3.88 mmHg. For females, BMI was the strongest association with SBP. After adjusting for covariate variables, a one SD higher BMI was associated with a 4.41 mmHg higher SBP. HtaW, HC, WHR, and WHtR had a relatively weaker association with SBP in both males and females. The main findings for SBP were also broadly applicable for DBP.

To verify the gender differences in adiposity and BP in our study, an analysis among men and women in urban and rural areas was performed. Figure 3 and Figure 4 show the associations between SBP and general or central adiposity among males and females in urban and rural areas. For males in both urban and rural areas, WC was the strongest predictor of SBP, with each 1 SD higher WC associated with about a 3.79-mm Hg and a 3.85-mm Hg higher SBP, respectively. In contrast, compared to central adiposity markers, BMI was observed to have the strongest association with SBP among females in both urban and rural areas, with an increase of 1 SD corresponding to an increase of

about 4.41 mmHg and 4.47 mmHg SBP, respectively. Among males and females in both areas, the association between HtaW, HC, WHR, WHtR, and BP was less significant for BMI and WC after adjusting for covariate variables. Similar findings were observed for DBP.

Discussion

In the present study, data from a nationally representative study of 135,825 Chinese adults was utilized to examine the relationship between different measures of adiposity and BP. Compared with measures of central adiposity, BMI was observed to be closely associated with BP, with significant gender differences inherent in this association. Specifically, central adiposity measurement such as WC in males and general adiposity parameter such as BMI in females had the strongest associations with SBP/DBP. On average, every 3.18kg/m² higher BMI was associated with about 4.04 mm Hg higher SBP and 2.51 mmHg higher DBP. For other markers of adiposity, including HtaW, HC, WHtR and WHR were relatively poor predictors of SBP/DBP in this Chinese adult population. We believe that these findings may also applicable to other Asian countries.

In this study, WC showed the strongest association with BP in males, and BMI showed the strongest association with BP in females, which are in line with findings from some previous studies^[15, 23-24]. It was also consistent with the statement from the AHA in 2015, that suggested using WC for evaluating the risk of cardiovascular disease^[17]. However, our findings differ from several other studies that reported BMI in men and WC in

women having the strongest associations with BP ^[25-27]. The difference may be explained by the differences in the criteria for evaluating obesity by BMI between Chinese and other populations. It was claimed that the average BMI of Chinese adults with metabolic abnormalities was lower than the WHO's current overweight cut-off point 25kg/m², so that an overweight cut-off point of 24kg/m² for Chinese has been suggested by WHO ^[16]. Therefore, the relationship between obesity and BP in Chinese could be different from that in other populations. Another explanation is that use of a larger sample, as in the current study, rather than a small population sample, may also underlie discrepant results.

The strength of associations of BP with BMI was stronger in this Chinese population compared with that previously reported in Western populations ^[14]. It is well known that there are considerable differences in body fat distribution by ethnicity. Compared with Europeans, Asians tend to have more generalized adiposity ^[24]. Generalized adiposity is associated with higher BP and other metabolic risk factors ^[28]. Thus, different body fat distribution may partly account for these between-population differences. Relatively weak associations found in the Western populations might be the result of antihypertensive treatment that is common in western countries, even though such information was not reported in any studies ^[29]. Furthermore, compared with Western populations, dietary sodium intake is higher in adults in China ^[30], whereas sodium intake predicts SBP independently of BMI ^[31], and variation in sodium intake could also explain the differences between these populations.

Interestingly, there are significant gender differences for the relationship between

different measures of adiposity and BP. WC, an indicator of central adiposity, showed the strongest association with BP in males, and BMI, a measure of general obesity, showed the strongest association with BP in females. The potential reasons for these gender-specific differences are uncertain, although it may be partly explained by the gender difference in body fat distribution. Visceral fat is predominant in men, and subcutaneous fat is predominant in women ^[32]. Visceral fat is more likely to result in metabolic abnormalities than subcutaneous fat ^[33]. In addition, visceral adiposity has the positive correlation with BP ^[34]. In Asian men with less subcutaneous fat, WC are more appropriate than BMI to indicate visceral fat. Besides, in Asian women with more subcutaneous fat, BMI may be more suitable than waist circumference to indicate total and abdominal fat accumulation ^[15]. Additionally, compared with females, daily work stress and economic pressure may stimulate males to produce more stress hormones and adrenaline, resulting in more fat into the abdomen ^[23]. Because WC is a better indicator of abdominal fat ^[35], it was more strongly associated with BP in males. Further population and laboratory researches are clearly warranted to study the different biological mechanisms between obesity and BP in both males and females.

Proposed mechanisms behind the association of adiposity with BP are complex. Sympathetic nervous system (SNS) up-regulation has been an important mechanism of obesity-related blood pressure. Adipose tissue, especially perivascular adipose tissue, yield leptin, effects on the hypothalamic neuronal systems include reduced food consumption and upregulated thermogenesis and energy expenditure, and stimulation of sympathetic activity. Obesity increases plasma renin and angiotensinogen, which

activates the renin - Angiotensin System (RAS). RAS could also make the SNS upregulating^[36-37].

Strengths and limitations

The present study has several strengths. Firstly, to the best of our knowledge, this is the first study with a national sample of Chinese adults to compare the relationship between different measures of adiposity with BP. Secondly, this analysis included nearly 140000 adults in China, with adequate statistical power to compare different obesity measures. Limitations of this study were that this was a cross-sectional design, and we cannot infer causality from these results. In addition, all study participants were aged 40 or older, thus results may not be generalizable to younger populations, although similar results have been observed in younger subjects^[14, 23].

Conclusion

General obesity is more closely associated with BP than central adiposity, and that gender differences exist in these relationships. Based on the evidence from the present study and the statement of the AHA in 2015, we suggest that in addition to BMI, WC, an index that takes 1-2 minutes to measure, should be added for the general health checkups for Chinese populations. Further prospective population-based studies and mechanistic studies are still needed to validate our findings.

Acknowledgements

We thank the 726,451 study participants of the CSPP and all staff members involved in this study for their painstaking efforts in conducting the data collection.

Sources of Funding

This study was sponsored by the Ministry of Finance of the People's Republic of China (Issued by Finance and Social Security [2011] Document No. 61, Ministry of Finance).

Authors' contributions

WNF and ZXL conceived and designed the study. LDW is in charge of this project. SYC, BL, FJS, YG, WZL, SJY, WC, XJW and HL participated in the acquisition of data. WNF and SYC analyzed the data. BL and YG gave advice on methodology. WNF drafted the manuscript, and YG, WNF, BL and ZXL revised the manuscript. All authors read and approved the final manuscript. ZXL is the guarantor of this work and had full access to all the data in the study and take responsibility for its integrity and the accuracy of the data analysis.

References

1. Lu Y, Hajifathalian K, Ezzati M, Woodward M, Rimm EB, Danaei G, et al. Metabolic mediators of the effects of body-mass index, overweight, and obesity on coronary heart disease and stroke: a pooled analysis of 97 prospective cohorts with 1.8 million participants. *Lancet*. 2014; 383 (9921):970-83.
2. Ng M, Fleming T, Robinson M, Thomson B, Graetz N, Margono C, et al. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet*. 2014; 384 (9945):766-81.
3. Bell AC, Adair LS, Popkin BM. Ethnic Differences in the Association between Body Mass Index and Hypertension. *American Journal of Epidemiology*. 2002; 155 (4):346.
4. Brown CD, Higgins M, Donato KA, Rohde FC, Garrison R, Obarzanek E, et al. Body mass index and the prevalence of hypertension and dyslipidemia. *Obesity Research*. 2000; 8 (9):605-19.
5. Timpson NJ, Harbord R, Davey SG, Zacho J, Tybjaerghansen A, Nordestgaard BG. Does greater adiposity increase blood pressure and hypertension risk?: Mendelian randomization using the FTO/MC4R

- genotype. *Hypertension*. 2009; 54 (1):84-90.
6. Dorresteyn JA, Visseren FL, Spiering W. Mechanisms linking obesity to hypertension. *Obesity Reviews An Official Journal of the International Association for the Study of Obesity*. 2012; 13 (1):17-26.
 7. Hall JE, do Carmo JM, Da SA, Wang Z, Hall ME. Obesity-induced hypertension: interaction of neurohumoral and renal mechanisms. *Circulation Research*. 2015; 116 (6):991-1006/
 8. Buss J. Limitations of body mass index to assess body fat. *Workplace Health & Safety*. 2014; 62 (6):264.
 9. Fletcher I. Defining an epidemic: the body mass index in British and US obesity research 1960-2000. *Sociol Health Illn*. 2014; 36 (3):338-53.
 10. Huxley R, James WP, Barzi F, Patel JV, Lear SA, Suriyawongpaisal P, et al. Ethnic comparisons of the cross-sectional relationships between measures of body size with diabetes and hypertension. *Obesity Reviews An Official Journal of the International Association for the Study of Obesity*. 2008; 9 Suppl 1 (s1):53..
 11. Lee CM, Huxley RR, Wildman RP, Woodward M. Indices of abdominal obesity are better discriminators of cardiovascular risk factors than BMI: a meta-analysis. *Journal of Clinical Epidemiology*. 2008; 61 (7):646.
 12. Wang ZM, Heshka S, Heo M. Waist circumference and obesity-associated risk factors among whites in the third National Health and Nutrition Examination Survey: clinical action thresholds. *American Journal of Clinical Nutrition*. 2002; 76 (4):743.
 13. Janssen I, Katzmarzyk PT, Ross R. Waist circumference and not body mass index explains obesity-related health risk. *American Journal of Clinical Nutrition*. 2004; 79 (3):379.
 14. Gnatiuc L, Alegrediaz J, Halsey J, Herrington WG, Lewington S, Collins R, et al. Adiposity and Blood Pressure in 110 000 Mexican Adults. *Hypertension*. 2017; 69 (4):608-14.
 15. Sakurai M, Miura K, Takamura T, Ota T, Ishizaki M, Morikawa Y, et al. Gender Differences in the Association between Anthropometric Indices of Obesity and Blood Pressure in Japanese. *Hypertension Research Official Journal of the Japanese Society of Hypertension*. 2006; 29 (2):75.
 16. Consultation WE. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet*. 2004; 363 (9403):157.
 17. Rao G, Powell-Wiley TM, Ancheta I, Hairston K, Kirley K, Lear SA, et al. Identification of Obesity and Cardiovascular Risk in Ethnically and Racially Diverse Populations: A Scientific Statement From the American Heart Association. *Circulation*. 2015; 132 (5):457.
 18. Arnett DK, Evans GW, Riley WA. Arterial stiffness: a new cardiovascular risk factor? *Am J Epidemiol*. 1994; 140(8):669-82.
 19. Bönner G. Risk evaluation and therapeutical implications of pulse pressure in primary arterial hypertension. *Dtsch Med Wochenschr*. 2002; 127(45):2396-9.
 20. Zhang YX1, Wang SR. The relationship of waist circumference distribution to blood pressure levels among children and adolescents in Shandong, China. *Int J Cardiol*. 2013; 168(2):1516-20.
 21. Benmohammed K1, Valensi P2, Nguyen MT, et al. Influence of waist circumference on blood pressure status in non-obese adolescents. *Int J Adolesc Med Health*. 2018;15.
 22. Lewington S, Clarke R, Qizilbash N, Peto R, Collins R. Age-specific relevance of usual blood pressure to vascular mortality: a meta-analysis of individual data for one million adults in 61 prospective studies. *Lancet*. 2003; 361 (9366):1389; author reply 91.
 23. Dobbelssteyn CJ, Joffres MR, Maclean DR, Flowerdew G. A comparative evaluation of waist circumference, waist-to-hip ratio and body mass index as indicators of cardiovascular risk factors. The canadian heart health surveys. *International Journal of Obesity & Related Metabolic Disorders Journal*

- of the International Association for the Study of Obesity. 2001; 25: 652.
24. Lear SA, Humphries KH, Kohli S, Chockalingam A, Frohlich JJ, Birmingham CL. Visceral adipose tissue accumulation differs according to ethnic background: results of the Multicultural Community Health Assessment Trial (M-CHAT). *American Journal of Clinical Nutrition*. 2007; 86 (2):353.
 25. Dalton M, Cameron AJ, Zimmet PZ, Shaw JE, Jolley D, Dunstan DW, et al. Waist circumference, waist-hip ratio and body mass index and their correlation with cardiovascular disease risk factors in Australian adults. *Journal of Internal Medicine*. 2003; 254 (6):555-63.
 26. Zhu SK, Wang ZM, Heshka S, Heo M, Faith MS, Heymsfield SB. Waist circumference and obesity-associated risk factors among whites in the third National Health and Nutrition Examination Survey: clinical action thresholds. *American Journal of Clinical Nutrition*. 2002; 76 (4):743.
 27. Foucan L, Hanley J, Deloumeaux J, Suissa S. Body mass index (BMI) and waist circumference (WC) as screening tools for cardiovascular risk factors in Guadeloupean women. *Journal of Clinical Epidemiology*. 2002; 55 (10):990-6.
 28. Fox CS, Massaro JM, Hoffmann U, Pou KM, Maurovichhorvat P, Liu CY, et al. Abdominal Visceral and Subcutaneous Adipose Tissue Compartments Association With Metabolic Risk Factors in the Framingham Heart Study. *Circulation*. 2007; 116 (1):39.
 29. Whitlock, Lewington G, Sherliker S, Clarke P, Emberson R, Halsey J, et al. Body-mass index and cause-specific mortality in 900 000 adults: collaborative analyses of 57 prospective studies. *Lancet*. 2009; 373 (9669):1083-96.
 30. Dyer AR, Elliott P, Shipley M, Stamler R, Stamler J. Body mass index and associations of sodium and potassium with blood pressure in INTERSALT. *Hypertension*. 1994; 23 (1):729-36.
 31. He FJ, Macgregor GA. Effect of longer-term modest salt reduction on blood pressure: John Wiley & Sons, Ltd; 2008.
 32. Tchernof A. Pathophysiology of Human Visceral Obesity: An Update. *Physiological Reviews*. 2013; 93 (1):359.
 33. Nakamura T, Tokunaga K, Shimomura I, Nishida M, Yoshida S, Kotani K, et al. Contribution of visceral fat accumulation to the development of coronary artery disease in non-obese men. *Atherosclerosis*. 1994; 107 (2):239-46.
 34. Hayashi T, Boyko EJ, Leonetti DL, Mcneely MJ, Newell-Morris L, Kahn SE, et al. Visceral adiposity and the prevalence of hypertension in Japanese Americans. *Circulation*. 2003; 108 (14):1718-23.
 35. Pouliot MC, Després JP, Lemieux S, Moorjani S, Bouchard C, Tremblay A, et al. Waist circumference and abdominal sagittal diameter: best simple anthropometric indexes of abdominal visceral adipose tissue accumulation and related cardiovascular risk in men and women. *American Journal of Cardiology*. 1994; 73 (7):460-8.
 36. Armitage JA, Burke SL, Prior LJ, Barzel B, Eikelis N, Lim K, et al. Rapid onset of renal sympathetic nerve activation in rabbits fed a high-fat diet. *Hypertension*. 2012; 60 (1):163.
 37. Leopold JA. Cellular and molecular mechanisms of arterial stiffness associated with obesity. *Hypertension*. 2013; 62 (6):1003-4.

Table1 Baseline characteristics of the study subjects stratified by genders

Variables	Total(n=135825)	Males(n=62451)	Females(n=73374)	P
Age groups, n (%)				<0.01
40-49	42535 (31.3)	19943 (31.9)	22592 (30.8)	
50-59	40226 (29.6)	18014 (28.8)	22212 (30.3)	
60-69	33473 (24.60)	15259 (24.4)	18214 (24.8)	
≥70	19591 (14.4)	9235 (14.8)	10356 (14.1)	
Height (cm)	163±8.0	169±6.3	159±6.2	
Weight (kg)	64.7±10.3	69.2±9.9	60.9±9.0	<0.01
BMI (kg/m ²)	24.3±3.18	24.3±3.01	24.2±3.31	<0.01
WC (cm)	84.0±8.88	85.8±8.89	82.4±8.57	<0.01
HC (cm)	95.3±9.35	96.5±9.27	94.2±9.29	<0.01
WHR	0.883±0.061	0.891±0.061	0.876±0.061	<0.01
WHtR	0.515±0.056	0.509±0.053	0.520±0.057	<0.01
SBP (mmHg)	129±16	130±12	129±17	<0.01
DBP (mmHg)	80±10	81±10	79±10	<0.01
Smoking n (%)				<0.01
Yes	22830 (16.8)	20560 (32.9)	2270 (9.9)	
No	112995 (83.2)	41891 (67.1)	71104 (96.9)	
Drinking status n (%)				<0.01
Yes	19287 (14.2)	18110 (29)	2127 (2.7)	
No	116538 (85.7)	44341 (71)	71247 (97)	
Physical activity n (%)				
Yes	33394 (24.6)	14916 (23.9)	18478 (25.2)	<0.01
No	102431 (75.5)	47535 (76.1)	54896 (74.8)	

Categorical data were described as percentages and continuous data were described as means ± SD.

BMI: body mass index; HtaW: height-adjusted weight; WC: waist circumference; HC: hip circumference;

WHR: waist-hip ratio; WHtR: Waist-height ratio; SBP: systolic blood pressure; DBP: diastolic blood pressure.

Figure legends

Figure 1 Association between markers of adiposity and systolic blood pressure

95% CI = 95% confidence interval. Adjust for age, smoking, drinking and physical activity.

Figure 2 Association between markers of adiposity and systolic blood pressure in both men and women

95% CI = 95% confidence interval. Adjust for age, smoking, drinking and physical activity.

Figure 3 Association between markers of adiposity and systolic blood pressure in urban areas

95% CI = 95% confidence interval. Adjust for age, smoking, drinking and physical activity.

Figure 4 Association between markers of adiposity and systolic blood pressure in rural areas

95% CI = 95% confidence interval. Adjust for age, smoking, drinking and physical activity.