

1 **PGC-1 α driven mitochondrial biogenesis in stromal cells underpins**
2 **mitochondrial trafficking to leukemic blasts**

3

4 Christopher R Marlein¹, Lyubov Zaitseva¹, Rachel E Piddock¹, Livia Raso-Barnett²,
5 Michael A Scott², Christopher J Ingham³, Angela Collins⁴, Kristian M Bowles^{1,4*}, Stuart
6 A Rushworth^{1*}

7

8 ¹Department of Molecular Haematology, Norwich Medical School, The University of
9 East Anglia, Norwich Research Park, NR4 7TJ, United Kingdom

10 ²Haematopathology and Oncology Diagnostic Service, Addenbrooke's Hospital, Hills
11 Rd, Cambridge, CB2 0QQ, United Kingdom

12 ³Department of Trauma and Orthopaedic Surgery, Norfolk and Norwich University
13 Hospitals NHS Trust, Colney Lane, Norwich, NR4 7UY, United Kingdom

14 ⁴Department of Haematology, Norfolk and Norwich University Hospitals NHS Trust,
15 Colney Lane, Norwich, NR4 7UY, United Kingdom

16

17 * denotes joint corresponding authors

18

19 email: s.rushworth@uea.ac.uk / k.bowles@uea.ac.uk

20

21 Department of Molecular Haematology,
22 Norwich Medical School,
23 University of East Anglia,
24 Norwich Research Park,
25 Norwich, NR4 7TJ,
26 United Kingdom: Tel: 01603 591802

27

28 **Word count: 1480**

29

30 **Running title: Mitochondrial biogenesis in BMSC is a prerequisite for**
31 **mitochondrial transfer to AML.**

32

33 **Keywords: Mitochondrial transfer, AML, PGC-1 α**

34 Acute myeloid leukemia (AML) is a disease known to be heavily reliant on its bone
35 microenvironment (BMM) to survive and proliferate ^{1,2}. We have previously shown that
36 AML disease progression is enabled by the transfer of functional mitochondria to the
37 malignant cell from bone marrow stromal cells (BMSC) ^{3,4}. This process was shown
38 to be stimulated by superoxide generated by NADPH oxidase-2 (NOX2) on the AML
39 blast ³. However, beyond the stimulation of reactive oxygen species in BMSC, the
40 mechanisms controlling mitochondrial transfer in BMSC have yet to be elucidated.

41

42 There are no apparent adverse effects on BMSC after donation of mitochondria to
43 AML blasts, implying the presence of a mechanism whereby the BMSC can recover
44 their metabolic potential. The master regulator of mitochondrial biogenesis ⁵,
45 peroxisome proliferator-activated receptor gamma coactivator 1-alpha (PGC-1 α), has
46 been implicated in cancer progression and metabolism ^{6,7}. In these studies, PGC-1 α
47 is up-regulated and causes an increased accumulation of functional mitochondria.
48 Here, we investigate the effect AML has on the mitochondrial mass, bio-energetic
49 potential and PGC-1 α expression in BMSC.

50

51 First, using MitoTracker Green staining and flow cytometry, we determined the
52 mitochondrial levels in primary BMSC (n=8) after co-culture with AML blasts. In Figure
53 1A, we found significantly elevated mitochondrial levels in BMSC after co-culture,
54 implying that AML blasts stimulate BMSC to produce more mitochondria. We next
55 wanted to see if this increase in mitochondrial mass caused an increase in
56 mitochondrial based metabolism. To do this we analysed BMSC oxygen consumption
57 rate using the Seahorse extracellular flux assay. Increased mitochondrial respiration
58 was observed in BMSC (n=4) after co-culture with AML blasts (Figure 1B and C).
59 Moreover, in Figure 1D, we show that BMSC from patients with AML had increased
60 mitochondrial respiration compared to BMSC from healthy individuals (n=3). Together,
61 these results show that BMSC from patients with AML have increased mitochondrial
62 mass and functional bio-energetic consequence in BMSC metabolism.

63

64 As the transcription factor PGC-1 α is known to cause increased mitochondrial
65 biogenesis ⁵, we examined PGC1 α expression in BMSC after co-culture with AML
66 blasts. RNA expression of PGC-1 α in BMSC (n=5) was increased in BMSC after co-

67 culture with AML blasts compared to BMSC cultured alone (Figure 1F). Next, we
68 showed that total PGC-1 α protein was elevated in BMSC after AML co-culture
69 compared to control (Figure 1F). Moreover, we show that BMSC have increased
70 nuclear levels of PGC-1 α after co-culture with AML compared to BMSC cultured alone
71 (Figure 1F), an effect which was reversed upon the addition of N-acetylcysteine to the
72 co-culture (Supplementary figure 1). Previous studies have shown that AMPK can be
73 stimulated by ROS⁸ and in turn can stimulate PGC-1 α ⁹. Therefore as NOX-2 derived
74 ROS stimulates mitochondrial transfer to AML³, we assessed whether AMPK is
75 activated in BMSC after culture with AML blasts. We found increased phosphorylation
76 of Thr182 in AMPK from BMSC after co-culture with AML blasts (Supplementary figure
77 2). Together, results from RNA and Western blotting highlight that AML blasts cause
78 an increase in PGC-1 α expression and localization in BMSC suggesting that PGC-1 α
79 becomes activated via AMPK, in response to AML co-culture.

80

81 We next wanted to determine if elevated PGC-1 α expression and nuclear localization
82 and subsequent mitochondrial biogenesis was required for the mitochondrial transfer
83 from BMSC to AML blasts. Figure 2A shows that mitochondrial transfer occurs from
84 BMSC to the primary AML blasts. Next, we knocked down (KD) PGC-1 α in BMSC with
85 shRNA (Figure 2B). A MitoTracker Green based staining assay was then used to
86 analyse the levels of mitochondrial transfer to AML blasts cultured on control KD and
87 PGC-1 α KD BMSC. Figure 2C shows that mitochondrial transfer from BMSC to AML
88 is impaired when cultured on PGC-1 α KD BMSC compared with control KD BMSC
89 (Figure 2C). This data shows that PGC-1 α activation is prerequisite for pro-tumoral
90 mitochondrial transfer from BMSC to blasts in AML.

91

92 To investigate the effect of PGC-1 α on ROS and oxidative stress in BMSC, ROS levels
93 in PGC-1 α KD and control KD BMSC were analysed with and without AML co-culture.
94 Basal ROS levels were elevated in BMSC when PGC-1 α was knocked down
95 (Supplementary figure 3A), however upon the addition of AML reduced ROS levels
96 compared with control KD BMSC were observed (Supplementary figure 3B).
97 Therefore, AML blasts are unable to stimulate ROS in PGC-1 α KD BMSC to the same
98 extent as control KD BMSC, which would account for the reduced mitochondrial
99 transfer observed.

100 Finally, we wanted to examine the effect PGC-1 α KD in BMSC has on the disease
101 progression of AML. To do this we used an NSG mouse model whereby we
102 transplanted BMSC and AML blasts subcutaneously. Using this model, OCI-AML3
103 cells tagged with a luciferase construct ¹⁰ and then subcutaneously injected with
104 BMSC (into the right flank) and without BMSC (into the left flank), only proliferate in
105 the presence of BMSC (Supplementary Figure 1). We modified this model for use in
106 the PGC-1 α KD study, where we injected OCI-AML3 or MV4-11 luciferase cells with
107 control KD BMSC (left flank) or PGC-1 α KD BMSC (right flank). Figure 2D and 2E
108 show that AML combined with PGC-1 α KD BMSC has reduced tumor volume
109 compared with animals with control KD BMSC. Figure 2F shows the bioluminescence
110 from live animal imaging matches with excised tumors, where the tumors are reduced
111 in the PGC-1 α KD flank. Histologic analysis showed no difference between the AML
112 tumors grown with PGC-1 α KD BMSC compared to those that developed with control
113 KD BMSC; with respect to the type or frequency of inflammatory cells or other non-
114 malignant cells (Supplementary figure 5). Overall it was observed that PGC-1 α KD in
115 BMSC has a negative effect on AML disease progression *in vivo*.

116
117 In conclusion, this study provides a novel insight into the mechanisms controlling pro-
118 tumoral mitochondrial transfer in AML. We have shown that AML increases oxidative
119 stress in the BMSC ³, and this causes an increase in PGC-1 α expression and
120 mitochondrial biogenesis in BMSC. This process is prerequisite for the pro-tumoral
121 mitochondrial transfer from BMSC to leukemic blasts observed in AML. Inhibition of
122 PGC-1 α in BMSC reduces the trafficking of mitochondria and thus limits the
123 proliferative capacity of the tumor. As pro-tumoral mitochondrial transfer is increasingly
124 recognised as part of the malignant phenotype in multiple cancers ¹¹⁻¹³, this study
125 provides a novel mechanistic insight as to how PGC-1 α may be targeted in the
126 microenvironment as a means to limiting mitochondrial transfer to cancer. Treatments
127 inhibiting mitochondrial metabolism and function in AML blasts, including IDH1/2
128 mutant inhibitors ¹⁴ and the Bcl-2 inhibitor venetoclax, have recently been shown to be
129 clinically effective ¹⁵. This study also provides an important step in understanding the
130 complex nature of tumor metabolism, not only in the malignant cell, but also within the
131 microenvironment which supports it.

132

133 **Conflict of interest**

134 All authors declare no conflict of interest.

135

136 **Acknowledgements**

137 The authors thank the Rosetrees Trust and the Norwich Research Park Doctoral
138 Training Program for funding. They also thank Professor Richard Ball, Dr Mark
139 Wilkinson, Mr Iain Sheriffs, and Ms Sue Steel, Norwich Biorepository (UK) for help with
140 sample collection and storage. pCDH-luciferase-T2A-mCherry was kindly gifted by
141 Professor Irmela Jeremias, MD, from Helmholtz Zentrum München, Munich, Germany.
142 The authors also thank Dr Allyson Tyler, Dr Ian Thirkettle and Dr Karen Ashurst from
143 the Laboratory Medicine department at the Norfolk and Norwich University Hospital
144 for technical assistance.

145

146 **Authorship contributions**

147 CRM, KMB and SAR designed the research; CRM performed the research; CRM and
148 REP carried out *in vivo* work; LZ, LRB, MAS, CJI, AC and KMB provided essential
149 knowledge and reagents; CRM, KMB and SAR wrote the paper

150 **References**

- 151 1. Abdul-Aziz AM, Shafat MS, Mehta TK, Di Palma F, Lawes MJ, Rushworth SA, *et al.*
152 MIF-Induced Stromal PKCbeta/IL8 Is Essential in Human Acute Myeloid Leukemia.
153 *Cancer Res* 2017 Jan 15; **77**(2): 303-311.
- 154
155 2. Shafat MS, Oellerich T, Mohr S, Robinson SD, Edwards DR, Marlein CR, *et al.*
156 Leukemic blasts program bone marrow adipocytes to generate a protumoral
157 microenvironment. *Blood* 2017 Mar 09; **129**(10): 1320-1332.
- 158
159 3. Marlein CR, Zaitseva L, Piddock RE, Robinson SD, Edwards DR, Shafat MS, *et al.*
160 NADPH oxidase-2 derived superoxide drives mitochondrial transfer from bone marrow
161 stromal cells to leukemic blasts. *Blood* 2017 Oct 05; **130**(14): 1649-1660.
- 162
163 4. Moschoi R, Imbert V, Nebout M, Chiche J, Mary D, Prebet T, *et al.* Protective
164 mitochondrial transfer from bone marrow stromal cells to acute myeloid leukemic cells
165 during chemotherapy. *Blood* 2016 Jul 14; **128**(2): 253-264.
- 166
167 5. Fernandez-Marcos PJ, Auwerx J. Regulation of PGC-1alpha, a nodal regulator of
168 mitochondrial biogenesis. *Am J Clin Nutr* 2011 Apr; **93**(4): 884S-890.
- 169
170 6. LeBleu VS, O'Connell JT, Gonzalez Herrera KN, Wikman H, Pantel K, Haigis MC, *et al.*
171 PGC-1alpha mediates mitochondrial biogenesis and oxidative phosphorylation in
172 cancer cells to promote metastasis. *Nat Cell Biol* 2014 Oct; **16**(10): 992-1003, 1001-
173 1015.
- 174
175 7. Andrzejewski S, Klimcakova E, Johnson RM, Tabaries S, Annis MG, McGuirk S, *et al.*
176 PGC-1alpha Promotes Breast Cancer Metastasis and Confers Bioenergetic Flexibility
177 against Metabolic Drugs. *Cell Metab* 2017 Nov 7; **26**(5): 778-787 e775.
- 178
179 8. Rabinovitch RC, Samborska B, Faubert B, Ma EH, Gravel SP, Andrzejewski S, *et al.*
180 AMPK Maintains Cellular Metabolic Homeostasis through Regulation of Mitochondrial
181 Reactive Oxygen Species. *Cell Rep* 2017 Oct 3; **21**(1): 1-9.
- 182
183 9. Jager S, Handschin C, St-Pierre J, Spiegelman BM. AMP-activated protein kinase
184 (AMPK) action in skeletal muscle via direct phosphorylation of PGC-1alpha. *Proc Natl*
185 *Acad Sci U S A* 2007 Jul 17; **104**(29): 12017-12022.
- 186
187 10. Vick B, Rothenberg M, Sandhofer N, Carlet M, Finkenzeller C, Krupka C, *et al.* An
188 advanced preclinical mouse model for acute myeloid leukemia using patients' cells of
189 various genetic subgroups and in vivo bioluminescence imaging. *PLoS One* 2015;
190 **10**(3): e0120925.
- 191
192 11. Lou E, Fujisawa S, Morozov A, Barlas A, Romin Y, Dogan Y, *et al.* Tunneling
193 nanotubes provide a unique conduit for intercellular transfer of cellular contents in
194 human malignant pleural mesothelioma. *PLoS One* 2012; **7**(3): e33093.

196 12. Lu J, Zheng X, Li F, Yu Y, Chen Z, Liu Z, *et al.* Tunneling nanotubes promote
197 intercellular mitochondria transfer followed by increased invasiveness in bladder
198 cancer cells. *Oncotarget* 2017 Feb 28; **8**(9): 15539-15552.

199
200 13. Pasquier J, Guerrouahen BS, Al Thawadi H, Ghiabi P, Maleki M, Abu-Kaoud N, *et al.*
201 Preferential transfer of mitochondria from endothelial to cancer cells through tunneling
202 nanotubes modulates chemoresistance. *J Transl Med* 2013 Apr 10; **11**: 94.

203
204 14. Stein EM, DiNardo CD, Pollyea DA, Fathi AT, Roboz GJ, Altman JK, *et al.* Enasidenib
205 in mutant IDH2 relapsed or refractory acute myeloid leukemia. *Blood* 2017 Aug 10;
206 **130**(6): 722-731.

207
208 15. Dinardo CD, Pratz KW, Potluri J, Pullarkat VA, Jonas BA, Wei AH, *et al.* Durable
209 response with venetoclax in combination with decitabine or azacitadine in elderly
210 patients with acute myeloid leukemia (AML). *J Clin Oncol*, 36 2018; **suppl**;(abstract
211 7010).

212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233

234 **Figure legends**

235

236 **Figure 1. AML stimulates mitochondrial biogenesis in BMSC through PGC-1 α**

237 (A) BMSC were cultured with AML blasts for 24 hours, then BMSC (n=8) were stained
238 with 200 nM MitoTracker Green FM. Mitochondrial levels were analysed using
239 MitoTracker Green mean fluorescence intensity by flow cytometry. (B) A
240 representative plot of oxygen consumption rate from the Seahorse MitoStress assay.
241 Oligomycin (O), FCCP (F) and rotenone (R) were injected periodically and oxygen
242 consumption rate was measured. (C and D) Basal and maximum mitochondrial
243 respiration in BMSC cultured with and without AML blasts (C) and from AML and
244 healthy patients (D) (n=3). (E) RNA qPCR analysis of BMSC (n=5) with and without
245 co-culture with AML blasts. (F) Western blot analysis of nuclear, cytosolic and total
246 PGC-1 α protein from BMSC (n=2) cultured with and without AML blasts (n=2).

247

248 **Figure 2. PGC-1 α is crucial for mitochondrial transfer and AML disease**
249 **progression.**

250 (A) MitoTracker Green based transfer assay showing that AML blasts, used in this
251 study, have acquired mitochondria from BMSC. (B) PGC-1 α RNA expression is
252 significantly reduced in BMSC after specific lentiviral targeting. (C) Mitochondrial
253 transfer levels to AML blasts are reduced when cultured on PGC-1 α KD BMSC. (D)
254 Schematic representation of the NSG mouse model used. (E) Bioluminescent live
255 animal images showing OCI-AML3/MV4-11 AML disease progression, when injected
256 subcutaneously with control KD or PGC-1 α BMSC. (F) Quantification of
257 bioluminescent images seen in E (OCI-AML3; n=5. MV4-11; n=4). (G) Excised tumors
258 were measured using calipers.

259