

Reducibility of equivalence relations arising from non-stationary ideals under large cardinal assumptions

David Asperó
Tapani Hyttinen Vadim Kulikov and Miguel Moreno

Abstract Working under large cardinal assumptions such as supercompactness, we study the Borel-reducibility between equivalence relations modulo restrictions of the non-stationary ideal on some fixed cardinal κ . We show the consistency of $E_{\lambda\text{-club}}^{\lambda^{++}, \lambda^{++}}$, the relation of equivalence modulo the non-stationary ideal restricted to $S_{\lambda}^{\lambda^{++}}$ in the space $(\lambda^{++})^{\lambda^{++}}$, being continuously reducible to $E_{\lambda^{++}\text{-club}}^{2, \lambda^{++}}$, the relation of equivalence modulo the non-stationary ideal restricted to $S_{\lambda^{++}}^{\lambda^{++}}$ in the space $2^{\lambda^{++}}$. Then we show that for κ ineffable $E_{\text{reg}}^{2, \kappa}$, the relation of equivalence modulo the non-stationary ideal restricted to regular cardinals in the space 2^{κ} , is Σ_1^1 -complete. We finish by showing, for Π_2^1 -indescribable κ , that the isomorphism relation between dense linear orders of cardinality κ is Σ_1^1 -complete.

1 Introduction

Throughout this article we assume that κ is an uncountable cardinal that satisfies $\kappa^{<\kappa} = \kappa$. The equivalence relations modulo (restrictions of) the non-stationary ideal have provided a very useful tool, and a main focus of study, in generalized descriptive set theory. In [1] it was shown that the relation of equivalence modulo the non-stationary ideal is not a Borel relation, and that if $V = L$, then it is not Δ_1^1 . The equivalence relation modulo the non-stationary ideal restricted to a stationary set S , denoted $E_S^{2, \kappa}$ (see Definition 1.3), is useful when it comes to studying the complexity of the isomorphism relations of first order theories (\cong_T , see Definition 1.5). In [1] it was proved that, under some cardinality assumptions, $E_{S_{\delta}^{\kappa}}^{2, \kappa}$ is Borel reducible to \cong_T for every first order stable unsuperstable theory T , where S_{δ}^{κ} is the set of λ -cofinal

2010 Mathematics Subject Classification: Primary 03E15; Secondary 03C45
Keywords: Large cardinals, Generalized Baire spaces, Equivalence relations

ordinals below κ . Similar results were obtained in [1] for the other non-classifiable theories. This motivates the study of the Borel-reducibility properties of $E_S^{2,\kappa}$.

Theorem 1.1 ([1], Theorem 56) *The following is consistent: For all stationary S and S' , $E_S^{2,\kappa}$ is Borel reducible to $E_{S'}^{2,\kappa}$ if and only if $S \subseteq S'$.*

Theorem 1.2 ([1], Theorem 55) *The following is consistent: $E_{S_\omega^2}^{2,\omega_2}$ is Borel reducible to $E_{S_\omega^1}^{2,\omega_2}$.*

In [6] the authors used the Borel-reducibility properties of the equivalence relation modulo the non-stationary ideal to prove that in L , all Σ_1^1 equivalence relations are reducible to \cong_{DLO} , where DLO is the theory of dense linear orderings without end points, which means that this equivalence relation is on top of the Borel-reducibility hierarchy among Σ_1^1 -equivalence relations, i.e. it is Σ_1^1 -complete. This result stands in contrast to the classical, countable case, $\kappa = \omega$, for which it is known that all other isomorphism relations are reducible to \cong_{DLO} [3], but far from all Σ_1^1 -equivalence relations are reducible to it; even some Borel-equivalence relations such as E_1 are not reducible to any isomorphism relations in the countable case. So the question remained: is the Σ_1^1 -completeness of \cong_{DLO} just a manifestation of the pathological behaviour of L or is it a more robust property in the generalised realm? One of the contributions of this paper is that the Σ_1^1 -completeness of \cong_{DLO} is indeed a rather robust phenomenon and holds whenever κ has certain large cardinal properties (Theorem 3.10).

It was asked in [2] and in [7, Question 3.46] whether or not the equivalence relation modulo the non-stationary ideal on the Baire space can be reduced to the Cantor space for some fixed cofinality: in our notation, whether or not $E_{S_\mu^\kappa}^{\kappa,\kappa} \leq E_{S_\mu^\kappa}^{2,\kappa}$. We approach the problem by proving several results in this direction. Our results have the forms

$$E_{S_\mu^\kappa}^{\kappa,\kappa} \leq E_{S_{\mu^*}^\kappa}^{2,\kappa},$$

$$E_{S_\mu^\kappa}^{\kappa,\kappa} \leq E_{\text{reg}(\kappa)}^{2,\kappa},$$

and

$$E_{\text{reg}(\kappa)}^{\kappa,\kappa} \leq E_{\text{reg}(\kappa)}^{2,\kappa},$$

where μ^* is larger than μ and $\text{reg}(\kappa)$ is the set of regular cardinals below κ , for κ Mahlo. These results are obtained under various assumptions and sometimes in forcing extensions.

Many of the results in the area of reducibility of equivalence relations modulo non-stationary ideals use combinatorial principles, like \diamond , and other reflection principles. In this paper we bring also some large cardinal principles into the picture.

The generalized Baire space is the set κ^κ with the bounded topology. For every $\zeta \in \kappa^{<\kappa}$, the set

$$[\zeta] = \{\eta \in \kappa^\kappa \mid \zeta \subset \eta\}$$

is a basic open set. The open sets are of the form $\bigcup X$ where X is a collection of basic open sets. The collection of κ -Borel subsets of κ^κ is the smallest set which contains the basic open sets and is closed under unions and intersections of length κ . Since in this paper we do not consider any other kind of Borel sets besides κ -Borel,

we will omit the prefix “ κ -”.

The generalized Cantor space is the subspace $2^\kappa \subset \kappa^\kappa$ with the relative subspace topology. For $X, Y \in \{\kappa^\kappa, 2^\kappa\}$, we say that a function $f: X \rightarrow Y$ is *Borel* if for every open set $A \subseteq Y$ the inverse image $f^{-1}[A]$ is a Borel subset of X . Let E_1 and E_2 be equivalence relations on X and Y respectively. We say that E_1 is *Borel reducible* to E_2 if there is a Borel function $f: X \rightarrow Y$ that satisfies $(\eta, \xi) \in E_1 \Leftrightarrow (f(\eta), f(\xi)) \in E_2$. We call f a *reduction* of E_1 to E_2 . This is denoted by $E_1 \leq_B E_2$, and if f is continuous, then we say that E_1 is *continuously reducible* to E_2 , which is denoted by $E_1 \leq_c E_2$.

For every stationary $S \subset \kappa$, we define the equivalence relation modulo the non-stationary ideal restricted to a stationary set S , on the space λ^κ for $\lambda \in \{2, \kappa\}$:

Definition 1.3 For every stationary $S \subset \kappa$ and $\lambda \in \{2, \kappa\}$, we define $E_S^{\lambda, \kappa}$ as the relation

$$E_S^{\lambda, \kappa} = \{(\eta, \xi) \in \lambda^\kappa \times \lambda^\kappa \mid \{\alpha < \kappa \mid \eta(\alpha) \neq \xi(\alpha)\} \cap S \text{ is not stationary}\}.$$

Note that $E_S^{2, \kappa}$ can be identified with the equivalence relation on the power set of κ in which two sets A and B are equivalent if their symmetric difference restricted to S is non-stationary. This can be done by identifying a set $A \subset \kappa$ with its characteristic function.

For every regular cardinal $\mu < \kappa$, we denote $\{\alpha < \kappa \mid cf(\alpha) = \mu\}$ by S_μ^κ . A set C is μ -club if it is unbounded and closed under μ -limits. For brevity, when $S = S_\mu^\kappa$, we will denote $E_{S_\mu^\kappa}^{\lambda, \kappa}$ by $E_{\mu\text{-club}}^{\lambda, \kappa}$. Note that $(f, g) \in E_{\mu\text{-club}}^{\lambda, \kappa}$ if and only if the set $\{\alpha < \kappa \mid f(\alpha) \neq g(\alpha)\}$ contains a μ -club.

For a Mahlo cardinal κ , the set $\text{reg}(\kappa) = \{\alpha < \kappa \mid \alpha \text{ a regular cardinal}\}$ is stationary. We will denote the equivalence relation $E_{\text{reg}(\kappa)}^{\lambda, \kappa}$ by $E_{\text{reg}}^{\lambda, \kappa}$.

Given an equivalence relation E on $X \in \{\kappa^\kappa, 2^\kappa\}$, we can define the λ -product relation of E for any $0 < \lambda < \kappa$. The λ -product relation $\Pi_\lambda E$ is the relation defined on $X^\lambda \times X^\lambda$ by $\eta \Pi_\lambda E \xi$ if $\eta_\gamma E \xi_\gamma$ holds for every $\gamma < \lambda$, where $\eta = (\eta_\gamma)_{\gamma < \lambda}$ and $\xi = (\xi_\gamma)_{\gamma < \lambda}$. We endow the space X^λ , $X \in \{\kappa^\kappa, 2^\kappa\}$, with the box topology generated by the basic open sets:

$$\{\Pi_{\alpha < \lambda} \mathcal{O}_\alpha \mid \forall \alpha < \lambda (\mathcal{O}_\alpha \text{ is an open set in } X)\}.$$

One of the motivations to study Borel reducibility in generalized Baire spaces is the connection with model theory. This connection consists in the possibility to study the Borel reducibility of the isomorphism relation of theories by coding structures with universe κ via elements of κ^κ . We may fix this coding, relative to a given countable relational vocabulary $\mathcal{L} = \{P_n \mid n < \omega\}$, as in the following definition.

Definition 1.4 Fix a bijection $\pi: \kappa^{<\omega} \rightarrow \kappa$. For every $\eta \in \kappa^\kappa$ define the \mathcal{L} -structure \mathcal{A}_η with universe κ as follows: For every relation P_m with arity n , every tuple (a_1, a_2, \dots, a_n) in κ^n satisfies

$$(a_1, a_2, \dots, a_n) \in P_m^{\mathcal{A}_\eta} \iff \eta(\pi(m, a_1, a_2, \dots, a_n)) \geq 1.$$

When we describe a complete theory T in a vocabulary $\mathcal{L}' \subseteq \mathcal{L}$, we think of it as a complete \mathcal{L} -theory extending $T \cup \{\forall \bar{x} \neg P_n(\bar{x}) \mid P_n \in \mathcal{L} \setminus \mathcal{L}'\}$.

Definition 1.5 (The isomorphism relation) Assume T is a complete first order theory in a countable vocabulary. We define \cong_T as the relation

$$\{(\eta, \xi) \in \kappa^\kappa \times \kappa^\kappa \mid (\mathcal{A}_\eta \models T, \mathcal{A}_\xi \models T, \mathcal{A}_\eta \cong \mathcal{A}_\xi) \text{ or } (\mathcal{A}_\eta \not\models T, \mathcal{A}_\xi \not\models T)\}.$$

In the second section we will study the reducibility between different cofinalities, and in the last section we will study the reducibility of $E_{\text{reg}}^{\kappa, \kappa}$ and $E_{\text{reg}}^{2, \kappa}$. Here is the list of the main results in this article:

- (Theorem 2.11) Suppose κ is a $\Pi_1^{\lambda^+}$ -indescribable cardinal for some $\lambda < \kappa$ and $V = L$. Then there is a forcing extension where κ is collapsed to become λ^{++} and $E_{\lambda\text{-club}}^{\lambda^{++}, \lambda^{++}} \leq_c E_{\lambda^+\text{-club}}^{2, \lambda^{++}}$.
- (Corollary 2.14) Let $\kappa_2 < \kappa_3 < \dots < \kappa_n < \dots$ be a sequence of supercompact cardinals. There is a generic extension $V[G]$ in which $\kappa_n = \aleph_n$ for all $n \geq 2$ and such that: $E_{\omega\text{-club}}^{\omega_2, \omega_2} \leq_c E_{\omega_1\text{-club}}^{\omega_2, \omega_2}$, and for every $n > 2$ and every $0 \leq k \leq n - 3$, $E_{\omega_k\text{-club}}^{\omega_n, \omega_n} \leq_c E_{\omega_{n-1}\text{-club}}^{\omega_n, \omega_n}$.

This corollary follows from [[8], Theorem 1.3] and gives a model (different from L or the one in Theorem 1.2) in which reducibility between different cofinalities holds.

- (Theorem 3.5) Suppose $S = S_\lambda^\kappa$ for some regular cardinal $\lambda < \kappa$, or $S = \text{reg}(\kappa)$ and κ weakly compact. If κ has the weakly compact diamond (Definition 3.2), then $E_S^{\kappa, \kappa} \leq_c E_{\text{reg}}^{2, \kappa}$.
- (Corollary 3.6) Suppose $V = L$ and κ is weakly compact. Then $E_{\text{reg}}^{2, \kappa}$ is Σ_1^1 -complete.
- (Corollary 3.7) Suppose κ is a weakly ineffable cardinal. Then $E_{\text{reg}}^{\kappa, \kappa} \leq_c E_{\text{reg}}^{2, \kappa}$.
- (Theorem 3.8) If κ is a Π_2^1 -indescribable cardinal, then $E_{\text{reg}}^{\kappa, \kappa}$ is Σ_1^1 -complete.
- (Corollary 3.9) Suppose κ is an ineffable cardinal (or weakly ineffable and Π_2^1 -indescribable). Then $E_{\text{reg}}^{2, \kappa}$ is Σ_1^1 -complete.
- (Theorem 3.10) Let DLO be the theory of dense linear orderings without end points. If κ is a Π_2^1 -indescribable cardinal, then \cong_{DLO} is Σ_1^1 -complete.

2 Reducibility between different cofinalities

In [1] the authors studied the reducibility between the relations $E_{\mu\text{-club}}^{2, \kappa}$ and showed in particular the consistency of $E_{\lambda\text{-club}}^{2, \lambda^{++}} \leq_c E_{\lambda^+\text{-club}}^{2, \lambda^{++}}$. In this section we continue along these lines.

Definition 2.1 We say that a set $X \subset \kappa$ strongly reflects to a set $Y \subset \kappa$ if for all stationary $Z \subset X$ there exist stationary many $\alpha \in Y$ with $Z \cap \alpha$ stationary in α .

In [1, Theorem 55] it is proved that: If κ is a weakly compact cardinal, then S_λ^κ strongly reflects to $\text{reg}(\kappa)$, for any regular cardinal $\lambda < \kappa$. This result can be generalized to Π_1^λ -indescribable cardinals:

Definition 2.2 A cardinal κ is Π_1^λ -indescribable (for $\lambda < \kappa$) if whenever $A \subset V_\kappa$ and σ is a Π_1 sentence such that

$$(V_{\kappa+\lambda}, \in, A, (V_{\kappa+\xi} \mid \xi < \lambda)) \models \sigma,$$

then for some $\alpha < \kappa$,

$$(V_{\alpha+\lambda}, \in, A \cap V_\alpha, (V_{\alpha+\xi} : \xi < \lambda)) \models \sigma$$

Note that, in Definition 2.2, the existence of some $\alpha < \kappa$ at which the required reflection is effected is equivalent to the existence of stationary many such $\alpha < \kappa$.

Lemma 2.3 *Suppose κ is a Π_1^λ -indescribable cardinal. There are λ many disjoint stationary subsets of κ , $\langle S_\gamma \rangle_{\gamma < \lambda}$, such that for every $\gamma < \lambda$, $S_\gamma \subseteq \text{reg}(\kappa)$ and κ strongly reflects to S_γ .*

Proof Let S_β^* denote the set of all the Π_1^β -indescribable cardinals below κ . Since “ κ is Π_1^β -indescribable” is a Π_1 property of the structure $(V_{\kappa+\lambda}, \in, (V_{\kappa+\xi} \mid \xi < \lambda))$, the set S_β^* is stationary for every $\beta < \lambda$.

Let us show that for every stationary set $X \subseteq \kappa$,

$$B = \{\alpha \in S_\beta^* \mid X \cap \alpha \text{ is stationary in } \alpha\}$$

is stationary. Let C be a club in κ . The sentence

$$(C \text{ is unbounded in } \kappa) \wedge (X \text{ is stationary in } \kappa) \wedge (\kappa \text{ is } \Pi_1^\beta\text{-indescribable})$$

is a Π_1 property of the structure $(V_{\kappa+\lambda}, \in, X, C, (V_{\kappa+\xi} \mid \xi < \lambda))$. By reflection, there is $\gamma < \kappa$ such that $C \cap \gamma$ is unbounded in γ , and hence $\gamma \in C$, $X \cap \gamma$ is stationary in γ , and γ is Π_1^β -indescribable. We conclude that $C \cap B \neq \emptyset$.

Let us denote $S_\beta^* \setminus S_{\beta+1}^*$ by S_β . Let us show that for every stationary set $X \subseteq \kappa$,

$$\{\alpha \in S_\beta \mid X \cap \alpha \text{ is stationary in } \alpha\}$$

is stationary. Let C be a club in κ . Since $\{\alpha \in S_\beta^* \mid X \cap \alpha \text{ is stationary in } \alpha\}$ is stationary, we can pick $\gamma \in C \cap \{\alpha \in S_\beta^* \mid X \cap \alpha \text{ is stationary in } \alpha\}$ such that γ is minimal.

Claim 2.3.1 *γ is not $\Pi_1^{\beta+1}$ -indescribable.*

Proof Suppose, towards a contradiction, that γ is $\Pi_1^{\beta+1}$ -indescribable. The sentence

$$(C \cap \gamma \text{ is unbounded in } \gamma) \wedge (X \cap \gamma \text{ is stationary in } \gamma) \wedge (\gamma \text{ is } \Pi_1^\beta\text{-indescribable})$$

is a Π_1 property of the structure $(V_{\gamma+\beta+1}, \in, X \cap \gamma, C \cap \gamma, (V_{\gamma+\xi} \mid \xi < \beta + 1))$. By reflection, there is $\gamma' < \gamma$ such that $C \cap \gamma'$ is unbounded in γ' , $X \cap \gamma'$ is stationary in γ' , and γ' is Π_1^β -indescribable. This contradicts the minimality of γ . \square

We conclude that S_β is stationary and $\{\alpha \in S_\beta \mid X \cap \alpha \text{ is stationary in } \alpha\}$ is stationary, for every $\beta < \lambda$. \square

The notion of \diamond -reflection was introduced in [1] in order to find reductions between equivalence relations modulo non-stationary ideals (see below).

Definition 2.4 (\diamond -reflection) *Let X, Y be subsets of κ and suppose Y consists of ordinals of uncountable cofinality. We say that X \diamond -reflects to Y if there exists a sequence $\langle D_\alpha \rangle_{\alpha \in Y}$ such that:*

- $D_\alpha \subset \alpha$ is stationary in α for all $\alpha \in Y$.
- if $Z \subset X$ is stationary, then $\{\alpha \in Y \mid D_\alpha = Z \cap \alpha\}$ is stationary.

Theorem 2.5 ([1], Theorem 59) *Suppose $V = L$ and that $X \subseteq \kappa$ and $Y \subseteq \text{reg}(\kappa)$. If X strongly reflects to Y , then X \diamond -reflects to Y .*

Theorem 2.6 ([1], Theorem 58) *If X \diamond -reflects to Y , then $E_X^{2, \kappa} \leq_c E_Y^{2, \kappa}$.*

\diamond -reflection also implies some reductions for the relations $E_{\mu\text{-club}}^{\kappa,\kappa}$ on the space κ^κ . To show this, we first need to introduce some definitions.

Definition 2.7 For every $\alpha < \kappa$ with $\gamma < cf(\alpha)$ define $E_{\gamma\text{-club}}^{\kappa,\kappa} \upharpoonright \alpha$ by:

$$E_{\gamma\text{-club}}^{\kappa,\kappa} \upharpoonright \alpha = \{(\eta, \xi) \in \kappa^\kappa \times \kappa^\kappa \mid \exists C \subseteq \alpha \text{ a } \gamma\text{-club}, \forall \beta \in C, \eta(\beta) = \xi(\beta)\}.$$

Proposition 2.8 Suppose $\gamma < \lambda < \kappa$ are regular cardinals. If S_γ^κ strongly reflects to S_λ^κ , then $E_{\gamma\text{-club}}^{\kappa,\kappa} \leq_c E_{\lambda\text{-club}}^{\kappa,\kappa}$.

Proof Suppose that for every stationary set $S \subset S_\gamma^\kappa$ it holds that

$$\{\alpha \in S_\lambda^\kappa \mid S \cap \alpha \text{ is stationary in } \alpha\}$$

is a stationary set, and define $F: \kappa^\kappa \rightarrow \kappa^\kappa$ by

$$F(\eta)(\alpha) = \begin{cases} f_\alpha(\eta), & \text{if } cf(\alpha) = \lambda \\ 0, & \text{otherwise.} \end{cases}$$

where $f_\alpha(\eta)$ is a code in $\kappa \setminus \{0\}$ for the $(E_{\gamma\text{-club}}^{\kappa,\kappa} \upharpoonright \alpha)$ -equivalence class of η .

Let us prove that if $(\eta, \xi) \in E_{\gamma\text{-club}}^{\kappa,\kappa}$, then $(F(\eta), F(\xi)) \in E_{\lambda\text{-club}}^{\kappa,\kappa}$. Suppose $(\eta, \xi) \in E_{\gamma\text{-club}}^{\kappa,\kappa}$. There is a γ -club where η and ξ coincide and so there is a club C such that for all $\alpha \in C \cap S_\lambda^\kappa$ the functions η and ξ are $(E_{\gamma\text{-club}}^{\kappa,\kappa} \upharpoonright \alpha)$ -equivalent. Thus, by the definition of F , for all $\alpha \in C \cap S_\lambda^\kappa$, $F(\eta)(\alpha) = F(\xi)(\alpha)$. We conclude that $(F(\eta), F(\xi)) \in E_{\lambda\text{-club}}^{\kappa,\kappa}$.

Let us prove that if $(\eta, \xi) \notin E_{\gamma\text{-club}}^{\kappa,\kappa}$, then $(F(\eta), F(\xi)) \notin E_{\lambda\text{-club}}^{\kappa,\kappa}$. Suppose that $(\eta, \xi) \notin E_{\gamma\text{-club}}^{\kappa,\kappa}$. Then there is a stationary $S \subset S_\gamma^\kappa$ on which $\eta(\alpha) \neq \xi(\alpha)$. Since $A = \{\alpha \in S_\lambda^\kappa \mid S \cap \alpha \text{ is stationary in } \alpha\}$ is stationary and for all $\alpha \in A$, $f_\alpha(\eta) \neq f_\alpha(\xi)$, we conclude that $(F(\eta), F(\xi)) \notin E_{\lambda\text{-club}}^{\kappa,\kappa}$. \square

Corollary 2.9 Suppose $\gamma < \lambda < \kappa$ are regular cardinals. If S_γ^κ \diamond -reflects to S_λ^κ , then

1. $E_{\gamma\text{-club}}^{2,\kappa} \leq_c E_{\lambda\text{-club}}^{2,\kappa}$.
2. $E_{\gamma\text{-club}}^{\kappa,\kappa} \leq_c E_{\lambda\text{-club}}^{\kappa,\kappa}$.

Proof 1. Follows from Theorem 2.6.

2. By the definition of \diamond -reflection, S_γ^κ \diamond -reflecting to S_λ^κ implies that for all $S \subseteq S_\gamma^\kappa$ the set $\{\alpha \in S_\lambda^\kappa \mid S \cap \alpha \text{ is stationary in } \alpha\}$ is a stationary set. The result follows from Proposition 2.8. \square

In [1], the consistency of $S_\lambda^{\lambda^{++}}$ \diamond -reflecting to $S_{\lambda^+}^{\lambda^{++}}$ was shown. This gives a model in which $E_{\lambda\text{-club}}^{2,\kappa} \leq_c E_{\lambda^+\text{-club}}^{2,\kappa}$ and $E_{\lambda\text{-club}}^{\lambda^{++},\lambda^{++}} \leq_c E_{\lambda^+\text{-club}}^{\lambda^{++},\lambda^{++}}$.

Theorem 2.10 ([1], Theorem 55) Suppose that κ is a weakly compact cardinal and $V = L$. Then:

1. $E_{\lambda\text{-club}}^{2,\kappa} \leq_c E_{\text{reg}}^{2,\kappa}$ holds for all regular $\lambda < \kappa$.
2. For every regular $\lambda < \kappa$ there is a forcing extension where κ is collapsed to become λ^{++} and $E_{\lambda\text{-club}}^{2,\lambda^{++}} \leq_c E_{\lambda^+\text{-club}}^{2,\lambda^{++}}$.

The proof of this theorem can be generalised using Lemma 2.3 to show the consistency of $E_{\lambda\text{-club}}^{\lambda^{++}, \lambda^{++}} \leq_c E_{\lambda^+-\text{club}}^{2, \lambda^{++}}$:

Theorem 2.11 *Suppose κ is a $\Pi_1^{\lambda^+}$ -indescribable cardinal and that $V = L$. Then there is a forcing extension where κ is collapsed to become λ^{++} and $E_{\lambda\text{-club}}^{\lambda^{++}, \lambda^{++}} \leq_c E_{\lambda^+-\text{club}}^{2, \lambda^{++}}$.*

Proof Let us collapse κ to λ^{++} with the Levy collapse

$$\mathbb{P} = \{f: \text{reg}(\kappa) \rightarrow \kappa^{<\lambda^+} \mid \text{rang}(f(\mu)) \subset \mu, |\{\mu \mid f(\mu) \neq \emptyset\}| \leq \lambda\}$$

where $f \geq g$ if and only if $f(\mu) \subseteq g(\mu)$ for all $\mu \in \text{reg}(\kappa)$. Let us define \mathbb{P}_μ and \mathbb{P}^μ for all μ by: $\mathbb{P}_\mu = \{f \in \mathbb{P} \mid \text{sprt}(f) \subset \mu\}$ and $\mathbb{P}^\mu = \{f \in \mathbb{P} \mid \text{sprt}(f) \subset \kappa \setminus \mu\}$. It is known that all regular $\lambda < \mu \leq \kappa$ satisfy:

- (i) if $\mu > \lambda^+$, then \mathbb{P}_μ has the μ -c.c.,
- (ii) \mathbb{P}_μ and \mathbb{P}^μ are $<\lambda^+$ -closed,
- (iii) $\mathbb{P} = \mathbb{P}_\kappa \Vdash \lambda^{++} = \check{\kappa}$,
- (vi) if $\mu < \kappa$, then $\mathbb{P} \Vdash c f(\check{\mu}) = \lambda^+$,
- (v) if $p \in \mathbb{P}$, σ a name, and $p \Vdash \text{“}\sigma \text{ is a club in } \lambda^{++}\text{”}$, then there is a club $E \subset \kappa$ such that $p \Vdash \check{E} \subset \sigma$.

Claim 2.11.1 *There is a sequence $\langle S_\gamma \rangle_{\gamma < \lambda^+}$ of disjoint stationary subsets of $S_{\lambda^+}^{\lambda^{++}}$ such that in $V[G]$ $S_{\lambda^+}^{\lambda^{++}}$ \diamond -reflects to S_γ for every $\gamma < \lambda^+$.*

Proof Let G be a \mathbb{P} -generic over V , and define $G_\mu = G \cap \mathbb{P}_\mu$ and $G^\mu = G \cap \mathbb{P}^\mu$. So G_μ is \mathbb{P}_μ -generic over V , G^μ is \mathbb{P}^μ -generic over $V[G_\mu]$, and $V[G] = V[G_\mu][G^\mu]$. Let S_β^* denote the set of all Π_1^β -indescribable cardinals below κ and $S_\beta = S_\beta^* \setminus S_{\beta+1}^*$. We will show that $S_{\lambda^+}^{\lambda^{++}}$ \diamond -reflects to S_β^V for all $\beta < \lambda^+$. Let us fix $\beta < \lambda^+$ and denote by Y the set S_β^V . By Lemma 2.3 we know that S_β^V is stationary and by (v), it remains stationary in $V[G]$. By (i) we know that there are no antichains of length μ in \mathbb{P}_μ , and since $|\mathbb{P}_\mu| = \mu$ we conclude that there are at most μ antichains. On the other hand, there are μ^+ many subsets of μ . Hence, there is a bijection

$$h_\mu: \mu^+ \rightarrow \{\sigma \mid \sigma \text{ is a nice } \mathbb{P}_\mu \text{ name for a subset of } \mu\}$$

for each $\mu \in \text{reg}(\kappa)$ such that $\mu > \lambda^+$, where a nice \mathbb{P}_μ name for a subset of $\check{\mu}$ is of the form $\bigcup \{\check{\alpha} \times A_\alpha \mid \alpha \in B\}$ with $B \subset \check{\mu}$ and A_α an antichain in \mathbb{P}_μ . Notice that the nice \mathbb{P}_μ names for subsets of $\check{\mu}$ are subsets of V_μ . Let us define

$$D_\mu = \begin{cases} [h_\mu((\bigcup G)(\mu^+))(0)]_G & \text{if this set is stationary} \\ \mu & \text{otherwise.} \end{cases}$$

We will show that $\langle D_\mu \rangle_{\mu \in Y}$ is the needed \diamond -sequence in $V[G]$.

Suppose, towards a contradiction, that there are a stationary set $S \subset S_{\lambda^+}^{\lambda^{++}}$ and a club $C \subset \lambda^{++}$ (in $V[G]$) such that for all $\alpha \in C \cap Y$, $D_\alpha \neq S \cap \alpha$. By (v) there is a club $C_0 \subset C$ such that $C_0 \in V$. Let \check{S} be a nice name for S and p a condition such that p forces that \check{S} is stationary. We will show that

$$H = \{q < p \mid q \Vdash D_\mu = \check{S} \cap \check{\mu} \text{ for some } \mu \in C_0\}$$

is dense below p , which is a contradiction. Let us slightly redefine \mathbb{P} .

Let $\mathbb{P}^* = \{q \mid \exists r \in \mathbb{P} (r \upharpoonright \text{sprt}(r) = q)\}$. Clearly $\mathbb{P} \cong \mathbb{P}^*$, $\mathbb{P}^* \subseteq V_\kappa$, and $\mathbb{P}_\mu^* = \mathbb{P}^* \cap V_\mu$, where $\mathbb{P}_\mu^* = \{q \mid \exists r \in \mathbb{P}_\mu (r \upharpoonright \text{sprt}(r) = q)\}$. It can be verified that the properties mentioned above also hold for \mathbb{P}_μ^* . From now on denote \mathbb{P}_μ^* by \mathbb{P}_μ . Let r be a condition stronger than p and

$$R = (\mathbb{P} \times \{0\}) \cup (\dot{S} \times \{1\}) \cup (C_0 \times \{2\} \cup (\{r\} \times \{3\})).$$

Let $\forall A\varphi$ be the formula:

If A is closed and unbounded and $t < r$ are arbitrary, then there exists $q < r$ and $\alpha \in A$ such that $q \Vdash_{\mathbb{P}} \check{\alpha} \in \dot{S}$.

Clearly, $\forall A\varphi$ says $r \Vdash \dot{S}$ is stationary. By (v) it is enough to quantify over club sets in V . Notice that $t < r$, $q < t$, A is a club, and $\alpha \in A$ are first order expressible using R as a parameter. The definition of $\check{\alpha}$ is recursive in α :

$$\check{\alpha} = \{(\check{\gamma}, 1_{\mathbb{P}}) \mid \gamma < \alpha\}$$

and it is absolute for V_κ . Then $q \Vdash_{\mathbb{P}} \check{\alpha} \in \dot{S}$ is equivalent to saying that for each $q' < q$ there exists $q'' < q'$ with $(\check{\alpha}, q'') \in \dot{S}$, and this is first order expressible using R as a parameter. Therefore $\forall A\varphi$ is a Π_1 property of the structure (V_κ, \in, R) , even more

$$(\forall A\varphi) \wedge (\kappa \text{ is } \Pi_1^\beta\text{-indescribable})$$

is a Π_1 property of the structure $(V_{\kappa+\lambda^+}, \in, R, (V_{\mu+\xi} \mid \xi < \lambda^+))$. By reflection, there is $\mu < \kappa$ Π_1^β -indescribable, such that $\mu \in C_0$, $r \in \mathbb{P}_\mu$, and

$$(V_{\mu+\lambda^+}, \in, R, (V_{\mu+\xi} \mid \xi < \lambda^+)) \models \forall A\varphi.$$

In the same way as in Claim 2.3.1, we can show that there is there is $\mu < \kappa$ Π_1^β -indescribable that is not $\Pi_1^{\beta+1}$ -indescribable, i.e. $(\check{\mu}_G \in Y)^{V[G]}$, such that $\mu \in C_0$, $r \in \mathbb{P}_\mu$, and $(V_{\mu+\lambda^+}, \in, R, (V_{\mu+\xi} \mid \xi < \lambda^+)) \models \forall A\varphi$. Notice that $\alpha \in S \cap \mu$ implies that $(\check{\alpha}, \check{q}) \in \dot{S}$ for some $q \in \mathbb{P}_\mu$. Let $\dot{S}_\mu = \dot{S} \cap V_\mu$, thus $r \Vdash_{\mathbb{P}_\mu} (\dot{S}_\mu \text{ is stationary})$. Let us define q as follows: $\text{dom}(q) = \text{dom}(r) \cup \{\mu^+\}$, $q \upharpoonright \mu = r \upharpoonright \mu$ and $q(\mu^+) = f$, $\text{dom}(f) = \{0\}$, and $f(0) = h_\mu^{-1}(\dot{S}_\mu)$. Since \mathbb{P}^μ is $<\lambda^+$ -closed and does not kill stationary subsets of $S_\lambda^{\lambda^+}$, $(\dot{S}_\mu)_{G_\mu}$ is stationary in $V[G]$, and by the way we chose μ , $(\dot{S}_\mu)_{G_\mu} = (\dot{S}_\mu)_G$. Therefore $q \Vdash_{\mathbb{P}} (\dot{S}_\mu \text{ is stationary})$, and by the definition of D_μ (in $V[G]$) we conclude that $q \Vdash_{\mathbb{P}} \dot{S}_\mu = D_\mu$. Finally, by the way we chose μ , we get that $(\dot{S}_\mu)_G = S \cap \mu$. We conclude that H is dense below p , a contradiction. \square

From now on in this proof, we will work in $V[G]$. In particular, κ will be λ^{++} .

Claim 2.11.2 $E_{\lambda\text{-club}}^{\kappa,\kappa} \leq_c \Pi_{\lambda^+} E_{\lambda\text{-club}}^{2,\kappa}$.

Proof Let H be a bijection from κ to 2^{λ^+} . Define $\mathcal{F}: \kappa^\kappa \rightarrow (2^\kappa)^{\lambda^+}$ by $\mathcal{F}(f) = (f_\gamma)_{\gamma < \lambda^+}$, where $f_\gamma(\alpha) = H(f(\alpha))(\gamma)$ for every $\gamma < \lambda^+$ and $\alpha < \kappa$. Let us show that \mathcal{F} is a reduction of $E_{\lambda\text{-club}}^{\kappa,\kappa}$ to $\Pi_{\lambda^+} E_{\lambda\text{-club}}^{2,\kappa}$.

Clearly $f(\alpha) = g(\alpha)$ implies $H(f(\alpha)) = H(g(\alpha))$ and $f_\gamma(\alpha) = g_\gamma(\alpha)$ for every $\gamma < \lambda^+$. Therefore, $f E_{\lambda\text{-club}}^{\kappa,\kappa} g$ implies that for all $\gamma < \lambda^+$, $f_\gamma E_{\lambda\text{-club}}^{2,\kappa} g_\gamma$ holds. So $f \Pi_{\lambda^+} E_{\lambda\text{-club}}^{2,\kappa} g$.

Suppose that for every $\gamma < \lambda^+$ there is C_γ , a λ -club, such that $f_\gamma(\alpha) = g_\gamma(\alpha)$ holds for every $\alpha \in C_\gamma$. Since the intersection of less than κ λ -club sets is a λ -club set, there is a λ -club C on which the functions f_γ and g_γ coincide for every $\gamma < \lambda^+$. Therefore $H(f(\alpha))(\gamma) = H(g(\alpha))(\gamma)$ holds for every $\gamma < \lambda^+$ and every $\alpha \in C$, so

$H(f(\alpha)) = H(g(\alpha))$ for every $\alpha \in C$. Since H is a bijection, we can conclude that $f(\alpha) = g(\alpha)$ for every $\alpha \in C$, and hence $f \mathop{E}_{\lambda\text{-club}}^{\kappa,\kappa} g$. \square

By Claim 2.11.1, there is a sequence $\langle S_\gamma \rangle_{\gamma < \lambda^+}$ of disjoint stationary subsets of $S_{\lambda^+}^\kappa$ such that $S_\lambda^\kappa \diamond$ -reflects to S_γ for all $\gamma < \lambda^+$. Let $\langle D_\alpha^\gamma \rangle_{\alpha \in S_\gamma}$ be a sequence that witnesses that $S_\lambda^\kappa \diamond$ -reflects to S_γ .

For every $\eta \in \kappa^\kappa$ define $F(\eta)$ by:

$$F(\eta)(\alpha) = \begin{cases} 1 & \text{if there is } \gamma < \lambda^+ \text{ with } \alpha \in S_\gamma \text{ and } \mathcal{F}(\eta)_\gamma^{-1}[1] \cap D_\alpha^\gamma \text{ stationary in } \alpha \\ 0 & \text{otherwise} \end{cases}$$

where $(\mathcal{F}(\eta)_\gamma)_{\gamma < \lambda^+} = \mathcal{F}(\eta)$ and where \mathcal{F} is the reduction given by Claim 2.11.2.

Suppose η, ξ are not $E_{\lambda\text{-club}}^{\kappa,\kappa}$ -equivalent. By Claim 2.11.2 there exists $\gamma < \lambda^+$ such that $\mathcal{F}(\eta)_\gamma^{-1}[1] \Delta \mathcal{F}(\xi)_\gamma^{-1}[1]$ is stationary. Therefore, either $\mathcal{F}(\eta)_\gamma^{-1}[1] \setminus \mathcal{F}(\xi)_\gamma^{-1}[1]$ or $\mathcal{F}(\xi)_\gamma^{-1}[1] \setminus \mathcal{F}(\eta)_\gamma^{-1}[1]$ is stationary. Without loss of generality, let us assume that $\mathcal{F}(\eta)_\gamma^{-1}[1] \setminus \mathcal{F}(\xi)_\gamma^{-1}[1]$ is stationary. Since $S_\lambda^\kappa \diamond$ -reflects to S_γ ,

$$A = \{ \alpha \in S_\gamma \mid (\mathcal{F}(\eta)_\gamma^{-1}[1] \setminus \mathcal{F}(\xi)_\gamma^{-1}[1]) \cap \alpha = D_\alpha^\gamma \}$$

is stationary and D_α^γ is stationary in α , and therefore $A \subseteq F(\eta)^{-1}[1]$. On the other hand, for every α in A we have $\mathcal{F}(\xi)_\gamma^{-1}[1] \cap D_\alpha^\gamma = \emptyset$, so $A \cap F(\xi)^{-1}[1] = \emptyset$ and we conclude that $A \subseteq F(\eta)^{-1}[1] \Delta F(\xi)^{-1}[1]$. Therefore $F(\eta)^{-1}[1] \Delta F(\xi)^{-1}[1]$ is stationary, and $F(\eta)$ and $F(\xi)$ are not $E_{\lambda^+\text{-club}}^{2,\lambda^+}$ -equivalent.

Suppose $F(\eta)$ and $F(\xi)$ are not $E_{\lambda^+\text{-club}}^{2,\lambda^+}$ -equivalent, so $F(\eta)^{-1}[1] \Delta F(\xi)^{-1}[1]$ is stationary. Since $\lambda^+ < \kappa$, by Fodor's lemma we know that there exists $\gamma < \lambda^+$ such that $\{ \alpha \in S_\gamma \mid F(\eta)(\alpha) \neq F(\xi)(\alpha) \}$ is stationary. Hence, the symmetric difference of the sets $\{ \alpha \in S_\gamma \mid \mathcal{F}(\eta)_\gamma^{-1}[1] \cap D_\alpha^\gamma \text{ is stationary in } \alpha \}$ and $\{ \alpha \in S_\gamma \mid \mathcal{F}(\xi)_\gamma^{-1}[1] \cap D_\alpha^\gamma \text{ is stationary in } \alpha \}$ is stationary. For simplicity, let us denote by A_η the set $\{ \alpha \in S_\gamma \mid \mathcal{F}(\eta)_\gamma^{-1}[1] \cap D_\alpha^\gamma \text{ is stationary in } \alpha \}$ and A_ξ the set $\{ \alpha \in S_\gamma \mid \mathcal{F}(\xi)_\gamma^{-1}[1] \cap D_\alpha^\gamma \text{ is stationary in } \alpha \}$. Therefore, either $A_\eta \setminus A_\xi$ or $A_\xi \setminus A_\eta$ is stationary. Without loss of generality we can assume that $A_\eta \setminus A_\xi$ is stationary. Hence, $\bigcup_{\alpha \in A_\eta \setminus A_\xi} (\mathcal{F}(\eta)_\gamma^{-1}[1] \cap D_\alpha^\gamma) \setminus \mathcal{F}(\xi)_\gamma^{-1}[1]$ is stationary and is contained in $\mathcal{F}(\eta)_\gamma^{-1}[1] \Delta \mathcal{F}(\xi)_\gamma^{-1}[1]$. By Claim 2.11.2 we conclude that η and ξ are not $E_{\lambda\text{-club}}^{\kappa,\kappa}$ -equivalent. \square

Notice that Theorem 2.11 implies the consistency of

$$E_{\lambda\text{-club}}^{2,\lambda^+} \leq_c E_{\lambda\text{-club}}^{\lambda^+,\lambda^+} \leq_c E_{\lambda^+\text{-club}}^{2,\lambda^+} \leq_c E_{\lambda^+\text{-club}}^{\lambda^+,\lambda^+}.$$

In particular, for $\lambda = \omega$ we get the expression

$$E_{\omega\text{-club}}^{2,\omega_2} \leq_c E_{\omega\text{-club}}^{\omega_2,\omega_2} \leq_c E_{\omega_1\text{-club}}^{2,\omega_2} \leq_c E_{\omega_1\text{-club}}^{\omega_2,\omega_2}.$$

Question 2.12 *Is it consistent that*

$$E_{\gamma\text{-club}}^{2,\kappa} \leq_c E_{\gamma\text{-club}}^{\kappa,\kappa} \leq_c E_{\lambda\text{-club}}^{2,\kappa}$$

holds for all $\gamma, \lambda < \kappa$ and $\gamma < \lambda$?

We will finish this section by showing that the reduction $E_{\omega\text{-club}}^{\omega_2,\omega_2} \leq_c E_{\omega_1\text{-club}}^{\omega_2,\omega_2}$ can be obtained using other reflection principles. Specifically, full reflection implies this reduction. For stationary subsets S and A of κ , we say that S *reflects fully* in A if

the set $\{\alpha \in A \mid S \cap \alpha \text{ is non-stationary in } \alpha\}$ is non-stationary. Notice that if $S \subset S_\gamma^\kappa$ reflects fully in S_λ^κ , then the set $\{\alpha \in S_\lambda^\kappa \mid S \cap \alpha \text{ is stationary in } \alpha\}$ is a stationary set.

Theorem 2.13 ([8], Theorem 1.3) *Let $\kappa_2 < \kappa_3 < \dots < \kappa_n < \dots$ be a sequence of supercompact cardinals. There is a generic extension $V[G]$ in which $\kappa_n = \aleph_n$ for all $n \geq 2$ and such that:*

1. *Every stationary set $S \subset S_\omega^{\omega_2}$ reflects fully in $S_{\omega_1}^{\omega_2}$.*
2. *For every $2 < n$ and every $0 \leq k \leq n-3$, every stationary set $S \subset S_{\omega_k}^{\omega_n}$ reflects fully in $S_{\omega_{n-1}}^{\omega_n}$.*

In the generic extension of 2.13 it holds that $\omega_i^{<\omega_i} = \omega_i$ for all $i < \omega$ (see [[8], Theorem 1.3]).

Corollary 2.14 *Let $\kappa_2 < \kappa_3 < \dots < \kappa_n < \dots$ be a sequence of supercompact cardinals. There is a generic extension $V[G]$ in which $\kappa_n = \aleph_n$ for all $n \geq 2$ and such that: $E_{\omega\text{-club}}^{\omega_2, \omega_2} \leq_c E_{\omega_1\text{-club}}^{\omega_2, \omega_2}$, and for every $n > 2$ and every $0 \leq k \leq n-3$, $E_{\omega_k\text{-club}}^{\omega_n, \omega_n} \leq_c E_{\omega_{n-1}\text{-club}}^{\omega_n, \omega_n}$.*

In [8] it was also proved that Theorem 2.13 (ii) is optimal, in the sense that it cannot be improved to include the case $k = n-2$ [8, Proposition 1.6]. The best possible reduction we can get using only full reflection is the one in Corollary 2.14. By a Σ_1^1 -completeness result, it is known that the following is consistent:

$$\forall k < n-1 (E_{\omega_k\text{-club}}^{\omega_n, \omega_n} \leq_c E_{\omega_{n-1}\text{-club}}^{\omega_n, \omega_n}),$$

see Theorem 3.1 below.

3 Σ_1^1 -completeness

An equivalence relation E on $X \in \{\kappa^\kappa, 2^\kappa\}$ is Σ_1^1 if E is the projection of a closed set in $X^2 \times \kappa^\kappa$ and it is Σ_1^1 -complete if every Σ_1^1 equivalence relation is Borel reducible to it. The study of Σ_1^1 and Σ_1^1 -complete equivalence relations is an important area of generalised descriptive set theory, because e.g. the isomorphism relation on classes of models is always Σ_1^1 . The same holds, in fact, in classical descriptive set theory, but the behaviour of Σ_1^1 complete relations there is different. For example, in the classical setting ($\kappa = \omega$) the isomorphism relation is never Σ_1^1 -complete, while in generalised descriptive set theory this is often the case (see for example [6, 1]).

Theorem 3.1 ([6], Theorem 7) *Suppose $V = L$ and $\kappa > \omega$. Then $E_{\mu\text{-club}}^{\kappa, \kappa}$ is Σ_1^1 -complete for every regular $\mu < \kappa$.*

We know that $E_{\lambda\text{-club}}^{\kappa, \kappa} \upharpoonright \alpha$ is an equivalence relation for every $\alpha < \kappa$ with $cf(\alpha) > \lambda$. Let us define the following relation:

$$(\eta, \xi) \in E_{\text{reg}}^{\kappa, \kappa} \upharpoonright \alpha \Leftrightarrow \{\beta \in \text{reg}(\alpha) \mid \eta(\beta) \neq \xi(\beta)\} \text{ is not stationary.}$$

It is easy to see that $E_{\text{reg}}^{\kappa, \kappa} \upharpoonright \alpha$ is an equivalence relation.

Definition 3.2 (Weakly compact diamond) *This notion was originally defined in [9]. Let $\kappa > \omega$ be a cardinal. The weakly compact ideal is generated by the sets of the form $\{\alpha < \kappa \mid \langle V_\alpha, \in, U \cap V_\alpha \rangle \models \neg \varphi\}$ where $U \subset V_\kappa$ and φ is a Π_1^1 -sentence such that $\langle V_\kappa, \in, U \rangle \models \varphi$. One can define a diamond principle with respect to this ideal (rather than the non-stationary ideal). A set $A \subset \kappa$ is said to be weakly compact, if it does not belong to the weakly compact ideal. Note that κ is weakly compact if and only if there exists $A \subset \kappa$ which is weakly compact, i.e. the weakly compact ideal is*

proper. For weakly compact $S \subset \kappa$, the S -weakly compact diamond, $\text{WC}_\kappa(S)$, is the statement that there exists a sequence $(A_\alpha)_{\alpha < \kappa}$ such that for every $A \subset S$ the set

$$\{\alpha < \kappa \mid A \cap \alpha = A_\alpha\}$$

is weakly compact. We denote $\text{WC}_\kappa = \text{WC}_\kappa(\kappa)$.

For a survey on weakly compact diamonds, see [5].

Fact 3.3 *The main facts that we will use are the following:*

- If κ is weakly compact and $V = L$, then WC_κ holds.
- If κ is weakly ineffable (same as almost ineffable), then WC_κ holds.

See [5] for proofs and references.

Lemma 3.4 *Let κ be a weakly compact cardinal. The weakly compact diamond WC_κ implies the following principle WC_κ^* . There exists a sequence $\langle f_\alpha \rangle_{\alpha \in \text{reg}(\kappa)}$ such that*

- $f_\alpha: \alpha \rightarrow \alpha$,
- for all $g \in \kappa^\kappa$ and stationary $Z \subset \kappa$ the set

$$\{\alpha \in \text{reg}(\kappa) \mid g \upharpoonright \alpha = f_\alpha \wedge \alpha \cap Z \text{ is stationary}\}$$

is stationary.

Proof For the sake of this proof we view functions $f: \alpha \rightarrow \alpha$ as subsets of $\alpha \times \alpha$.

Let $(A_\alpha)_{\alpha < \kappa}$ be the WC_κ -sequence and let $\pi: \kappa \times \kappa \rightarrow \kappa$ be a bijection. Let C_π be the set $\{\alpha < \kappa \mid \pi[\alpha \times \alpha] = \alpha\}$. It is standard to verify that C_π is a club. For all $\alpha \in \text{reg}(\kappa)$ let $f_\alpha = \pi^{-1}[A_\alpha]$ if $\alpha \in C_\pi$ and $\pi^{-1}[A_\alpha]$ is a function (i.e. for all $\beta < \alpha$ there exists exactly one γ such that $(\beta, \gamma) \in \pi^{-1}[A_\alpha]$) and otherwise set f_α to be arbitrary. Let us show that this sequence is as desired. Let $g \in \kappa^\kappa$ be a function and Z stationary. Let C_g be the set $\{\alpha < \kappa \mid g \upharpoonright \alpha \subset \alpha\}$ which is again a club. The set

$$\{\alpha < \kappa \mid \pi[g \upharpoonright \alpha] \cap \alpha = A_\alpha\}$$

is weakly compact and so is

$$\{\alpha \in C_g \cap C_\pi \mid \pi[g \upharpoonright \alpha] \cap \alpha = A_\alpha\}.$$

But since $\alpha \in C_\pi \cap C_g$, we have $\pi[g \upharpoonright \alpha] \cap \alpha = \pi[g \upharpoonright (\alpha \times \alpha)]$, so this set is equal to

$$\begin{aligned} S &= \{\alpha \in C_g \cap C_\pi \mid g \upharpoonright (\alpha \times \alpha) = \pi^{-1}[A_\alpha]\} \\ &= \{\alpha \in C_g \cap C_\pi \mid g \upharpoonright \alpha = f_\alpha\}. \end{aligned}$$

By the weak compactness of S , the stationarity of Z is reflected to a stationary subset $S' \subset S$, so $Z \cap \alpha$ is stationary for all $\alpha \in S'$. \square

Theorem 3.5 *Suppose $S = S_\lambda^\kappa$ for some λ regular cardinal, or $S = \text{reg}(\kappa)$ and κ is a weakly compact cardinal. If κ has the weakly compact diamond, then $E_S^{\kappa, \kappa} \leq_c E_{\text{reg}}^{2, \kappa}$.*

Proof Let $\langle f_\alpha \rangle_{\alpha < \kappa}$ be a sequence that witnesses WC_κ^* of Lemma 3.4. Let $g_\alpha: \kappa \rightarrow \kappa$ be the function defined by $g_\alpha \upharpoonright \alpha = f_\alpha$ and $g_\alpha(\beta) = 0$ for all $\beta \geq \alpha$. Let us define $F: \kappa^\kappa \rightarrow 2^\kappa$ by

$$F(\eta)(\alpha) = \begin{cases} 1 & \text{if } \alpha \in \text{reg}(\kappa), E_S^{\kappa, \kappa} \upharpoonright \alpha \text{ is an equivalence relation, and } (\eta, g_\alpha) \in E_S^{\kappa, \kappa} \upharpoonright \alpha \\ 0 & \text{otherwise.} \end{cases}$$

(Recall Definition 2.7 for $E_S^{\kappa,\kappa} \upharpoonright \alpha$.) Let us prove that if $(\eta, \xi) \in E_S^{\kappa,\kappa}$, then $(F(\eta), F(\xi)) \in E_{\text{reg}}^{2,\kappa}$. Suppose $(\eta, \xi) \in E_S^{\kappa,\kappa}$. Note that $F(\eta)(\alpha) = F(\xi)(\alpha) = 0$ for all $\alpha \notin \text{reg}(\kappa)$, so it is sufficient to show that the set

$$\{\alpha \in \text{reg}(\kappa) \mid F(\eta)(\alpha) \neq F(\xi)(\alpha)\}$$

is non-stationary. Now, there is a club D such that $D \cap \{\alpha \in S \mid \eta(\alpha) \neq \xi(\alpha)\}$ is non-stationary. So, letting C be the club of the limit points of D , it holds that for all $\alpha \in C \cap \text{reg}(\kappa)$, the functions η and ξ are $E_S^{\kappa,\kappa} \upharpoonright \alpha$ -equivalent. Thus, by the definition of F , at the points of the set $C \cap \text{reg}(\kappa)$ the functions $F(\eta)$ and $F(\xi)$ will get the same value.

Now let us prove that if $(\eta, \xi) \notin E_S^{\kappa,\kappa}$, then $(F(\eta), F(\xi)) \notin E_{\text{reg}}^{2,\kappa}$. Suppose that $(\eta, \xi) \notin E_S^{\kappa,\kappa}$. Then there is a stationary $Z \subset S$ on which $\eta(\alpha) \neq \xi(\alpha)$. By Lemma 3.4, there is a stationary set $A \subseteq \text{reg}(\kappa)$ such that for all $\alpha \in A$ we have that $Z \cap \alpha$ is stationary and $\eta \upharpoonright \alpha = f_\alpha$. This means that

$$\{\beta < \alpha \mid \eta(\beta) \neq \xi(\beta)\}$$

is stationary, and so $(\eta, \xi) \notin E_S^{\kappa,\kappa} \upharpoonright \alpha$ holds for all $\alpha \in A$. However $\eta \upharpoonright \alpha = f_\alpha$ implies that $(\eta, g_\alpha) \in E_S^{\kappa,\kappa} \upharpoonright \alpha$, and so by transitivity $(\xi, g_\alpha) \notin E_S^{\kappa,\kappa} \upharpoonright \alpha$. Hence we get that $F(\eta)(\alpha) = 1$, but $F(\xi)(\alpha) = 0$. This holds for all $\alpha \in A$ and A is stationary, so $(F(\eta), F(\xi)) \notin E_{\text{reg}}^{2,\kappa}$. \square

Corollary 3.6 *Suppose $V = L$ and κ is weakly compact. Then $E_{\text{reg}}^{2,\kappa}$ is Σ_1^1 -complete.*

Proof This follows from Theorem 3.1, Fact 3.3 and Theorem 3.5. \square

Corollary 3.7 *Suppose κ is a weakly ineffable cardinal. Then $E_{\text{reg}}^{\kappa,\kappa} \leq_c E_{\text{reg}}^{2,\kappa}$.*

Proof The result follows from Theorem 3.5 and Fact 3.3 \square

Theorem 3.8 *If κ is a Π_2^1 -indescribable cardinal, then $E_{\text{reg}}^{\kappa,\kappa}$ is Σ_1^1 -complete.*

Remark *Here the notion of Π_2^1 -indescribability is the usual one, not to be confused with the Π_1^1 -indescribability from Definition 2.2.*

Proof Let E be a Σ_1^1 equivalence relation on κ^κ . Then there is a closed set C on $\kappa^\kappa \times \kappa^\kappa \times \kappa^\kappa$ such that $\eta E \xi$ if and only if there exists $\theta \in \kappa^\kappa$ such that $(\eta, \xi, \theta) \in C$. Let us define $U = \{(\eta \upharpoonright \alpha, \xi \upharpoonright \alpha, \theta \upharpoonright \alpha) \mid (\eta, \xi, \theta) \in C \wedge \alpha < \kappa\}$, and for every $\gamma < \kappa$ define

$$C_\gamma = \{(\eta, \xi, \theta) \in \gamma^\gamma \times \gamma^\gamma \times \gamma^\gamma \mid \forall \alpha < \gamma (\eta \upharpoonright \alpha, \xi \upharpoonright \alpha, \theta \upharpoonright \alpha) \in U\}.$$

Let $E_\gamma \subset \gamma^\gamma \times \gamma^\gamma$ be the relation defined by $(\eta, \xi) \in E_\gamma$ if and only if there exists $\theta \in \gamma^\gamma$ such that $(\eta, \xi, \theta) \in C_\gamma$. Notice that E_γ is not necessarily an equivalence relation. Let us define the reduction by

$$F(\eta)(\alpha) = \begin{cases} f_\alpha(\eta) & \text{if } E_\alpha \text{ is an equivalence relation and } \eta \upharpoonright \alpha \in \alpha^\alpha \\ 0 & \text{otherwise.} \end{cases}$$

where $f_\alpha(\eta)$ is a code in $\kappa \setminus \{0\}$ for the E_α -equivalence class of η .

Let us prove that if $(\eta, \xi) \in E$, then $(F(\eta), F(\xi)) \in E_{\text{reg}}^{\kappa,\kappa}$. Suppose $(\eta, \xi) \in E$. Then there is $\theta \in \kappa^\kappa$ such that $(\eta, \xi, \theta) \in C$ and for all $\alpha < \kappa$ we have that $(\eta \upharpoonright \alpha, \xi \upharpoonright \alpha, \theta \upharpoonright \alpha) \in U$. On the other hand, we know that there is a club D such that for all $\alpha \in D \cap \text{reg}(\kappa)$, $\eta \upharpoonright \alpha, \xi \upharpoonright \alpha, \theta \upharpoonright \alpha \in \alpha^\alpha$. We conclude that for all

$\alpha \in D \cap \text{reg}(\kappa)$, if E_α is an equivalence relation, then $(\eta, \xi) \in E_\alpha$. Therefore, for all $\alpha \in D \cap \text{reg}(\kappa)$, $F(\eta)(\alpha) = F(\xi)(\alpha)$, so $(F(\eta), F(\xi)) \in E_{\text{reg}}^{\kappa, \kappa}$. Let us prove that if $(\eta, \xi) \notin E$, then $(F(\eta), F(\xi)) \notin E_{\text{reg}}^{\kappa, \kappa}$. Suppose $\eta, \xi \in \kappa^\kappa$ are such that $(\eta, \xi) \notin E$. We know that there is a club D such that for all $\alpha \in D \cap \text{reg}(\kappa)$, $\eta \restriction \alpha, \xi \restriction \alpha \in \alpha^\alpha$.

Notice that because C is closed $(\eta, \xi) \notin E$ is equivalent to

$$\forall \theta \in \kappa^\kappa (\exists \alpha < \kappa (\eta \restriction \alpha, \xi \restriction \alpha, \theta \restriction \alpha) \notin U),$$

so the sentence $(\eta, \xi) \notin E$ is a Π_1^1 property of the structure $(V_\kappa, \in, U, \eta, \xi)$. On the other hand, the sentence $\forall \zeta_1, \zeta_2, \zeta_3 \in \kappa^\kappa [((\zeta_1, \zeta_2) \in E \wedge (\zeta_2, \zeta_3) \in E) \rightarrow (\zeta_1, \zeta_3) \in E]$ is equivalent to the sentence $\forall \zeta_1, \zeta_2, \zeta_3, \theta_1, \theta_2 \in \kappa^\kappa [\exists \theta_3 \in \kappa^\kappa (\psi_1 \vee \psi_2 \vee \psi_3)]$, where ψ_1, ψ_2 and ψ_3 are, respectively, the formulas $\exists \alpha_1 < \kappa (\zeta_1 \restriction \alpha_1, \zeta_2 \restriction \alpha_1, \theta_1 \restriction \alpha_1) \notin U$, $\exists \alpha_2 < \kappa (\zeta_2 \restriction \alpha_2, \zeta_3 \restriction \alpha_2, \theta_2 \restriction \alpha_2) \notin U$, and $\forall \alpha_3 < \kappa (\zeta_1 \restriction \alpha_3, \zeta_3 \restriction \alpha_3, \theta_3 \restriction \alpha_3) \in U$. Therefore, the sentence $\forall \zeta_1, \zeta_2, \zeta_3 \in \kappa^\kappa [((\zeta_1, \zeta_2) \in E \wedge (\zeta_2, \zeta_3) \in E) \rightarrow (\zeta_1, \zeta_3) \in E]$ is a Π_2^1 property of the structure (V_κ, \in, U) .

The sentence $\forall \zeta_1, \zeta_2 \in \kappa^\kappa [(\zeta_1, \zeta_2) \in E \rightarrow (\zeta_2, \zeta_1) \in E]$ is equivalent to the sentence $\forall \zeta_1, \zeta_2, \theta_1 \in \kappa^\kappa [\exists \theta_2 \in \kappa^\kappa (\psi_1 \vee \psi_2)]$, where ψ_1 and ψ_2 are, respectively, the formula $\exists \alpha_1 < \kappa (\zeta_1 \restriction \alpha_1, \zeta_2 \restriction \alpha_1, \theta_1 \restriction \alpha_1) \notin U$, and the formula $\forall \alpha_2 < \kappa (\zeta_2 \restriction \alpha_2, \zeta_1 \restriction \alpha_2, \theta_2 \restriction \alpha_2) \in U$.

Therefore, the sentence $\forall \zeta_1, \zeta_2 \in \kappa^\kappa [(\zeta_1, \zeta_2) \in E \rightarrow (\zeta_2, \zeta_1) \in E]$ is a Π_2^1 property of the structure (V_κ, \in, U) .

The sentence $\forall \zeta \in \kappa^\kappa [(\zeta, \zeta) \in E]$ is equivalent to the following sentence

$$\forall \zeta \in \kappa^\kappa [\exists \theta \in \kappa^\kappa (\forall \alpha < \kappa (\zeta \restriction \alpha, \zeta \restriction \alpha, \theta \restriction \alpha) \in U)].$$

Therefore, the sentence $\forall \zeta \in \kappa^\kappa [(\zeta, \zeta) \in E]$ is a Π_2^1 property of the structure (V_κ, \in, U) .

It follows that the sentence

$$(D \text{ is unbounded in } \kappa) \wedge ((\eta, \xi) \notin E) \wedge (E \text{ is an equivalence relation}) \wedge (\kappa \text{ is regular})$$

is a Π_2^1 property of the structure $(V_\kappa, \in, U, \eta, \xi)$. By Π_2^1 reflection, we know that there are stationary many $\gamma \in \text{reg}(\kappa)$ such that γ is a limit point of D , E_γ is an equivalence relation, and $(\eta \restriction \gamma, \xi \restriction \gamma) \notin E_\gamma$. We conclude that there are stationary many $\gamma \in \text{reg}(\kappa)$ such that $f_\gamma(\eta) \neq f_\gamma(\xi)$, and hence $(F(\eta), F(\xi)) \notin E_{\text{reg}}^{\kappa, \kappa}$. \square

Corollary 3.9 *Suppose κ is an ineffable cardinal, or weakly ineffable and Π_2^1 indescribable. Then $E_{\text{reg}}^{2, \kappa}$ is Σ_1^1 -complete.*

Proof An ineffable cardinal is both weakly ineffable and Π_2^1 -indescribable. So the result follows by combining Corollary 3.7 and Theorem 3.8. \square

We will finish this article with a model theoretic result.

Theorem 3.10 *Let DLO be the theory of dense linear orderings without end points. If κ is a Π_2^1 -indescribable cardinal, then \cong_{DLO} is Σ_1^1 -complete.*

Proof By Theorem 3.8 it is enough to show that $E_{\text{reg}}^{\kappa, \kappa} \leq_c \cong_{\text{DLO}}$. To show this, first we will construct models of DLO, $\mathcal{A}^{\mathcal{F}(f)}$, for every $f: \kappa \rightarrow \kappa$, such that $f \in E_{\text{reg}}^{\kappa, \kappa}$ if and only if $\mathcal{A}^{\mathcal{F}(f)} \cong \mathcal{A}^{\mathcal{F}(g)}$. After that we construct the reduction of $E_{\text{reg}}^{\kappa, \kappa}$ to \cong_{DLO} .

Let us take the language $\mathcal{L}^l = \{L, C, <, R\}$, with L and C as unary predicates, and $<$ and R as binary relations. Let K be the class of \mathcal{L}^l -structures $\mathcal{A} = (\text{dom}(\mathcal{A}), L, C, <, R)$ that satisfy the following conditions:

- $L \cap C = \emptyset$.
- $L \cup C = \text{dom}(\mathcal{A})$.
- $< \subseteq L \times L$ is a dense linear order without end points on L .
- $R \subseteq L \times C$.
- Let us denote by $R^-(y, x)$ the formula $\neg R(y, x)$. For all $x \in C$, it holds that $R(\mathcal{A}, x) \cup R^-(\mathcal{A}, x) = L$, $R(\mathcal{A}, x)$ has no largest element, and $R^-(\mathcal{A}, x)$ has no least element and they are non-empty.

Let us define the following partial order \preceq on K . We say that $\mathcal{A} \preceq \mathcal{B}$ iff:

- $\mathcal{A} \subseteq \mathcal{B}$,
- for all $x \in C^{\mathcal{A}}$, $R(\mathcal{B}, x) = \{y \in L^{\mathcal{B}} \mid \exists z \in R(\mathcal{A}, x), y < z\}$ and $R^-(\mathcal{B}, x) = \{y \in L^{\mathcal{B}} \mid \exists z \in R^-(\mathcal{A}, x), z < y\}$,
- for all $x \in C^{\mathcal{B}} \setminus C^{\mathcal{A}}$ there are $y \in R(\mathcal{B}, x)$ and $z \in R^-(\mathcal{B}, x)$ such that for all $a \in L^{\mathcal{A}}$, $a < y \vee a > z$.

Notice that it is possible to have a chain $\mathcal{A}_0 \preceq \mathcal{A}_1 \preceq \dots$ of length α in K , and a structure $\mathcal{C} \in K$, such that $\bigcup_{i < \alpha} \mathcal{A}_i \in K$, $\mathcal{A}_i \preceq \mathcal{C}$ holds for all $i < \alpha$, and $\bigcup_{i < \alpha} \mathcal{A}_i \not\preceq \mathcal{C}$. But all other requirements of AEC's are satisfied, as one can easily see, in particular for every chain $\mathcal{A}_0 \preceq \mathcal{A}_1 \preceq \dots$ of length α in K , $\bigcup_{i < \alpha} \mathcal{A}_i \in K$.

Claim 3.10.1 (K, \preceq) has the amalgamation property and the joint embedding property.

Proof The joint embedding property is easily seen to follow from the amalgamation property. For the amalgamation property, let $\mathcal{A}, \mathcal{B}, \mathcal{C} \in K$ be such that $\mathcal{A} \preceq \mathcal{B}$ and $\mathcal{A} \preceq \mathcal{C}$ hold. Without loss of generality, we can assume that $\text{dom}(\mathcal{B}) \cap \text{dom}(\mathcal{C}) = \text{dom}(\mathcal{A})$. Let us construct \mathcal{D} with $\text{dom}(\mathcal{D})$ equals to $\text{dom}(\mathcal{B}) \cup \text{dom}(\mathcal{C})$, $L^{\mathcal{D}} = L^{\mathcal{B}} \cup L^{\mathcal{C}}$, and $C^{\mathcal{D}} = C^{\mathcal{B}} \cup C^{\mathcal{C}}$. To define $<^{\mathcal{D}}$ and $R^{\mathcal{D}}$, first define $<' = <^{\mathcal{B}} \cup <^{\mathcal{C}}$. For every two elements $b, c \in L^{\mathcal{D}}$ define $b <^{\mathcal{D}} c$ if either $b <' c$, or there is $a \in L^{\mathcal{A}}$ such that $b <' a <' c$, or $b \in L^{\mathcal{B}}$, $c \in L^{\mathcal{C}}$ and there is no $a \in L^{\mathcal{A}}$ such that $c <' a <' b$. For every $x \in C^{\mathcal{A}}$, $R(\mathcal{D}, x) = R(\mathcal{B}, x) \cup R(\mathcal{C}, x)$. For all $x \in C^{\mathcal{B}} \setminus C^{\mathcal{A}}$, $y \in R(\mathcal{D}, x)$ if and only if there exists $z \in L^{\mathcal{B}}$ such that $z \in R(\mathcal{B}, x)$ and $y <^{\mathcal{D}} z$. For all $x \in C^{\mathcal{C}} \setminus C^{\mathcal{A}}$, $y \in R(\mathcal{D}, x)$ if and only if there exists $z \in L^{\mathcal{C}}$ such that $z \in R(\mathcal{C}, x)$ and $y <^{\mathcal{D}} z$. It is clear that $\mathcal{D} \in K$, and $\mathcal{B} \preceq \mathcal{D}$ and $\mathcal{C} \preceq \mathcal{D}$. \square

Let us denote by $\mathcal{A}_1 \oplus_{\mathcal{A}_0} \mathcal{A}_2$ the structure \mathcal{D} , in Claim 3.10.1, that witnesses the amalgamation property for the structures $\mathcal{A}_0 \preceq \mathcal{A}_1$ and $\mathcal{A}_0 \preceq \mathcal{A}_2$. For every ordinal α , let us denote by α^* the set α ordered by the reverse order $<^*$, i.e., $\beta <^* \gamma$ if $\gamma \in \beta$. Let us order the members of $\mathbb{Q} \times \alpha^*$ by: $(r_1, \alpha_1) <^{*\alpha} (r_2, \alpha_2)$ iff $\alpha_1 <^* \alpha_2$, or $\alpha_1 = \alpha_2$ and $r_1 <^{\mathbb{Q}} r_2$.

Let $K_{<\kappa}$ be the collection of all members of K of size less than κ . For every $\mathcal{A} \in K_{<\kappa}$, denote by $\{\mathcal{A}(i)\}_{i < \kappa}$ an enumeration of all the strong extensions of \mathcal{A} , i.e. $\mathcal{A} \preceq \mathcal{B}$, of size less than κ (up to isomorphism over \mathcal{A}). Let $\Pi: \kappa \rightarrow \kappa \times \kappa$, $\Pi(\alpha) = (pr_1(\Pi(\alpha)), pr_2(\Pi(\alpha)))$ be a bijection such that $pr_1(\Pi(i)) \leq i$ for all i . Given a function $f: \kappa \rightarrow \text{reg}(\kappa)$, let us construct the following sequence of models:

- $\mathcal{A}_0^f = (\mathbb{Q}, \emptyset, <, \emptyset)$.
- For a successor ordinal, let $\mathcal{D} = \mathcal{A}_i^f \oplus_{\mathcal{A}^f} \mathcal{A}_{pr_1(\Pi(i))}^f (pr_2(\Pi(i)))$. Define $L^{\mathcal{A}_{i+1}^f} = L^{\mathcal{D}} \cup \mathbb{Q}$, $C^{\mathcal{A}_{i+1}^f} = C^{\mathcal{D}}$, $<^{\mathcal{A}_{i+1}^f} = <^{\mathcal{D}} \cup <^{\mathbb{Q}} \cup \{(x, y) \mid x \in L^{\mathcal{D}} \wedge y \in \mathbb{Q}\}$, and $R^{\mathcal{A}_{i+1}^f} = R^{\mathcal{D}}$. Clearly $\mathcal{A}_{i+1}^f \in K$.

- For i a limit ordinal, let $\mathcal{D} = \bigcup_{j < i} \mathcal{A}_j^f$. Define $L^{\mathcal{A}_i^f} = L^{\mathcal{D}} \cup (\mathbb{Q} \times f(i)^*)$, $C^{\mathcal{A}_i^f} = C^{\mathcal{D}} \cup \{x\}$, $<^{\mathcal{A}_i^f} = <^{\mathcal{D}} \cup <^{*f(i)} \cup \{(a, b) \mid a \in L^{\mathcal{D}} \wedge b \in \mathbb{Q} \times f(i)^*\}$, and $R^{\mathcal{A}_i^f} = R^{\mathcal{D}} \cup \{(y, x) \mid y \in L^{\mathcal{D}}\}$. Clearly $\mathcal{A}_i^f \in K$.

Define \mathcal{A}_κ^f by $\bigcup_{j < \kappa} \mathcal{A}_j^f$. Then $\mathcal{A}^f = (L^{\mathcal{A}_\kappa^f}, <^{\mathcal{A}_\kappa^f})$ is a model of DLO.

Notice that if $i < \kappa$ and $\mathcal{C} \in K$, $|\mathcal{C}| < \kappa$, are such that $\mathcal{A}_i^f \preceq \mathcal{C}$, then there is $j < \kappa$ such that $\mathcal{A}_i^f(j) = \mathcal{C}$. Therefore there is $l < \kappa$ such that $\Pi(l) = (i, j)$, $\mathcal{A}_{pr_1(\Pi(l))}^f = \mathcal{A}_i^f$, and $\mathcal{A}_{pr_2(\Pi(l))}^f(pr_2(\Pi(l))) = \mathcal{C}$. We conclude that if $i < \kappa$ and $\mathcal{C} \in K_{< \kappa}$ are such that $\mathcal{A}_i^f \preceq \mathcal{C}$, then there is $j < \kappa$ and a strong embedding $F: \mathcal{C} \rightarrow \mathcal{A}_j^f$ such that $F(\mathcal{C}) \preceq \mathcal{A}_j^f$ and $F \upharpoonright \mathcal{A}_i^f = id$. Now we will show that if f and g are functions from κ into $\text{reg}(\kappa)$ such that $f \upharpoonright (\kappa \setminus \text{reg}(\kappa)) = g \upharpoonright (\kappa \setminus \text{reg}(\kappa))$, then $f E_{\text{reg}}^{\kappa, \kappa} g$ if and only if $\mathcal{A}^f \cong \mathcal{A}^g$. First of all, let us prove that $(f, g) \in E_{\text{reg}}^{\kappa, \kappa}$ implies $\mathcal{A}^f \cong \mathcal{A}^g$. Suppose $(f, g) \in E_{\text{reg}}^{\kappa, \kappa}$. Then there is a club C such that for all $\alpha \in C \cap \text{reg}(\kappa)$, $f(\alpha) = g(\alpha)$. Since $f \upharpoonright (\kappa \setminus \text{reg}(\kappa)) = g \upharpoonright (\kappa \setminus \text{reg}(\kappa))$, we have that for all $\alpha \in C$, $f(\alpha) = g(\alpha)$. By the way the models \mathcal{A}_α^f and \mathcal{A}_α^g were constructed for α a limit ordinal, we know that if α is such that $f(\alpha) = g(\alpha)$ and there is an isomorphism $F: \bigcup_{i < \alpha} \mathcal{A}_i^f \rightarrow \bigcup_{i < \alpha} \mathcal{A}_i^g$, then there is an isomorphism $G: \mathcal{A}_\alpha^f \rightarrow \mathcal{A}_\alpha^g$ such that $F \subseteq G$. For all $i < \kappa$ construct $\alpha_i < \kappa$ and a strong embedding F_i such that the following hold:

- For every $i < \kappa$ there is some $\gamma \in C$ such that $\alpha_i < \gamma < \alpha_{i+1}$.
- For all $i < j < \kappa$, $f_i \subseteq f_j$.
- The following holds for every limit ordinal $\beta < \kappa$:
 - for every even $0 < i < \omega$, $\text{dom}(F_{\beta+i}) = \mathcal{A}_{\alpha_{\beta+i}}^f$, and $F_{\beta+i}(\mathcal{A}_{\alpha_{\beta+i}}^f) \preceq \mathcal{A}_{\alpha_{\beta+i+1}}^g$,
 - for every odd $0 < i < \omega$, $\text{rang}(F_{\beta+i}) = \mathcal{A}_{\alpha_{\beta+i}}^g$, and $F_{\beta+i}^{-1}(\mathcal{A}_{\alpha_{\beta+i}}^g) \preceq \mathcal{A}_{\alpha_{\beta+i+1}}^f$,
 - $\alpha_\beta = \bigcup_{i < \beta} \alpha_i$, $\text{dom}(F_\beta) = \mathcal{A}_{\alpha_\beta}^f$, and $\text{rang}(F_\beta) = \mathcal{A}_{\alpha_\beta}^g$.

We will construct these sequences by induction. For $i = 0$, take $\alpha_0 = 0$ and $F_0 = id$.

Successor case: Suppose β is a limit ordinal or zero, and $0 \leq i < \omega$ are such that $\alpha_{\beta+i}$ and $F_{\beta+i}$ are constructed such that (i), (ii), and (iii) are satisfied. Let us start with the case when i is odd. Choose $\alpha_{\beta+i+1}$ such that (i) holds. Since $F^{-1}(\mathcal{A}_{\alpha_{\beta+i}}^g) \preceq \mathcal{A}_{\alpha_{\beta+i+1}}^f$, there are $\mathcal{C} \in K_{< \kappa}$ and $F \supseteq F_{\beta+i}$ such that $\mathcal{A}_{\alpha_{\beta+i}}^g \preceq \mathcal{C}$ and $F: \mathcal{A}_{\alpha_{\beta+i+1}}^f \rightarrow \mathcal{C}$ is an isomorphism. By the observation we made above, there is $j < \kappa$ and a strong embedding $G: \mathcal{C} \rightarrow \mathcal{A}_j^g$ such that $G(\mathcal{C}) \preceq \mathcal{A}_j^g$ and $G \upharpoonright \mathcal{A}_{\alpha_{\beta+i}}^g = id$. Define $F_{\alpha_{\beta+i+1}} = G \circ F_{\alpha_{\beta+i}}$. Clearly $F_{\alpha_{\beta+i+1}}$ satisfies conditions (ii) and (iii). The case when i is even is similar to the odd case.

Limit case: Suppose β is a limit ordinal such that for all $i < \beta$, α_i and F_i are constructed such that (i), (ii), and (iii) are satisfied. By (i), we know that $\alpha_\beta = \bigcup_{i < \beta} \alpha_i$ is a limit point of C , so $f(\alpha_\beta) = g(\alpha_\beta)$. On the other hand, by conditions (ii) and (iii) we know that

$$\bigcup_{i < \beta} F_i: \bigcup_{i < \beta} \mathcal{A}_{\alpha_i}^f \rightarrow \bigcup_{i < \beta} \mathcal{A}_{\alpha_i}^g$$

is an isomorphism. Therefore, there is an isomorphism $G: \mathcal{A}_\beta^f \rightarrow \mathcal{A}_\beta^g$ such that $\bigcup_{i < \beta} F_i \subseteq G$. We conclude that $F_{\alpha_\beta} = G$ satisfies (ii) and (iii).

Finally, notice that

$$\bigcup_{i < \kappa} F_i: \bigcup_{i < \kappa} \mathcal{A}_{\alpha_i}^f \rightarrow \bigcup_{i < \kappa} \mathcal{A}_{\alpha_i}^g$$

is an isomorphism. We conclude that \mathcal{A}^f and \mathcal{A}^g are isomorphic.

Let us prove that $\mathcal{A}^f \cong \mathcal{A}^g$ implies $(f, g) \in E_{\text{reg}}^{\kappa, \kappa}$. Suppose, towards a contradiction, that $(f, g) \notin E_{\text{reg}}^{\kappa, \kappa}$ and there is an isomorphism $F: \mathcal{A}^f \rightarrow \mathcal{A}^g$. Since F is an isomorphism, there is a club C such that $F(\bigcup_{i < \alpha} \mathcal{A}_i^f) = \bigcup_{i < \alpha} \mathcal{A}_i^g$ holds for all $\alpha \in C$. Since $(f, g) \notin E_{\text{reg}}^{\kappa, \kappa}$, $C \cap \{\alpha \in \text{reg}(\kappa) \mid f(\alpha) \neq g(\alpha)\}$ is nonempty. Take $\alpha \in C \cap \{\gamma \in \text{reg}(\kappa) \mid f(\gamma) \neq g(\gamma)\}$. We know that $F(\bigcup_{i < \alpha} \mathcal{A}_i^f) = \bigcup_{i < \alpha} \mathcal{A}_i^g$ and $f(\alpha) \neq g(\alpha)$. Hence, the co-initiality of $\{a \in \mathcal{A}^f \mid \forall b \in \bigcup_{i < \alpha} \mathcal{A}_i^f (b <^{\mathcal{A}^f} a)\}$ with respect to $<^{\mathcal{A}^f}$ is $f(\alpha)$. Since F is an isomorphism and $F(\bigcup_{i < \alpha} \mathcal{A}_i^f) = \bigcup_{i < \alpha} \mathcal{A}_i^g$, the co-initiality of $\{a \in \mathcal{A}^g \mid \forall b \in \bigcup_{i < \alpha} \mathcal{A}_i^g (b <^{\mathcal{A}^g} a)\}$ with respect to $<^{\mathcal{A}^g}$ is also $f(\alpha)$. We conclude that $f(\alpha) = g(\alpha)$, a contradiction. To finish with the construction of the models, let us define $\mathcal{A}^{\mathcal{F}(f)}$ for all $f: \kappa \rightarrow \kappa$. Fix a bijection $G: \kappa \rightarrow \text{reg}(\kappa)$. Define $\mathcal{F}: \kappa^\kappa \rightarrow \kappa^\kappa$ by

$$\mathcal{F}(f)(\alpha) = \begin{cases} G(f(\alpha)) & \text{if } \alpha \in \text{reg}(\kappa) \\ 0 & \text{otherwise} \end{cases}$$

Clearly $f E_{\text{reg}}^{\kappa, \kappa} g$ if and only if $\mathcal{F}(f) E_{\text{reg}}^{\kappa, \kappa} \mathcal{F}(g)$, and $\mathcal{F}(f) E_{\text{reg}}^{\kappa, \kappa} \mathcal{F}(g)$ if and only if $\mathcal{A}^{\mathcal{F}(f)}$ and $\mathcal{A}^{\mathcal{F}(g)}$ are isomorphic. Now we will construct a reduction of $E_{\text{reg}}^{\kappa, \kappa}$ to \cong_{DLO} by coding the models $\mathcal{A}^{\mathcal{F}(f)}$ by functions $\eta: \kappa \rightarrow \kappa$.

Clearly the models $\mathcal{A}^{\mathcal{F}(f)}$ satisfy that

$$\mathcal{F}(f) \upharpoonright \alpha = \mathcal{F}(g) \upharpoonright \alpha \Leftrightarrow \mathcal{A}_\alpha^{\mathcal{F}(f)} = \mathcal{A}_\alpha^{\mathcal{F}(g)}.$$

For every $f \in \kappa^\kappa$ define $C_f \subseteq \text{Card} \cap \kappa$ such that for all $\alpha \in C_f$, it holds that for every $\beta < \alpha$, $|\mathcal{A}_\beta^{\mathcal{F}(f)}| < |\mathcal{A}_\alpha^{\mathcal{F}(f)}|$. For every $f \in \kappa^\kappa$ and $\alpha \in C_f$ choose a bijection $E_f^\alpha: \text{dom}(\mathcal{A}_\alpha^{\mathcal{F}(f)}) \rightarrow |\mathcal{A}_\alpha^{\mathcal{F}(f)}|$ such that for all $\beta < \alpha$ in C_f it holds that $E_f^\beta \subseteq E_f^\alpha$. Then $\bigcup_{\alpha \in C_f} E_f^\alpha = E_f$ is such that $E_f: \text{dom}(\mathcal{A}^{\mathcal{F}(f)}) \rightarrow \kappa$ is a bijection, and for every $f, g \in \kappa^\kappa$ and $\alpha < \kappa$ the following holds: If $\mathcal{F}(f) \upharpoonright \alpha = \mathcal{F}(g) \upharpoonright \alpha$, then $E_f \upharpoonright \text{dom}(\mathcal{A}_\alpha^{\mathcal{F}(f)}) = E_g \upharpoonright \text{dom}(\mathcal{A}_\alpha^{\mathcal{F}(g)})$. Let π be the bijection in Definition 1.4. Define the function \mathcal{G} by:

$$\mathcal{G}(\mathcal{F}(f))(\alpha) = \begin{cases} 1 & \text{if } \alpha = \pi(m, a_1, \dots, a_n) \text{ and } \mathcal{A}^{\mathcal{F}(f)} \models P_m(E_f^{-1}(a_1), \dots, E_f^{-1}(a_n)) \\ 0 & \text{in the other case.} \end{cases}$$

To show that \mathcal{G} is continuous, let $[\eta \upharpoonright \alpha]$ be a basic open set and $\xi \in \mathcal{G}^{-1}[[\eta \upharpoonright \alpha]]$. There is $\beta \in C_\xi$ such that for all $\gamma < \alpha$, if $\gamma = \pi(m, a_1, a_2, \dots, a_n)$, then $E_\xi^{-1}(a_i)$ is an element of $\text{dom}(\mathcal{A}_\beta^\xi)$ for all $i \leq n$. Since for all $\zeta \in [\xi \upharpoonright \beta]$ it holds that $\mathcal{A}_\beta^\xi = \mathcal{A}_\beta^\zeta$, for every $\gamma < \alpha$ such that $\gamma = \pi(m, a_1, a_2, \dots, a_n)$, it holds that

$$\mathcal{A}_\beta^\xi \models P_m(E_\xi^{-1}(a_1), E_\xi^{-1}(a_2), \dots, E_\xi^{-1}(a_n))$$

if and only if

$$\mathcal{A}_\beta^\zeta \models P_m(E_\zeta^{-1}(a_1), E_\zeta^{-1}(a_2), \dots, E_\zeta^{-1}(a_n))$$

We conclude that $\mathcal{G}(\zeta) \in [\eta \upharpoonright \alpha]$, and $\mathcal{G} \circ \mathcal{F}$ is a continuous reduction of $E_{\text{reg}}^{\kappa, \kappa}$ to \cong_{DLO} . \square

4 Further research

In this paper we established the Σ_1^1 -completeness of a range of equivalence relations in various circumstances. Some of these theorems are proved in ZFC, some are consistency results, and some are relative consistency results. In particular the equivalence relation modulo the non-stationary ideal is Σ_1^1 -complete if κ is an ineffable cardinal. This, and related equivalence relations, play a role in model theory as exemplified by Theorem 3.10 which shows how generalized descriptive set theory is different from the classical study where $\kappa = \omega$ and the isomorphism relation of countable structures is never Σ_1^1 -complete. This was also the original motivation for studying such fine-grained questions as whether $E_{\mu\text{-club}}^{\kappa, \kappa}$ can be reduced to $E_{\mu\text{-club}}^{2, \kappa}$ for some $\mu < \kappa$. How much more can one prove in ZFC for $\kappa > \omega$? For successor cardinals the answer is partially known [4] starting from $V = L$ for every successor cardinal κ there exists a GCH and cardinal preserving forcing notion such that in the extension the equivalence relation modulo the non-stationary ideal is not Σ_1^1 -complete. The following questions remain open.

Question 4.1 *Is it consistent that the isomorphism relation on graphs or dense linear orders is not Σ_1^1 -complete for some $\kappa > \omega$? Of course κ cannot be Π_2^1 -indescribable by Theorem 3.10.*

Question 4.2 *Is it consistent for some cardinal κ and a regular $\mu < \kappa$ that $E_{\mu}^{\kappa, \kappa}$ is not reducible to $E_{\mu}^{2, \kappa}$? Note: it has been shown [1] that it is consistent that $E_S^{2, \kappa}$ is not reducible to $E_{S'}^{2, \kappa}$ for $S' \setminus S$ stationary which implies the consistency of e.g. $E_{\mu}^{\kappa, \kappa} \not\leq_B E_{\mu'}^{2, \kappa}$ for $\mu \neq \mu'$.*

Question 4.3 *Is it consistent that κ is inaccessible and $E_S^{2, \kappa}$ is not Σ_1^1 -complete for some stationary $S \subset \kappa$? What about κ weakly compact and $S = S_{\mu}^{\kappa}$ for some regular $\mu < \kappa$? Note: it follows from the result of [4] that it is consistent that $E_{\kappa}^{2, \kappa}$ is not Σ_1^1 -complete (in fact Δ_1^1) for successor κ .*

References

- [1] Friedman, S.D., Hyttinen, T., and Kulikov, V. “Generalized descriptive set theory and classification theory,” *Memories of the American Mathematical Society*, vol. 230 (2014). [1](#), [2](#), [4](#), [5](#), [6](#), [10](#), [17](#)
- [2] Friedman, S.D., Hyttinen, T., and Kulikov, V. *On Borel reducibility in Generalized Baire space*, *Fundamenta Mathematicae*, vol. 203 (2015), pp. 285 – 298. [2](#)
- [3] Friedman, H., and Stanley, L. *A Borel reducibility theory for classes of countable structures*, *The Journal of Symbolic Logic*, vol. 54 (1989), pp. 894 – 914. [2](#)
- [4] Friedman, S.-D., Wu, L., and Zdomskyy, L. Δ_1 -Definability of the non-stationary ideal at successor cardinals., *Fundamenta Mathematicae* 229 (2015): 231-254. [17](#)

- [5] Hellsten, A. *Diamonds on large cardinals*, *Finnish Academy of Science and Letters* vol. 134 (2003) <https://www.acadsci.fi/mathematica/e-theses/hellsten.pdf> 11
- [6] Hyttinen, T., and Kulikov, V. *On Σ_1^1 -complete equivalence relations on the generalized Baire space*, *Mathematical Logic Quarterly*, vol. 61 (2015), pp. 66 – 81. 2, 10
- [7] Khomskii, Y., Laguzzi, G., Löwe, B., and Sharankou, I. *Questions on generalised Baire spaces*, *Mathematical Logic Quarterly*, vol. 62 (2016), pp. 439 – 456. 2
- [8] Jech, T., and Shelah, S. *Full reflection of stationary sets below \aleph_ω* , *The Journal of Symbolic Logic*, vol. 55 (1990), pp. 822 – 830. 4, 10
- [9] Sun, W. *Stationary cardinals* *Arch. Math. Logic*, vol. 32, nr. 6, (1993), pp. 429 – 442. 10

Asperó
School of Mathematics
University of East Anglia
Norwich Research Park
Norwich, Norfolk, NR4 7TJ
UK
d.aspero@uea.ac.uk

Hyttinen
Department of Mathematics and Statistics
University of Helsinki
P.O. Box 68
Helsinki
FINLAND
tapani.hyttinen@helsinki.fi

Kulikov
Department of Mathematics and Statistics
University of Helsinki
P.O. Box 68
Helsinki
FINLAND
vadim.kulikov@helsinki.fi

Moreno
Department of Mathematics and Statistics
University of Helsinki
P.O. Box 68
Helsinki
FINLAND
miguel.moreno@helsinki.fi