

This is the peer-reviewed version of the following article:

Sutter A, Travers LM, Weedon M, Oku K, Price TAR, Wedell N. No selection for change in polyandry under experimental evolution. *J Evol Biol.* 2019; 0 0:1–14, which has been published in final form at <https://doi.org/10.1111/jeb.13476>

No selection for change in polyandry under experimental evolution

Andreas Sutter^{†§1}, Laura M. Travers^{†2}, Melanie Weedon[†], Keiko Oku[†], Thomas A. R. Price[‡], & Nina Wedell[†]

[†] Centre for Ecology & Conservation, College of Life and Environmental Sciences, University of Exeter, Cornwall Campus, TR10 9FE, Penryn, UK.

[‡] Institute of Integrative Biology, University of Liverpool, Crown Street, Liverpool, L69 7ZB, UK

§ corresponding author details: email a.sutter@uea.ac.uk; telephone: +44 (0) 1603 456161; fax: n/a

Running title:

No experimental evolution of polyandry

Abstract

What drives mating system variation is a major question in evolutionary biology. Female multiple mating (polyandry) has diverse evolutionary consequences, and there are many potential benefits and costs of polyandry. However, our understanding of its evolution is biased towards studies enforcing monandry in polyandrous species. What drives and maintains variation in polyandry between individuals, genotypes, populations and species remains poorly understood.

¹ Current address: Centre for Ecology, Evolution and Conservation, School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich, NR4 7TJ, UK

² Current address: Centre for Ecology, Evolution and Conservation, School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich, NR4 7TJ, UK

Genetic variation in polyandry may be actively maintained by selection, or arise by chance if polyandry is selectively neutral. In *Drosophila pseudoobscura*, there is genetic variation in polyandry between and within populations. We used isofemale lines to found replicate populations with high or low initial levels of polyandry, and tracked polyandry under experimental evolution over seven generations. Polyandry remained relatively stable, reflecting the starting frequencies of the experimental populations. There were no clear fitness differences between high versus low polyandry genotypes, and there was no signature of balancing selection. We confirmed these patterns in direct comparisons between evolved and ancestral females, and found no consequences of polyandry for female fecundity. The absence of differential selection even when initiating populations with major differences in polyandry casts some doubt on the importance of polyandry for female fitness.

Keywords

multiple mating, monandry, genetic variation, balancing selection, directional selection, isofemale lines, *Drosophila pseudoobscura*

Introduction

Female multiple mating (polyandry) has many important consequences for sexual selection (Parker, 1970; Birkhead & Moller, 1998; Simmons, 2001), population viability (Price *et al.*, 2010a; Holman & Kokko, 2013; Lumley *et al.*, 2015), genetic variation (Balloux & Lehmann, 2003), genome evolution (Mank *et al.*, 2013), and may even drive speciation (Gavrilets, 2014). Polyandry is extremely widespread across the animal kingdom, with evidence for multiple paternity from 89% of all natural populations investigated across animal taxa (Taylor *et al.*,

2014). Much research has focused on the costs and benefits of polyandry (Zeh & Zeh, 1996; Arnqvist & Nilsson, 2000; Jennions & Petrie, 2000; Slatyer *et al.*, 2012), finding substantial support for direct, and mixed support for indirect benefits of multiple mating for females. Nonetheless, given the many factors that potentially influence the dynamics of polyandry, polyandry remains a puzzling trait.

If polyandry is beneficial, how is variation between populations maintained? An intriguing observation shows that polyandry appears to correlate with latitude in many taxa (Taylor *et al.*, 2014), but the reasons for this remain elusive (Price *et al.*, 2012; Taylor *et al.*, 2016). Nevertheless, this points towards a strong role of ecology for regulating a population's mating frequency, either directly by altering the costs/benefits of polyandry (Välimäki *et al.*, 2008), or indirectly by altering the intensity of sexual conflict (Arbuthnott *et al.*, 2014). Sexual conflict over mating rate is very common, and realised mating rates will reflect the outcome of male persistence at making mating attempts and female resistance to such attempts (Parker, 2006). The costs and benefits of accepting or resisting multiple matings can take many forms given a set of ecological circumstances, and females are likely to adjust their mating strategy to optimise their fitness, balancing the costs and benefits of multiple mating (Arnqvist & Nilsson, 2000). Thus, directional selection should lead the frequency of polyandry towards an externally derived local optimum (Emlen & Oring, 1977; Candolin & Heuschele, 2008). Support for a role of ecological drivers of polyandry come from observations of laboratory adaptation with evolution towards higher or lower frequencies of polyandry (Harano & Miyatake, 2005; Burton-Chellew *et al.*, 2007), presumably because the costs and benefits of (multiple) mating are altered in the lab relative to the wild (Markow, 2011).

The costs and benefits of polyandry are typically assumed to be uniform for all females, such that the same strategy maximises fitness for all females (for reviews see Jennions & Petrie, 2000; Slatyer *et al.*, 2012). Most laboratory experiments on the benefits of polyandry involve drastic manipulations, where females are moved away from evolved optima. Because monandrous species typically cannot be forced to remate (but see e.g. Arnqvist & Andrés, 2006; King & Bressac, 2010), experimenters commonly deny females from polyandrous species any opportunity for remating, and then assess the fitness consequences (e.g. Newcomer *et al.*, 1999; Evans & Magurran, 2000; Gowaty *et al.*, 2010). However, these studies can only explain why monandry does not evolve in polyandrous species but not vice versa. Other studies have used experimental evolution while manipulating the number of males a female mates with, and have revealed adaptations to mating systems both in males and females (e.g. Martin *et al.*, 2004; Wigby & Chapman, 2004; Crudgington *et al.*, 2010; Demont *et al.*, 2014; Perry *et al.*, 2016). In comparison, relatively few studies have experimentally manipulated aspects of the evolving populations to observe how the frequency of polyandry evolves in response (e.g. sex ratio distorter: Price *et al.*, 2008; inbreeding Michalczyk *et al.*, 2011; male sterility: Kuriwada *et al.*, 2014). Studies demonstrating experimental evolution of polyandry highlight that genetic variation within the starting population is an essential requirement for an adaptive response in polyandry to the local conditions. In natural populations, the costs and benefits of polyandry are likely to change dynamically, and females may adopt a flexible strategy that relies on phenotypic plasticity (Gowaty & Hubbell, 2009; Gowaty, 2013). However, evidence that genetic variation in polyandry is commonly present within populations is accumulating (Solymar & Cade, 1990; Sgrò *et al.*, 1998; Wedell, 2001; Torres-Vila *et al.*, 2001, 2002; Simmons, 2003; Shuker *et al.*, 2007; Torres-Vila, 2013; Price *et al.*, 2014; Taylor *et al.*, 2014; Travers *et al.*, 2016). This evidence of standing genetic variation for polyandry opens questions about what maintains it. If there is a

single optimum for females, what maintains genetic variation once that optimum has been reached? To better understand polyandry evolution, we need to understand its fitness consequences in situations that better incorporate selective forces that act in natural populations, including social interactions (e.g. Takahashi & Kawata, 2013).

Most previous studies have simply addressed the question whether polyandry is subject to directional selection, manifested as a fitness difference between monandrous and polyandrous females. However, directional selection should lead to the depletion of genetic variation, and does not explain the presence of genetic variation in polyandry within populations (Taylor *et al.*, 2014). Balancing selection under negative frequency dependence (nFDS) is a pervasive force for maintaining genetic variation (Clarke, 1979; but see Brisson, 2018). Under nFDS, the fitness of a certain genotype or phenotype depends on its frequency in the population, increasing at low frequencies and decreasing when high frequencies are reached (Ayala & Campbell, 1974). In the context of polyandry, the fitness effects of multiple mating may depend on what other females in the population do. Traditionally, evidence for nFDS on reproductive strategies has come from males (e.g. Sinervo & Lively, 1996), but has more recently included female mating strategies (Neff & Svensson, 2013). A thoroughly demonstrated example is female colour-dependent harassment by male *Ischnura* damselflies (Svensson *et al.*, 2005; see also Takahashi & Kawata, 2013). More generally, Svensson and Råberg (2010) suggested that sexual conflict could generally lead to nFDS on female mating strategies, if females avoid the costs of male harassment by tolerance rather than by resistance. Sexual conflict over remating is common, with males trying to manipulate females away from reaching their optimum remating rate. However, females will in turn counteract these manipulations (Arnqvist & Rowe, 2005). If the majority of females mate with multiple males, males may respond to increased levels of sperm competition

by increasing attempts to prevent females from remating, including seminal fluids that decrease female longevity (Chapman *et al.*, 2003). This may give females that mate only once an advantage over polyandrous females through reduced cost of receiving male ejaculates, especially if the costs of mating increase more than linearly (Kuijper *et al.*, 2006). As female mating frequency decreases, males may reduce costs to females (Hollis *et al.*, 2014, 2016), in turn favouring polyandrous females that gain potential benefits of polyandry with reduced exposure to mating costs. At equilibrium, different female mating strategies may have equal net fitness.

Alternatively, genetic variation in polyandry need not be actively maintained through selection. Instead, genetic variation could be maintained by random mutation, especially if polyandry is a highly polygenic trait (e.g. Torres-Vila *et al.*, 2001). Polyandry may be selectively neutral and the frequency of polyandry might change only through genetic drift. This could be true especially in benign conditions such as laboratory environments, where reduced exposure to predators, pathogens and competing species might limit the benefits and costs of multiple mating.

Studying the fitness consequences of polyandry and its evolution in a population context is notoriously difficult, and is not possible in many experimental systems. Here, we use naturally occurring genetic variation in polyandry in the fruit fly *Drosophila pseudoobscura* to investigate selection on polyandry through experimental evolution over multiple generations in a laboratory population context. Using genetic variation in polyandry enabled us to test for fitness consequences of multiple mating in a population setting without manipulating the adult sex ratio or females' access to mates. *D. pseudoobscura* shows remarkable genetic variation in polyandry, both between and within populations. There is genetic variation in average degree of polyandry

between populations across a latitudinal cline across North America (Price *et al.*, 2014). Moreover, genetic variation exists within populations, revealed by comparisons of wild-caught females with their descendants (Price *et al.*, 2011) and through variation between isofemale lines (Herrera *et al.*, 2014; Taylor *et al.*, 2016) that represent a snapshot of the genetic variation in a population (David *et al.*, 2005; Nouhaud *et al.*, 2016). Laboratory experiments show that genetic variation in polyandry is stable with respect to temperature variation (Taylor *et al.*, 2016), and is largely under female control (Price *et al.*, 2008; but see Crudgington *et al.*, 2009 and Price *et al.*, 2010b). Except for in very long-lived females, males provide no direct fitness benefits to females (Turner & Anderson, 1983). Polyandry can however provide indirect benefits for offspring survival (Gowaty *et al.*, 2010). In the presence of a naturally occurring sex ratio distorter, polyandry can have strong fitness benefits by allowing females to avoid fertilisation by distorter-carrying males (Price *et al.*, 2010a). In the presence of this sex ratio distorter, polyandry showed a clear increase within nine generations in experimental evolution (Price *et al.*, 2008). In nature, the distorter correlates negatively with the latitudinal polyandry cline, likely due to polyandry regulating the frequency of the distorter by reduced transmission success (Price *et al.*, 2014). However, what drives and maintains variation in polyandry between populations, and especially within populations, remains unknown (Price *et al.*, 2014; Taylor *et al.*, 2016).

Here, we investigated whether in the absence of the sex ratio distorter, balancing or directional selection acts on polyandry in evolving populations where we eliminated differences in the abiotic environment, but started with an initially high or low representation of polyandrous genotypes. If balancing selection is the main force maintaining variation in polyandry, we would expect all populations to evolve towards an intermediate frequency of polyandry. If polyandry is consistently beneficial or costly, all populations should evolve towards a high or low frequency

of polyandry, irrespective of their initial starting frequency. Finally, if polyandry is selectively neutral, polyandry should remain the same as its initial high or low frequency. We first characterised isofemale lines for female mating behaviour and selected lines that represented differences in the genetic predisposition to mate multiply. Variation in polyandry was continuous, but to create contrasting backgrounds, we grouped isolines into two categories with more polyandrous versus relatively monandrous lines, respectively. Using the selected isolines, we then initiated replicate populations that differed in their initial average frequency of polyandry, and tracked the frequency of polyandry over seven consecutive generations during experimental evolution. Finally, after a generation of common garden breeding, we compared the evolved populations directly with the ancestral isolines with regards to female remating behaviour and fecundity, and male ability to inhibit female remating. Using tester flies that had not co-evolved, we tested female and male effects on polyandry independently. This allowed us to compare the observed patterns to those predicted under different scenarios regarding the evolution of polyandry.

Material and Methods

Establishment of isofemale isogenic lines

Collection and maintenance

We established isofemale isogenic lines using wild female *D. pseudoobscura* from three populations across the Western USA (Lewistown Montana, Show Low Arizona, and Shaver Lake California). We reared full-sib inbred offspring of wild caught females for 15 or more generations, maintaining flies under standardised laboratory conditions throughout. We give a

schematic overview of our methods in Figure 1, and describe full details for our methods in the electronic supplementary material (ESM).

Preliminary assays

We first quantified variation in genetic predisposition for polyandry in 29 isolines using a remating assay routinely performed in our laboratory (Price *et al.*, 2011; Herrera *et al.*, 2014; Taylor *et al.*, 2016). We aspirated sexually mature virgin females from each isoline individually into a vial containing a single male from the same isoline. Males had been separated into individual vials the day before the mating assay to reduce effects arising from prior male-male interactions. We observed matings by scan sampling, and after two hours we discarded all males, as well as females that had not mated. Scan sampling was performed by one or two observers (depending on the size of the assay) who checked vials for mating pairs, observing every vial for a few seconds approximately every two minutes. Females were left to oviposit for four days, after which we aspirated them into the vial of a second male from their isoline and observed them for two hours by scan sampling. Female *D. pseudoobscura* do not remate within 24h (Snook & So, 2000), such that females had a maximum of two matings across the two assay days. We confirmed first matings by presence of larvae in the oviposition vial, but were not able to ascertain sperm transfer in second matings. The proportion of females that remated ranged from 0 to 0.83 for individual isolines (mean 0.28; 28 ± 10 females tested per isoline; Figure 1b and Table S1). A likelihood ratio test between binomial GLMMs including or excluding isoline identity as a random effect confirmed that this variation between isolines was substantial and statistically significant ($\chi^2 = 42.1$, $df = 1$, $N = 821$, $p = 8.7 \times 10^{-11}$).

Selecting focal isolines

To establish our experimental evolution replicates, we chose 16 isolines from the three populations fulfilling the following three criteria: i) eight isolines had to have a relatively high (i.e. more polyandrous *P* lines) versus relatively low (i.e. relatively monandrous *M* lines) frequency of polyandry (see Figure 1), ii) *P* and *M* isolines had to be balanced with regards to population of origin, and iii) polyandry had to have been tested for a satisfactory number of females (N = 21–41). While this meant that the exact threshold that separated *P* from *M* isolines was arbitrary, our method helped avoid biases with respect to representation of the three populations of origin. We repeated the polyandry assay for the 16 chosen isolines before starting experimental evolution, this time giving females two mating opportunities with outbred tester males (population from Chiricahua, Arizona) to minimise male effects on polyandry estimates. The remating proportion of isolines was significantly correlated between this and the prior assay (linear regression weighted by sample size: $R^2 = 0.43$, $F_{1,14} = 12.15$, $p = 0.004$; see Table S1).

Experimental evolution

Population setup and maintenance

We established six replicate experimental evolution populations for each of two treatments. We used all 16 isolines (eight *P*, eight *M* isolines) in all 12 replicates, but varied the relative representation of the isolines between the treatments. We initiated *low polyandry* replicate populations with twelve females and twelve males from each of the eight *M* isolines, and three females and three males from each of the eight *P* isolines. In contrast, we founded *high polyandry* replicate populations with three flies of both sexes from each *M* isoline and twelve flies of both sexes from each *P* isoline (Figure 1c). Thus, we founded all 12 replicate populations with 120 virgin females and 120 virgin males, maintained in large plastic tubs within a single incubator

under standard conditions. From day one to five, flies mated freely for four days. On day five we removed males and left females to oviposit for further six days across three sets of vials (Figure 1d). Adult offspring eclosing from these vials were collected as virgins across multiple days and used to create the next generation. Population identity was blinded for all procedures after the initial population setup. See our supplementary methods for detailed procedures.

Every generation, we obtained an estimate of the frequency of polyandry for each of the twelve experimentally evolving populations as described in detail above and in the supplementary methods. We used tester males from the unrelated Chiricahua population, and allowed a minimum of 90 minutes of observation in each assay.

Statistical analyses

We used R version 3.4.2 (R Core Team, 2018) for all statistical analyses and figures, running linear mixed effects models (LMM) and generalised linear mixed effects models (GLMM) implemented in the *lme4* package version 1.1-14 (Bates *et al.*, 2015). We extracted effect sizes and p values from full models to avoid biasing effect sizes through the removal of non-significant terms (Forstmeier & Schielzeth, 2011). P values from LMMs were obtained from F-tests using the Kenward-Roger approximation for denominator degrees of freedom implemented in *lmerTest* (Kuznetsova *et al.*, 2016). We centred all covariates to a mean of zero to facilitate the interpretation of main effects in the presence of interactions and to aid model convergence. Age covariates were mean-centred, and order was centred and scaled to a standard deviation of one. We centred contrasts between two factors (*high* and *low* populations, *P* and *M* isolines) by coding factor levels as -0.5 and 0.5, respectively (Schielzeth, 2010). We calculated approximate 95%

confidence intervals (*CI*) for effect sizes as twice the standard error either side of the mean (Crawley, 2007).

We analysed the evolution of the frequency of polyandry using female remating as our binary response variable in a binomial GLMM. Our main interest was in how the frequency of polyandry changed over generations from the two respective starting frequencies, i.e. backgrounds (*low* versus *high*). Thus, our fixed effects were background, generation and their interaction. Generation was centred at the experimental evolution mid-point of four generations. We included as further fixed effects the age of the female and both males (first and second mate), as well as the order in the assay to control for potential variation arising from age variation and time available for mating in a given assay. To control for sources of non-independence between measurements and for stochastic day effects, we modelled random intercepts for female post-eclosion vial ID (4.7 \pm 1.3 females from the same post-eclosion vial were used in an assay), population replicate as well as assay day, and random slopes over the seven generations for each population replicate (Schielzeth & Forstmeier, 2009). We removed females (N = 74) for which we could not confirm fertilisation during their first mating through the presence of larvae in their oviposition vial.

Assays after experimental evolution

After seven generations of experimental evolution, we subjected all experimental populations to one generation of common garden breeding and used the offspring for our final assays described below. Because polyandry assays can be subject to substantial block effects, comparisons of absolute estimates of the frequency of polyandry cannot be made across assays conducted on different days. Thus, to make direct comparisons not only between experimentally evolved

replicate populations, but also between the ancestral isolines and the experimentally evolved populations, we simultaneously assayed flies from the twelve replicate evolved populations and from the 16 original ancestral isolines (see Nouhaud *et al.*, 2016).

Female remating latency

To refine our comparisons, here we used female latency to remating (Price *et al.*, 2008) as a more precise measure of polyandry that correlates with the proportion of females remating given one opportunity (Price *et al.*, 2008, 2011). All 12 populations and 16 isolines were simultaneously tested in each of two experimental blocks. Mating assays followed our general methods for remating assays described above, with the difference that here females were given a remating opportunity every day from two to five days after their first mating, or until they remated. Due to logistical limitations in obtaining several hundreds of virgin tester males for every mating day, we re-used some males for remating opportunities, such that our assays included some non-virgin tester males that had been sexually rested for at least two days. We found that female remating was not affected by mating status of tester males (data not shown).

Because data for remating latency were right-censored (23% of females did not remate in any of their four opportunities), we analysed remating analogous to death in survival models, using mixed effects cox models implemented in the *coxme* package (Therneau, 2015). We used days to remating as a right-censored response variable. As fixed effects, we included focal female background (two-levels: *P/high* and *M/low*), female age, age of the first male and order in the assay. Fixed effects were centred and scaled as described above. Female post-eclosion vial, nested within population replicate or isoline, as well as experimental block were included as random effects. We first ran separate models on ancestral isolines and evolved populations,

respectively. To ask whether populations had evolved polyandry levels different from their initial setup, we then simulated resampling of our setup of the 12 population replicates from the 16 ancestral isolines before experimental evolution, using *for* loops in R. We ran *coxme* models on 1000 simulated datasets to obtain a distribution of the inferred initial difference between *low* versus *high* polyandry population replicates, with the sample size reflecting our remating latency assay (see supplementary methods). We compared the observed difference between evolved *low* and *high* polyandry populations to that distribution under the null hypothesis that the difference in polyandry between the populations did not change during experimental evolution. Similarly, we compared the simulated populations (i.e. inferred remating latencies in the population replicates before experimental evolution) with the observed remating latencies of the experimentally evolved populations.

Remating inhibition by males

To investigate potential male effects on female remating, we assessed variation in the ability of males from the 12 populations and 16 isolines to induce a refractory period (i.e. male remating inhibition) in females from the tester (Chiricahua) population. We used variation in the proportion of tester females that remated with tester males four days after mating with focal males as our proxy for variation in remating inhibition by focal males. We conducted the experiment across two blocks and used the same methods as for our polyandry assays during experimental evolution. In the second block, we quantified reproductive output after the first mating to test for its association with remating inhibition (see ESM).

In this assay, higher tester female remating would indicate lower remating inhibition by focal males. Our main questions were whether our experimental evolution protocol had generally

changed male remating inhibition, whether experimental evolution under our *low* versus *high* polyandry regime had manifested in differences in males' ability to inhibit remating (Price *et al.*, 2010b), and if so, whether the difference already existed in the isolines used to initiate the populations. We used GLMMs with female remating as a binary response, and included focal male background, the ages of the female and both her (potential) mates as well as order in the assay as fixed effects. Random effects were female post-eclosion vial nested within experimental block and the genetic background (isoline/replicate population) of the focal first-to-mate male. Ancestral and evolved populations were compared in analogy to female remating latency, using resampling to simulate the experimental setup of the population replicates (see *Female remating latency*).

To explore a possible pre-existing genetic correlation between female mating behaviour and male remating inhibition, we first obtained predictions for isolines for both female remating latency and male remating inhibition. We used a linear model for remating latency and a generalised linear model for remating inhibition with isolate ID as well as age and order (centred and scaled) and block (centred) as fixed effects. Thus, we ignored variation between female post-eclosion vials, which was found to be very small in the previous mixed models (see Tables 2 and 3). To test for a correlation between female remating latency and male remating inhibition, we used linear regression on the predictions for the 16 isolines, backtransformed from the latent scale for male remating inhibition and weighted by the combined sample sizes of the female and male assays. We excluded evolved populations from this analysis to avoid pseudo-replication arising from repeated representation of isolate genotypes in the evolved population replicates.

Fecundity after experimental evolution

Finally, we measured fecundity of females that evolved in populations with relatively high versus relatively low levels of polyandry. We used the same methods as for our standardised polyandry assays, except that females were paired with males from their own replicate population. Females were subjected to different remating regimes to test for phenotypic effects of polyandry on fecundity. We randomly chose four to five females per population that were not given a remating opportunity (i.e., forced monandry), aspirating the male out of his vial before the female was introduced. The remaining females (12-15 per population) had one opportunity to remate four days after their initial mating. After their denied or realised remating opportunity, females oviposited for six days across two vials. We incubated vials under standard conditions and counted the total number of offspring eclosed nine days after the first eclosion in a given vial.

To explore variation in female fecundity, we pooled counts of eclosed offspring from the two vials in which females had oviposited for three days each after their second mating opportunity, thus matching the oviposition period used during experimental evolution. Our full LMM included female background (*low* versus *high*), remating regime (forced monandry, elected monandry and polyandry), their interaction, and age of the female and her first mate (both centred) as fixed effects. We included post-eclosion vial nested within replicate population as random effects.

Results

Experimental evolution of polyandry

The overall frequency of polyandry across all mating assays over seven generations was 34.1%, but there was substantial variation between generations and between replicate populations (Figure 2). Each generation, we aimed to test 35 females per population. However, failed first matings

(8%) mortality between the two assays (3%) and absence of larvae in the oviposition vial (2%) meant that we estimated the frequency of polyandry for each replicate population at every generation from an average of 30.5 females (N = 2559 across seven generations).

Inspection of our binomial GLMM on polyandry revealed that the interaction between generation and background was small and not significantly different from zero (effect size [approx. 95% CI] on the logit scale = 0.03 [-0.07;0.14]; $p = 0.517$; Table 1), meaning that there was neither evidence for convergence nor divergence of the frequency of polyandry between the populations with *high* and *low* polyandry backgrounds. There was a clear main effect of background indicating that polyandry was indeed lower in the *low* background (-0.30 [-0.52;-0.08]; $p = 0.006$) i.e., the population that had been set up with predominantly low polyandry genotypes. There was also a slight positive trend of generation showing a general increase in polyandry over time (0.06 [-0.02;0.13]; $p = 0.119$). The first male's age had a clear negative effect on remating, meaning that females mated to older males were less likely to remate four days later. The age of the female and of the second male had no significant impact on polyandry. The order in the assay showed a minor negative trend, with flies entering the assay later having a slightly lower probability of remating (Table 1).

Polyandry in isolines and after experimental evolution

We assessed latency to remating in females from each of the 12 populations and 16 isolines. Figure 3 illustrates differences between isolines and experimentally evolved populations, and between high polyandry and low polyandry isolines and populations, assigning females that did not remate a maximum remating latency of 6 days. In total, 156 pairs of virgin flies did not mate (total N = 894). Failed matings were heavily biased towards three of the four isolines that

originated from the Shaver Lake population (76–83% mating failure), resulting in small sample sizes for these isolines (N = 6–9 versus N = 18–36 for other lines). After removal of females that died before their first remating opportunity, our final sample size for remating latency was 734 females, of which 169 (isolines: 86 *M*, 33 *P*; populations: 30 *low*, 20 *high*) were right-censored, i.e., had not remated by day six. Not surprisingly, *M* isolines had a longer remating latency than *P* isolines (odds ratio for remating [approx. 95% *CI*]: 0.49 [0.27;0.92]; N = 419; p = 0.023; Table 2, Figure 3a & Figure S1). In our evolved population replicates, we found correspondingly that *low* populations had a longer latency to remating than *high* populations (odds ratio 0.72 [0.53;0.99]; N = 315; p = 0.037). Females initially mated to older males were slower to remate, female age did not matter, and females with a later order in the assay (i.e. less time allowed for remating) showed delayed remating, which was statistically significant in the population subset but not in the isoline subset (Table 2). The comparison of the observed evolved populations to the populations simulated based on resampling of isoline females revealed the observed difference between *low* and *high* population replicates (odds ratio) to be remarkably similar to that in the simulated datasets (odds ratio observed 0.72; simulated 0.71 [0.53;0.93]; p = 0.866). However, females from evolved population replicates generally remated faster than expected based on the simulated ancestral composition of population replicates (odds ratio 1.70 [1.47;1.95]; p < 0.001; Figure 3a).

Male influence on female remating?

Analogous to the assay on female latency to remating, failed mating trials between focal males and tester females were heavily biased towards three of the isolines originating from the Shaver Lake population (76-98% mating failure). Sample sizes for these isolines were consequently very small (N = 1-8 versus N = 19-33 for other isolines/populations; total N = 710).

There was no difference in the likelihood of tester female remating after mating with males from *M* versus *P* isolines (effect on logit scale 0.23 [-0.21;0.67]; N = 363; p = 0.301). Males from *low* polyandry population replicates showed a tendency to be less effective at reducing tester female remating relative to males from *high* polyandry populations, although this was marginally non-significant (effect on logit scale 0.43 [-0.02;0.89]; N = 347; p = 0.059). Male effects on female remating were not simply mediated through male effects on female reproductive output (see ESM). Additionally, there were effects of the age of females and both males on the probability of remating, with consistent effect signs but varying effect sizes between tests on isolines and evolved populations (Table 3). Generally, older females were more likely to remate, older first males reduced remating later on, and females were more likely to remate when presented with younger tester males. These results were robust to omitting pseudo-polyandrous females (i.e. females with no larvae in their oviposition vial), thus only focussing on fertilised females (N = 694).

The comparison of the observed evolved populations to the simulated populations based on resampling of remating inhibition by isoline males showed a minor trend for a greater difference between *high* and *low* population replicates after experimental evolution than expected based on the simulated initial population setup (observed 0.43; simulated 0.09 [-0.33;0.53]; p = 0.139). This was probably mainly driven by evolved *high* polyandry replicates (Figure 3), with males from evolved population replicates overall inhibiting female remating more efficiently than expected based on the simulated ancestral composition of population replicates (effect size for tester female remating on logit scale -0.20 [-0.41;0.02]; p < 0.033).

Finally, we found no evidence for a genetic correlation between female remating latency and male remating inhibition in our 16 original isolines. The correlation coefficient was positive but not significantly different from zero (0.05 [-0.02;0.12], $F_{1,14} = 2.17$, $p = 0.163$).

Fitness effects of polyandry?

We pooled counts of offspring eclosing from the two vials in which individual females (N = 226) from evolved population replicates had oviposited over a combined period of six days. There was no significant influence of any of the variables included in the full model, except for significant variation between population replicates ($p = 0.024$; Table S2 & Figure S5). Thus, there was no significant difference in fecundity between females from a *low* versus *high* polyandry background, nor was there an effect of mating phenotype, i.e. of whether the opportunity to remate was experimentally prevented, or refused or accepted by the female. Finally, there was no interaction between genetic background and mating phenotype.

Discussion

What drives and maintains variation in polyandry between and within populations is poorly understood. Here, we used naturally occurring genetic variation in polyandry and investigated whether experimental populations that started with a high versus low initial frequency of polyandry would show evidence for balancing or directional selection, or evolve neutrally. We found that the frequency of polyandry remained remarkably stable over time, remaining relatively low in populations with an initially lower frequency, and relatively high in populations with an initially higher frequency of polyandry. Thus, we found no clear evidence for directional or balancing selection on polyandry. Despite starting with a substantial difference in polyandry in the *high* versus *low* polyandry populations, remarkably we found no difference in fecundity between females from these populations, and no significant change in the difference between these populations over time which would have indicated fitness consequences of polyandry. Data on male inhibition of female remating showed a trend consistent with previous findings that males evolve enhanced remating inhibition in response to elevated female remating (Price *et al.*, 2010b). This indicates ongoing evolution in males in our experimental populations, but the absence of a correlation between polyandry and male remating inhibition in ancestral isolines suggests selection can operate independently on male and female traits. Overall, our findings are consistent with genetic control over female remating behaviour, but indicate that polyandry does not have strong fitness consequences under these conditions.

Neutral experimental evolution of polyandry?

Populations initiated with many polyandrous females maintained a higher frequency of polyandry than did populations initiated with relatively fewer polyandrous females (Figure 2). Our assay on female remating latency after one generation of common garden breeding allowed us to directly

compare experimentally evolved populations with ancestral isolines, and confirmed genetic differences between the *high* and *low* polyandry populations. Importantly, using tester males that had not co-evolved with females allowed us to assess selection on polyandry independent of selection acting on males. There was only a very minor tendency for populations to be more similar after experimental evolution than when they were initially founded; we found no clear evidence for convergence towards a common polyandry frequency. We experimentally evolved populations for only seven generations, admittedly limiting our power to detect convergence. Indeed, the best model estimates based on assays during experimental evolution (Table 1) suggested that high and low populations might indeed have converged after a few more generations. However, in our remating latency assays where we tested experimentally evolved and ancestral isolines simultaneously—arguably a more accurate comparison—the observed difference between high and low populations after seven generations of experimental was only very marginally smaller than expected based on our resampling simulation of the initial isoline composition (odds ratios 0.72 and 0.71, respectively), suggesting populations would only fully converge after more than 100 generations. This was in contrast with the trend observed for male remating inhibition (Figure 3b), which suggested that a rapid response was possible despite the limited timeframe. Rather than convergence in polyandry levels, the patterns from the female remating assays both during (Figure 2) and after experimental evolution (Figure 3a) suggested a parallel increase in polyandry in the evolved populations relative to the ancestral isolines. This increase was visible as a trend across seven assays during experimental evolution and reached statistical significance only in the direct comparison between ancestral and evolved females. The small number of matings between individuals from the Shaver Lake isolines and tester individuals from the Chiricahua population weakened our direct comparison between isolines and evolved populations. Generally, Shaver Lake flies appeared to have reduced compatibility with

flies from the other populations (see ESM for more details). However, Shaver Lake isolines represented average polyandry genotypes both within the P and M isoline groups (cf. Figure 1b) and our balanced design would have prevented a systematic bias in polyandry arising from selective disappearance of Shaver Lake genotypes. The observed increase in polyandry could indicate a selective advantage of polyandry alleles in all populations due to a superior fitness of highly polyandrous genotypes. Under this scenario however, selection should favour the high polyandry alleles both in *high* and *low* polyandry populations, and the populations to consequently converge towards a high frequency of polyandry. Alternatively, the increase in polyandry could be a manifestation of condition-dependent polyandry. Experimentally evolved females have high heterozygosity and might therefore have higher fecundity and remate more than highly inbred isoline females, for example due to reduced costs of mating (Perry *et al.*, 2009) or higher demands for sperm numbers. Whether the observed increase in polyandry reflects a change in the frequency of high polyandry alleles or represents a phenotypically plastic response that is independent of allele frequency changes is currently unknown. Although we acknowledge that the duration of our experiment meant limited power to detect convergence, we believe that the phenotypic plasticity explanation is more consistent with our observation that the increase in polyandry was parallel in both the *low* and *high* polyandry populations.

Experimentally investigating the evolution of polyandry without manipulating access to mates is challenging, because monandrous females can typically not be forced to mate multiply (but see Arnqvist & Andrés, 2006; King & Bressac, 2010). As a consequence, the majority of evidence for the benefits of polyandry has come from experiments where naturally polyandrous females were denied the possibility for multiple mating. While experimentally manipulating sex ratio may offer much insight into how selection from sperm competition acts on males, enforcing a

particular mating frequency on females may reveal little about why there is so much variation in female mating strategies (Taylor *et al.*, 2014). Our design allowed us to initiate replicate populations with substantial differences in the average frequency of polyandry without altering the sex ratio or manipulating female access to mates, allowing for a more realistic competition between different female strategies. To our knowledge, only one previous study has employed genetic variation in female mating behaviour to manipulate sexual selection. Using a sex peptide receptor knockout to render females hyper-promiscuous, the study highlighted that purely manipulating the mating frequency may have consequences for sexual selection that are different from those of sex ratio manipulations (Perry *et al.*, 2016). Genetic variation in polyandry is potentially very widespread (Taylor *et al.*, 2014), so utilising it offers an invaluable experimental tool for improving our understanding of the evolution of polyandry in semi-natural conditions.

Consequences of polyandry for males

Consistent with previous findings in *D. pseudoobscura*, we found that males had some effect on female remating behaviour. Across all experiments, age of the first male had a consistently negative effect on female remating (Tables 1-3). This effect could have been driven by age-dependent variation in male accessory gland size (Ruhmann *et al.*, 2016) and/or by older males allocating larger ejaculates during mating (Avent *et al.*, 2008). We cannot tell whether reduced remating after mating with older males represents male suppression of female remating decisions or adaptive female mate choice, given that females can benefit directly from mating with older males (Avent *et al.*, 2008; Verspoor *et al.*, 2015). However, we found no evidence for a preference for older males during rematings (in fact, there was a trend for the opposite effect), thus favoring the idea that reduced remating propensity reflects a male effect. Indeed, our results on experimentally evolved males were in agreement with previous results showing that more

frequent remating by females selects for improved remating inhibition in males (Crudgington *et al.*, 2005; Price *et al.*, 2010b; Figure 3b). Our direct comparison between isolines and evolved populations indicated that the tendency for higher remating inhibition by males that had experimentally evolved with high polyandry was not driven by a pre-existing genetic correlation between polyandry and male remating inhibition. In support of this interpretation, there was no difference in remating inhibition in *M* versus *P* isolines, and no correlation between female remating latency and male remating inhibition across the 16 isolines (Figure S2).

Polyandry does not affect fecundity

After seven generations of experimental evolution and one generation of common garden breeding, we found no evidence that genetic polyandry was associated with higher fecundity. Although we found variation between evolved populations (Figure S5), this variation did not covary with polyandry levels, suggesting polyandry does not evolve simply through a genetic correlation between polyandry and fecundity. Indeed, early life fecundity was neither linked to genetic variation in polyandry nor to phenotypic variation in polyandry (Table S2). Moreover, we found no evidence that females evolving with higher polyandry levels became dependent on polyandry, which would have manifested in increased costs of forced monandry. In combination, this means that the overall increase in polyandry after experimental evolution (see above) is unlikely to have been caused by a direct or correlated response to selection on fecundity. Unlike our fecundity assay after experimental evolution which focused on the effect of polyandry on a single fitness measure in isolated females, tracking polyandry during experimental evolution was an integrated measure of the costs and benefits of polyandry. Thus, potential costs of polyandry manifesting through injury, sexually transmitted diseases or foregone foraging opportunities would have operated simultaneously with potential direct benefits of fertility assurance, and

indirect genetic effects of good genes or sexy sperm (Arnqvist & Nilsson, 2000; Jennions & Petrie, 2000). The absence of clear changes in polyandry levels in our populations indicates that these costs and benefits are of small effect or that the costs and benefits are balanced, at least under our laboratory conditions.

What maintains genetic variation in polyandry?

Despite a considerable body of work on the costs and benefits of polyandry, and many empirical demonstrations of fitness effects, genetic variation in and experimental evolution of polyandry, what drives and maintains variation in polyandry between and within wild populations remains elusive. Given there are many factors that can influence multiple mating, including stochastic variation between females, phenotypic variation in polyandry rather than monandry may well be the null model (Gowaty, 2013; Kokko & Mappes, 2013). However, if polyandry is adaptively flexible, why should genetic variation in polyandry persist (Gowaty, 2013)? One potential answer is fluctuating selection imposed by fluctuating environmental conditions, which can favour the maintenance of alternative polyandry genotypes in butterflies (Wedell *et al.*, 2002; Välimäki *et al.*, 2008). Or perhaps genetic variation is simply the product of mutation-selection balance? Indeed, if polyandry is a highly polygenic trait that is largely selectively neutral in many females, then we might expect substantial genetic variation arising through random mutation that is not counteracted by strong selection. If so, then we might expect to find genetic variation predominantly in species and populations where polyandry has little effect on reproductive fitness. To understand the evolution of polyandry, we need to better understand the genetic basis of polyandry and the evolutionary processes that increase and decrease genetic variation in polyandry.

Summary

In this study, we confirmed strong genetic control over remating decisions in female *D. pseudoobscura*. Populations initiated with a high versus low frequency of alleles conferring a predisposition for polyandry maintained their genetic differences in polyandry over time. We found no evidence for balancing selection, and little evidence for positive selection on polyandry. Author contributions:

NW and TARP conceived the study; AS, TARP and NW designed the study; AS collected the data with help from LMT, MW and KO; AS performed the analyses; AS wrote the manuscript, with input from NW, TARP and LMT; all authors approved the manuscript.

Acknowledgements:

We thank Ali Skeats and Michelle Taylor for establishing isofemale isogenic lines. This work was supported by Swiss National Science Foundation fellowships awarded to AS (Grants P2ZHP3_164990 & P300PA_177906) and a Marie Skłodowska-Curie Fellowship to KO (Grant 746169 – IMMUNFUNC). We are grateful to editors and reviewers for constructive feedback on previous versions of this manuscript. The authors declare no conflict of interest.

Data accessibility:

All data supporting this manuscript will be deposited in dryad upon acceptance.

References

Arbuthnott, D., Dutton, E.M., Agrawal, A.F. & Rundle, H.D. 2014. The ecology of sexual conflict: Ecologically dependent parallel evolution of male harm and female resistance in

- Drosophila melanogaster*. *Ecol. Lett.* **17**: 221–228.
- Arnqvist, G. & Andrés, J.A. 2006. The effects of experimentally induced polyandry on female reproduction in a monandrous mating system. *Ethology* **112**: 748–756.
- Arnqvist, G. & Nilsson, T. 2000. The evolution of polyandry: multiple mating and female fitness in insects. *Anim. Behav.* **60**: 145–164.
- Arnqvist, G. & Rowe, L. 2005. *Sexual conflict*. Princeton University Press.
- Avent, T.D., Price, T.A.R. & Wedell, N. 2008. Age-based female preference in the fruit fly *Drosophila pseudoobscura*. *Anim. Behav.* **75**: 1413–1421.
- Ayala, F.J. & Campbell, C.A. 1974. Frequency-dependent selection. *Annu. Rev. Ecol. Syst.* **5**: 115–138.
- Balloux, F. & Lehmann, L. 2003. Random mating with a finite number of matings. *Genetics* **165**: 2313–2315.
- Bates, D., Maechler, M., Bolker, B. & Walker, S. 2015. Fitting linear mixed-effects models using lme4. *J. Stat. Softw.* **67**: 1–48.
- Birkhead, T.R. & Moller, A.P. 1998. *Sperm competition and sexual selection*. Academic Press, London, UK.
- Brisson, D. 2018. Negative frequency-dependent selection is frequently confounding. *Front. Ecol. Evol.* **6**: 1–9.
- Burton-Chellew, M.N., Beukeboom, L.W., West, S.A. & Shuker, D.M. 2007. Laboratory evolution of polyandry in the parasitoid wasp *Nasonia vitripennis*. *Anim. Behav.* **74**: 1147–1154.
- Candolin, U. & Heuschele, J. 2008. Is sexual selection beneficial during adaptation to environmental change? *Trends Ecol. Evol.* **23**: 446–452.
- Chapman, T., Arnqvist, G., Bangham, J. & Rowe, L. 2003. Sexual conflict. *Trends Ecol. Evol.*

18: 41–47.

Clarke, B. 1979. The evolution of genetic diversity. *Proc. R. Soc. London B* **205**: 453–474.

Crawley, M.J. 2007. *The R Book*. John Wiley & Sons, Chichester.

Crudgington, H.S., Beckerman, A.P., Brüstle, L., Green, K. & Snook, R.R. 2005. Experimental removal and elevation of sexual selection: does sexual selection generate manipulative males and resistant females? *Am. Nat.* **165**: S72–S87.

Crudgington, H.S., Fellows, S., Badcock, N.S. & Snook, R.R. 2009. Experimental manipulation of sexual selection promotes greater male mating capacity but does not alter sperm investment. *Evolution* **63**: 926–938.

Crudgington, H.S., Fellows, S. & Snook, R.R. 2010. Increased opportunity for sexual conflict promotes harmful males with elevated courtship frequencies. *J. Evol. Biol.* **23**: 440–446.

David, J.R., Gibert, P., Legout, H., Pétavy, G., Capy, P. & Moreteau, B. 2005. Isofemale lines in *Drosophila*: an empirical approach to quantitative trait analysis in natural populations. *Heredity (Edinb)*. **94**: 3–12.

Demont, M., Grazer, V.M., Michalczyk, Ł., Millard, A.L., Sbilordo, S.H., Emerson, B.C., *et al.* 2014. Experimental removal of sexual selection reveals adaptations to polyandry in both sexes. *Evol. Biol.* **41**: 62–70.

Emlen, S.T. & Oring, L.W. 1977. Ecology, sexual selection, and the evolution of mating systems. *Science (80-)*. **197**: 215–223.

Evans, J. & Magurran, A. 2000. Multiple benefits of multiple mating in guppies. *Proc. Natl. Acad. Sci. U. S. A.* **97**: 10074–10076.

Forstmeier, W. & Schielzeth, H. 2011. Cryptic multiple hypotheses testing in linear models: Overestimated effect sizes and the winner’s curse. *Behav. Ecol. Sociobiol.* **65**: 47–55.

Gavrilets, S. 2014. Is sexual conflict an “engine of speciation”? *Cold Spring Harb. Perspect.*

Biol. **6**: a017723.

Gowaty, P.A. 2013. Adaptively flexible polyandry. *Anim. Behav.* **86**: 877–884.

Gowaty, P.A. & Hubbell, S.P. 2009. Reproductive decisions under ecological constraints: it's about time. *Proc. Natl. Acad. Sci. U. S. A.* **106 Suppl**: 10017–10024.

Gowaty, P.A., Kim, Y.-K., Rawlings, J. & Anderson, W.W. 2010. Polyandry increases offspring viability and mother productivity but does not decrease mother survival in *Drosophila pseudoobscura*. *Proc. Natl. Acad. Sci. U. S. A.* **107**: 13771–13776.

Harano, T. & Miyatake, T. 2005. Heritable variation in polyandry in *Callosobruchus chinensis*. *Anim. Behav.* **70**: 299–304.

Herrera, P., Taylor, M.L., Skeats, A., Price, T.A.R. & Wedell, N. 2014. Can patterns of chromosome inversions in *Drosophila pseudoobscura* predict polyandry across a geographical cline? *Ecol. Evol.* **4**: 3072–3081.

Hollis, B., Houle, D. & Kawecki, T.J. 2016. Evolution of reduced post-copulatory molecular interactions in *Drosophila* populations lacking sperm competition. *J. Evol. Biol.* **29**: 77–85.

Hollis, B., Houle, D., Yan, Z., Kawecki, T.J. & Keller, L. 2014. Evolution under monogamy feminizes gene expression in *Drosophila melanogaster*. *Nat. Commun.* **5**: 3482. Nature Publishing Group.

Holman, L. & Kokko, H. 2013. The consequences of polyandry for population viability, extinction risk and conservation. *Philos. Trans. R. Soc. B Biol. Sci.* **368**: 20120053.

Jennions, M.D. & Petrie, M. 2000. Why do females mate multiply? A review of the genetic benefits. *Biol. Rev. Camb. Philos. Soc.* **75**: 21–64.

King, B.H. & Bressac, C. 2010. No fitness consequence of experimentally induced polyandry in a monandrous wasp. *Behaviour* **147**: 85–102.

Kokko, H. & Mappes, J. 2013. Multiple mating by females is a natural outcome of a null model

- of mate encounters. *Entomol. Exp. Appl.* **146**: 26–37.
- Kuijper, B., Stewart, A.D. & Rice, W.R. 2006. The cost of mating rises nonlinearly with copulation frequency in a laboratory population of *Drosophila melanogaster*. *J. Evol. Biol.* **19**: 1795–1802.
- Kuriwada, T., Kumano, N., Shiromoto, K., Haraguchi, D., Matsuyama, T. & Kohama, T. 2014. Female preference did not evolve under laboratory conditions in the solanaceous fruit fly *Bactrocera latifrons*. *Int. J. Pest Manag.* **60**: 160–165.
- Kuznetsova, A., Brockhoff, P.B. & Christensen, R.H.B. 2016. lmerTest: 2.0-33., Tests in Linear Mixed Effects Models. R package version 2.0-33.
- Lumley, A.J., Michalczyk, Ł., Kitson, J.J., Spurgin, L.G., Morrison, C.A., Godwin, J.L., *et al.* 2015. Sexual selection protects against extinction. *Nature* **522**: 470–473.
- Mank, J.E., Wedell, N. & Hosken, D.J. 2013. Polyandry and sex-specific gene expression. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **368**: 20120047.
- Markow, T.A. 2011. “Cost” of virginity in wild *Drosophila melanogaster* females. *Ecol. Evol.* **1**: 596–600.
- Martin, O.Y., Hosken, D.J. & Ward, P.I. 2004. Post-copulatory sexual selection and female fitness in *Scathophaga stercoraria*. *Proc. R. Soc. B Biol. Sci.* **271**: 353–359.
- Michalczyk, Ł., Millard, A.L., Martin, O.Y., Lumley, A.J., Emerson, B.C., Chapman, T., *et al.* 2011. Inbreeding promotes female promiscuity. *Science* **333**: 1739–1742.
- Neff, B.D. & Svensson, E.I. 2013. Polyandry and alternative mating tactics. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **368**: 20120045.
- Newcomer, S.D., Zeh, J. a & Zeh, D.W. 1999. Genetic benefits enhance the reproductive success of polyandrous females. *Proc. Natl. Acad. Sci. U. S. A.* **96**: 10236–10241.
- Nouhaud, P., Tobler, R., Nolte, V. & Schlötterer, C. 2016. Ancestral population reconstitution

- from isofemale lines as a tool for experimental evolution. *Ecol. Evol.* **6**: 7169–7175.
- Parker, G. 2006. Sexual conflict over mating and fertilization: an overview. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **361**: 235–59.
- Parker, G.A. 1970. Sperm competition and its evolutionary consequences in the insects. *Biol. Rev.* **45**: 525–567.
- Perry, J.C., Joag, R., Hosken, D.J., Wedell, N., Radwan, J. & Wigby, S. 2016. Experimental evolution under hyper-promiscuity in *Drosophila melanogaster*. *BMC Evol. Biol.* **16**: 1–14.
- Perry, J.C., Sharpe, D.M.T. & Rowe, L. 2009. Condition-dependent female remating resistance generates sexual selection on male size in a ladybird beetle. *Anim. Behav.* **77**: 743–748.
- Price, T.A.R., Hodgson, D.J., Lewis, Z., Hurst, G.D.D. & Wedell, N. 2008. Selfish genetic elements promote polyandry in a fly. *Science* **322**: 1241–1243.
- Price, T.A.R., Hoskyns, R.C., Rapley, H., Evans, J.C. & Wedell, N. 2012. No evidence that temperature-related fertility differences influence the distribution of a selfish genetic element. *Funct. Ecol.* **26**: 657–665.
- Price, T.A.R., Hurst, G.D.D. & Wedell, N. 2010a. Polyandry prevents extinction. *Curr. Biol.* **20**: 471–475.
- Price, T.A.R., Lewis, Z., Smith, D.T., Hurst, G.D.D. & Wedell, N. 2011. Remating in the laboratory reflects rates of polyandry in the wild. *Anim. Behav.* **82**: 1381–1386.
- Price, T.A.R., Lewis, Z., Smith, D.T., Hurst, G.D.D. & Wedell, N. 2010b. Sex ratio drive promotes sexual conflict and sexual coevolution in the fly *Drosophila pseudoobscura*. *Evolution* **64**: 1504–1509.
- Price, T., Bretman, A., Gradilla, A.C., Reger, J., Taylor, M.L., Giraldo-Perez, P., *et al.* 2014. Does polyandry control population sex ratio via regulation of a selfish gene? *Proc. R. Soc. B Biol. Sci.* **281**: 20133259.

- R Core Team. 2018. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.
- Ruhmann, H., Wensing, K.U., Neuhalfen, N., Specker, J.-H. & Fricke, C. 2016. Early reproductive success in *Drosophila* males is dependent on maturity of the accessory gland. *Behav. Ecol.* **27**: 1859–1868.
- Schielzeth, H. 2010. Simple means to improve the interpretability of regression coefficients. *Methods Ecol. Evol.* **1**: 103–113.
- Schielzeth, H. & Forstmeier, W. 2009. Conclusions beyond support: overconfident estimates in mixed models. *Behav. Ecol.* **20**: 416–420.
- Sgrò, C.M., Chapman, T. & Partridge, L. 1998. Sex-specific selection on time to remate in *Drosophila melanogaster*. *Anim. Behav.* **56**: 1267–1278.
- Shuker, D.M., Phillimore, A.J., Burton-Chellew, M.N., Hodge, S.E. & West, S.A. 2007. The quantitative genetic basis of polyandry in the parasitoid wasp, *Nasonia vitripennis*. *Heredity (Edinb)*. **98**: 69–73.
- Simmons, L.W. 2001. *Sperm competition and its evolutionary consequences in the insects*. Princeton University Press, Princeton, New Jersey.
- Simmons, L.W. 2003. The evolution of polyandry: patterns of genotypic variation in female mating frequency, male fertilization success and a test of the sexy-sperm hypothesis. **16**: 624–634.
- Sinervo, B. & Lively, C.M. 1996. The rock–paper–scissors game and the evolution of alternative male strategies. *Nature* **380**: 240–243.
- Slatyer, R. a, Mautz, B.S., Backwell, P.R.Y. & Jennions, M.D. 2012. Estimating genetic benefits of polyandry from experimental studies: a meta-analysis. *Biol. Rev. Camb. Philos. Soc.* **87**: 1–33.

- Snook, R.R. & So, Y.K. 2000. Associations between female remating behavior, oogenesis and oviposition in *Drosophila melanogaster* and *Drosophila pseudoobscura*. *J. Insect Physiol.* **46**: 1489–1496.
- Solymar, B.D. & Cade, W.H. 1990. Heritable variation for female mating frequency in field crickets, *Gryllus integer*. *Behav. Ecol. Sociobiol.* **26**: 73–76.
- Svensson, E.I., Abbott, J. & Hardling, R. 2005. Female polymorphism, frequency dependence, and rapid evolutionary dynamics in natural populations. *Am. Nat.* **165**: 567–576.
- Svensson, E.I. & Råberg, L. 2010. Resistance and tolerance in animal enemy-victim coevolution. *Trends Ecol. Evol.* **25**: 267–274.
- Takahashi, Y. & Kawata, M. 2013. A comprehensive test for negative frequency-dependent selection. *Popul. Ecol.* **55**: 499–509.
- Taylor, M.L., Price, T.A.R., Skeats, A. & Wedell, N. 2016. Temperature can shape a cline in polyandry, but only genetic variation can sustain it over time. *Behav. Ecol.* **27**: 462–469.
- Taylor, M.L., Price, T.A.R. & Wedell, N. 2014. Polyandry in nature: a global analysis. *Trends Ecol. Evol.* **29**: 376–383.
- Therneau, T. 2015. coxme: Mixed Effects Cox Models. R package version 2.2-5.
<https://CRAN.R-project.org/package=coxme>.
- Torres-Vila, L., Gragera, J., Rodríguez-Molina, M. & Stockel, J. 2002. Heritable variation for female remating in *Lobesia botrana*, a usually monandrous moth. *Anim. Behav.* **64**: 899–907.
- Torres-Vila, L.M. 2013. Polyandry-fecundity relationship in insects: Methodological and conceptual problems. *J. Evol. Biol.* **26**: 325–334.
- Torres-Vila, L.M., Rodríguez-Molina, M.C., Gragera, J. & Bielza-Lino, P. 2001. Polyandry in Lepidoptera: a heritable trait in *Spodoptera exigua* Hübner. *Heredity (Edinb)*. **86**: 177–183.

- Travers, L.M., Simmons, L.W. & Garcia-Gonzalez, F. 2016. Additive genetic variance in polyandry enables its evolution, but polyandry is unlikely to evolve through sexy or good sperm processes. *J. Evol. Biol.* **29**: 916–928.
- Turner, M.E. & Anderson, W.W. 1983. Multiple mating and female fitness in *Drosophila pseudoobscura*. *Evolution* **37**: 714–723.
- Välimäki, P., Kivelä, S.M., Jääskeläinen, L., Kaitala, A., Kaitala, V. & Oksanen, J. 2008. Divergent timing of egg-laying may maintain life history polymorphism in potentially multivoltine insects in seasonal environments. *J. Evol. Biol.* **21**: 1711–1723.
- Verspoor, R., Cuss, M. & Price, T.A.R. 2015. Age-based mate choice in the monandrous fruit fly *Drosophila subobscura*. *Anim. Behav.* **102**: 199–207. Elsevier Ltd.
- Wedell, N. 2001. Female remating in butterflies: Interaction between female genotype and nonfertile sperm. *J. Evol. Biol.* **14**: 746–754.
- Wedell, N., Wiklund, C. & Cook, P.A. 2002. Monandry and polyandry as alternative lifestyles in a butterfly. *Behav. Ecol.* **13**: 450–455.
- Wigby, S. & Chapman, T. 2004. Female resistance to male harm evolves in response to manipulation of sexual conflict. *Evolution* **58**: 1028–1037.
- Zeh, J.A. & Zeh, D.W. 1996. The evolution of polyandry I: intragenomic conflict and genetic incompatibility. *Proc. R. Soc. B Biol. Sci.* **263**: 1711–1717.

Figure legends:

Figure 1: Schematic overview of the experimental evolution setup (see main text for details). a) Establishing isofemale isogenic lines (isolines) from three US populations in Lewistown, Montana (green), Show Low, Arizona (light purple), and Shaver Lake, California (dark purple); b) selecting isolines with higher (P) and lower (M) than average levels of polyandry (selected lines are highlighted with squares and thicker lines; Table S1); c) founding populations with females (and males, not shown here) from predominantly low polyandry isolines (80% from *M* isolines = *low polyandry*) or predominantly high polyandry isolines (80% from *P* isolines = *high polyandry*). d) Experimental procedures during experimental evolution: females and males were allowed to interact freely for four days, after which males were removed and females were left to oviposit for another six days. The resulting offspring were used to initiate the next generation and additional daughters were collected for polyandry assays.

Figure 2: Experimental evolution of polyandry. The proportion of females that remated was tracked in twelve independent populations over seven generations (thin solid lines). Populations were initially set up with a high (blue) versus low (orange) relative representation of isolines with higher than average polyandry levels. For illustration, means (circles connected by dashed lines) and standard errors (vertical bars) were calculated across the six replicates within a background for each generation. Thick solid lines show the model predictions from a GLMM on polyandry in the two backgrounds across generations, with other fixed effects mean-centred (Table 1). Filled circles at generation zero indicate the initial frequency of polyandry in the two backgrounds based on preliminary assays (Figure 1b & Table S1). Our results indicated that the two backgrounds differed in their frequency of polyandry, and that this did not change over the course

of the experiment. Although not significant, the main effect of generation and its interaction with background are retained here for illustrative purposes.

Figure 3: a) Female latency to remate with tester males and b) male ability to inhibit tester female remating in ancestral isolines and after seven generations of experimental evolution. Shown are means (circles, with area proportional to sample size) for *P/high* (blue) and *M/low* (orange) isolines and evolved populations, respectively. Squares and bars show model predictions and 95% *CI*. Our main analyses on remating latency were based on *coxme* models (see Fig S1), but for illustrative purposes, for a) here we use predictions from LMMs on remating latency (assigning females that did not mate a maximum of 6 days), with fixed effects mean-centred. Diamonds represent predictions for evolved populations based on isoline means and accounting for the relative initial representation of isolines in *high* and *low* polyandry populations. Note that in a) higher polyandry means a shorter latency and in b) stronger remating inhibition means a lower proportion of tester females remating. Further note that sample sizes for three isolines were very small due to a low incidence of mating between individuals from these isolines and tester flies (see discussion).

Figures:

Figure 1:




Figure 2:




Figure3:




Table 1: Full model summary for experimental evolution of polyandry. Coefficients, standard errors, test statistics and variance components are taken from a GLMM on female remating (binary response) and are consequently on the logit scale. Continuous and factorial covariates were centred and scaled as described in the main text, such that the global intercept describes the prediction for the mid-point for all covariates. Effects associated with a p value smaller than 0.05 are highlighted in bold.

<i>Polyandry exp. evolution (N = 2517)</i>	glmer (logit scale)			
Fixed effects	Coef	se (coef)	z	p
Intercept	-0.690	0.072	-9.64	<0.001
female age (centred)	0.048	0.038	1.27	0.204
first male age (centred)	-0.199	0.053	-3.78	<0.001
second male age (centred)	0.039	0.027	1.45	0.146
order (centred & scaled)	-0.075	0.046	-1.63	0.103
generation (centred)	0.055	0.036	1.56	0.119
background (centred; <i>low v high</i>)	-0.302	0.111	-2.73	0.006
generation:background	0.035	0.054	0.65	0.517
Random effects	Var	SD		
Post-eclosion vial (545 levels)	<0.001	<0.001		
Replicate (12 levels)	0.117	0.342		
Generation:replicate (12 random slopes)	0.003	0.056		
Assay day (7 levels)	0.014	0.120		

Table 2: Full model summaries for female remating latency of the 16 ancestral isolines and the 12 replicate populations after experimental evolution. Remating latency was analysed analogous to survival using the *coxme* function, with females that did not remate entered as right-censored data points. Continuous and factorial covariates were centred as described in the main text. Effects associated with a p value smaller than 0.05 are highlighted in bold.

<i>Latency to remating</i>	Isoline females (N = 419)				Evolved females (N = 315)			
Fixed effects (<i>coxme</i>)	coef	se (coef)	z	p	coef	se (coef)	z	p
female age (centred)	0.004	0.047	0.08	0.930	0.015	0.046	0.32	0.750
first male age (centred)	-0.164	0.053	-3.10	0.002	-0.144	0.056	-2.58	0.010
order (centred & scaled)	-0.075	0.928	-1.10	0.270	-0.166	0.065	-2.54	0.011
background (centred; <i>low v high</i>)	-0.704	0.495	-2.28	0.023	-0.323	0.155	-2.09	0.037
Random effects	Var	SD			Var	SD		
Housing vial	0.058	0.242			0.045	0.211		
Isoline/Population	0.296	0.544			0.139	0.373		
Block (2 levels)	0.004	0.060			<0.001	0.019		

Table 3: Full model summary for tester female remating after mating to males from the 16 ancestral isolines and the 12 replicate populations after experimental evolution. Coefficients, standard errors, test statistics and variance components are taken from GLMMs on tester female remating (binary response) and are consequently on the logit scale. Continuous and factorial covariates were centred and scaled as described in the main text. Effects associated with a p value smaller than 0.05 are highlighted in bold.

<i>Tester female remating</i>	Isoline males (N = 363)				Evolved males (N = 347)			
	coef	se (coef)	z	p	coef	se (coef)	z	p
Fixed effects (binomial GLMM)								
Intercept	-0.117	0.115	-1.01	0.312	-0.301	0.119	-2.54	0.011
female age (centred)	0.272	0.111	2.44	0.015	0.185	0.101	1.82	0.069
first male age (centred)	-0.182	0.088	-2.08	0.038	-0.104	0.085	-1.23	0.218
second male age (centred)	-0.270	0.139	-1.94	0.052	-0.260	0.157	-1.66	0.097
order (centred & scaled)	0.129	0.127	1.02	0.307	-0.155	0.147	-1.05	0.293
background (centred; <i>low v high</i>)	0.228	0.220	1.04	0.301	0.434	0.229	1.89	0.059
Random effects	Var	SD			Var	SD		
Tester female housing vial	0.093	0.305			0.062	0.120		
Male isolate/population	<0.001	<0.001			0.002	0.041		
Block (2 levels)	<0.001	<0.001			<0.001	<0.001		

Electronic Supplementary material

No selection for change in polyandry under experimental evolution

Andreas Sutter^{1,3,@}, Laura M. Travers^{1,3}, Melanie Weedon¹, Keiko Oku¹, Thomas A. R. Price², & Nina Wedell¹

¹ Centre for Ecology & Conservation, College of Life and Environmental Sciences, University of Exeter, Cornwall Campus, TR10 9FE, Penryn, UK.

² Institute of Integrative Biology, University of Liverpool, Crown Street, Liverpool, L69 7ZB, UK

³ Current address: Centre for Ecology, Evolution and Conservation, School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich, NR4 7TJ, UK

@ a.sutter@uea.ac.uk

Contents:

- Supplementary methods
- Supplementary results and discussion
- Supplementary tables
 - *Table S1*: Preliminary polyandry assays
 - *Table S2*: Model summary for female fecundity after experimental evolution
- Supplementary figures
 - *Figure S1*: Remating latency of isolines and evolved populations
 - *Figure S2*: No intersexual genetic correlation among isolines
 - *Figure S3*: Tester female fecundity and remating propensity
 - *Figure S4*: Male effects on tester female fecundity
 - *Figure S5*: Female fecundity after experimental evolution

Supplementary methods:

Fly stocks and maintenance

We collected wild female *Drosophila pseudoobscura* from three populations (Lewistown, Montana, 47°03'N, 109°28'W, in 2008; Show Low, Arizona, 34°16'N, 110°00'W, in 2008 and 2012; Shaver Lake, California, 35°06'N, 119°19'W, in 2015). To establish isofemale isogenic lines, we took offspring from each wild caught female and full-sib inbred them each generation for a minimum of 15 generations. During the establishment of these isolines, we let pairs of siblings mate and oviposit freely, and discarded them before any new eclosions of offspring to maintain non-overlapping generations. We replicated single sibling pairs three times for each genotype to prevent losses. This results in isolines that are as fully homozygous as possible, so that flies within an isolate are effectively genetically identical (David *et al.*, 2005). Flies were maintained under a 14:10 light: dark cycle at 23°C in standard *Drosophila* food vials (75mm in height by 25mm in width) containing a porridge medium (12.5g agar, 25g yeast, 75g oats, 105g sugar, 2.5g nipagin dissolved in EtOH and 5.6mL propionic acid in 1L of water). The experiments reported here were conducted in 2016 and 2017.

Experimental evolution

Population setup and maintenance

Founder individuals in the twelve replicate populations were 6.1 ± 1.3 days old (mean \pm SD). The populations were maintained in 3.5L plastic tubs (250 x 140 x 100mm) within a single incubator under a 14:10 light: dark cycle at 23°C and fed our standard porridge medium (see above). On day one, we introduced the flies into the tubs using aspiration to avoid disruption of courtship and mating behaviour by CO₂ anaesthesia (Barron, 2000). Flies were then left to mate freely for four days, after which we removed males under light CO₂ anaesthesia. Females were then left to oviposit for six days in three sets of ten small *Drosophila* vials with porridge food and live yeast flakes for two days each (Figure 1d). The vials were incubated and adult offspring were collected and separated by sex within 18 hours of eclosion to ensure virginity, and kept in groups of 10–15 flies. Offspring collection started between 18 and 20 days after the first day of oviposition and was spread over five to six days, depending on the development speed and the number of offspring eclosing. Collected offspring aged 2–8 days were used to create the next generation, leading to a generation time of 30–32 days. After the initial setup, populations were randomly allocated identifiers (1–12) that were changed every generation, such that all procedures were performed blind with respect to population composition (*low* vs *high* polyandry) and populations were not handled in a systematic order.

Every generation, we obtained an estimate of the frequency of polyandry for each of the twelve replicate populations. Female offspring from the experimental evolution populations were collected as virgins as described above. When they were 4.8 ± 1.3 days old we aspirated females into the vial of a tester male at the laboratory temperature of 19–22°C and allowed a minimum of 90min for mating, during which we observed pairs as continuously as possible. Tester males came from an outbred population whose

founders were caught in Chiricahua, Arizona (31°54'N, 109°16'W) in 2012. After mating, males were removed and females left to oviposit in the vial. Pairs that did not mate (N = 228, 8% of all pairs) were discarded. Four days after their initial mating, females were paired with a new tester male and allowed 90–180 minutes for mating (depending on their order in the assay, which we balanced across populations). We retained the vial in which the female had oviposited after her first mating to confirm functional matings and to be able to distinguish between pseudopolyandry and true polyandry (Fisher *et al.*, 2013). For the assays, we used three to nine day old (mean 5.3 days) virgin tester males that were separated into individual vials the day before the mating assay. Occasionally, we were unable to collect enough virgin males for second matings (about 10% across all assays), in which case we used males that mated in the first mating and ensured males were sexually rested and not paired with familiar females.

Assays after experimental evolution

To subject populations to one generation of common garden breeding after experimental evolution, we set up ten vials with five females and ten virgin males each per population, which gave females ample opportunity for mate choice. After 24 hours, before *D. pseudoobscura* females remate (Snook & So, 2000), males were removed and females were transferred to a new vial to oviposit. Female groups were then transferred to new food every 48h for a period of two weeks. We used the offspring of these flies for our final assays on polyandry (female remating latency), male remating inhibition, and fecundity.

Simulating initial polyandry in experimental populations

We simulated resampling of our setup of the 12 population replicates from the 16 ancestral isolines before experimental evolution, using *for* loops in R. Thus, for each of 1000 simulations, 12 experimental populations were virtually constructed by sampling females randomly from within isolines, with twelve females from each of the eight *M* isolines, and three females from each of the eight *P* isolines used to create virtual *low polyandry* replicate populations and vice versa for *high polyandry* populations. From these simulated populations, we used the remating latency of 26 random females from each population replicate (12 x 26 = 312), thus reflecting the sample size for females from our real, experimentally evolved populations (N = 316). From the *coxme* models run on all simulated datasets we obtained a distribution of the effect sizes for the difference between simulated populations with *low* versus *high* initial levels of polyandry, and compared it to the observed difference between real, evolved *low* and *high* polyandry populations. Similarly, we combined the simulated datasets with our real dataset to compare the simulated remating latencies in the population replicates before experimental evolution with the observed remating latencies of the real, experimentally evolved populations.

Supplementary results and discussion:

Remating inhibition by males

In one of the two experimental blocks, we counted offspring from vials in which tester females had oviposited for four days after the first mating. We did this to investigate whether variation in remating inhibition was associated with reproductive output (see e.g. Crudgington *et al.*, 2005; Price *et al.*, 2010b). Female remating tended to increase with reproductive output of fertile females between the first and second mating (0.016 [−0.002;0.033], GLMM, N = 348, $z = 1.82$, $p = 0.069$); Figure S3). However, although female reproductive output showed a signature of male genetic background (LRT: $\chi^2 = 17.7$, $df = 1$, N = 350, $p < 0.001$), there was no association with whether males came from *low/M* versus *high/P* polyandry or isoline versus evolved backgrounds (both $p > 0.5$; Figure S4).

Incompatibility between isolines and tester population

Our direct comparisons between isolines and evolved populations were limited by small sample sizes for matings between individuals derived from the Shaver Lake population isolines and tester individuals derived from the Chiricahua population. The consequences of this apparent incompatibility for our experimental evolution results are however rather minor. On the one hand, additional data indicate that both the incidence of fertile matings and the reproductive output from fertile matings are reduced in crosses between these Shaver Lake isolines and flies from the other three populations used in this study, which suggests partial reproductive isolation (Andreas Sutter, unpublished data). Overall, random genetic drift is only expected to have very minor effects on polyandry genotypes in our experimental populations of more than 200 individuals. But initiating the populations with 16 isolines meant that genotypes representing individual isolines were likely to be lost through drift. In combination with apparent reproductive incompatibility between Shaver Lake and other populations, loss of Shaver Lake genotypes from some of the experimental populations in early generations seems a likely scenario. On the other hand, all isolines were included in all experimental populations, such that loss of Shaver Lake alleles through reproductive incompatibility between Shaver Lake and other isolines should not have been biased towards one experimental evolution background or the other. Additionally, in the setup of our experimental populations we ensured a balanced representation of *P* and *M* isolines for all three populations of origin, including Shaver Lake, and the Shaver Lake isolines represented average polyandry genotypes within the *P* and *M* isoline groups (Figure 1b & Table S1).

References:

- Barron, A.B. 2000. Anaesthetising *Drosophila* for behavioural studies. *J. Insect Physiol.* **46**: 439–442.
- Crudgington, H.S., Beckerman, A.P., Brüstle, L., Green, K. & Snook, R.R. 2005. Experimental removal and elevation of sexual selection: does sexual selection generate manipulative males and resistant females? *Am. Nat.* **165**: S72–S87.
- David, J.R., Gibert, P., Legout, H., Pétavy, G., Capy, P. & Moreteau, B. 2005. Isofemale lines in

Drosophila: an empirical approach to quantitative trait analysis in natural populations. *Heredity*. **94**: 3–12.

Fisher, D.N., Doff, R.J. & Price, T.A. 2013. True polyandry and pseudopolyandry: why does a monandrous fly remate? *BMC Evol. Biol.* **13**: 157.

Price, T.A., Lewis, Z., Smith, D.T., Hurst, G.D.D. & Wedell, N. 2010. Sex ratio drive promotes sexual conflict and sexual coevolution in the fly *Drosophila pseudoobscura*. *Evolution* **64**: 1504–1509.

Snook, R.R. & So, Y.K. 2000. Associations between female remating behavior, oogenesis and oviposition in *Drosophila melanogaster* and *Drosophila pseudoobscura*. *J. Insect Physiol.* **46**: 1489–1496.

Supplementary tables:

Table S1: Summary results from assays on polyandry in isofemale isogenic lines sourced from three different populations. Sample sizes are summed across four experimental blocks for the preliminary assay and across two blocks for the repeat assay (see main text). The bottom 13 isolines were not selected for setting up the experimental evolution populations and were thus not included in the repeat assay.

Population	Isoline	Preliminary assay			Repeat assay		
		Tested	Remated	Polyandry	Tested	Remated	Polyandry
Lewistown, MT	<i>LEW3</i>	29	24	83%	24	20	83%
Show Low, AZ	<i>2SLOD6</i>	34	22	65%	31	11	35%
Lewistown, MT	<i>LEW64</i>	28	14	50%	18	9	50%
Shaver Lake, CA	<i>SHAB1</i>	23	9	39%	3	0	0%
Shaver Lake, CA	<i>SHAE4</i>	30	11	37%	5	2	40%
Show Low, AZ	<i>2SLOD29</i>	27	9	33%	32	6	19%
Show Low, AZ	<i>2SLOD33</i>	41	11	27%	20	7	35%
Show Low, AZ	<i>2SLOD15</i>	35	9	26%	23	4	17%
Lewistown, MT	<i>LEW23</i>	29	7	24%	27	7	26%
Lewistown, MT	<i>LEW17</i>	37	8	22%	31	2	6%
Show Low, AZ	<i>SLOC48</i>	26	5	19%	27	12	44%
Shaver Lake, CA	<i>SHAA6</i>	26	4	15%	3	1	33%
Show Low, AZ	<i>2SLOD26</i>	34	5	15%	25	7	28%
Show Low, AZ	<i>2SLOC4</i>	33	4	12%	32	7	22%
Shaver Lake, CA	<i>SHAB5</i>	21	2	10%	25	1	4%
Show Low, AZ	<i>SLOB3</i>	37	2	5%	29	3	10%
Total		490	146	30%	355	99	28%
Show Low, AZ	<i>2SLOC11</i>	39	17	44%	–	–	–
Shaver Lake, CA	<i>SHAB14</i>	27	12	44%	–	–	–
Show Low, AZ	<i>SLOC2</i>	12	5	42%	–	–	–
Lewistown, MT	<i>LEW13</i>	8	3	38%	–	–	–
Shaver Lake, CA	<i>SHAC1</i>	28	10	36%	–	–	–
Shaver Lake, CA	<i>SHAB8</i>	11	3	27%	–	–	–
Shaver Lake, CA	<i>SHAE7</i>	35	9	26%	–	–	–
Show Low, AZ	<i>SLOC9</i>	23	5	22%	–	–	–
Show Low, AZ	<i>SLOC12</i>	44	8	18%	–	–	–
Show Low, AZ	<i>SLOA13</i>	38	7	18%	–	–	–
Show Low, AZ	<i>SLOB7</i>	35	4	11%	–	–	–
Shaver Lake, CA	<i>SHAA10</i>	23	2	9%	–	–	–
Show Low, AZ	<i>2SLOD10</i>	8	0	0%	–	–	–

Table S2: Full LMM summary for fecundity of females from the 12 experimentally evolved populations. Fecundity was measured as the number of offspring eclosing from 6 days of oviposition after the second (denied or realised) mating opportunity. Covariates were centred as described in the main text. Except for significant variation between population replicates (included as a random effect), none of the covariates explained a significant amount of variation.

<i>Female fecundity</i>	Evolved females (N = 226)				
Fixed effects (LMM)	coef	se (coef)	ddf	t	p
Intercept (phenotype_forced monandry)	89.8	4.15	35.6	21.7	<0.001
female age (centred)	0.37	3.48	30.4	0.11	0.916
first male age (centred)	-1.44	3.25	89.5	-0.4	0.659
background (centred; <i>low v high</i>)	-12.7	8.33	35.8	-1.5	0.135
phenotype_elected monandry	4.80	4.21	209.9	1.14	0.255
phenotype_polyandry	-4.30	4.89	203.2	-0.9	0.381
background:elected monandry	11.7	8.41	209.8	1.39	0.166
background:polyandry	5.72	9.77	201.8	0.59	0.559
Random effects	Var	SD	df	χ^2	p
Female housing vial	<0.001	<0.001	1	0	1
Population replicate	66.6	8.16	1	5.1	0.024

Supplementary figures:

Figure S1: Cox proportional hazard model predictions for female latency to remate when given daily remating opportunities with tester males. Age and order covariates were mean-centred and scaled as described in the main text. The difference between *M/low* (orange) and *P/high* (blue) was significant both for isolines (solid lines) and for evolved populations (dashed lines). Note that higher polyandry results in a faster drop in these lines. See Figure 3a for an alternative presentation of these results.




Figure S2: No genetic correlation between female remating latency and male remating inhibition ($F_{1,14} = 2.17$, $p = 0.163$). Circles represent isolines and triangles represent experimentally evolved populations. Predominantly monandrous *M* isolines and *low* polyandry populations are shown in orange, while blue represents more polyandrous *P* isolines and *high* polyandry populations (see main text). Predictions for means and standard errors for remating latency and tester female remating (proxy for remating inhibition) for isolines and population replicates were obtained from univariate linear (latency) and generalised linear models (remating inhibition, backtransformed). The estimate for the correlation between males and females was based only on isolines to avoid pseudoreplication (see main text).




Figure S3: Tester female remating tended to increase with reproductive output from oviposition over the four days between virgin mating and remating opportunity (GLMM, $N = 348$ females, $p = 0.069$; see main text). Individual females are shown as red ticks, and a backtransformed prediction from a GLMM including male and female age as well as focal male isolate/replicate identity. Horizontal stripes illustrate bins of five-offspring-increments for which mean polyandry is shown as grey squares, with area proportional to sample size.




Figure S4: Variation in male-induced tester female fecundity over four days following their virgin mating. Circles and squares represent P/high and M/low polyandry backgrounds, respectively, and colours indicate population of origin (EE: experimental evolution; LEW: Lewistown, SHA: Shaver Lake; SLO: Show Low). Note the very small sample sizes for three of the four Shaver Lake isolines (see main text). Means and approximate 95% CI are shown in solid points and bars, and raw data are shown as semi-transparent points.




Figure S5: Variation in fecundity after experimental evolution, but no association with genetic polyandry. Raw data for female fecundity resulting from 6 days of oviposition across two consecutive vials after the second (denied or realised) mating opportunity, i.e. days 5-10 after the first mating. Offspring counts were pooled for the three treatments (forced monandry, chosen monandry and chosen polyandry), which showed no significant effect on fecundity (see Table S3). Colour denotes experimental evolution background, with *high* polyandry replicates shown in blue. Means and approximate 95% CI are shown in solid points and bars, and raw data are shown as semi-transparent points.

