

1 **Developing pathways to improve smallholder agricultural productivity through**
2 **ecological intensification technologies in semi-arid Limpopo, South Africa**

3

4 **¹Farirai Rusere, ¹Siyabusa Mkuhlani, ¹Olivier Crespo, ²Lynn V. Dicks**

5 **¹Climate System Analysis Group, Department of Environmental and Geographical Science,**
6 **University of Cape Town, South Africa**

7 **²School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich, NR4**
8 **7TJ, UK**

9 **Email: farirairusere@gmail.com**

10

11 **Abstract**

12 Agricultural productivity in many rural areas in Sub Saharan Africa is low. This affects food
13 security and rural livelihoods. Understanding farm diversity is essential to delineate
14 recommendation domains for new technologies. Farm typologies are a useful tool to assist in
15 unpacking and understanding the wide diversity among smallholder farms to improve targeting
16 of agricultural intensification strategies. We studied a community of smallholder farmers in Ha
17 Lambani, a village, Limpopo South Africa. In this study, agricultural experts identified farmer
18 groupings through based on the crops grown, farm size and major the source in which gross
19 maximum income was earned. A survey was then carried out to identify farming patterns,
20 constraints and we linked these constraints and solutions to specific ecosystem services that
21 appear to be currently important to the farming systems. This enabled us to explore the potential
22 to enhance productivity through ecological intensification, and provides important information
23 about which specific ecological intensification measures are likely to gain traction or appeal to
24 a particular group of farmers in this community. We conclude that although expert based
25 typologies enhance local relevance and reality, they need to be combined with statistical
26 approaches for effective selection of farms, innovation targeting and out-scaling of
27 technologies.

28

29 **Key Words: farm types; smallholder agriculture; ecosystem services, ecological intensification**

30

31 **Contents**

32	Abstract.....	1
33	1.0 Introduction.....	2
34	2.0 Materials and methods	5
35	2.1 Study area.....	5
36	2.2 Expert based typology construction.....	6
37	2.3 Identifying challenges, constraints and opportunities for ecological intensification	6
38	3.0 Results.....	7
39	3.1 Farm types and household characterisation	7
40	3.2 Farm types and farming system patterns.....	10

41	3.3 Farmers perceptions to their current challenges and constraints	11
42	3.4 Perceived solutions to their farming constraints and challenges	11
43	3.5 Identification of ecosystem services as a framework for ecological intensification targeting....	12
44	4.0 Discussion.....	14
45	4.1 The diversity of the farm types and farming system patterns	14
46	4.2 Perspectives on underperformance of farming systems.....	15
47	4.3 Opportunities for improved production through ecological intensification.....	17
48	4.4 Typology limitations.....	Error! Bookmark not defined.
49	5.0 Conclusion	19
50	Acknowledgements.....	21
51	References.....	21
52		

53 **1.0 Introduction**

54 In Southern Africa, smallholder farming is dominated by dryland crop production. The
55 regional average grain yields ranged from 0.3 to 2.2 tha⁻¹ during the period 2008–2012 (FAO,
56 2014). South Africa is generally considered as a food secure nation (De Cock et al., 2013), but
57 many households in rural areas are food insecure (Pereira et al., 2014). About 35.2 % of the
58 South African population live in rural areas and practice subsistence agriculture. They rely on
59 agricultural activities for their livelihoods, and are amongst the poorest and most vulnerable in
60 the country (Tibesigwa et al., 2014; Ncube et al., 2016). The rural farming households are
61 particularly vulnerable to climate and other disaster risks because they are mostly dependent
62 on rain fed traditional agriculture (Mwenge Kahinda & Taigbenu, 2011; Kong et al., 2014)
63 and have a low adaptive capacity due to technical, financial and infrastructural constraints
64 (Gbetibouo et al., 2010).

65

66 In South Africa and most surrounding countries in Southern Africa, agriculture and agricultural
67 related activities contribute to most of the employment in rural areas (Dercon & Gollin, 2014).
68 Smallholder agriculture has the potential to generate more employment, income and improve
69 livelihood opportunities in rural areas of South Africa (Shisanya & Hendriks, 2011; Mpandeli
70 & Maponya, 2014). Therefore, improving agriculture is considered as a viable and sustainable
71 alternative in reducing rural poverty in South Africa and other Sub Saharan African (SSA)
72 countries (Adekunle, 2014; Thamaga-Chitja & Morojele, 2014; Shisanya & Mafongoya, 2016).
73 With proactive technical and policy support, smallholder farmers can realise their potential to
74 become competitive in their agricultural production activities. Thus, improvement of

75 smallholder farming is a high priority in South Africa's improvement of rural communities
76 (Aliber & Hall, 2012; Kepe & Tessaro, 2014).

77

78 The manner in which agricultural technologies/innovations should be promoted in SSA to
79 improve agricultural production and sustainable livelihoods through smallholder farming is
80 largely debated (Wainaina, et al., 2016). Technologies on sustainable land use and improved
81 agricultural productivity have been developed, promoted and scaled out in the past 30 years in
82 SSA (Bidogeza, et al., 2009). However some of these technologies have only been partially
83 adopted (Giller, et al 2009, 2015), indeed most have not been fully adopted (Wainaina et al.,
84 2016). This is because most interventions are not reflective of smallholder farmer
85 circumstances and fail to acknowledge the environmental realities of smallholder farmers,
86 their social views, their perceptions of their own environmental realities and the strategies used
87 to meet their food security needs (Nhantumbo et al., 2016). For example, Giller et al. (2009)
88 argue that, despite CA being promoted heavily in SSA, they question its suitability and
89 effectiveness, especially in smallholder agriculture in SSA, highlighting a possible mismatch
90 between the conditions required for all CA principles to be adopted by farmers and the
91 circumstances that characterize and constrain smallholder African farming systems. This
92 disconnect undermines effective engagement between farmers, extension services and
93 researchers for effective improvement of technologies for adoption. Therefore new pathways
94 of fostering agricultural interventions in South Africa and SSA are needed before scaling up
95 such interventions (Whitbread et al., 2010; Sanyang et al., 2016). Agricultural
96 technologies/interventions aiming to enhance production, income and household livelihoods,
97 must capture the contrasting biophysical circumstances within and across the heterogeneous
98 agro ecologies in smallholder agriculture in SSA (Baudron, et al., 2015; Giller et al., 2015).
99 This must include the differing socio-economic circumstances within the sector.

100

101 Effectively identifying and integrating major issues that guide smallholder farmers' decision
102 making is therefore important to unlock current low adoption rates of practices such as
103 conservation agriculture, in situ rain water harvesting among others (Nhantumbo et al., 2016).
104 A practical way to understand smallholder farmers' decision making is to identify performance,
105 efficiency levels, challenges/constraints and opportunities. Understanding the vulnerability of
106 the farming systems to climate, social, economic and biophysical shocks and their impacts
107 could also help. Different modelling frameworks can be used to achieve the above. However,
108 a successful farming system analysis model requires the establishment of farm typologies.

109 Amongst farm households with similar production goals, biophysical and resource
110 endowments, farm typologies effectively classify the heterogeneity of farmers' motivations
111 and socio-economic circumstances related to their farming systems (Bidogeza et al., 2009;
112 Chikowo et al., 2014; Chenoune et al., 2016). Two approaches could be used to identify farms
113 heterogeneity in smallholder agriculture in SSA. The first is a bottom-up participatory approach
114 where every farmer is consulted and engaged through field visits, discussions and interviews.
115 The second is a top-down approach is where key informants are used to identify heterogeneity
116 and generate typologies as shown by (Tittonell et al., 2005; Zingore et al., 2007). Whilst the
117 first approach cannot be implemented on a large scale (lack of human and time resources), it
118 allows for better description of farmer's anticipation, capacity and willingness to adopt new
119 management strategies and agro-technologies. The second approach has the potential for large
120 scale implementation in cases where time and other resources are limiting. The classification
121 criteria depend on the goal of the typology and the kind of data available. Furthermore,
122 agricultural scientists are being encouraged to develop farm typologies to support a more
123 tailored approach to agricultural development and innovation (Kuivanen et al., 2016).

124

125 In SSA two models of fostering agricultural development and innovation to improve
126 smallholder agriculture have gained momentum, namely sustainable and ecological
127 intensification (Petersen & Snapp, 2015). The two are closely linked in terms of definitions,
128 principles and practices thus creating some confusion in their meaning, interpretation and
129 implications, although it is often argued that ecological intensification is more clearly defined,
130 with a better theoretical basis (Petersen & Snapp, 2015; Wezel et al., 2015). The major
131 difference between these two models is that for ecological intensification, agricultural systems
132 are designed to benefit from ecological processes and functions, including biological control
133 of biotic stressors and efficient use of available resources and ecological services (Bommarco
134 et al., 2013; Tittonell, 2014; Kovács-Hostyánszki et al., 2017). Sustainable intensification, on
135 the other hand, does not have a focus on ecological processes, although these can be
136 incorporated if they contribute to reduced inputs, increased outputs or enhanced efficiency. In
137 recent literature, sustainable intensification has tended to have more of a focus on technological
138 innovation and increased production without environmental impacts (Loos et al., 2014; Kuyper
139 & Struik, 2014; Godfray, 2015).

140

141 In this study, we focus on and explore ecological intensification, a means of increasing
142 agricultural production and environmental services while reducing the need for external inputs

143 and capitalising on ecological processes that support and regulate primary productivity in agro
144 ecosystems (Tittone, 2014). Ecological intensification seeks to ensure long term productivity
145 and sustainability through restoration of biodiversity and a full array of ecosystem functions
146 and services that support food production and human well-being aims to achieve a healthy
147 environment that provides multiple ecosystem services (*i.e.*, clean water, soil fertility, pest
148 suppression, nutrient cycling, and climate regulation) (Bommarco et al., 2013; Geertsema et
149 al., 2016). Ecosystem services and functions have particular relevance in Sub Saharan Africa
150 (SSA), where the majority of the population live in rural areas and rely on ecosystem services
151 and functions for their living through smallholder farming, pastoralism and fisheries (Egoh, et
152 al., 2012). Despite the potential of ecological intensification to improve food production
153 systems in smallholder agricultural systems in SSA (Rusere & Crespo, 2017), it has rarely been
154 seriously addressed in the context of smallholder farming systems of rural Africa (Tittone &
155 Giller, 2013) and its research remains limited in SSA (Struik, 2017). In this paper, the objective
156 was to explore the structure of the smallholder farming system, constraints, solutions and link
157 the constraints and solutions to specific ecosystem services that appear important in developing
158 pathways to ecological intensification in smallholder farms in rural South Africa.

159

160 **2.0 Materials and methods**

161 **2.1 Study area**

162 This study was conducted in Ha Lambani, a village in Vhembe District in Limpopo province
163 South Africa. Limpopo province is the fourth largest province in South Africa (SSA, 2015). It
164 has the highest population growth rate of 3.9 % per annum and 90 % of the population live in
165 rural areas (De Cock et al., 2013). According to Mpandeli & Maponya, (2014) the main
166 contributor to employment and livelihoods in the Vhembe District is agriculture. Smallholder
167 agriculture accounts for 70 % of the farming activities in the district whilst the other 30 % is
168 commercial agriculture. The district is situated in a semi-arid area and experiences water
169 shortages from May to August. Most commercial farmers depend on irrigation systems for
170 farming while the subsistence or smallholder farmers rely on seasonal rainfall which is
171 normally received from November to March. The district average annual rainfall is
172 approximately 820 mm. The smallholder farmers predominantly grow maize, legumes and
173 some vegetables for their own consumption, with any surplus sold or loaned to neighbours or
174 relatives. Rain fed crop yields are generally poor due to low and erratic rainfall.

175

176 **2.2 Identifying different smallholder farm types in Ha Lambani**

177 In the context of the project, to identify farm types, we used expert knowledge. An introductory
178 meeting was held with the senior agricultural extension workers to request cooperation from
179 the field based extension workers and to present the research objectives, which were (i) to
180 classify farms and farmers in the study area, (ii) unravel and assess farming system
181 performance and efficiency levels, (iii) identify challenges and constraints and (iv) identify
182 opportunities to drive farming systems towards more sustainable ones through ecological
183 intensification technologies in rural areas of South Africa. Five key informants, field based
184 agricultural extension workers based in the study area were identified by the senior agricultural
185 extension workers. Four of the agricultural extension workers specialised in crops and one
186 specialised in livestock. The agricultural extension workers were informed that the objective
187 was to classify smallholder farmers based on predominant socio-economic characteristics,
188 resource endowments and production objectives. Thereafter, local experts (agricultural
189 extension workers), based on their knowledge, in-depth experience and considering the
190 structure of the farming system and landscape and by identifying the most important sources
191 of variation among farms in the area, three farmer types were identified from the classification
192 variables listed in table 1 below. According to Kuivanen, et al., (2016) the expert based
193 approach of classifying farmers captures context specific aspects of farm complexity and has
194 potential to enhance local relevance and socio-cultural sensitivity aspects of interventions.
195 Nevertheless, the degree to which an expert based approach based on these variables can
196 predict actual behaviour in a context of rural development has not been proven. The main
197 limitation of the expert based approach in classifying farmers and farms is that the reliance on
198 local experts as sources of information is not enough for comprehensive understanding and
199 analysis of complex and diverse farming systems as it can be potentially misleading and biased.
200 Therefore, the expert based approaches need to be combined with participatory and statistical
201 approaches to retain objectivity and reproducibility Acknowledging this limitation, this work
202 is only able to complete the initial steps of establishing a baseline to guide in exploring potential
203 strategies to promote or foster ecological intensification.

204

205 **2.3 Identifying challenges, constraints and opportunities for ecological intensification**

206 We used a snowball sampling approach to identify farmers representing each of the three farm
207 types, to take part in the face to face interviews. Snowball sampling is an approach for locating
208 information-rich key informants to participate in the study (Duan & Hoagwood, 2015). Using
209 this approach, agricultural extension officers identified potential farmer respondents to

210 represent the three farm types. Face to face questionnaire based structured interviews were
211 conducted with the help of the agricultural extension workers who assisted in the translation of
212 the questions and farmer responses. The interviews sought information on the estimates of farm
213 size, cropped area, types of crops grown, estimates of yields obtained, crop preferences and
214 production objectives. Farmers were asked to identify major constraints and challenges to their
215 current crop and livestock farming practices. Farmers' perceptions on their potential solutions
216 to their production constraints and objectives were sought by means of open ended questions.
217 Furthermore, through discussions with farmers, we could identify key ecosystem services
218 important to different farm types in the study area. Owing to the small sample size of farmers
219 interviewed in each class, descriptive statistical analysis was carried out on the survey data.
220 The results of the survey are summarized in table 2

221 **3.0 Results**

222 **3.1 Farm types and household characterisation**

223 Field based agricultural extension workers identified three types of farms in Ha lambani. The
224 farms were overlapping in many characteristics but differed in their main source of income.
225 The farm types identified were namely (1) the cereal and livestock based, (2) the horticultural
226 based and (3) the off-farm income dependent farms. The table1 shows the variables used to
227 build the typology.

228

229 **Type 1: Cereal and livestock based farms**

230 The cereal and livestock based farms were large farms (averaging more than 2 ha), with elderly
231 household heads (60 years old and more). Maize is the most cultivated crop whereas legumes
232 and vegetables are minor crops in this category. Livestock is a determinant factor, with farms
233 rearing mainly cattle and goats (10-15 cows and 5 goats on average). Cereal and livestock
234 activities contribute most to the household income (75%), while social grants and remittances
235 come as a complement (25%).

236

237 **Type 2: Horticultural based farms**

238 The horticulture based farms are small, often less than 1.5 ha. They comprise mainly young
239 household heads ranging from 18 to 35 years of age. Vegetables are mostly grown and maize
240 (green mealies) is cultivated as a minor crop. Most of the farmers in this category do not own
241 livestock. Income from horticultural activities is the major source of household income.

242

243 **Type 3: The off-farm income dependent farms**

244 The off-farm income farms are average size farms often between 1.5-2 ha. The household heads
245 are mainly farmers aged between 36 to 60 years. They mostly grow maize, vegetables and
246 legume as minor crops. They own a small herd of livestock biased towards ruminants (5 cows
247 and 5 goats own in average). The largest household income comes from salaries and part time
248 jobs they engage into in their local communities, complemented in small portion by agricultural
249 activities.

250

251

252

253

254 **Table 1: Variables used to construct the farm types**

255	Variable	(1) Cereal and livestock based	(2) Horticultural based	(3) Off farm income
256	dependent			
257	Household size	5	3	5
258	Average Number of Children	3	1	3
259	Age of household head	> 60	18-35	35-60
260	Level of education of household head	no education	matric grade 12	matric grade 12
261	Major source of income	farming	farming	salaries/ par time jobs
262	Other sources of income	grants/remittances	grants	farming
263	Average farm size	>2ha	<1.5ha	1.5-2ha
264	Average size of land cultivated	1.5 ha	1 ha	1ha
265	Major crops grown	maize	vegetables	maize
266	Minor crops grown	legumes, vegetables	green mealies	vegetables, legumes
267	Average maize yields	1 tonne/ha	0.25-0.5 tonnes/ha	> 0.5 tonnes/ha
268	Number of cattle	15	0	5
269	Number of goats	5	0	5
270	Use of chemical inputs	Low	Moderate	Low

271

272

273 **3.2 Farm types and farming system patterns**

274 We interviewed 40 farmers of which 16 were cereal and livestock based farmers, 7 horticultural
275 based farmers and 17 off farm income dependent farmers. The interview results revealed an
276 estimated average farm size of more than 2 ha for Type 1 and less than 2 ha but more 1.5 ha
277 for Types 3, with Type 1 farms exhibiting the largest average cropped area of 1.5 ha. Type 2
278 had the smallest average farm size of less than 1.5 ha which corresponded to the smallest
279 average cropped area of less than 1 ha. Maize being the major crop grown by Type 1 and 3
280 farmers, although no crop yields records were available in almost all the households
281 interviewed. The farmers estimated that the yields obtained were very poor averaging 1 t/ha
282 and just above 0.5 t/ha for Type 1 and 3 respectively. All farm types were involved in vegetable
283 production with only type 2 farms growing vegetables as their major crops and primarily as a
284 cash crop and a major source of income. The results indicated that Type 2 farmers preferred to
285 grow high value horticultural crops grown on a small area throughout the year. Type 1 and 3
286 farmers are involved in the production of legumes (mainly cowpea and groundnuts) and
287 vegetables on a small scale mainly for household consumption and rarely as cash crops.
288 Furthermore, results from the interviews further affirmed that Type 1 and 3 farms are involved
289 in livestock production with Type1 farms possessing the most animals and the largest cattle
290 herds. Type 2 farmers did not possess any cattle or small ruminants citing lack of capital to
291 purchase as well as lack of resources and labour to rear the animals. Type 2 farmers generally
292 lack access to animal traction, resulting in reduced crop area.

293

294 Chemical input use in all farm types was generally low with Type I and 3 farmers applying
295 between 1-3 50kg bags of inorganic fertiliser per hectare. Type 2 (horticulture based) farms
296 where chemical input use was moderate applied between 4 to 8 bags per hectare. This was
297 because horticultural crops are input demanding and their lack of livestock meant lack of
298 organic fertilisers such as manure as soil fertility amendments. Type 2 farmers highlighted that
299 they relied more on suboptimal application of inorganic fertilisers for soil fertility improvement
300 and sub optimal application agrochemicals for crop protection against pest and diseases.
301 Although income from social grants helped Type 1 (cereal and livestock based) farmers to
302 acquire some farming inputs, Type 1 and 3 low use is mostly due to fertilizer and herbicides
303 cost and access. Hence Type 1 and 3 farms relied on traditionally low resource input methods
304 of agriculture.

305

306 Type 1 farmers rely mostly on farming (sale of agricultural produce and livestock) for income
307 although they are recipients of government social grants of the elderly and remittances from
308 their children located in urban areas. Type 2 farmers rely on producing and selling high value
309 horticultural products although most are also recipients of child support grants. Type 2 farmers
310 also highlighted that financial returns from crops like maize, cowpea and groundnuts were
311 often not worth the effort when set against the risks of producing those crops under rain fed
312 conditions. Type 3 farms often engage in non-farm based strategies such as craft making, bead
313 work, carpentry, brick moulding, traditional beer selling and seasonal work as hired labour for
314 household income. They engage in agricultural activities to supplement household income.
315

316 **3.3 Farmers perceptions to their current challenges and constraints**

317 The interview results revealed that all farm types faced varying challenges and constraints in
318 their agricultural activities, although poor seasonal rainfall distribution, low precipitation
319 amounts and lack of and or poor irrigation infrastructure which was dilapidated were common
320 constraints among all the farm types. A significant proportion of Type 1 farmers also cited poor
321 access to inputs as well as high costs of input especially fertilizer as a major constraint.
322 Furthermore, they pointed out shortage of livestock feed, especially during the dry season and
323 drought years, leading to loss of livestock or crop damage by livestock during the dry season.
324 Type 2 farmers cited high incidences of pest and diseases in their fields as a dominant constraint
325 in their cropping fields. In addition, Type 2 farmers pointed out post-harvest losses and poor
326 access to markets as major constraints. Furthermore, they highlighted poor access to pesticides
327 despite having limited financial resources. Mechanization and draught power were their major
328 challenges to increase area under crops. Type 3 farmers considered lack of access to inputs,
329 lack of livestock feed during the dry season and drought years as well as damage of crops by
330 livestock during the dry season as significant constraints
331

332 **3.4 Perceived solutions to their farming constraints and challenges**

333 All farmers in all the farm types proposed government subsidies on agricultural inputs as well
334 as improved irrigation infrastructure as potential sustainable solution to their challenges and
335 constraints. Type 1 farmers proposed access to drought tolerant varieties of their cereal crops
336 to help achieve higher yields. Establishment of paddocks was cited to allow their livestock for
337 their livestock to graze. Lastly, they highlighted the need to access to financial institutions for
338 loans or grant as it would facilitate the acquisition of much needed irrigation systems

339 machinery and or inputs for improved crop production. Type 2 farmers see quick access to
340 markets and proper post-harvest handling facilities as direct improvements to cater for their
341 perishable horticultural products. Furthermore, they pointed out the need for training in local
342 horticultural crop production skills. Type 3 farmers, for whom farming is supplementary to off
343 farm income, consider fencing of fields and establishment of paddocks for their livestock to
344 graze as would greatly improve their agricultural activities.

345

346 **3.5 Identification of ecosystem services as a framework for ecological intensification** 347 **targeting**

348 Using the interview results, three key ecosystem services needed for each farm type to improve
349 agricultural productivity were identified (Table 2). All farm types, identified soil and water
350 conservation as a key ecosystem service they would benefit from to increase agricultural
351 production. Type 2 farmers (horticultural based) further emphasised the need for improved
352 water quality for improved horticultural production. A significant proportion of Type 1 (cereal
353 and livestock based) and Type 3 (off farm income dependent) farmers identified, nutrient
354 recycling as a key ecosystem service for improving agricultural production in their farming
355 landscapes. Type 1 farmers further emphasised the need for ecosystem services that improve
356 availability of forage and fodder for improved livestock production. Type 2 and Type 3
357 identified pest and disease suppression as key ecosystem service needed for improved
358 agricultural productivity.

359 **Table 2: Showing the challenges and constraints, solutions and the key ecosystem services needed to implement ecological intensification**
 360 **in the three farmer types in Ha Lambani, Vhembe District South Africa**

361 Farm Type	Cereal and livestock based	Horticultural based	Off farm income based
362 Initial problems	Poor rainfall	Poor rainfall	Poor rainfall
363	Access to inputs	High incidences of diseases	Poor irrigation infrastructure
364	Shortage of livestock feed	Poor mechanization	Shortage of livestock feed
365	High input costs	Inputs not easily accessible	Damage of crops by livestock
366	Poor irrigation infrastructure	Limited irrigation infrastructure	Access to inputs
367	Damage of crops by livestock	Poor access to markets	
368			
369 Proposed solutions	Government subsidies	Government subsidies	Government subsidies
370	Irrigation infrastructure	Irrigation infrastructure	Irrigation infrastructure
371	Tolerant varieties	Access to markets	Fencing of fields
372	Access to finance	Knowledge and skills	Paddocking of livestock
373	Paddocking of livestock	Proper post-harvest handling	
374			
375 Ecosystems services	Soil and water conservation	Soil and water conservation	Soil water conservation
376 related issues	Nutrient recycling	Pest and disease suppression	Nutrient recycling
377	Forage and fodder	water quality	Pest and disease suppression
378			

379 **4.0 Discussion**

380 **4.1 The diversity of the farm types and farming system patterns**

381 The typology developed in this study combined both expert knowledge and participatory
382 approaches to unravel the complexity and diversity in heterogeneous smallholder farming
383 systems. The clear differentiating factors identified among farm types were farm size, the farm
384 objective and the major contributor to household income, which resulted in three farm types
385 (Table 1). Results have shown that farming systems are driven by different farming objective
386 that in turn are shaped by various factors. These different objectives influence the different
387 farming system patterns exhibited in different smallholder farm types. Of these farm types, we
388 found that Type 2 (horticultural based) farms was well distinguished from Type 1 (Cereal and
389 livestock based) and Type 3 (averaged sized farms with off farm income dependent farms).
390 These two types showed intermediate properties, hence less distinctiveness.

391
392 Cereal and livestock based farm types have capacity to grow cereal and leguminous crops, use
393 best agronomic practices, including early planting, weeding and application of organic manures
394 fertilizers, and this enhances the yield difference when compared with off farm income and
395 horticultural based farm types who have limited land and labour. Furthermore, the rearing of
396 livestock is very important for satisfying food security in South Africa. Livestock represent the
397 most important store of value for farmers and the wealth of a household can be measured by
398 the number and type of animals owned (Chaminuka, Udo, Eilers, & Zijpp, 2014). Livestock
399 herds owned by Type 1 (cereal and livestock based) farmers and Type 3 (off farm income
400 dependent) farmers provide animal traction and manure, thus putting these farmers at an
401 advantage in terms of agronomic performance, improved soil fertility and planting large area
402 when compared to Type 2 (horticultural based) farmers.

403
404 Farmer income affects most decisions, including those regarding adoption of farming practices
405 which can require financial investment and can reduce short term profitability. The extension
406 workers we consulted in Ha Lambani segregated on farm and off-farm income because the
407 source of income influences its connection to farm business investment decisions. Type 1 farms
408 rely mostly on farming (sale of produce and livestock) for income although they are recipients
409 of government social grants of the elderly and remittances from their children located in urban
410 areas. Financial and resource limitation in Type 3 farms may often induce a shift in livelihood
411 strategies towards a higher dependence on off-farm income. This influences decision-making,
412 cropping patterns and farming practices. Engagement in non-farm activities limits the amount

413 of time Type 3 farmers engage in cropping activities. Delays in farming operations are common
414 in Type 3 farms resulting in poor yields. In semi-arid environments, where there is only a
415 narrow window for getting the right balance of agronomic practices that facilitate high yields.
416 To improve livelihoods and household income Type 3 farms often engage in non-farm based
417 strategies such as craft making, bead work, carpentry, brick moulding, traditional beer selling
418 and seasonal work as hired labour to supplement household income. These findings suggest
419 income from farming, off farm income generating activities and social grants play an important
420 role in the livelihoods of people in the study area. This is because the income from these
421 activities determine the livelihood strategies to be adopted by the households.

422

423 A very small proportion of the rural population in Ha Lambani make a significant income from
424 growing crops like maize, cowpea and groundnuts. This has led the few young people involved
425 in farming to specialise in horticultural crops which are of high value with high returns for
426 income in addition to the child support grants they receive. Hence type 2 farms are
427 horticulturally based and derive most of their income from sale of horticultural produce. An
428 important finding of this study that agrees with other studies is that very few young people
429 want to engage in cereal and legume crop production in rural areas. This is because agriculture
430 is often perceived as an occupation of the poor, hence young people have little desire to be
431 involved in it (Leavy & Hossain, 2014). Furthermore, these findings highlight the importance
432 of taking a comprehensive survey of the production envelope, rather than focusing only on
433 blanket recommendations when targeting and tailoring agricultural interventions to local
434 contexts. Technological interventions, development strategies and policies to address the
435 problem of poor productivity and reduce poverty in smallholder agricultural systems must be
436 designed to target socially diverse and spatially heterogeneous farms and farming systems in
437 rural South Africa.

438

439 **4.2 Perspectives on underperformance of farming systems**

440 As shown by the results, the different farm types tend to experience the same major constraints
441 in general. Poor seasonal rainfall distribution and amount, and poor or lack of irrigation
442 infrastructure were common constraints among all farm types. This is because most
443 smallholders, if not all farmers in Ha Lambani depend on rainfall for their agricultural
444 activities. The unreliable and limited availability of water and infrastructure for irrigation,
445 increases unpredictability thus affecting farmers' ability to plan what, when and where to plant
446 their crops and other farm related decisions. The low mean annual rainfall of 500-800 mm,

447 high annual evaporation of 2000-2500 mm in Ha Lambani (Botha et al., 2014) and recurring
448 droughts indicate severe crop water stress during most seasons. Limited irrigation
449 infrastructure that is dilapidated and malfunctional further exacerbates the problem.

450

451 Furthermore, the limited access to seed, farming equipment, fertilizers and agrochemicals by
452 poorer households translate into a limited capacity to diversify their livelihood strategies by
453 growing more demanding crops. In many aspects of smallholder production in Ha Lambani,
454 declining soil fertility is a major constraint. Although Type 1 and Type 3 farms relied on animal
455 manure for soil fertility improvement, the low nutrient content of manure tend to mean that
456 very large quantities of manure are needed. The average quantity of manure applied to crops
457 was insufficient to achieve good yields. Furthermore, manure alone may be an unsatisfactory
458 source of nutrients, especially for nitrogen and phosphorus which are required by plants in
459 large quantities, and therefore rarely provide the productivity needed for high yielding cereal
460 crops. It has therefore been suggested that to sustain high crop yields, manure may need to be
461 combined with nitrogen fixing legumes in resource constrained low input farming systems.

462

463 The fact that weed, insect pest and disease problems are amongst the major constraints being
464 experienced by Type 2 farmers, lead us to suspect that they have an impact on their livelihoods
465 due to the susceptibility of horticultural crops to these biotic stressors. Smallholder farmers in
466 Ha Lambani operate in a resource constrained environment in terms of access to inputs such as
467 pesticides and fertilisers. Furthermore, the demand for constant labour, herbicides, pesticides
468 and the lack of a strong technical resource base for crop protection available further exacerbates
469 the problem. Technical agronomic and horticultural information relating to cultivar and seed
470 choice, soil fertility, water management and pest management using cultural, biological and
471 chemical methods is also still lacking. The smallholder horticultural sector therefore requires
472 support in the form of improved access to technical pest management information (in an appropriate
473 form). Furthermore, research targeting knowledge gaps through in which ecological intensification
474 can help manage pest and diseases via biological control methods such as, use of natural enemies,
475 plant extracts and other sustainable integrated pest management (IPM) methods is needed.

476

477 Despite numerous efforts to promote production of high value cash crops in smallholder
478 agriculture as a crucial step in solving food security problems in Africa, most famers including
479 Type 2 farmers cannot easily access profitable cash crop markets for their high value
480 horticultural produce. Their burden is further made worse due to lack of proper or poor storage

481 facilities resulting in severe post-harvest losses. This indicates that most smallholder farmers
482 are still excluded and marginalized with regards to markets access and market information.
483 Moreover, farmers who can produce surpluses remain trapped in the poverty cycle and more
484 often these farmers are forced to sell their produce at low prices to unscrupulous buyers who
485 dictate market prices.

486

487 However, the low quality and quantity of available forages during the dry season is a major
488 constraint for improved livestock production in Ha Lambani. Like in many rural areas of South
489 Africa, the available grazing is not generally sufficient to meet the maintenance requirements
490 of grazing animals (Matlebyane et al 2010) during dry periods. Although Type 1 (cereal and
491 livestock based) and Type 3 (off farm income dependent) farmers use different types of feed
492 to supplement for their livestock during the dry season and drought years, issues of availability,
493 quantity and quality of feed resources tends of affect them. Feed problems are mainly attributed
494 to land shortage, lack of improved forage technologies and awareness problem. Introduction
495 of improved forage technologies that can fit into the existing land use system coupled with
496 improved feeding systems would be necessary to resolve the feed related problems. At the same
497 time, other problems affecting livestock production in the area should be addressed
498 simultaneously to realize the potential benefits to be accrued from livestock.

499

500 Among the solutions mentioned by all the farm types, to the above-mentioned constraints and
501 challenges were increased government subsidies for agricultural inputs and rehabilitation or
502 improvement of existing irrigation infrastructure. This indicates that most of the agricultural
503 activities are currently low input systems relying on supporting and regulating ecological
504 process. Therefore, improving these farming systems in smallholder agriculture through
505 improved ecologically based management strategies might represent a viable and sustainable
506 pathway to increase productivity and resilience of smallholder agricultural systems given the
507 limited financial support of smallholder farmers from government.

508

509 **4.3 Opportunities for improved production through ecological intensification**

510 Enhanced ecosystem service provision is therefore critical for building resilience and
511 improving food and nutrition security for smallholder farmers in SSA. The farmer interviews
512 identified four key ecosystem services needed to improve agricultural productivity in Ha
513 Lambani. All farm types, identified soil and water conservation as a key ecosystem service
514 they would benefit from to increase agricultural production. Insufficient rainfall over the years

515 has resulted in severe water shortages for both domestic and agricultural purposes. Thus,
516 managing and harnessing ecosystem services linked to soil and water conservation offer
517 potential to increase agricultural production. This presents tremendous opportunities for
518 ecological intensification practices and interventions like minimum tillage, mulching, water
519 harvesting among others (Kassam et al., 2014) which make use of natural capital within the
520 soil to promote soil and water conservation in agricultural landscapes. For instance,
521 (Thierfelder et al., 2015) collated and summarised evidence on effects of minimum tillage, and
522 various soil amendments on soil water storage in smallholder agriculture in southern Africa.
523 Type 2 farmers mostly grow their horticultural crops under some form of irrigation.
524 Horticultural crops are highly dependent on water quality therefore a clean and constant water
525 supply is very important and would highly benefit Type 2 farmers.

526

527 A significant proportion of the Type 1 (cereal and livestock based) and Type 3 (off farm income
528 improved agricultural production. This emerges from high nutrient demanding main cereal
529 crops and would benefit from nutrient recycling ecosystem services to improve soil fertility.
530 Furthermore, depletion of soil fertility because of low fertilizer use and high rates of nutrient
531 mining are common challenges among smallholder farmers in South Africa and the region
532 beyond. (Shamie Zingore, 2016). Ecosystem services and processes that increase soil fertility
533 in their fields are therefore critical. This presents an opportunity for ecological intensification
534 through practices and interventions that promote ecological processes and biological diversity
535 in farming systems. The supporting and regulating ecosystem services and processes can be
536 incorporated into cropping systems, such that production is improved, nutrient flow and soil
537 fertility is enhanced and at the same time reducing the need for external inputs such as fertiliser.
538 This further presents an opportunity for ecological intensification practices and interventions
539 like intercropping, crop rotations to maximize production, nutrient flow and improve soil
540 fertility in resource constrained farms in Ha Lambani.

541

542 Furthermore, provision of forage and fodder was identified as key ecosystem services they
543 would benefit from to improve livestock productivity. Low quality and quantity of feeds are a
544 major constraint limiting livestock productivity among smallholder farmers. Ecosystem
545 services and processes that provide forage and fodder are important and could benefit Type 1
546 (cereal and livestock based) and Type 3 (off farm income dependent) farms. Although
547 ecological intensification is widely documented in field crops (Gomes, et al., 2014), it is less
548 well documented in animal production. However, it presents an opportunity for the

549 development and operationalisation of ecological processes and services in resource
550 constrained smallholder livestock systems. To foster such a development and
551 operationalisation, we propose the introduction of improved forage technologies such as forage
552 legumes and crop residues that can fit into the existing land use system coupled with improved
553 feeding systems would be necessary to resolve the feed related problems.

554

555 Lastly Type 2 (horticultural based) farmers identified proposed pest and disease suppression as
556 key ecosystem services needed for them to improve productivity. Weeds, insects and pathogens
557 infestation a major challenge to their horticultural farming activities, demand constant labour
558 and pesticides to treat them. In Ha Lambani where farmers access and ability to purchase
559 chemical pesticides is limited, ecosystem services that enhance natural pest control are very
560 critical. This presents an opportunity for ecological intensification to enhance crop protection
561 in resource constrained farmers. Dicks et al., (2016) summarised evidence that identified
562 practices that enhanced natural pest control in agriculture. In this regard, ecological
563 intensification approaches that make use of biological processes (such as use of natural
564 enemies, push-pull systems, crop rotations among others) to regulate pest population may
565 enhance pest suppression and regulating ecosystem services thus contributing to crop
566 protection. There is quite clear evidence that some of these interventions work, especially the
567 push-pull systems (Khan et al., 2008; Midega et al., 2014). This could be a beneficial low costs
568 and environmentally friendly crop protection strategy in resource constrained farms.

569

570 **5.0 Conclusion**

571 This study was in response to the need to identify the heterogenous farming system patterns
572 and diversity in smallholder farmers in South Africa to target ecological intensification in the
573 design and implementation of agricultural development interventions and technologies. The
574 farmer classification is the first step to identify diversity of the 3 farm types in Ha Lambani, a
575 village in Vhembe district, Limpopo, South Africa. Farmers can be distinguished based on their
576 sources of income, household involvement in both on and off farm activities and the diversity
577 of the farmers' agricultural land use. The farmer classification offered a more contextualized
578 representation of farming system heterogeneity in terms of challenges, constraints and
579 opportunities faced by farmers of the 3 identified farm types. Different types of farmers are
580 expected to pursue different trajectories in farm system design for targeting ecological
581 intensification to harness ecosystem services that flow from the agroecosystems under study.

582

583 **List of Abbreviations**

584 ACCESS: Alliance for Collaboration for Climate and Earth System Sciences

585 ADCI: African Climate Development Initiative

586 NRF: National Research Foundation

587 SSA: Sub Saharan Africa

588 WRC: Water Research Commission

589 **Declarations**

590 **Ethics approval and consent to participate**

591 Not applicable

592 **Consent for publication**

593 Not applicable

594 **Availability of Data and materials**

595 Not applicable

596 **Competing interests**

597 The authors declare that they have no financial or personal relationships which may have
598 inappropriately influenced them in writing this article.

599 **Funding**

600 This research was supported with funding from National Research Foundation (NRF), Water
601 Research Commission (WRC) and African Climate Development Initiative (ACDI) and
602 Alliance for Collaboration for Climate and Earth System Sciences (ACCESS). Lynn Dicks was
603 funded by the Natural Environment Research Council (NERC) (grant code: NE/N014472/1).
604 Interpretation of the findings and conclusion drawn from the study were the responsibilities of
605 the authors and not on any part of National Research Foundation (NRF), Alliance for
606 Collaboration for Climate and Earth System Sciences (ACCESS), Water Research
607 Commission (WRC), African Climate Development Initiative (ACDI) and Natural
608 Environment Research Council (NERC).

609 **Authors contributions**

610 Farirai Rusere and Siyabusa Mkuhlani was responsible for developing the initial content of the
611 manuscript, including literature search, data collection and analysis. Olivier Crespo and Lynn
612 Dicks were the supervisors for the research and provided guidance in terms of the article
613 structure and directed the retrieval of relevant literature and finalization of the manuscript
614

615 **Acknowledgements**

616 The authors are very grateful to Alliance for Collaboration for Climate and Earth System
617 Sciences (ACCESS), South African Water Research Commission (WRC), African Climate
618 Development Initiative (ACDI) and Natural Environment Research Council (NERC) (grant
619 code: NE/N014472/1) for funding the work, the Limpopo Department of Agriculture and the
620 Institute of Rural Development, University of Venda for their technical support.

621 **References**

- 622 Adekunle, A. K. (2014). Exploring the Roles of Agricultural Extension in Promoting, Food
623 Security in Kwazulu Natal Province, South Africa. *International Journal of Agricultural*
624 *Science, Research and Technology in Extension and Education System* 3(3), 147-158.
- 625 Aliber, M., & Hall, R. (2012). Support for smallholder farmers in South Africa: Challenges of
626 scale and strategy. *Development Southern Africa*, 29(4), 548–562.
627 <https://doi.org/10.1080/0376835X.2012.715441>
- 628 Baudron, F., Thierfelder, C., Nyagumbo, I., & Gérard, B. (2015). Where to Target
629 Conservation Agriculture for African Smallholders? How to Overcome Challenges
630 Associated with its Implementation? Experience from Eastern and Southern Africa.
631 *Environments*, 2(3), 338–357. <https://doi.org/10.3390/environments2030338>
- 632 Benis Egoh, Evangelia G. Drakou, Martha B. Dunbar, Joachim Maes, L. W., Egoh, B.,
633 Drakou, E. G., Dunbar, M. B., & Maes, J. (2012). *Indicators for mapping ecosystem*
634 *services: a review*. <https://doi.org/10.2788/41823>
- 635 Bidogeza, J. C., Berentsen, P. B. M., Graaff, J., & Oude Lansink, a. G. J. M. (2009). A
636 typology of farm households for the Umutara Province in Rwanda. *Food Security*, 1(3),
637 321–335. <https://doi.org/10.1007/s12571-009-0029-8>
- 638 Bommarco, R., Kleijn, D., & Potts, S. G. (2013). Ecological intensification: Harnessing
639 ecosystem services for food security. *Trends in Ecology and Evolution*, 28(4), 230–238.

- 640 <https://doi.org/10.1016/j.tree.2012.10.012>
- 641 Botha, J. J., Anderson, J. J., Theron, J. F., Taljaard, D. J., Venter, I. S., & Koatla, T. A. B.
642 (n.d.). *Guidelines on Best Management Practices for Rainwater Harvesting and*
643 *Conservation (RWH&C) for Cropland and Rangeland Productivity in Communal Semi-*
644 *Arid Areas of South Africa.*
- 645 Chaminuka, P., Udo, H. M. J., Eilers, K. C. H. A. M., & Zijpp, A. Van Der. (2014). Land Use
646 Policy Livelihood roles of cattle and prospects for alternative land uses at the wildlife /
647 livestock interface in South Africa. *Land Use Policy*, 38, 80–90.
648 <https://doi.org/10.1016/j.landusepol.2013.10.007>
- 649 Chenoune, R., Belhouchette, H., Paloma, S. G. y, & Capillon, A. (2016). Assessing the
650 diversity of smallholder rice farms production strategies in Sierra Leone. *NJAS -*
651 *Wageningen Journal of Life Sciences*, 76, 7–19.
652 <https://doi.org/10.1016/j.njas.2015.10.001>
- 653 Chikowo, R., & Zingore, S. (2014). Farm typologies , soil fertility variability and nutrient
654 management in smallholder farming in Sub-Saharan Africa, 1–18.
655 <https://doi.org/10.1007/s10705-014-9632-y>
- 656 De Cock, N., D’Haese, M., Vink, N., van Rooyen, C. J., Staelens, L., Sch??nfeldt, H. C., &
657 D’Haese, L. (2013). Food security in rural areas of Limpopo province, South Africa.
658 *Food Security*, 5(2), 269–282. <https://doi.org/10.1007/s12571-013-0247-y>
- 659 Dercon, S., & Gollin, D. (2014). Agriculture in African Development: Theories and
660 Strategies. *Annual Review of Resource Economics*, 6(1), 471–492.
661 <https://doi.org/10.1146/annurev-resource-100913-012706>
- 662 Dicks, L. V, Wright, H. L., Ashpole, J. E., Hutchison, J., McCormack, C. G., Livoreil, B., ...
663 Sutherland, W. J. (2016). What works in conservation ? Using expert assessment of
664 summarised evidence to identify practices that enhance natural pest control in
665 agriculture. *Biodiversity and Conservation*, 25(7), 1383–1399.
666 <https://doi.org/10.1007/s10531-016-1133-7>
- 667 Duan, N., & Hoagwood, K. (2015). Purposeful Sampling for Qualitative Data Collection and
668 Analysis in Mixed Method Implementation Research. *Administration and Policy in*
669 *Mental Health and Mental Health Services Research*, 533–544.

670 <https://doi.org/10.1007/s10488-013-0528-y>

671 Food and Agriculture Organization of the United Nations (FAO). (2014). *FAO Statistical*
672 *Yearbook 2014- Asia and the Pacific Food and Agriculture*.
673 <https://doi.org/http://www.fao.org/docrep/018/i3107e/i3107e.PDF>

674 Gbetibouo, G. A., Ringler, C., & Hassan, R. (2010). Vulnerability of the South African
675 farming sector to climate change and variability : An indicator approach Vulnerability of
676 the South African farming sector to climate change and variability : An indicator
677 approach, *34*(September 2015), 175–187. [https://doi.org/10.1111/j.1477-](https://doi.org/10.1111/j.1477-8947.2010.01302.x)
678 [8947.2010.01302.x](https://doi.org/10.1111/j.1477-8947.2010.01302.x)

679 Geertsema, W., Rossing, W. A., Landis, D. A., Bianchi, F. J., Van Rijn, P. C., Schaminée, J.
680 H., ... Van Der Werf, W. (2016). Actionable knowledge for ecological intensification of
681 agriculture. *Frontiers in Ecology and the Environment*, *14*(4), 209–216.
682 <https://doi.org/10.1002/fee.1258>

683 Giller, K. E., Andersson, J. A., Corbeels, M., Kirkegaard, J., Mortensen, D., Erenstein, O., &
684 Vanlauwe, B. (2015). Beyond conservation agriculture. *Frontiers in Plant Science*,
685 *6*(October), 1–14. <https://doi.org/10.3389/fpls.2015.00870>

686 Giller, K. E., Witter, E., Corbeels, M., & Tittonell, P. (2009). Conservation agriculture and
687 smallholder farming in Africa: The heretics' view. *Field Crops Research*, *114*(1), 23–
688 34. <https://doi.org/10.1016/j.fcr.2009.06.017>

689 Godfray, H. C. J. (2015). The debate over sustainable intensification, 199–208.
690 <https://doi.org/10.1007/s12571-015-0424-2>

691 Gomes, L. C., Pailleux, J., Dedieu, B., Regina, C., & Cournut, S. (n.d.). systems, (v), 1352–
692 1363.

693 Kassam, A., Derpsch, R., & Friedrich, T. (2014). Global achievements in soil and water
694 conservation : The case of Conservation Agriculture 1 Introduction. *International Soil*
695 *and Water Conservation Research*, *2*(1), 5–13. [https://doi.org/10.1016/S2095-](https://doi.org/10.1016/S2095-6339(15)30009-5)
696 [6339\(15\)30009-5](https://doi.org/10.1016/S2095-6339(15)30009-5)

697 Kepe, T., & Tessaro, D. (2014). Trading-off: Rural food security and land rights in South
698 Africa. *Land Use Policy*, *36*, 267–274. <https://doi.org/10.1016/j.landusepol.2013.08.013>

699 Khan, Z. R., Midega, C. A. O., Amudavi, D. M., Hassanali, A., & Pickett, J. A. (2008). On-

- 700 farm evaluation of the “ push – pull ” technology for the control of stemborers and striga
701 weed on maize in western Kenya, *106*, 224–233.
702 <https://doi.org/10.1016/j.fcr.2007.12.002>
- 703 Kong, T. M., Austin, D. E., Kellner, K., & Orr, B. J. (2014). The interplay of knowledge,
704 attitude and practice of livestock farmers’ land management against desertification in the
705 South African Kalahari. *Journal of Arid Environments*, *105*, 12–21.
706 <https://doi.org/10.1016/j.jaridenv.2014.02.002>
- 707 Kovács-Hostyánszki, A., Espíndola, A., Vanbergen, A. J., Settele, J., Kremen, C., & Dicks,
708 L. V. (2017). REVIEW AND Ecological intensification to mitigate impacts of
709 conventional intensive land use on pollinators and pollination, 673–689.
710 <https://doi.org/10.1111/ele.12762>
- 711 Kuivanen, K. S., Alvarez, S., Michalscheck, M., Adjei-nsiah, S., & Descheemaeker, K.
712 (2016). NJAS - Wageningen Journal of Life Sciences Characterising the diversity of
713 smallholder farming systems and their constraints and opportunities for innovation : A
714 case study from the Northern Region , Ghana. *NJAS - Wageningen Journal of Life*
715 *Sciences*, *78*, 153–166. <https://doi.org/10.1016/j.njas.2016.04.003>
- 716 Kuyper, T. W., & Struik, P. C. (2014). Epilogue : global food security , rhetoric , and the
717 sustainable intensification debate. *Current Opinion in Environmental Sustainability*, *8*,
718 71–79. <https://doi.org/10.1016/j.cosust.2014.09.004>
- 719 Leavy, J., & Hossain, N. (2014). *IDS WORKING PAPER Volume 2014 No 439 Who Wants to*
720 *Farm ? Youth Aspirations , Opportunities and Rising Food Prices* (Vol. 2014).
- 721 Loos, J., Abson, D. J., Chappell, M. J., Hanspach, J., Mikulcak, F., Tichit, M., & Fischer, J.
722 (2014). Putting meaning back into “ sustainable intensification ” In a nutshell :, 356–
723 361. <https://doi.org/10.1890/130157>
- 724 Midega, C. A. O., Jonsson, M., Khan, Z. R., & Ekbom, B. (2014). Effects of landscape
725 complexity and habitat management on stemborer colonization, parasitism and damage
726 to maize. *Agriculture, Ecosystems and Environment*, *188*, 289–293.
727 <https://doi.org/10.1016/j.agee.2014.02.028>
- 728 Mpandeli, S., & Maponya, P. (2014). Constraints and Challenges Facing the Small Scale
729 Farmers in Limpopo Province, South Africa. *Journal of Agricultural Science*, *6(4)*, 135–

- 730 143. <https://doi.org/10.5539/jas.v6n4p135>
- 731 Mwenge Kahinda, J., & Taigbenu, A. E. (2011). Rainwater harvesting in South Africa:
732 Challenges and opportunities. *Physics and Chemistry of the Earth*, 36(14–15), 968–976.
733 <https://doi.org/10.1016/j.pce.2011.08.011>
- 734 Ncube, M., Madubula, N., Ngwenya, H., Zinyengere, N., Zhou, L., Francis, J., ...
735 Madzivhandila, T. (2016). Climate change, household vulnerability and smart
736 agriculture: The case of two South African provinces. *Jamba: Journal of Disaster Risk
737 Studies*, 8(2), 6–8. <https://doi.org/10.4102/jamba.v8i2.182>
- 738 Nhantumbo, N. S., Zivale, C. O., Nhantumbo, I. S., & Gomes, A. M. (2016). Making
739 agricultural intervention attractive to farmers in Africa through inclusive innovation
740 systems. *World Development Perspectives*, 4, 19–23.
741 <https://doi.org/10.1016/j.wdp.2016.12.003>
- 742 Pereira, L. M., Cuneo, C. N., & Twine, W. C. (2014). Food and cash: Understanding the role
743 of the retail sector in rural food security in South Africa. *Food Security*, 6(3), 339–357.
744 <https://doi.org/10.1007/s12571-014-0349-1>
- 745 Petersen, B., & Snapp, S. (2015). What is sustainable intensification? Views from experts.
746 *Land Use Policy*, 46, 1–10. <https://doi.org/10.1016/j.landusepol.2015.02.002>
- 747 Rusere, F., & Crespo, O. (2018). A review on the potential of ecological intensification to
748 improve food production systems in smallholder agriculture in Sub Saharan Africa. In
749 press submitted in *Agriculture & Food Security Journal*
- 750 Sanyang, S., Taonda, S. J., Kuisu, J., & Coulibaly, N. T. (2016). A paradigm shift in African
751 agricultural research for development : the role of innovation platforms, 5903(February
752 2017). <https://doi.org/10.1080/14735903.2015.1070065>
- 753 Security, F., Province, K., Africa, S., Adegunle, A. K., & West, N. (2014). Exploring the
754 Roles of Agricultural Extension in Promoting, 3(3).
- 755 Shisanya, S., & Mafongoya, P. (2016). Adaptation to climate change and the impacts on
756 household food security among rural farmers in uMzinyathi District of Kwazulu-Natal,
757 South Africa. *Food Security*, 8(3), 597–608. <https://doi.org/10.1007/s12571-016-0569-7>
- 758 Shisanya, S. O., & Hendriks, S. L. (2011). The contribution of community gardens to food
759 security in the Maphephetheni uplands. *Development Southern Africa*, 28(4), 509–526.

- 760 <https://doi.org/10.1080/0376835X.2011.605568>
- 761 Struik, P. C. (2017). Sustainable intensification in agriculture : the richer shade of green . A
762 review. <https://doi.org/10.1007/s13593-017-0445-7>
- 763 Thamaga-Chitja, J. M., & Morojele, P. (2014). The Context of Smallholder Farming in South
764 Africa: Towards a Livelihood Asset Building Framework. *Journal of Human Ecology*,
765 45(2), 147–155.
- 766 Thierfelder, C., Rusinamhodzi, L., Ngwira, A. R., Mupangwa, W., Nyagumbo, I., Kassie, G.
767 T., & Cairns, J. E. (2015). Conservation agriculture in Southern Africa: Advances in
768 knowledge. *Renewable Agriculture & Food Systems*, 30(4), 328–348.
769 <https://doi.org/10.1017/S1742170513000550>
- 770 Tibesigwa, B., Visser, M., & Turpie, J. (2014). The impact of climate change on net revenue
771 and food adequacy of subsistence farming households in South Africa. *Environment and*
772 *Development Economics*, (May), 327–353.
773 <https://doi.org/10.1017/S1355770X14000540>
- 774 Tittonell, P. (2014). Ecological intensification of agriculture-sustainable by nature. *Current*
775 *Opinion in Environmental Sustainability*, 8, 53–61.
776 <https://doi.org/10.1016/j.cosust.2014.08.006>
- 777 Tittonell, P., & Giller, K. E. (2013). When yield gaps are poverty traps: The paradigm of
778 ecological intensification in African smallholder agriculture. *Field Crops Research*, 143,
779 76–90. <https://doi.org/10.1016/j.fcr.2012.10.007>
- 780 Tittonell, P., Vanlauwe, B., Leffelaar, P. A., Shepherd, K. D., & Giller, K. E. (2005).
781 Exploring diversity in soil fertility management of smallholder farms in western Kenya
782 II . Within-farm variability in resource allocation , nutrient flows and soil fertility status,
783 110, 166–184. <https://doi.org/10.1016/j.agee.2005.04.003>
- 784 Wainaina, P., Tongruksawattana, S., & Qaim, M. (2016). Tradeoffs and complementarities in
785 the adoption of improved seeds, fertilizer, and natural resource management
786 technologies in Kenya. *Agricultural Economics (United Kingdom)*, 47(3), 351–362.
787 <https://doi.org/10.1111/agec.12235>
- 788 Wezel, A., Soboksa, G., McClelland, S., Delespesse, F., & Boissau, A. (2015). The blurred
789 boundaries of ecological , sustainable , and agroecological intensification : a review,

790 1283–1295. <https://doi.org/10.1007/s13593-015-0333-y>

791 Whitbread, A. M., Robertson, M. J., Carberry, P. S., & Dimes, J. P. (2010). How farming
792 systems simulation can aid the development of more sustainable smallholder farming
793 systems in southern Africa. *European Journal of Agronomy*, 32(1), 51–58.
794 <https://doi.org/10.1016/j.eja.2009.05.004>

795 Zingore, S. (2016). Influence of soil fertility variability and nutrient source on maize
796 productivity and nitrogen use efficiency on smallholder farms in Zimbabwe,
797 (December), 4–8.

798 Zingore, S., Murwira, H. K., Delve, R. J., & Giller, K. E. (2007). Influence of nutrient
799 management strategies on variability of soil fertility, crop yields and nutrient balances
800 on smallholder farms in Zimbabwe. *Agriculture, Ecosystems and Environment*, 119(1–
801 2), 112–126. <https://doi.org/10.1016/j.agee.2006.06.019>

802

803

804