

# The human imperative of stabilizing global climate change at 1.5°C.

## Authors:

O. Hoegh-Guldberg<sup>1,2</sup>, D. Jacob<sup>3</sup>, M. Taylor<sup>4</sup>, T. Guillén Bolaños<sup>3</sup>, M. Bindi<sup>5</sup>, S. Brown<sup>6,7</sup>, I. A. Camilloni<sup>8</sup>, A. Diedhiou<sup>9</sup>, R. Djalante<sup>10,11</sup>, K. Ebi<sup>12</sup>, F. Engelbrecht<sup>13</sup>, J. Guiot<sup>14</sup>, Y. Hijioka<sup>15</sup>, S. Mehrotra<sup>16</sup>, C. W. Hope<sup>17</sup>, A. J. Payne<sup>18</sup>, H-O Pörtner<sup>19</sup>, S. I. Seneviratne<sup>20</sup>, A. Thomas<sup>21,22</sup>, R. Warren<sup>23</sup>, G. Zhou<sup>24</sup>.

## Affiliations:

1. Global Change Institute, University of Queensland, St Lucia 4072 QLD Australia
2. School of Biological Sciences, University of Queensland, St Lucia 4072 QLD, Australia
3. Climate Service Center Germany (GERICS), Helmholtz-Zentrum Geesthacht, Hamburg, Germany
4. Department of Physics, University of the West Indies, Kingston, Jamaica
5. Department of Agriculture, Food, Environment and Forestry (DAGRI) -University of Florence, Piazzale delle Cascine 18, 50144 Italy
6. Faculty of Engineering and Physical Sciences, University of Southampton, Boldrewood Innovation Campus, Burgess Road, Southampton. SO16 7QF, UK
7. Department of Life and Environmental Sciences, Faculty of Science and Technology, Bournemouth University, Fern Barrow, Poole, Dorset, BH12 5BB, UK
8. Centro de Investigaciones del Mar y la Atmósfera (UBA-CONICET), UMI-IFAECI/CNRS, and Departamento de Ciencias de la Atmósfera y los Océanos (FCEN, UBA), Buenos Aires, Argentina
9. Université Grenoble Alpes, French National Research Institute for Sustainable Development (IRD), CNRS, Grenoble INP, IGE, F-38000 Grenoble, France United Nations University
10. United Nations University - Institute for the Advanced Study of Sustainability (UNU-IAS), Tokyo, Japan
11. Halu Oleo University, Kendari, South East Sulawesi, Indonesia
12. Center for Health and the Global Environment, University of Washington, Seattle WA, USA
13. Global Change Institute, University of the Witwatersrand, Johannesburg 2193, South Africa

14. Aix Marseille University, CNRS, IRD, INRA, College de France, CEREGE, Aix-en-Provence, France
15. Center for Climate Change Adaptation, National Institute for Environmental Studies, 16-2, Onogawa, Tsukuba, Ibaraki, 305–8506, Japan
16. The World Bank, Washington DC, United States.
17. Cambridge Judge Business School, University of Cambridge, Cambridge, UK
18. University of Bristol, Bristol, UK
19. Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven, Germany
20. Institute for Atmospheric and Climate Science, ETH Zurich, Zurich, Switzerland
21. Climate Analytics, 10961 Berlin, Germany
22. Environmental and Life Sciences, University of The Bahamas, Nassau 76905, The Bahamas
23. Tyndall Centre for Climate Change Research and School of Environmental Sciences, University of East Anglia, Norwich NR4 7TJ, UK
24. State Key Laboratory of Severe Weather, Chinese Academy of Meteorological Sciences, Beijing 100081, China

# **Abstract:**

Global mean surface temperature is now 1.0°C higher than the pre-industrial period due to increasing atmospheric greenhouse gases. Here, we expand on the recent IPCC Special Report on global warming of 1.5°C and review the additional risks associated with higher levels of warming, each having major implications for multiple geographies, climates and ecosystems. Limiting warming to 1.5°C rather than 2.0°C is very beneficial, maintaining significant proportions of ecosystems as well as having clear benefits for human health and economies. These conclusions are relevant for people everywhere, particularly in low- and middle-income countries, where climate related risks are escalating with major implications for the achievement of the United Nations Sustainable Development Goals.

**One Sentence Summary:** Climate change is already driving dangerous impacts that will be progressively less manageable at 1.5°C of global warming or higher.

**Main text:**

Climate change is one of the greatest challenges for humanity. Global mean surface temperature (GMST) is increasing at the rate of  $0.2^{\circ}\text{C} \pm 0.1^{\circ}\text{C}$  per decade, reaching  $1.0^{\circ}\text{C}$  above the pre-industrial period (reference period 1850–1900) in 2017 (1). GMST is projected to reach  $1.5^{\circ}\text{C}$  above the pre-industrial period between 2030 and 2052, depending on the model and assumptions regarding projected changes to atmospheric greenhouse gas (GHG) levels and climate sensitivity (1). At the same time, growing awareness of impacts beyond  $1.5^{\circ}\text{C}$  has focused international attention on the feasibility and implications of stabilizing temperatures at this level (2).

In broad terms, limiting warming to  $1.5^{\circ}\text{C}$  will require a total investment in the energy sector of 1.46–3.51 trillion (US\$2010) in energy supply and 0.64–0.91 trillion (US\$2010) in energy demand measures in order to reach net zero GHG emissions by 2050 (3)(p154). On the other hand, the mean net present value (in 2008) of the avoided damages resulting from this action is estimated as totalling \$496 trillion (US\$2010) by the year 2200 (3–5). This, together with other damages that are difficult to fully cost and include (e.g. disruption and migration of human communities; reductions in ecosystem services associated with biodiversity loss), suggests that potential economic benefits arising from limiting warming to  $1.5^{\circ}\text{C}$  may be at least four or five times larger than the investments needed to stabilize GMST to  $1.5^{\circ}\text{C}$  (SM1)(3).

Here, we explore the near-term mostly unmonetized impacts projected for  $1.5^{\circ}\text{C}$  of global warming, and the associated risks and adaptation options for natural and human (managed) systems. In order to understand the implications of reaching  $1.5^{\circ}\text{C}$ , we compare it to recent conditions (i.e.  $1.0^{\circ}\text{C}$  warming above the pre-industrial period, Fig 1), and to those that are projected to emerge as we approach  $2.0^{\circ}\text{C}$  of warming. This comparison helps understand the benefits or not of stabilizing GMST at  $1.5^{\circ}\text{C}$  as compared to  $2.0^{\circ}\text{C}$  or higher, as well as providing a framework for societal responses and consequences.

94 [Insert Figure 1 here]

95 **Crossing the 1.0°C threshold has already severely impacted natural and human systems**

96  
97 The incidence of extremes has increased sharply as GMST has warmed from 0.5°C to 1.0°C  
98 (~1980 – 2018) relative to the Pre-industrial period, with the intensity and/or frequency of  
99 extremes projected to change further with another 0.5°C of warming (5). As GMST has  
100 increased, for example, the average temperature of cold days and nights (i.e. the coldest 10%)  
101 has also increased overall, as has the average temperature of warm days and nights (i.e. the  
102 warmest 10%) globally (5). These changes have also been accompanied by increases in the  
103 frequency and/or duration of heatwaves for large parts of Europe, North America and Australia.  
104 Increases in GMST have been accompanied by increases in the frequency, intensity and/or  
105 amount of heavy precipitation in more regions than those with decreases, especially in North-  
106 Hemisphere mid-latitude and high-latitude areas (5, 6). There is also evidence of increasing  
107 rainfall associated with recent tropical cyclones (6, 7) and increasingly heavy precipitation  
108 during storms in the Central Sahel (8, 9). The total annual number of tropical cyclones has  
109 decreased, while the number of very intense cyclones has increased, for many areas (5). There is  
110 less confidence regarding trends in the length of drought, although a significant increasing trend  
111 has been detected in the Mediterranean region (particularly Southern Europe, North Africa and  
112 the near-East) (10–12).

113  
114 As on land, coastal and marine habitats have also experienced an increased frequency, intensity  
115 and duration of underwater heatwaves, with a threefold increase in the number of marine  
116 heatwave days globally since 1980 (13). The differential heating of the water column has also led  
117 to increased thermal stratification in some coastal and oceanic regions which decreases ocean-  
118 atmosphere gas exchange as well the turnover of nutrients between the photic layer and deeper  
119 layers of the ocean. The annual mean Arctic sea ice extent decreased by 3.5 - 4.1% per annum  
120 from 1979 to 2012 (6). The melting of land-based ice includes potentially unstable regions such  
121 as the Western Antarctic Ice Sheet (WAIS, Fig 1B), which contributed  $6.9 \pm 0.6$  mm over 1979-  
122 2017 to global mean sea level (GMSL). Together with glacial melt water, thermal expansion of  
123 the ocean has accelerated the rate of GMSL increase by up to 0.013 [0.007-0.019] mm yr<sup>-2</sup> since

the early 20th Century (14). Changes in ocean temperature have also decreased the oxygen concentration of the bulk ocean, interacting with coastal pollution to increase the number and extent of low oxygen dead zones in many deep-water coastal habitats (15). In addition to increasing GMST, anthropogenic CO<sub>2</sub> also enters the ocean causing a reduction in pH (ocean acidification) which negatively impacts processes such as early development, calcification, photosynthesis, respiration, sensory systems, and gas exchange in organisms from algae to fish (5).

Changing weather patterns (e.g. temperature, rainfall, dryness, storms) have increased negative impacts on natural and managed systems (Fig 1A-D). Changes to coral reefs (5), forests (e.g. changing drought/fire regimes) (16, 17), low-lying islands and coasts (5), and impacts on agriculture production and yield (18, 19) are threatening resources for dependent human communities. There are also many gradual changes that have occurred as GMST has increased, with many being no less important than the more abrupt changes. Land-based biomes (i.e. major natural and agricultural ecosystem types) have also shifted to higher latitudes and elevation in boreal, temperate and tropical regions (5, 15), with similar shifts reported for marine and freshwater organisms. Marine organisms and some ecosystems have also shifted their biogeographical ranges to higher latitudes at rates up to 40 km yr<sup>-1</sup>. Rates are highest for pelagic organisms and ecosystems such as plankton, and are lowest for more sedentary benthic organisms and ecosystems such as seaweeds and kelp forests (5, 15). These types of changes (e.g. temperature, storms, circulation) have also affected the structure and function of ocean ecosystems with respect to its biodiversity, food-webs, incidence disease and invasive species (5).

Other changes to biological systems include changes to the phenology of marine, freshwater and terrestrial organisms (e.g. timing of key events such as reproduction and migration) (5, 15). The phenology of plants and animals in the Northern-Hemisphere, for example, has advanced by 2.8 ± 0.35 days per decade due to climate change, with similar changes in the flowering and pollination of plants and crops, and the egg-laying and migration times of birds (5, 20). There are indications that climate change has already contributed to observed declines in insects and arthropods in some regions (21, 22). Variations in these types of changes have also been

observed in the phenology of tropical forests, which have been more responsive to changes in moisture stress rather than to the direct changes in temperature (5). While the intention here is not to catalogue all of the changes that have occurring in natural systems, it is important to acknowledge that deep and fundamental changes are underway in biological systems with just 1°C of global warming so far (5).

Changes in GMST of 1.0°C have also directly and indirectly affected human communities, many of which depend on natural and managed systems for food, clean water, coastal defence, safe places to live, and livelihoods among many other ecosystem goods and services (5). Coral reefs clearly illustrate the linkage between climate change, ecosystem services and human well-being. At 1.0°C, large-scale mortality events driven by lengthening marine heatwaves have already reduced coral populations in many places (5), with prominent coral reef ecosystems such as the Great Barrier Reef in Australia losing as much as 50% of their shallow water corals in the last four years alone (5, 23, 24). These changes have potential implications for millions of people given their dependency on coral reefs for food, livelihoods and well-being (5).

## **Understanding climate change over the next few decades: methods and assumptions**

There is a range of strategies for quantifying risks for natural and human systems at 1.5°C and 2.0°C above the pre-industrial period. This requires calculating the future exposure of systems to changes in climatic hazards. Some methods rely on the fact that an equivalent amount of warming (e.g. 0.5°C) occurred in the recent past (e.g. ca. 1950 to 2000, or ca. 1980 to 2018, Fig 2A; (3)) potentially providing insights into how risks might change in the near future. In this case, the associated risks of the next 0.5°C of global warming (Fig 2A) are linearly extrapolated from the impacts associated with the previous 0.5°C increase (ca. 1980-2018). This method of projecting future risk is likely to be conservative given (i) the pace of climate change is increasing (25) and (ii) the impacts per unit of temperature are likely to increase as conditions are pushed increasingly beyond the optimal conditions for a particular organism or physiological process (Fig 2B)(26). Responses by natural and human systems are likely to also differ if temperature pathways involve a gradual increase to 1.5°C above the pre-industrial period (no ‘overshoot’) as opposed to pathways that first exceed 1.5°C before later declining to 1.5°C,

which is referred to as an ‘overshoot’ (5) (Fig 2A). High levels of overshoot involve exceeding 1.5°C by 0.1°C (Figure 2A) (3).

[Insert Figure 2 here]

Other approaches for understanding how the world may change at 1.5°C and 2.0°C of global warming draw on laboratory, mesocosm, and field experiments. These approaches simulate projected conditions for different levels of warming and, in the case of marine systems, levels of acidification (e.g. changes in pH, carbonate, pollution levels (5, 26, 27). These experimental approaches also provide calibration as well as insight into future conditions and responses (i.e. 1.5°C versus 2.0°C). Some caution is also required given that global increases of 1.5°C or 2.0°C may involve a broad range of regional responses. This arises due to uncertainties in (for example) the likelihood of overshoot, land-atmosphere interactions, biophysical effects of land use changes, and interannual climate variability (28). Several lines of evidence for understanding these complex problems include the analysis of the frequency and intensity of extremes as well as projections based on existing climate simulations and empirical scaling relationships for 1.5°C and 2.0°C of global warming (5). Lines of investigation may also include dedicated experiments prescribing sea surface conditions consistent with these levels of warming, as done in the HAPPI (Half a degree Additional warming, Prognosis and Projected Impacts) project (5). Furthermore, fully-coupled climate model experiments can be achieved using GHG forcing consistent with 1.5°C or 2.0°C scenarios (5). These multiple yet different lines of evidence (above) underpin the development of qualitatively consistent results regarding how temperature means and extremes could change at 1.5°C as compared to 2.0°C of global warming.

## **Projected changes in climate at 1.5°C versus 2.0°C of global warming**

Understanding the potential advantages of restraining global warming to 1.5°C requires an understanding of the risks associated with the exposure of natural and human systems to climatic hazards, and how they change at 1.5°C relative to 2.0°C (Fig 3)(29). Increases of GMST to 1.5°C will further increase the intensity and frequency of hot days and nights, and decrease the



intensity and frequency of cold days and nights (Fig 3 C.D.E). Warming trends are projected to be highest over land, in particular for temperature extremes, with increases of up to 3°C in the mid-latitude warm season and up to 4.5°C in cold seasons at high latitudes. These increases are projected to be greater at 2.0°C of global warming, with increases of up to 4°C in the mid-latitude warm season and up to 6°C in the high-latitude cold season (e.g. Fig 3 A.C.D.E.) (29). Heatwaves on land, which are already increasing pressure on health and agricultural systems, are projected to become more frequent and longer (Fig 3 C.D.).

There is considerable evidence that dryness will increase in some regions, especially the Mediterranean as well as southern Africa (5, 30–32). Risks of drought, dryness and precipitation deficits are projected to increase at 1.5°C and even further at 2.0°C for some regions relative to the pre-industrial period (Fig 3B,F)(5, 33). Recent studies also suggest similar projections for the western Sahel and southern Africa, as well as the Amazon, north-eastern Brazil, and Central Europe (5, 34). Projected trends in dryness are uncertain in several regions, however, and some regions are projected to become wetter(Fig 3 B,F) (5). Reaching GMST of 1.5°C and 2.0°C, for example, would lead to a successive increase in the frequency, intensity and/or amount of heavy rainfall when averaged over global land area (Fig 3 B,F). Global warming of 2.0°C versus 1.5°C increases exposure to fluvial flood risk particularly at higher latitudes and in mountainous regions, as well as in East Asia, China (35) and eastern North America overall (5). The prevalence of subsequent intense wet and dry spells, in which a prolonged drought is immediately followed by heavy precipitation at the same location (potentially leading to flooding) or vice versa, is projected to be greater at 2.0°C global warming versus 1.5°C (36). These large changes between coupled wet and dry conditions represent a major challenge for adaptation as they will affect water quality and availability as well as increased soil erosion along many coastal areas. Sea level rise can also amplify problems through damage to coastal infrastructure and the salinization of water supplies for drinking and agriculture (5).

Relatively few studies have directly explored the effect of 1.5°C versus 2.0°C of global warming on tropical cyclones (5). These studies consistently reveal a decrease in the global number of tropical cyclones at 1.5°C vs 1.0°C of global warming, with further decreases under 2.0°C vs 1.5°C of global warming. Simultaneously, very intense cyclones are likely to occur more



frequently at 2.0°C vs 1.5°C of global warming, with associated increases in heavy rainfall and damage, further emphasizing the advantages of not exceeding 1.5°C (5).

[Insert Figure 3 here]

Coastal and oceanic regions are also projected to increase in temperature as GMST increases to 1.5°C, and further to 2.0°C, above the pre-industrial period. Absolute rates of warming are only slightly lower in the ocean than on land although the shallower spatial gradient of ocean temperature will mean that the velocity of climate change may be higher in many regions of the ocean (5, 37). Increases in ocean temperature associated with 1.5°C and 2.0°C of global warming will increase the frequency and duration of marine heatwaves, as well as reducing the extent of ocean mixing due to the greater thermal stratification of the water column (13, 15). Sea ice is projected to continue to decrease in the Arctic, although restraining warming to 1.5°C will mean an ice free Arctic summer will only occur every 100 years, while warming to 2.0°C above the pre-industrial period will mean an ice free Arctic summer is likely to occur every 10 years by 2100 (5, 38). These and other models indicate that there will be no long-term consequences for sea ice coverage in the Arctic (i.e. no hysteresis) if GMST is stabilised at or below 1.5°C (3).

### **Impacts on ecosystems at 1.5°C versus 2.0°C of global warming**

Multiple lines of evidence (5) indicate that reaching and exceeding 1.5°C will further transform both natural and human systems, leading to reduced ecosystem goods and services for humanity. Importantly, risks for terrestrial and wetland ecosystems such as increasing coastal inundation, fire intensity and frequency, extreme weather events, and the spread of invasive species and diseases are lower at 1.5°C as compared to 2.0°C of global warming (5). In this regard, the global terrestrial land area that is predicted to be affected by ecosystem transformations at 2.0°C (13%, interquartile range 8-20%) is approximately halved at 1.5°C (4%, interquartile range 2-7%). Risks for natural and managed ecosystems are higher on drylands as compared to humid lands (5). The number of species that are projected to lose at least half of their climatically determined geographic range at 2.0°C of global warming (18% of insects, 16% of plants, 8% of

vertebrates) would be significantly reduced at global warming of 1.5°C (i.e. to 6% of insects, 8% of plants, and 4% of vertebrates)(5). In this regard, species loss and associated risks of extinction are much lower at 1.5°C than 2°C. Tundra and boreal forests at high latitudes are particularly at risk, with woody shrubs having already encroached on tundra, which will increase with further warming (5). Constraining global warming to 1.5°C would reduce risks associated with the thawing of an estimated 1.5-2.5 million km<sup>2</sup> of permafrost (over centuries) compared to the extent of thawing expected at 2.0°C (5).

Ecosystems in the ocean are also experiencing large-scale changes, with critical thresholds projected to be increasingly exceeded at 1.5°C and higher global warming. Increasing water temperatures are driving the relocation of many species (e.g. fish, plankton) while sedentary organisms, such as kelp and corals, are relatively less able to move. In these cases, there are multiple lines of evidence that indicate that 70-90% of warm water tropical corals present today are at risk of being eliminated even if warming is restrained to 1.5°C. Exceeding 2.0°C of global warming will drive the loss of 99% of reef-building corals (5). These non-linear changes in survivorship are a consequence of the increasing impact of changes as they move away from optimal conditions (Fig 2B) (26). Impacts on oceanic ecosystems are expected to increase at global warming of 1.5°C relative to today, with losses being far greater at 2.0°C of global warming. Significant compound or secondary risks exist with respect to declining ocean productivity, loss of coastal protection, damage to ecosystems, shifts of species to higher latitudes, and the loss of fisheries productivity (particularly at low latitudes)(15). There is substantial evidence that these changes to coastal risks will increasingly threaten the lives and livelihoods of millions of people throughout the world (5).

### **Increasing risks for human (managed) systems at 1.5°C and 2.0°C of global warming**

Many risks for society will increase as environmental conditions change. Water, for example, is often central to the success or failure of human communities. The projected frequency and scale of floods and droughts in some regions will be smaller under 1.5°C global warming as opposed to 2°C, with risks to water scarcity being greater at 2.0°C than at 1.5°C of global warming for many regions (5). Salinization of freshwater resources on small islands and along low-lying coastlines is a major risk that will become successively more important as sea levels rise,

particularly as they will continue to increase even if temperatures stabilise. (5). Depending on future socio-economic conditions, limiting warming to 1.5°C is projected to reduce the proportion of the world's population exposed to climate induced water stress by up to 50% as compared at 2°C (5), although there is considerable variability among regions as already discussed. Most regions, including the Mediterranean and Caribbean regions, are projected to experience significant benefits from restraining global warming to 1.5°C (39), although socio-economic drivers are expected to play a dominant role relative to climate change for these communities over the next 30-40 years.

Limiting global warming to 1.5°C is projected to result in smaller reductions in the yield of maize, rice, wheat and potentially other cereal crops than at 2.0°C, particularly in sub-Saharan Africa, Southeast Asia, and Central and South America (40–42). A loss of 7-10% of rangeland stock globally is also projected to occur at an increase of 2.0°C above the pre-industrial period, which will have considerable economic consequences for many communities and regions. Reduced food availability at 2.0°C as compared to 1.5°C of global warming is projected for many regions including the Sahel, Southern Africa, the Mediterranean, Central Europe and the Amazon. Few examples exist where crop yields are increasing and hence food security is at increasing risk in many regions (41). Although food systems in future economic and trade environments may provide important options for mitigating hunger risk and disadvantage (43, 44)(5), assuming that solutions are found to the decline in the nutritional quality of major cereal crops from higher CO<sub>2</sub> concentrations (5).

Food production from marine fisheries and aquaculture is of growing importance to global food security but is facing increasing risks from ocean warming and ocean acidification (5). These risks increase at 1.5°C of global warming and ocean acidification, and are projected to impact key organisms such as finfish, corals, crustaceans and bivalves (e.g. oysters) especially at low latitudes (5). Small-scale fisheries that depend on coastal ecosystems such as coral reefs, seagrass, kelp forests and mangroves, are expected to face growing risks at 1.5°C of warming as a result of the loss of habitat (5). Risks of impacts, and subsequent risks to food security, are projected to become greater as global warming reaches 1.5°C (5, 43, 44) Tropical cyclones have major impacts on natural and human systems, and are projected to increase in intensity in many

regions, with the damage exacerbated by rapid sea level rise (14, 45). The tropical cyclones in the North Atlantic basin in 2017 had significant and widespread effects on the small islands of Caribbean as well as the United States, resulting in many deaths, displacement of communities, elevated rates of morbidity and mental health issues, as well as the long-term loss of electricity generation and distribution. These impacts have resulted in significant economic damage, which has exceeded the annual GDP of some small island developing States (46, 47).

Millions of people are already exposed to coastal flooding due to sea level rise and storms, particularly in cities. Projections of sea level rise remain uncertain (5), and may include significant non-linear responses, in part due to the contribution of land-based ice (48–50). Due to the time lag between increased emissions and higher sea levels, differences in mitigation at 1.5°C and 2.0°C, are relatively small compared with the uncertainty in the projections at 2050 or even 2100. Small differences can, however, have big impacts: an increase of 0.1m of sea level rise, for example, will expose an additional 10 million people to flooding (5) particularly those living in low-lying deltas and small islands (5, 51). Even with mitigation, adaptation remains essential, particularly as multi-metre sea level rise remains possible over several centuries for higher levels of temperature rise (5). Estimates of the net present value in 2008 of global aggregate damage costs (which would be incurred by 2200 if global warming is limited to 2.0°C) reach \$69 trillion (5). Damages from sea level rise alone contributes several trillion of dollars per annum (52). The net present value in 2008 of global aggregate damage costs associated with 1.5°C warming which would be incurred by 2200 if global warming is limited to 1.5C are less than those at 2.0°C, with comparable estimates around \$54 trillion in total (5).

Warming of 1°C has increased the frequency and scale of impacts on human health through changes to the intensity and frequency of heatwaves, droughts, floods and storms, as well as impacts on food quantity and nutritional quality (through increasing CO<sub>2</sub> concentrations) resulting in undernutrition or malnutrition in some regions (5, 43, 44). Multiple lines of evidence indicate that any further increases in GMST could have negative consequences for human health, mainly through the intensification of these risks (5, 53). Lower risks are projected at 1.5°C than 2.0°C of global warming for heat-related morbidity and mortality, and for ozone-related mortality if ozone precursor emissions remain high. Limiting global warming to 1.5°C would

result in 420 million fewer people being frequently exposed to ‘extreme heatwaves’ (defined by duration and intensity (54)) and about 65 million fewer people being exposed to ‘exceptional’ heatwaves as compared to conditions at 2.0°C GMST warming (55). Human health will also be affected by changes in the distribution and abundance of vector-borne diseases such as dengue fever and malaria, which are projected to increase with warming of 1.5°C and further at 2.0°C in most regions (5). Risks vary by human vulnerability, development pathways, and adaptation effectiveness (43, 44, 56). In some cases, human activities can lead to local amplification of heat risks from urban heat island effects in large cities (57, 58). More specific impacts of, and solutions to, climate change on cities are provided elsewhere (43, 56)

Global warming of 1.5°C will also affect human well-being through impacts on agriculture, industry and employment opportunities. For example, increased risks are projected for tourism in many countries, whereby changes in climate have the potential to affect the attractiveness and/or safety of destinations, particularly those dependent on seasonal tourism including sun, beach and snow sport destinations (5, 15). Businesses that have multiple locations or markets may reduce overall risk and vulnerability, although these options are likely to be reduced as stress and impacts increase in frequency and areal extent. Risks and adaptation options may lie in developing alternative business activities that are less dependent on environmental conditions. These risks become greater as warming increases to 2.0°C and pose serious challenges for a large number of countries dependent on tourism and related activities for national income (5).

Multiple lines of evidence also reveal that poverty and disadvantage are also correlated with warming to 1.0°C above pre-industrial period, with the projection of increasing risks as GMST increases from 1.0°C (today) to 1.5°C and higher (43, 44). In this regard, out-migration from agriculturally-dependent communities is positively correlated with global temperature although our understanding of the links between human migration and further warming of 1.5°C and 2.0°C is at an early stage (5). Similarly, risks to global aggregate economic growth due to climate change impacts are projected to be lower at 1.5°C than 2.0°C by the end of the century (5). The largest reduction in economic growth at 2.0°C compared to 1.5°C are projected for low- and middle-income countries and regions (the African continent, Southeast Asia, India, Brazil and Mexico). Countries in the tropics and Southern Hemisphere subtropics, are projected to

experience the largest negative impacts on economic growth if global warming increases from 1.5°C to 2.0°C above the pre-industrial period (5, 43, 44). The most perceptible impacts of climate change are likely to occur in tropical regions as GMST increases to 1.5 °C and eventually to 2°C above the pre-industrial period (59).

Table 1 summarizes the emergence of potential climate change ‘hotspots’ (i.e. areas where risks are large and growing rapidly) for a range of geographies and sectors (5). In all cases, these vulnerable regions show increasing risks as warming approaches 1.5°C and higher. Not all regions, however, face the same challenges. In the Arctic, for example, habitat loss is paramount, while changing temperature and precipitation regimes represent primary risks in the Mediterranean, Southern Africa, West Africa and the Sahel. These rapidly changing locations represent interactions across climate systems, ecosystems and socio-economic human systems, and are presented here to illustrate the extent to which risks can be avoided or reduced by achieving the 1.5°C global warming goal (as opposed to 2.0°C).

[Insert Table 1 here]

Trajectories toward hotspots can also involve significant non-linearities or tipping points. Tipping points refer to critical thresholds in a system that result in rapid systemic change when exceeded (5). The risks associated with 1.5°C or higher levels of global warming reveal relatively low risks for tipping points at 2.0°C but a substantial and growing set of risks as global temperature increases to 3°C or more above the pre-industrial (Table 2) (5). For example, increasing GMST to 3°C above the pre-industrial period substantially increases the risk of tipping points such as permafrost collapse, Arctic sea ice habitat loss, major reductions in crop production in Africa as well as globally, and persistent heat stress that is driving sharp increases in human morbidity and mortality (Table 2) (5).

[Insert Table 2 here]

## Solutions: scalability, feasibility and ethics

GMST will increase by 0.5°C between 2030 and 2052 and will multiply and intensify risks for natural and human systems across different geographies, vulnerabilities, development pathways, as well as adaptation and mitigation options (1, 43, 44, 56). To keep GMST to no more than 1.5°C above the pre-industrial period, the international community will need to bring GHG emissions to net zero by 2050 while adapting to the risks associated with an additional 0.5°C being added to GMST (3, 5). The impacts associated with limiting warming to 1.5°C, however, will be far less than those at 2.0°C or higher (Table 1, 2). Aiming to limit warming to 1.5°C is now a human imperative if escalating risks of dangerous if not catastrophic tipping points and climate change hotspots are to be avoided (2, 5).

An important conclusion of the IPCC special report on 1.5°C is that limiting GMST to 1.5°C or less is still possible (3, 60). This will require limiting GHG emissions to a budget of 420 Gt CO<sub>2</sub> for a 66% or higher probability of not exceeding 1.5°C (44). As global emissions are currently around 42 Gt CO<sub>2</sub> per year, pathways should bring CO<sub>2</sub> emissions to net zero over the next few decades (i.e. phase out fossil fuel use) alongside a substantial reduction (~35% relative to 2010) in emissions of methane and black carbon over the same time scale (44). The current set of national voluntary emission reduction pledges (Nationally Determined Contributions or NDCs), however, will not achieve the goals of the Paris Agreement (2, 61), particularly when considering the land-use sector (62). Instead, GMST is projected to increase by 3-4°C above the pre-industrial period (1, 44), posing serious levels of risk for natural and human systems (3, 5, 20).

The majority of pathways for achieving 1.5°C also require the carbon dioxide removal (CDR) from the atmosphere. Delays in bringing CO<sub>2</sub> emissions to net zero over the next 20-30 years will also increase the likelihood of pathways that exceed 1.5°C (so-called ‘overshoot’ scenarios) and hence a greater reliance on net negative emissions after mid-century if GMST to return to 1.5°C (Fig 2A). Technologies designed to remove CO<sub>2</sub> from the atmosphere are at an early stage of development, with many questions as to their feasibility and scalability (5). For example, bioenergy with carbon capture and storage (BECCS), afforestation and reforestation, blue carbon



(i.e. carbon sequestration by marine ecosystems and processes), soil carbon sequestration, direct capture, biochar (i.e. charcoal for burial in soils), and enhanced weathering, variously struggle from issues such as feasibility, scalability, and acceptability. These strategies are potentially in competition with each other. For example, BECCS would require approximately 18% of global land to sequester 12 Gt CO<sub>2</sub>/yr (5). This requirement is likely, however, to drive an accelerating the loss of primary forest and natural grassland which would increase GHG emissions (5). Early emission reductions plus measures to conserve land carbon stocks may reduce these effects. Policy options might limit the expansion of agriculture at the expense of natural ecosystems, and/or safeguard agricultural productivity from reductions due to BECCS and/or biofuel production (5).

There are CDR options, however, that do not rely as extensively on BECCS, but rather focus on afforestation and/or the restoration of natural ecosystems. It is feasible, for example, to limit warming to 1.5°C using strategies such as changing diets and promoting afforestation to remove CO<sub>2</sub> (3, 5, 43, 44). Negative consequences of afforestation such as monoculture plantations on local biodiversity might be countered by preferentially restoring natural ecosystems, re-establishing the ability of native grasslands, peatlands, forests, mangroves, kelp forests, and saltmarshes to sequester carbon. This creates a ‘win-win’ scenario in which both climate and biodiversity benefit, contributing to SDG 15 ‘Life on Land’: and hence, simultaneously making an enormous contribution to the goals of both CBD and UNFCCC. Compatible with this idea is the recent UN establishment of the 2020s as the ‘Decade of Restoration’, with the intention to build a global resolve to conserve biodiversity, increase its resilience to climate change, and use it to sequester up to a total of 26 GtC (63).

Extensive adaptation to 1.5°C of global warming or higher will be very important, especially if we have underestimated climate sensitivity. Developing socially-just and sustainable adaptation responses will be increasingly necessary to help natural and human systems to prepare and respond to rapid and complex changes in risk (43). The global adaptation stocktake instigated by the Paris Agreement will help accountability through documentation and mechanisms that inform enhancement at national levels (64, 65). It must also be acknowledged that there are limits to adaptation for natural and human systems (66) and hence subsequent loss and damage

(5, 67–69). For example, actions to restore ecosystems may not always be possible given available resources and it may not be feasible to protect all coastal regions from erosion and loss of land. These challenges mean that identifying, assessing, prioritizing and implementing adaptation options are very important for reducing the overall vulnerability to increasing climate-related risks as GMST increases. It has become increasingly clear that long-term solutions to climate change must also reduce disadvantage and poverty. Consequently, the recent IPCC Special Report pursued its findings in the context of ‘strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty’ (3). While previous reports recognized the importance of not aggravating disadvantage, few have specifically focused on solutions that involve multiple elements of climate change, sustainable development and poverty alleviation. For example, greater insights and knowledge are required to understand how multiple Sustainable Development Goals (SDGs) interact with each other, although many of these interactions are beneficially synergistic (70). Importantly, SDGs are far more easily reached at 1.5°C versus 2.0°C or more of global warming (43).

The important issue of ‘loss and damage’ also highlights the inequity between nations that have largely caused climate change (and have received the greatest benefits) and those who have not. This inequity is particularly important for least developed countries (LDCs) and small island developing States (SIDSs) that have contributed relatively little to global GHG emissions but now face disproportionate risks and harm from climate change, even at 1.5°C (67–69, 71). UNESCO has also emphasized the importance of ethics within a non-binding Declaration of Ethical Principles in Relation to Climate Change in 2017 (72). Specifically, this declaration states that “decision-making based on science is critically important for meeting the mitigation and adaptation challenges of a rapidly changing climate. Decisions should be based on, and guided by, the best available knowledge from natural and social sciences including interdisciplinary and transitionary science and by considering (as appropriate) local, traditional and indigenous knowledge”. These types of initiatives are especially important in the development of policies and actions that avoid inequalities that arise through exclusion and misinformation (61). A transformation toward climate-resilient and low-carbon societies needs to be done in a way that addresses the issue of justice and equity, through ensuring that trade-offs and synergies are identified and actioned (43).

## Conclusion

Warming of 1.0°C since the mid-20th century has fundamentally transformed our planet and its natural systems. Multiple lines of evidence reveal that a 1.5°C world will entail larger risks to both human and natural systems. The risks of a 2°C world are much greater. This places us at a critical time in human history where proportionate action taken today will almost certainly minimize the dangerous impacts of a changing climate for hundreds of millions of people. Our preliminary estimates suggest that the benefits of avoided damage by the year 2200 may exceed the costs of mitigation by a factor of four or five. Current NDCs for 2030 are insufficient to drive this even if followed by ‘very challenging increases in the scale and ambition of mitigation after 2030’ (44)(p 95), because models based on the current understanding of economic and technical dynamics cannot identify how to reduce GHG emissions to net zero by 2050 from the current NDC starting point in 2030. Rather, these ambitions are consistent with a global warming level of 3-4°C which means that immediate and transformative action is required between now and 2030 in order to greatly scale up current nationally stated plans for GHG reductions. Strategies for responding to climate change must be scalable to the challenges of climate change being faced today and into the future, while at the same time being feasible and fair. Given the scope and threats associated with climate change, there is an increasing need for large scale strategies such as the UN Climate Resilient Development Pathways (CRDP) or ‘Green New Deal’ (UNEP) if society is to avoid potentially catastrophic circumstances over the next few decades.

542 **References and Notes:**

- 543 1. M. R. Allen *et al.*, in *Framing and Context, In: Global warming of 1.5°C. An IPCC*  
544 *Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and*  
545 *related global greenhouse gas emission pathways, in the context of strengthening the*  
546 *global response to the* , V. Masson-Delmotte *et al.*, Eds. (World Meteorological  
547 Organization, Geneva, Switzerland, 2018).
  
- 548 2. UNFCCC, “Decision 1/CP.21 Adoption of the Paris Agreement” (Paris, France, 2015),  
549 (available at  
550 <https://unfccc.int/sites/default/files/resource/docs/2015/cop21/eng/10a01.pdf>).
  
- 551 3. IPCC, in *Global warming of 1.5°C. An IPCC Special Report on the impacts of global*  
552 *warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission*  
553 *pathways, in the context of strengthening the global response to the threat of climate*  
554 *change*, J. B. R. Matthews, Ed. (World Meteorological Organization, Geneva,  
555 Switzerland, 2018).
  
- 556 4. C. Hope, The Social Cost of Co2 from the Page09 Model. *SSRN Electron. J.* (2011),  
557 doi:10.2139/ssrn.1973863.
  
- 558 5. O. Hoegh-Guldberg *et al.*, in *Impacts of 1.5°C Global Warming on Natural and Human*  
559 *Systems In: Global warming of 1.5°C. An IPCC Special Report on the impacts of global*  
560 *warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission*  
561 *pathways, in the context of s*, V. Masson-Delmotte *et al.*, Eds. (World Meteorological  
562 Organization, Geneva, Switzerland, 2018).
  
- 563 6. IPCC, “Climate Change 2013: The Physical Science Basis. Working Group I Contribution  
564 to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change”  
565 (Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA,  
566 2013), , doi:<http://www.ipcc.ch/report/ar5/wg1/>.

- 567 7. M. D. Risser, M. F. Wehner, Attributable Human-Induced Changes in the Likelihood and  
568 Magnitude of the Observed Extreme Precipitation during Hurricane Harvey. *Geophys.*  
569 *Res. Lett.* **44**, 12457–12464 (2017).
  
- 570 8. G. Panthou, T. Vischel, T. Lebel, Recent trends in the regime of extreme rainfall in the  
571 Central Sahel. *Int. J. Climatol.* **34**, 3998–4006 (2014).
  
- 572 9. C. M. Taylor *et al.*, Frequency of extreme Sahelian storms tripled since 1982 in satellite  
573 observations. *Nature*. **544**, 475–478 (2017).
  
- 574 10. L. Gudmundsson, S. I. Seneviratne, X. Zhang, Anthropogenic climate change detected in  
575 European renewable freshwater resources. *Nat. Clim. Chang.* **7**, 813–816 (2017).
  
- 576 11. S. Mathbout *et al.*, Observed Changes in Daily Precipitation Extremes at Annual  
577 Timescale Over the Eastern Mediterranean During 1961–2012. *Pure Appl. Geophys.* **175**,  
578 3875–3890 (2018).
  
- 579 12. F. Raymond, P. Drobinski, A. Ullmann, P. Camberlin, Extreme dry spells over the  
580 Mediterranean Basin during the wet season: Assessment of HyMeX/Med-CORDEX  
581 regional climate simulations (1979–2009). *Int. J. Climatol.* **38**, 3090–3105 (2018).
  
- 582 13. D. A. Smale *et al.*, Marine heatwaves threaten global biodiversity and the provision of  
583 ecosystem services. *Nat. Clim. Chang.* **9**, 306–312 (2019).
  
- 584 14. J. A. Church *et al.*, in *Climate Change 2013: The Physical Science Basis. Contribution of*  
585 *Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on*  
586 *Climate Change*, T. F. Stocker *et al.*, Eds. (Cambridge University Press, Cambridge,  
587 United Kingdom and New York, NY, USA, 2013), pp. 1137–1216.
  
- 588 15. O. Hoegh-Guldberg *et al.*, in *Climate Change 2014: Impacts, Adaptation, and*  
589 *Vulnerability. Part B: Regional Aspects. Contribution of Working Group II to the Fifth*

- 590        *Assessment Report of the Intergovernmental Panel on Climate Change*, V. R. Barros et al.,  
591        Eds. (Cambridge University Press, Cambridge, United Kingdom and New York, NY,  
592        USA, 2014), pp. 1655–1731.
- 593    16.    Y. Yin, D. Ma, S. Wu, Climate change risk to forests in China associated with warming.  
594        *Sci. Rep.* **8**, 1–13 (2018).
- 595    17.    M. Turco *et al.*, Exacerbated fires in Mediterranean Europe due to anthropogenic warming  
596        projected with non-stationary climate-fire models. *Nat. Commun.* **9**, 1–9 (2018).
- 597    18.    B. Sultan *et al.*, Assessing climate change impacts on sorghum and millet yields in the  
598        Sudanian and Sahelian savannas of West Africa. *Environ. Res. Lett.* **8**, 014040 (2013).
- 599    19.    X. Ren, Y. Lu, B. C. O'Neill, M. Weitzel, Economic and biophysical impacts on  
600        agriculture under 1.5 °C and 2 °C warming. *Environ. Res. Lett.* **13**, 115006 (2018).
- 601    20.    IPCC, *Climate Change 2013 - The Physical Science Basis* (Cambridge, United Kingdom  
602        and New York, NY, USA, 2014).
- 603    21.    B. C. Lister, A. Garcia, Climate-driven declines in arthropod abundance restructure a  
604        rainforest food web. *Proc. Natl. Acad. Sci. U. S. A.* **115**, E10397–E10406 (2018).
- 605    22.    B. Martay *et al.*, An indicator highlights seasonal variation in the response of Lepidoptera  
606        communities to warming. *Ecol. Indic.* **68**, 126–133 (2016).
- 607    23.    T. P. Hughes *et al.*, Coral reefs in the Anthropocene. *Nature.* **546**, 82–90 (2017).
- 608    24.    T. P. Hughes, J. T. Kerry, T. Simpson, Large-scale bleaching of corals on the Great  
609        Barrier Reef. *Ecology.* **99**, 501 (2018).
- 610    25.    L. Cheng, J. Abraham, Z. Hausfather, K. E. Trenberth, How fast are the oceans warming?

- 611 *Science*. **363**, 128–129 (2019).
- 612 26. H.-O. Pörtner, C. Bock, F. C. Mark, Oxygen- and capacity-limited thermal tolerance:  
613 bridging ecology and physiology. *J. Exp. Biol.* **220**, 2685–2696 (2017).
- 614 27. S. G. Dove *et al.*, Future reef decalcification under a business-as-usual CO<sub>2</sub> emission  
615 scenario. *Proc. Natl. Acad. Sci.* **110**, 15342–15347 (2013).
- 616 28. S. I. Seneviratne *et al.*, The many possible climates from the Paris Agreement’s aim of 1.5  
617 °C warming. *Nature*. **558**, 41–49 (2018).
- 618 29. S. I. Seneviratne, M. G. Donat, A. J. Pitman, R. Knutti, R. L. Wilby, Allowable CO<sub>2</sub>  
619 emissions based on regional and impact-related climate targets. *Nature*. **529**, 477–83  
620 (2016).
- 621 30. P. Greve, L. Gudmundsson, S. I. Seneviratne, Regional scaling of annual mean  
622 precipitation and water availability with global temperature change. *Earth Syst. Dyn.* **9**,  
623 227–240 (2018).
- 624 31. T. Ozturk, Z. P. Ceber, M. Türkeş, M. L. Kurnaz, Projections of climate change in the  
625 Mediterranean Basin by using downscaled global climate model outputs. *Int. J. Climatol.*  
626 **35**, 4276–4292 (2015).
- 627 32. S. D. Polade, A. Gershunov, D. R. Cayan, M. D. Dettinger, D. W. Pierce, Precipitation in  
628 a warming world: Assessing projected hydro-climate changes in California and other  
629 Mediterranean climate regions. *Sci. Rep.* **7**, 10783 (2017).
- 630 33. G. Naumann *et al.*, Global Changes in Drought Conditions Under Different Levels of  
631 Warming. *Geophys. Res. Lett.* **45**, 3285–3296 (2018).
- 632 34. W. Liu *et al.*, Global Freshwater Availability Below Normal Conditions and Population



- 633            Impact Under 1.5 and 2 degrees C Stabilization Scenarios. *Geophys. Res. Lett.* **45**, 9803–  
634            9813 (2018).
  
- 635    35.    L. Lin *et al.*, Additional Intensification of Seasonal Heat and Flooding Extreme Over  
636            China in a 2 degrees C Warmer World Compared to 1.5 degrees C. *EARTH'S Futur.* **6**,  
637            968–978 (2018).
  
- 638    36.    G. D. Madakumbura *et al.*, Event-to-event intensification of the hydrologic cycle from  
639            1.5 °C to a 2 °C warmer world. *Sci. Rep.* **9**, 3483 (2019).
  
- 640    37.    J. García Molinos, M. T. Burrows, E. S. Poloczanska, Ocean currents modify the coupling  
641            between climate change and biogeographical shifts. *Sci. Rep.* **7**, 1332 (2017).
  
- 642    38.    J. A. Screen, Arctic sea ice at 1.5 and 2 °C. *Nat. Clim. Chang.* **8**, 362–363 (2018).
  
- 643    39.    K. B. Karnauskas *et al.*, Freshwater Stress on Small Island Developing States: Population  
644            Projections and Aridity Changes at 1.5°C and 2°C. *Reg. Environ. Chang.*, 1–10 (2018).
  
- 645    40.    A. C. Ruane, M. M. Phillips, C. Rosenzweig, Climate shifts within major agricultural  
646            seasons for +1.5 and +2.0 °C worlds: HAPPI projections and AgMIP modeling scenarios.  
647            *Agric. For. Meteorol.* **259**, 329–344 (2018).
  
- 648    41.    B. Liu *et al.*, Global wheat production with 1.5 and 2.0°C above pre-industrial warming.  
649            *Glob. Chang. Biol.* **25**, 1428–1444 (2019).
  
- 650    42.    K. Rhiney, A. Eitzinger, A. D. Farrell, S. D. Prager, Assessing the implications of a 1.5 °C  
651            temperature limit for the Jamaican agriculture sector. *Reg. Environ. Chang.* **18**, 2313–  
652            2327 (2018).
  
- 653    43.    J. Roy *et al.*, in *Sustainable development, poverty eradication and reducing inequalities*;  
654            *In: Global warming of 1.5°C. An IPCC Special Report on the impacts of global warming*

- 655        *of 1.5°C above pre-industrial levels and related global greenhouse gas emission*  
656        *pathways, in the c*, V. Masson-Delmotte et al., Eds. (World Meteorological Organization,  
657        Geneva, Switzerland, 2018).
- 658    44.    J. Rogelj *et al.*, in *Mitigation pathways compatible with 1.5°C in the context of sustainable*  
659        *development; In: Global warming of 1.5°C. An IPCC Special Report on the impacts of*  
660        *global warming of 1.5°C above pre-industrial levels and related global greenhouse gas*  
661        *emission pathw*, V. Masson-Delmotte et al., Eds. (World Meteorological Organization,  
662        Geneva, Switzerland, 2018).
- 663    45.    K. Bhatia *et al.*, Projected Response of Tropical Cyclone Intensity and Intensification in a  
664        Global Climate Model. *J. Clim.* **31**, 8281–8303 (2018).
- 665    46.    J. M. Shultz, J. P. Kossin, C. Ettman, P. L. Kinney, S. Galea, The 2017 perfect storm  
666        season, climate change, and environmental injustice. *Lancet Planet. Heal.* **2**, e370–e371  
667        (2018).
- 668    47.    J. M. Shultz *et al.*, Risks, Health Consequences, and Response Challenges for Small-  
669        Island-Based Populations: Observations From the 2017 Atlantic Hurricane Season.  
670        *Disaster Med. Public Health Prep.* **13**, 5–17 (2019).
- 671    48.    T. L. Edwards *et al.*, Revisiting Antarctic ice loss due to marine ice-cliff instability.  
672        *Nature.* **566**, 58–64 (2019).
- 673    49.    D. F. Martin, S. L. Cornford, A. J. Payne, Millennial-Scale Vulnerability of the Antarctic  
674        Ice Sheet to Regional Ice Shelf Collapse. *Geophys. Res. Lett.* **46**, 1467–1475 (2019).
- 675    50.    F. Pattyn *et al.*, The Greenland and Antarctic ice sheets under 1.5 °C global warming. *Nat.*  
676        *Clim. Chang.* **8**, 1053–1061 (2018).
- 677    51.    M. I. Vousdoukas *et al.*, Global probabilistic projections of extreme sea levels show

- 678 intensification of coastal flood hazard. *Nat. Commun.* **9**, 2360 (2018).
- 679 52. S. Jevrejeva, L. P. Jackson, A. Grinsted, D. Lincke, B. Marzeion, Flood damage costs  
680 under the sea level rise with warming of 1 . 5 ° C and 2 ° C. *Environ. Res. Lett.* **13** (2018).
- 681 53. M. B. Sylla, A. Faye, F. Giorgi, A. Diedhiou, H. Kunstmann, Projected Heat Stress Under  
682 1.5 degrees C and 2 degrees C Global Warming Scenarios Creates Unprecedented  
683 Discomfort for Humans in West Africa. *EARTHS Futur.* **6**, 1029–1044 (2018).
- 684 54. S. Russo, J. Sillmann, E. M. Fischer, Top ten European heatwaves since 1950 and their  
685 occurrence in the coming decades. *Environ. Res. Lett.* **10**, 124003 (2015).
- 686 55. A. Dosio, L. Mentaschi, E. M. Fischer, K. Wyser, Extreme heat waves under 1.5 °C and 2  
687 °C global warming. *Environ. Res. Lett.* **13**, 054006 (2018).
- 688 56. H. de Coninck *et al.*, in *Strengthening and implementing the global response. In: Global*  
689 *warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C*  
690 *above pre-industrial levels and related global greenhouse gas emission pathways, in the*  
691 *context of strengthen*, V. Masson-Delmotte *et al.*, Eds. (World Meteorological  
692 Organization, Geneva, Switzerland, 2018).
- 693 57. J. Mika *et al.*, Impact of 1.5 K global warming on urban air pollution and heat island with  
694 outlook on human health effects. *Curr. Opin. Environ. Sustain.* **30** (2018), pp. 151–159.
- 695 58. C. R. O’Lenick *et al.*, Urban heat and air pollution: A framework for integrating  
696 population vulnerability and indoor exposure in health risk analyses. *Sci. Total Environ.*  
697 **660**, 715–723 (2019).
- 698 59. A. D. King, L. J. Harrington, The Inequality of Climate Change From 1.5 to 2°C of Global  
699 Warming. *Geophys. Res. Lett.* **45**, 5030–5033 (2018).

- 700 60. K. Zickfeld *et al.*, Current fossil fuel infrastructure does not yet commit us to 1.5 °C  
701 warming. *Nat. Commun.* **10**, 1–10 (2019).
- 702 61. C. Brown, P. Alexander, A. Arneth, I. Holman, M. Rounsevell, Achievement of Paris  
703 climate goals unlikely due to time lags in the land system. *Nat. Clim. Chang.* **9**, 203–208  
704 (2019).
- 705 62. A. B. Harper *et al.*, Land-use emissions play a critical role in land-based mitigation for  
706 Paris climate targets. *Nat. Commun.* **9**, 2938 (2018).
- 707 63. UN Environment, New UN Decade on Ecosystem Restoration offers unparalleled  
708 opportunity for job creation, food security and addressing climate change (2019),  
709 (available at [https://www.unenvironment.org/news-and-stories/press-release/new-un-](https://www.unenvironment.org/news-and-stories/press-release/new-un-decade-ecosystem-restoration-offers-unparalleled-opportunity)  
710 [decade-ecosystem-restoration-offers-unparalleled-opportunity](https://www.unenvironment.org/news-and-stories/press-release/new-un-decade-ecosystem-restoration-offers-unparalleled-opportunity)).
- 711 64. B. Craft, S. Fisher, Measuring the adaptation goal in the global stocktake of the Paris  
712 Agreement. *Clim. Policy.* **18**, 1203–1209 (2018).
- 713 65. E. L. Tompkins, K. Vincent, R. J. Nicholls, N. Suckall, Documenting the state of  
714 adaptation for the global stocktake of the Paris Agreement. *Wiley Interdiscip. Rev. Clim.*  
715 *Chang.* **9**, e545 (2018).
- 716 66. K. Dow *et al.*, Limits to adaptation. *Nat. Clim. Chang.* **3**, 305–307 (2013).
- 717 67. K. E. McNamara, G. Jackson, Loss and damage: A review of the literature and directions  
718 for future research. *Wiley Interdiscip. Rev. Clim. Chang.* **10**, e564 (2019).
- 719 68. R. Mechler, L. M. Bouwer, T. Schinko, S. Surminski, J. Linnerooth-Bayer, Eds., *Loss and*  
720 *Damage from Climate Change* (Springer International Publishing, Cham, 2019;  
721 <http://link.springer.com/10.1007/978-3-319-72026-5>), *Climate Risk Management, Policy*  
722 *and Governance*.

69. E. Boyd, R. A. James, R. G. Jones, H. R. Young, F. E. L. Otto, A typology of loss and damage perspectives. *Nat. Clim. Chang.* **7**, 723–729 (2017).
70. M. Nilsson *et al.*, Mapping interactions between the sustainable development goals: lessons learned and ways forward. *Sustain. Sci.* **13**, 1489–1503 (2018).
71. A. Thomas, L. Benjamin, Management of loss and damage in small island developing states: implications for a 1.5 °C or warmer world. *Reg. Environ. Chang.* **18**, 2369–2378 (2018).
72. UNESCO, “Declaration of Ethical Principles Climate Change” (2017), (available at <https://unesdoc.unesco.org/ark:/48223/pf0000260129>).
73. R. Wartenburger *et al.*, Changes in regional climate extremes as a function of global mean temperature: an interactive plotting framework. *Geosci. Model Dev.* **10**, 3609–3634 (2017).

## Acknowledgments:

The authors are grateful for the support provided by the Intergovernmental Panel on Climate Change (IPCC), particularly that of the Technical Support Units for Working Groups I and II, as well as the large number of Contributing Authors and Science Officers involved in the IPCC Special Report on the “Impacts of Global Warming of 1.5°C above Pre-Industrial Levels and Related Global Greenhouse Gas Emission Pathways, in the Context of Strengthening the Global Response to the Threat of Climate Change, Sustainable Development, and Efforts to Eradicate Poverty” (3). **Author contributions:** All authors volunteered their time and contributed their time in producing this publication and the underlying Special Report from the IPCC (3). **Funding:** OHG acknowledges support from the Australia Research Council (ARC Laurette Fellowship and ARC Centre for Excellence in Coral Reef Studies). SIS acknowledges partial support from the European Research Council (ERC) ‘DROUGHT-HEAT’ project funded by the European Community’s Seventh Framework Programme (grant agreement FP7-IDEAS-ERC-617518). **Competing Interests:** The findings, interpretations, and conclusions expressed in the work do not necessarily reflect the views of The World Bank, its Board of Executive Directors, or the governments they represent. KE established a sole proprietor consulting company (ClimAdapt, LLC) more than a decade ago, and occasionally consults with international entities and development partners on managing the health risks of climate change. No other competing interests have been declared by the authorship.

## List of Supplementary Materials:

Material and Methods

767 Table 1: Climate change ‘hotspots’ are expanding. Emergence and intensity of climate change ‘hotspots’ under different degrees of  
768 global warming (summary, updated, Table 3.6 from (5), see text in 3.5.4 for supporting literature and discussion; not intended to be all  
769 inclusive). Calibrated uncertainty language is as defined by the Intergovernmental Panel on Climate Change (3).  
770

Region and/or Phenomenon	Warming of 1.5°C or less	Warming of 1.5°C–2°C	Warming of up to 3°C
<b>Arctic sea ice</b>	<u>Arctic summer sea ice</u> is <i>likely</i> to be maintained  <u>Habitat losses</u> for organisms such as polar bears, whales, seals and sea birds  <u>Benefits</u> for Arctic fisheries	<u>The risk of an ice-free Arctic</u> in summer is about 50% or higher  <u>Habitat losses</u> for organisms such as polar bears, whales, seals and sea birds may be critical if summers are ice free.  <u>Benefits</u> for Arctic fisheries	<u>The Arctic</u> is <i>very likely</i> to be ice free in summer  <u>Critical habitat losses</u> for organisms such as polar bears, whales, seals and sea birds  <u>Benefits</u> for Arctic fisheries
<b>Arctic land regions</b>	<u>Cold extremes warm</u> by a factor of 2–3, reaching up to 4.5°C ( <i>high confidence</i> )  <u>Biome shifts</u> in the tundra and permafrost deterioration are <i>likely</i>	<u>Cold extremes warm</u> by as much as 8°C ( <i>high confidence</i> )  <u>Larger intrusions of trees and shrubs</u> in the tundra than under 1.5°C of warming are likely; larger but constrained losses in permafrost <i>are likely</i>	<u>Drastic regional warming</u> is <i>very likely</i>  <u>A collapse in permafrost may occur</u> ( <i>low confidence</i> ); a drastic biome shift from tundra to boreal forest is possible ( <i>low confidence</i> )
<b>Alpine regions</b>	<u>Severe shifts</u> in biomes are <i>likely</i>	<u>Even more severe shifts</u> are <i>likely</i>	<u>Critical losses</u> in alpine habitats are <i>likely</i>
<b>Southeast Asia</b>	<u>Risks for increased flooding</u> related to sea level rise  <u>Increases, heavy precipitation</u> events  <u>Significant risks</u> of crop yield reductions are avoided	<u>Higher risks of increased flooding</u> related to sea level rise ( <i>medium confidence</i> )  <u>Stronger increases, heavy precipitation</u> events ( <i>medium confidence</i> )  <u>One-third decline</u> in per capita crop production ( <i>medium confidence</i> )	<u>Substantial increases in risks</u> related to flooding from sea level rise  <u>Substantial increase</u> in heavy precipitation and high-flow events  <u>Substantial reductions</u> in crop yield



<b>Mediterranean</b>	<p><u>Increase in probability of extreme drought</u> (<i>medium confidence</i>)</p> <p><i>Medium confidence</i> in reduction in runoff of about 9% (likely range 4.5–15.5%)</p> <p><u>Risk of water deficit</u> (<i>medium confidence</i>)</p>	<p><u>Robust increase</u> in probability of extreme drought (<i>medium confidence</i>)</p> <p><i>Medium confidence</i> in further reductions (about 17%) in runoff (likely range 8–28%)</p> <p><u>Higher risks of water deficit</u> (<i>medium confidence</i>)</p>	<p><u>Robust and large increases</u> in extreme drought.</p> <p><u>Substantial reductions in precipitation</u> and in runoff (<i>medium confidence</i>)</p> <p><u>Very high risks</u> of water deficit (<i>medium confidence</i>)</p>
<b>West Africa &amp; the Sahel</b>	<p><u>Increases in the number</u> of hot nights and longer and more frequent heatwaves are <i>likely</i></p> <p><u>Reduced maize and sorghum</u> production is <i>likely</i>, with area suitable for maize production reduced by as much as 40%</p> <p><u>Increased risks of undernutrition</u></p>	<p>Further increases in number of hot nights and longer and more frequent heatwaves are likely</p> <p>Negative impacts on maize and sorghum production likely larger than at 1.5°C; <i>medium confidence</i> that vulnerabilities to food security in the African Sahel will be higher at 2.0°C compared to 1.5°C</p> <p><u>Higher risks of undernutrition</u></p>	<p>Substantial increases in the number of hot nights and heatwave duration and frequency (<i>very likely</i>)</p> <p>Negative impacts on crop yield may result in major regional food insecurities (<i>medium confidence</i>)</p> <p><u>High risks of undernutrition</u></p>
<b>Southern Africa</b>	<p><u>Reductions in water availability</u> (<i>medium confidence</i>)</p> <p><u>Increases in number of hot nights</u> and longer and more frequent heatwaves (<i>high confidence</i>),</p> <p><u>High risks of increased mortality</u> from heatwaves</p> <p><u>High risk of undernutrition</u> in communities dependent on dryland agriculture and livestock</p>	<p><u>Larger reductions in rainfall</u> and water availability (<i>medium confidence</i>)</p> <p><u>Further increases in number of hot nights</u> and longer and more frequent heatwaves (<i>high confidence</i>), associated increases in risks of <u>increased mortality from heatwaves</u> compared to 1.5°C warming (<i>high confidence</i>)</p> <p><u>Higher risks of undernutrition</u> in communities dependent on dryland agriculture and livestock</p>	<p><u>Large reductions in rainfall</u> and water availability (<i>medium confidence</i>)</p> <p><u>Drastic increases</u> in the number of hot nights, hot days and heatwave duration and frequency to impact substantially on agriculture, livestock and human health and mortality (<i>high confidence</i>)</p> <p><u>Very high risks of undernutrition</u> in communities dependent on dryland agriculture and livestock</p>

<b>Tropics</b>	<p><u>Increases in the number of hot days</u> and hot nights as well as longer and more frequent heatwaves (<i>high confidence</i>)</p> <p><u>Risks to tropical crop yields</u> in West Africa, Southeast Asia and Central and South America are significantly less than under 2.0°C of warming</p>	<p><u>The largest increase in hot days</u> under 2.0°C compared to 1.5°C is projected for the tropics.</p> <p><u>Risks to tropical crop yields in West Africa, Southeast Asia and Central and South America</u> could be extensive</p>	<p><u>Oppressive temperatures</u> and accumulated heatwave duration <i>very likely</i> to directly impact human health, mortality and productivity</p> <p><u>Substantial reductions</u> in crop yield <i>very likely</i></p>
<b>Small islands</b>	<p><u>Land of 60,000 less people</u> exposed by 2150 on SIDS compared to impacts under 2.0°C of global warming</p> <p><u>Risks for coastal flooding</u> reduced by 20–80% for SIDS compared to 2.0°C of global warming</p> <p><u>Freshwater stress</u> reduced by 25% as compared to 2.0°C</p> <p><u>Increase in the number of warm days</u> for SIDS in the tropics</p> <p><u>Persistent heat stress</u> in cattle avoided</p> <p><u>Loss of 70–90% of coral reefs</u></p>	<p><u>Tens of thousands of people displaced</u> owing to inundation of SIDS</p> <p><u>High risks</u> for coastal flooding and increased frequency of extreme water-level events</p> <p><u>Freshwater stress</u> from projected aridity</p> <p><u>Further increase</u> of ca. 70 warm days/year</p> <p><u>Persistent heat stress</u> in cattle in SIDS</p> <p><u>Loss of most coral reefs</u> and weaker remaining structures owing to ocean acidification (i.e. less coastal protection)</p>	<p><u>Substantial and widespread impacts</u> through inundation of SIDS, coastal flooding, freshwater stress, persistent heat stress and loss of most coral reefs (<i>very likely</i>)</p> <p><u>Risk of multi-meter sea level</u> rise due to ice sheet instability</p>
<b>Fynbos biome</b>	<p><u>About 30% of suitable climate area</u> lost (<i>medium confidence</i>)</p>	<p><u>Increased losses (about 45%)</u> of suitable climate area (<i>medium confidence</i>)</p>	<p><u>Up to 80% of suitable climate area</u> lost (<i>medium confidence</i>)</p>

Table 2: Increased warming increases risks of exceeding tipping points. Summary of enhanced risks in the exceedance of regional tipping points under different global temperature goals (summary, Table 3.7 from(5), see text in 3.5.5 for supporting literature and discussion; updated, not intended to be exhaustive).

Tipping point	Warming of 1.5°C or less	Warming of 1.5°C–2°C	Warming of up to 3°C
<b>Arctic sea ice</b>	<u>Arctic summer sea ice</u> is <i>likely</i> to be maintained  <u>Sea ice changes</u> reversible under suitable climate restoration	<u>The risk of an ice-free Arctic</u> in summer is about 50% or higher  <u>Sea ice changes</u> reversible under suitable climate restoration	<u>Arctic</u> is <i>very likely</i> to be ice free in summer  <u>Sea ice changes</u> reversible under suitable climate restoration
<b>Tundra</b>	<u>Decrease</u> in number of growing degree days below 0°C  <u>Abrupt</u> increases in tree cover are <i>unlikely</i>	<u>Further</u> decreases in number of growing degree days below 0°C  <u>Abrupt</u> increases in tree cover are <i>unlikely</i>	     <u>Potential</u> for an abrupt increase in tree fraction ( <i>low confidence</i> )
<b>Permafrost</b>	17–44% reduction in permafrost <u>Approximately 2 million km<sup>2</sup></u> more permafrost maintained than under 2.0°C of global warming ( <i>medium confidence</i> )  Irreversible loss of stored carbon	28–53% reduction in permafrost with     <u>Irreversible loss</u> of stored carbon	     <u>Potential</u> for permafrost collapse ( <i>low confidence</i> )
<b>Asian monsoon</b>	<u>Low confidence</u> in projected changes	<u>Low confidence</u> in projected changes	<u>Increases in the intensity of</u> monsoon precipitation <i>likely</i>
<b>West African monsoon &amp; Sahel</b>	<u>Uncertain changes</u> ; <i>unlikely</i> that a tipping point is reached	<u>Uncertain changes</u> ; <i>unlikely</i> that tipping point is reached	<u>Strengthening of monsoon</u> with wettening and greening of the Sahel and Sahara ( <i>low confidence</i> )  <u>Negative associated impacts</u> through increases in extreme temperature events
<b>Rainforests</b>	<u>Reduced biomass</u> , deforestation and fire increases pose uncertain risks to forest dieback	<u>Larger biomass reductions</u> than under 1.5°C of warming; deforestation and fire increases pose uncertain risks to forest dieback	<u>Reduced extent of tropical rainforest</u> in Central America and large replacement of rainforest and savanna grassland  <u>Potential tipping point</u> leading to pronounced forest dieback ( <i>medium confidence</i> )

<b>Coral reefs</b>	<u>Increased mass coral bleaching and mortality</u> – decline in abundance to 10-30% of values of present day by 1.0°C ( <i>high confidence</i> )	<u>High mortality</u> - corals decrease to very low levels (<1%), impacts on organisms that dependent on coral reefs for habitat (fish, biodiversity, <i>high confidence</i> ).	<u>Irreversible changes occur</u> with tipping point around 2°C–2.5°C – reefs are no longer resemble coral reef ecosystems – recovery potential very low ( <i>medium confidence</i> ).
<b>Boreal forests</b>	<u>Increased tree mortality</u> at southern boundary of boreal forest ( <i>medium confidence</i> )	<u>Further increases in tree mortality</u> at southern boundary of boreal forest ( <i>medium confidence</i> )	<u>Potential tipping point</u> at 3°C–4°C for significant dieback of boreal forest ( <i>low confidence</i> )
<b>Heatwaves, unprecedented heat and human health</b>	<u>Continued increase</u> in occurrence of potentially deadly heatwaves ( <i>likely</i> )	<u>Substantial increase</u> in potentially deadly heatwaves ( <i>likely</i> )  <u>More than 350 million more people</u> exposed to deadly heat by 2050 under a midrange population growth scenario ( <i>likely</i> )  <u>Annual occurrence</u> of heatwaves similar to the deadly 2015 heatwaves in India and Pakistan ( <i>medium confidence</i> )	<u>Further increases</u> in potentially deadly heatwaves ( <i>very likely</i> )
<b>Agricultural systems: key staple crops</b>	<u>Global maize crop reductions</u> of about 10%	<u>Larger reductions in maize crop</u> production than under 1.5°C of about 15%	<u>Drastic reductions in maize crop globally</u> and in Africa ( <i>high confidence</i> ) <u>potential tipping point</u> for collapse of maize crop in some regions ( <i>low confidence</i> )
<b>Livestock in the tropics and subtropics</b>	<u>Increased heat stress</u>	<u>Onset of persistent heat stress</u> ( <i>medium confidence</i> )	<u>Persistent heat stress</u> <i>likely</i>

777 **Figure captions:**

778 Figure 1. Impacts are already serious at 1.0°C of global warming. Increases in Global Mean  
 779 Surface Temperature (GMST) of 1.0°C have already had major impacts on natural and human  
 780 systems. Examples include: A. Increased temperatures and dryness in the Mediterranean region  
 781 is driving longer and more intense fire seasons with serious impacts on people, infrastructure and  
 782 natural ecosystems. Image shows tragic devastation of fire in the Greek village of Mati Greece in  
 783 July 25, 2018. B. Evidence of ice sheet disintegration is increasing (here showing a 30 km  
 784 fracture across the Pine Island Glacier which is associated with the Western Antarctic Ice sheet,  
 785 WAIS). The fracture (see arrow) appeared in mid-October 2011 and has increased concern that  
 786 we may be approaching a tipping point with respect to disintegration of the WAIS. C. Many low-  
 787 lying countries such as the Maldives experience flooding and will be at an increased threat from  
 788 sea level rise and strengthening storms over time. D. Many insects and birds have shifted  
 789 reproductive events or migration to early times in the season as conditions have warmed. Image  
 790 credits: A. ‘Lotus R’, <https://www.flickr.com/photos/66012345@N00/964251167/>; B. Image  
 791 credits: NASA/GSFC/METI/ERSDAC/JAROS and U.S./Japan ASTER Science Team Last  
 792 Updated: Aug. 7, 2017, C. Male, Maldives (O. Hoegh-Guldberg) and D. Semipalmated Sand  
 793 Piper (*Calidris pusilla*, Creative Commons (CC BY-SA 3.0, GNU Free Documentation License)





Figure 2. There is still time to restrain global warming to 1.5°C. A. Action on climate change can still result in stable or even decreasing global temperatures, although variability around projections is substantial. Strategies that include ‘overshoot’ (red dashed line, illustrative of a very high level of overshoot) require as yet early stage technologies to ensure that overshoot is kept as short as possible. Also, the larger overshoot, the higher the risk of irreversible change in affected systems. B. Responses to changing conditions (shown here as a thermal performance curve) are typically tilted to the right with a steep decline in performance such as growth, towards high temperature extremes. Beyond a thermal optimum,  $T_{opt}$ , performance begins to decline beyond the *Pejus* temperature,  $T_p$ . A critical temperature,  $T_c$ , characterizes a low level of performance and time limited passive endurance when, as in ectothermic animals, oxygen supply capacity becomes insufficient to cover oxygen supply, or, as in corals, a symbiosis between corals and their dinoflagellate symbionts suddenly breaks down (coral bleaching) and corals go from appearing healthy to experiencing large scale mortality over days-to-weeks. Accordingly, the high  $T_c$  characterizes a temperature of high responsiveness to small increases in temperature extremes, such as by 0.5°C, especially, if some life stages have a narrow thermal range indicating high vulnerability. Figure 2B redrawn from with permission of the Journal of Experimental Biology (26).



813

814



815 Figure 3 Changes varies geographically at GMST of 1.5°C. Projected changes are shown in A.  
 816 Mean temperature, B. Mean precipitation, C. Number of hot days (NHD; 10% warmest days), D.  
 817 Temperature of hottest day (TXx), E. Temperature of coldest night (TNn), and F. Change in  
 818 extreme precipitation (Rx5day). Conditions are projected for 1.5°C (left-hand column) and  
 819 2.0°C (middle-hand column) of global warming compared to the pre-industrial period (1861–  
 820 1880), with the difference between 1.5°C and 2.0°C of global warming being shown in the third  
 821 column. Cross-hatching highlights areas where at least two-thirds of the models agree on the sign  
 822 of change as a measure of robustness (18 or more out of 26). Values were assessed from the  
 823 transient response over a 10-year period at a given warming level, based on Representative  
 824 Concentration Pathway (RCP) 8.5 Coupled Model Intercomparison Project Phase 5 (CMIP5)  
 825 model simulations (5)(3); adapted from (29, 73); see Supplementary Material 3.SM.2 in (5).  
 826 Figure is a composite of Figure 3.3 (here, Fig 3A-B), Figure 3.7 (here, Fig 3C), and Figure 3.4  
 827 (here, Fig 3D-F) in (5). Use of figures with permission of the IPCC.

