

C: Physical Processes in Nanomaterials and Nanostructures

**Time-Resolved Structural Dynamics of
Extended π -Electron Porphyrin Nanoring**

Giovanni Bressan, Martin D. Peeks, Harry L. Anderson, Stephen R. Meech, and Ismael Heisler

J. Phys. Chem. C, **Just Accepted Manuscript** • DOI: 10.1021/acs.jpcc.9b07494 • Publication Date (Web): 15 Oct 2019Downloaded from pubs.acs.org on October 17, 2019**Just Accepted**

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

1
2
3
4
5
6
7
8
9

Time-Resolved Structural Dynamics of Extended π -Electron Porphyrin Nanoring

10 Giovanni Bressan¹, Martin D. Peeks², Harry L. Anderson,² Stephen R. Meech¹ and Ismael A.
11
12 Heisler^{3,*}
13

14
15
16 ¹*School of Chemistry, Norwich Research Park, University of East Anglia, Norwich*
17
18 *NR4 7TJ, United Kingdom*
19

20
21 ²*Department of Chemistry, University of Oxford, Chemistry Research Laboratory,*
22
23 *Oxford OX1 3TA, United Kingdom*
24

25
26 ³*Departamento de Física, Universidade Federal do Paraná, Caixa Postal 19044,*
27
28 *81531-990 Curitiba, Parana, Brazil*
29

30
31
32
33
34
35
36
37
38
39
40
41
42
43 *Corresponding author. Email: heisler@fisica.ufpr.br
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Molecular structure design inspired by naturally occurring light harvesting systems has been intensely pursued over the last couple of decades. Interesting new structures include the π -conjugated porphyrin nanorings, which show promising features such as ultrafast excited-state delocalization, leading to suppressed radiative rates, superradiance with increasing temperature and energy transfer times comparable to their natural counterparts. An important question to be addressed in such systems is the role and time scale of structural motions and how they affect excited-state delocalization. Here it is shown that porphyrin nanorings which are not rigidified by a template are structurally heterogeneous in the ground state and evolve dynamically on a tens of picoseconds timescale. In the excited state a structural relaxation of the porphyrin nanorings is observed, on a picosecond timescale. Furthermore ultrafast excitation delocalization is observed, by anisotropy measurements, being insensitive to structural motions of the nanorings.

TOC graphics

Introduction

Molecular structures based on π -conjugated chromophores present attractive features such as semiconducting character^{1,2,3} and nonlinear optical properties^{4,5,6,7}. Various applications rely on such features including electroluminescent devices^{8,9}, solar cells^{10,11}, optical switches^{12,13}, and fluorescent biosensors^{14,15}. The synthesis of conjugated porphyrin multi-chromophore structures tailored for specific applications has witnessed great progress during the last couple of decades^{16,17}. A significant amount of work has been done on π -conjugated molecular systems comprised of extended linear one-dimensional oligomers^{18,19,20}, for which their physical properties correlate well with the length of the conjugated chain²¹. However, due to their finite size, linear oligomers present end-effects that can alter their physical behaviour^{7,22,23}. These effects can be overcome by working with shape persistent and cyclically conjugated two-dimensional macrocycles. Such structures can behave like idealized, infinite π -conjugated oligomers with completely new photophysical characteristics^{24,25}. Significantly, nature exploits such structures to harvest sunlight^{26,27}. A number of photosynthetic organisms have evolved to use a wide variety of macrocycles comprising π -conjugated units to efficiently harvest solar energy and transfer it to reaction centers.^{28,29}

Therefore, there is great interest in understanding the photophysics and excited state dynamics of conjugated nanorings. Excitations in such structures tend to exhibit highly delocalized character²⁵. However, large extended chromophoric structures inevitably present conformational heterogeneity that can act as a perturbation and localize the excitation on a given chromophore or region of the macrocycle³⁰. This is detrimental to the efficient operation envisaged for photonic structures and biomimetic light harvesting applications. An essential source of perturbation is provided by linking groups that affect and modulate the dihedral angle formed between the ring chromophoric units. The dihedral angle minimization (also called

1
2
3 “planarization”) enhances the conjugation length, generally increasing excitation delocalization
4
5 along the porphyrin oligomer structure.
6
7

8
9 The most important physical and chemical properties of π -conjugated chromophores
10 involve their excited electronic states. However, excited states have relatively short lifetimes
11 and therefore it is necessary to apply time resolved spectroscopy to infer detailed information
12 and therefore it is necessary to apply time resolved spectroscopy to infer detailed information
13 about their photophysical and photochemical processes. Furthermore, given the high energy
14 content in visible electronic excitation, structural motions can significantly affect the dynamics
15 in π -conjugated chromophores and a better understanding of how such motions are coupled to
16 electronic transitions is of great interest.
17
18
19
20
21
22
23
24

25
26 In this work we present a detailed study of ultrafast excited state dynamics of π -conjugated
27 six-membered porphyrin nanorings, with (*c*-P6•T6) and without (*c*-P6) a template. The
28 molecular structure of such compounds, together with a parent porphyrin dimer (P2) is shown
29 in Figure 1. The aim is to investigate and characterize structural heterogeneity by monitoring
30 changes in the electronic spectra. Ultrafast excitation delocalization and picosecond-timescale
31 structural dynamics are revealed.
32
33
34
35
36
37
38
39

40 **Experimental and Theoretical Methods**

41
42
43 The time resolved measurements have been carried out on a conventional ultrafast transient
44 absorption apparatus³¹. Briefly, an amplified Ti:Sapphire laser system (Spectra Physics Spitfire
45 ACE), that generates 120 fs pulses centred at 800 nm with a repetition rate of 1 kHz, drives a
46 commercial noncollinear optical parametric amplifier (NOPA, Light Conversion, TOPAS
47 White), producing pulses around 20 fs duration, and also two collinear commercial OPAs (Light
48 Conversion, TOPAS Prime) capable of giving 75 fs pulses tuneable from the UV to the IR. The
49 NOPA pulses, used as pump, were recompressed close to the Fourier transform limit at the
50 sample position and characterized by transient grating frequency resolved optical gating (TG-
51 FROG) using a 1 mm fused silica window as the nonlinear medium. To generate broadband
52
53
54
55
56
57
58
59
60

1
2
3 probe pulses in the vis - near IR region, a 30 mm focal length lens was used to focus around 10
4 μJ , originating from an OPA centred at 1200 nm, onto a 3 mm thick sapphire window, to
5
6 generate a supercontinuum spectrum spanning a spectral region from 550 nm until 1200 nm.
7
8 Reflective optics were used (predominantly) in the experiment to direct the pump and probe
9
10 pulses to the sample position, thus keeping the dispersion as low as possible. The pump and
11
12 probe pulses were focused to spots with FWHM of $158 \pm 7 \mu\text{m}$ and $37 \pm 2 \mu\text{m}$. The probe beam
13
14 had its polarization set at 45° with regards to the pump beam. After the sample, the probe beam
15
16 is split into two with a polarizing beam splitter and each polarization is measured
17
18 simultaneously. The spectrum of both probe polarizations are measured at 1 kHz using two
19
20 back-thinned full frame transfer (FFT) CCD cameras (Entwicklungsbuero Stresing)
21
22 incorporated into a home-built two-channel prism-based spectrometer. An optical chopper was
23
24 used to modulate the pump pulses at 500 Hz allowing transient absorption difference spectra to
25
26 be calculated for consecutive probe pulses. The isotropic response is determined by calculating
27
28 $S_{iso} = 1/3(S_{\parallel} + 2S_{\perp})$ whereas the anisotropy³¹ is given by $S_{aniso} = (S_{\parallel} - S_{\perp})/(S_{\parallel} + 2S_{\perp})$.
29
30
31
32
33
34
35

36 The porphyrin nanorings were synthesised as reported previously^{32,33} and dissolved in
37
38 toluene with 1% v/v pyridine to prevent aggregation. The resulting solutions were placed in 1
39
40 mm optical path static cells for both the steady-state and time-resolved measurements, adjusting
41
42 their concentration in order to obtain OD around 0.3 at their steady-state maxima.
43
44 Measurements performed at lower concentration gave the same results, confirming that no
45
46 aggregation was present.
47
48
49

50 **Results and Discussion**

51
52
53 The porphyrin structures studied here consist of nanorings composed by six meso-
54
55 substituted zinc porphyrin units (aryl groups (3,5-bis(trihexylsilyl)benzene) at meso-positions)
56
57 forming a circular structure through butadiyne links. Their structures are shown in Figure 1.
58
59 The circular structure can be rigidified by the support of an inner hexapyridyl template (shown
60

1
2
3 in blue in Figure 1). Importantly, the butadiyne interconnecting links ensure complete π -
4 conjugation of the nanoring, as evidenced by a large red shift of the lowest singlet absorption
5 band (Q-band) relative to that of the bisalkynylporphyrin monomer³⁴. The visible to near UV
6 absorption spectrum of *c*-P6 and *c*-P6•T6 with their respective band assignments are shown in
7 Figure S1. Similarly to other substituted porphyrin structures³⁴, the lowest energy spectral
8 amplitude is related to transitions from the ground state into the so-called Q-band. Due to the
9 asymmetric substitution present in the nanoring porphyrin units, the degeneracy of the transition
10 dipole moments along the *x* and *y* axes is lifted³⁵.

21
22 Determination of the lowest energy transition of the nanoring structure is complicated.
23 Nanoring systems, including the light-harvesting LH2 system, have a circular symmetry.
24 Therefore, by analogy, the excited state energy-levels can be conveniently described by the
25 Hückel molecular orbital theory for π -conjugated molecules.³⁶ The main outcome of the model
26 is that there will be two nondegenerate $k = 0$ (S_1) and $k = 3$ (S_4) as well as two pairwise
27 degenerate $k = \pm 1$ (S_2) and $k = \pm 2$ (S_3) electronic excited states. Furthermore, Hückel theory
28 also indicates that only transition dipole moments to the two degenerate $k = \pm 1$ (S_2) states of
29 orthogonal polarization (*x* and *y* transitions) carry significant oscillator strength. Additionally,
30 according to the Kasha model, for a circular structure the transition dipole moment
31 contributions will sum to zero, i.e., the lowest $k = 0$ (S_1) has a transition dipole moment strength
32 equal to zero.³² Once the molecule has relaxed into its lowest energy state, emission is expected
33 to be strongly suppressed compared to the emission observed in the porphyrin monomer. This
34 was indeed reported for *c*-P6 and *c*-P6•T6³³, which present a small fluorescence quantum yield
35 (0.43% and 0.12%, respectively) and small radiative rate constants (0.029 ns⁻¹ and 0.0048 ns⁻¹
36 respectively). Thus, the intense absorption spectrum is assigned to the strongly allowed $S_0 \rightarrow$
37 S_2 transition followed by a fast internal conversion to the S_1 state, where emission is suppressed.
38 Given that *c*-P6 is not held in place by the template, it has some flexibility, so cancellation of
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 the transition dipoles is expected to be less complete, explaining a larger transition moment
4
5 than for *c*-P6•T6.²⁵
6
7

8 The dark state nature of the lowest excited singlet state in nanorings, makes the
9
10 identification of the ground to first excited state ($S_0 \rightarrow S_1$) transition difficult. Recently time
11
12 dependent density functional theory (TDDFT) estimated this transition to be at ~ 945 nm.³³ A
13
14 further challenge remains the assignment of the sequence of similarly-spaced strong absorption
15
16 peaks clearly visible in the templated nanoring *c*-P6•T6, at 777, 813, and 858 nm (Figure S1).
17
18 One assignment was made in terms of a Franck–Condon progression of a vibrational mode
19
20 coupled to the electronic transition³³, with a wavenumber around ~ 605 cm^{-1} . An alternative
21
22 interpretation assigns electronic states to the peaks that appear in the absorption spectrum. This
23
24 assignment was based on two-dimensional electronic spectroscopy measurements³⁷, where the
25
26 analysis and classification of coherences of different origins allowed for the identification of
27
28 electronic transitions present in these nanorings. However, an unambiguous and complete
29
30 assignment of the origin of the peaks appearing in the absorption and emission spectra remains
31
32 an elusive question.
33
34
35
36
37

38 A series of time-resolved pump-probe spectra for *c*-P6 and for *c*-P6•T6 are shown in Figures
39
40 2a and 2c (main graphs), together with the linear absorption and pump laser spectra (top graphs,
41
42 shown in red and blue, respectively). For both structures the pump-probe signal consists of a
43
44 strong ground state bleach (GSB) (680 nm – 890 nm, related to $S_0 \rightarrow S_2$ transitions) and a broad
45
46 excited state absorption (ESA) which extends into the IR region as well as to wavelengths
47
48 shorter than 680 nm. Those signals are established within the time resolution of our experiment
49
50 (~ 50 fs) as can be seen in Figures 2b and 2d, an important observation indicating that the $S_2 \rightarrow$
51
52 S_1 internal conversion is very efficient, as was also observed for other cyclic structures.
53
54 Theoretical calculations performed on conjugated cycloparaphenylenes, made up of 14
55
56 monomers, have shown that the internal energy transfer to the lowest singlet excited state
57
58
59
60

1
2
3 happens on a 50 fs timescale³⁸. The results presented on Figure 2b and 2d give evidence that
4 such fast internal conversion also occurs for the porphyrin nanorings. There is no risetime or
5 fast decay time observable for the ESA contribution, which would indicate any S₁ population
6 formation or S₂ relaxation. Therefore, this process must be occurring during pulse overlap
7 around time zero or on a timescale which is too fast to be resolved with our experimental setup.
8
9

10
11
12
13
14
15 As the inset of figures 2c and 2d shows, the ESA signal fully relaxes in a couple of
16 nanoseconds whereas the GSB does not completely recover within the time range of our
17 experiment. This can be assigned to triplet state formation, which relaxes back to the ground
18 state over much longer timescales. The nonradiative rate constant for *c*-P6•T6 ($k_{nr} = 1.6 \text{ ns}^{-1}$) is
19 smaller than for *c*-P6 ($k_{nr} = 2.03 \text{ ns}^{-1}$) and, therefore, the efficiency in accumulating population
20 in the triplet state is reduced, producing a smaller offset in the final bleach values. Apart from
21 a long time decay, there is a fast spectral evolution, mainly in the 820 – 1100 nm spectral region,
22 during the first 500 fs, which is observed in both structures.
23
24
25
26
27
28
29
30
31
32

33
34 To quantify the time evolution of the whole pump-probe signal, we applied a global analysis
35 fitting procedure to the data.³⁹ The resulting decay associated difference spectra (DADS) are
36 shown in Figure 3 (normalized DADS are presented in Figure S6). To access quality of the
37 fitting, time resolved curves at specific wavelengths together with fit are presented in Figure
38 S2. For *c*-P6•T6 at least three components were necessary to completely fit the data whereas
39 for *c*-P6 one additional time component was necessary. The shortest component, with a time
40 constant around 150 fs, accounts for a fast spectral evolution already mentioned (in the region
41 of the 820 – 1100 nm). This contribution can be assigned either to excited state energy
42 equilibration within the vibronic energy levels or downhill vibrational energy relaxation. The
43 much slower contributions are due to ESA decay and GSB recovery (462 ps for *c*-P6 and 241
44 ps for *c*-P6•T6) and triplet state formation (~ ns) that is not resolved in our experiment. The
45 triplet states of *c*-P6 and *c*-P6•T6 have previously been observed with transient electron
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 paramagnetic resonance (EPR) spectroscopy by Tait et al⁴⁰, where they had lifetimes of
4
5 hundreds of microseconds (at 20 K) and were fully delocalized on the EPR timescale.
6
7 Furthermore, the longer relaxation time for *c*-P6 (462 ps) compared to *c*-P6•T6 (241 ps) can be
8
9 explained by the fact that structural inhomogeneity breaks perfect circular symmetry of the *c*-
10
11 P6 rings enabling slight increase in transition dipole moment. This is reflected in the higher
12
13 radiative rate for *c*-P6 (0.029 ns⁻¹) when compared to *c*-P6•T6 (0.0048 ns⁻¹). An extra
14
15 component, with a time constant of 200 ps, is required to describe the spectral evolution of *c*-
16
17 P6. This component can be assigned to the molecular planarization in the excited state, i.e.,
18
19 minimization of the dihedral angles between the porphyrin monomers composing the nanoring
20
21 as reported before.^{41,42,43} Figure 4 compares curves at 900 nm and 910 nm (these wavelengths
22
23 were chosen because the rising signal is maximized at those specific wavelengths),
24
25 corresponding to ESA, for *c*-P6 and *c*-P6•T6, respectively, clearly evidencing the absence of a
26
27 rising ESA signal for *c*-P6•T6. This is so because the *c*-P6•T6 structure is rigidified by the
28
29 support of an inner hexapyridyl template, which keeps the porphyrin rings planarized.
30
31 Furthermore, in the same figure, a comparison is made with a measurement done for the *c*-P6
32
33 parent dimer, P2, for which the planarization in the excited state was already well
34
35 characterized⁴⁴. The curves for *c*-P6 and P2 overlap almost perfectly, providing strong evidence
36
37 for the origin of this contribution to the pump-probe signal.
38
39
40
41
42
43
44

45 Further evidence for the structural inhomogeneity present in the *c*-P6 solution is provided
46
47 in Figure 5. A comparison is made between pump-probe curves obtained when exciting *c*-P6 at
48
49 high energy (pump centred at 720 nm, blue curve) with excitation of *c*-P6 set at a low energy
50
51 (pump centre at 850 nm, red curve). Figure 4, corresponding to an early time ($T = 100$ fs)
52
53 transient spectra, shows that when excitation is set at low energy, the transient absorption shows
54
55 the well resolved peaks as for the rigidified *c*-P6•T6 structure. Whereas for the same early times,
56
57 when excitation is set at higher energies, a much smoother transient spectrum is obtained. The
58
59
60

1
2
3 explanation proposed for this is that *c*-P6 within the solution has a considerable amount of static
4 disorder, caused by the distribution of dihedral angles between the porphyrin units that make
5 up the nanoring. This assignment is confirmed by comparison to an equivalent set of
6 measurements performed on templated *c*-P6•T6, as shown in Figure S3. For *c*-P6•T6, the
7 spectral shape of pump-probe spectra is independent from excitation. Structures which have,
8 on average, larger dihedral angles present higher transition energies whereas structures with
9 smaller dihedral angles^{42,45}, i.e., more planarized, will have smaller transition energies. Thus
10 excitation at lower energies will give transient spectra resembling more *c*-P6•T6, in which the
11 dihedral angles are kept fixed at small values by the hexapyridyl inner template. The transient
12 behaviour is in agreement with what is observed for *c*-P6 and *c*-P6•T6 linear absorption spectra,
13 where the rigidified structure *c*-P6•T6, which has small dihedral angles between its comprising
14 porphyrin units, absorbs at lower energies when compared to the non-templated *c*-P6 nanoring.
15 Therefore, what Figure 5a shows is that the low energy excitation preferentially excites the
16 planarized structures among the distribution. In other words, changing the wavelength of the
17 excitation pulse, selective excitation (hole-burning) of different subsets of nanorings is
18 performed, with a relatively narrow distribution of dihedral angles (more twisted on the blue
19 side and more planar on the red side) among the inhomogeneously broadened distribution of *c*-
20 P6 nanorings in solution.
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44

45 As time evolves, there is a dynamical evolution of the structures and the dihedral angles can
46 change, altering the distribution's absorption spectrum. This is shown in Figure 5b and 5c. As
47 the delay time between pump and probe increases, the peaks start to be "washed out", resulting
48 in transient spectra for different excitations to become very similar. The time constant retrieved
49 for this dihedral angle randomization in the ground state (given that the GSB is being observed)
50 is 55 ps. This time constant is obtained by fitting the residual of the difference between pump-
51 probe curves obtained when exciting *c*-P6 at high energy (pump centred at 720nm) with
52
53
54
55
56
57
58
59
60

1
2
3 excitation of *c*-P6 set at a low energy (pump centre at 850nm) (Figure S4). Therefore, this
4 process is almost 4 times faster than the excited state planarization. Interestingly, the
5 planarization in the excited state of the untemplated ring is almost four times slower than the
6 random fluctuations taking place in its ground state. This is in contrast to what was observed in
7 the parent dimer P2, in which structural dynamics occurs on a slower timescale in the ground
8 state. These differences are related to the nature of the electronic excited states in which
9 electronic dynamics are taking place, with the S_1 excited state of P2 resembling more S_2 of *c*-
10 P6, which is initially populated. Due to the ultrafast $S_2 \rightarrow S_1$ population transfer in *c*-P6,
11 planarization takes place in S_1 which has a different character than S_1 state in P2.
12
13
14
15
16
17
18
19
20
21
22
23

24 Finally, the wavelength and time resolved anisotropy was determined, as shown for *c*-P6 in
25 Figure 6a and for *c*-P6•T6 in Figure S5. The two-dimensional plot shows the anisotropy as a
26 colour scale for a given wavelength and time. The top graph shows the laser spectrum (blue),
27 steady-state absorption spectrum (red) and an isotropic transient absorption curve at $T=200$ fs
28 (green). When the isotropic signal crosses zero the anisotropy becomes undefined (at around
29 675 and 900 nm). The main feature of this graph is that there is no major anisotropy variation
30 across the spectral region studied. The anisotropy value is around 0.1, apart from very early
31 times. This is so for *c*-P6 (Figure 6a) and for *c*-P6•T6 (Figure S5), implying that anisotropy is
32 not affected by structural inhomogeneity. Figure 6b shows the anisotropy time evolution for a
33 wavelength equal to 1050 nm, located in the ESA region. It is possible to resolve a fast (~ 50
34 fs) decay during the first 200 fs, after which the anisotropy value stabilizes at around 0.1. As
35 the inset graph shows, this value persists until the excited state signal has fully relaxed, even
36 for *c*-P6. An anisotropy value of 0.1 is expected for a system which has no preferred excitation
37 polarization direction as well no preferred emission dipole moment. This is exactly the case for
38 transitions in the nanoring plane, where the polarization memory is completely lost in a two-
39 dimensional plane. However, energy transfer among the ring chromophores could also lead to
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 this randomization, it is however unlikely. It would have to be extremely fast given that it has
4
5 to cover a distance of ~ 15 nm (circumference length of a 2.4 nm ring) in 50 fs.
6
7

8 Wynne and Hochstrasser have shown⁴⁶ that, when exciting degenerate states that have their
9
10 transition dipole moment orthogonal, the electronic coherence properties of the two states
11
12 involved need to be considered. In such a case the anisotropy value can be as high as 0.7 but
13
14 quickly decays to a value of 0.1 as the coherence between states is lost. Therefore, the fast 50
15
16 fs decay measured here, which is within the instrument time resolution, indicates a decoherence
17
18 between $k = \pm 1$ states. Importantly, it also indicates that the excitation delocalizes along the
19
20 nanoring within this fast timescale. This result confirms the earlier suggestion³⁰ that both the
21
22 absorbing and the emitting states must be fully delocalized over the nanoring.
23
24
25
26
27
28
29

30 **Conclusions**

31
32
33 The photophysics of two different π -electron porphyrin nanorings was investigated through
34
35 time resolved pump-probe spectroscopy. The non-templated *c*-P6 nanoring shows significant
36
37 structural dynamics. In the excited state, planarization occurs on a time scale of 200 ps whereas
38
39 in the ground state dihedral angle randomization occurs faster, with a timescale of 55 ps. On
40
41 the other hand, the addition of an inner template, rigidifies the nanoring structure. Therefore,
42
43 for *c*-P6•T6 no significant structural motions are observed. In both nanorings, ultrafast excited
44
45 state delocalization occurs along the whole nanoring, as evidenced by anisotropy
46
47 measurements. The delocalization is faster than 50 fs, the instrumental response function used
48
49 in this work. Interestingly, the excitation remains delocalized for the whole excited state
50
51 population time and is not affected by structural motions as present in *c*-P6.
52
53
54
55
56
57
58
59
60

1
2
3 **Acknowledgments**
4
5

6 This work was supported by EPSRC grants EP/J009148/1 and J021431/1.
7
8
9

10
11 **Supporting Information**
12

13 Additional figures including UV-Vis absorption, residual difference curve, *c*-P6•T6 anisotropy
14 measurement.
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure Captions

Figure 1. Porphyrin structures consist of nanorings composed by six meso-substituted zinc porphyrin units (bearing aryl groups (3,5-bis(trihexylsilyl)benzene) at meso-positions) forming a circular structure through butadiyne links. *c*-P6•T6 is the templated ring, *c*-P6 is the same structure without the hexapyridyl template (shown in blue). P2 is the parent dimer.


Figure 2. Transient absorption spectra at several probe delay times for (a) *c*-P6 and (c) *c*-P6•T6. Graphs (b) and (d) show time resolved transient absorption curves taken at selected probing wavelengths of 1083 (red) and 776 nm (blue) for *c*-P6 and 1100 (red) and 809 nm (blue) for *c*-P6•T6.


Figure 3. Decay associated difference spectra (DADS) for (a) *c*-P6 and (b) *c*-P6•T6.


Figure 4. Comparison of time resolved transient absorption curves taken at selected probing wavelengths: 900 nm for *c*-P6 (black), 910 nm for *c*-P6•T6 (red) and 1160 nm for P2 (blue). The right hand side scale is related only to the P2 (blue) curve.


Figure 5. Transient absorption spectra for *c*-P6 at two different excitation conditions, as shown in the top graph, centred at 700 nm (blue) and 830 nm (red). The different graphs correspond to different pump-probe delay times as for (a) T=0.1 ps, (b) T=10 ps and (c) T=120 ps.


Figure 6. (a) Two-dimensional plot showing the anisotropy as a colour scale. The top graph shows the laser spectrum (blue), absorption spectrum (red) and an isotropic transient absorption curve at T=200fs (green). (b) Anisotropy as a function of time taken at a selected probing wavelength of 1051 nm (black) and fit to the data (red).


**Figure 1**

**Figure 2**

**Figure 3**

**Figure 4**

**Figure 5**

**Figure 6**

References

- (1) Hafner, R. J.; Tian, L.; C. Brauer, J.; Schmaltz, T.; Sienkiewicz, A.; Balog, S.; Flauraud, V.; Brugger, J.; Frauenrath, H. Unusually Long-Lived Photocharges in Helical Organic Semiconductor Nanostructures. *ACS Nano* **2018**, *12* (9), 9116–9125.
- (2) Huang, Y.; Xu, Z.; Jin, S.; Li, C.; Warncke, K.; A. Evangelista, F.; Lian, T.; Egap, E. Conjugated Oligomers with Stable Radical Substituents: Synthesis, Single Crystal Structures, Electronic Structure, and Excited State Dynamics. *Chem. Mater.* **2018**, *30* (21), 7840–7851.
- (3) Walzer, K.; Maennig, B.; Pfeiffer, M.; Leo, K. Highly Efficient Organic Devices Based on Electrically Doped Transport Layers. *Chem. Rev.* **2007**, *107* (4), 1233–1271.
- (4) McEwas, K. J.; Bourhill, G.; Robertson, J. M.; Anderson, H. L. The Nonlinear Optical Characterization Of Meso-Substituted Porphyrin Dyes. *J. Nonlinear Opt. Phys. Mater.* **2000**, *09* (04), 451–468.
- (5) B. Teran, N.; S. He, G.; Baev, A.; Shi, Y.; T. Swihart, M.; N. Prasad, P.; J. Marks, T.; R. Reynolds, J. Twisted Thiophene-Based Chromophores with Enhanced Intramolecular Charge Transfer for Cooperative Amplification of Third-Order Optical Nonlinearity. *J. Am. Chem. Soc.* **2016**, *138* (22), 6975–6984.
- (6) W. Becker, M.; S. Sapochak, L.; Ghosen, R.; Xu, C.; R. Dalton, L.; Shi, Y.; H. Steier, W.; K.-Y. Jen, A. Large and Stable Nonlinear Optical Effects Observed for a Polyimide Covalently Incorporating a Nonlinear Optical Chromophore. *Chem. Mater.* **2002**, *6* (2), 104–106.
- (7) Ogawa, K.; Zhang, T.; Yoshihara, K.; Kobuke, Y. Large Third-Order Optical Nonlinearity of Self-Assembled Porphyrin Oligomers. *J. Am. Chem. Soc.* **2001**, *124* (1), 22–23.
- (8) Qian, G.; Dai, B.; Luo, M.; Yu, D.; Zhan, J.; Zhang, Z.; Ma, D.; Yuan Wang, Z. Band Gap Tunable, Donor–Acceptor–Donor Charge-Transfer Heteroquinoid-Based Chromophores: Near Infrared Photoluminescence and Electroluminescence. *Chem. Mater.* **2008**, *20* (19), 6208–6216.
- (9) Gankin, A.; Mervinetsky, E.; Alshanski, I.; Buchwald, J.; Dianat, A.; Gutierrez, R.; Cuniberti, G.; Sfez, R.; Yitzchaik, S. ITO Work Function Tunability by Polarizable Chromophore Monolayers. *Langmuir* **2019**, *35* (8), 2997–3004.
- (10) Chandrabose, S.; Chen, K.; J. Barker, A.; J. Sutton, J.; K. K. Prasad, S.; Zhu, J.; Zhou, J.; C. Gordon, K.; Xie, Z.; Zhan, X.; et al. High Exciton Diffusion Coefficients in Fused Ring Electron Acceptor Films. *J. Am. Chem. Soc.* **2019**, *141* (17), 6922–6929.
- (11) Kay, A.; Graetzel, M. Artificial Photosynthesis. 1. Photosensitization of Titania Solar Cells with Chlorophyll Derivatives and Related Natural Porphyrins. *J. Phys. Chem.* **2002**, *97* (23), 6272–6277.
- (12) S. He, G.; Zhu, J.; Baev, A.; Samoć, M.; L. Frattarelli, D.; Watanabe, N.; Facchetti, A.; Ågren, H.; J. Marks, T.; N. Prasad, P. Twisted π -System Chromophores for All-Optical Switching. *J. Am. Chem. Soc.* **2011**, *133* (17), 6675–6680.
- (13) Ambroise, A.; W. Wagner, R.; Dharma Rao, P.; A. Riggs, J.; Hascoat, P.; R. Diers, J.;

- 1
2
3 Seth, J.; K. Lammi, R.; F. Bocian, D.; Holten, D.; et al. Design and Synthesis of
4 Porphyrin-Based Optoelectronic Gates. *Chem. Mater.* **2001**, *13* (3), 1023–1034.
5
6 (14) Rong, Y.; Wu, C.; Yu, J.; Zhang, X.; Ye, F.; Zeigler, M.; Elena Gallina, M.; Wu, I.-C.;
7 Zhang, Y.; Chan, Y.-H.; et al. Multicolor Fluorescent Semiconducting Polymer Dots
8 with Narrow Emissions and High Brightness. *ACS Nano* **2013**, *7* (1), 376–384.
9
10 (15) Wang, B.; Yu, C. Fluorescence Turn-on Detection of a Protein through the Reduced
11 Aggregation of a Perylene Probe. *Angew. Chemie - Int. Ed.* **2010**, *49* (8), 1485–1488.
12
13 (16) J. M. Hoeben, F.; Jonkheijm, P.; W. Meijer, E.; P. H. J. Schenning, A. About
14 Supramolecular Assemblies of π -Conjugated Systems. *Chem. Rev.* **2005**, *105* (4),
15 1491–1546.
16
17 (17) Bols, P. S.; Anderson, H. L. Template-Directed Synthesis of Molecular Nanorings and
18 Cages. *Acc. Chem. Res.* **2018**, *51* (9), 2083–2092.
19
20 (18) Joong Lee, S.; T. Hupp, J.; T. Nguyen, S. Growth of Narrowly Dispersed Porphyrin
21 Nanowires and Their Hierarchical Assembly into Macroscopic Columns. *J. Am. Chem.*
22 *Soc.* **2008**, *130* (30), 9632–9633.
23
24 (19) Fathalla, M.; Neuberger, A.; Li, S.-C.; Schmehl, R.; Diebold, U.; Jayawickramarajah, J.
25 Straightforward Self-Assembly of Porphyrin Nanowires in Water: Harnessing
26 Adamantane/ β -Cyclodextrin Interactions. *J. Am. Chem. Soc.* **2010**, *132* (29), 9966–
27 9967.
28
29 (20) Bressan, G.; Green, D.; Chan, Y.; Bulman Page, P. C.; Jones, G. A.; Meech, S. R.;
30 Heisler, I. A. One- to Two-Exciton Transitions in Perylene Bisimide Dimer Revealed
31 by Two-Dimensional Electronic Spectroscopy. *J. Phys. Chem. A* **2019**, *123* (8), 1594–
32 1601.
33
34 (21) Liu, X.; Sun, Y.; A. Perez, L.; Wen, W.; F. Toney, M.; J. Heeger, A.; C. Bazan, G.
35 Narrow-Band-Gap Conjugated Chromophores with Extended Molecular Lengths. *J.*
36 *Am. Chem. Soc.* **2012**, *134* (51), 20609–20612.
37
38 (22) Ma, W.; Yang, C.; Gong, X.; Lee, K.; Heeger, A. J. Thermally Stable, Efficient
39 Polymer Solar Cells with Nanoscale Control of the Interpenetrating Network
40 Morphology. *Adv. Funct. Mater.* **2005**, *15* (10), 1617–1622.
41
42 (23) Mena-Osteritz, E. Superstructures of Self-Organizing Thiophenes. *Adv. Mater.* **2002**,
43 *14* (8), 609–616.
44
45 (24) Donehue, J. E.; Varnavski, O. P.; Cemborski, R.; Iyoda, M.; Goodson, T. Probing
46 Coherence in Synthetic Cyclic Light-Harvesting Pigments. *J. Am. Chem. Soc.* **2011**,
47 *133* (13), 4819–4828.
48
49 (25) Yong, C.-K.; Parkinson, P.; Kondratuk, D. V.; Chen, W.-H.; Stannard, A.;
50 Summerfield, A.; Sprafke, J. K.; O’Sullivan, M. C.; Beton, P. H.; Anderson, H. L.; et
51 al. Ultrafast Delocalization of Excitation in Synthetic Light-Harvesting Nanorings.
52 *Chem. Sci.* **2015**, *6* (1), 181–189.
53
54 (26) McDermott, G.; Prince, S. M.; Freer, a. a.; Hawthornthwaite-Lawless, a. M.; Papiz,
55 M. Z.; Cogdell, R. J.; Isaacs, N. W. Crystal Structure of an Integral Membrane Light-
56 Harvesting Complex From Photosynthetic Bacteria. *Nature.* **1995**, *374*, 517–521.
57
58
59
60

- 1
2
3 (27) D. Scholes, G. Quantum-Coherent Electronic Energy Transfer: Did Nature Think of It
4 First? *J. Phys. Chem. Lett.* **2010**, *1* (1), 2–8.
5
6 (28) Wang, L.; Allodi, M. A.; Engel, G. S. Quantum Coherences Reveal Excited-State
7 Dynamics in Biophysical Systems. *Nat. Rev. Chem.* **2019**, *3*, 477–490.
8
9 (29) Kondo, T.; Jia Chen, W.; S. Schlau-Cohen, G. Single-Molecule Fluorescence
10 Spectroscopy of Photosynthetic Systems. *Chem. Rev.* **2017**, *117* (2), 860–898.
11
12 (30) Parkinson, P.; Kondratuk, D. V.; Menelaou, C.; Gong, J. Q.; Anderson, H. L.; Herz, L.
13 M. Chromophores in Molecular Nanorings: When Is a Ring a Ring? *J. Phys. Chem.*
14 *Lett.* **2014**, *5* (24), 4356–4361.
15
16 (31) Berera, R.; van Grondelle, R.; Kennis, J. T. M. Ultrafast Transient Absorption
17 Spectroscopy: Principles and Application to Photosynthetic Systems. *Photosynth. Res.*
18 **2009**, *101* (2–3), 105–118.
19
20 (32) Hoffmann, M.; Kärnbratt, J.; Chang, M. H.; Herz, L. M.; Albinsson, B.; Anderson, H.
21 L. Enhanced π -Conjugation around a Porphyrin[6] Nanoring. *Angew. Chemie - Int. Ed.*
22 **2008**, *47* (27), 4993–4996.
23
24 (33) Sprafke, J. K.; Kondratuk, D. V.; Wykes, M.; Thompson, A. L.; Hoffmann, M.;
25 Drevinskas, R.; Chen, W. H.; Yong, C. K.; Karnbratt, J.; Bullock, J. E.; et al. Belt-
26 Shaped π -Systems: Relating Geometry to Electronic Structure in a Six-Porphyrin
27 Nanoring. *J. Am. Chem. Soc.* **2011**, *133* (43), 17262–17273.
28
29 (34) V. A. Camargo, F.; L. Anderson, H.; R. Meech, S.; A. Heisler, I. Full Characterization
30 of Vibrational Coherence in a Porphyrin Chromophore by Two-Dimensional Electronic
31 Spectroscopy. *J. Phys. Chem. A* **2014**, *119* (1), 95–101.
32
33 (35) Anderson, H. L. Building Molecular Wires from the Colors of Life: Conjugated
34 Porphyrin Oligomers. *Chem. Commun.*, **1999**, *23*, 2323–2330.
35
36 (36) Gong, J. Q.; Favereau, L.; Anderson, H. L.; Herz, L. M. Breaking the Symmetry in
37 Molecular Nanorings. *J. Phys. Chem. Lett.* **2016**, *7* (2), 332–338.
38
39 (37) Butkus, V.; Alster, J.; Bašinskaite, E.; Augulis, R.; Neuhaus, P.; Valkunas, L.;
40 Anderson, H. L.; Abramavicius, D.; Zigmantas, D. Discrimination of Diverse
41 Coherences Allows Identification of Electronic Transitions of a Molecular Nanoring. *J.*
42 *Phys. Chem. Lett.* **2017**, *8* (10), 2344–2349.
43
44 (38) Adamska, L.; Nayyar, I.; Chen, H.; Swan, A. K.; Oldani, N.; Fernandez-Alberti, S.;
45 Golder, M. R.; Jasti, R.; Doorn, S. K.; Tretiak, S. Self-Trapping of Excitons, Violation
46 of Condon Approximation, and Efficient Fluorescence in Conjugated
47 Cycloparaphenylenes. *Nano Lett.* **2014**, *14* (11), 6539–6546.
48
49 (39) Snellenburg, J. J.; Laptinok, S. P.; Seger, R.; Mullen, K. M.; van Stokkum, I. H. M.
50 Glotaran : A Java -Based Graphical User Interface for the R Package TIMP. *J. Stat.*
51 *Softw.* **2012**, *49* (3), 1–10.
52
53 (40) Tait, C. E.; Neuhaus, P.; Peeks, M. D.; Anderson, H. L.; Timmel, C. R. Transient EPR
54 Reveals Triplet State Delocalization in a Series of Cyclic and Linear π -Conjugated
55 Porphyrin Oligomers. *J. Am. Chem. Soc.* **2015**, *137* (25), 8284–8293.
56
57 (41) Chang, M.-H.; Hoffmann, M.; L. Anderson, H.; M. Herz, L. Dynamics of Excited-State
58
59
60

- 1
2
3 Conformational Relaxation and Electronic Delocalization in Conjugated Porphyrin
4 Oligomers. *J. Am. Chem. Soc.* **2008**, *130* (31), 10171–10178.
5
- 6 (42) U. Winters, M.; Kärnbratt, J.; Eng, M.; J. Wilson, C.; L. Anderson, H.; Albinsson, B.
7 Photophysics of a Butadiyne-Linked Porphyrin Dimer: Influence of Conformational
8 Flexibility in the Ground and First Singlet Excited State. *J. Phys. Chem. C* **2007**, *111*
9 (19), 7192–7199.
10
- 11 (43) V. A. Camargo, F.; L. Anderson, H.; R. Meech, S.; A. Heisler, I. Time-Resolved
12 Twisting Dynamics in a Porphyrin Dimer Characterized by Two-Dimensional
13 Electronic Spectroscopy. *J. Phys. Chem. B* **2015**, *119* (46), 14660–14667.
14
- 15 (44) Camargo, F. V. A.; Hall, C. R.; Anderson, H. L.; Meech, S. R.; Heisler, I. A. Time
16 Resolved Structural Dynamics of Butadiyne-Linked Porphyrin Dimers. *Struct. Dyn.*
17 **2016**, *3* (2), 023608.
18
- 19 (45) Peeks, M. D.; Neuhaus, P.; Anderson, H. L. Experimental and Computational
20 Evaluation of the Barrier to Torsional Rotation in a Butadiyne-Linked Porphyrin
21 Dimer. *Phys. Chem. Chem. Phys.* **2016**, *18* (7), 5264–5274.
22
- 23 (46) Wynne, K.; Hochstrasser, R. M. Anisotropy as an Ultrafast Probe of Electronic
24 Coherence in Degenerate Systems Exhibiting Raman Scattering, Fluorescence,
25 Transient Absorption and Chemical Reactions. *J. Raman Spectrosc.* **1995**, *26* (7), 561–
26 569.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

TOC graphic:

