

1 **Flexible polyandry in female flies is an adaptive response to infertile males**

2 Andreas Sutter<sup>1,2,†</sup>, Laura M Travers<sup>1,2</sup>, Keiko Oku<sup>1</sup>, Kynan Delaney<sup>1</sup>, Stefan Store<sup>1</sup>,  
3 Tom A. R. Price<sup>3</sup>, & Nina Wedell<sup>1</sup>

4 1. Centre for Ecology & Conservation, College of Life and Environmental Sciences,  
5 University of Exeter, Cornwall Campus, TR10 9FE, Penryn, UK.

6 2. School of Biological Sciences, Norwich Research Park, University of East Anglia,  
7 Norwich NR4 7TJ, UK

8 3. Institute of Integrative Biology, University of Liverpool, Crown Street, Liverpool, L69  
9 7ZB, UK

10 † [a.sutter@uea.ac.uk](mailto:a.sutter@uea.ac.uk)

11 **Abbreviated title**

12 Male infertility increases polyandry

13 **Funding**

14 This work was supported by a Swiss National Science Foundation fellowship awarded  
15 to AS (grant number P300PA\_177906) a Marie Skłodowska-Curie Fellowship to KO  
16 (grant number 746169 – IMMUNFUNC).

17 **Acknowledgements**

18 We thank Ali Skeats and Michelle Taylor for establishing the isofemale lines, Steven  
19 Parratt and Katherine Roberts for help with heat-exposure protocols, and Zeynep  
20 Karahan for help with wing measurements.

21 **Data Accessibility Statement**

22 Analyses reported in this article can be reproduced using the data provided by Sutter *et*  
23 *al.*, (2019a).

24 **Abstract**

25 Infertility is common in nature despite its obvious cost to individual fitness. Rising  
26 global temperatures are predicted to decrease fertility, and male sterility is frequently  
27 used in attempts to regulate pest or disease vector populations. When males are infertile,  
28 females may mate with multiple males to ensure fertilisation, and changes in female  
29 mating behaviour in turn could intensify selection on male fertility. Fertility assurance is  
30 a potentially wide-spread explanation for polyandry, but whether and how it actually  
31 contributes to the evolution of polyandry is not clear. Moreover, whether a drop in male  
32 fertility would lead to a genetic increase in polyandry depends on whether females  
33 respond genetically or through behavioural plasticity to male infertility. Here, we  
34 experimentally manipulate male fertility through heat-exposure in *Drosophila*  
35 *pseudoobscura*, and test female discrimination against infertile males before and after  
36 mating. Using isogenic lines, we compare the roles of behaviourally plastic versus  
37 genetically fixed polyandry. We find that heat-exposed males are less active and  
38 attractive, and that females are more likely to remate after mating with these males.  
39 Remating rate increases with reduced reproductive output, indicating that females use  
40 current sperm storage threshold to make dynamic remating decisions. After remating  
41 with fertile males, females restore normal fecundity levels. Our results suggest that male  
42 infertility could explain the evolution of adaptively flexible polyandry, but is less likely  
43 to cause an increase in genetic polyandry.

44 **Keywords:** sexual selection, male sterility, multiple mating, phenotypic plasticity,  
45 temperature, sterile insect technique

## 46 **Introduction**

47 Mating failure, defined as adult females remaining unmated (Rhainds, 2010) or as the  
48 failure to convert matings into reproductive success (Greenway *et al.*, 2015), is  
49 pervasive in nature (Garcia-Gonzalez, 2004; Rhainds, 2010). In insects, as many as two-  
50 thirds of all matings do not result in any offspring production, and the median for  
51 mating failure across 30 species is 22% (Garcia-Gonzalez, 2004). Fertilisation failure  
52 can also be considerable in birds (Adkins-Regan, 2015; Schmoll *et al.*, 2016) and  
53 reptiles (Olsson & Shine, 1997), though estimates from wild populations remain rare.  
54 Male infertility may often be responsible for mating failure. Male fertility is often  
55 impaired at high temperatures (David *et al.*, 2005; Setchell, 2006; Hurley *et al.*, 2018;  
56 Sales *et al.*, 2018; but see Janowitz & Fischer, 2011), and increased occurrence of heat  
57 waves due to climate change (Meehl, 2004) may cause higher sterility rates (Reinhardt  
58 *et al.*, 2015; Walsh *et al.*, 2019). Further, selfish genetic elements such as meiotic  
59 drivers favourably target male gametes (Taylor & Ingvarsson, 2003; Price & Wedell,  
60 2008), and mito-nuclear incompatibilities can devastate sperm function (Dowling *et al.*,  
61 2015), meaning intra-genomic conflict is another potentially common source for a  
62 reduction in male fertility. Finally, mass-sterilisation of males is a common strategy for  
63 human pest control (Knipling, 1955; Dyck *et al.*, 2005).

64 Given the wide variety of factors that can create complete or partial infertility in males,  
65 how should females respond? Females show adaptations that help minimise failure to  
66 copulate and become inseminated (Rhainds, 2010). But copulating and/or receiving an

67 ejaculate alone will not guarantee a female successful reproduction if some males are  
68 infertile. In contrast, actively choosing fertile over infertile males could allow females to  
69 secure some reproductive output. The phenotype-linked fertility hypothesis posits that  
70 male signals and fertility are positively correlated, allowing females to simply choose  
71 attractive males to avoid reduced fertility (Sheldon, 1994). While some studies have  
72 found positive correlations between male attractiveness indicators and semen quality  
73 parameters (Malo *et al.*, 2005; Forstmeier *et al.*, 2017), a recent meta-analysis found no  
74 general support for a link between male secondary sexual signals and tentative indices  
75 of ejaculate quality (Mautz *et al.*, 2013). Even when intrinsic male fertility correlates  
76 with male attractiveness, more attractive males may become sperm depleted because of  
77 their increased mating success, making intrinsically more fertile males temporarily less  
78 fertile (Preston *et al.*, 2001), and thus undermining the fertility benefit of female choice  
79 for attractive males. The paucity of evidence for an association between male external  
80 phenotype and fertility may explain why discrimination against sub-fertile or infertile  
81 males is rare. For example, despite mating failure being attributable to individual seed  
82 bug males (Greenway & Shuker, 2015), females do not choose fertile males (Greenway  
83 *et al.*, 2017).

84 When females do not discriminate between fertile and sterile males before mating,  
85 females may safeguard against mating failure simply by mating with multiple males,  
86 thus making multiple mating (polyandry) an alternative to precopulatory choice (e.g.  
87 Sakaluk & Cade, 1980; Gibson & Jewell, 1982; Sheldon, 1994; Arnqvist & Nilsson,  
88 2000; Mossinson & Yuval, 2003; Forbes, 2014). Importantly, polyandrous females can  
89 benefit even without being able to detect fertile males, as long as infertile males' sperm  
90 are outcompeted by fertile males' sperm, or females remate more after mating with

91 sterile males (Lorch & Chao, 2003; Barclay, 2005; Champion de Crespigny *et al.*, 2008;  
92 Hasson & Stone, 2009). Hence, increased fertility assurance for females might be a  
93 major reason why polyandry is so common. Across animal taxa, 89% of all natural  
94 populations investigated showed evidence for multiple paternity (Taylor *et al.*, 2014).  
95 The theory underlying the evolution of polyandry for fertility assurance is well  
96 developed (Hasson & Stone, 2009), and correlative studies support the notion that  
97 females remate more after receiving small or infertile ejaculates (Wetton & Parkin,  
98 1991; Delisle & Hardy, 1997; Torres-Vila *et al.*, 1997; Krokene *et al.*, 1998; Uller &  
99 Olsson, 2005). Support through experimentally impaired male fertility, often in the  
100 context of the sterile insect technique (SIT), comes from many (Miyatake *et al.*, 1999;  
101 Kraaijeveld & Chapman, 2004; Gavriel *et al.*, 2009; Friesen *et al.*, 2014; Landeta-  
102 Escamilla *et al.*, 2016) but not all studies (Harmer *et al.*, 2006; Abraham *et al.*, 2013;  
103 Haq *et al.*, 2013; Krüger *et al.*, 2019).

104 One common limitation is that researchers have typically measured the mean response  
105 of target females (Calkins & Parker, 2005 and references above). While this assesses the  
106 present potential for population control through the release of sterile males, it largely  
107 ignores the possibility of a dynamic female response that evolves over multiple  
108 generations. Indeed, field studies of releases of sterile males into natural populations  
109 have observed the evolution of precopulatory behavioural discrimination against sterile  
110 males (Hibino & Iwahashi, 1991; Mcinnis *et al.*, 1996), which demonstrates the  
111 importance of considering genetic variation in female mating behaviours when aiming  
112 to predict evolutionary responses. Similarly, male infertility could lead to an  
113 evolutionary increase in polyandry. Selection could favour either genes controlling a  
114 behaviourally plastic increase in female remating after mating with infertile males, or

115 genes underlying generally polyandrous behaviour without behavioural plasticity. If  
116 male sterility in natural populations is consistently high, these two scenarios have the  
117 same outcome. However, if male fertility is compromised only over a temporally  
118 limited period (for example, after a heatwave), a genetic response would lead to a  
119 persisting increase in polyandry in the population, whereas behavioural plasticity would  
120 only increase polyandry during the period of increased male sterility. To our knowledge  
121 only one empirical study has explicitly addressed the evolution of female remating  
122 behaviour in response to sterilised males, and did not find evidence neither for increased  
123 behavioural plasticity nor increased genetically fixed polyandry after 12 generations of  
124 experimental evolution in Tephritid fruit flies (Kuriwada *et al.*, 2014). However, the  
125 authors concluded that insufficient genetic variation in the starting population may have  
126 limited the potential for an evolutionary response (Kuriwada *et al.*, 2014).

127 Here, we investigated whether females of the fly *Drosophila pseudoobscura* mate  
128 multiply to ensure successful fertilisation. Experimentally manipulating the fertility of a  
129 female's first mate through heat-exposure, we measured female reproductive output in  
130 the first four days following the mating, and assessed whether females are more likely  
131 to remate after an infertile/sub-fertile mating. We also assessed male attractiveness and  
132 courtship vigour in an attempt to infer what cues from first mates females may use to  
133 make remating decisions. Importantly, using isolines that genetically differ in polyandry  
134 enabled us to examine the relative roles of behavioural plasticity and genetic  
135 predisposition in shaping the remating response, and hence the evolutionary potential  
136 for polyandry to evolve in response to male infertility.

## 137 **Material and Methods**

138 Fly stocks

139 We used *D. pseudoobscura* that were collected from two populations in the Western  
140 USA (Lewistown, Montana, 47°03'N, 109°28'W; Show Low, Arizona, 34°16'N,  
141 110°00'W) in 2008 and 2012. We maintained all flies under a 14:10 light: dark cycle at  
142 23°C, with standard *Drosophila* food vials (75 mm in height by 25 mm in width)  
143 containing commercial Jazz-Mix™ *Drosophila* food (Fisher Scientific) for feeding and  
144 oviposition. The experiments described here were performed between March and May  
145 2018 across two experimental blocks that were shifted by three days.

146 To explicitly address the roles of behavioural plasticity and genetic variation in female  
147 remating behaviour, we sourced females from ten isofemale isogenic lines that differ in  
148 polyandry and that had been established using wild-caught females as described in  
149 detail elsewhere (Taylor *et al*., 2016; Sutter *et al*., 2019b). Briefly, offspring of wild-  
150 caught females were full-sib inbred for 15 or more generations, after which flies within  
151 an isoline are virtually genetically identical, and after which these isolines were  
152 maintained under less-restrictive breeding conditions. Before the experiment, isolines  
153 were subjected to one generation of common garden breeding. We set up five vials per  
154 isoline with five virgin females and five males each, which gave females opportunity for  
155 mate choice. After 24 hours, before *D. pseudoobscura* females remate (Snook & So,  
156 2000), males were removed and females were transferred to a new vial to oviposit.  
157 Female groups were then transferred to new food every 24 hours for 7 consecutive days.  
158 We used the daughters of these females in experimental mating assays.

159 Males were derived from the same populations as the isofemale lines, but were  
160 maintained across several standard *Drosophila* vials as small outbred laboratory

161 populations with overlapping generations and fluctuating population sizes. Flies  
162 collected in 2008 were kept separately from flies collected in 2012, such that we  
163 maintained four laboratory populations, two from both localities. Before the start of the  
164 experiment, we mass-bred these small populations into large 3.5L population cages.  
165 Focal males were collected from standard vials that had been left for oviposition in the  
166 population cages for up to 24h.

#### 167 Male heat-exposure treatment

168 To reduce male fertility, we exposed males to an increased temperature for a few days.  
169 Heat-exposure was achieved by submerging standard vials with groups of ten males into  
170 a water bath that was maintained at either the elevated temperature of 31°C or at the  
171 control temperature of 23°C. About 90% of the vial volume was submerged under  
172 water, such that gas exchange through a foam plug at the top of the vial was still  
173 possible, but the bottom of the foam plug forced all flies to remain below the level of  
174 the water surface. For logistic reasons, water baths were kept on a lab bench and thus  
175 exposed to a natural diurnal light cycle. All males used had been collected within 18h of  
176 eclosion and separated into single sex groups of up to 20 males. To obtain the large  
177 number of virgin males needed for our mating assays we had to pool males collected  
178 over several days. Thus, we heat-exposed two separate cohorts of males for each  
179 experimental block. The first cohort of males (cohort A) had been kept in standard  
180 conditions for 1–2 days, before they were exposed to 31°C for 72h, and finally  
181 separated into individual vials and left at 23°C on the evening before the day of their  
182 mating trial (i.e. around 15h before the mating trial). Because they were collected only  
183 three days prior to the mating assays, the second cohort of males (cohort B) was  
184 subjected to heat-exposure immediately after collection on the day of eclosion for about

185 62h until two hours before their mating trial. Thus, male cohort A was older (5–6d  
186 versus 3d), exposed to heat for longer (72h versus 62h), and given more time to recover  
187 from heat exposure (15h versus 2h) than cohort B. To obtain a measure of how  
188 physiologically stressful our heat-exposure was to males, we measured male survival  
189 during heat-exposure. To do this we counted the number of alive and dead males when  
190 separating them into their individual vials at the end of their heat-exposure treatment.  
191 Further, we checked whether mortality during heat-exposure led to a bias in male size,  
192 i.e. favouring smaller or larger males in the heat-exposure versus the control treatment,  
193 because a male size bias could in turn have affected female (re)mating patterns. As a  
194 proxy for male size we measured the length of the third longitudinal vein (Taylor *et al.*,  
195 2008) of one wing using *Fiji* (Schindelin *et al.*, 2012).

#### 196 Mating assays

197 To avoid fertilisation failure, females might discriminate against sterile males before or  
198 after mating by refusing to mate with sterile males or by increasing remating after  
199 having mated with sterile males, respectively. Alternatively, males may provide females  
200 with cues about their fertility during mating, and females may use these to make future  
201 remating decisions. We used a mating assay routinely performed in our laboratory  
202 (Price *et al.*, 2011; Herrera *et al.*, 2014; Taylor *et al.*, 2016) to address whether heat-  
203 exposed males were less likely or slower to mate, indicating reduced male vigour or  
204 attractiveness. We also determined whether heat-exposed males copulated for a shorter  
205 duration, possibly indicating reduced ejaculate transfer (Price *et al.* 2008), and whether  
206 these behaviours predicted female remating behaviour, potentially informing about  
207 proximate mechanisms underlying polyandry. We used females from each of ten

208 isolines and males from the two populations, the temperature treatments and male  
209 cohorts in a fully-factorial design.

210 We aspirated sexually mature, virgin females that were five or six days old individually  
211 into vials into which a single male had been aspirated the previous day or earlier that  
212 morning, depending on its cohort (see above). We took note of the time when the  
213 female was introduced, and two observers scan-sampled for initiation and termination  
214 of mating to record copulation latency, the duration from female introduction to the first  
215 observed stable mount (i.e. the pair being relatively immobile), indicating successful  
216 copulation, and copulation duration, the time from that first stable mount until the pair  
217 separated. Scan-sampling meant that flies were not continuously observed, but checked  
218 for copulation in short (~2min) intervals. In the second experimental block we  
219 additionally recorded *ad libitum* observations of the onset of male courtship to obtain  
220 data on latency to initiate courtship and time between courtship initiation and mating.  
221 Observers were always blind with regards to male heat-exposure treatment and female  
222 isoline identity. We used a combination of randomisation and stratification to determine  
223 order in the assay to avoid time-of-day effects on mating parameters. After giving pairs  
224 a minimum of two hours to mate, we removed males and froze them for later size  
225 measurements. We left females that had mated to oviposit for four days, and discarded  
226 females that had not mated.

227 We gave females a single opportunity to remate four days after their first mating. Again,  
228 we aspirated a female into a vial containing a single 5-day old virgin male from the  
229 same population as the female's first mate. These males had been kept in incubators at  
230 the control temperature of 23°C. Two observers regularly scanned pairs for mating.

231 After allowing a minimum of 90min for remating, we discarded all males. To examine  
232 the consequences of enforced monandry on female fitness, we denied a subset (~15%)  
233 of females the opportunity to remate by aspirating the male out of the vial immediately  
234 before the female was introduced. We left females to oviposit for another four days,  
235 after which they were transferred to a third vial for a further four days and finally  
236 discarded.

### 237 Fitness consequences

238 To assess the consequences of male heat-exposure and female remating for female  
239 fitness, we quantified female reproductive output over 12 days, which has been shown  
240 previously to correlate with lifetime reproductive success under control conditions  
241 (Avent *et al.*, 2008). We counted the number of eclosed offspring from these vials 23  
242 days after the first day of oviposition.

243 To obtain additional data on male fertility and mating capacity, we left a single male in  
244 a vial with five virgin females for 24h, after which females were isolated and left to  
245 oviposit for four days, following offspring counts after 23 days. For this small  
246 experiment, we only used males from one of the populations (Show Low) from cohort B  
247 in the first and from cohort A in the second experimental block, and used a haphazard  
248 selection of virgin females from the ten isolines.

### 249 Statistical analyses

250 To test the physiological impact of heat-exposure on males and its consequences for  
251 females we analysed the impact of heat-exposure on multiple aspects of male and  
252 female reproductive behaviour and fitness: i) male heat-exposure survival, ii) mating

253 success, copulation latency and duration, as well as iii) female reproductive output and  
254 iv) polyandry. We used R version 3.5.1 (R Core Team, 2018) for all statistical analyses  
255 and figures, running binomial generalised linear mixed effects models (GLMM) and  
256 linear mixed effects models (LMM) implemented in *lme4* version 1.1-14 (Bates *et al.*,  
257 2015), and zero-inflated mixed models in *glmmTMB* (Brooks *et al.*, 2017). Descriptive  
258 statistics and sample sizes for the different response variables are summarised in  
259 Table 1. Here we give an overview of the fixed and random predictor variables included  
260 in the different models (see also Tables 2, 3 & S1–S4).

261 i) We first measured male survival to assess how physiologically stressful our heat-  
262 exposure treatment was: We ran a binomial GLMM with heat-exposure, male  
263 cohort, their interaction and block as fixed effects, and post-eclosion housing vial  
264 and population as random intercepts. To ask whether survival was biased with  
265 respect to male size, we ran an LMM on the wing size of surviving males, with  
266 heat-exposure, male cohort and their interaction as fixed effects, and male  
267 collection batch (16 unique block, population, and collection day combinations) as  
268 a random effect.

269 ii) We measured male mating success, copulation latency and duration to test for  
270 effects of heat-exposure on male reproductive performance: We ran a binomial  
271 GLMM for mating success as well as LMMs on log-transformed copulation  
272 latency and duration with heat-exposure, male cohort, their interaction, block,  
273 female age, male size and temporal order within the mating assay as fixed effects,  
274 and random intercepts for female post-eclosion housing vial, female isoline and  
275 male population.

276 iii) We then tested the consequences of mating with a heat-exposed male with or  
277 without successive remating with control males for female reproductive output:  
278 Because many of the oviposition vials contained no offspring, we used zero-  
279 inflated models with a Gaussian distribution for the conditional part implemented  
280 in *glmmTMB* (Brooks *et al.*, 2017), and examined residuals with *DHARMA*  
281 (Hartig, 2018). Our conditional full model included heat-exposure, female  
282 remating, male size, laying vial and two- and three-way interactions as fixed  
283 effects. We included random intercepts for female ID, female isoline, male  
284 collection batch (see above), and random slopes for individual females to account  
285 for repeated measures across a female's three laying vials. (Schielzeth &  
286 Forstmeier, 2009). Our zero-inflated full model included heat-exposure, female  
287 remating, male cohort, laying vial and two- and three-way interactions as fixed  
288 effects.

289 iv) Finally, we asked what explained variation in polyandry: We ran a binomial  
290 GLMM with fixed effects for heat-exposure, reproductive output from the first  
291 oviposition vial and male size including two-way interactions with heat-exposure,  
292 and female age and temporal order within the mating assay. We included random  
293 intercepts for female isoline and male collection batch as random intercepts.  
294 Because of our explicit interest in distinguishing between behavioural plasticity  
295 and genetic polyandry, we additionally included the interaction between first male  
296 temperature treatment and female isoline as an additional random effect (i.e.  
297 random slopes for isolines).

298 Whenever possible, we extracted effect sizes and p values from full models to avoid  
299 biasing effect sizes through the removal of non-significant terms (Forstmeier &

300 Schielzeth, 2011). P values for fixed effects from LMMs were obtained from *t*-tests  
301 using the Kenward-Roger approximation for denominator degrees of freedom  
302 implemented in *lmerTest* (Kuznetsova *et al.*, 2016). For reproductive output, we ran a  
303 large albeit not exhaustive selection of combinations of full and reduced conditional and  
304 zero-inflation models, and selected the best model based on the lowest AIC value. To  
305 facilitate the interpretation of main effects in the presence of interactions and to aid  
306 model convergence, we centred covariates to a mean of zero. Age covariates were  
307 mean-centred, and temporal order within an assay (pairs that were set-up earlier had  
308 more time available for mating/remating) was centred and scaled to a standard deviation  
309 of one. For models on mating behaviour, we additionally centred contrasts between two  
310 factors (male cohorts A and B, first and second experimental blocks) by coding factor  
311 levels as minus 0.5 and 0.5, respectively (Schielzeth, 2010). Approximate 95%  
312 confidence intervals (*CI*) for effect sizes were taken as twice the standard error either  
313 side of the mean (Crawley, 2007). We tested significance of random effects using  
314 likelihood ratio tests between models including and excluding the variable of interest  
315 (Bolker *et al.*, 2009). Additionally, we estimated among-isoline variances and the  
316 covariance between polyandry after mating with control and heat-exposed males using a  
317 Bayesian approach implemented in *MCMCglmm* (Hadfield, 2010; see the  
318 supplementary material).

## 319 **Results**

### 320 Heat-exposure reduces male survival and mating success

321 Heat exposure decreased male survival substantially in male cohort A, but only  
322 marginally in cohort B (Table 1; Fig S3). Survival was lower than 50% in cohort A

323 heat-exposed males but higher than 97% in the three other treatment-cohort  
324 combinations, manifested as a highly significant interaction between treatment and male  
325 cohort (GLMM, N = 1515, effect size  $\beta$  [95%CI] on logit scale = -3.8 [-5.9;-1.7],  $z = -$   
326 3.58,  $p < 0.001$ ; Table S2). There was no indication that heat-exposure caused size-  
327 dependent mortality, as the interaction between temperature and male cohort did not  
328 significantly explain variation in body size of surviving males (i.e. wing length; LMM,  
329 N = 925,  $\beta = -0.01$  [-0.03;0.02],  $t_{1,907.6} = -0.58$ ,  $p = 0.565$ ; Table S3). Substantial  
330 variation in body size was explained by pre-eclosion conditions (unique combinations  
331 of populations, male cohorts and experimental blocks; likelihood ratio test LRT,  
332  $\chi^2(14) = 2.6$ ,  $p < 0.001$ ) but not by post-eclosion treatment (heat-exposure;  $p > 0.5$ ).

333 Males that had been heat-exposed were much less likely to mate (binomial GLMM,  
334 N = 916,  $\beta = -3.1$  [-3.6;-2.7],  $z = -14.2$ ,  $p < 0.001$ ; Table S2). Mating success was 86%  
335 in control males but only 30% in heat-exposed males (Fig 1, Table 1). In conjunction  
336 with a decrease in mating success, copulation latency of successful males was longer for  
337 heat-exposed males (log-transformed latency in minutes; LMM, N = 496,  $\beta = 1.1$   
338 [0.8;1.3],  $t_{1,459.2} = 9.4$ ,  $p < 0.001$ ; Fig 1 & S1; Table S1). Copulations with heat-exposed  
339 males were shorter than those with control males (LMM, N = 487,  $\beta = -0.25$  [-0.35;-  
340 0.15],  $t_{1,451.1} = 9.4$ ,  $p < 0.001$ ; Table 1 & S1; Fig S1). Additional data on male courtship  
341 collected only in the second experimental block indicated that heat-exposed males were  
342 slower and less likely to initiate courtship, and that their courtship quality or intensity  
343 may have been inferior to that of control males (see supplementary Results, Table S1 &  
344 Fig S1).

345 Male heat-exposure reduces female reproductive fitness

346 Females mated to heat-exposed males had lower reproductive fitness than females  
347 mated to control males. This was true both for the likelihood of failing to produce any  
348 offspring over four days after mating as well as for the number of offspring produced  
349 among the subset of females that did produce offspring (Fig 2). In our main dataset, this  
350 was evidenced by a significant baseline effect of male heat exposure treatment on the  
351 zero-inflation model ( $N = 498$ ,  $\beta = 6.5$ ,  $z = 8.4$ ,  $p < 0.001$ ) as well as the conditional  
352 model ( $\beta = -28.9$  [ $-44.6$ ;  $-13.2$ ],  $z = -3.9$ ,  $p < 0.001$ ; Table 2). In our additional, small  
353 dataset, where we housed males with five females for 24h, heat-exposed males  
354 successfully reproduced with fewer females (binomial GLM,  $\beta = -2.9$ ,  $z = -6.8$ ,  
355  $p < 0.001$ ), and sired marginally fewer offspring per fertile mating (LM,  $\beta = -13.0$  [ $-$   
356  $25.5$ ;  $0.5$ ],  $t_{1,29} = 4.3$ ,  $p = 0.046$ ; Table 1).

### 357 Polyandry restores female reproductive fitness in the face of male infertility

358 Polyandry had a beneficial effect on reproductive fitness of females previously mated to  
359 heat-exposed males (Table 3), mainly through reducing the incidence of complete  
360 reproductive failure (Fig 2 & Table 2). In contrast, polyandry had no substantial effect  
361 on fecundity under control conditions (Fig 2), consistent with a recent study (Sutter *et*  
362 *al*), 2019b). Females with higher initial reproductive output were less likely to remate  
363 (chosen monandry; see below), but appeared to run out of sperm over the next 4–8 days  
364 (Fig 2). The temporal decline in reproductive fitness of facultatively monandrous  
365 females and the reproductive increase in polyandrous females within the male heat-  
366 exposure treatment contrasted with the consistent temporal patterns within the control  
367 treatment. This explained the three-way interaction between heat-exposure treatment,  
368 remating phenotype and oviposition vial.

369 Phenotypically plastic polyandry

370 Four days after their first mating, females that had mated with a heat-exposed male were  
371 twice as likely to remate (84%) as were females that had mated with control males  
372 (42%; Table 1). The relationship between polyandry and reproductive output after the  
373 first mating suggests the difference in mating behaviour is causally related to reduced  
374 fertility and fecundity. Females were more likely to remate if they had produced fewer  
375 offspring after the first mating (binomial GLMM,  $N = 427$ ,  $\beta = -0.4$  [-0.7;-0.1],  $z = -$   
376  $2.5$ ,  $p = 0.012$ ; Table 3). However, when matched for fecundity, females mated to heat-  
377 exposed males still had a higher remating likelihood ( $\beta = 1.9$  [1.1;2.7],  $z = 4.7$ ,  
378  $p < 0.001$ ; Table 3). Polyandry tended to decrease after mating with larger males and to  
379 increase with female age (Table 3).

380 The increase in polyandry after mating with heat-exposed males was consistent in  
381 females from all ten isolines, indicated by the interaction between female isoline and  
382 heat-exposure of the first mate not explaining a significant amount of variation in  
383 polyandry (Fig 3; LRT,  $\chi^2(2) = 0.85$   $p = 0.654$ ). In contrast, significant variation  
384 between isolines confirmed genetic variation in polyandry (LRT,  $\chi^2(1) = 10.4$   
385  $p = 0.001$ ). However, our additional analyses using MCMCglmm (Hadfield, 2010)  
386 showed this genetic variation was substantial in control females but negligible in  
387 females mated to heat-exposed males (supplementary Results). Moreover, there was no  
388 clear correlation between polyandry of isolines after mating with control versus heat-  
389 exposed males. In combination, this meant we were unable to confidently reject that  
390 there is genetic variation in behavioural plasticity, nor could we confidently conclude  
391 that the response of isolines was quantitatively consistent. Our results indicate

392 behavioural plasticity in polyandry, and genetic variation in polyandry, but show no  
393 clear evidence for genetic variation in behavioural plasticity.

## 394 **Discussion**

395 Here we show that females representing distinct genotypes consistently use polyandry  
396 as a behaviourally flexible strategy to mitigate the potential fitness loss arising from  
397 male sterility, using cues from stored ejaculates. We found no clear evidence for genetic  
398 variation in how females respond to male infertility, but the flexible female response we  
399 describe here could intensify selection on male fertility, and aid population resilience.

### 400 Adaptively flexible polyandry

401 After mating with heat-exposed males with severely compromised fertility, female  
402 remating doubled from 42% to 84%. Safeguarding against male infertility is a potential  
403 adaptive explanation for the ubiquity of female multiple mating, and a number of  
404 studies have reported increased polyandry after mating with experimentally sterilised  
405 males (e.g. medfly: Miyatake *et al.*, 1999; Kraaijeveld & Chapman, 2004; Gavriel *et al.*,  
406 2009; red garter snake: Friesen *et al.*, 2014; *Anastrepha serpentina*: Landeta-Escamilla  
407 *et al.*, 2016), further supported by correlational data (Sakaluk & Cade, 1980; Wetton &  
408 Parkin, 1991; Uller & Olsson, 2005; Reding, 2015; but see Morrow *et al.*, 2002). Other  
409 experiments however found no effect of male sterility on female remating behaviour  
410 (Queensland fruit fly: Harmer *et al.*, 2006; *Anastrepha fraterculus*: Abraham *et al.*,  
411 2013; melon fly: Haq *et al.*, 2013; *Drosophila suzukii*: Krüger *et al.*, 2018). A potential  
412 explanation for this discrepancy is that the latter studies used artificial techniques such  
413 as genetic manipulation and irradiation to induce male sterility, and these males may

414 lack the cues present in naturally sterile males, with which female remating behaviour  
415 has coevolved.

416 Heat-induced male infertility is likely to be relevant in nature (Sales *et al.*, 2018; Walsh  
417 *et al.*, 2019), and should create a strong incentive for female multiple mating. Here,  
418 more than half of the females that mated with heat-exposed males produced no  
419 offspring following mating, indicating high rates of male sterility, compared to a mere  
420 five percent in the control group. Among these females with failed early reproduction,  
421 remating rates were as high as the proportion of virgin females that mated with control  
422 males, meaning the effect of heat-exposure on polyandry could have been driven by  
423 pseudopolyandry rather than true polyandry (Fisher *et al.*, 2013). However, when  
424 focusing on the subsets of females that had non-zero early reproductive output, the  
425 difference in polyandry between females mated to heat-exposed versus control males  
426 was again almost two-fold (76% [N = 66] versus 39% [N = 274]). More formally, in our  
427 analysis on polyandry where we included early reproductive output as a predictor  
428 variable, male heat-exposure showed a very strong effect on polyandry (Table 3, see  
429 also Fig S2).

430 Females may have used information obtained during the first mating to make remating  
431 decisions. Heat-exposure decreased survival only in male cohort A, but had pronounced  
432 sub-lethal effects on sexual behaviour that were similar in both male cohorts. Heat-  
433 exposed males were slower to initiate courtship, took longer to be accepted by females  
434 and copulated for a shorter duration, possibly because heat-exposure had negative  
435 effects on male condition, thus providing females with additional pre- and peri-  
436 copulatory cues about male fertility. However, remating likelihood was not related to

437 copulation latency or duration of a female's first mating (Table S6), making it more  
438 likely that females used cues from stored ejaculates. Our experimental design did not  
439 distinguish between whether changes in sperm or seminal fluids were responsible for  
440 the increase in polyandry. Either mechanism is plausible, but the effects are likely to be  
441 species-specific. For example, sperm-less males can induce a refractory period in  
442 female Queensland fruit flies and Medflies (Harmer *et al.*, 2006; Gabrieli *et al.*, 2016),  
443 but both seminal fluids and sperm are required for inhibiting remating in *Anastrepha*  
444 *fraterculus* and *A. ludens* (Abraham *et al.*, 2016), and *Drosophila melanogaster* flies  
445 (Liu & Kubli, 2003). Independent of the precise mechanism, our results suggest that  
446 polyandry is not simply a response to the absence of fertile sperm but that females take  
447 current semen storage into account when making remating decisions (Manning, 1967;  
448 Crudgington *et al.*, 2005).

449 Behavioural plasticity appeared to be more important than genetic variation in  
450 polyandry. Polyandry increases with latitude across *D. pseudoobscura* populations in  
451 North America, consistent with the proximate effect of lower temperature increasing  
452 polyandry (Taylor *et al.*, 2016). But variation in polyandry between populations is  
453 genetic and not simply explained by these proximate effects (Taylor *et al.*, 2016).  
454 Similarly, the genetic cline is opposite to that expected if polyandry had evolved in  
455 response to higher rates of heat-induced male sterility. More generally, variation in male  
456 fertility could have favoured the evolution of behavioural plasticity in polyandry. Using  
457 females from distinct genetic backgrounds that differ in polyandry (Taylor *et al.*, 2016;  
458 Sutter *et al.*, 2019b), we found that females from all backgrounds substantially  
459 elevated polyandry levels after mating with sub-fertile males, suggesting behavioural  
460 plasticity was largely independent of genetic variation in polyandry. Including

461 reproductive output as a covariate meant our tests were controlled for variation in  
462 reproductive output among isolines (see above). Unfortunately, our power to detect a  
463 potential subtle genotype-by-treatment interaction for polyandry was limited by the low  
464 mating success of heat-exposed males (Fig 3 & Table S5). This means we cannot  
465 comprehensively rule out that there may be genetic variation in behavioural plasticity of  
466 polyandry. Selection may in general favour females that make reproductive decisions  
467 dynamically and flexibly (Gowaty, 2013; Ah-King & Gowaty, 2016). In the context of  
468 male infertility, females appear to update their remating decisions according to their  
469 current state (Gowaty & Hubbell, 2009), and to indeed dynamically lower their mate  
470 acceptance threshold when sperm storage is low.

#### 471 Consequences for populations

472 Plastically elevated polyandry levels have important implications for population  
473 viability (Holman & Kokko, 2013), particularly for populations under threat due to  
474 rising male infertility, and for targets of the sterile insect technique (SIT). First, climate  
475 change means that many organisms are likely to face increased male fertility problems  
476 (Walsh et al., 2019). If females increase remating after mating with infertile males, heat-  
477 induced male infertility may have little impact on population productivity as long as  
478 there are enough fertile males. Little is known about the heritability of temperature  
479 sensitivity of male fertility (Walsh et al., 2019). But, if variation in male fertility is  
480 heritable and continuous, more intense postcopulatory sexual selection due to increased  
481 polyandry (Morimoto *et al.*, 2019) will increase reproductive skew towards fully fertile  
482 males, which may accelerate adaptation to increasing temperatures and delay population  
483 extinction (Parrett & Knell, 2018). Second, plastically elevated polyandry thwarts  
484 population control attempts through SIT (Kraaijeveld & Chapman, 2004; Barclay,

485 2005). Thus, understanding short-term plasticity in polyandry as well as the amount of  
486 genetic variation underlying this plasticity is important for predicting the potential of  
487 SIT. For example, even if the average female shows no increased remating after mating  
488 with sterile males, populations may still harbour genetic variation in female remating  
489 behaviour. This would lead to an increase in polyandry in response to SIT across  
490 generations, hence hampering SIT effectiveness.

#### 491 Conclusions

492 Mating failure is common, and represents a potential explanation for the ubiquity of  
493 female multiple mating. Male fertility is often compromised by natural processes and  
494 human intervention. Here, we have shown that females flexibly adjusted their remating  
495 rate according to their demands for fertile sperm, consistent with behavioural plasticity  
496 that was largely independent of genetic variation in polyandry. Polyandry allowed  
497 females to buffer against fitness costs associated with mating with heat-exposed males  
498 with low fertility, which may hamper the impact of release of sterile males for  
499 population control, but may increase selection on male fertility and assist adaptation to  
500 increasing global temperatures.

#### 501 **References**

- 502 Abraham, S., Lara-Pérez, L.A., Rodríguez, C., Contreras-Navarro, Y., Nuñez-Beverido,  
503 N., Ovruski, S., *et al.* 2016. The male ejaculate as inhibitor of female remating in  
504 two tephritid flies. *J. Insect Physiol.* **88**: 40–47.
- 505 Abraham, S., Liendo, M.C., Devescovi, F., Peralta, P.A., Yusef, V., Ruiz, J., *et al.* 2013.  
506 Remating behavior in *Anastrepha fraterculus* (Diptera: Tephritidae) females is  
507 affected by male juvenile hormone analog treatment but not by male sterilization.

508 *Bull. Entomol. Res.* **103**: 310–317.

509 Adkins-Regan, E. 2015. Hit or miss: Fertilization outcomes of natural inseminations by  
510 Japanese quail. *PLoS One* **10**: 1–20.

511 Ah-King, M. & Gowaty, P.A. 2016. A conceptual review of mate choice: stochastic  
512 demography, within-sex phenotypic plasticity, and individual flexibility. *Ecol.*  
513 *Evol.* **6**: 4607–4642.

514 Arnqvist, G. & Nilsson, T. 2000. The evolution of polyandry: multiple mating and  
515 female fitness in insects. *Anim. Behav.* **60**: 145–164.

516 Avent, T.D., Price, T.A.R. & Wedell, N. 2008. Age-based female preference in the fruit  
517 fly *Drosophila pseudoobscura*. *Anim. Behav.* **75**: 1413–1421.

518 Barclay, H. 2005. Mathematical Models for the Use of Sterile Insects. In: *Sterile Insect*  
519 *Technique: Principles and Practice in Area-Wide Integrated Pest Management* (V.  
520 Dyck, J. Hendrichs, & A. Robinson, eds), pp. 147–174. Springer, Dordrecht, The  
521 Netherlands.

522 Bates, D., Maechler, M., Bolker, B. & Walker, S. 2015. Fitting linear mixed-effects  
523 models using lme4. *J. Stat. Softw.* **67**: 1–48.

524 Bolker, B.M., Brooks, M.E., Clark, C.J., Geange, S.W., Poulsen, J.R., Stevens, M.H.H.,  
525 *et al.* 2009. Generalized linear mixed models: a practical guide for ecology and  
526 evolution. *Trends Ecol. Evol.* **24**: 127–135.

527 Brooks, M.E., Kristensen, K., van Benthem, , Koen J. Magnusson, A., Berg, C.W.,  
528 Nielsen, A., Skaug, H.J., *et al.* 2017. glmmTMB Balances Speed and Flexibility  
529 Among Packages for Zero-inflated Generalized Linear Mixed Modeling. *R J.* **9**:  
530 378–400.

531 Calkins, C. & Parker, A. 2005. Sterile Insect Quality. In: *Sterile Insect Technique:*

532 *Principles and Practice in Area-Wide Integrated Pest Management* (V. A. Dyck, J.  
533 Hendrichs, & A. Robinson, eds), pp. 269–296. Springer, Dordrecht, The  
534 Netherlands.

535 Champion de Crespigny, F.E., Hurst, L.D. & Wedell, N. 2008. Do *Wolbachia*-  
536 associated incompatibilities promote polyandry? *Evolution* **62**: 107–122.

537 Crawley, M.J. 2007. *The R Book*. John Wiley & Sons, Chichester.

538 Crudginton, H.S., Beckerman, A.P., Brüstle, L., Green, K. & Snook, R.R. 2005.  
539 Experimental removal and elevation of sexual selection: does sexual selection  
540 generate manipulative males and resistant females? *Am. Nat.* **165**: S72–S87.

541 David, J.R., Araripe, L.O., Chakir, M., Legout, H., Lemos, B., Pétavy, G., *et al.* 2005.  
542 Male sterility at extreme temperatures: A significant but neglected phenomenon for  
543 understanding *Drosophila* climatic adaptations. *J. Evol. Biol.* **18**: 838–846.

544 Delisle, J. & Hardy, M. 1997. Male nutrition influences the reproductive success of both  
545 sexes of the spruce budworm *Choristoneura fumiferana* (Lepidoptera: Tortricidae).  
546 *Funct. Ecol.* **11**: 451–463.

547 Dowling, D.K., Tompkins, D.M. & Gemmell, N.J. 2015. The Trojan Female Technique  
548 for pest control: A candidate mitochondrial mutation confers low male fertility  
549 across diverse nuclear backgrounds in *Drosophila melanogaster*. *Evol. Appl.* **8**:  
550 871–880.

551 Dyck, V.A., Hendrichs, J. & Robinson, A.S. 2005. *Sterile Insect Technique: Principles*  
552 *and Practice in Area-Wide Integrated Pest Management* (V. A. Dyck, J.  
553 Hendrichs, & A. S. Robinson, eds). Springer, Dordrecht, The Netherlands.

554 Fisher, D.N., Doff, R.J. & Price, T.A.R. 2013. True polyandry and pseudopolyandry:  
555 why does a monandrous fly remate? *BMC Evol. Biol.* **13**: 157.

556 Forbes, S. 2014. Partial fertility and polyandry: a benefit of multiple mating hiding in  
557 plain sight? *Behav. Ecol. Sociobiol.* **68**: 1329–1334.

558 Forstmeier, W., Ihle, M., Opatová, P., Martin, K., Knief, U., Albrechtová, J., *et al.*  
559 2017. Testing the phenotype-linked fertility hypothesis in the presence and absence  
560 of inbreeding. *J. Evol. Biol.* **30**: 968–976.

561 Forstmeier, W. & Schielzeth, H. 2011. Cryptic multiple hypotheses testing in linear  
562 models: Overestimated effect sizes and the winner’s curse. *Behav. Ecol. Sociobiol.*  
563 **65**: 47–55.

564 Friesen, C.R., Uhrig, E.J. & Mason, R.T. 2014. Females remate more frequently when  
565 mated with sperm-deficient males. *J. Exp. Zool. Part A Ecol. Genet. Physiol.* **321**:  
566 603–609.

567 Gabrieli, P., Scolari, F., Di Cosimo, A., Savini, G., Fumagalli, M., Gomulski, L.M., *et*  
568 *al.* 2016. Sperm-less males modulate female behaviour in *Ceratitis capitata*  
569 (Diptera: Tephritidae). *Insect Biochem. Mol. Biol.* **79**: 13–26.

570 Garcia-Gonzalez, F. 2004. Infertile matings and sperm competition: the effect of  
571 ‘‘nonsperm representation’’ on intraspecific variation in sperm precedence  
572 patterns. *Am. Nat.* **164**: 457–472.

573 Gavriel, S., Gazit, Y. & Yuval, B. 2009. Remating by female Mediterranean fruit flies  
574 (*Ceratitis capitata*, Diptera: Tephritidae): Temporal patterns and modulation by  
575 male condition. *J. Insect Physiol.* **55**: 637–642.

576 Gibson, R.M. & Jewell, P.A. 1982. Semen quality, female choice and multiple mating  
577 in domestic sheep: A test of Trivers’ Sexual Competence Hypothesis. *Behaviour*  
578 **80**: 9–31.

579 Gowaty, P.A. 2013. Adaptively flexible polyandry. *Anim. Behav.* **86**: 877–884.

580 Gowaty, P.A. & Hubbell, S.P. 2009. Reproductive decisions under ecological  
581 constraints: it's about time. *Proc. Natl. Acad. Sci. U. S. A.* **106 Suppl**: 10017–  
582 10024.

583 Greenway, E., Dougherty, L.R. & Shuker, D.M. 2015. Mating failure. *Curr. Biol.* **25**:  
584 R534–R536.

585 Greenway, E. & Shuker, D. 2015. The repeatability of mating failure in a polyandrous  
586 bug. *J. Evol. Biol.* **28**: 1578–1582.

587 Greenway, E.V., Balfour, V.L. & Shuker, D.M. 2017. Can females choose to avoid  
588 mating failure in the seed bug *Lygaeus simulans*? *Anim. Behav.* **129**: 61–69.

589 Hadfield, J.D. 2010. MCMC Methods for Multi-Response Generalized Linear Mixed  
590 Models: The MCMCglmm R Package. *J. Stat. Softw.* **33**: 1–22.

591 Haq, I.U., Vreysen, M.J.B., Abd-Alla, A. & Hendrichs, J. 2013. Ability of Genetic  
592 Sexing Strain male melon flies (Diptera: Tephritidae) to suppress wild female  
593 remating: Implications for SIT. *Florida Entomol.* **96**: 839–849.

594 Harmer, A.M.T., Radhakrishnan, P. & Taylor, P.W. 2006. Remating inhibition in  
595 female Queensland fruit flies: Effects and correlates of sperm storage. *J. Insect*  
596 *Physiol.* **52**: 179–186.

597 Hartig, F. 2018. DHARMA: Residual Diagnostics for Hierarchical (Multi-Level /  
598 Mixed) Regression Models. R package version 0.2.0. , doi: 10.1111/j.1753-  
599 0407.2009.00036.x.

600 Hasson, O. & Stone, L. 2009. Male infertility, female fertility and extrapair copulations.  
601 *Biol. Rev.* **84**: 225–244.

602 Herrera, P., Taylor, M.L., Skeats, A., Price, T.A.R. & Wedell, N. 2014. Can patterns of  
603 chromosome inversions in *Drosophila pseudoobscura* predict polyandry across a

604 geographical cline? *Ecol. Evol.* **4**: 3072–3081.

605 Hibino, Y. & Iwahashi, O. 1991. Appearance of wild females unreceptive to sterilized  
606 males on Okinawa Is. in the eradication program of the Melon fly, *Dacus*  
607 *cucurbitae* Coquillett (Diptera: Tephritidae). *Appl. Entomol. Zool.* **26**: 265–270.

608 Holman, L. & Kokko, H. 2013. The consequences of polyandry for population viability,  
609 extinction risk and conservation. *Philos. Trans. R. Soc. B Biol. Sci.* **368**: 20120053.

610 Hurley, L.L., McDiarmid, C.S., Friesen, C.R., Griffith, S.C. & Rowe, M. 2018.  
611 Experimental heatwaves negatively impact sperm quality in the zebra finch. *Proc.*  
612 *R. Soc. B Biol. Sci.* **285**.

613 Janowitz, S.A. & Fischer, K. 2011. Opposing effects of heat stress on male versus  
614 female reproductive success in *Bicyclus anynana* butterflies. *J. Therm. Biol.* **36**:  
615 283–287.

616 Knipling, E.F. 1955. Possibilities of insect control or eradication through the use of  
617 sexually sterile males. *J. Econ. Entomol.* **48**: 459–462.

618 Kraaijeveld, K. & Chapman, T. 2004. Effects of male sterility on female remating in the  
619 Mediterranean fruitfly, *Ceratitis capitata*. *Proc. R. Soc. B Biol. Sci.* **271**: S209–  
620 S211.

621 Krokene, C., Rigstad, K., Dale, M. & Lifjeld, J.T. 1998. The function of extrapair  
622 paternity in blue tits and great tits: good genes or fertility insurance? *Behav. Ecol.*  
623 **9**: 649–656.

624 Krüger, A.P., Schlesener, D.C.H., Martins, L.N., Wollmann, J., Deprá, M. & Garcia,  
625 F.R.M. 2019. Radiation effects on *Drosophila suzukii* (Diptera: Drosophilidae)  
626 reproductive behaviour. *J. Appl. Entomol.* **143**: 88–94.

627 Kuriwada, T., Kumano, N., Shiromoto, K., Haraguchi, D., Matsuyama, T. & Kohama,

628 T. 2014. Female preference did not evolve under laboratory conditions in the  
629 solanaceous fruit fly *Bactrocera latifrons*. *Int. J. Pest Manag.* **60**: 160–165.

630 Kuznetsova, A., Brockhoff, P.B. & Christensen, R.H.B. 2016. lmerTest: 2.0-33., Tests  
631 in Linear Mixed Effects Models. R package version 2.0-33.

632 Landeta-Escamilla, A., Hernández, E., Arredondo, J., Díaz-Fleischer, F. & Pérez-  
633 Staples, D. 2016. Male irradiation affects female remating behavior in *Anastrepha*  
634 *serpentina* (Diptera: Tephritidae). *J. Insect Physiol.* **85**: 17–22.

635 Liu, H. & Kubli, E. 2003. Sex-peptide is the molecular basis of the sperm effect in  
636 *Drosophila melanogaster*. *Proc. Natl. Acad. Sci.* **100**: 9929–9933.

637 Lorch, P.D. & Chao, L. 2003. Selection for multiple mating in females due to mates that  
638 reduce female fitness. *Behav. Ecol.* **14**: 679–686.

639 Malo, A.F., Roldan, E.R., Garde, J., Soler, A.J. & Gomendio, M. 2005. Antlers honestly  
640 advertise sperm production and quality. *Proc. R. Soc. B Biol. Sci.* **272**: 149–57.

641 Manning, A. 1967. The control of sexual receptivity in female *Drosophila*. *Anim.*  
642 *Behav.* **15**: 239–250.

643 Mautz, B.S., Møller, A.P. & Jennions, M.D. 2013. Do male secondary sexual characters  
644 signal ejaculate quality? A meta-analysis. *Biol. Rev.* **88**: 669–682.

645 Mcinnis, D.O., Lance, D.R. & Jackson, C.G. 1996. Behavioral resistance to the sterile  
646 insect technique by Mediterranean fruit fly (Diptera: Tephritidae) in Hawaii. *Ann.*  
647 *Entomol. Soc. Am.* **89**: 739–744.

648 Meehl, G.A. 2004. More Intense, More Frequent, and Longer Lasting Heat Waves in  
649 the 21st Century. *Science (80-. )*. **305**: 994–997.

650 Miyatake, T., Chapman, T. & Partridge, L. 1999. Mating-induced inhibition of remating  
651 in female mediterranean fruit flies *Ceratitis capitata*. *J. Insect Physiol.* **45**: 1021–

652 1028.

653 Morimoto, J., McDonald, G.C., Smith, E., Smith, D.T., Perry, J.C., Chapman, T., *et al.*  
654 2019. *Sex peptide receptor*-regulated polyandry modulates the balance of pre- and  
655 post-copulatory sexual selection in *Drosophila*. *Nat. Commun.* **10**: 283.

656 Morrow, E.H., Arnqvist, G. & Pitcher, T.E. 2002. The evolution of infertility: Does  
657 hatching rate in birds coevolve with female polyandry? *J. Evol. Biol.* **15**: 702–709.

658 Mossinson, S. & Yuval, B. 2003. Regulation of sexual receptivity of female  
659 Mediterranean fruit flies: Old hypotheses revisited and a new synthesis proposed.  
660 *J. Insect Physiol.* **49**: 561–567.

661 Olsson, M. & Shine, R. 1997. Advantages of multiple matings to females: a test of the  
662 infertility hypothesis using lizards. *Evolution* **51**: 1684–1688.

663 Parrett, J.M. & Knell, R.J. 2018. The effect of sexual selection on adaptation and  
664 extinction under increasing temperatures. *Proc. R. Soc. B Biol. Sci.* **285**: 20180303.

665 Preston, B.T., Stevenson, I.R., Pemberton, J.M. & Wilson, K. 2001. Dominant rams  
666 lose out by sperm depletion. *Nature* **409**: 681–682.

667 Price, T.A.R., Lewis, Z., Smith, D.T., Hurst, G.D.D. & Wedell, N. 2011. Remating in  
668 the laboratory reflects rates of polyandry in the wild. *Anim. Behav.* **82**: 1381–1386.

669 Price, T.A.R. & Wedell, N. 2008. Selfish genetic elements and sexual selection: their  
670 impact on male fertility. *Genetica* **132**: 295–307.

671 R Core Team. 2018. R: A language and environment for statistical computing. R  
672 Foundation for Statistical Computing, Vienna, Austria. URL [http://www.R-](http://www.R-project.org/)  
673 [project.org/](http://www.R-project.org/).

674 Reding, L. 2015. Increased hatching success as a direct benefit of polyandry in birds.  
675 *Evolution* **69**: 264–270.

676 Reinhardt, K., Dobler, R. & Abbott, J. 2015. An Ecology of Sperm: Sperm  
677 Diversification by Natural Selection. *Annu. Rev. Ecol. Evol. Syst.* **46**: 435–459.

678 Rhainds, M. 2010. Female mating failures in insects. *Entomol. Exp. Appl.* **136**: 211–  
679 226.

680 Sakaluk, S.K. & Cade, W.H. 1980. Female mating frequency and progeny production in  
681 singly and doubly mated house and field crickets. *Can. J. Zool.* **58**: 404–411.

682 Sales, K., Vasudeva, R., Dickinson, M.E., Godwin, J.L., Lumley, A.J., Michalczyk, Ł.,  
683 *et al.* 2018. Experimental heatwaves compromise sperm function and cause  
684 transgenerational damage in a model insect. *Nat. Commun.* **9**: 4771.

685 Schielzeth, H. 2010. Simple means to improve the interpretability of regression  
686 coefficients. *Methods Ecol. Evol.* **1**: 103–113.

687 Schielzeth, H. & Forstmeier, W. 2009. Conclusions beyond support: overconfident  
688 estimates in mixed models. *Behav. Ecol.* **20**: 416–420.

689 Schindelin, J., Arganda-Carreras, Ignacio Frise, E., Kaynig, V., Longair, M., Pietzsch,  
690 T., Preibisch, S., *et al.* 2012. Fiji: an open-source platform for biological-image  
691 analysis. *Nat. Methods* **9**: 676–682.

692 Schmoll, T., Kleven, O. & Reynolds, J. 2016. Functional infertility in a wild passerine  
693 bird. *Ibis (Lond. 1859)*. **158**: 670–673.

694 Setchell, B.P. 2006. The effects of heat on the testes of mammals. *Anim. Reprod.* **3**: 81–  
695 91.

696 Sheldon, B.C. 1994. Male phenotype, fertility, and the pursuit of extra-pair copulations  
697 by female birds. *Proc. R. Soc. B Biol. Sci.* **257**: 25–30.

698 Snook, R.R. & So, Y.K. 2000. Associations between female remating behavior,  
699 oogenesis and oviposition in *Drosophila melanogaster* and *Drosophila*

700 *pseudoobscura*. *J. Insect Physiol.* **46**: 1489–1496.

701 Sutter, A., Travers, L.M., Oku, K., Delaney, K., Store, S., Price, T.A.R., *et al.* 2019a.  
702 Data from: Flexible polyandry in female flies is an adaptive response to infertile  
703 males. *Dryad Digital Repository* <http://dx.doi.org/10.5061/dryad.ct8r61v>.

704 Sutter, A., Travers, L.M., Weedon, M., Oku, K., Price, T.A.R. & Wedell, N. 2019b. No  
705 selection for change in polyandry under experimental evolution. *J. Evol. Biol.*  
706 *jeb.13476*.

707 Taylor, D.R. & Ingvarsson, P.K. 2003. Common features of segregation distortion in  
708 plants and animals. *Genetica* **117**: 27–35.

709 Taylor, M.L., Price, T.A.R., Skeats, A. & Wedell, N. 2016. Temperature can shape a  
710 cline in polyandry, but only genetic variation can sustain it over time. *Behav. Ecol.*  
711 **27**: 462–469.

712 Taylor, M.L., Price, T.A.R. & Wedell, N. 2014. Polyandry in nature: a global analysis.  
713 *Trends Ecol. Evol.* **29**: 376–383.

714 Taylor, M.L., Wedell, N. & Hosken, D.J. 2008. Sexual selection and female fitness in  
715 *Drosophila simulans*. *Behav. Ecol. Sociobiol.* **62**: 721–728.

716 Torres-Vila, L.M., Stockel, J. & Rodríguez-Molina, M.C. 1997. Physiological factors  
717 regulating polyandry in *Lobesia botrana* (Lepidoptera: Tortricidae). *Physiol.*  
718 *Entomol.* **22**: 387–393.

719 Uller, T. & Olsson, M. 2005. Multiple copulations in natural populations of lizards:  
720 evidence for the fertility assurance hypothesis. *Behaviour* **142**: 45–56.

721 Walsh, B.S., Parratt, S.R., Hoffmann, A.A., Atkinson, D., Snook, R.R., Bretman, A., *et*  
722 *al.* 2019. The Impact of Climate Change on Fertility. *Trends Ecol. Evol.* **34**: 249–  
723 259.

724 Wetton, J.H. & Parkin, D.T. 1991. An association between fertility and cuckoldry in the  
725 house sparrow, *Passer domesticus*. *Proc. R. Soc. B Biol. Sci.* **245**: 227–233.  
726

727 **Figure legends**

728 **Figure 1:** Male mating success and latency. Heat-exposed males (red) had a longer  
729 copulation latency and reduced mating success compared to control males (blue; see  
730 main text and Table S1). Thin lines represent approximate 95% confidence intervals  
731 from a cox proportional hazard model on right-censored mating latency with other fixed  
732 effects centred. Note the log-scale of the x axis.

733 **Figure 2:** Male heat-exposure reduces female reproductive output, but polyandry can  
734 restore fitness. Framed circles and error bars depict mean and approximate 95%  
735 confidence intervals. Faint circles represent raw data, with circle area proportional to the  
736 number of observations. Under enforced monandry, females mated to heat-exposed  
737 males had consistently low reproductive fitness (left panel). Females often chose not to  
738 remate when initial reproductive output was substantial after mating with heat-exposed  
739 males, but soon after showed reduced reproductive output (central panel). Remating  
740 with fertile males fully restored subsequent reproductive fitness in females that had  
741 mated with heat-exposed males (right panel).

742 **Figure 3:** Females increase polyandry after mating with heat-exposed males through  
743 behavioural plasticity. Isolines were assigned a colour gradient according to polyandry  
744 at the control temperature. Polyandry was consistently higher after mating with heat-  
745 exposed males (right) versus control males (left; Table 3). The area of circles is  
746 proportional to the sample size. Raw values and sample sizes are given in Table S5.  
747 Note the smaller sample sizes for females first mated to heat-exposed males due to low  
748 mating success of heat-exposed males, limiting the power to detect genetic variation in  
749 behavioural plasticity.

750 **Table 1:** Summary statistics and sample sizes.

Temperature	Control (23°C)		Heat-exposure (31°C)		Heat effect	Full model	Illustration
	A	B	A	B			
Male cohort							
<i>main experiment</i>							
Male mortality (N)	2% (285)	0.8% (260)	52% (460)	2% (510)	(↑)	Table S2	Fig S3
Mating success (N)	91% (163)	84% (230)	24% (148)	33% (381)	↓	Table S1	Fig 1
Copulation latency [min] (N)	3.4±3.2 (147)	8.2±17.5 (192)	16.5±25.8 (36)	18.5±23.6 (124)	↑	Table S1	Fig 1 & S1
Copulation duration [sec] (N)	6.6±2.3 (148)	5.9±1.8 (194)	4.9±2.0 (36)	5.7±5.1 (123)	↓	Table S1	Fig S1
4d fecundity (N)	42.3±19.8 (147)	43.2±19.0 (192)	33.3±25.0 (35)	9.8±18.6 (125)	↓	Table 2	Fig 2
Polyandry (N)	44% (147)	40% (136)	77% (35)	85% (109)	↑	Table 3	Fig 3
<i>additional males</i>							
Male fertility (N)	4.8±0.7 (9)	3.1±1.2 (8)	2.0±1.3 (11)	0.5±0.7 (11)	↓		
4d fecundity (N)	229±65 (9)	131±63 (8)	78±38 (9)	41±26 (5)	↓		

751 Given are mean, standard deviation and sample sizes for survival, mating behaviours and reproductive output. The effect of male heat-  
 752 exposure is indicated by arrows. For detailed results see the full models as indicated in the last column.

753 **Table 2:** Model summary for female reproductive output.

<i>Fixed/Random effects</i>	<b>Conditional model</b>				<b>Zero-inflation model</b>					
	<b>Coef</b>	<b>SE (Coef)</b>	<b>z</b>	<b>p</b>	<b>Var</b>	<b>SD</b>	<b>Coef</b>	<b>SE (Coef)</b>	<b>z</b>	<b>p</b>
Intercept [control; forced monandry; Vial A (d1–5)]	42.777	2.775	15.41	<b>&lt;0.001</b>			–5.624	0.942	–5.97	<b>&lt;0.001</b>
Heat-exposure	–28.826	7.456	–3.87	<b>&lt;0.001</b>			6.509	0.778	8.37	<b>&lt;0.001</b>
Chosen monandry (Mono)	3.169	2.544	1.25	0.213			–0.648	1.020	–0.64	0.525
Chosen polyandry (Poly)	0.894	2.705	0.33	0.741			3.876	0.904	4.29	<b>&lt;0.001</b>
First mate's size (centred & scaled)	–1.106	0.565	–1.96	<b>0.050</b>						
Vial B (d5–9)	–16.407	3.038	–5.40	<b>&lt;0.001</b>			2.395	0.934	2.57	<b>0.010</b>
Vial C (d9–13)	–6.657	3.186	–2.09	<b>0.037</b>			3.512	0.918	3.83	<b>&lt;0.001</b>
Male cohort (A)							–0.705	0.226	–3.12	<b>0.002</b>
Heat:Mono	30.004	8.566	3.50	<b>&lt;0.001</b>			–0.632	0.725	–0.87	0.383
Heat:Poly	17.664	7.976	2.22	<b>0.027</b>			–4.237	0.696	–6.09	<b>&lt;0.001</b>
Heat:Vial B	28.646	11.099	2.58	<b>0.010</b>			–2.425	0.636	–3.81	<b>&lt;0.001</b>
Heat:Vial C	0.711	12.416	0.06	0.954			–3.140	0.558	–5.63	<b>&lt;0.001</b>
Mono:Vial B	–5.725	3.514	–1.63	0.103			0.772	0.949	0.81	0.416
Mono:Vial C	–11.299	3.683	–3.07	<b>0.002</b>			0.897	0.977	0.92	0.358
Poly:Vial B	–4.031	3.725	–1.08	0.279			–3.312	0.854	–3.88	<b>&lt;0.001</b>

Table 2 (continued)


Fixed/Random effects	Conditional model				Zero-inflation model					
	Coef	SE (Coef)	z	p	Var	SD	Coef	SE (Coef)	z	p
Poly:Vial C	-6.831	3.914	-1.75	0.081			-3.287	0.850	-3.87	<0.001
Heat:Mono:Vial B	-32.379	12.860	-2.52	<b>0.012</b>						
Heat:Mono:Vial C	-21.605	14.467	-1.49	0.135						
Heat:Poly:Vial B	-15.155	11.634	-1.30	0.193						
Heat:Poly:Vial C	12.732	12.929	0.99	0.325						
<i>Individual female</i>					7.70	2.77				
<i>Female:Vial (random slopes)</i>					<0.01	0.02				
<i>Female isoline (10 levels)</i>					26.17	5.12				
<i>Male collection batch (16 levels)</i>					3.59	1.90				
<i>Residual</i>					245.20	15.66				


754 The conditional model describes the Gaussian component of female reproductive output (498 females) while the zero-inflation model  
755 accounts for the likelihood of reproductive failure. The model with the lowest AIC value was chosen as the best model. See Table S4 for an  
756 overview of models and associated AIC values.


757 **Table 3:** Full model summary for polyandry.

binomial GLMM (N = 427)						
<b>Fixed/Random effects</b>	<b>Coef</b>	<b>SE (Coef)</b>	<b>z</b>	<b>p</b>	<b>Var</b>	<b>SD</b>
Intercept (control temperature)	-0.163	0.246	-0.66	0.508		
Heat-exposure	1.910	0.408	4.68	<b>&lt;0.001</b>		
4d reproductive output (centred & scaled)	-0.411	0.164	-2.51	<b>0.012</b>		
First mate's size (centred & scaled)	-0.220	0.136	-1.62	0.105		
Female age (centred)	0.513	0.271	1.89	0.059		
Order in assay (centred & scaled)	0.097	0.145	0.67	0.504		
Heat:Reproductive_output	-0.337	0.295	-1.14	0.253		
Heat:First_mate_size	-0.650	0.348	-1.87	0.062		
<i>Male collection batch (16 levels)</i>					<0.001	<0.001
<i>Female isoline (10 levels)</i>					0.38	0.62
<i>Heat:Female_isoline (random slopes)</i>					0.21	0.46

758 Four-day reproductive output corresponds to the number of offspring eclosed from the  
 759 vial in which a female was housed between her first mating and the remating  
 760 opportunity. Random slopes for female isolines were included to test for genetic  
 761 variation in behavioural plasticity (G x E; see Fig 3).







# Electronic supplementary material

## Adaptively flexible polyandry by female flies when males are infertile

*Andreas Sutter*<sup>1,2,†</sup>, *Laura M Travers*<sup>1,2</sup>, *Keiko Oku*<sup>1</sup>, *Kynan Delaney*<sup>1</sup>, *Stefan Store*<sup>1</sup>, *Tom A. R. Price*<sup>3</sup>,  
& *Nina Wedell*<sup>1</sup>

1. Centre for Ecology & Conservation, College of Life and Environmental Sciences, University of Exeter, Cornwall Campus, TR10 9FE, Penryn, UK.
2. School of Biological Sciences, Norwich Research Park, University of East Anglia, Norwich NR4 7TJ, UK
3. Institute of Integrative Biology, University of Liverpool, Crown Street, Liverpool, L69 7ZB, UK

† [a.sutter@uea.ac.uk](mailto:a.sutter@uea.ac.uk)

## Supplementary results:

### Male courtship behaviour

To investigate whether reduced mating success for heat-exposed males was caused by female discrimination against heat-exposed males or reduced courtship by heat-exposed males, we recorded and analysed data on courtship latency in the second experimental block. Courtship latency was longer for heat-exposed males. This was true both for eventually successful and unsuccessful males (Table 2; Fig S1). Additionally, of the males that did not mate, heat-exposed males were more likely not to have been observed courting (71% versus 50% for control males). And in the subset of males that were observed to both court and mate, latency from courtship initiation to mating tended to be longer for heat-exposed males (3.6 versus 1.5min), though the effect was not statistically significant, probably because of the small sample size (N = 52). In combination, these results suggest that heat-exposed males were slower and less likely to initiate courtship, and that their courtship quality or intensity may have been inferior to that of control males.

### Estimating genetic variation in polyandry and behavioural plasticity

In addition to simply testing for significant effects of female isoline and its interaction with male treatment on polyandry (described in the main text), we used a bivariate model in MCMCglmm to estimate among-isoline variances, and covariance between polyandry of females that had mated with control or with heat-exposed males. For fixed effects (specified in Table 3) and the random effect associated with female isoline, we fitted an unstructured variance-covariance matrix that allows estimation of covariances between parts of the model. We fitted variances but no covariances for the random effect male collection batch. We fixed the residual variance for polyandry (binary outcome) at 10, and rescaled random effect variance estimates as  $Var/(1 + c2 * 10)$ , where  $c2 = ((16 * \sqrt{3})/(15 * \pi))^2$ , following Jarrod Hadfield's MCMCglmm course notes (<https://cran.r-project.org/web/packages/MCMCglmm/vignettes/CourseNotes.pdf>). The model was run for 4,050,000 iterations with a thinning interval of 2000 and a burn-in of 50,000 with parameter-inflated priors. This resulted in 2000 samples from the posterior for which autocorrelation between successive samples for parameters was less than 0.1.

Female isoline explained a substantial proportion of variation in polyandry of females after mating with control males (posterior mode [95% credible interval] = 0.12 [0.025, 0.39]), whereas very little variation in polyandry of females mated with heat-exposed males was explained by isoline identity (0.002 [ $<0.0001$ , 0.42]). Finally, there was no clear correlation (estimated from (co)variances) between isoline female behaviour after mating with control or heat-exposed males (0.31 [-0.55, 0.88]; note the very large credible interval).

**Table S1: Model summaries for mating behaviours**

Fixed/Random effects	Mating success (binomial GLMM; N = 916)						Copulation latency (log LMM; N = 496)						Copulation duration (log LMM; N = 487)					
	Coef	SE (Coef)	z	p	Var	SD	Coef	SE (Coef)	t	p	Var	SD	Coef	SE (Coef)	t	p	Var	SD
Intercept (control)	2.078	0.223	9.31	<0.001	-	-	5.248	0.104	50.70	<0.001	-	-	5.866	0.060	97.71	<0.001	-	-
Heat-exposure	-3.116	0.219	-14.21	<0.001	-	-	1.056	0.113	9.37	<0.001	-	-	-0.252	0.051	-4.99	<0.001	-	-
Cohort (B→A; centred)	0.357	0.340	1.05	0.294	-	-	-0.446	0.113	-3.94	<0.001	-	-	0.109	0.051	2.14	0.033	-	-
Heat:Cohort	-0.980	0.406	-2.41	0.016	-	-	0.233	0.221	1.05	0.293	-	-	-0.168	0.099	-1.69	0.092	-	-
Block (centred)	-0.548	0.185	-2.96	0.003	-	-	0.101	0.099	1.03	0.308	-	-	0.110	0.044	2.49	0.014	-	-
Female age (centred)	-0.040	0.218	-0.18	0.855	-	-	-0.182	0.121	-1.51	0.134	-	-	-0.082	0.054	-1.51	0.134	-	-
Male size (centred & scaled)	0.324	0.097	3.36	0.001	-	-	-0.035	0.053	-0.65	0.514	-	-	-0.036	0.024	-1.51	0.132	-	-
Order in assay (centred & scaled)	-0.346	0.092	-3.75	<0.001	-	-	0.055	0.048	1.15	0.255	-	-	0.007	0.021	0.34	0.735	-	-
Mating (yes vs no)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female housing vial ( $\leq 93$ levels)	-	-	-	-	0.059	0.244	-	-	-	-	<0.001	<0.001	-	-	-	-	<0.001	<0.001
Female isoline (10 levels)	-	-	-	-	0.061	0.247	-	-	-	-	0.009	0.096	-	-	-	-	<0.001	<0.001
Male population (4 levels)	-	-	-	-	0.052	0.227	-	-	-	-	0.027	0.164	-	-	-	-	0.012	0.109
Residual	-	-	-	-	-	-	-	-	-	-	0.987	0.994	-	-	-	-	0.197	0.444

Fixed/Random effects	Courtship latency (log LMM; N = 127)						Courtship duration (log LMM; N = 52)					
	Coef	SE (Coef)	t	p	Var	SD	Coef	SE (Coef)	t	p	Var	SD
Intercept (control)	5.924	0.266	22.29	<0.001	-	-	4.531	0.341	13.29	<0.001	-	-
Heat-exposure	1.191	0.289	4.11	<0.001	-	-	0.764	0.562	1.36	0.180	-	-
Cohort (B→A; centred)	-0.521	0.376	-1.38	0.169	-	-	-	-	-	-	-	-
Heat:Cohort	1.079	0.521	2.07	0.041	-	-	-	-	-	-	-	-
Block (centred)	-	-	-	-	-	-	-	-	-	-	-	-
Female age (centred)	-0.107	0.315	-0.34	0.743	-	-	-	-	-	-	-	-
Male size (centred & scaled)	-0.098	0.143	-0.69	0.496	-	-	-0.219	0.290	-0.76	0.454	-	-
Order in assay (centred & scaled)	0.313	0.124	2.53	0.013	-	-	-	-	-	-	-	-
Mating (yes vs no)	-0.933	0.276	-3.38	0.001	-	-	-	-	-	-	-	-
Female housing vial ( $\leq 93$ levels)	-	-	-	-	-	-	-	-	-	-	-	-
Female isoline (10 levels)	-	-	-	-	0.016	0.128	-	-	-	-	-	-
Male population (4 levels)	-	-	-	-	<0.001	<0.001	-	-	-	-	<0.001	<0.001
Residual	-	-	-	-	1.438	1.199	-	-	-	-	3.767	1.941

**Table S2:** Full model summary for male survival under heat-exposure and control temperature. Vials containing groups of up to ten males were submerged in water baths at 23°C or 31°C for two-and-a-half (cohort B; see main text) or three days (cohort A). Experimental block was centred as described in the main text. Effects associated with a p value smaller than 0.05 are highlighted in bold.

binomial GLMM (N = 1645)						
<b>Fixed/Random effects</b>	<b>Coef</b>	<b>SE (Coef)</b>	<b>z</b>	<b>p</b>	<b>Var</b>	<b>SD</b>
Intercept (control; cohort B)	5.570	0.813	6.85	<b>&lt;0.001</b>	-	-
Heat-exposure	-1.053	0.869	-1.21	0.226	-	-
Cohort (A)	-0.833	0.941	-0.89	0.376	-	-
Heat:Cohort	-3.770	1.053	-3.58	<b>&lt;0.001</b>	-	-
Block (centred)	0.929	0.349	2.67	<b>0.008</b>	-	-
<i>Housing vial (155 levels)</i>	-	-	-	-	1.51	1.23
<i>Population (4 levels)</i>	-	-	-	-	<0.001	<0.001

**Table S3:** Full model summary for male size (length of wing L3 [mm]). Only males surviving the heat-exposure/control were measured. Virgin collection batch corresponds to unique combinations of virgin collection day and population cage.

LMM (N = 925)							
<b>Fixed/Random effects</b>	<b>Coef</b>	<b>SE (Coef)</b>	<b>ddf</b>	<b>t</b>	<b>p</b>	<b>Var</b>	<b>SD</b>
Intercept (control; cohort B)	1.435	0.020	15.2	73.182	<b>&lt;0.001</b>	-	-
Heat-exposure	-0.004	0.010	907.0	-0.569	0.569	-	-
Cohort (A)	0.021	0.028	15.8	0.783	0.445	-	-
Heat:Cohort	-0.008	0.013	907.6	-0.576	0.565	-	-
<i>Virgin collection batch (16 levels)</i>	-	-	-	-	-	0.003	0.053
<i>Residual</i>	-	-	-	-	-	0.008	0.091

**Table S4:** Overview of models for reproductive output. Models were run using glmmTMB and were sorted along ascending AIC values. All conditional models included random intercepts for female ID, female isoline, male collection batch (unique combinations of population, cohort and block), and random slopes for individual females across the three laying vials (see main text).

	Conditional model										Zero-inflation model								df	AIC	ΔAIC	
	Intercept	Heat-exposure	Mating (FM, Mono, Poly)	First mate's size	Vial (A-C)	Heat:Mating	Heat:Vial	Mating:Vial	Heat:Size	Heat:Mating:Vial	Intercept	Heat-exposure	Mating (FM, Mono, Poly)	Vial (A-C)	Male cohort	Heat:Mating	Heat:Vial	Mating:Vial				Heat:Mating:Vial
Model 1	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	39	11648.5	0.0
Model 2	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	40	11650.1	1.6
Model 3	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	38	11650.3	1.9
Model 4	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	42	11654.6	6.1
Model 5	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	41	11662.5	14.0
Model 6	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	34	11664.5	16.0
Model 7	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	32	11668.0	19.5
Model 8	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	38	11668.7	20.2
Model 9	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	32	11671.9	23.4
Model 10	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	36	11672.3	23.8
Model 11	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	30	11674.6	26.1
Model 12	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	36	11676.2	27.7
Model 13	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	30	11677.0	28.6
Model 14	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	28	11678.7	30.2
Model 15	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	34	11679.3	30.9
Model 16	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	34	11681.5	33.0
Model 17	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	32	11683.4	34.9
Model 18	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	28	11686.6	38.1
Model 19	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	24	11689.0	40.5
Model 20	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	23	11690.5	42.0
Model 21	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	32	11691.7	43.2
Model 22	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	26	11692.8	44.3
Model 23	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	25	11693.9	45.4
Model 24	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	30	11697.9	49.4
Model 25	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	29	11705.9	57.4
Model 26	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	30	11769.4	120.9
Model 27	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	29	11792.6	144.1
Model 28	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	19	11805.1	156.6
Model 29	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	19	11808.8	160.3
Model 30	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	26	11839.8	191.3
Model 31	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	17	11848.1	199.7
Model 32	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	15	11861.8	213.3
Model 33	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	15	11879.2	230.7
Model 34	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	33	11879.7	231.2
Model 35	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	13	11891.8	243.3
Model 36	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	22	11915.6	267.1
Model 37	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	24	11918.9	270.4
Model 38	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	23	11923.8	275.3
Model 39	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	21	11923.8	275.3
Model 40	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	21	11923.8	275.3
Model 41	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	27	11946.7	298.2
Model 42	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	27	11967.7	319.2
Model 43	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	25	11971.9	323.4
Model 44	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	23	12006.9	358.4
Model 45	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	16	12010.8	362.3
Model 46	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	14	12020.6	372.1
Model 47	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	14	12036.6	388.1
Model 48	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	7	not converged	

**Table S5:** Summary statistics for isofemale isolines. Percentages and sample sizes for mating and remating, and early fecundity of females paired with a control (23°C) or a heat-exposed (31°C) male. Note the smaller sample sizes for polyandry and fecundity due to low mating success of heat-exposed males.

Population	Isoline	Mating				Polyandry				4d Fecundity			
		23C	N	31C	N	23C	N	31C	N	23C	N	31C	N
Show Low	SLOB3	90%	41	42%	55	55%	31	94%	18	41.6±21.8	37	9.2±18.4	22
Show Low	2SLOC4	85%	41	47%	55	14%	28	91%	23	43.5±21.2	35	5.8±11.7	26
Show Low	SLOC48	90%	40	32%	56	52%	31	65%	17	49.1±20.2	35	27.1±27.2	18
Show Low	2SLOD29	85%	40	38%	55	15%	27	68%	19	48.3±17.9	34	16.2±27.2	21
Show Low	2SLOD33	85%	40	13%	56	57%	28	100%	7	46.5±18.9	34	0±0	7
Show Low	2SLOD6	83%	40	25%	55	41%	27	92%	13	47.5±16.8	33	16±20.9	14
Lewistown	LEW17	83%	40	37%	54	27%	30	71%	17	32.5±15.1	33	18.6±22.8	20
Lewistown	LEW23	88%	41	20%	55	57%	30	78%	9	42.2±16.1	36	15.6±21.2	11
Lewistown	LEW3	93%	40	18%	56	50%	30	100%	10	37.8±17.7	37	15.5±22.9	10
Lewistown	LEW64	83%	40	30%	56	46%	26	94%	16	37.3±20.9	33	20.1±24.9	17


**Table S6:** Model summary for polyandry (*cf* Table 3), additionally including copulatory behaviour from a female's first mating

binomial GLMM (N = 416)						
<b>Fixed/Random effects</b>	<b>Coef</b>	<b>SE (Coef)</b>	<b>z</b>	<b>p</b>	<b>Var</b>	<b>SD</b>
Intercept (control temperature)	-0.200	0.256	-0.78	0.437		
Heat-exposure	1.820	0.426	4.27	<b>&lt;0.001</b>		
4d reproductive output (centred & scaled)	-0.393	0.167	-2.35	<b>0.019</b>		
First mate's size (centred & scaled)	-0.234	0.139	-1.68	0.093		
Female age (centred)	0.456	0.275	1.66	0.097		
Order in assay (centred & scaled)	0.090	0.147	0.61	0.540		
Log copulation latency (centred & scaled)	0.005	0.135	0.03	0.973		
Log copulation duration (centred & scaled)	0.074	0.136	0.55	0.585		
Heat:Reproductive_output	-0.518	0.347	-1.49	0.135		
Heat:First_mate_size	-0.437	0.302	-1.45	0.148		
<i>Male collection batch (16 levels)</i>					<0.001	<0.001
<i>Female isoline (10 levels)</i>					0.40	0.64
<i>Heat:Female_isoline (random slopes)</i>					0.21	0.46


Copulation latency and copulation duration were log-transformed and then scaled and centred to aid model convergence.

## Supplementary figures:

**Fig S1:** Male heat-exposure affects multiple aspects of sexual behaviour. Courtship latencies (note the log-scale) of males that did not mate are shown as open circles. Bars illustrate approximate 95% confidence intervals, taken as twice the standard error calculated on the log-scale. Heat-exposed males (red) were less likely to court and mate, took longer to initiate courtship and to procure a mating, and mated for a shorter duration than control males (blue; see Tables 1 & S2).



**Fig S2:** Lower reproductive output after the first mating is associated with increased polyandry. Ticks represent individual females, initially mated to heat-exposed (red) or control males (blue). Individual females are represented by ticks. Circles illustrate average polyandry for females within ranges of similar reproductive output (shaded horizontal bars), with surface area proportional to sample size. Irrespective of reproductive output, polyandry was higher after mating with heat-exposed males (main effect of male heat-exposure). The interaction between reproductive output and male heat-exposure was not significant (see Table 3) but is retained here for illustrative purposes.



**Fig S3:** Heat-exposure decreased survival only in the male cohort A. Vials containing groups of up to 10 males were transferred into water baths set to 23°C (blue) or 31°C (red) one to two days (cohort A) or immediately (cohort B) after eclosion. Compared to survival (Table S2), mating performance was more similar for both male cohorts after heat-exposure (see Tables 1 & S1). Solid Bars illustrate approximate 95% confidence intervals and point surface area is proportional to the number of vials tested.

