Supplementary Table 1. Results from the general clinical & research Round 2 survey or experts (N = 23)

	HEALTHCARE PROFESSIONALS Mode, Median (IQR)	FAMILY CAREGIVERS and PAID CAREGIVERS Mode, Median (IQR)
Have specific techniques to apply in their work / use in daily life with patients with	Very Important, 3 (0)	Very Important, 3 (0)
MND who have cognitive impairment		
Have specific techniques to apply in their work / use in daily life with patients with	Very Important, 3 (0)	Very Important, 3 (0)
MND who have behaviour impairment		
Have specific techniques to apply in their work / use in daily life with patients with	Very Important, 3 (0)	Very Important, 3 (0)
MND who have concomitant FTD		
Have examples on how cognitive and behavioural impairment might impact the	Very Important, 3 (0)	
caregiver or families the person with MND		
Know about changes in day-to-day practice taking into account cognitive and	Very Important, 3 (1)	
pehaviour impairment in MND		
Know about changes in communication style with the person with MND who has	Very Important, 3 (0)	
cognitive and behaviour impairment		
Know about changes in communication style with the family or paid caregiver of	Very Important, 3 (0)	
he person with MND who has cognitive and behaviour impairment		
Have general information on cognitive and behavioural impairment in MND		Very Important, 3 (1)
Have examples of what cognitive and behavioural impairment look like and how it		Very Important, 3 (1)
mpacts people with MND		
Have a list of helpful health and social care contacts		Very Important, 3 (0)

IQR = interquartile range, MND = motor neurone disease, FTD =. Frontotemporal dementia. Not applicable fields in grey.