

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24

**Functional dynamics of a single tryptophan residue in a BLUF protein revealed by
fluorescence spectroscopy**

Kristof Karadi^{1,2}, Sofia M. Kapetanaki^{1,2}, Katalin Raics¹, Ildiko Pecs¹, Robert Kapronczai¹,
Zsuzsanna Fekete¹, James Iuliano³, Jinnette Tolentino Collado³, Agnieszka Gil³, Jozsef
Orban¹, Miklos Nyitrai^{1,2}, Greg M. Greetham⁴, Marten H. Vos⁵, Peter J. Tonge³, Stephen R.
Meech⁶, Andras Lukacs^{1,2}

¹ Department of Biophysics, Medical School, University of Pécs, 7624 Pécs, Hungary

² Szentagothai Research Center, University of Pecs, 7624 Pécs, Hungary

³ Department of Chemistry, Stony Brook University, Stony Brook, NY, 11794-3400, USA

⁴ Central Laser Facility, Harwell Science and Innovation Campus, Didcot, Oxfordshire, UK.

⁵ LOB, CNRS, INSERM, Ecole Polytechnique, Institut Polytechnique de Paris, 91128
Palaiseau Cedex, France

⁶ School of Chemistry, University of East Anglia, Norwich NR4 7TJ, UK

Correspondence and requests for materials should be addressed to A. L. (e-mail:
andras.lukacs@aok.pte.hu)


1 **Abstract**

2
3 Blue Light Using Flavin (BLUF) domains are increasingly being adopted for use in
4 optogenetic constructs. Despite this, much remains to be resolved on the mechanism of their
5 activation. The advent of unnatural amino acid mutagenesis opens up a new toolbox for the
6 study of protein structural dynamics. The tryptophan analogue, 7-aza-Trp (7AW) was
7 incorporated in the BLUF domain of the Activation of Photopigment and PUC A (AppA)
8 photoreceptor in order to investigate the functional dynamics of the crucial W104 residue
9 during photoactivation of the protein. The 7-aza modification to Trp makes selective
10 excitation possible using 310 nm excitation and 380 nm emission, separating the signals of
11 interest from other Trp and Tyr residues. We used Förster energy transfer (FRET) between
12 7AW and the flavin to estimate the distance between Trp and flavin in both the light- and
13 dark-adapted states in solution. Nanosecond fluorescence anisotropy decay and picosecond
14 fluorescence lifetime measurements for the flavin revealed a rather dynamic picture for the
15 tryptophan residue. In the dark-adapted state, the major population of the tryptophan 104 is
16 pointing away from the flavin and can move freely, in contrast to previous results reported in
17 the literature. Upon blue-light excitation, the tryptophan population is reorganized, the
18 dominant population moves closer to the flavin occupying a rigidly bound state participating
19 in the hydrogen-bond network around the flavin molecule.

20
21 **Introduction**

22
23 Flavins are found in more than 370 enzymes¹ but only a few of them are photoactive^{2,3}.
24 Three major families of photoreceptors which utilize flavin as a cofactor and whose functions
25 are triggered by absorption of light are the photolyase/cryptochromes, the light oxygen voltage
26 (LOV) domains and the blue light sensors using flavin (BLUF) proteins. Their photochemistry,
27 though is rather diverse. In photolyases and cryptochromes, FAD (flavin adenine dinucleotide)
28 is reduced via electron transfer through a tryptophan triad⁴⁻⁶. Photolyases use light to repair

1 UV-damaged DNA⁷ whereas the proposed functions of cryptochromes range from setting the
2 circadian clock in insects to sensing the weak magnetic field of Earth in migrating birds⁶. In the
3 LOV domains, the flavin cofactor is excited to a triplet state upon blue light absorption,
4 followed by formation of a signalling state, characterized by a covalent bond between the flavin
5 and a nearby cysteine residue, leading to the enhancement of the phototropin kinase activity³.
6 In BLUF domains, blue light excitation results in a signalling state (light-adapted state) that is
7 characterized by a reorganization of the hydrogen bond network around FAD and the Tyr-Gln-
8 Trp (Met) tetrad (Fig. 1). This is revealed by a characteristic 10-15 nm red-shift of the first
9 $\pi \rightarrow \pi^*$ transition and a 20 cm^{-1} downshift of the flavin $\text{C}_4=\text{O}_4$ stretching vibration compared to
10 the dark-adapted state^{8,9}. Site directed mutagenesis has shown that Y21 and Q63 play a crucial
11 role during photoactivation as the red shift upon illumination disappears if one of these residues
12 is replaced⁸⁻¹². A strong and specific chemical coupling between the flavin and the protein by
13 means of glutamine tautomerization has been proposed to provide a basis for light sensing¹³.
14



15 **Figure 1.** Ribbon diagrams of the BLUF domain crystal structures from WT (A, pdb:1yrx) and C20S (B, pdb:2iyg)
16 AppA_{BLUF} domains showing W104 in the vicinity of the flavin in the Trp_{in} conformation (A) and away from the
17 flavin in the Trp_{out} conformation (B). Important residues involved in an H-bond network around the flavin (Y21,
18 H44, N45, Q63) are shown. Dashed lines represent H-bonds between the flavin and the residues.
19

1 W104 is also a key player in the photocycle in communicating the electronic excitation
2 of the flavin ring to the protein backbone¹⁴. Recent time-resolved IR experiments¹⁵ have shown
3 that W104A AppA_{BLUF} undergoes reduced structural changes on the μ s timescale along with
4 an 80-fold increase in the rate of dark-state recovery, compared to that of the wild-type¹⁶. A
5 1.5-fold increase of the quantum yield of signalling state formation with a slight increase of the
6 ground state recovery have been observed for the W104F AppA_{BLUF} mutant¹⁷. Mutation of the
7 W104 residue (W104A) does suppress the red-shift in the full-length protein¹⁸ and renders the
8 protein insensitive to blue light *in vivo*¹⁶ underlying the importance of this residue for the
9 downstream signalling events.

10 The exact conformation of W104 during the photoactivation process in AppA and other
11 BLUF domains has been a controversial topic in the field. The first crystal structure of AppA¹⁹
12 (pdb:1yrx) showed that W104 in the dark-adapted state is located close to flavin in the so called
13 Trp_{in} conformation (Fig.1A), whereas subsequent crystal structures from Schlichting's group²⁰
14 (pdb: 2iyg, 2iyi) presented a different picture where the tryptophan is pointing away from the
15 flavin both in the dark- and light-adapted state (Trp_{out} conformation) (Fig.1B). Crystal
16 structures of other BLUF domains like PixD^{21,22}, BlrB²³ and OaPAC²⁴ have also contributed
17 to the confusion as tryptophan is located in a solvent exposed position in a Trp_{out} conformation.
18 Interestingly, Slr1694 exhibits both conformations with one of the 10 crystallographic subunits
19 adopting the Trp_{in} conformation and the others the Trp_{out} conformation²².

20 Computational calculations and spectroscopic studies have tried to address the question
21 of the tryptophan flip in AppA, leading to opposite conclusions. Two independent studies by
22 Udvarhelyi et al.²⁵ and Collette et al.²⁶ concluded that the Trp_{out} represents the dark-adapted
23 state of the protein whereas a recent study by Goyal and Hammes-Schiffer²⁷ revealed that both
24 the *in* and *out* conformations are allowed at equilibrium but the *in* conformation is more stable.
25 The solution NMR structure (pdb: 2bun) has also the majority of the snapshots in the Trp_{in}

1 conformation²⁸. UV resonance Raman studies have supported the Trp_{in} conformation in the
2 dark state²⁹. Tryptophan fluorescence spectroscopic studies have suggested that Trp does not
3 become fully solvent exposed as suggested by some crystal structures but were conflicting
4 regarding the conformation of the Trp^{18,30}.

5 As there is still a controversy about the conformation and the role of tryptophan W104
6 during photoactivation of AppA, we revisited the question using a different approach.
7 Specifically, we exploit the spectroscopic properties of the 7-aza Trp (7AW) unnatural amino
8 acid as an intrinsic fluorescent probe, able to separate the signals of interest from other Trp and
9 Tyr residues. The absorption and fluorescence maxima of 7AW are red-shifted compared to
10 canonical Trp allowing the exclusive excitation of the Trp analogue. We incorporated 7AW at
11 position 104 and applied a wide range of fluorescence techniques to probe the functional
12 dynamics of Trp104 in AppA_{BLUF}. The results of our study provide strong evidence that W104
13 adopts a Trp_{out} conformation in the dark-adapted state and a Trp_{in} conformation in the light-
14 adapted state.

15

16 **Results**

17

18 *FRET measurements*

19 To provide quantitative information on the position of Trp104 during the photoactivation
20 process in AppA, we used fluorescence resonance energy transfer (FRET) measurements.

21 Fluorescence resonance energy transfer, is a long-range non-radiative energy transfer process
22 that takes place between two fluorophores, one in the excited state and one in the ground
23 state^{31,32}. The FRET efficiency-which is defined as the fraction of energy absorbed by the donor
24 that is subsequently transferred to the acceptor depends on the distance between the donor and
25 the acceptor^{33,34} and can be calculated by equation (1)

26

$$E = \frac{1}{1 + \left(\frac{R}{R_0}\right)^6} (1)$$

1 where E is the FRET efficiency, R is the distance between the donor and the acceptor
2 fluorophores and R_0 is the distance between the two fluorophores in the case of 50% transfer
3 efficiency. This strong distance dependence of FRET efficiency makes possible the
4 measurement of molecular distances in the range of 10-100 Å and the term “spectroscopic ruler”
5 has been coined for the method³⁵.

6 FRET efficiency between W104 and FAD (Fig.2A) has been obtained by two methods:
7 measuring the fluorescence lifetime of the donor (7AW in position 104) in the absence and in
8 the presence of the acceptor (FAD) and by using the method of acceptor enhancement. In this
9 latter case the increase of the acceptor fluorescence intensity is monitored when the donor is
10 present: if the donor transfers energy to the acceptor, enhancement of the fluorescence intensity
11 of the acceptor is observed (see Supplementary Information).


12

13

14

15

1



2

3 **Figure 2.**

4 (A) Ribbon diagram of the crystal structure of C20S AppA_{BLUF} (pdb: 2iyg), showing the FRET pair (FAD, W104).
 5 (B) Emission spectra of W64F AppA_{BLUF} in the dark- (black) and light- (red) adapted states ($\lambda_{exc}=295\text{nm}$). (C)
 6 Chemical structures showing ionization of tyrosine to tyrosinate. (D) Emission spectrum of W64F/W104A
 7 AppA_{BLUF} ($\lambda_{exc}=295\text{nm}$). The narrow peak $\sim 330\text{ nm}$ is the contribution of Raman scattering of water (E) pH
 8 dependent formation of tyrosinates monitored at 295 nm for the W64F/W104A AppA_{BLUF} mutant and for L-
 9 tyrosine in solution (inset). Fitting of the sigmoidal curve with a Boltzmann equation reveals a $pK=8.0$ for the two
 10 tyrosine residues present in AppA_{BLUF} and a $pK=10.5$ for L-tyrosine in solution (inset).
 11

12 *The 7-aza-Trp analogue (7AW) enables FRET measurements between residue 104 and flavin*
 13 *in the presence of tyrosines*

1 Besides W104, AppA_{BLUF} contains one more tryptophan residue, W64. To eliminate a FRET
2 contribution arising from W64, we have used the W64F mutant in our studies. In addition, to
3 avoid fluorescence emission from the two tyrosine residues (Y21 and Y56) in AppA_{BLUF}, we
4 used 295 nm as the excitation wavelength, where tyrosines do not absorb (Fig. S1A). However,
5 the emission spectra of W64F (Fig. 2B) and of the tryptophanless mutant W64F/W104A (Fig.
6 2D) display a substantial fluorescence emission around 345 nm, suggesting the existence of a
7 fluorophore (other than a tryptophan residue) and hence the presence of another component that
8 may contribute to FRET in addition to W104. The 345 nm emission component is shown below
9 to arise from tyrosinates.

10 Tyrosine residues are well known to deprotonate at high pH to form tyrosinates (Fig. 2C) which
11 have very similar fluorescent features to tryptophan. In particular, an increase of the pH results
12 in an increase of the absorption at 295 nm with concomitant decrease at 270 nm and a
13 fluorescence emission shift from 303 nm to 340 nm^{36,37}. This generally occurs at pH>10.5
14 which is the pK_a of free tyrosine³⁸. However, as it has been reported for a series of proteins^{39–}
15 ⁴² tyrosine residues can have lower pK_a values which results in the formation of tyrosinates and
16 hence fluorescence emission appears around 340 nm, even at pH<10.5. For example, tyrosinate
17 fluorescence was observed from Y99 in calmodulin from bovine testes and the pK_a of this
18 tyrosine residue was shown to be as low as 7.0⁴¹. Tyrosinate fluorescence was also observed in
19 photoactive yellow protein (PYP) where the pK_a changes substantially, depending on whether
20 the tyrosine is solvent exposed, buried or hydrogen bonded,⁴³ and in chicken ovomucoid where
21 the pK_a changes upon pressure dependent unfolding resulting in an increase of the fluorescence
22 emission at 340 nm³⁷.

23 To test whether tyrosinates are also formed in the AppA W64F/W104A mutant at pH=8.0 we
24 performed pH dependent measurements of the absorbance at 295 nm and determined a pK_a
25 value of 8.0 (Fig.2e). This finding suggests that at pH = 8.0, half of the tyrosine residues exists

1 as tyrosinates giving rise to the strong fluorescence emission at ~ 345 nm. This will necessarily
2 result in a distorted picture for any fluorescence spectroscopy measurement where tryptophan
3 fluorescence is monitored. To eliminate any contribution from tyrosinates, we used 7AW
4 (Fig.3A) and exploited its spectroscopic properties.


5 7AW (Fig. 3A) was first incorporated in bacterial proteins 50 years ago⁴⁴. It is an ideal
6 non-invasive *in situ* probe of the structure and dynamics of proteins, as it introduces minimal
7 structural and functional modifications to the protein and in addition has favourable spectral
8 properties compared to canonical Trp and other Trp analogues^{45,46}. In particular, the
9 absorption and fluorescence maxima of 7AW are red-shifted by 10 nm and 50 nm,
10 respectively compared to that of canonical Trp (Fig.S1A, Fig.S1B. and⁴⁵). This red shift in
11 the absorbance allows us to excite selectively the tryptophan analogue 7AW, using $\lambda_{exc} =$
12 310nm (canonical Trp has no absorbance at 310 nm) and to avoid the tyrosinates' absorbance
13 at 295 nm (see previous paragraph).

14

15

16

17



1

2 **Figure 3.** A) Chemical structures showing 7AW (left) and canonical tryptophan (right). B) Emission spectra of
 3 W64F (blue line) and 7aza-W104/W64F AppA_{BLUF} mutant (red line) ($\lambda_{exc}=310$ nm). W64F does not show
 4 fluorescence emission with 310 nm excitation; around 350 nm the spectrum shows remains of the Raman peak
 5 (not shown on this scale). C) Absorption spectra of W64F (black line) and 7aza-W104/W64F AppA_{BLUF}
 6. Spectra are vertically translated for visualization purposes. D) Absorption spectra of 7aza-W104/W64F AppA_{BLUF}
 7 in the dark- (black line) and light-adapted (red line) states showing the characteristic red-shift. The inset shows the
 8 recovery kinetics of the dark-adapted state monitored at 490 nm after illumination at 385 nm. E) Fluorescence

1 decay of the 7aza-W104/W64F AppA_{BLUF} in the dark (green) - and light-adapted (red) states after excitation at
2 321nm, ($\lambda_{em}=380nm$) in the presence of FAD and of the 7aza-W104/W64F AppA_{BLUF} in the deflavinated, donor-
3 only (black) state after excitation at 321nm ($\lambda_{em}=380nm$). The instrument response function is shown in blue. F)
4 Emission spectra of 7aza-W104/W64F AppA_{BLUF} in the dark-adapted state G) Emission spectra of 7aza-
5 W104/W64F AppA_{BLUF} in the light-adapted state. The acceptor only spectra (blue) are calculated as explained in
6 the Supplementary Information. Excitation was set at 310 nm. A significant fluorescence enhancement in the light-
7 adapted state (red) is observed for 7aza-W104/W64F AppA_{BLUF} due to FRET transfer from 7AW to the flavin.
8 The fluorescent enhancement is lesser in the dark-adapted state (F).
9

10 Using tryptophan auxotroph cell lines (as described in the Methods section), we incorporated
11 the 7AW analogue in the place of W104 in the W64F mutant of AppA_{BLUF}. The successful
12 incorporation of the analogue in W64F AppA_{BLUF} was confirmed by the characteristic
13 emission band at 380 nm ($\lambda_{exc}=310$ nm) which is absent in the W64F mutant (Fig. 3B). The
14 structural modification to the protein, due to the more hydrophilic character of the side chain
15 of 7AW and potential new contacts by means of hydrogen bonds compared to canonical Trp
16 is minimal, as reflected in (1) the flavin absorption spectrum which is similar to that of W64F
17 (Fig. 3C) (2) the characteristic red shift of the isoalloxazine absorption maximum in the light-
18 adapted state is similar to the corresponding shift observed in the W64F mutant (Fig. 3D and
19 Fig.S2A) and (3) the identical dark state recovery rate (~14 min) compared to that of W64F
20 (Fig.3D inset.) **The hydrogen bonding environment due to the extra N- atom of the 7aza-
21 W104/W64F AppABLUF mutant is expected to be different between the wild-type and the
22 7aza analogue, resulting in an electronically modified structure of the flavin which is reflected
23 as a small shift of the maximum and change in its vibronic structure (Figure 3D). In the light-
24 adapted state, Trp104 points towards the flavin. As explained above, the extra N atom of the
25 indole ring of the tryptophan results in a different hydrogen-bonding environment compared
26 to the wild type, resulting to different atom displacements in the excited state and hence to
27 different shifts of the maximum for the S₀ to S₁ transition.**
28

29 *FRET: Fluorescence lifetime measurements of 7aza-W104/W64F AppA_{BLUF}*
30

1 To determine the FRET efficiency (E) between 7AW and FAD we performed time correlated
2 single photon counting (TCSPC) measurements on the tryptophan in the case of the apoprotein
3 and on the protein in the dark-adapted and light-adapted states (Fig. 3E). FRET efficiency was
4 calculated using equation:

$$5 \quad E = 1 - \frac{\tau_{DA}}{\tau_D} \quad (2),$$

6 where τ_{DA} is the fluorescence lifetime of the donor (7AW) in the presence of the acceptor
7 (FAD) and τ_D is the fluorescence lifetime of the donor alone. In all three cases the decay of the
8 excited tryptophan analogue is described best using three lifetime components (Table 1) as
9 previously reported for the canonical tryptophan. Two of the fluorescence lifetimes of canonical
10 tryptophan (0.2-0.9 ns and 1.2-3.6 ns) are inherent to the tryptophan structure itself independent
11 of the surrounding environment whereas the third lifetime (3.7-9.2 ns) is generated by
12 interactions between the tryptophan and neighbouring residues⁴⁷. In the case of 7AW, the most
13 striking difference compared to the canonical tryptophan is that the third lifetime component is
14 longer (ca. 14-16 ns), whereas the other two components are observed in the same range as for
15 tryptophan. The lifetime of the tryptophan analogue is shorter when FAD is present, suggesting
16 the existence of Förster-type energy transfer. Calculating the FRET efficiency – using the
17 average lifetimes – we obtained 23% efficiency in the dark-adapted case and 40 % in the light-
18 adapted case (see Supplementary information). An estimate of the average distance between
19 the tryptophan analogue and the flavin can be obtained by

$$20 \quad R = R_0 \sqrt[6]{\frac{1-E}{E}} \quad (3).$$

21 In the case of 7AW and FAD, R_0 is 16.8 Å using the assumption that the tryptophan can freely
22 rotate thus the value of the orientation factor (see SI) $\kappa^2 = 2/3$. In support of free rotation of the
23 tryptophan ($\kappa^2=2/3$) the anisotropy decay of the dark-adapted state (see next section) is
24 substantially faster when compared to the light-adapted state. Hence in the dark-adapted state,
25 the distance between 7AW and the flavin is estimated to be 20.5 Å (see Supplementary

1 Information). The slower rotational correlation time observed in the case of the light-adapted
 2 state of the protein (see below) suggests that the tryptophan is not able to freely rotate as it is
 3 hydrogen-bonded in the Trp_{in} position. This locked position imposes a strong constraint on the
 4 orientation of the transition dipole of the tryptophan, implying that $\kappa^2=2/3$ used for the freely
 5 rotating dipoles cannot be used. From the crystal structure (pdb:1yrx) we can observe that the
 6 orientation of the transition dipole moments of the flavin and the tryptophan 104 are close to
 7 perpendicular (see Fig. S6) thus κ^2 is expected to be close to zero, the theoretical value for
 8 perpendicular orientation; we have actually calculated $\kappa^2=0.02$ (see Supplementary
 9 Information). In this case, the value of R_0 is modified to 8.9 Å, and the distance calculated
 10 between tryptophan and flavin in the light-adapted state is 9.5 Å.

11

12 **Table 1.** Fluorescence lifetimes (ns) of 7AW and corresponding amplitudes in the 7aza-
 13 W104/W64F mutant (no FAD present), and the 7aza-W104/W64Fmutant (FAD present) in the
 14 dark- and light-adapted states.

7azaTrp- W64F/W104	τ_1	α_1	τ_2	α_2	τ_3	α_3	τ_{average}	reference
apoprotein	0.4	0.2	4.1	0.3	16.2	0.4	8.7	this work
holoprotein dark	1.1	0.4	4.4	0.3	16	0.3	6.8	this work
holoprotein light	1.3	0.4	3.5	0.3	14.7	0.2	5.2	this work
canonical Trp	0.2-0.9		1.2-3.6		3.7-9.2			⁴⁷

15

16 *FRET: acceptor enhancement method*

17 FRET efficiency between 7AW104 and the flavin can also be calculated by measuring the
 18 intensity loss of the emission of the tryptophan or the intensity enhancement of the emission of
 19 the flavin due to the resonance energy transfer. In the latter case, known as the acceptor
 20 enhancement approach^{31,32}, the FRET efficiency is calculated by equation (4)

1

$$2 \quad E = \frac{\varepsilon_A(\lambda_D^{ex})}{\varepsilon_D(\lambda_D^{ex})} \left[\frac{I_{AD}(\lambda_A^{em})}{I_A(\lambda_A^{em})} - 1 \right] \quad (4)$$

3 where $\varepsilon_A(\lambda_D^{ex})$ and $\varepsilon_D(\lambda_D^{ex})$ are the extinction coefficients of the acceptor and donor at the
4 donor excitation wavelength (λ_D^{ex}), and they were calculated using the absorption spectrum of
5 FAD (Fig. S4). $I_{AD}(\lambda_A^{em})$ and $I_A(\lambda_A^{em})$ are the acceptor intensity in the presence and the
6 absence of the donor, respectively.

7 The enhancement of the fluorescence intensity due to the FRET between tryptophan
8 and flavin was measured in the case of the 7aza-W104/W64F mutant in order to avoid the
9 fluorescence from the tyrosinates as previously discussed.

10 Figures 3F and 3G show the emission spectra of the 7aza-W104/W64F mutant in the dark and
11 light states, respectively using $\lambda_{exc}=310\text{nm}$. Inspection of the data reveals a significant
12 increase of the emission of FAD in the presence of the donor both in the dark-adapted and
13 light-adapted states indicating efficient FRET from 7AW to FAD. This fluorescence
14 enhancement is even more pronounced in the light-adapted state. We calculate a FRET
15 efficiency of $E_{\text{dark}}=42\%$ and $E_{\text{light}}=59\%$, which corresponds to a distance $R_{\text{dark}}=17.7 \text{ \AA}$ and
16 $R_{\text{light}}=8.3 \text{ \AA}$ for the dark- and light-adapted states, respectively^{8,48}.

17

18

19

20

21 *Comparison of the FRET obtained distances with those from crystal structures*

22

23 The estimated distances between Trp 104 and FAD obtained from lifetime
24 measurements and the acceptor enhancement method are in relatively good agreement for the
25 dark-adapted state (20.5 \AA and 17.7 \AA , respectively).

1 The estimated distances from our fluorescence measurements are also in good
2 agreement with the distances obtained from the crystal structures of AppA_{BLUF} (pdb: 2iyg,
3 2iyi) and AppA full-length without the Cys-rich region (pdb:4hh0, 4hh1) (Table 2) where
4 W104 has been reported to be located on the surface/distal to FAD (Trp_{out} conformation).
5 However, they significantly deviate from those in the first solved crystal structure of
6 AppA_{BLUF} (pdb:1yrx, 5.9 Å) and the NMR solution structure (pdb:2bun, 6.2Å) for which
7 W104 exhibits the Trp_{in} conformation (close to FAD). It should be noted that local differences
8 between the AppA1-124C20S dark structure (pdb:2iyg) and the AppA17-133 (pdb:1yrx) have
9 been earlier attributed to the presence of bound detergent molecules²⁰, and hence considered
10 for the structure of AppA1-124C20S to represent better AppA_{BLUF} in solution. The
11 significantly (9.4-11 Å) shorter distance between W104 and FAD for the light-adapted state
12 from FRET and fluorescence lifetime measurements suggest that in the light-adapted state
13 W104 adopts a Trp_{in} conformation. It should be mentioned that such a difference was not
14 observed between the dark-adapted and light-adapted states of the crystal structures of
15 AppA1-124C20S²⁰ (Table 1) potentially originating from constraints in the movement of the
16 indole ring of W104 inside the crystal. This is also reflected in the smaller light-induced red
17 shift in the crystal (5-6 nm) compared to that in the solution (14 nm) suggesting the formation
18 of a preceding intermediate of the full signalling light-adapted state²⁰. Interestingly, no
19 significant conformational changes have been observed in the corresponding tryptophan
20 residue (W90) in the photoactivated state of the OaPAC BLUF protein despite the ~10 nm
21 light-induced red-shift observed in the crystal⁴⁹.

22
23 **Table 2.** Comparison of the distance between C7 of the indole ring of Trp (W104 in
24 AppA_{BLUF}) and N3 of the isoalloxazine ring of flavin from the various crystal structures
25 available for AppA_{BLUF}, AppA full-length domains and other BLUF domains. An average
26 distance is provided for those structures that contain more than one subunit. *In Slr1694_{BLUF}

1 (2-140), a shorter distance (5.7 Å) is observed in one of the ten subunits. FRET: FL refers to
 2 our FRET measurements using fluorescent lifetimes and FRET: AEM to FRET measurements
 3 using the acceptor enhancement method.

4
 5

Protein	pdb/FRET	Distance (Å)	references
AppA _{BLUF} (17-133)	1yrx	5.9	50
AppA _{BLUF} (5-125)	2bun	6.2	28
AppA _{BLUF} C20S (1-124)	2iyg (dark)	16.0	20
AppA _{BLUF} C20S (1-124)	2iyi (light)	16.2	20
AppA C20S Δ399	4hh0	15.7	50
wt AppA Δ399	4hh1	15.7	50
TePixD _{BLUF} (2-143)	1x0p	16.1	21
Slr1694 _{BLUF} (2-140)	2hfn	15.7, 5.7*	22
OAPAC (1-366)	4yus	16.6	24
BrlB _{BLUF} (1-140)	2byc	16.0	23
7azaW104W64F (dark)	FRET:FL	20.5	this work
7azaW104W64F (light)	FRET:FL	9.5	this work
7azaW104W64F (dark)	FRET: AEM	17.7	this work
7azaW104W64F (light)	FRET: AEM	8.3	this work

6

7 *7-aza W104 has restricted movement in the light-adapted state as revealed by fluorescence*
 8 *anisotropy decay measurements.*


9 To probe the conformational dynamics of W104 and steric restrictions, we have applied time-
 10 resolved fluorescence anisotropy decay measurements, which have been used for decades to

1 characterize protein dynamics. The relaxation of anisotropy induced by a polarized excitation
 2 pulse reflects both the dynamics of the fluorophore itself and the protein or segment to which
 3 it is attached to. Time dependent anisotropy $r(t)$ can be fitted as a sum of exponentials³²,

$$4 \quad r(t) = \sum_i r_{0i} \exp(-\frac{t}{\theta_i}) \quad (5),$$

5 where θ_i values are the rotational correlation times, and r_{0i} are the limiting anisotropies in the
 6 absence of rotational diffusion. In principle, fast rotational correlation times (~ 1 ns) suggest a
 7 fast movement of the fluorophore whereas longer correlation times (tens of nanoseconds) may
 8 reflect a slower rotation of the fluorophore or the entire protein complex³².

9



10

11 **Figure 4.** Decay of the fluorescence anisotropy of 7aza-W104/W64F AppA_{BLUF} in the dark-(A) and light-
 12 adapted states (B). C) Fluorescence decay of the Y21F/Y56F/W64F/W104F AppA_{BLUF} (blue) and free FMN (red)
 13 after excitation at 455 nm, ($\lambda_{em}=510$ nm). The instrument response function (~ 1 ns) is shown in black. D) Kinetics
 14 at the emission maximum ($\lambda_{em}=520$ nm) after excitation at 390 nm for AppA W64F dark-adapted state (navy), light-
 15 adapted state (red) and Y21F/Y56F/W64F AppA BLUF (green). The instrument response function was ~ 1 ps.

16

1 Figure 4A shows the decay of the fluorescence anisotropy of the 7aza-W104/W64F AppA_{BLUF}
2 mutant in the dark- and light- adapted states. Single exponential fitting reveals a decay constant
3 for the dark-adapted state, $\tau_{dark}=1.5 \pm 0.06$ ns and for the light-adapted state, $\tau_{light}=11.1 \pm 0.5$
4 ns. The short rotational correlation time in the dark-adapted state reflects a less restricted
5 rotation of the tryptophan residue, and it is close to the rotational correlation time of L-
6 tryptophan in buffer ($\theta= \sim 0.6$ ns) implying an almost free rotation of the indole moiety.
7 The observed 11 ns long rotational correlation time in the light adapted state corresponds to the
8 rotation of the whole protein and agrees well with the value expected based on the empirical
9 formula proposed by Visser⁵¹: $\Phi = 3.84 \cdot 10^{-4} M_r$, where Φ is the rotational correlation time in
10 ns and M_r is the molecular weight of the protein in Da. This observation suggests that upon blue
11 light absorption, the W104 tryptophan residue moves closer to the flavin and gets hydrogen
12 bonded to an adjacent amino acid residue, most probably Q63.
13 Fluorescence anisotropy decay measurements were performed in the Y21F/Y56F/W64F AppA
14 mutant as well, in which case the protein is locked in a state which is very similar to the light
15 adapted. The rotational correlation time was calculated to be ~ 10 ns (see Figure S5), hence
16 very similar to the one observed in the light-adapted state of the 7aza-W104/W64F mutant.
17 That finding suggests that the tryptophan at 104 position is rigidly bound to the protein, in the
18 triple mutant.
19 These results are in line with the findings from our FRET measurements: W104 in AppA_{BLUF}
20 is characterized by flexible dynamics in the dark- adapted state whereas in the light-adapted
21 state, it is present in a more restricted environment, probably due to a stronger hydrogen bond
22 network around the flavin molecule, originating from the nitrogen atoms of the tryptophan that
23 are expected to form additional hydrogen bonds with Q63 and O14 of the flavin. It should be
24 pointed out that the flexibility of W104 in the dark-adapted state is a novel finding that hasn't

1 been observed in previous studies which have favoured a position of the W104 close to the
2 flavin in a Trp_{in} conformation^{18,19,29}.

3

4 *Fluorescence lifetimes of the flavin in the dark- and light-adapted state*

5 Ultrafast electron transfer from the neighbouring aromatic amino acid residues (Trp and Tyr)
6 to the electronically excited flavin chromophore has been observed in many flavoproteins
7 (including glucose oxidase, flavodoxin, riboflavin binding protein, photolyase/cryptochromes
8 and ThyX)⁵²⁻⁵⁶. This photoinduced electron transfer (PET) results in substantial shortening of
9 the fluorescence lifetime of the flavin. Measurements of the fluorescence lifetime of the flavin
10 can therefore provide significant information on the flavin environment. Here, we applied
11 TCSPC and Kerr-gated fluorescence spectroscopy⁵⁷ to determine the fluorescence lifetimes of
12 the flavin in the dark- and light-adapted state in order to obtain further information on the
13 position of W104 during photoactivation, and its influence on the quenching mechanism.

14 As expected, in the AppA_{BLUF} mutant where all four tyrosine and tryptophan residues have been
15 mutated (Y21F/Y56F/W64F/W104F) and hence all possible electron donors are removed, the
16 fluorescence lifetime of the flavin was 3.8 ns (Fig.4B). This is in the range expected for the
17 fluorescence lifetime of the flavin in solution ~3-5 ns: the fluorescence lifetime (the longer
18 component) of FAD is around 3 ns⁵⁸, free FMN in water is around 4.8 ns, lumiflavin is 5.2
19 ns^{59,60}. As expected from the lower quantum yield of FAD fluorescence in AppA W64F (8.5%
20 and 4.8% in dark- and light-adapted state respectively) a substantially faster fluorescence decay
21 with respect to that of the quadruple mutant was observed both in the dark-and light-adapted
22 case (Fig.4C and Table 3).

23 Transient picosecond fluorescence spectra (Fig. S3A-C) and kinetics at the emission maximum
24 ($\lambda_{\text{max}}=520$ nm) (Fig. 4D) are shown for AppA W64F (dark- and light-adapted states) and AppA
25 Y21F/Y56F/W64F. Global analysis reveals a biphasic fluorescence decay for the flavin. (Table

1 3). In the dark-adapted state of the 7aza-W104/W64F, there is a dominant phase with a decay
2 constant $\tau_{dark}=500$ ps that is around fifty times slower than the dominant phase of the light-
3 adapted state, $\tau_{light}=11$ ps. The faster decay in the light-adapted state, suggests that the excited
4 state of the flavin is quenched by electron transfer from close-by aromatic residues that should
5 be positioned closer to the flavin compared to their position in the dark-adapted state.

6 We measured the fluorescence lifetime of the flavin in the Y21F/Y56F/W64F AppA mutant
7 where all the potential electron donors were removed except for W104. In this case we also
8 observed a significantly (~ 4-5 times) shortened relaxation time compared to the time constant
9 measured in the dark-adapted state of AppA W64F. This indicates that the tryptophan in this
10 mutant is closer to the flavin than in dark-adapted state of AppA W64F and it is able to quench
11 efficiently the excited state of the flavin. This result lines up well with the findings from the
12 anisotropy decay measurements altogether indicating that W104 is in a rigidly bound position
13 close to the flavin.

14 We therefore assign the drastic acceleration of the fluorescence decay in the light-adapted state
15 of AppA W64F to the movement of the W104 towards the flavin and its acting as the main
16 quencher in the light-adapted state (see Discussion). **Despite the fact that in PixD the respective**
17 **tyrosine (Y8) is involved in proton coupled electron transfer (PCET) we don't anticipate that**
18 **Y21 is an efficient quencher in the case of AppA_{BLUF}. We have already demonstrated⁹ the lack**
19 **of an efficient ET during the forward reaction in AppA_{BLUF} which we attribute to the low pKa**
20 **of Tyr21. Our previous work⁷³ on PixD where we replaced Tyr8 with fluorinated tyrosines,**
21 **lowering the pKa of the tyrosine (by fluorination) led to an increase of the lifetime of the excited**
22 **state. The same effect was observed in another BLUF protein (unpublished results) implying**
23 **that the ability of the respective tyrosine to quench the flavin decreases with low pKa. The**
24 **possible reason is that despite the higher driving force for ET, tyrosinate formation prevents the**
25 **PCET which is clearly needed in PixD.**

1 In addition, the emission spectra of the W64F AppA_{BLUF} (Figs. 2B and S3B) lose the
2 characteristic vibrational shoulder at ~500 nm in the light-adapted state. These changes are also
3 accompanied by a decrease in the fluorescence (Fig. 2B) and are indicative of an enhanced
4 electron transfer in the case of the light-adapted state.

5 Using the simple empirical expression known as the Dutton ruler (equation 6)⁶¹

$$6 \log k_{ET} = 15 - 0.6R - 3.1 \frac{(\Delta G + \lambda)^2}{\lambda} \quad (6)$$

7 that relates the electron transfer rate constant (k_{ET}) to the edge-to-edge distance (R in Å) between
8 two reaction centres, we roughly estimated the distances between flavin and the tryptophan for
9 the dark-adapted state ($R_{\text{dark}}=9.5$ Å) and the light-adapted state ($R_{\text{light}}=6.7$ Å) (Supplementary
10 information), assuming a barrierless electron transfer ($\Delta G = -\lambda$), where ΔG is the driving force
11 for the electron transfer between the electron donor (presumably W104 at least for the light-
12 adapted state) and the excited flavin and λ is the reorganization energy.

13 The triple mutant (Y21F/Y56F/W64F) where both tyrosines and W64 tryptophan were removed
14 is photoinactive, no red shift is observed upon blue light illumination. **The absorption spectra**
15 **of this mutant and the Y21F/Y56F/W64F/W104F mutant can be found in the Supplementary**
16 **Information (Fig S7). Both mutants are very stable even at high concentrations (300-500 μM)**
17 **and able to retain the flavin despite the multiple mutations. In the case of the Y21F/Y56F/W64F**
18 **mutant it seems that W104 is locked in a position that is characterized by similar photophysics**
19 **with the light-adapted state of WT AppABLUF as the fluorescence decay of the flavin is very**
20 **fast.** Interestingly, the two decay phases of the flavin have similar contributions indicating that
21 W104 exists in two conformations relative to its proximity to the flavin chromophore. Based
22 on the Dutton ruler, the following estimated distances were calculated, $R_{in}=7.6$ Å ($\tau=36$ ps) and
23 $R_{out}=8.8$ Å ($\tau=200$ ps).

24

1 **Table 3:** Fluorescence decay constants of the flavin and corresponding amplitudes in various
 2 AppA_{BLUF} mutants. Estimated distances between the flavin and W104 based on electron
 3 transfer calculations. The major contribution is shown in bold.

Protein	A1	T1(ps)	A2	T2(ps)	R1 (Å)	R2 (Å)
W64F dark	0.7	500	0.3	70	9.5	8.03
W64F light	0.13	230	0.87	11	8.9	6.7
Y21F/Y56F/W64F	0.43	200	0.57	36	8.8	7.6


4
5

6 *Transient infrared measurements*

7

8 To study the protein dynamics in the AppA_{BLUF} domain after blue light absorption we
 9 performed transient infrared measurements on the dark- and light-adapted states of AppA_{BLUF}
 10 W64F and on the AppA_{BLUF} Y21F/Y56F/W64F mutants (Fig. 5A). Transient infrared spectra
 11 recorded at time delays between 1ps and 2 ns after 450 nm excitation of the flavin show
 12 similarities with the spectra observed for the wild type AppA_{BLUF}^{62,63}. The negative features
 13 (bleaches) of the recorded spectra associate with depletion of the ground state population or
 14 with photoinduced changes in the vibrational spectrum of the protein, occurring either directly
 15 through electronic excitation at $t = 0$ or as a result of subsequent structural dynamics; positive
 16 bands associated with vibrations of the electronically excited state of the flavin, or with modes
 17 of the protein which shift as a result of electronic excitation, or of products formed
 18 subsequently. In AppA_{BLUF} (and the majority of flavoproteins) four dominant bleaches are
 19 observed in the 1300-1800 cm⁻¹ region: the bands at 1547 cm⁻¹ and 1581 cm⁻¹ arise from ring
 20 modes of the isoalloxazine ring whereas the ~ 1650 cm⁻¹ and ~ 1710 cm⁻¹ bands are related to
 21 the carbonyl stretches C2=O and C4=O of the FAD ground state, respectively, and are
 22 sensitive to the H-bond environment^{62,64}.

23



1
 2 **Figure 5.** A) TRIR spectra of dark- and light-adapted (blue and red respectively) W64F and the Y21F/Y56F/W64F
 3 AppA mutants (green) recorded at 10 ps; B) Kinetics of the excited state of the flavin in the dark- and light-adapted
 4 (blue and red respectively) states of W64F AppA_{BLUF} and in the Y21F/Y56F/W64F AppA_{BLUF} (green) observed
 5 at 1380 cm⁻¹.
 6

7 Comparison of the dark- and light adapted spectra of the W64F AppA mutant reveals a ~ 9 cm⁻¹
 8 ¹ downshift for the C4=O carbonyl band in the light adapted state, that is similar to the shift
 9 observed for the WT AppA⁶³ and attributed to the formation of new hydrogen bond to C4=O.
 10 That result demonstrates the arrangement of a rigid environment for W104 in the light-adapted
 11 state due to the formation of a H-bond to the flavin, and supports our FRET measurements that
 12 reveal a change of the location of W104 close to the flavin in that state. It is also in line with
 13 our anisotropy decay measurements that demonstrate a slower rotation (~11 ns) for W104 in
 14 the light-adapted state, reflecting rotation of W104 together with the whole protein due to the
 15 more restricted environment of W104. Similarly, the vibrational mode related to the C4=O
 16 carbonyl group is downshifted in the Y21F/Y56F/W64F AppA_{BLUF} compared to the dark-
 17 adapted W64F AppA_{BLUF} suggesting a bond strengthening. Figure 5B shows the kinetics
 18 observed at 1380 cm⁻¹ which reflect the decay of the excited state of the flavin⁹. The kinetics in
 19 Table 4 are in line with our findings from the picosecond fluorescence lifetime measurements
 20 for the flavin: the decay of the excited state is slower in the dark-adapted state and faster in the
 21 light-adapted state reflecting the efficient quenching of the excited state in the latter case. The
 22 kinetics for the Y21F/Y56F/W64F reflect also a significantly faster relaxation compared to the

1 AppA_{BLUF} W64F dark-adapted state which allows us to conclude that W104 (the only possible
2 electron donor in this case) is in a rigidly bound and closer position to the flavin compared to
3 the dark-adapted state.

4 **Table 4.** Decay time constants at 1380 cm⁻¹

5

6

	τ_1 (ps)	α_1	τ_2 (ps)	α_2
Dark W64F	41	0.27	482	0.73
Light W64F	23	0.86	221	0.14
Y21F/Y56F/W64F	23	0.36	160	0.64

7

8

9 Discussion

10 W104, a residue present in the majority of BLUF containing photoreceptors [AppA, Slr1694,
11 OaPAC, TePxD, BlrB]⁴⁹ is of special interest because of the inconsistencies reported
12 concerning its conformation in the dark- adapted states, based on crystal structures and its
13 potential role in the long-range signalling. It is well known that X-ray crystal structures of
14 proteins capture only those conformations that fit into the crystal lattice, which may or may not
15 be relevant to function.

16 Earlier spectroscopic studies^{18,29} have favoured the Trp_{in} conformation for the dark-adapted
17 state based on indirect evidence for a buried and hydrophobic environment around the
18 tryptophan. A fluorescence spectroscopy approach was used by Dragnea et al.¹⁸ where they
19 used acrylamide quenching to characterize the dynamics of the respective tryptophan.
20 Acrylamide quenching was proved to be a useful method to obtain information about the
21 tryptophan accessibility. The quenching of the fluorescence of a buried tryptophan is less
22 efficient than the quenching of a solvent exposed one. In the previous experiments though, the

1 results of acrylamide quenching were influenced by the unexpected appearance of tyrosinate
2 fluorescence, revealed here. Incorporating the unnatural amino acid 7AW we were able to avoid
3 the fluorescence of tyrosinate, in this way our work provides unequivocal evidence, as it relies
4 on a novel approach and the application of advanced fluorescence techniques to address the
5 orientation of Trp104 in the BLUF functional states.

6 Fluorescence spectroscopy methods – namely fluorescence anisotropy decay and Förster-type
7 energy transfer– were used to both characterize the mobility and position of W104 tryptophan
8 during photoactivation, and its distance from flavin in dark- and light- adapted states. In the
9 case of the FRET experiments we monitored the changes of the fluorescence intensity of the
10 acceptor as there was no possibility to measure the FRET efficiencies from the donor side;
11 removing the flavin would have led to a photoinactive protein. In order to measure the FRET
12 efficiency using the acceptor enhancement method we first made the W64F/W104A mutant
13 which was anticipated to have a fluorescence intensity arising only from the acceptor following
14 excitation at 295 nm. However, an unexpectedly strong fluorescence signal around 340 nm was
15 observed, despite the lack of the tryptophans, pointing to substantial fluorescence from the
16 tyrosinate form of tyrosines in AppA_{BLUF}. In order to eliminate this fluorescence emission,
17 which overlaps with the fluorescence coming from the tryptophans, we used the 7AW
18 tryptophan analogue which has shifted absorption and fluorescence spectra compared to the
19 canonical one.

20 Substitution of W104 with the unnatural amino acid residue 7AW resulted in a tailor made
21 BLUF protein with useful spectral properties that allowed us to obtain clear and distinct
22 fluorescent signals, making possible measurement of the fluorescent anisotropy decay of 7AW
23 in the 104 position of AppA_{BLUF} in the dark- and light-adapted state of the protein. The outcome
24 of the fluorescence anisotropy decay measurements resulted in an unexpected picture compared
25 to our presumption: the fast anisotropy decay (~ 1 ns) of 7AW in the dark- adapted state reflects

1 the nearly free rotation of the tryptophan suggesting that it is therefore not hydrogen bonded to
2 the flavin and the adjacent glutamine (Q63). In contrast a slow rotation (~ 11 ns) was observed
3 in the light- adapted state, which shows that the tryptophan is rotating together with the protein,
4 most probably because it is tightly hydrogen bonded to Q63.

5 The substitution of tryptophan W104 by the tryptophan analogue 7AW paved the way for the
6 FRET experiments using the “spectroscopic ruler” to determine the tryptophan-flavin distance
7 in both the dark- and light-adapted states. The estimated distance obtained from the acceptor
8 enhancement method for the dark-adapted state (17.7 \AA) is in good agreement with the distance
9 obtained from the crystal structures of the AppA_{BLUF} (16 \AA) and full-length protein (15.7 \AA)
10 (Table 2). A slightly longer distance (20.5 \AA) was obtained from the fluorescence lifetime
11 measurements probably due to the uncertainty in the estimation of the fluorescence lifetime of
12 7AW in position 104 in the apoprotein. In the absence of the flavin, the protein may adopt a
13 different conformation compared to when flavin is present which may result in a modified
14 fluorescence lifetime as the tryptophan is very sensitive to the protein environment. The FRET
15 measurements for the dark-adapted state point towards a more distant position for the
16 tryptophan compared to the light-adapted state in which the tryptophan is very close ($8.3/9.5$
17 \AA , Table 2) to FAD. The estimates of $8.3/9.5 \text{ \AA}$ center-to-center between FAD and 7AW-
18 Trp104 from the FRET experiments are in good agreement with the estimate of 6.7 \AA (edge-to-
19 edge) between FAD and W104 from the electron transfer quenching experiments. This
20 assessment also confirms that W104 acts as the predominant electron donor to FAD* (excited
21 FAD) in the light-adapted state.

22 The change in the orientation and the distance of W104 during the photoactivation was
23 reflected in earlier fluorescence and transient absorption spectroscopy measurements. Streak
24 camera measurements on the dark-adapted AppA_{BLUF} have shown that the fluorescence lifetime
25 of flavin is around 600 ps^{10} in line with our measurements (Table 3). This value drops by a

1 factor of ten (Table 3) in the case of the light-adapted state of the protein suggesting that flavin
2 fluorescence is strongly quenched most probably by the nearby tryptophan, hence supporting a
3 different location for the tryptophan residue closer to the flavin molecule.

4 Results from visible and infrared transient absorption measurements are also running
5 alongside the above measurements: in the light-adapted state a flavin and tryptophan radical
6 was formed, which is not the case for the dark-adapted state^{9,58} (in agreement with our
7 assessment that W104 is not a main quencher in the dark-adapted state). These observations
8 can be explained by the simple mechanistic scheme that in the light-adapted state efficient
9 electron transfer takes place by the closely positioned tryptophan that can provide an electron
10 to the excited flavin, quench it and form the flavin radical. As discussed above, the distances
11 obtained from the electron transfer calculations are in good agreement with those obtained from
12 the FRET measurements supporting an electron transfer from the tryptophan residue. This
13 finding is underlined by the strong quenching of the excited state observed in the
14 Y21F/Y56F/W64F mutant where the only possible electron donor is W104 and it is in a close
15 position (7.6 Å) to the flavin. Fluorescence anisotropy decay measurements performed on this
16 mutant show a similar rotational correlation time (see Fig S5) to that measured in the light-
17 adapted case, indicating that the tryptophan in this case is locked – most probably via hydrogen
18 bonding –in a light-like state.

19 The TRIR data measured on the light-adapted W64F and Y21FY56FW64F mutant show a
20 downshift of frequency of C4=O carbonyl which can be explained based on the model
21 suggested by Iwata et al.⁶⁵ (Model 7) which assumes a hydrogen bond formed between the enol
22 form of Q63 and W104.

23 All these findings agree with recent calculations²⁷ which show-that the close position of W104
24 is favourable for electron transfer from tryptophan to flavin and this pathway is dominating
25 when the transfer is from Y21 to the flavin is repressed by the lack of the conducive hydrogen-

1 bonding network. Overall our study addresses previous discrepancies on the subtle
2 conformational differences of W104 that characterize the dark and signalling states in BLUF
3 containing photoreceptor proteins, and provides significant information that allow us to re-
4 examine the existing mechanistic models of photoactivation⁶³. The hierarchical relaxation
5 pathway after flavin photoexcitation in the AppA_{BLUF} domain occurs on the pico- to nanosecond
6 time scale for the flavin localized dynamics while subsequent protein structural reorganization
7 is observed over microseconds⁶⁶. Rearrangement of the hydrogen bond network around the
8 flavin is believed to be associated with conformational changes involving the β 5 strand^{18,28}.
9 The movement of the W104 residue which resides on the β 5-strand is therefore not unexpected
10 and can result in further structural changes in the C-terminal α -helices affecting the interaction
11 of the protein with downstream components. Understanding the dynamic behaviour of
12 photoreceptors has a major impact on the field of optogenetics as it can help to control gene
13 expression and engineered biosynthetic pathways.

14

15

16

17 **Conclusions**

18 A wide range of fluorescence techniques, time-resolved infrared spectroscopy and unnatural
19 amino acid incorporation have been used to probe the functional dynamics of W104 during the
20 photoactivation process in AppA_{BLUF}. Fluorescence resonance transfer (FRET) measurements
21 and replacement of W104 with the unnatural amino acid 7AW have provided quantitative
22 information on the position of W104 in the dark- and light-adapted states. In the light-adapted
23 state, W104 is in a restricted environment with an enhanced H-bond network compared to that
24 in the dark-adapted state, as revealed by the long rotational correlation time (~11 ns) for W104
25 from fluorescence anisotropy decay measurements and the downshift of the C4=O carbonyl

1 band from infrared measurements. In addition, in the light-adapted state the short fluorescence
2 lifetime of the flavin suggests that efficient electron transfer takes place to the excited flavin
3 from the closely placed W104. Overall, our study addresses previous discrepancies on the
4 position of W104 during the photocycle of AppA and supports a conformation of W104 close
5 to the flavin in the light-adapted state whereas in the dark-adapted state W104 is present in a
6 less restricted environment pointing away from the flavin.

1 **Methods**

2 **Materials** Glycogen (Type III: from rabbit liver) was purchased from Sigma-Aldrich. 7AW
3 was purchased from Aldrich.

4 **Expression and purification of W64F, W64F/W104A, 7aza-W104/W64F,** 5 **Y21F/Y56F/W64F and Y21F/Y56F/W64F/W104F AppA_{BLUF} proteins.**

6 The BLUF domain of AppA (AppA_{BLUF}: residues 5-125) was expressed in BL21(DE3)
7 *Escherichia coli* cells. Protein expression and purification were performed in the dark as
8 described previously⁶³. The constructs for the site-directed mutants W64F, W64F/W104A and
9 Y21F/Y56F/W64F were generated by PCR amplification (overlap extension method) from
10 wild-type pET15b_AppA_{BLUF} using the NdeI and BamHI restriction enzymes. The primers used
11 are described in Supplementary Table 1. The DNA sequence for expressing the
12 Y21F/Y56F/W64F/W104F AppA_{BLUF} mutant was purchased from Thermo Fisher Scientific.
13 All constructs were verified by DNA sequencing.

14 The 7aza-W104/W64F construct was expressed in a Tryptophan auxotroph M5219
15 *Escherichia coli* strain, purchased from the Belgian Coordinated Collection of Micro-
16 Organisms (<http://bccm.belspo.be/>). The W64F AppA_{BLUF} contained construct was transformed
17 by heat shock into M5219 *E.coli* strain. A single colony was inoculated into 25mL Luria Broth
18 (LB) media incubated overnight at 200 rpm and 37 °C. The overnight culture was used to
19 inoculate 1 L M9 medium in a 4L flask and cells were allowed to grow at 30°C until OD_{600nm}
20 reached ~0.6. At this point, the cells were washed three times with washing buffer (M9 mineral
21 salts solution and 20% glucose) to remove all Trp from the medium harvested, resuspended in
22 1L M9 media and cultured for additional 30 minutes at 30°C to ensure full Trp depletion⁶⁶.
23 7AW was then added and incubated for 15 minutes to allow the cells to uptake it after which
24 the temperature was decreased to 18°C for 30 min followed by addition of 0.8 mM isopropyl
25 β-D-1-thiogalactopyranoside (IPTG) to induce protein expression. Cells were harvested, lysed

1 and purified using Ni-NTA affinity resin as previously described⁶³. The apoprotein is produced
2 during protein expression and was isolated by size exclusion chromatography.

3 **Steady state optical and fluorescence measurements**

4 Absorption spectra were measured on a Perkin Elmer Lambda XLS+ and Jasco V-660
5 spectrophotometer. Fluorescence emission spectra were obtained with a Horiba Jobin Yvon
6 Fluorolog spectrofluorometer using three excitation wavelengths ($\lambda_{exc}=295\text{nm}$, $\lambda_{exc}=350\text{ nm}$,
7 $\lambda_{exc}=455\text{ nm}$). The applied slit width was set at 5 nm for both the excitation side and the
8 emission side. Unless otherwise indicated all fluorescence spectra were measured in the dark at
9 22°C in a 10 mm x 1 mm quartz cuvette.

10 **Fluorescence lifetime and anisotropy decay measurements.**

11 *Fluorescence anisotropy decay measurements:* Fluorescence anisotropy (r) measures the
12 depolarization of the fluorescence emission of a fluorophore due to energy transfer to another
13 molecule with a different orientation or molecular rotation caused by Brownian motion. As
14 molecular motion depends on the size of the fluorophore and on local environment factors
15 such as viscosity and molecular confinement, fluorescence anisotropy measurements can
16 provide useful information on the molecular size and the mobility of the fluorophore³². In the
17 case of continuous excitation, the steady-state anisotropy \bar{r} is given by equation (7)

$$18 \quad \bar{r} = \frac{I_{\parallel} - I_{\perp}}{I_{\parallel} + 2I_{\perp}} \quad (7)$$

19 where I_{\parallel} and I_{\perp} are the intensities measured with a linear polarizer for emission parallel and
20 perpendicular, respectively, to the electric vector of linearly polarized incident light.

21 In a fluid environment, the initial uniaxial distribution of photo-selected molecules is
22 usually randomized by rotational diffusion, leading to depolarization of fluorescence. In the
23 case of free rotation, the fluorescence anisotropy decays to zero, while for hindered rotation
24 the anisotropy reaches a time-independent value at long times, r_{∞} , called the *residual*
25 *anisotropy*. The analysis of the temporal decay of the anisotropy yields direct information on

1 the orientational dynamics, which depends on the size and shape of the rotating species and
2 on the fluidity of its microenvironment^{31,32}.

3 For an excited species in a single type of isotropic environment, $r(t)$ is, in general,
4 given by a linear combination of exponentially decaying functions

$$5 \quad r(t) = \sum_i \beta_i \exp\left(-\frac{t}{\theta_i}\right) \quad (8)$$

6 where θ_i are the rotational correlation times. Obviously, the sum of the factors β_i yields
7 the initial emission anisotropy r_0 .

8 *Nanosecond time-resolved fluorescence measurements:* Time-resolved fluorescence
9 measurements in the nanosecond time range were performed on a Horiba Jobin Yvon Nanolog
10 FL3-2Ihr spectrofluorometer operating in the time-correlated single photon counting (TCSPC)
11 mode and coupled to a R928P Hamamatsu photomultiplier. The applied detector voltage was
12 set at 950 V. The measurement range of the TCSPC system was 200 ns, the repetition rate of
13 the excitation pulses was 1 MHz, and the sync delay was 50 ns. Decay curves were collected in
14 4096 channels of a multi-channel analyser using a channel width of 55 ps. For the deconvolution
15 procedure, the dynamic instrument response function (IRF) was determined using a freshly
16 made solution of glycogen in water (half-bandwidth ~1 ns).

17 To measure the fluorescence lifetimes, the excitation wavelength was set at 321 nm (7aza-
18 tryptophan excitation) and 455 nm (FAD excitation) using pulsed nanoLEDs (Horiba) with
19 pulse duration <1ns and 1.2 ns, respectively. The fluorescence emission was collected at 380
20 nm ($\lambda_{\text{exc}}=321$ nm) and at 520 nm ($\lambda_{\text{exc}}=455$ nm). The appropriate emission wavelength was set
21 by a built-in the system monochromator. All measurements were performed using the Data
22 Station v 2.6 software and the lifetime and anisotropy data were analysed using the Jobin Yvon
23 DAS6 v6.6 and FluoFit softwares.

24 To generate and maintain the light-adapted state of the BLUF proteins, a Thorlabs M385 L2-
25 C1 UV Led ($\lambda_{\text{exc}}=385$ nm) was placed above the sample compartment and covered by a black

1 cloth. The light-adapted state was reached after 10 minutes of illumination. Sample
2 concentration was in the 5-10 μM range. All samples were illuminated continuously during the
3 fluorescence measurements. To eliminate the 385 nm scattered light, an UG11 filter was placed
4 between the sample and the detector. 10 mm x 1 mm quartz cuvettes were used for the
5 measurements. UV-vis absorption spectra were obtained before and after the fluorescence
6 measurements to monitor the integrity of the light-adapted state, using a Thermo Scientific
7 Evolution 600 UV-vis spectrophotometer.

8 *Picosecond time-resolved fluorescence measurements:* Time-resolved fluorescence
9 experiments in the ps time range were performed using a spectrally resolved Kerr-Gate
10 femtosecond fluorometer. The setup employs a Kerr shutter^{67,68} and allows measuring
11 fluorescence spectra with a resolution of ~ 100 fs and up to the nanoseconds timescale⁵⁷. The
12 setup was described elsewhere^{52,72} but briefly, the excitation pulse centered at 390 nm is
13 obtained by frequency-doubling, using a BBO crystal, part of the 780 nm pulse from the
14 Ti:sapphire laser/amplifier system (Quantronix Integra-C) operating at 1 kHz. The remaining
15 780 nm beam is led through a motorized delay-line and focused into the Kerr medium where it
16 spatially overlapped the fluorescence from the sample. The Kerr medium used was CS₂
17 (response function width ~ 1.2 ps). The sample was flowed through the 1mm pathlength optical
18 cell using a peristaltic pump. To generate and maintain the light-adapted state the sample
19 reservoir was illuminated using a ThorLabs M450LP1 LED ($\lambda_{\text{exc}}=450$ nm). Transient
20 fluorescence spectra were measured on time windows up to 1500 ps for all samples. Global
21 analysis of the time and spectrally resolved data sets in terms of a linear combination of a
22 discrete number of components, each with a distinct exponential rate constant and decay-
23 associated spectrum⁶⁹, was completed using Glotaran⁷⁰.

24 *Transient infrared spectroscopy*

1 All transient infrared absorption (TRIR) measurements reported here used the 10 kHz ULTRA
2 facility developed at the Central Laser Facility of the Rutherford Appleton Laboratories which
3 are described in detail elsewhere.⁷¹ As described earlier^{9,15,71} ULTRA offers wide tunability in
4 the visible region, a broad bandwidth in the IR probe, sub 100 fs time resolution and excellent
5 stability. In the present experiments it was used in the visible pump – IR probe geometry. The
6 excitation (pump) wavelength was 450 nm with a pulse energy of a few hundred nJ focussed to
7 a 100 micron spot size. It was checked that the spectra and kinetics were independent of the
8 pump wavelength and pulse energy. IR transmission was measured sequentially for pump-on
9 and pump-off using a 5 kHz mechanical chopper, and the data were processed to provide the
10 TRIR difference spectra.

11

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

References

1. MacHeroux, P., Kappes, B. & Ealick, S. E. Flavogenomics - A genomic and structural view of flavin-dependent proteins. *FEBS J.* **278**, 2625–2634 (2011).
2. Losi, A. & Gärtner, W. Old chromophores, new photoactivation paradigms, trendy applications: flavins in blue light-sensing photoreceptors. *Photochem. Photobiol.* **87**, 491–510 (2011).
3. Conrad, K. S., Manahan, C. C. & Crane, B. R. Photochemistry of flavoprotein light sensors. *Nat. Chem. Biol.* **10**, 801–809 (2014).
4. Lukacs, A., Eker, A. P., Byrdin, M., Brettel, K. & Vos, M. H. Electron hopping through the 15 Å triple tryptophan molecular wire in DNA photolyase occurs within 30 ps. *J. Am. Chem. Soc.* **130**, 14394–14395 (2008).
5. Aubert, C., Vos, M. H., Mathis, P., Eker, A. P. & Brettel, K. Intraprotein radical transfer during photoactivation of DNA photolyase. *Nature* **405**, 586–90 (2000).
6. Chaves, I. *et al.* The cryptochromes: blue light photoreceptors in plants and animals. *Annu Rev Plant Biol* **62**, 335–64 (2011).
7. Sancar, A. Structure and function of DNA photolyase and cryptochrome blue-light photoreceptors. *Chem Rev* **103**, 2203–37 (2003).
8. Masuda, S., Hasegawa, K. & Ono, T. Light-induced structural changes of apoprotein and chromophore in the sensor of blue light using FAD (BLUF) domain of AppA for a signaling state. *Biochemistry* **44**, 1215–24 (2005).
9. Lukacs, A. *et al.* BLUF domain function does not require a metastable radical intermediate state. *J. Am. Chem. Soc.* **136**, 4605–15 (2014).
10. Gauden, M. *et al.* Photocycle of the flavin-binding photoreceptor AppA, a bacterial transcriptional antirepressor of photosynthesis genes. *Biochemistry* **44**, 3653–62 (2005).

- 1 11. Unno, M., Masuda, S., Ono, T. A. & Yamauchi, S. Orientation of a key glutamine residue in the
2 BLUF domain from AppA revealed by mutagenesis, spectroscopy, and quantum chemical
3 calculations. *J. Am. Chem. Soc.* **128**, 5638–9 (2006).
- 4 12. Obanayama, K., Kobayashi, H., Fukushima, K. & Sakurai, M. Structures of the Chromophore
5 Binding Sites in BLUF Domains as Studied by Molecular Dynamics and Quantum Chemical
6 Calculations. *Photochem. Photobiol.* **84**, 1003–1010 (2008).
- 7 13. Domratcheva, T., Hartmann, E., Schlichting, I. & Kottke, T. Evidence for Tautomerisation of
8 Glutamine in BLUF Blue Light Receptors by Vibrational Spectroscopy and Computational
9 Chemistry. *Sci. Rep.* **6**, 1–14 (2016).
- 10 14. Masuda, S., Hasegawa, K. & Ono, T. Tryptophan at position 104 is involved in transforming light
11 signal into changes of beta-sheet structure for the signaling state in the BLUF domain of AppA.
12 *Plant Cell Physiol.* **46**, 1894–901 (2005).
- 13 15. Brust, R., Lukacs, A. & Haigney, A. Proteins in Action: Femtosecond to Millisecond Structural
14 Dynamics of a Photoactive Flavoprotein. *J. Am. Chem. Soc.* **135**, 16168–16174 (2013).
- 15 16. Masuda, S., Tomida, Y., Ohta, H. & Takamiya, K. The Critical Role of a Hydrogen Bond between
16 Gln63 and Trp104 in the Blue-Light Sensing BLUF Domain That Controls AppA Activity. 1223–
17 1230 *J. Mol. Biol.* (2007)
- 18 17. Laan, W. *et al.* On the mechanism of activation of the BLUF domain of AppA. *Biochemistry* **45**,
19 51–60 (2006).
- 20 18. Dragnea, V., Arunkumar, a I., Yuan, H., Giedroc, D. P. & Bauer, C. E. Spectroscopic studies of the
21 AppA BLUF domain from *Rhodobacter sphaeroides*: addressing movement of tryptophan 104 in
22 the signaling state. *Biochemistry* **48**, 9969–9979 (2009).
- 23 19. Anderson, S. *et al.* Structure of a novel photoreceptor, the BLUF domain of AppA from
24 *Rhodobacter sphaeroides*. *Biochemistry* **44**, 7998–8005 (2005).

- 1 20. Jung, A., Reinstein, J., Domratcheva, T., Shoeman, R. L. & Schlichting, I. Crystal Structures of the
2 AppA BLUF Domain Photoreceptor Provide Insights into Blue Light-mediated Signal Transduction.
3 *J. Mol. Biol.* **362**, 717–732 (2006).
- 4 21. Kita, A., Okajima, K., Morimoto, Y., Ikeuchi, M. & Miki, K. Structure of a cyanobacterial BLUF
5 protein, Tll0078, containing a novel FAD-binding blue light sensor domain. *J Mol Biol* **349**, 1–9
6 (2005).
- 7 22. Yuan, H. *et al.* Crystal structures of the *Synechocystis* photoreceptor Slr1694 reveal distinct
8 structural states related to signaling. *Biochemistry* **45**, 12687–94 (2006).
- 9 23. Jung, A. *et al.* Structure of a bacterial BLUF photoreceptor: Insights into blue light-mediated
10 signal transduction. *Proc. Natl. Acad. Sci. USA* **102**, 12350–12355 (2005).
- 11 24. Ohki, M. *et al.* Structural insight into photoactivation of an adenylate cyclase from a
12 photosynthetic cyanobacterium. *Proc. Natl. Acad. Sci. USA* **113**, 6659–64 (2016).
- 13 25. Udvarhelyi, A. & Domratcheva, T. Glutamine rotamers in BLUF photoreceptors: A mechanistic
14 reappraisal. *J. Phys. Chem. B* **117**, 2888–2897 (2013).
- 15 26. Collette, F., Renger, T. & Schmidt am Busch, M. Revealing the functional states in the active site
16 of BLUF photoreceptors from electrochromic shift calculations. *J Phys Chem B* **118**, 11109–11119
17 (2014).
- 18 27. Goyal, P. & Hammes-Schiffer, S. Role of active site conformational changes in photocycle
19 activation of the AppA BLUF photoreceptor. *Proc. Natl. Acad. Sci. USA* **114**, 1480–1485 (2017).
- 20 28. Grinstead, J. S. *et al.* The solution structure of the AppA BLUF domain: insight into the
21 mechanism of light-induced signaling. *ChemBioChem* **7**, 187–93 (2006).
- 22 29. Unno, M., Kikuchi, S. & Masuda, S. Structural refinement of a key tryptophan residue in the BLUF
23 photoreceptor AppA by ultraviolet resonance Raman spectroscopy. *Biophys. J.* **98**, 1949–56
24 (2010).
- 25 30. Toh, K. C. *et al.* On the signaling mechanism and the absence of photoreversibility in the AppA
26 BLUF domain. *Biophys. J.* **95**, 312–21 (2008).

- 1 31. Valeur, B. *Molecular Fluorescence: Principles and applications*. (Wiley-VCH Verlag GmbH, 2001).
- 2 32. Joseph R. Lakowicz. *Principles of Fluorescence Spectroscopy*. (Springer, 2006).
- 3 33. Förster, T. Zwischenmolekulare Energiewanderung und Fluoreszenz. *Annalen der Physik* **437**, 55–
4 75 (1948).
- 5 34. Okamoto, K. & Sako, Y. Recent advances in FRET for the study of protein interactions and
6 dynamics. *Curr Opin Struct Biol* **46**, 16–23 (2017).
- 7 35. Stryer, L. & Haugland, R. P. Energy transfer: a spectroscopic ruler. *Proc. Natl. Acad. Sci. USA* **58**,
8 719–26 (1967).
- 9 36. Antosiewicz, J. M. & Shugar, D. UV-Vis spectroscopy of tyrosine side-groups in studies of protein
10 structure. Part 1: basic principles and properties of tyrosine chromophore. *Biophys Rev* **8**, 151–
11 161 (2016).
- 12 37. Maeno, A., Matsuo, H. & Akasaka, K. Tyrosine/tyrosinate fluorescence at 700 MPa: a pressure
13 unfolding study of chicken ovomucoid at pH 12. *Biophys Chem* **183**, 57–63 (2013).
- 14 38. Martin, R. B., Edsall, J. T., Wetlaufer, D. B. & Hollingworth, B. R. A complete ionization scheme for
15 tyrosine, and the ionization constants of some tyrosine derivatives. *J Biol Chem* **233**, 1429–35
16 (1958).
- 17 39. Szabo, A. G., Lynn, K. R., Krajcarski, D. T. & Rayner, D. M. Tyrosinate fluorescence maxima at 345
18 nm in proteins lacking tryptophan at pH 7. *FEBS Lett* **94**, 249–52 (1978).
- 19 40. Prendergast, F. G., Hampton, P. D. & Jones, B. Characteristics of tyrosinate fluorescence emission
20 in alpha- and beta-purothionins. *Biochemistry* **23**, 6690–6697 (1984).
- 21 41. Pundak, S. & Roche, R. S. Tyrosine and tyrosinate fluorescence of bovine testes calmodulin:
22 calcium and pH dependence. *Biochemistry* **23**, 1549–1555 (1984).
- 23 42. Alexander Ross, J. B., Laws, W. R., Rousslang, K. W. & Wyssbrod, H. R. Tyrosine fluorescence and
24 phosphorescence from proteins and polypeptides. in *Topics in Fluorescence Spectroscopy*,
25 *Biomedical Applications* vol. 3 1–64 (Kluwer Academic Publishers, 2002).

- 1 43. Oktaviani, N. A. *et al.* Comprehensive Determination of Protein Tyrosine p K a Values for
2 Photoactive Yellow Protein Using Indirect ¹³ C NMR Spectroscopy. *Biophys. J.* **102**, 579–586
3 (2012).
- 4 44. Schlesinger, S. & Schlesinger, M. J. The effect of amino acid analogues on alkaline phosphatase
5 formation in Escherichia coli K-12. I. Substitution of triazolealanine for histidine. *J Biol Chem* **242**,
6 3369–72 (1967).
- 7 45. Twine, S. M. & Szabo, A. G. Fluorescent amino acid analogs. *Methods Enzymol.*, **360**, 104–27
8 (2003).
- 9 46. Alexander Ross, J. B., Rusinova, E., Luck, L. A. & Rousslang, K. W. Spectral Enhancement of
10 proteins in vivo incorporation of tryptophan analogues. in *Topics in Fluorescence, Protein*
11 *fluorescence* vol. 6 17–43 (Kluwer Academic Publishers, 2002).
- 12 47. Albani, J. R. Origin of tryptophan fluorescence lifetimes. Part 2: fluorescence lifetimes origin of
13 tryptophan in proteins. *J. Fluoresc.* **24**, 105–117 (2014).
- 14 48. Laan, W., van der Horst, M. A., van Stokkum, I. H. & Hellingwerf, K. J. Initial characterization of
15 the primary photochemistry of AppA, a blue-light-using flavin adenine dinucleotide-domain
16 containing transcriptional antirepressor protein from Rhodobacter sphaeroides: a key role for
17 reversible intramolecular proton transfer from the flavin adenine dinucleotide chromophore to a
18 conserved tyrosine? *Photochem. Photobiol.* **78**, 290–7 (2003).
- 19 49. Ohki, M. *et al.* Molecular mechanism of photoactivation of a light-regulated adenylate cyclase.
20 *Proc. Natl. Acad. Sci. U.S.A.* **114**, 8562–8567 (2017).
- 21 50. Winkler, A. *et al.* A ternary AppA-PpsR-DNA complex mediates light regulation of photosynthesis-
22 related gene expression. *Nat Struct Mol Biol* **20**, 859–67 (2013).
- 23 51. Dijkstra, D. S., Broos, J., Visser, A. J., van Hoek, A. & Robillard, G. T. Dynamic fluorescence
24 spectroscopy on single tryptophan mutants of Ell(mtl) in detergent micelles. Effects of substrate
25 binding and phosphorylation on the fluorescence and anisotropy decay. *Biochemistry* **36**, 4860–6
26 (1997).

- 1 52. Nag, L., Lukacs, A. & Vos, M. H. Short-Lived Radical Intermediates in the Photochemistry of
2 Glucose Oxidase. *ChemPhysChem* **20**, 1793–1798 (2019).
- 3 53. Liu, Z. *et al.* Determining complete electron flow in the cofactor photoreduction of oxidized
4 photolyase. *Proc. Natl. Acad. Sci. USA* **110**, 12966–71 (2013).
- 5 54. Mataga, N. *et al.* Dynamics and Mechanisms of Ultrafast Fluorescence Quenching Reactions of
6 Flavin Chromophores in Protein Nanospace. *J. Phys. Chem. B* **104**, 10667–10677 (2000).
- 7 55. Laptinok, S. P. *et al.* Ultrafast real-time visualization of active site flexibility of flavoenzyme
8 thymidylate synthase ThyX. *Proc. Natl. Acad. Sci. USA* **110**, 8924–9 (2013).
- 9 56. Mataga, N., Chosrowjan, H., Shibata, Y. & Tanaka, F. Ultrafast Fluorescence Quenching Dynamics
10 of Flavin Chromophores in Protein Nanospace. *J. Phys. Chem. B* **102**, 7081–7084 (1998).
- 11 57. Laptinok, S. P., Nuernberger, P., Lukacs, A. & Vos, M. H. Subpicosecond Kerr-gate
12 spectrofluorometry. *Methods Mol Biol* **1076**, 321–36 (2014).
- 13 58. Brazard, J. *et al.* New insights into the ultrafast photophysics of oxidized and reduced FAD in
14 solution. *J Phys Chem A* **115**, 3251–62 (2011).
- 15 59. Leenders, R., Van Hoek, A., Van Iersel, M., Veeger, C. & Visser, A. J. Flavin dynamics in oxidized
16 *Clostridium beijerinckii* flavodoxin as assessed by time-resolved polarized fluorescence. *Eur. J.*
17 *Biochem.* **218**, 977–984 (1993).
- 18 60. Leenders, H. R., Vervoort, J., van Hoek, A. & Visser, A. J. Time-resolved fluorescence studies of
19 flavodoxin. Fluorescence decay and fluorescence anisotropy decay of tryptophan in *Desulfovibrio*
20 flavodoxins. *Eur. Biophys. J.* **18**, 43–55 (1990).
- 21 61. Page, C. C., Moser, C. C., Chen, X. & Dutton, P. L. Natural engineering principles of electron
22 tunnelling in biological oxidation–reduction. *Nature* **402**, 47–52 (1999).
- 23 62. Haigney, A. *et al.* Ultrafast Infrared Spectroscopy of an Isotope-Labeled Photoactivatable
24 Flavoprotein. *Biochemistry* **50**, 1321–1328 (2011).
- 25 63. Lukacs, A. *et al.* Photoexcitation of the Blue Light Using FAD Photoreceptor AppA Results in
26 Ultrafast Changes to the Protein Matrix. *J. Am. Chem. Soc.* **133**, 16893–16900 (2011).

- 1 64. Haigney, A. *et al.* Vibrational assignment of the ultrafast infrared spectrum of the
2 photoactivatable flavoprotein AppA. *J. Phys. Chem. B* **116**, 10722–10729 (2012).
- 3 65. Iwata, T. *et al.* Hydrogen Bonding Environments in the Photocycle Process around the Flavin
4 Chromophore of the AppA-BLUF domain. *J. Am. Chem. Soc.* **140**, 11982–11991 (2018).
- 5 66. Broos, J. Biosynthetic incorporation of tryptophan analogs in proteins. *Methods Mol Biol* **1076**,
6 359–70 (2014).
- 7 67. Arzhantsev, S. & Maroncelli, M. Design and characterization of a femtosecond fluorescence
8 spectrometer based on optical Kerr gating. *Appl Spectrosc* **59**, 206–20 (2005).
- 9 68. Schmidt, B., Laimgruber, S., Zinth, W. & Gilch, P. A broadband Kerr shutter for femtosecond
10 fluorescence spectroscopy. *Appl. Phys. B* **76**, 809–814 (2003).
- 11 69. van Stokkum, I. H., Larsen, D. S. & van Grondelle, R. Global and target analysis of time-resolved
12 spectra. *Biochim Biophys Acta* **1657**, 82–104 (2004).
- 13 70. Snellenburg, J. J., Laptенок, S., Seger, R., Mullen, K. M. & Stokkum, I. H. M. van. Glotaran: A Java-
14 Based Graphical User Interface for the R Package TIMP. *J. Stat. Softw.* **49**, 1–22 (2012).
- 15 71. Greetham, G. M. *et al.* ULTRA: A Unique Instrument for Time-Resolved Spectroscopy. *Appl*
16 *Spectrosc* **64**, 1311–1319 (2010).
- 17 72. Nag L., Sournia P., Myllykallio H., Liebl U. and Marten H. Vos, Identification of the TyrOH•+
18 Radical Cation in the Flavoenzyme TrmFO, *J. Am. Chem. Soc.* **139**, 11500-11505,(2017)
- 19 73. Gil, A *et al.* Photoactivation of the BLUF Protein PixD Probed by the Site-Specific Incorporation of
20 Fluorotyrosine Residues, *J. Am. Chem. Soc.* 2017, 139, 41, 14638-14648

21

1
2
3
4
5
6
7
8
9
10
11

Acknowledgments: A.L. acknowledges funding from EFOP-3.6.2-16-2017-00005. A.L. and M.H.V. are grateful to the “Balaton” project (NKFIH 2017-2.2.5-TÉT-FR-2017-00005 and PHC Balaton 40173VE) for funding the travel exchanges. S.R.M. is grateful to EPSRC for financial support (EP/N033647/1 and EP/M001997/1). P.J.T. thanks NSF for financial support (MCB-1817837)

We thank Éva Hoffmanné Simon for excellent technical assistance in protein expression and purification.

12 **Author contributions**

13 All raw data and metadata from experiments and simulations are available from the
14 corresponding author on request. A.L., K.R., S.R.M. and P.J.T. designed the experiments.
15 I.P. developed the protocol for the production of the 7aza-W104/W64F App_{ABLUF} mutant and
16 Z.F. made the constructs of all App_{ABLUF} mutants. Experimental work was conducted by
17 K.K., S.M.K., K.R. I.P., R.K., Z.F., J.I., J. O., G.G., M.H.V. and A.L. Experimental data were
18 analysed by K.K., S.M.K., K.R., M.N., M.H.V. and A.L. and results discussed by all authors.
19 S.M.K and A.L. wrote the paper. The project was conceived by A.G., J.T.C., S.R.M., P.J.T.
20 and A.L.

21