

Chemistry A European Journal

 **Chemistry
Europe**
European Chemical
Societies Publishing

Accepted Article

Title: Exploring Multi-Subsite Binding Pockets in Proteins: DEEP-STD NMR Fingerprinting and Molecular Dynamics Unveil a Cryptic Subsite at the GM1 Binding Pocket of Cholera Toxin B

Authors: Serena Monaco, Samuel Walpole, Hassan Doukani, Ridvan Nepravishta, Ridvan Nepravishta, Macarena Martínez-Bailén, Ana Teresa Carmona, Javier Ramos-Soriano, Maria Bergström, Inmaculada Robina, Jesus Angulo, Jesus Angulo, and Jesus Angulo

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: *Chem. Eur. J.* 10.1002/chem.202001723

Link to VoR: <https://doi.org/10.1002/chem.202001723>

WILEY-VCH

FULL PAPER

Exploring Multi-Subsite Binding Pockets in Proteins: DEEP-STD NMR Fingerprinting and Molecular Dynamics Unveil a Cryptic Subsite at the GM1 Binding Pocket of Cholera Toxin B

Serena Monaco,^[a] Samuel Walpole,^[a] Hassan Doukani,^[a] Ridvan Nepravishta,^[a,b] Macarena Martínez-Bailén,^[c] Ana T. Carmona,^[c] Javier Ramos-Soriano,^[c] Maria Bergström,^[d] Inmaculada Robina,^{*,[c]} Jesus Angulo^{*,[a,c,e]}

Dedicated to Prof. Jesús Jiménez-Barbero on the occasion of his 60th birthday

School of Pharmacy, University of East Anglia, Norwich Research Park, NR4 7TJ Norwich, UK

Department of Organic Chemistry, Faculty of Chemistry, University of Seville, 41012 Seville, Spain

- [a] Dr. Serena Monaco, Dr. Samuel Walpole, Hassan Doukani, Dr. Ridvan Nepravishta, Dr. Jesús Angulo
School of Pharmacy
University of East Anglia
Norwich Research Park, NR4 7TJ, Norwich, UK
E-mail: j.angulo@uea.ac.uk
- [b] Dr. Ridvan Nepravishta
Department of Biochemistry & Molecular Biology, Sealy Center for Structural Biology & Molecular Biophysics
University of Texas Medical Branch
301 University Blvd, Galveston, TX, 77555-1068 USA
- [c] Dr. Macarena Martínez-Bailén, Dr. Ana T. Carmona, Dr. Javier Ramos-Soriano, Prof. Inmaculada Robina, Dr. Jesús Angulo
Department of Organic Chemistry, Faculty of Chemistry
University of Seville
41012 Seville, Spain
E-mail: robina@us.es, janqulo@us.es
- [d] Dr. Maria Bergström
Department of Chemistry and Biomedical Sciences
Linnaeus University
391 82 Kalmar, Sweden
- [e] Dr. Jesús Angulo
Instituto de Investigaciones Químicas (CSIC-US)
Avda. Américo Vespucio, 49
41092 Sevilla, Spain

Supporting information for this article is given via a link at the end of the document

Abstract: Ligand-based NMR techniques to study protein-ligand interactions are potent tools in drug design. STD NMR stands out as one of the most versatile, allowing screening of fragments libraries and providing structural information on binding modes. Recently, we have shown that a multi-frequency STD NMR approach, DEEP-STD NMR, can provide additional information on the orientation of small ligands within the binding pocket. Here, we have expanded the approach by what we call *DEEP-STD NMR fingerprinting* to explore the binding subsites of cholera toxin subunit B (CTB). To that aim, we present the synthesis of a set of new ligands which have been subject to a thorough study of their interactions with CTB by WAC and NMR. Remarkably, the combination of DEEP-STD NMR fingerprinting and Hamiltonian Replica Exchange Molecular Dynamics has proved to be an excellent approach to explore the geometry, flexibility, and ligand occupancy of multi-subsite binding pockets. In the particular case of CTB, it allowed to reveal the existence of a hitherto unknown binding subsite adjacent to the GM1 binding pocket, paving the way to the design of novel leads for inhibition of this relevant toxin.

Introduction

In the context of drug discovery, fragment-based drug discovery (FBDD) has gained momentum along the last decade as a method for lead generation.^[1] The technique relies on screening small compounds against a pharmacologically relevant target to find fragments with low affinity towards some of the adjacent binding *subsites* in which the target's binding pocket can be formally thought as composed of. The binding fragments are then combined together to obtain potent leads. In the FBDD approach, detailed information on the target structural features is in fact necessary.^[2]

On these premises, it is not surprising that NMR is one of the election methods to rationally design novel drugs using FBDD, particularly for low affinity binders, which tend to generate protein-ligand complexes difficult to crystallize. Back in 1997, the first NMR methodology for efficient ligand screening and lead design was proposed: "SAR by NMR" (structure-activity relationship by NMR).^[3] In SAR by NMR, isotopically labelled proteins are studied with each fragment of a particular library, and chemical shift perturbations allow, upon protein assignment, to identify different subsite binders. Analysis of further NOESY experiments allows to

FULL PAPER

give rational directions to connect fragments, in order to obtain very high affinity leads.^[3] A complementary approach, based on observation of NMR signals of the binder, is the use of ligand-based NMR techniques.^[4] Within this approach, one decade later, Pellecchia and co-workers proposed a method based on transfer-NOESY experiments: "SAR by Inter-Ligand NOE" (ILOE). Without the need of labelling and spectroscopic assignment of the targets, it is possible to study a small pool of fragments, accommodated in different adjacent binding subsites within a binding pocket, and provide points of contact between them in order to obtain inhibitors of nanomolar potency.^[5]

Among ligand-based NMR approaches, Saturation Transfer Difference NMR (STD NMR)^[6] spectroscopy has proved itself to be an efficient and reliable tool in FBDD, allowing to discriminate between specific and unspecific binders, even when they are analyzed as a pool.^[7] We have recently proposed an expansion of the methodology termed Differential Epitope Mapping STD NMR (DEEP-STD NMR) based on the new concept of multi-frequency STD NMR experiments.^[8] In DEEP-STD NMR, two binding epitope mappings are determined, following the use of two different frequencies for protein saturation, in contrast to a single one as in standard STD NMR. By changing the on-resonance frequency, different types of side chains in the binding pocket will reach slightly different levels of saturation and those differences can be picked up by building a differential epitope mapping, where the differences between the two mappings are revealed. Those differences indicate proximity to the type of protein side chain directly irradiated in each experiment, providing information on the type of amino acid side chain surrounding a proton of the ligand, as well as it hints about the orientation of the ligand relative to the side chains lining the binding pocket. This way, further than identifying selective binders, the technique allows to orient the fragments in the binding subsites. This is an extremely valuable information when it comes to drug design, and previous results from our laboratory have shown DEEP-STD NMR to be a very informative tool in the discovery of WWP2 ubiquitin ligase inhibitors.^[9]

Here, we have applied DEEP-STD NMR, combined with advanced molecular dynamics simulations to explore the plasticity and dynamics of the multi-subsite binding pocket of cholera toxin B (CTB). The interaction between CTB and its natural substrate, the glycolipid ganglioside GM1, is one of the strongest and best studied protein-carbohydrate interactions known, and is biologically highly relevant, as it mediates the onset of the cholera infection.^[10] Currently, still, cholera treatment mainly resorts to antibiotics, but the appearance of antibiotic resistant *V. cholera* strains^[11] have triggered the development of alternative cures for this endemic disease.^[12] Large efforts have been reported towards the design of inhibitors of the GM1/CTB interaction. The structural and thermodynamic studies on this complex showed that high specificity is given by the presence of two adjacent binding subsites, accommodating the two "arms" of the GM1 ligand, i.e. the galactose and sialic acid non-reducing ends.^{[13], [14], [15]} Grounded on this knowledge, many carbohydrate-based scaffolds have been designed that aim to target both binding subsites.^{[16], [17], [18], [19], [20]}

Despite these glycomimetics showing very good affinity levels, the challenge has remained to overcome the problem of hydrolyzability. In our labs, we have reported that CTB ligands based on thio-galactosides bearing a polyhydroxyalkylfuroate moiety (PHF), showed promise as non-hydrolyzable inhibitors.^[21]

The PHF moiety was demonstrated to contribute effectively to the affinity for CTB, and STD NMR studies on three representatives of this family of CTB ligands (Figure 1, compounds 1-3) supported a bidentate binding mode, so that both the galactose ring and the PHF moiety are the main contacts with the protein in the bound state of ligand 3.

In this work, we present the synthesis of a new set of CTB ligands to explore the GM1 binding pocket of CTB and propose the DEEP-STD NMR fingerprinting protocol, to gain orientational information of the ligands in the subsites by comparison to DEEP-STD NMR data of a binder with a known 3D structure of its complex to CTB. The combination of STD NMR competition experiments^[22], ILOEs^[23], docking^{[24], [25], [26]}, advanced molecular dynamics^[27] and STD NMR intensity prediction from 3D molecular models of the complexes using CORCEMA-ST^[28], has allowed us to answer the question of how large and flexible ligands like 3 can be accommodated in the GM1 binding pocket of CTB, demonstrating the existence of a hitherto unknown binding subsite adjacent to the GM1 binding pocket, paving the way to the design of novel non-carbohydrate leads for CTB inhibition.




Figure 1. Structure of ligands 1, 2, and 3^[21] and the novel set of thio-galactoside CTB ligands 4-9 described in this work


Results and Discussion

A novel set of promising small molecule CTB inhibitors, to explore the binding subsites of the GM1 binding pocket of CTB by DEEP-STD NMR fingerprinting. The STD NMR results previously obtained for CTB ligand 3^[21] indicated that both (galactose and benzyl) ends of the binder established the closest contacts with CTB in the bound state, whereas the polyhydroxyalkyl spacer made less contacts with the protein. As no 3D structure of the 3/CTB complex is available, this result could be interpreted as a simultaneous binding of both ends to the galactose and sialic acid subsites within the GM1 binding pocket

FULL PAPER

of CTB (bidentate bound mode). Based on those results, we first wanted to experimentally probe the role in CTB binding of the polyhydroxyalkyl spacer linking both ends in **3**, to inform size optimization of novel CTB ligands. To that aim, we designed a novel set of shorter thio-galactoside ligands (compounds **4-9**, Figure 1) lacking the triazole-polyhydroxyalkyl linker present in **2** and **3** and have instead the thio-galactose residue directly connected to the benzyl furoate moiety. The variability across the compound set is given by the anomeric configuration (α or β), and the oxidation state of the glycosidic sulphur (present as a thioether, sulphone or sulphoxide functional group), for a total of 6 new compounds (Figure 1).

The synthesis of α -galactosides **5**, **7** and **9** (Scheme 1) started from peracetylated thioglycosides **13**.^[29] Selective S-deacetylation followed by reaction with chloro derivative **11b** afforded α -thioglycoside **15** in 48% yield. Compound **11b** was prepared from polyhydroxyalkylfuroate **10**,^[21] after oxidation with NaIO₄ followed by reduction with NaBH₄ and chlorination. Subsequent nucleophilic displacement with KSAc and deacetylation afforded thiol **12b** in good overall yield, which after reaction with penta-O-acetyl- α -D-galactopyranosyl bromide **14** exclusively gave β -thioglycoside **16**. Oxidation of **15** and **16** with MCPBA under different reaction conditions afforded sulphone derivatives **17** and **18** and sulphoxides **19** and **20**. Final deacetylation of compounds **15-20** under Zemplén conditions gave compounds **4-9** in 65-99% yield.



Scheme 1. Reagents and conditions: (a) 1. NaIO₄, H₂O/MeOH, 0 °C→r.t.; 2. NaBH₄, MeOH, r.t.; (b) NCS, Me₂S, CH₂Cl₂, -20 °C→0 °C; (c) KSAc, DMF, 50 °C; (d) NaOMe, MeOH, 0 °C; (e) EtNH₂, DMF, r.t.; (f) aq. Na₂CO₃, EtOAc, TBAHS, r.t.; (g) MCPBA, CH₂Cl₂, r.t.; (h) MCPBA, CH₂Cl₂, -78 °C→-30 °C.

Table 1. K_D for the binding of ligands **4-9** to CTB, as determined by WAC.

Compounds	Glycosidic bond	X, X'	K _D (mM)
4	β	none, none	0.35
5	α	none, none	0.26
6	β	O, O	0.34
7	α	O, O	0.34
8	β	O, none	0.45
9	α	O, none	0.48

The interactions of **4-9** with CTB were first studied by weak affinity chromatography (WAC),^[30] and K_D values are shown in Table 1. The new ligands showed K_D ranging from 0.35 mM to 0.48 mM, coming across as very promising CTB binders as compared to **1-3**, whose K_D values were in the low mM range.^[21]

STD NMR experiments were carried out to confirm the binding to CTB for all the members of the set **4-9**, and to get structural information on their binding modes.




Figure 2. STD NMR binding epitope mapping of ligands **4-9**, based on the normalized STD values from the initial slope of each proton, relative to the most intense one (for which a 100% is arbitrarily assigned). Legend indicates weak (blue), medium (yellow) and strong (red) intensities (closer contacts with CTB). Proton assignment of the ligands, raw build-up curves data and normalized STD values are reported in the Supporting Information.

The determined binding group epitope mappings of their interactions with CTB showed a significant degree of variability among the six ligands (Figure 2), even at the common galactose ring. This was surprising, as the GM1 binding pocket of CTB provides a specific galactose subsite, which called for further investigation. To that aim, we resorted to our recently developed DEEP-STD NMR method^[8] to elucidate ligand orientation in the bound state.

FULL PAPER

“DEEP-STD NMR fingerprinting” characterizes the bound orientations of 4-7 in the GM1 binding pocket of CTB. DEEP-STD NMR (Differential Epitope Mapping STD NMR) is a novel tool for revealing the bioactive orientation of ligands in a protein binding pocket, if the 3D geometry of the binding pocket is known.^[6] The availability of a crystal structure of the complex of CTB with a small galactoside derivative, 3-nitrophenyl- α -D-galactopyranoside, 3NPG (PDB ID: 1EEI),^[31] encouraged us to propose a novel method that we call “DEEP-STD NMR fingerprinting”. First, the DEEP-STD NMR pattern of 3NPG in complex with CTB is determined, providing DEEP-STD factors, Δ STD, which can be directly correlated with the orientation of 3NPG in the crystal structure (i.e. which ligand protons are close to a given type of amino acid side chain^[6]). Then, the patterns of DEEP-STD factors of the “unknown” ligands (**4-9**) are determined and compared to that of 3NPG. This approach, first, allows us to confirm binding of unknown ligands to a known binding pocket and, second, it sheds light on their relative orientation, in comparison to the orientation of known ligands. We carried out the DEEP-STD NMR fingerprinting of the GM1 binding pocket of CTB with 3NPG^[6] and ligands **4-7** (ligands **8** and **9** were excluded due to degradation over the time of the experiments). We irradiated the different samples containing the CTB/ligand complexes at 0.60 ppm and 2.25 ppm (on-resonance frequencies), as already reported for the 3NPG/CTB complex.^[6] At 2.25 ppm, the glutamate and glutamine side chains in the galactose subsite are directly irradiated, and at 0.60 ppm the irradiation is on aliphatic side chains.




Figure 3. Differential Epitope Mapping Δ STD(2.25ppm/0.60ppm) of (a) ligands **4-7** and (b) 3NPG in complex with CTB (reference sample), obtained at 2 s saturation time. For each ligand, the three largest positive Δ STD values are shown as orange bars. (c) Crystal structure of the complex 3NPG/CTB (PDB ID: 1EEI).^[31] Protein protons directly irradiated at 2.25 ppm are highlighted in orange surface.

The determined Δ STD (2.25 ppm/0.60 ppm) for each complex are shown in Figure 3 (see also Supporting Information, Figure S2). The overall positive Δ STD values at the galactose protons for all the ligands **4-7** (Figure 3a) indicate that the thio-galactose ring is allocated in the galactose binding subsite of the GM1 pocket, by

comparison with the positive Δ STDs observed for the 3NPG/CTB complex (Figure 3b). Still, differences in Δ STDs were noticeable for some galactose protons of **4-7**, suggesting some reorientation of the galactose moiety in the galactose subsite, which results from the chemical differences at the benzyl furoate side of the molecules, in good agreement with the experimental binding epitope data (Figure 2).

Molecular docking: ligands 4-7 bind CTB occupying the GM1 galactose and sialic acid subsites. We performed rigid molecular docking of ligands **4-7** using the crystal structure of CTB in the GM1/CTB complex (PDB ID: 3CHB).^[32] For each ligand, the lowest energy solutions converged to a binding mode in which the thio-galactose moiety sits in the galactose binding subsite and the furoate-benzyl tail lies in the sialic acid binding subsite within the GM1 binding pocket of CTB (Figure 4). Remarkably, the galactose ring of the four ligands showed different orientations within the subsite, in good agreement with the experimental NMR observations (Figures 2 and 3). On the contrary, the benzyl groups of all the four ligands converged into a single orientation in the adjacent sialic acid subsite (blue surface in Figure 4a), where a stabilizing π -stacking with Tyr12 can take place (Figure 4c).




Figure 4. (a) Lowest energy docking solutions for Ligand **4** (grey), Ligand **6** (pink), Ligand **5** (green) and Ligand **7** (violet) in complex with CTB. The residues lining the galactose and sialic acid binding subsites are highlighted in orange, and blue surfaces, respectively. (b) On a semi-transparent surface, the sulphone oxygens of **6** and **7** are shown pointing towards Asn14 and Trp88 (in turquoise sticks), and the methyl group on the furoate pointing parallel to His13 (red sticks), rather than towards Gln61 (turquoise sticks). (c) Zoom on the phenyl moieties of Ligands **4-7**, showing the interaction with Tyr12 (turquoise sticks).

These results agree with previous studies showing that CTB ligands with aromatic rings can efficiently π -stack with Tyr12 in the sialic acid binding subsite.^[31, 33] On the other hand, ligands **4-**

FULL PAPER

7 fine tune the orientation of their thio-galactose moiety within the binding subsite by favorable contacts with the amide group of Asn14 and/or the NH group of the side chain of Trp88 present at the protein surface (Figure 4b). The impact on galactose orientation is substantial for the sulphone epimers, which can form H-bonds with Asn14 and Trp88, through their two oxygens (Figure 4b), disrupting the H-bond network of the galactose ring in the natural ligand GM1 and altering its orientation.

Globally, the 3D molecular models of the complexes of CTB with ligands **4-7**, in agreement with the DEEP-STD NMR fingerprinting approach, demonstrate that small galactoside-based ligands armed with a benzyl furoate tail bind the GM1 pocket in CTB in a bidentate mode, simultaneously occupying the galactose and sialic acid subsites. Decorations at the glycosidic linkage have a sensitive impact on the binding mode of the galactose ring. This is relevant for the design of CTB inhibitors based on mimicking the GM1 galactose, as sulphone affects mimicry of the galactose orientation in the GM1 natural ligand.

How do larger ligands **2 and **3** occupy the GM1 binding pocket? STD NMR competition experiments suggest the presence of an unknown subsite.** In the complexes of CTB with ligands **4-7**, the whole GM1 binding pocket is very efficiently occupied (Figure 4) with all parts of the relatively small ligand being intimately recognized by the protein (Figures 2 and 4). This result prompted the question of how a larger ligand, like **3**, is able to be accommodated in the pocket. Binding group epitope mappings of ligands **2** and **3** for their interactions with CTB have already been reported,^[21] supporting a bidentate mode of binding and showing that the spacer linking both ligand ends also establishes contacts with the protein (Supporting Information Figure S1).^[21] Interestingly, ligand **2** and the PHF tail of ligand **3** showed similar binding epitope mappings, strongly suggesting that both bind into the same CTB subsite (Supporting Information Figure S1).

Due to the non-carbohydrate nature of the PHF end present in ligands **2** and **3**, and their larger size, in comparison to ligands **4-7**, the question arises on whether these ligands might be interacting in a binding site other than sialic acid subsite of the GM1 pocket. Here, we tested this hypothesis and aimed to provide 3D models of the complexes of **2** and **3** with CTB, validated by NMR spectroscopy.




Figure 5. Chemical formulae for 3-Nitrophenyl α-D-galactopyranoside (3NPG; reporter for the Gal subsite) and 3' Sialyllactose (3'SL; reporter Neu5Ac subsite).

We started by exploring the occupancy of the two (galactose/sialic acid) binding subsites of CTB by STD NMR competition experiments. To that aim, we selected STD-active probe ligands known to occupy either subsite in the GM1 binding pocket of CTB (we call them subsite "reporters") and probed whether any of the ligands **2** or **3** was able to displace any of the reporters in competition experiments. 3NPG was used as the galactose subsite reporter and 3'-sialyllactose (3'SL), the Neu5Ac-containing branch of GM1, as the reporter for the sialic acid subsite (Figure 5).

First (Table 2, experiment a), STD NMR signals of a sample containing ligand **3** and CTB were observed and its competition with the galactose subsite reporter 3NPG was monitored. In this experiment, 3NPG displaced ligand **3** when added to the solution in equimolar amount (Supporting Information Figure S3a). Upon addition of 3NPG, the STD signals of ligand **3** were halved, in agreement with their similar K_D (3NPG 1.10 mM vs. ligand **3** 1.05 mM)^[21], confirming that the thio-galactose moiety of ligand **3** binds to the galactose subsite within the GM1 binding pocket.

After that, we probed competition between ligands **2** and **3**, so ligand **2** was added at an equimolar amount to a sample of ligand **3** in complex with CTB (Table 2, experiment b). In this case, the STD signals of ligand **3** dropped upon addition of ligand **2** (Supporting Information Figure S3b), which has a K_D =1.35 mM^[21], proving that both ligands **3** and **2** occupy a common subsite, most likely via their common PHF tail.

Table 2. Table of results of the competition experiments (for the STD spectra see Supporting Information Figure S3).

Experiment	Ligand	Competitor	Displacement
a)	3	3NPG	✓
b)	3	2	✓
c)	2	3NPG	✗
d)	3'SL	2	✗

Then, ligand **2** was tested in competition with 3NPG (Table 2, experiment c). No competition was observed in this case, showing that binding of ligand **2** to CTB does not involve the galactose subsite (Supporting Information Figure S3c).

Finally, to investigate if the PHF tail binds to CTB in the sialic acid subsite, competition between 3'SL and ligand **2** was carried out (Table 2, experiment d). Binding of 3'SL was detected (Supporting Information Figure S5), but the STD signals of 3'SL were not affected by the addition of ligand **2** (Supporting Information Figure S3d). As a control, adding 3'SL to a sample containing CTB and ligand **2**, did not affect the signals of **2** either (Supporting Information Figure S3e), confirming that binding of ligands **3** and **2** does not involve the sialic acid subsite. Globally, the STD NMR competition experiments strongly support the existence of an unknown subsite on CTB, able to accommodate the PHF tails in ligands **2** and **3**.

ILOE experiments explore subsites proximity in the GM1 binding pocket of CTB. The scenario depicted by the STD NMR competition experiments did not exclude that ligands **3** and **2** could compete for a subsite outside the GM1 binding pocket. On the other hand, as previously observed (experiment c, Table 2), 3NPG and Ligand **2** can bind simultaneously to CTB. On this basis, we decided to test experimentally if they bind in proximity

FULL PAPER

to each other (adjacent binding subsites) by generating the 2/CTB/3NPG ternary complex in solution and resorting to an ILOE (tr-NOESY) experiment^[23] (Figure 6).

In the tr-NOESY spectrum of the ternary complex we did observe an *inter*-ligand 2-3NPG cross peak between the triazole proton (H_{triaz}) of ligand 2 (7.67 ppm) and the H_{c,d} protons of 3NPG (positions *meta* and *para* to the nitro group, both at 7.27 ppm), along with the expected intra-ligand cross peaks for both ligands. As a control, to confirm the inter-ligand nature of that NOE, the same tr-NOESY experiment was performed on the binary 3NPG/CTB and ligand 2/CTB complexes as well as on CTB alone (see Supporting Information Figures S5 and S6 for control and full spectral width spectra of the ternary complex). The 7.67 ppm/7.27 ppm cross peak was absent in the three control spectra, confirming that the cross peak is not due to any protein-ligand or protein-protein NOE, but it genuinely reports a close contact (ILOE) between the triazole moiety of ligand 2 and the nitrophenyl group of 3NPG in the bound state.




Figure 6. Top, expansion of the superposition of tr-NOESY experiments of the 2/3NPG/CTB ternary complex (in black) and of the 3NPG/CTB control complex (in red). Diagonal and cross peaks are assigned and the ILOE peak correlating H_{c,d} of 3NPG and H_{triaz} of ligand 2 is highlighted in blue. Bottom, magnification of the ILOE cross peak. Signals belonging to ligand 2 and 3NPG are marked with a red asterisk and a black hashtag, respectively.

The observation of this *inter*-ligand 2-3NPG NOE in the state bound to CTB is of paramount importance in our study, as it proves the existence of a subsite accommodating the furoate-

benzyl chain, adjacent to the galactose subsite, but different to the sialic acid subsite. To generate 3D models of the complexes with 2 and 3, we then focused our structural search on the proximities of the GM1 binding pocket. Furthermore, the observed ILOE gave us information on the orientation of ligand 2 in this novel subsite, which is such that its triazole is in proximity of the galactose subsite.

Hamiltonian Replica Exchange Molecular Dynamics (HREMD) reveals a novel subsite in the GM1 binding pocket that accommodates the polyhydroxyalkylfuroate tail of 2 and 3. In contrast to the smaller ligands 4-7, docking calculations of ligands 2 and 3 into the GM1 binding pocket of CTB failed to generate 3D models that could agree with the experimental data (see Supporting Information). STD predictions from the docking models were obtained using CORCEMA-ST.^[28] Predicted STD intensities are compared to experimental NMR data through the so-called NOE R-factor, with a low value indicating a good matching. In our experience, values ≤ 0.3 report on good agreement between the 3D model and the experimental data. The PHF tail of 3 was predicted to occupy the sialic acid subsite, in disagreement with STD NMR competition experiments and binding epitopes, as quantitated by the high NOE R-factor (Supporting Information Figure S11). Similarly, docking solutions for ligand 2 also showed high NOE R-factors (Supporting Information Figure S12).

Taking into account the ILOE results, we then decided to investigate the dynamics of the GM1 binding pocket in CTB to probe the flexibility of side chains in the vicinity of the galactose and sialic acid subsites. To that aim, a 100 ns molecular dynamics (MD) simulation of free CTB was performed. In order to enhance sampling of the free state, an HREMD-based method was chosen in which the non-bonding interaction between the solvent and hydrophobic regions of the protein were modified in each replica, which has previously been shown to be effective in identifying druggable transient binding sites.^[27]




Figure 7. Opening of a novel subsite in the GM1 binding pocket of CTB. Top, in green, the conformations of the side chains of Ile58 and Lys34 in the crystal structure of CTB (PDB ID:3CHB); in turquoise, overlaid, the open state of the novel subsite from HREMD simulations. The red arrows indicate the χ_1 torsional rotation from closed to open states for the two residues involved (χ_1 values reported for both). Bottom, surface representation of the closed (left) and open conformations (right), corresponding to initial and final frames of the HREMD apo CTB simulation. Galactose binding subsite in orange; sialic acid binding subsite in blue; Ile 58 and Lys 34 in green and labelled.

FULL PAPER

Analysis of the trajectory over all five subunits of the CTB pentamer revealed the existence of a transient subsite close to the galactose and sialic acid subsites (Figure 7 and Supporting Information Figure S13). While this subsite is closed in the crystal structure, opening is facilitated by rotation of the Ile59 and Lys138 side chains (Figure 7). Since no main chain rearrangement is observed, the energy barrier between the 'closed' and 'open' conformations is expected to be minimal such that both states are readily sampled in the unbound protein.

Furthermore, repeating the simulation with CTB bound to GM1 revealed that opening of this cryptic subsite still takes place in the presence of the native ligand (Supporting Information Figures S14 and S15). This shows that this subsite is accessible even when the galactose and sialic acid subsites are engaged in binding, opening the door to further CTB inhibitors design that could exploit all the three subsites.

We then carried out docking calculations of ligand **3** to this 'open' conformation. The best docking solutions showed indeed that the PHF moiety occupies the novel subsite, in addition to good convergence at the galactose end towards the galactose subsite (Figure 8, ligand **3** in yellow). Furthermore, a long MD simulation of this complex indicated that this conformation is dynamically stable, and, most importantly, the novel subsite is occupied for most of the simulation (Supporting Information Figures S16 and S17).




Figure 8. Superposition of the representative MD frames of the most populated clusters of the complexes of CTB with ligand **3** (pale yellow) and ligand **2** (light red) bound to CTB in the "open" conformation. The galactose subsite is in orange and the sialic acid subsite in blue; Lys34 and Ile58, defining the novel subsite, in green.

The same approach was followed for ligand **2** to generate a 3D model of ligand **2** bound in the open conformation of CTB (Figure 8, ligand **2** in light red). Long MD simulation revealed even greater dynamic stability than the **3**/CTB complex, with the furoate-benzyl chain exclusively occupying the novel subsite throughout the simulation. Without a galactose moiety to act as an anchor, ligand **2** buries itself further into the hydrophobic binding subsite (Figure 8). Moreover, the basic side chain of Lys138 is able to form several favorable hydrogen bonding interactions with the furoate-benzyl moiety, which are not possible for ligand **3**.

Finally, the 3D models of the complexes were validated against the experimental NMR data by using CORCEMA-ST. Instead of

considering a single docking solution, STD NMR intensities were predicted running CORCEMA-ST on a set of 100 frames from each of the MD simulations of the complexes of ligands **2** and **3** with CTB, in order to reflect the flexibility of the systems. The agreement with the experimental data was very good, with NOE R-factors of 0.25 and 0.31, respectively (Supporting Information Figures S18 and S19). These results confirm that, in solution, the novel binding subsite in the GM1 binding pocket of CTB is populating the open state and can accommodate the PHF tail in the bound state to ligands **2** and **3**.

Conclusion

In this work, we have shown that our novel DEEP-STD NMR fingerprinting approach, combined with MD simulations, is a very powerful tool to explore the multi-subsite architecture of protein binding pockets. Here we have applied it to an important drug target as cholera toxin subunit B (CTB), but the approach is expected to be of broad applicability in the investigation of ligand binding to other multi-subsite relevant targets.

To explore the binding subsites of CTB, six novel thio-galactosidic CTB inhibitors (**4-9**) were synthesized and studied. These showed a much higher affinity for CTB, relative to their precursors.^[21] Structural analysis revealed that, for the six ligands, the aromatic moiety sits in the sialic acid binding subsite of GM1, π -stacking with the underlying side chain of Tyr12. On the other hand, the decoration at the thio-glycosidic linkage and the variability of the anomeric configuration disrupt mimicry of the galactose ring in the binding subsite of GM1.

For the larger ligands **2** and **3**,^[21] a combination of STD NMR competition and ILOE experiments, as well as docking and Hamiltonian Replica Exchange Molecular Dynamics (HREMD) allowed us to unveil the existence of a novel cryptic binding subsite on the GM1 binding pocket of CTB. This subsite was not present in the rigid crystal structure of CTB bound to GM1, but molecular dynamics revealed that it can be readily formed by rearrangements in the side chains of Ile59 and Lys138, which are adjacent to the well-known galactose and sialic acid binding subsites at the GM1 binding pocket. 3D models of the complexes, generated by MD simulations combined with STD NMR and validated using CORCEMA-ST, demonstrated that ligands **2** and **3** occupy this subsite with their polyhydroxyalkylfuroate tails. The existence of this new non-carbohydrate binding subsite in CTB, accessible even when the two known subsites are occupied, paves the way for future design of three-finger thio-galactose mimetics as potentials CTB inhibitors.

Experimental Section

Materials 3-nitrophenyl α -D-galactopyranoside (3NPG), Cholera Toxin Subunit B, deuterium oxide (99.9% ^2H), dimethyl sulphoxide- d_6 (DMSO- d_6), disodium hydrogen phosphate (Na_2HPO_4), potassium dihydrogen phosphate (KH_2PO_4), sodium chloride (NaCl) and potassium chloride (KCl) were purchased from Sigma Aldrich. Ligands were prepared in stock solutions 3% DMSO- d_6 .

Synthesis of Ligands 4-9

((4-Benzyloxycarbonyl-5-methylfuran-2-yl)methyl)-1-thio- β -D-galactopyranoside (4**):** To a solution of **16** (40 mg, 0.066 mmol) in anhydrous MeOH (2 mL), NaOMe (13 μL , 0.5 M in MeOH) was added and the mixture stirred at 0 $^\circ\text{C}$ for 2 h. Then the mixture was neutralized with

FULL PAPER

Amberlite IR-120H⁺, filtered, and washed with MeOH. The filtered solution was concentrated to give pure **4** (26 mg, 0.061 mmol, 92%) as a pale yellow oil. $[\alpha]_D^{25} = -104.8$ (c 0.64, MeOH). IR ($\bar{\nu}$) 3381 (OH), 2907, 1705 (C=O), 1072 cm⁻¹. ¹H NMR (300 MHz, CD₃OD) δ 7.43-7.31 (m, 5H, H-Ar), 6.52 (s, 1H, H-3'), 5.26 (s, 2H, CH₂Ph), 4.28 (d, 1H, $J_{1,2} = 9.6$, H-1), 4.03 (d, 1H, $J_{H,H} = 14.5$, CH₂S), 3.87 (dd, 1H, $J_{4,3} = 3.2$, $J_{4,5} = 0.6$, H-4), 3.78 (d, 1H, $J_{H,H} = 14.5$, CH₂S), 3.76 (dd, 1H, $J_{6a,6b} = 11.5$, $J_{6a,5} = 7.1$, H-6a), 3.68 (dd, 1H, $J_{6b,5} = 5.1$, H-6b), 3.57 (t, 1H, $J_{2,3} = 9.5$, H-2), 3.49 (td, 1H, H-5), 3.41 (dd, 1H, H-4), 2.52 (s, 3H, Me). ¹³C NMR (75.4 MHz, CD₃OD) δ 165.2 (COOBn), 160.4, 151.4 (C-2', C-5'), 137.7 (C_q-Ar), 129.6, 129.2, 129.1 (C-Ar), 115.0 (C-4'), 109.3 (C-3'), 86.1 (C-1), 80.8 (C-5), 76.3 (C-3), 71.4 (C-2), 70.5 (C-4), 67.0 (CH₂Ph), 62.7 (C-6), 26.2 (CH₂S), 13.9 (Me). HRESIMS m/z found 447.1079, calcd. For C₂₀H₂₄O₈NaS (M+Na)⁺: 447.1084.

((4-Benzyloxycarbonyl-5-methylfuran-2-yl)methyl)-1-thio- α -D-galactopyranoside (5): To a solution of **15** (47 mg, 0.079 mmol) in anhydrous MeOH (1.5 mL), NaOMe (16 μ L, 0.5 M in MeOH) was added and the mixture stirred at 0 °C for 1.5 h. Then the mixture was neutralized with Amberlite IR-120H⁺, filtered, and washed with MeOH. The filtered solution was concentrated to give pure **5** (34 mg, 0.079, quantitative) as a white solid. $[\alpha]_D^{25} = 249.8$ (c 0.68, MeOH). IR ($\bar{\nu}$) 3382 (OH), 2927, 1708 (C=O), 1069 cm⁻¹. ¹H NMR (300 MHz, CD₃OD) δ 7.43-7.29 (m, 5H, H-Ar), 6.50 (s, 1H, H-3'), 5.35 (d, 1H, $J_{1,2} = 5.7$, H-1), 5.25 (s, 2H, CH₂Ph), 4.16 (t, 1H, $J_{5,6} = 6.2$, H-5), 4.09 (dd, 1H, $J_{2,3} = 10.0$, H-2), 3.89 (d, 1H, $J_{4,3} = 2.4$, H-4), 3.82 (d, 1H, $J_{H,H} = 14.9$, CH₂S), 3.76-3.68 (m, 2H, H-6), 3.65-3.59 (m, 2H, H-3, CH₂S), 2.52 (s, 3H, Me). ¹³C NMR (75.4 MHz, CD₃OD) δ 165.2 (COOBn), 160.4, 151.4 (C-2', C-5'), 137.7 (C_q-Ar), 129.6, 129.21, 129.19 (C-Ar), 115.0 (C-4'), 109.1 (C-3'), 85.8 (C-1), 73.0 (C-5), 72.3 (C-3), 70.9 (C-4), 69.5 (C-2), 67.0 (CH₂Ph), 62.7 (C-6), 25.2 (CH₂S), 13.8 (Me). HRESIMS m/z found 447.1072, calcd. For C₂₀H₂₄O₈NaS (M+Na)⁺: 447.1084.

((4-Benzyloxycarbonyl-5-methylfuran-2-yl)methyl)-1-sulfonyl- β -D-galactopyranoside (6): To a solution of **18** (60 mg, 0.096 mmol) in anhydrous MeOH (2 mL), NaOMe (20 μ L, 0.5 M in MeOH) was added and the mixture stirred at 0 °C for 1 h. Then the mixture was neutralized with Amberlite IR-120H⁺, filtered, and washed with MeOH. The filtered solution was concentrated and the residue was purified by column chromatography on silica gel (CH₂Cl₂:MeOH 10:1→8:1) to give **6** (33 mg, 0.072 mmol, 75%) as a white solid. $[\alpha]_D^{25} = -25.0$ (c 0.84, MeOH). IR ($\bar{\nu}$) 3382 (OH), 2920, 1711 (C=O), 1082 cm⁻¹. ¹H NMR (300 MHz, CD₃OD) δ 7.44-7.29 (m, 5H, H-Ar), 6.86 (s, 1H, H-3'), 5.28 (s, 2H, CH₂Ph), 4.76 (d, 1H, $J_{H,H} = 15.0$, CH₂S), 4.49 (d, 1H, CH₂S), 4.33 (d, 1H, $J_{1,2} = 9.5$, H-1), 4.12 (t, 1H, $J_{2,3} = J_{2,1} = 9.4$, H-2), 3.89-3.81 (m, 2H, H-4, H-6a), 3.75-3.64 (m, 2H, H-5, H-6b), 3.55 (dd, 1H, $J_{3,4} = 3.3$, H-3), 2.56 (s, 3H, Me). ¹³C NMR (75.4 MHz, CD₃OD) δ 164.8 (COOBn), 162.0, 142.3 (C-2', C-5'), 137.7 (C_q-Ar), 129.6, 129.3, 129.2 (C-Ar), 115.9 (C-4'), 114.2 (C-3'), 91.2 (C-1), 82.2 (C-5), 75.7 (C-3), 70.2 (C-4), 67.5 (C-2), 67.1 (CH₂Ph), 62.8 (C-6), 50.9 (CH₂S), 13.9 (Me). HRESIMS m/z found 479.0976, calcd. For C₂₀H₂₄O₁₀NaS (M+Na)⁺: 479.0982.

((4-Benzyloxycarbonyl-5-methylfuran-2-yl)methyl)-1-sulfonyl- α -D-galactopyranoside (7): To a solution of **17** (56 mg, 0.090 mmol) in anhydrous MeOH (1.5 mL), NaOMe (18 μ L, 0.5 M in MeOH) was added and the mixture stirred at 0 °C for 2 h. Then the mixture was neutralized with Amberlite IR-120H⁺, filtered, and washed with MeOH. The filtered solution was concentrated to give pure **7** (32 mg, 0.070 mmol, 78%) as a white solid. $[\alpha]_D^{26} = 91.8$ (c 0.98, MeOH). IR ($\bar{\nu}$) 3382 (OH), 2925, 1713 (C=O), 1080 cm⁻¹. ¹H NMR (300 MHz, CD₃OD) δ 7.43-7.29 (m, 5H, H-Ar), 6.80 (s, 1H, H-3'), 5.27 (s, 2H, CH₂Ph), 5.09 (d, 1H, $J_{1,2} = 6.0$, H-1), 4.70 (d, 1H, $J_{H,H} = 14.9$, CH₂S), 4.50 (d, 1H, CH₂S), 4.37-4.24 (m, 3H, H-2, H-3, H-4), 4.00-3.99 (m, 1H, H-5), 3.82 (dd, 1H, $J_{6a,6b} = 11.8$, $J_{6a,5} = 7.3$, H-6a), 3.71 (dd, 1H, $J_{6b,5} = 4.4$, H-6b), 2.55 (s, 3H, Me). ¹³C NMR (75.4 MHz, CD₃OD) δ 164.7 (COOBn), 161.9, 142.6 (C-2', C-5'), 137.6 (C_q-Ar), 129.6, 129.3, 129.2 (C-Ar), 115.8 (C-4'), 114.0 (C-3'), 90.9 (C-1), 78.7, 70.7, 69.2 (C-2, C-3, C-4), 70.0 (C-5), 67.1 (CH₂Ph), 62.6 (C-6), 52.8 (CH₂S), 13.9

(Me). HRESIMS m/z found 479.0971, calcd. For C₂₀H₂₄O₁₀NaS (M+Na)⁺: 479.0982.

((4-Benzyloxycarbonyl-5-methylfuran-2-yl)methyl)-1-sulfinyl- β -D-galactopyranoside (8): To a -78 °C solution of **16** (142 mg, 0.24 mmol) in CH₂Cl₂ (5 mL), MCPBA (59 mg, 0.24 mmol) was added and the mixture stirred at -78 °C→ -30 °C for 3 h. Then, sat. aq. Soln. of NaHCO₃ was added and the mixture was extracted with CH₂Cl₂. The combined organic phases were washed with sat. aq. Soln. of NaHCO₃ and brine, dried over Na₂SO₄, filtered and evaporated. The resulting residue was purified by column chromatography on silica gel (EtOAc:cyclohexane 1:1→1:1) to give **20** (100 mg, 0.164 mmol, 68%) as a white solid. To a solution of this compound (60 mg, 0.099 mmol) in anhydrous MeOH (2 mL), NaOMe (20 μ L, 0.5 M in MeOH) was added and the mixture stirred at 0 °C for 1 h. Then the mixture was neutralized with Amberlite IR-120H⁺, filtered, and washed with MeOH. The filtered solution was concentrated and the residue was purified by column chromatography on silica gel (CH₂Cl₂:MeOH 10:1→8:1) to give **8** (28 mg, 0.064 mmol, 65%) as a white solid. $[\alpha]_D^{27} = -73.5$ (c 0.82, MeOH). IR ($\bar{\nu}$) 3341 (OH), 2929, 1713 (C=O), 1072 cm⁻¹. ¹H NMR (300 MHz, CD₃OD, 1.5:1 mixture of diastereoisomers A:B) δ 7.43-7.28 (m, H-Ar(A), H-Ar(B)), 6.76 (s, H-3'(B)), 6.75 (s, H-3'(A)), 5.31-5.23 (m, CH₂Ph(A), CH₂Ph(B)), 4.42-4.33 (m, H-1(A), CH₂S(A), CH₂S(B)), 4.03-3.55 (m, H-1(B), H-2(A), H-2(B), H-3(A), H-3(B), H-4(A), H-4(B), H-5(A), H-5(B), H-6(A), H-6(B)), 2.56 (s, Me(A)) 2.55 (s, Me(B)). ¹³C NMR (75.4 MHz, CD₃OD, mixture of diastereoisomers) δ 164.8, 164.7 (COOBn(A), COOBn(B)), 161.83, 161.77, 144.7 (C-2'(A), C-2'(B), C-5'(A), C-5'(B)), 137.7 (C_q-Ar), 129.6, 129.3, 129.2 (C-Ar), 115.8 (C-4'), 113.4, 113.2 (C-3'(A), C-3'(B)), 93.8 (C-1(A)), 90.8 (C-1(B)), 82.4, 82.3, 76.3, 76.1, 70.8, 70.3, 68.2, 66.5, (C-2(A), C-2(B), C-3(A), C-3(B), C-4(A), C-4(B), C-5(A), C-5(B)), 67.1 (CH₂Ph), 62.9, 62.8 (C-6(A), C-6(B)), 46.6, 46.1 (CH₂S(A), CH₂S(B)), 13.9 (Me). HRESIMS m/z found 463.1024, calcd. For C₂₀H₂₄O₉NaS (M+Na)⁺: 463.1033.

((4-Benzyloxycarbonyl-5-methylfuran-2-yl)methyl)-1-sulfinyl- α -D-galactopyranoside (9): To a -78 °C solution of **15** (100 mg, 0.169 mmol) in CH₂Cl₂ (3.5 mL), MCPBA (42 mg, 0.17 mmol) was added and the mixture stirred at -78 °C→ -30 °C for 2 h. Then, sat. aq. Soln. of NaHCO₃ was added and the mixture was extracted with CH₂Cl₂. The combined organic phases were washed with sat. aq. Soln. of NaHCO₃, water and brine, dried over Na₂SO₄, filtered and evaporated. The resulting residue was purified by column chromatography on silica gel (Et₂O:cyclohexane 4:1→3:1) to give **19** (66 mg, 0.11 mmol, 65%) as a white solid. To a solution of this compound (20 mg, 0.034 mmol) in anhydrous MeOH (0.7 mL), NaOMe (7 μ L, 0.5 M in MeOH) was added and the mixture stirred at 0 °C for 1 h. Then the mixture was neutralized with Amberlite IR-120H⁺, filtered, and washed with MeOH. The filtered solution was concentrated and the residue was purified by column chromatography on silica gel (EtOAc:MeOH 5:1) to give **9** (11 mg, 0.025 mmol, 74%) as a pale yellow oil. $[\alpha]_D^{27} = 195.5$ (c 0.67, MeOH). IR ($\bar{\nu}$) 3369 (OH), 2923, 1710 (C=O), 1079 cm⁻¹. ¹H NMR (300 MHz, CD₃OD, single diastereoisomer) δ 7.43-7.29 (m, 5H, H-Ar), 6.78 (s, 1H, H-3'), 5.27 (s, 2H, CH₂Ph), 4.75 (d, 1H, $J_{1,2} = 4.2$, H-1), 4.44 (d, 1H, $J_{H,H} = 14.5$, CH₂SO), 4.28 (dd, 1H, $J_{2,3} = 7.7$, H-2), 4.24 (d, 1H, CH₂SO), 4.06 (t, 1H, $J = 2.9$, H-5), 4.00 (dd, 1H, $J_{3,4} = 3.3$, H-3), 3.96-3.88 (m, 2H, H-4, H-6a), 3.76-3.68 (m, 1H, H-6b), 2.56 (s, 3H, Me). ¹³C NMR (75.4 MHz, CD₃OD, single diastereoisomer) δ 164.8 (COOBn), 161.8, 144.6 (C-2', C-5'), 137.7 (C_q-Ar), 129.6, 129.22, 129.19 (C-Ar), 115.7 (C-4'), 113.5 (C-3'), 92.0 (C-1), 80.3 (C-4), 72.3 (C-3), 69.9 (C-2), 68.8 (C-5), 67.1 (CH₂Ph), 61.5 (C-6), 47.8 (CH₂SO), 13.9 (Me). HRESIMS m/z found 463.1024, calcd. For C₂₀H₂₄O₉NaS (M+Na)⁺: 463.1033.

Weak affinity chromatography: Recombinant CTB (SBL vaccines, Stockholm, Sweden) was immobilized onto Nucleosil silica (10 μ m, 300 Å) and packed into a 50x2.1 mm column. The number of active groups of CTB was estimated to be 261 nmol. All chromatographic experiments were performed on an Agilent-1100/Agilent-1200 HPLC system. The mobile phase was 10 mM sodium phosphate, 0.15 mM sodium chloride, pH 7. The flow rate was 0.1 mL min⁻¹ and the temperature 22 °C. The sample

FULL PAPER

volume was 5 μL and the sample concentration 35 μM . Detection was performed at 220 nm. The retention factor (k') and the affinity (K_d) of the derivatives were calculated as described previously,^{[21][30]} by using 3NPG as reference compound (K_d 1.06 ± 0.04 mM).

Nuclear Magnetic Resonance: Ligands **4-9** were assigned by performing COSY, TOCSY and HSQC experiments on a 2 mM solution of the ligands dissolved in D_2O , d6-DMSO 5%. For the STD NMR build-up curves, sample containing 1 mM substrate (Ligands **4-9**) and 5 μM CTB (each pentamer contains 5 equivalent binding sites), *i.e.*, protein:ligand ([P]:[L]) ratio of 1:40 were prepared. The build-up curves were acquired with irradiation frequency 0.60 ppm, at 0.5 s, 1 s, 2 s, 3 s, 4 s and 5s, with a delay between experiment of 5 s and 256 scans. For the DEEP-STD NMR fingerprinting approach and on-resonance scanning of 3 NPG and ligands **4-7**, STD NMR experiments at 2 s saturation time and 32 scans were performed at the irradiation frequencies of 0.60 ppm and 2.25 ppm.

The STD NMR competitions experiments were performed on samples constituted of 1 mM ligand and 5 μM CTB (each pentamer contains 5 equivalent binding sites), *i.e.*, protein:ligand ([P]:[L]) ratio of 1:40. First, the STD NMR experiment of the ligand under investigation (Ligand **2** and **3**) was performed (saturation time 2 s, delay between experiments 5s and 512 scans; the competitor ligand in equimolar concentration was freeze dried and added to the sample, then the experiment was repeated. Exchange-transferred-NOESY (tr-NOESY) experiments were performed with [P]:[L] 1:10 on the ternary complex of Ligand **3** and 3NPG with CTB (1 mM of each ligand and 20 μM CTB), with mixing time 1.2 s, delay 2 s and 160 scans. As a control, the same experiment was performed on 3NPG/CTB, Ligand **3**/CTB and CTB alone in the same conditions. All the NMR experiments were performed at 278 K.

An STD pulse sequence that included 2.5 ms and 5 ms trim pulses and a 3 ms spoil gradient and water suppression by excitation sculpting with gradients was used (*stdiffesp.3*). Saturation was achieved applying a train of 50 ms Gaussian pulses (0.40 mW) on the f_2 channel, at 0 ppm or 7.27 ppm (on-resonance experiments) and 40 ppm (off-resonance experiments). The broad protein signals were removed using a 40 ms spinlock ($T_{1\rho}$) filter (*stdiff.3*). Tr-NOESY experiments were performed using a phase-sensitive pulse program with gradient pulses in the mixing time and a relaxation delay of 2 s, with water suppression using 3-9-19 pulse sequence with gradients (*noesyfpgpph19*).^{[34],[35],[36]} All the experiments were recorded at ^1H frequency of 800.23 MHz on a Bruker Avance III spectrometer equipped with a 5-mmD probe TXI 800 MHz H-C/N-D-05 Z BTO.

Protein-Ligand docking calculations: All molecular modelling was performed with the module Glide within Schrödinger's Maestro modelling suite^{[24],[25],[26]}. Coordinates for CTB were obtained from the Protein Data Bank (3CHB) for the closed conformation and from a representative frame of our HREMD trajectory of the GM1/CTB complex for the open conformation. Where necessary, coordinates for missing atoms were added according to known protein chemistry and sidechain protonation was optimized for neutral pH. A short energy minimization was run using the OPLS3 force field, converging heavy atoms to a RMSD of 0.3 Å. The receptor grid was then calculated, centering on the centroid of GM1 and with a length of 30 Å in order to encompass all subsites. Three-dimensional structures of all ligands were generated using a conformational search, implementing Monte-Carlo torsional sampling, keeping only unique structures (RMSD > 0.5 Å) and eliminating all structure with an energy 21 kJ mol^{-1} greater than the lowest energy structure. All resulting structures were then energy minimized using conjugate gradient minimization, converging on a threshold of 0.05 $\text{kJ mol}^{-1} \text{Å}^{-1}$. For each ligand, the 10 lowest energy conformations were used to initiate docking. The docking consisted of further conformer sampling (generation of ring conformations switched off, to avoid potential generation of distorted galactopyranose ring conformations out of the well-known low energy range around the $^4\text{C}_1$ conformation), docking and then energy minimization. Conformers were generated with 4x enhanced sampling and, during docking, the non-

bonded term of the potential energy function was softened for nonpolar ligand atoms (charge < |0.15|) by applying a scaling factor of 0.8. Finally, energy minimization was performed using implicit solvent with a distance dependent dielectric constant of 4.

Molecular dynamics: Derivation of Charges The charges of all protein and carbohydrate atoms were obtained from AMBER ff14SB and GLYCAM_06j libraries respectively. The charges of organics moieties were derived by using the RESP fitting method implemented on the RED server^[37], following a similar protocol to previous publications^{[38],[39],[40]}. The non-carbohydrate regions of both ligands **2** and **3** were each split into two fragments in order to increase computational efficiency, by disconnecting the bond between the sulphur and the anomeric carbon. In place of this bond, both ends were capped with a methyl group. Charges of individual fragments were computed using the HF/6-31G* level of theory with a weight factor of 0.01. All aliphatic protons were constrained to a charge of 0. During the fitting, the combined charge of the starred methyl and hydroxyl groups was set to zero, removing these groups in the final fitting and merging the fragments to give the full-length non-carbohydrate moiety. For ligand **2** the charge of the sulphide methyl group was set to 0.194 throughout the fitting stages before removing to give a final fragment with net charge -0.194, in keeping with the modularity of the GLYCAM forcefield used for galactose charges.

Molecular Dynamics: Initial Coordinates. Initial coordinates for simulations of free CTB and the GM1/CTB complex were obtained from the Protein Data Bank (3CHB). Unresolved atoms were added to the coordinate files using *tleap*, according to known protein chemistry and protonation state at physiological pH. Coordinates for simulations involving ligands **2** and **3** were generated by molecular docking as described above.

Molecular Dynamics: Simulation Protocol. All systems were parameterized using AMBER ff14SB, GLYCAM_06j and GAFF forcefields for the protein, carbohydrate and organic moieties respectively. Model systems were generated by solvating with explicit TIP4PEW water molecules within a truncated octahedral bounding box such that no solute atom was less than 10 Å from any face. An appropriate number of chloride ions were added to each system in order to neutralize the total charge. Conjugate gradient energy minimization was run with 100 $\text{kcal mol}^{-1} \text{Å}^{-2}$ restraints on solute atoms, converging on a threshold of $1 \times 10^{-4} \text{ kcal mol}^{-1} \text{Å}^{-1}$, before repeating with no restraints. The system was then heated at constant volume to 300 K over a period of 100 ps using a weak coupling algorithm with a time constant of 1 ps, employing periodic boundary conditions (PBC) with the particle mesh Ewald (PME) method. A constant temperature and pressure (1 atm) simulation was then run for 200 ps, using PBC with PME, to equilibrate the solvent density. A Langevin thermostat with a collision frequency of 2 ps^{-1} and a Berendsen barostat with a relaxation time of 2 ps were used. The SHAKE algorithm was used to restrain all bonds involving hydrogen, allowing a timestep of 2 fs to be used. A cutoff of 8 Å was used for non-bonded interactions at all simulation stages. Using the same settings as for equilibration, classical molecular dynamics simulations were run for 100ns, saving trajectory coordinates every 5 ps. For HREMD simulations, systems were prepared as above, differing only in the production dynamics stage. 8 replicas were generated from the same initial coordinates following equilibration and were each run for 12.5 ns. The non-bonded potential between solvent atoms and non-polar protein atoms (defined as carbon and sulphur) were scaled by a factor between 1 and 1.35 at intervals of 0.05 and exchanges between replicas was allowed every 2 ps. All trajectories were analyzed with *cptraj*. All molecular graphics were generated with Schrödinger Maestro 11 version 2016-4.

CORCEMA-ST predictions:^{[28],[41]} Based on the experimental conditions, the concentration of ligand was 1 mM and the [P]:[L] ratio was kept fixed at 25:1. The cut distance around the binding pocket was 13 Å. k_{on} was set to $1 \times 10^{-8} \text{ M}^{-1}$ and the irradiation frequency to the range -0.8 ppm to 0.8 ppm. The bound ligand correlation time (τ) was 65 ns for both ligands, whereas the free correlation time was 1.2 ns, 1 ns and 1 ns respectively

FULL PAPER

for ligands **2**, **1** and **3**. The equilibrium constant used for ligand **2** was 2000 M⁻¹, 1000 M⁻¹ for ligand **1** and 1500 M⁻¹ for ligand **3**, with p leak of 0.25. For ligands **2** and **3**, CORCEMA-ST calculations were run in parallel on the High-Performance Computing Unit, and then averaged to provide a representative build up curve for each ligand. The NOE R-factor was calculated on the averaged simulated data obtained accordingly.

Acknowledgements


This work was supported by the Biotechnology and biological Sciences Research Council (BBSRC) through a New Investigator grant awarded to J.A. (BB/P010660/1). S.M, R.N., and S.W. also acknowledge support from BBSRC through the same grant and a DTP studentship to S.W. J.A. also acknowledges financial support from the Universidad de Sevilla (Acciones Especiales del VI Plan Propio de Investigación y Transferencia). This work was supported by the Ministerio de Economía y Competitividad of Spain (CTQ2012-31247, CTQ2016-77270-R) and the Junta de Andalucía (FQM-345). We thank CITIUS-University of Seville (MS and NMR services) and S. Carrión Jiménez for technical assistance. We also thank Prof. Sten Ohlson for helpful discussions.

Keywords: Multi-subsite binding pockets • DEEP-STD NMR • Cholera toxin inhibitors • Protein-Ligand Interactions • Ligand-based NMR

- [1] D. A. Erlanson, S. W. Fesik, R. E. Hubbard, W. Jahnke and H. Jhoti, *Nat. Rev. Drug Discov.* **2016**, *15*, 605.
- [2] P. J. Hajduk, J. C. Mack, E. T. Olejniczak, C. Park, P. J. Dandliker and B. A. Beutel, *J. Am. Chem. Soc.* **2004**, *126*, 2390-2398.
- [3] P. Hajduk, G. Sheppard, D. Nettekheim, E. Olejniczak, S. Shuker, R. Meadows, D. Steinman, G. Carrera, P. Marcotte and J. Severin, *J. Am. Chem. Soc.* **1997**, *119*, 5818-5827.
- [4] L. Unione, S. Galante, D. Díaz, F. J. Cañada and J. Jiménez-Barbero, *Med. Chem. Commun.* **2014**, *5*, 1280-1289.
- [5] B. Becattini and M. Pellecchia, *Chem. Eur. J.* **2006**, *12*, 2658-2662.
- [6] a) M. Mayer and B. Meyer, *Angew. Chem. Int. Ed.* **1999**, *38*, 1784-1788; b) M. Mayer and B. Meyer, *J. Am. Chem. Soc.* **2001**, *123*, 6108-6117.
- [7] O. Cala and I. Krimm, *J. Med. Chem.* **2015**, *58*, 8739-8742.
- [8] S. Monaco, L. E. Tailford, N. Juge and J. Angulo, *Angew. Chem. Int. Ed.* **2017**, *56*, 15289-15293.
- [9] J. E. Watt, G. R. Hughes, S. Walpole, S. Monaco, G. R. Stephenson, P. C. Bulman Page, A. M. Hemmings, J. Angulo and A. J. C. A. E. J. Chantry, *Chem. Eur. J.* **2018**, *24*, 17677-17680.
- [10] G. M. Kuziemko, M. Stroh and R. C. Stevens, *Biochemistry* **1996**, *35*, 6375-6384.
- [11] J. Harris, *Cholera. Lancet* **2012**, *379*, 2466-2476.
- [12] K. J. Baldauf, J. M. Royal, K. T. Hamorsky and N. Matoba, *Toxins* **2015**, *7*, 974-996.
- [13] W. B. Turnbull, B. L. Precious and S. W. Homans, *J. Am. Chem. Soc.* **2004**, *126*, 1047-1054.
- [14] E. A. Merritt, S. Sarfaty, F. V. D. Akker, C. L'Hoir, J. A. Martial and W. G. Hol, *Protein Sci.* **1994**, *3*, 166-175.
- [15] P. Brocca, P. Berthault and S. Sonnino, *Biophys. J.* **1998**, *74*, 309-318.
- [16] D. Arosio, S. Baretta, S. Cattalio, D. Potenza and A. Bernardi, *Bioorg. Med. Chem. Lett.* **2003**, *13*, 3831-3834.
- [17] A. Bernardi, L. Raimondi and F. Zuccotto, *J. Med. Chem.* **1997**, *40*, 1855-1862.
- [18] A. Bernardi, D. Arosio, L. Manzoni, D. Monti, H. Poster, D. Potenza, S. Mari and J. Jiménez-Barbero, *Org. Biomol. Chem.* **2003**, *1*, 785-792.
- [19] T. R. Branson and W. B. Turnbull, *Chem. Soc. Rev.* **2013**, *42*, 4613-4622.
- [20] Y. Maeda, K. Motoyama, T. Higashi, Y. Horikoshi, T. Takeo, N. Nakagata, Y. Kurauchi, H. Katsuki, Y. Ishitsuka and Y. Kondo, *J. Pharm. Pharmacol.* **2015**, *67*, 1133-1142.
- [21] J. Ramos - Soriano, U. Niss, J. Angulo, M. Angulo, A. J. Moreno - Vargas, A. T. Carmona, S. Ohlson and I. Robina, *Chem. Eur. J.* **2013**, *19*, 17989-18003.
- [22] C. Dalvit, M. Flocco, S. Knapp, M. Mostardini, R. Perego, B. J. Stockman, M. Veronesi and M. Varasi, *J. Am. Chem. Soc.* **2002**, *124*, 7702-7709.
- [23] B. Becattini and M. Pellecchia, *Chem. Eur. J.* **2006**, *12*, 2658-2662.
- [24] R. A. Friesner, R. B. Murphy, M. P. Repasky, L. L. Frye, J. R. Greenwood, T. A. Halgren, P. C. Sanschagrin and D. T. Mainz, *J. Med. Chem.* **2006**, *49*, 6177-6196.
- [25] T. A. Halgren, R. B. Murphy, R. A. Friesner, H. S. Beard, L. L. Frye, W. T. Pollard and J. L. Banks, *J. Med. Chem.* **2004**, *47*, 1750-1759.
- [26] R. A. Friesner, J. L. Banks, R. B. Murphy, T. A. Halgren, J. J. Klicic, D. T. Mainz, M. P. Repasky, E. H. Knoll, M. Shelley and J. K. Perry, *J. Med. Chem.* **2004**, *47*, 1739-1749.
- [27] V. Oleinikovas, G. Saladino, B. P. Cossins and F. L. Gervasio, *J. Am. Chem. Soc.* **2016**, *138*, 14257-14263.
- [28] V. Jayalakshmi and N. R. Krishna, *J. Magn. Reson.* **2002**, *155*, 106-118.
- [29] K. Yamamoto, N. Watanabe, H. Matsuda, K. Oohara, T. Araya, M. Hashimoto, K. Miyairi, I. Okazaki, M. Saito, T. J. B. Shimizu, *Bioorg. Med. Chem. Lett.* **2005**, *15*, 4932-4935.
- [30] M. Bergström, S. Liu, K. L. Kiick and S. Ohlson, *Chem. Biol. Drug Des.* **2009**, *73*, 132-141.
- [31] E. Fan, E. A. Merritt, Z. Zhang, J. C. Pickens, C. Roach, M. Ahn and W. G. Hol, *Acta Crystallogr. D* **2001**, *57*, 201-212.
- [32] E. A. Merritt, P. Kuhn, S. Sarfaty, J. L. Erbe, R. K. Holmes and W. G. Hol, *J. Mol. Biol.* **1998**, *282*, 1043-1059.
- [33] A. Bernardi, D. Arosio, D. Potenza, I. Sánchez - Medina, S. Mari, F. J. Cañada and J. Jiménez - Barbero, *Chem. Eur. J.* **2004**, *10*, 4395-4406.
- [34] M. Piotto, V. Saudek and V. Sklenář, *J. Biomol. NMR* **1992**, *2*, 661-665.
- [35] V. Sklenar, M. Piotto, R. Leppik and V. Saudek, *J. Magn. Reson., Series A* **1993**, *102*, 241-245.
- [36] V. Sklenar, *J. Magn. Reson., Series A* **1995**, *114*, 132-135.
- [37] E. Vanqualef, S. Simon, G. Marquant, E. Garcia, G. Klimerek, J. C. Delepine, P. Cieplak and F.-Y. Dupradeau, *Nucleic Acids Res.* **2011**, *39*, W511-W517.
- [38] J. C. Muñoz-García, E. Chabrol, R. R. Vivès, A. Thomas, J. L. De Paz, J. Rojo, A. Imberty, F. Fieschi, P. M. Nieto and J. Angulo, *J. Am. Chem. Soc.* **2015**, *137*, 4100-4110.
- [39] N. Sapay, E. Cabannes, M. Petitou and A. Imberty, *Glycobiology* **2011**, *21*, 1181-1193.
- [40] S. G. Gouin, E. Vanqualef, J. M. García Fernández, C. Ortiz Mellet, F.-Y. Dupradeau and J. Kovensky, *J. Org. Chem.* **2007**, *72*, 9032-9045.
- [41] V. Jayalakshmi and N. R. Krishna, *J. Magn. Reson.* **2004**, *168*, 36-45.

FULL PAPER

Entry for the Table of Contents



Fingerprinting multi-subsite binding pockets. A powerful protocol using NMR and Molecular Dynamics to study multi-subsite binding pockets has been developed. For cholera toxin subunit B (CTB), the combination of DEEP-STD NMR, STD competition experiments, transfer-NOESY and HREMD unveiled the presence of a novel subsite adjacent to the well-known subsites of GM1 on CTB.

Institute and/or researcher Twitter usernames: @Angulo_Lab (Jesús Angulo)