

# Draft Genome Sequence of a Necrotoxicogenic *Escherichia coli* Isolate

Sadjia Bekal,<sup>a,b</sup> Alex Lin,<sup>a</sup> André Vincent,<sup>c</sup> Chrystal Berry,<sup>d</sup> Matthew Gilmour,<sup>d</sup> Éric Fournier,<sup>a</sup> Jean-Charles Côté,<sup>a</sup> Cécile Tremblay<sup>a,b</sup>

Laboratoire de santé publique du Québec, Sainte-Anne-de-Bellevue, Québec, Canada<sup>a</sup>; Département de Microbiologie, infectiologie et immunologie, Université de Montréal, Montréal, Québec, Canada<sup>b</sup>; Centre Hospitalier Affilié Universitaire, Hôtel-Dieu de Lévis, Lévis, Québec, Canada<sup>c</sup>; Public Health Agency of Canada, National Microbiology Laboratory, Winnipeg, Manitoba, Canada<sup>d</sup>

Here, we present the draft genome sequence of a necrotoxicogenic *Escherichia coli* strain isolated from a patient following a very rapidly evolving, lethal necrotizing fasciitis.

Received 20 August 2015 Accepted 21 August 2015 Published 1 October 2015

Citation Bekal S, Lin A, Vincent A, Berry C, Gilmour M, Fournier É, Côté J-C, Tremblay C. 2015. Draft genome sequence of a necrotoxicogenic *Escherichia coli* isolate. *Genome Announc* 3(5):e01152-15. doi:10.1128/genomeA.01152-15.

Copyright © 2015 Bekal et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/4.0/).

Address correspondence to Sadjia Bekal, Sadjia.Bekal@inspq.qc.ca.

*Escherichia coli* is a bacterial commensal of the intestinal tract of warm-blooded animals, including humans. It is Gram-negative, facultative anaerobe, and rod-shaped (1). Although most *E. coli* strains are harmless, several others are the etiological agents of various diseases. The latter can be divided into intestinal (diarrheagenic) and extraintestinal pathogenic *E. coli* (ExPEC) strains (2, 3). The intestinal pathogenic group comprises at least six pathotypes: enteropathogenic (EPEC), Shiga-toxin producing enterohemorrhagic (STEC/EHEC), enterotoxigenic (ETEC), enteroaggregative (EAEC), enteroinvasive (EIEC), and diffusely adherent *E. coli* (DAEC). The ExPEC comprises at least three named pathotypes: uropathogenic (UPEC), newborn meningitis-causing *E. coli* (MENEC), an unnamed pathotype which encompasses the strains that cause septicemia in humans and animals, and necrotoxicogenic *E. coli* (NTEC) (2, 3). A characteristic cytotoxic necrotizing factor (Cnf1 or Cnf2) is synthesized by uropathogenic and necrotoxicogenic *E. coli* strains (4, 5).

Necrotizing fasciitis, commonly referred to as “flesh-eating disease” is a rare but severe disease characterized by the necrosis of the subcutaneous tissues and fascia (6, 7). It is usually caused by *Streptococcus pyogenes* (8, 9), sometimes *Staphylococcus aureus* (10) or by a mixture of microorganisms including *Streptococcus*, *S. aureus*, *Enterobacteriaceae*, and some anaerobes (11, 12). On rare occasions, necrotoxicogenic *E. coli* was identified as the etiologic agent in chronically ill patients (13–20) or infants following surgery (21, 22). Recently, we reported a lethal case of very rapidly evolving necrotizing fasciitis. The bacterial isolate was identified as *E. coli* and designated LSPQ A134697 (Laboratoire de Santé Publique du Québec, strain A134697). DNA microarray revealed the presence of several toxin genes, including the cytotoxic necrotizing factor 1 gene *cnf1* (23).

We present here the draft genome sequence of LSPQ A134697. A sequencing library was prepared using the Nextera XT library preparation kit (Illumina, Inc., San Diego, CA, United States). Sequencing was performed on an Illumina MiSeq platform with the MiSeq Reagent Kit v2, 500 cycles to achieve 83× average genome coverage. The quality of the raw sequence data was checked using FastQC (<http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>). The sequence reads were *de novo* assembled into contigs using SPAdes

v3.5 (<http://bioinf.spbau.ru/spades> [24]). A total of 81 contigs, ranging in size from 505 bp to 621,560 bp, with a  $N_{50}$  of 244,924 bp, for a total length of 5,228,778 bp with an average G+C content of 50.6% was generated. They were annotated using the NCBI Prokaryotic Genome Automatic Annotation Pipeline v2.10 (PGAAP) (25). A total of 5,126 protein-coding sequences, 81 tRNAs, 17 rRNAs (5S, 16S, and 23S), and 81 pseudogenes are predicted. The *cnf1* gene is located on contig 24, at positions 42,141 to 45,185. Further analysis of this genome and comparison with others will be presented elsewhere.

This is the first annotated draft genome sequence of a necrotoxicogenic *E. coli* isolate.

**Nucleotide sequence accession number.** The annotated genome sequence was deposited in GenBank under accession no. [LELX00000000](https://www.ncbi.nlm.nih.gov/nuclseq/LELX00000000).

## ACKNOWLEDGMENTS

This work was supported in-house by Laboratoire de Santé Publique du Québec and by the National Microbiology Laboratory.

We thank Matthew Walker for excellent technical assistance.

We have no conflict of interest.

## REFERENCES

- Neidhardt FC, Curtiss R (ed). 1996. *Escherichia coli* and *Salmonella*: cellular and molecular biology. ASM Press, Washington, DC.
- Kaper JB, Nataro JP, Mobley HL. 2004. Pathogenic *Escherichia coli*. *Nat Rev Microbiol* 2:123–140. <http://dx.doi.org/10.1038/nrmicro818>.
- Russo TA, Johnson JR. 2000. Proposal for a new inclusive designation for extraintestinal pathogenic isolates of *Escherichia coli*: ExPEC. *J Infect Dis* 181:1753–1754. <http://dx.doi.org/10.1086/315418>.
- Boquet P. 2001. The cytotoxic necrotizing factor 1 (CNF1) from *Escherichia coli*. *Toxicon* 39:1673–1680. [http://dx.doi.org/10.1016/S0041-0101\(01\)00154-4](http://dx.doi.org/10.1016/S0041-0101(01)00154-4).
- De Rycke J, González EA, Blanco J, Oswald E, Blanco M, Boivin R. 1990. Evidence for two types of cytotoxic necrotizing factor in human and animal clinical isolates of *Escherichia coli*. *J Clin Microbiol* 28:694–699.
- Anaya DA, Dellinger EP. 2007. Necrotizing soft-tissue infection: diagnostic and management. *Clin Infect Dis* 44:705–710. <http://dx.doi.org/10.1086/511638>.
- Angoules AG, Kontakis G, Drakoulakis E, Vrentzos G, Granick MS, Giannoudis PV. 2007. Necrotizing fasciitis of upper and lower limb: a systematic review. *Injury* 38:S19–S26. <http://dx.doi.org/10.1016/j.injury.2007.10.030>.

8. Bisno AL, Stevens DL. 1996. Streptococcal infections of skin and soft tissues. *N Engl J Med* 334:240–245. <http://dx.doi.org/10.1056/NEJM199601253340407>.
9. Johansson L, Thulin P, Low DE, Norrby-Teglund A. 2010. Getting under the skin: the immunopathogenesis of *Streptococcus pyogenes* deep tissue infections. *Clin Infect Dis* 51:58–65. <http://dx.doi.org/10.1086/653116>.
10. Cheng NC, Wang JT, Chang SC, Tai HC, Tang YB. 2011. Necrotizing fasciitis caused by *Staphylococcus aureus*: the emergence of methicillin-resistant strains. *Ann Plast Surg* 67:632–636. <http://dx.doi.org/10.1097/SAP.0b013e31820b372b>.
11. Elliott D, Kufera JA, Myers RA. 2000. The microbiology of necrotizing soft tissue infections. *Am J Surg* 179:361–366. [http://dx.doi.org/10.1016/S0002-9610\(00\)00360-3](http://dx.doi.org/10.1016/S0002-9610(00)00360-3).
12. Wong CH, Chang HC, Pasupathy S, Khin LW, Tan JL, Low CO. 2003. Necrotizing fasciitis: clinical presentation, microbiology, and determinants of mortality. *J Bone Joint Surg Am* 85-A:1454–1460.
13. Castanet J, Lacour JP, Perrin C, Bodokh I, Dor JF, Ortonne JP. 1992. *Escherichia coli* cellulitis: two cases. *Acta Derm Venereol* 72:310–311.
14. Horowitz Y, Sperber AD, Almog Y. 2004. Gram-negative cellulitis complicating cirrhosis. *Mayo Clin Proc* 79:247–250. <http://dx.doi.org/10.4065/79.2.247>.
15. Li DM, Lun LD, Chen XR. 2006. Necrotising fasciitis with *Escherichia coli*. *Lancet Infect Dis* 6:456. [http://dx.doi.org/10.1016/S1473-3099\(06\)70526-6](http://dx.doi.org/10.1016/S1473-3099(06)70526-6).
16. Yoon TY, Jung SK, Chang SH. 1998. Cellulitis due to *Escherichia coli* in three immunocompromised subjects. *Br J Dermatol* 139:885–888. <http://dx.doi.org/10.1046/j.1365-2133.1998.02519.x>.
17. Liu CT, Chen YC, Chen TH, Ou TY. 2012. Necrotizing fasciitis of thigh associated with *Escherichia coli* bacteremia in a patient on chronic hemodialysis. *Hemodial Int* 16:564–567. <http://dx.doi.org/10.1111/j.1542-4758.2011.00658.x>.
18. Tsai MH, Leu JG, Fang Y, Hsieh SC. 12 January 2015. Necrotizing fasciitis and infective endocarditis caused by *Escherichia coli* in a hemodialysis patient. *Hemodial Int*. <http://dx.doi.org/10.1111/hdi.12264>.
19. Grimaldi D, Bonacorsi S, Roussel H, Zuber B, Poupet H, Chiche J-D, Poyart C, Mira J-P. 2010. Unusual “flesh-eating” strain of *Escherichia coli*. *J Clin Microbiol* 48:3794–3796.
20. Shaked H, Samra Z, Paul M, Madar-Shapiro L, Cohen J, Pitlik S, Bishara J. 2012. Unusual “flesh-eating” Strains of *Escherichia coli*. *J Clin Microbiol* 50:4008–4011. <http://dx.doi.org/10.1128/JCM.02316-12>.
21. Alsaif MA, Robinson JL. 2015. *Escherichia coli* necrotizing fasciitis in Hirschsprung’s disease. *J Pediatr Surg Case Rep* 3:174–175. <http://dx.doi.org/10.1016/j.epsc.2015.03.002>.
22. Barker L, Pringle K, Cusack J. 2013. Images in paediatrics: necrotizing fasciitis with *Escherichia coli* in a newborn infant after abdominal surgery. *Arch Dis Child Fetal Neonatal Ed* 98:F404. <http://dx.doi.org/10.1136/archdischild-2013-303771>.
23. Bekal S, Vincent A, Lin A, Harel J, Côté JC, Tremblay C. 2015. A case of fatal necrotizing fasciitis caused by a highly virulent *Escherichia coli* strain. *J Med Microbiol Case Rep*, in press.
24. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshtkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <http://dx.doi.org/10.1089/cmb.2012.0021>.
25. Tatusova T, DiCuccio M, Badretdin A, Ciufu S, Li W. 2013. Prokaryotic genome annotation pipeline. In *The NCBI handbook* [Internet], 2nd ed. National Center for Biotechnology Information, Bethesda, MD, <http://www.ncbi.nlm.nih.gov/books/NBK174280/>.