

TITLE PAGE

Title

The association between psychosocial factors and mental health symptoms to cervical spine pain with or without radiculopathy on health outcomes: systematic review protocol.

Author names and affiliations

Michael Mansfield MSc BSc (Hons) ¹

¹ School of Health and Social Care, Division of Allied Health Sciences, London South Bank University, London, United Kingdom.

Email: Michael.Mansfield@lsbu.ac.uk

Twitter: @MM_Physio

Tel: 020 7815 7815

Nicolas Spahr PhD MSc MA Grad Dip Phys ^{2 3}

² Guy's and St Thomas Hospital NHS Foundation Trust, Physiotherapy Department, St Thomas Hospital, Westminster Bridge Road, London, United Kingdom.

³ Pain Section, Neuroimaging. Institute of Psychiatry. Kings College London, United Kingdom.

Email: Nicolas.Spahr@gstt.nhs.uk

Twitter: @NicolasSpahr

Toby Smith PhD MA MSc BSc (Hons) ^{4 5}

⁴ Nuffield Department of Orthopaedics, Rheumatology and Musculoskeletal Sciences, University of Oxford, Botnar Research Centre, Oxford, United Kingdom.

⁵ Faculty of Medicine and Health Sciences, University of East Anglia, Norwich NR4 7TJ, United Kingdom.

Email: Toby.Smith@ndorms.ox.ac.uk; Toby.Smith@uea.ac.uk

Twitter: @TobyOSmith

Brendon Stubbs PhD MSc BSc (Hons) ^{6 7}

⁶ Department of Psychological Medicine, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London SE5 8AF, United Kingdom.

⁷ South London and Maudsley NHS Foundation Trust, London, United Kingdom.

Email: Brendon.stubbs@kcl.ac.uk

Twitter: @BrendonStubbs

Lesley Haig D.PT MSc Grad Dip Phys ⁸

⁸ AECC University College, Parkwood Campus, Parkwood Rd, Bournemouth BH5 2DF, United Kingdom.

Email: LHaig@aecc.ac.uk

Twitter: @ProfLesleyHaig

Michael Thacker PhD MSc Grad Dip Phys Grad Dip MNMSD ¹

¹ School of Health and Social Care, Division of Allied Health Sciences, London South Bank University, London, United Kingdom.

Email: Michael.Thacker@lsbu.ac.uk

Twitter: @dibbygibby

Corresponding author:

Michael Mansfield MSc BSc (Hons)

School of Health and Social Care, Division of Allied Health Sciences, London South Bank University, London, United Kingdom.

Email: Michael.Mansfield@lsbu.ac.uk

Twitter: @MM_Physio

Senior author:

Michael Thacker PhD MSc Grad Dip Phys Grad Dip MNMSD ¹

¹ School of Health and Social Care, Division of Allied Health Sciences, London South Bank University, London, United Kingdom.

Email: Michael.Thacker@lsbu.ac.uk

Twitter: @dibbygibby

Funding & Acknowledgement:

Nil funding.

This study forms part of Michael Mansfield's PhD study programme.

Conflict of Interest:

There are no conflicts of interest.

Significance Statement:

This is a first systematic review protocol investigating the association of psychosocial factors and mental health symptoms and outcomes in adults with cervical spine pain with or without upper limb radiculopathy. The findings of this review are intended to enhance assessment and management decisions of clinicians and healthcare services for this complex and disabling condition.

Key Words:

Cervical spine pain; Cervical spine radiculopathy; Systematic review; Psychosocial; Mental health.

PROSPERO Registration Number:

CRD42020169497

Author Contributions:

Conceptualisation: Michael Mansfield and Mick Thacker

Methodology: Michael Mansfield, Toby Smith, Nicolas Spahr, Brendon Stubbs, Lesley Haig and Mick Thacker

Project administration: Michael Mansfield

Writing and original draft: Michael Mansfield

Writing, review and editing: Michael Mansfield, Toby Smith, Nicolas Spahr, Brendon Stubbs, Lesley Haig and Mick Thacker

MANUSCRIPT

Introduction

Cervical spine pain is characterised as pain perceived anywhere in the posterior region of the cervical spine, from the superior nuchal line to the first thoracic spinous process [5]. Any lesion causing a mechanical compromise to a spinal nerve or its root can cause radiculopathy directly through compression or indirectly through the interruption of the blood supply or nutrition to a nerve axon or its root [44]. Cervical spine radiculopathy is identified as an objective loss of sensory and/or motor function as a result of conduction block to a spinal nerve or its root, this can occur with or without cervical spine pain [14]. People with CSp±R may also describe lancinating symptoms along narrow bands, which is similar but not identical to “dermatomal distribution”, this is known as radicular pain [16, 40].

Cervical Spine Pain with or without radiculopathy (CSp±R) has significant negative impacts to a person’s physical and mental health and wellbeing and is an enormous burden for individuals, families and societies [7, 11, 19]. The one year incidence of cervical spine pain ranges between 10.4% and 21.3% [20, 21] and Cervical spine radiculopathy prevalence values range between 1.1% to 6.3% [34, 37]. CSp±R is a leading cause of years lived with disability [30]. The global prevalence of cervical spine pain and years lived with disability has increased by 18.7% and 18.6% respectively over the last 10 years [30].

Multiple systematic reviews have established presence of psychological and/or mental health symptoms are associated with low back pain and negatively impact health outcomes and quality of life [33, 47]. This is not yet established for CSp±R and warrants further attention. Psychosocial factors encompass a wide range of characteristics such as patient's beliefs, emotions, behaviours, and family and workplace aspects [27]. Mental health symptoms or conditions are an extension of such factors. Examples of mental health conditions include major depressive disorders and anxiety. These conditions are among the leading causes of years lived with disability globally and less than a 10% positive improvement between 1990-2016 [30]. Stress, anxiety, depression and negative coping behaviours can adversely impact prognosis with musculoskeletal conditions such as low back pain [35], neck pain [39, 42], knee osteoarthritis [43, 49], carpal tunnel syndrome [18] and shoulder pain [8]. Psychosocial factors and/or mental health symptoms should be considered as part of a clinical reasoning framework to positively affect health outcomes and support prognosis [25].

Persistent pain and associative mental health symptoms are well established in low back pain [15, 32, 35]. Up to 20% of people with spinal pain, including CSp±R, will experience symptoms associated with depression and anxiety [3, 26]. Pain and disability associated with CSp±R limits a person's participation in social activities, working life and consequently negatively impacting quality of life [10, 31, 48].

Worryingly, there can be a devastating impact to a person's lived experience with CSp±R. People report their 'life is on hold', question their 'life purpose' and consider suicidal thoughts [36]. This potentially distressing association between CSp±R and

psychosocial factors or mental health symptoms requires enhanced attention from researchers, clinicians and healthcare services.

To date, no systematic review has examined the association between psychosocial factors and/or mental health symptoms in adults with CSp±R on health outcomes. A robust systematic review will enhance understanding and improve quality of healthcare services for these burdensome conditions. Therefore, our objective is to conduct a systematic review to assess the association between psychosocial factors and/or mental health symptoms and health outcomes in adults with CSp±R.

Methods

Protocol

The Preferred Reporting Items for Systematic Review and Meta-Analysis Protocols (PRISMA-P) [29] checklist guided the development and reporting of this review protocol. The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement [28] will be used on reporting the findings. The protocol for this systematic review has been registered with the International Prospective Register of Systematic Reviews (PROSPERO) database (CRD42020169497).

Eligibility criteria – Population

This systematic review will retrieve studies with samples of adults aged 18 years and over with CSp±R. It is anticipated that there will be variance in the definition of CSp±R across geographical location, clinical settings and from a historical standpoint. For example, The International Association of the Study of Pain [22] define cervical spine pain as pain perceived anywhere in the posterior region of the cervical spine, from the superior nuchal line to the first thoracic spinous process (Merskey and Bogduk, 1994). Whereas the Bone and Joint Decade 2000-2010 Task Force on Neck Pain describe cervical spine pain as the posterior neck region from the superior nuchal line to the spine of the scapula and the side region down to the superior border of the clavicle and the suprasternal notch [31].

Two reviewers will review all potentially eligible studies and a consensus decision will be sought on the CSp±R diagnosis. The CSp±R diagnosis criteria of all included studies will be described. From our previous work (Mansfield et al., *unpublished*) we anticipate a variance in cervical spine radiculopathy diagnostic criteria. Therefore this review we will take a pragmatic approach and include studies with *probable* or *definite* cervical spine radiculopathy diagnosis adapted from IASP and North American Spine Society [6, 22, 38] (**Figure 1**). Patients with CSp±R as a result of an upper motor neuron lesion, fracture, radiculitis, myelopathy, post-surgery, whiplash associated disorder (WAD), systemic pathology or metabolic diseases such as diabetes (including pre-diabetes) will be excluded.

Eligibility criteria - Exposure

Studies will be included if they assess psychosocial factors or mental health symptoms as an exposure. Studies must investigate one or more psychosocial or mental health symptoms (or condition). Psychosocial factors including cognitive (e.g. neuropsychological functioning), affective (e.g. distress, mood), behavioural (e.g. coping strategies), vocational (e.g. job satisfaction, self-perceived work ability) or interpersonal processes (e.g. social support) (Upton, 2013) will be considered. Mental health symptoms and conditions such as depressive symptoms, clinical depression, anxiety symptoms, perceived stress, personality, psychotic, traumatic and/or eating disorders will also be considered. Self-reported, objective standardised questionnaires (e.g. Beck Depression Index, Karasek's Job Control Questionnaire, GHQ-12) and/or psychosocial factors or mental health symptoms using dichotomous data ("yes/no") will be considered.

Eligibility criteria - Comparators

Studies will also be considered if the study population compares different severities of mental health symptoms or conditions or psychosocial factors. The “cut-off” values for mental health condition severity used in the included studies will be detailed in the data extraction tables.

Eligibility criteria - Outcome

Studies that include health outcomes informed by, but not limited to, core outcome domains for Outcome Measures in Rheumatology (OMERACT) for osteoarthritis [4], core outcome measurements for clinical trials with non-specific low back pain [9] and the Initiative on Methods Measurement and Pain Assessment in Clinical Trials for pain [12] will be considered. These domains are not designed specifically for CSp±R but intend to guide this review’s data collection and analysis in the absence of a core health outcome set for CSp±R. The domains included for this review will include, pain for example Visual Analogue Scale (VAS), Numeric Pain Rating Scale (NPRS) [24]; disability such as Neck disability Index (NDI) [45], Disability of Arm and Shoulder Index (DASH) [2], International Physical Activity Questionnaire (IPAQ) [17]; Health Status for example 36-item Short Form Survey (SF-36) [23], 12 item Short Form Survey (SF-12) [46]. Health care utilisation such as clinician visits, spinal imaging, hospital length of stay will also be reviewed. The outcomes at different follow up periods for study inclusion (e.g. 1 month, 3 months, 6 months) will be reviewed. We will contact lead authors for missing or incomplete data to inform

analysis. Where data is incomplete or missing the authorship will take a pragmatic decision based on data available and whether to include for narrative review. A list of studies with missing or incomplete data sets will be listed as an appendix on publication.

Study characteristics

Studies must meet the population, exposure, comparator and outcome criteria to be included. Studies must be in the English language (or can be translated to English) and be either case control, cross-sectional, or cohort study design. No restriction on publication date will be applied. All articles suitable for inclusion will be referenced in the final manuscript. Studies will be excluded if they are animal or cadaveric studies, commentaries, editorials, single case study, reports or laboratory data, books or book chapters, letters, conference posters or proceedings that do not have full methodology and data sets available on request, lecture slides, study protocols, social media postings (including blogs) or online learning material (e.g. massive online learning sites, videos).

Information sources

The search strategy will be primarily developed by one author (MM) and reviewed by all corresponding authors to reach agreement and consensus. The draft search strategy is presented in **Figure 2**. A systematic search of electronic databases EMBASE, CINAHL, MEDLINE (PubMed) from inception to April 2020 will be completed by one reviewer (NS). The full PubMed search strategy will be made available as a supplementary file on dissemination. EndNote (X9.2 for MacOS) will be the reference manager software to store records, identify and remove duplicates. All included studies will undergo reference checking. When formal study inclusion has taken place, we will agree on “expert” authors in the field to contact by email requesting any pending articles and/or missing data. An unpublished (grey) literature search and trial registry will also be completed (e.g. WHO.It, ZETOC, British library higher education thesis deposits).

Data collection - Study selection

Results of the search strategy will be uploaded into Microsoft Excel spreadsheet. This will be securely remote-stored for all authors to access. Two reviewers will independently review, check titles and abstracts and document decisions on which of the studies should be included. A third reviewer will independently review decisions made by the two reviewers. The third reviewer will adjudicate any disagreements and discussion to reach final consensus.

An assessment of reliability will take place between MM and NS of 10% potentially eligible articles for full text order using a weighted Kappa statistic. The between

reviewer agreement data will be available in the final manuscript for the overall agreement. Following the title and abstract screening, potential eligible articles will be ordered full text. MM and NS will independently review full text articles against the inclusion criteria. Independent decisions on the inclusion criteria will be stored in table format and will be made available on request. MM and NS will reach consensus through discussion, where consensus cannot be reached a third author (MT) will act as an adjudicator.

Data collection – Data collection and extraction

Data extraction forms will be designed by the lead author (MM). This form will be reviewed and agreed by all authors part of this review. Two authors (MM and TS) will independently extract data from the included studies. Two authors (MM and TS) will meet to discuss the data extracted, any disagreements and reach consensus. If consensus cannot be achieved, a third author (MT) will be consulted and will act as adjudicator. Data extracted will include lead author and date of publication; study design; study demographics (country, sample size, age range or mean, gender ratio); definition of exposure; definition of comparator; outcome measure description; Risk estimates (Risk ratios, Hazard ratios, Odds ratio and/or mean differences including 95% Confidence Intervals (CI) where available or can be calculated post-hoc).

Methodological quality

Two authors (MM, NS) will independently assess the quality of each included study using a Newcastle-Ottawa Quality Scale (NOS) Assessment quality appraisal tool [41]. This appraisal tool is recommended by the Cochrane Handbook for Systematic Review. The NOS checklist assesses quality of articles across 3 domains: selection of the studies groups; comparability of the groups and control for confounding factors; and exposure. MM and NS will discuss quality appraisal disagreements to reach consensus. MM and NS will independently determine whether outcomes are very low, low, moderate or high certainty based on the Grading of Recommendations, Assessment, Development and Evaluations [1]. GRADE will facilitate understanding outcomes quality and transparent grading of certainty in the included studies [1]. GRADE has five domains assessing the certainty of evidence: Risk of bias; Imprecision; Inconsistency; Indirectness; Publication bias.

Synthesis

Two authors (MM, TS) will assess included papers from a clinical perspective (e.g. diagnosis, variability in population characteristics) and study methodology to determine whether studies could be pooled together for synthesis. Where indicated, statistical heterogeneity assessments will be completed. If there is significant clinical heterogeneity, studies will be described narratively. For example, patient populations, clinical diagnosis. When there is minimal or no clinical heterogeneity, a random effects meta-analysis will be undertaken to compare the prevalence, odds or the risk of different psychosocial and mental health risk factors in CSp±R.

Additionally, statistical heterogeneity testing will be completed through the inconsistency value (I^2) and Cochran Q statistic. It is anticipated that there will be variability on assessment across included studies therefore a random-effect model will be adopted. A calculation of publication bias will be conducted using a Galbraith plot [13]. Data analysis will follow guidelines in the Cochrane handbook (Section 9.4.5). We will complete an inverse-variance random-effects method by assessing the standardised mean difference and present with 95% confidence intervals and forest plots

Mixed sample populations are anticipated. An analysis of studies with cervical spine pain and radiculopathy will be completed as an additional sub-group analysis.

Additionally, we will stratify results by duration of CSp±R (acute <3 months in duration and persistent ≥ 3 months in duration). Point estimates (Risk ratio, hazard ratio and odds ratio) will also be analysed separately. A further sub-analysis on cervical spine radiculopathy *probable* and *definite* diagnostic criteria and their association with psychosocial and/or mental health factors will be completed.

Furthermore, analyses will be conducted separately for cohort and case-control studies, and for unadjusted associations. Sensitivity analysis will be completed by an analysis of the study quality (risk of bias).

Discussion

This will be a first systematic review investigating the association between psychosocial factors or mental health symptoms in adults with CSp±R. It will be a comprehensive synthesis review with the aim to enhance understanding of the association of psychosocial factors and CSp±R on health outcomes.

It is acknowledged that there will be limitations to this review. First, there are no core outcome sets for CSp±R which may mean a variability of measurements used to assess outcomes in the literature. We have recognised this and have identified closely related core outcome sets for osteoarthritis, chronic pain and low back pain to inform our analysis. Furthermore, we will complete sub-group analyses as detailed above. Second, there is no universally accepted diagnostic criteria for cervical spine radiculopathy. Therefore we have adopted an approach reflecting contemporary clinical practice to moderate radiculopathy diagnostic variance. A strength of our review will be including studies with standardised validated health outcome questionnaires. However, questionnaires may not be generalisable across all healthcare settings globally. This will be discussed narratively in our analysis.

We are expecting that the topic theme, methodology and results will have interest for a wide range of audiences. We will submit the final piece of work to an international multi-disciplinary, peer reviewed journal (e.g. *PAIN Reports*, *PAIN*) with open access. We aim to present findings at an international health science conference (e.g. International Association of the Study of Pain, World Conference of Physical Therapists). The findings will also form part of the knowledge exchange and

research informed teaching strategies with health and social care students in our affiliated university posts and associated clinical departments. All authors have a social media presence freely available to the general public. We plan to detail plain English findings through this medium (e.g. Twitter).

In summary, our findings will have relevance to patients, healthcare clinicians, researchers and policy makers. Enhancing our understanding of how psychosocial factors and/or mental health symptoms are associated with health outcomes in people with CSp±R will support the formulation of prognosis and collaborative management decisions. The review will identify gaps in research, thereby informing future experimental and observational study design. The results will also support the guidance of healthcare resources and aim to enhance overall health outcomes for patients.

Reference List

1. Atkins D, Best D, Briss PA, Eccles M, Falck-Ytter Y, Flottorp S, Guyatt GH, Harbour RT, Haugh MC, Henry D, Hill S, Jaeschke R, Leng G, Liberati A, Magrini N, Mason J, Middleton P, Mrukowicz J, O'Connell D, Oxman AD, Phillips B, Schunemann HJ, Edejer T, Varonen H, Vist GE, Williams JW, Jr., Zaza S, Group GW. Grading quality of evidence and strength of recommendations. *BMJ*. 2004;328(7454):1490.
2. Beaton DE, Katz JN, Fossel AH, Wright JG, Tarasuk V, Bombardier C. Measuring the whole or the parts? Validity, reliability, and responsiveness of the Disabilities of the Arm, Shoulder and Hand outcome measure in different regions of the upper extremity. *J Hand Ther*. 2001;14(2):128-46.
3. Blozik E, Laptinskaya D, Herrmann-Lingen C, Schaefer H, Kochen MM, Himmel W, Scherer M. Depression and anxiety as major determinants of neck pain: a cross-sectional study in general practice. *BMC musculoskeletal disorders*. 2009;10:13.
4. Boers M, Kirwan JR, Wells G, Beaton D, Gossec L, d'Agostino MA, Conaghan PG, Bingham CO, 3rd, Brooks P, Landewe R, March L, Simon LS, Singh JA, Strand V, Tugwell P. Developing core outcome measurement sets for clinical trials: OMERACT filter 2.0. *Journal of clinical epidemiology*. 2014;67(7):745-53.
5. Bogduk N. The anatomy and pathophysiology of neck pain. *Physical medicine and rehabilitation clinics of North America*. 2011;22(3):367-82, vii.
6. Bono CM, Ghiselli G, Gilbert TJ, Kreiner DS, Reitman C, Summers JT, Baisden JL, Easa J, Fernand R, Lamer T, Matz PG, Mazanec DJ, Resnick DK, Shaffer WO, Sharma AK, Timmons RB, Toton JF, North American Spine S. An evidence-based clinical guideline for the diagnosis and treatment of cervical radiculopathy from degenerative disorders. *The spine journal : official journal of the North American Spine Society*. 2011;11(1):64-72.
7. Carroll LJ, Hogg-Johnson S, van der Velde G, Haldeman S, Holm LW, Carragee EJ, Hurwitz EL, Cote P, Nordin M, Peloso PM, Guzman J, Cassidy JD. Course and prognostic factors for neck pain in the general population: results of the Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders. *Spine*. 2008;33(4 Suppl):S75-82.
8. Chester R, Jerosch-Herold C, Lewis J, Shepstone L. Psychological factors are associated with the outcome of physiotherapy for people with shoulder pain: a multicentre longitudinal cohort study. *Br J Sports Med*. 2018;52(4):269-75.
9. Chiarotto A, Boers M, Deyo RA, Buchbinder R, Corbin TP, Costa LOP, Foster NE, Grotle M, Koes BW, Kovacs FM, Lin CC, Maher CG, Pearson AM, Peul WC, Schoene ML, Turk DC, van Tulder MW, Terwee CB, Ostelo RW. Core outcome measurement instruments for clinical trials in nonspecific low back pain. *Pain*. 2018;159(3):481-95.
10. Chou R, Cote P, Randhawa K, Torres P, Yu H, Nordin M, Hurwitz EL, Haldeman S, Cedraschi C. The Global Spine Care Initiative: applying evidence-based guidelines on the non-invasive management of back and neck pain to low- and middle-income communities. *Eur Spine J*. 2018;27(Suppl 6):851-60.
11. Cote P, Cassidy JD, Carroll LJ, Kristman V. The annual incidence and course of neck pain in the general population: a population-based cohort study. *Pain*. 2004;112(3):267-73.
12. Dworkin RH, Turk DC, Farrar JT, Haythornthwaite JA, Jensen MP, Katz NP, Kerns RD, Stucki G, Allen RR, Bellamy N, Carr DB, Chandler J, Cowan P, Dionne R, Galer BS, Hertz S, Jadad AR, Kramer LD, Manning DC, Martin S, McCormick CG, McDermott MP, McGrath P, Quessy S, Rappaport BA, Robbins W, Robinson JP, Rothman M, Royal MA, Simon L, Stauffer JW, Stein W, Tollett J, Wernicke J, Witter J, Impact. Core outcome measures for chronic pain clinical trials: IMMPACT recommendations. *Pain*. 2005;113(1-2):9-19.

13. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ*. 1997;315(7109):629-34.
14. Finnerup NB, Haroutounian S, Kamerman P, Baron R, Bennett DL, Bouhassira D, Cruccu G, Freeman R, Hansson P, Nurmikko T, Raja SN, Rice AS, Serra J, Smith BH, Treede RD, Jensen TS. Neuropathic pain: an updated grading system for research and clinical practice. *Pain*. 2016;157(8):1599-606.
15. Foster NE, Anema JR, Cherkin D, Chou R, Cohen SP, Gross DP, Ferreira PH, Fritz JM, Koes BW, Peul W, Turner JA, Maher CG, Lancet Low Back Pain Series Working G. Prevention and treatment of low back pain: evidence, challenges, and promising directions. *Lancet (London, England)*. 2018;391(10137):2368-83.
16. Haanpaa M, Attal N, Backonja M, Baron R, Bennett M, Bouhassira D, Cruccu G, Hansson P, Haythornthwaite JA, Iannetti GD, Jensen TS, Kauppila T, Nurmikko TJ, Rice AS, Rowbotham M, Serra J, Sommer C, Smith BH, Treede RD. NeuPSIG guidelines on neuropathic pain assessment. *Pain*. 2011;152(1):14-27.
17. Hagstromer M, Oja P, Sjostrom M. The International Physical Activity Questionnaire (IPAQ): a study of concurrent and construct validity. *Public Health Nutr*. 2006;9(6):755-62.
18. Harris-Adamson C, Eisen EC, Dale AM, Evanoff B, Hegmann KT, Thiese MS, Kapellusch J, Garg A, Burt S, Silverstein B, Bao S, Merlino L, Gerr F, Rempel D. Personal and workplace psychosocial risk factors for carpal tunnel syndrome: a pooled study cohort: author response. *Occup Environ Med*. 2014;71(4):303-4.
19. Hogg-Johnson S, van der Velde G, Carroll LJ, Holm LW, Cassidy JD, Guzman J, Cote P, Haldeman S, Ammendolia C, Carragee E, Hurwitz E, Nordin M, Peloso P. The burden and determinants of neck pain in the general population: results of the Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders. *Journal of manipulative and physiological therapeutics*. 2009;32(2 Suppl):S46-60.
20. Hoy D, March L, Woolf A, Blyth F, Brooks P, Smith E, Vos T, Barendregt J, Blore J, Murray C, Burstein R, Buchbinder R. The global burden of neck pain: estimates from the global burden of disease 2010 study. *Annals of the rheumatic diseases*. 2014;73(7):1309-15.
21. Hoy DG, Protani M, De R, Buchbinder R. The epidemiology of neck pain. *Best practice & research Clinical rheumatology*. 2010;24(6):783-92.
22. IASP. Classification of chronic pain. Descriptions of chronic pain syndromes and definitions of pain terms. Prepared by the International Association for the Study of Pain, Subcommittee on Taxonomy. *Pain Suppl*. 1986;3:S1-226.
23. Jenkinson C, Wright L, Coulter A. Criterion validity and reliability of the SF-36 in a population sample. *Quality of life research : an international journal of quality of life aspects of treatment, care and rehabilitation*. 1994;3(1):7-12.
24. Jensen MP, McFarland CA. Increasing the reliability and validity of pain intensity measurement in chronic pain patients. *Pain*. 1993;55(2):195-203.
25. Jones M, Edwards I, Gifford L. Conceptual models for implementing biopsychosocial theory in clinical practice. *Manual therapy*. 2002;7(1):2-9.
26. Liu F, Fang T, Zhou F, Zhao M, Chen M, You J, Jin Y, Xie J, Liu Z. Association of Depression/Anxiety Symptoms with Neck Pain: A Systematic Review and Meta-Analysis of Literature in China. *Pain Res Manag*. 2018;2018:3259431.
27. Martikainen P, Bartley M, Lahelma E. Psychosocial determinants of health in social epidemiology. *Int J Epidemiol*. 2002;31(6):1091-3.
28. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ*. 2009;339:b2535.
29. Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, Shekelle P, Stewart LA, Group P-P. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Systematic reviews*. 2015;4:1.

30. Murray CJ, Barber RM, Foreman KJ, Abbasoglu Ozgoren A, Abd-Allah F, Abera SF, Aboyans V, Abraham JP, Abubakar I, Abu-Raddad LJ, Abu-Rmeileh NM, Achoki T, Ackerman IN, Ademi Z, Adou AK, Adsuar JC, Afshin A, Agardh EE, Alam SS, Alasfoor D, Albittar MI, Alegretti MA, Alemu ZA, Alfonso-Cristancho R, Alhabib S, Ali R, Alla F, Allebeck P, Almazroa MA, Alsharif U, Alvarez E, Alvis-Guzman N, Amare AT, Ameh EA, Amini H, Ammar W, Anderson HR, Anderson BO, Antonio CA, Anwari P, Arnlov J, Arsic Arsenijevic VS, Artaman A, Asghar RJ, Assadi R, Atkins LS, Avila MA, Awuah B, Bachman VF, Badawi A, Bahit MC, Balakrishnan K, Banerjee A, Barker-Collo SL, Barquera S, Barregard L, Barrero LH, Basu A, Basu S, Basulaiman MO, Beardsley J, Bedi N, Beghi E, Bekele T, Bell ML, Benjet C, Bennett DA, Bensenor IM, Benzian H, Bernabe E, Bertozzi-Villa A, Beyene TJ, Bhala N, Bhalla A, Bhutta ZA, Bienhoff K, Bikbov B, Biryukov S, Blore JD, Blosser CD, Blyth FM, Bohensky MA, Bolliger IW, Bora Basara B, Bornstein NM, Bose D, Boufous S, Bourne RR, Boyers LN, Brainin M, Brayne CE, Brazinova A, Breitborde NJ, Brenner H, Briggs AD, Brooks PM, Brown JC, Brugha TS, Buchbinder R, Buckle GC, Budke CM, Bulchis A, Bulloch AG, Campos-Nonato IR, Carabin H, Carapetis JR, Cardenas R, Carpenter DO, Caso V, Castaneda-Orjuela CA, Castro RE, Catala-Lopez F, Cavalleri F, Cavlin A, Chadha VK, Chang JC, Charlson FJ, Chen H, Chen W, Chiang PP, Chimed-Ochir O, Chowdhury R, Christensen H, Christophi CA, Cirillo M, Coates MM, Coffeng LE, Coggeshall MS, Colistro V, Colquhoun SM, Cooke GS, Cooper C, Cooper LT, Coppola LM, Cortinovis M, Criqui MH, Crump JA, Cuevas-Nasu L, Danawi H, Dandona L, Dandona R, Dansereau E, Dargan PI, Davey G, Davis A, Davitoiu DV, Dayama A, De Leo D, Degenhardt L, Del Pozo-Cruz B, Dellavalle RP, Deribe K, Derrett S, Des Jarlais DC, Dessalegn M, Dharmaratne SD, Dherani MK, Diaz-Torne C, Dicker D, Ding EL, Dokova K, Dorsey ER, Driscoll TR, Duan L, Duber HC, Ebel BE, Edmond KM, Elshrek YM, Endres M, Ermakov SP, Erskine HE, Eshrati B, Esteghamati A, Estep K, Faraon EJ, Farzadfar F, Fay DF, Feigin VL, Felson DT, Fereshtehnejad SM, Fernandes JG, Ferrari AJ, Fitzmaurice C, Flaxman AD, Fleming TD, Foigt N, Forouzanfar MH, Fowkes FG, Paleo UF, Franklin RC, Furst T, Gabbe B, Gaffikin L, Gankpe FG, Geleijnse JM, Gessner BD, Gething P, Gibney KB, Giroud M, Giussani G, Gomez Dantes H, Gona P, Gonzalez-Medina D, Gosselin RA, Gotay CC, Goto A, Gouda HN, Graetz N, Gughani HC, Gupta R, Gupta R, Gutierrez RA, Haagsma J, Hafezi-Nejad N, Hagan H, Halasa YA, Hamadeh RR, Hamavid H, Hammami M, Hancock J, Hankey GJ, Hansen GM, Hao Y, Harb HL, Haro JM, Havmoeller R, Hay SI, Hay RJ, Heredia-Pi IB, Heuton KR, Heydarpour P, Higashi H, Hajar M, Hoek HW, Hoffman HJ, Hosgood HD, Hossain M, Hotez PJ, Hoy DG, Hsairi M, Hu G, Huang C, Huang JJ, Husseini A, Huynh C, Iannarone ML, Iburg KM, Innos K, Inoue M, Islami F, Jacobsen KH, Jarvis DL, Jassal SK, Jee SH, Jeemon P, Jensen PN, Jha V, Jiang G, Jiang Y, Jonas JB, Juel K, Kan H, Karch A, Karema CK, Karimkhani C, Karthikeyan G, Kassebaum NJ, Kaul A, Kawakami N, Kazanjan K, Kemp AH, Kengne AP, Keren A, Khader YS, Khalifa SE, Khan EA, Khan G, Khang YH, Kieling C, Kim D, Kim S, Kim Y, Kinfu Y, Kinge JM, Kivipelto M, Knibbs LD, Knudsen AK, Kokubo Y, Kosen S, Krishnaswami S, Kuate Defo B, Kucuk Bicer B, Kuipers EJ, Kulkarni C, Kulkarni VS, Kumar GA, Kyu HH, Lai T, Lalloo R, Lallukka T, Lam H, Lan Q, Lansingh VC, Larsson A, Lawrynowicz AE, Leasher JL, Leigh J, Leung R, Levitz CE, Li B, Li Y, Li Y, Lim SS, Lind M, Lipshultz SE, Liu S, Liu Y, Lloyd BK, Lofgren KT, Logroscino G, Looker KJ, Lortet-Tieulent J, Lotufo PA, Lozano R, Lucas RM, Lunevicius R, Lyons RA, Ma S, Macintyre MF, Mackay MT, Majdan M, Malekzadeh R, Marcenes W, Margolis DJ, Margono C, Marzan MB, Masci JR, Mashal MT, Matzopoulos R, Mayosi BM, Mazorodze TT, McGill NW, McGrath JJ, McKee M, McLain A, Meaney PA, Medina C, Mehndiratta MM, Mekonnen W, Melaku YA, Meltzer M, Memish ZA, Mensah GA, Meretoja A, Mhimbira FA, Micha R, Miller TR, Mills EJ, Mitchell PB, Mock CN, Mohamed Ibrahim N, Mohammad KA, Mokdad AH, Mola GL, Monasta L, Montanez Hernandez JC,

Montico M, Montine TJ, Mooney MD, Moore AR, Moradi-Lakeh M, Moran AE, Mori R, Moschandreas J, Moturi WN, Moyer ML, Mozaffarian D, Msemburi WT, Mueller UO, Mukaigawara M, Mullany EC, Murdoch ME, Murray J, Murthy KS, Naghavi M, Naheed A, Naidoo KS, Naldi L, Nand D, Nangia V, Narayan KM, Nejjari C, Neupane SP, Newton CR, Ng M, Ngalesoni FN, Nguyen G, Nisar MI, Nolte S, Norheim OF, Norman RE, Norrving B, Nyakarahuka L, Oh IH, Ohkubo T, Ohno SL, Olusanya BO, Opio JN, Ortblad K, Ortiz A, Pain AW, Pandian JD, Panelo CI, Papachristou C, Park EK, Park JH, Patten SB, Patton GC, Paul VK, Pavlin BI, Pearce N, Pereira DM, Perez-Padilla R, Perez-Ruiz F, Perico N, Pervaiz A, Pesudovs K, Peterson CB, Petzold M, Phillips MR, Phillips BK, Phillips DE, Piel FB, Plass D, Poenaru D, Polinder S, Pope D, Popova S, Poulton RG, Pourmalek F, Prabhakaran D, Prasad NM, Pullan RL, Qato DM, Quistberg DA, Rafay A, Rahimi K, Rahman SU, Raju M, Rana SM, Razavi H, Reddy KS, Refaat A, Remuzzi G, Resnikoff S, Ribeiro AL, Richardson L, Richardus JH, Roberts DA, Rojas-Rueda D, Ronfani L, Roth GA, Rothenbacher D, Rothstein DH, Rowley JT, Roy N, Ruhago GM, Saeedi MY, Saha S, Sahraian MA, Sampson UK, Sanabria JR, Sandar L, Santos IS, Satpathy M, Sawhney M, Scarborough P, Schneider IJ, Schottker B, Schumacher AE, Schwebel DC, Scott JG, Seedat S, Sepanlou SG, Serina PT, Servan-Mori EE, Shackelford KA, Shaheen A, Shahrz S, Shamah Levy T, Shangquan S, She J, Sheikhabaei S, Shi P, Shibuya K, Shinohara Y, Shiri R, Shishani K, Shiue I, Shrimel MG, Sigfusdottir ID, Silberberg DH, Simard EP, Sindi S, Singh A, Singh JA, Singh L, Skirbekk V, Slepak EL, Sliwa K, Soneji S, Soreide K, Soshnikov S, Sposato LA, Sreeramareddy CT, Stanaway JD, Stathopoulou V, Stein DJ, Stein MB, Steiner C, Steiner TJ, Stevens A, Stewart A, Stovner LJ, Stroumpoulis K, Sunguya BF, Swaminathan S, Swaroop M, Sykes BL, Tabb KM, Takahashi K, Tandon N, Tanne D, Tanner M, Tavakkoli M, Taylor HR, Te Ao BJ, Tediosi F, Temesgen AM, Templin T, Ten Have M, Tenkorang EY, Terkawi AS, Thomson B, Thorne-Lyman AL, Thrift AG, Thurston GD, Tillmann T, Tonelli M, Topouzis F, Toyoshima H, Traebert J, Tran BX, Trillini M, Truelsen T, Tsilimbaris M, Tuzcu EM, Uchendu US, Ukwaja KN, Undurraga EA, Uzun SB, Van Brakel WH, Van De Vijver S, van Gool CH, Van Os J, Vasankari TJ, Venketasubramanian N, Violante FS, Vlassov VV, Vollset SE, Wagner GR, Wagner J, Waller SG, Wan X, Wang H, Wang J, Wang L, Warouw TS, Weichenthal S, Weiderpass E, Weintraub RG, Wenzhi W, Werdecker A, Westerman R, Whiteford HA, Wilkinson JD, Williams TN, Wolfe CD, Wolock TM, Woolf AD, Wulf S, Wurtz B, Xu G, Yan LL, Yano Y, Ye P, Yentur GK, Yip P, Yonemoto N, Yoon SJ, Younis MZ, Yu C, Zaki ME, Zhao Y, Zheng Y, Zonies D, Zou X, Salomon JA, Lopez AD, Vos T. Global, regional, and national disability-adjusted life years (DALYs) for 306 diseases and injuries and healthy life expectancy (HALE) for 188 countries, 1990-2013: quantifying the epidemiological transition. *Lancet* (London, England). 2015;386(10009):2145-91.

31. Nordin M, Carragee EJ, Hogg-Johnson S, Weiner SS, Hurwitz EL, Peloso PM, Guzman J, van der Velde G, Carroll LJ, Holm LW, Cote P, Cassidy JD, Haldeman S, Bone, Joint Decade - Task Force on Neck Pain, Its Associated Disorders. Assessment of neck pain and its associated disorders: results of the Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders. *Spine*. 2008;33(4 Suppl):S101-22.

32. Pincus T, Burton AK, Vogel S, Field AP. A systematic review of psychological factors as predictors of chronicity/disability in prospective cohorts of low back pain. *Spine*. 2002;27(5):E109-20.

33. Pinheiro MB, Ferreira ML, Refshauge K, Maher CG, Ordonana JR, Andrade TB, Tsathas A, Ferreira PH. Symptoms of depression as a prognostic factor for low back pain: a systematic review. *The spine journal : official journal of the North American Spine Society*. 2016;16(1):105-16.

34. Radhakrishnan K, Litchy WJ, O'Fallon WM, Kurland LT. Epidemiology of cervical radiculopathy. A population-based study from Rochester, Minnesota, 1976 through 1990. *Brain : a journal of neurology*. 1994;117 (Pt 2):325-35.
35. Ramond A, Bouton C, Richard I, Roquelaure Y, Baufreton C, Legrand E, Huez JF. Psychosocial risk factors for chronic low back pain in primary care--a systematic review. *Fam Pract*. 2011;28(1):12-21.
36. Ryan C, Roberts L. 'Life on hold': The lived experience of radicular symptoms. A qualitative, interpretative inquiry. *Musculoskelet Sci Pract*. 2019;39:51-7.
37. Schoenfeld AJ, George AA, Bader JO, Caram PM, Jr. Incidence and epidemiology of cervical radiculopathy in the United States military: 2000 to 2009. *Journal of spinal disorders & techniques*. 2012;25(1):17-22.
38. Scholz J, Finnerup NB, Attal N, Aziz Q, Baron R, Bennett MI, Benoliel R, Cohen M, Cruccu G, Davis KD, Evers S, First M, Giamberardino MA, Hansson P, Kaasa S, Korwisi B, Kosek E, Lavand'homme P, Nicholas M, Nurmikko T, Perrot S, Raja SN, Rice ASC, Rowbotham MC, Schug S, Simpson DM, Smith BH, Svensson P, Vlaeyen JWS, Wang SJ, Barke A, Rief W, Treede RD, Classification Committee of the Neuropathic Pain Special Interest G. The IASP classification of chronic pain for ICD-11: chronic neuropathic pain. *Pain*. 2019;160(1):53-9.
39. Shahidi B, Curran-Everett D, Maluf KS. Psychosocial, Physical, and Neurophysiological Risk Factors for Chronic Neck Pain: A Prospective Inception Cohort Study. *J Pain*. 2015;16(12):1288-99.
40. Smyth MJ, Wright V. Sciatica and the intervertebral disc; an experimental study. *J Bone Joint Surg Am*. 1958;40-A(6):1401-18.
41. Stang A. Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. *Eur J Epidemiol*. 2010;25(9):603-5.
42. Sterud T, Johannessen HA, Tynes T. Work-related psychosocial and mechanical risk factors for neck/shoulder pain: a 3-year follow-up study of the general working population in Norway. *International archives of occupational and environmental health*. 2014;87(5):471-81.
43. Stubbs B, Aluko Y, Myint PK, Smith TO. Prevalence of depressive symptoms and anxiety in osteoarthritis: a systematic review and meta-analysis. *Age Ageing*. 2016;45(2):228-35.
44. Treede RD, Jensen TS, Campbell JN, Cruccu G, Dostrovsky JO, Griffin JW, Hansson P, Hughes R, Nurmikko T, Serra J. Neuropathic pain: redefinition and a grading system for clinical and research purposes. *Neurology*. 2008;70(18):1630-5.
45. Vernon H, Mior S. The Neck Disability Index: a study of reliability and validity. *Journal of manipulative and physiological therapeutics*. 1991;14(7):409-15.
46. Ware J, Jr., Kosinski M, Keller SD. A 12-Item Short-Form Health Survey: construction of scales and preliminary tests of reliability and validity. *Med Care*. 1996;34(3):220-33.
47. Wertli MM, Burgstaller JM, Weiser S, Steurer J, Kofmehl R, Held U. Influence of catastrophizing on treatment outcome in patients with nonspecific low back pain: a systematic review. *Spine*. 2014;39(3):263-73.
48. Wibault J, oberg B, Dederling A, Lofgren H, Zsigmond P, Persson L, Peolsson A. Individual factors associated with neck disability in patients with cervical radiculopathy scheduled for surgery: a study on physical impairments, psychosocial factors, and life style habits. *Eur Spine J*. 2014;23(3):599-605.
49. Wise BL, Niu J, Zhang Y, Wang N, Jordan JM, Choy E, Hunter DJ. Psychological factors and their relation to osteoarthritis pain. *Osteoarthritis Cartilage*. 2010;18(7):883-7.