



# Chemical warfare between fungus-growing ants and their pathogens

Sibyl F. D. Batey<sup>1</sup>, Claudio Greco<sup>1</sup>,  
Matthew I. Hutchings<sup>1,2</sup> and Barrie Wilkinson<sup>1</sup>

## Abstract

Fungus-growing attine ants are under constant threat from fungal pathogens such as the specialized mycoparasite *Escovopsis*, which uses combined physical and chemical attack strategies to prey on the fungal gardens of the ants. In defence, some species assemble protective microbiomes on their exoskeletons that contain antimicrobial-producing Actinobacteria. Underlying this network of mutualistic and antagonistic interactions are an array of chemical signals.

*Escovopsis weberi* produces the shearinine terpene-indole alkaloids, which affect ant behaviour, diketopiperazines to combat defensive bacteria, and other small molecules that inhibit the fungal cultivar. *Pseudonocardia* and *Streptomyces* mutualist bacteria produce depsipeptide and polyene macrolide antifungals active against *Escovopsis* spp. The ant nest metabolome is further complicated by competition between defensive bacteria, which produce antibacterials active against even closely related species.

## Addresses

<sup>1</sup> Department of Molecular Microbiology, John Innes Centre, Norwich, NR4 7UH, United Kingdom

<sup>2</sup> School of Biological Sciences, University of East Anglia, Norwich, NR4 7TU, United Kingdom

Corresponding authors: Wilkinson, Barrie ([barrie.wilkinson@jic.ac.uk](mailto:barrie.wilkinson@jic.ac.uk)); Hutchings, Matthew I ([matt.hutchings@jic.ac.uk](mailto:matt.hutchings@jic.ac.uk))

Current Opinion in Chemical Biology 2020, 59:172–181

This review comes from a themed issue on **Mechanistic biology**

Edited by Katherine S. Ryan and Tomohisa Kuzuyama

For a complete overview see the [Issue](#) and the [Editorial](#)

<https://doi.org/10.1016/j.cbpa.2020.08.001>

1367-5931/© 2020 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

## Keywords

Fungus-growing ants, Mutualism, Antagonism, Specialized metabolites, *Escovopsis*, *Pseudonocardia*, *Streptomyces*, Antimicrobials.

## Introduction: farmers at war

Ants of the tribe Attini cultivate basidiomycete fungi in a process akin to human agriculture that evolved around

50–60 million years ago [1]. The Attini can be divided according to their fungal cultivars, which include species across the tribe Leucocoprineae. The lower attines tend to have less specialized fungal cultivars, which they feed with dead biomass, including vegetative debris and insect corpses [2]. The higher attines have specialized obligate fungal cultivars which have developed hyphal swellings known as gongylidia that provide a rich source of nutrients for the colony [3]. The leafcutter ants are the most highly derived attines and comprise the genera *Atta* and *Acromyrmex* [2]. They mainly cultivate *Leucogarricus gongylophorus* [4], a polyploid clone, which is transmitted vertically between nests [5]. The ants actively cut fresh leaf material to feed to their cultivar, which provides the sole food source for the ant larvae [2].

The most significant threat to this mutualistic endeavour comes from specialized fungal pathogens in the genus *Escovopsis*, which have co-evolved with attines and are highly adapted for a mycoparasitic lifestyle [6]. More recently, *Escovopsiodes* have been identified as a further distinct mycoparasitic genus [7]. If left unchecked, *Escovopsis* spp. can overrun a nest leading to devastating colony collapse [8]. The most well-studied species is *Escovopsis weberi* [9], commonly found in the nests of leafcutter ants. The ants carefully groom and weed their fungal gardens to remove *Escovopsis* and other pathogens [2], modifying their hygiene strategy according to the growth stage of the pathogen [10]. Many attines also house and feed defensive bacteria on their cuticles, initially a single vertically transmitted *Pseudonocardia* phylotype transferred to newly eclosed workers and virgin queens from their sisters and/or the fungal cultivar [11]. Other actinomycetes, such as *Streptomyces* species, are horizontally acquired from the environment to form a protective microbiome [12]. These bacteria produce antifungal compounds that are active against *Escovopsis* spp., as well as antibacterials to outcompete other defensive bacteria (Table 1) [13]. Within the leafcutters, there has been an interesting divergence, with only *Acromyrmex* maintaining a protective microbiome, whereas *Atta* rely on their own endogenous chemical defences [14]. However, *Escovopsis* comes armed with a chemical war chest of its own that includes antibacterial, insecticidal and antifungal specialized

Table 1

## Summary of bioactive compounds isolated from fungus-growing ant mutualists and pathogens.

Producer	Ant Species	Compound	Bioactivity	References
<b>Escovopsis</b>				
<i>E. weberi</i>	<i>Acromyrmex octospinosus</i>	Shearinine L (1)	- <i>A. octospinosus</i> ants chose not to eat 1-impregnated oat flakes	[15]
<i>E. weberi</i>	<i>Acromyrmex echinator</i>	Shearinine D (2)	- Adverse effect on <i>A. echinator</i> ant behaviour, lethal at high concentrations. - Active against <i>Pseudonocardia echinator</i> and <i>Pseudonocardia octospinosus</i> strains isolated from <i>A. echinator</i> colonies.	[15,16]
<i>E. weberi</i> <i>E. aspergilloides</i>	<i>A. echinator</i> <i>Trachomyrmex cornetzi</i>	Melinacidin IV (3)	- Active against <i>P. echinator</i> and <i>P. octospinosus</i> strains isolated from <i>A. echinator</i> colonies.	[16]
<i>E. weberi</i> <i>E. aspergilloides</i>	<i>A. echinator</i> & <i>A. octospinosus</i> <i>T. cornetzi</i>	Emodin (4)	- Active against <i>Leucoagaricus gongylophorus</i> and <i>Streptomyces</i> strains isolated from leafcutter ant nests.	[15,16]
<i>E. weberi</i>	<i>A. echinator</i>	Cycloarthropsone (5)	- Active against <i>L. gongylophorus</i> .	[15]
<b>Streptomyces</b>				
S4, Ao10	<i>A. octospinosus</i>	Candicidin D (9)	- Active against <i>E. weberi</i> but not against <i>L. gongylophorus</i> .	[36,38,40]
Ae32_2, S4, Ao10 Av28_2, Av28_3 Av25_1	<i>A. echinator</i> <i>A. octospinosus</i> <i>Acromyrmex volcanus</i>	Antimycin A1-A4 (10a-d)	- Active against <i>E. weberi</i> and <i>L. gongylophorus</i> .	[39,40]
ICBG292	<i>Cyphomyrmex</i> sp.	Mer-A2026B (11) /Piericidin-A1 (12)	- Active against different <i>Escovopsis</i> spp. - Antileishmanial activity.	[41]
Av25_2	<i>A. volcanus</i>	Actinomycin D (16) Actinomycin X2 (17)	- Active against <i>Pseudonocardia</i> and <i>Streptomyces</i> spp.	[39]
<b>Pseudonocardia</b>				
CC011120-4	<i>Apterostigma dentigerum</i>	Dentigerumycin A (6)	- Antifungal activity against <i>E. weberi</i>	[32]
<i>P. octospinosus</i> P1	<i>A. octospinosus</i>	Nystatin P1 (7)	- Antifungal activity against <i>E. weberi</i>	[29,36]
HH130629-09, HH130630-07	<i>Apterostigma</i> sp.	Selvamicin (8)	- Antifungal activity, activity against <i>E. weberi</i> not reported	[37]
BCI2	<i>A. dentigerum</i>	9-methoxyrebeccamycin (13)	- Inhibits other <i>Pseudonocardia</i> strains.	[47]
17SE-9	<i>Trachymyrmex septentrionalis</i>	GE37468 (14)	- Inhibits <i>Pseudonocardia</i> spp. from the same ant nest.	[48]
EC080529-01	<i>A. dentigerum</i>	6-deoxy-8-O-methylrabelomycin (15)	- Antibacterial and antimalarial activity	[49]

metabolites that target each component of the ant-microbe symbiosis (Table 1) [15,16].

Thus, there exists a highly co-evolved, complex network of mutualistic and antagonistic relationships within the nests of fungus-farming ants, Figure 1. Here, we review recent advances in our understanding of the sophisticated attack and defence strategies utilised, focussing on the complex array of chemical signals underpinning each interaction.


## Under attack

### Physical attack

The mechanism of parasitism by *Escovopsis* and *Escovopsioides* species requires the pathogens to have physical contact with the cultivar fungus. Early observations reported that *E. weberi* grows faster in the presence of

*L. gongylophorus* and causes hyphal degradation. In addition, different *Escovopsis* isolates vary in the level of selectivity toward the cultivar fungus [17]. Building on this work, Marfetán and colleagues [18] showed that all *E. weberi* strains were, at different levels, virulent towards *L. gongylophorus* and that the most virulent isolates could develop hook-like structures that were used to attach to the fungal cultivar. More recently, several *Escovopsis* species and two *Escovopsioides nivea* strains were reported to inhibit the growth of *L. gongylophorus* in co-cultivation experiments where both the pathogen and the garden fungus had a drastic change in pigmentation [9,19]. In addition, when *Escovopsis* spp. were grown on water agar with or without a small colony of *L. gongylophorus* it was found that the *Escovopsis* spp. grew only on co-cultivation plates and the pathogen grew directly towards *L. gongylophorus* forming hyphal bridges, and outgrew it within 48 h [18].

Figure 1



**A network of mutualistic and antagonistic interactions exists in the nests of fungus-growing ants.** Representative examples of each member of the symbiotic network are shown, with *Pseudonocardia* hyphae imaged by scanning electron microscopy (SEM). Red lines indicate antagonistic interactions, green arrows indicate mutualistic interactions and the green dashed arrow indicates indirect mutualism.


### Chemical attack

While the threat posed by *Escovopsis* species to fungus-growing ants has been well-established, the specialized metabolites it utilises to modulate this interaction have only recently begun to be uncovered, Figure 2. Indirect evidence came from the first genome sequence of *E. weberi* and associated RNA-sequencing data which identified that a biosynthetic gene cluster (BGC) for a polyketide synthase-derived specialized metabolite was upregulated when *E. weberi* and *L. gongylophorus* are co-cultured [9]. Rodrigues and colleagues reported that chemical extracts from *Escovopsis* species and *E. nivea* cultures were able to inhibit the growth of *L. gongylophorus* and that the extracts obtained from co-cultures of pathogen and cultivar were generally more active than extracts from axenic plates of the pathogen [19]. Another recent study identified the production of the terpene-indole alkaloid shearinines by *Escovopsis* sp. TZ49, using a combination of imaging mass spectrometry and MS/MS molecular networking [20]. In subsequent separate studies, shearinine metabolites were shown to be upregulated when *E. weberi* was grown in the presence of *L. gongylophorus* [16], and several shearinine congeners were isolated and characterised from a range of *Escovopsis* strains (six *E. weberi* isolates

and one *Escovopsis aspergilloides*) [15,16]. Worker ants learned not to choose oat flakes impregnated with shearinine L (1), but there was no obvious effect on waste production or worker mortality in the colony [15]. However, shearinine D (2) supplied as a glucose solution in the absence of the fungal cultivar adversely affected the behaviour of worker ants and was ultimately lethal. Compound 2 was also found to be elevated in the tissues of worker ants in a captive colony that suffered a natural *E. weberi* outbreak [16]. This is consistent with the previously documented roles of terpene-indole alkaloids acting as feeding deterrents and modulators of ion channels in various insects [21]. Furthermore, 2 was active against mutualist *Pseudonocardia* strains isolated from *Acromyrmex echinator* [16] whereas 1 was not active against *L. gongylophorus* or *Pseudonocardia* Ao19 isolated from *Acromyrmex octospinosus* [15]. The variation of responses observed for *Pseudonocardia* species to these different shearinine congeners may be due to their structural variation or could be attributed to the different methods of antimicrobial testing used.

Production of the epipolythiodiketopiperazine (ETP) melinacidin IV (3), a known antimicrobial and cytotoxic agent, was also increased when *E. weberi* was co-cultured

Figure 2



**a) Antagonistic interactions of *Escovopsis weberi* with other organisms.** Representative images of each symbiont are shown with red lines indicating antagonistic interactions. Specialized metabolites are represented with coloured shapes, as shown in **b. b) Chemical compounds isolated from *E. weberi*.** Main classes identified thus far are the shearinine terpene-indole alkaloids and the melinacidin epipolythiodiketopiperazines (ETPs), as well as other polyketide metabolites such as cycloarthropsone and the anthraquinone emodin.

with *L. gongylophorus* and was shown to inhibit the growth of *A. echinator* mutualist *Pseudonocardia* species [16]. Two other metabolites were isolated from *E. weberi*, the anthraquinone emodin (4) and the polyketide cycloarthropsone (5) [15,16], both of which inhibit the growth of the cultivar fungus *L. gongylophorus*. Compound 4 is also produced by *E. aspergilloides* and inhibits the growth of several streptomycete leafcutter ant mutualists [15].

There have been several reports of the isolation of black yeast-like fungi in the order Chaetothiales from the cuticles of fungus-growing ants, a potential additional player in the symbiotic network [21–24]. It has been proposed that the black yeasts could indirectly benefit the fungal pathogen as they were shown to inhibit the growth of ant mutualist *Pseudonocardia in vitro* [25]. However, phylogenetic analyses suggest they are not


particularly specialized in their interactions with fungus-growing ants [26].

## Defence

### Microbial defences

A highly effective defensive strategy utilised by fungus-growing ants is the assembly of a protective microbiome on the external surface of their exoskeletons, which is dominated by antimicrobial-producing actinobacteria [27]. In addition to protection of the fungal cultivar from *Escovopsis* attack, cuticular microbiomes have been shown to protect worker ants from infection by the entomopathogenic fungus *Metarhizium anisopliae* [28] and are proposed to help shape the cuticular microbiome by excluding non-mutualist bacteria [29]. The presence of the cuticular microbiome may also serve as a physical barrier against fungal infection [30].

Figure 3



**a) Interactions between defensive Actinobacteria and *Escovopsis weberi*.** The protective microbiome is visible as a white coating on the integument of an *Acromyrmex* ant. Representative images of each symbiont are shown with red lines indicating antagonistic interactions. Compounds are produced by mutualistic bacteria as defence/attack against both the fungal pathogen and competitors. Specialized metabolites are represented with coloured shapes according to **b**. Production of compounds active against streptomycete bacteria by *Pseudonocardia* spp. has not been reported. **b) Specialized metabolites produced by mutualistic bacteria.** The main classes are non-ribosomal peptides (NRPs), such as the depsipeptides, which include dentigerumycin A **6** and the actinomycins **16** and **17**. The antifungal polyene macrolides include nystatin P1 **7**, candididin D **9** and selvamicin **8**, the latter having an apparently novel mode of action. The activity of **8** and 6-deoxy-8-O-methylrabelomycin **15** against species from the nests of fungus-growing ants has not been reported.

The most well-studied bacterial symbionts belong to the *Pseudonocardia* genus, and many attines vertically transmit a single strain of *Pseudonocardia* [12]. The bacteria grow in and around the openings of specialized crypts on the ant cuticle and are proposed to feed on secretions from subcuticular glands [27]. The presence of *Pseudonocardia* species greatly improves the suppression of *E. weberi* infections in leafcutter colonies [30]. The bioactivity of mutualistic *Pseudonocardia* strains against *Escovopsis* spp. has been recapitulated *in vitro* [31], and several antifungal compounds have been identified.

*Pseudonocardia* produce two main classes of antifungal molecules: cyclic depsipeptides and polyene macrolides, Figure 3. The cyclic piperazine-containing depsipeptide dentigerumycin A (6) is produced by a *Pseudonocardia* strain isolated from the lower attine *Apterostigma dentigerum*. Dentigerumycin A 6 is of mixed polyketide/non-ribosomal peptide origin and is active against *Escovopsis* pathogens but not the fungal cultivar [32]. Interestingly, dentigerumycin analogues were also identified in *Macrotermes* fungus-growing termites [33], and the closely related gerumycins were isolated from another *A. dentigerum*-derived *Pseudonocardia* species, as well as strains associated with the higher attine *Trachymyrmex cornetzi*, but these did not show antifungal activity against *Escovopsis* species [34].

The other main type of compounds are polyketide polyene macrolides, a well-known class of antifungal that includes the clinically important drugs nystatin A1 and amphotericin B, which act by binding ergosterol [35]. A novel polyene antifungal named nystatin P1 (7) was isolated from the *A. octospinosus* derived *Pseudonocardia octospinosus* P1 strain. While the structure of 7 has not been fully elucidated, MS/MS and BGC analysis strongly suggest that it has an additional hexose compared to nystatin A1, likely D-mannose linked to the D-mycosamine moiety via a  $\beta$ -1,4 linkage [29,36]. Selvamycin (8) is another novel polyene macrolide, identified from a *Pseudonocardia* strain associated with *A. dentigerum*. It exhibits activity against a range of fungal bioindicator strains; however, its activity against *Escovopsis* has not been reported. The unusual 4-O-methyl-digitoxose glycosylation and the lack of charged groups (Figure 3) makes 8 distinct from other antifungal polyene compounds, and as it showed no evidence of ergosterol binding, it appears to have a novel mechanism of action [37].

Other Actinobacteria have also been recovered from the exoskeletons of fungus-growing ants, most notably *Streptomyces* species. The production of the antifungals candicidin D (9) and antimycin A<sub>1</sub>-A<sub>4</sub> (10a-d) has been

observed for *Streptomyces* species derived from *Acromyrmex* nests [36,38–40]. Interestingly, 9 was highly active against *E. weberi* but not the *L. gongylophorus* cultivar [38], whereas 10a-d compounds showed more generalised activity against both fungal strains [39]. More recently, *Streptomyces* species isolated from the exoskeletons of *Cyphomyrmex* and *Acromyrmex rugosus* workers were found to produce the mono-hydroxypyridines Mer-A2026B (11) and piericidin A (12), along with the ionophores nigericin and dynactin, all of which inhibited the growth of various *Escovopsis* spp. in challenge experiments on agar. Interestingly, these compounds were also found to have anti-leishmanial activity, the first report of antiprotozoal activity by ant-nest compounds [41].

Beyond *Pseudonocardia* and *Streptomyces* species, yeasts isolated from *Atta texana* nests inhibited the growth of *E. weberi*, along with generalist pathogenic and entomopathogenic fungi [42]. *Bulkholderia* species have also been found in fungus-growing ant nests, and one strain isolated from an *Atta sexdens rubropilosa* colony inhibited the growth of *E. weberi* and other fungal pathogens, but not *L. gongylophorus* [43]. These reports hint at the much more complex interplay of different species in the nests of fungus growing ants, beyond the well-studied mutualists and pathogens.

The fungal-symbionts themselves have been implicated in defence against *Escovopsis*; however, reports of their antifungal capabilities have been variable [14]. The unique yeast cultivar maintained by the lower attine *Cyphomyrmex minutus* produced antifungal diketopiperazines [44]. More recently, leafcutter cultivars were shown to inhibit *Escovopsis* strains from lower attine ants, such as *Apterostigma* [45].

### Infighting

As well as producing antifungals to suppress host pathogens, defensive symbionts produce antibacterials to outcompete microbial competitors [29]. Indeed, antagonism between *Pseudonocardia* strains is common, and in challenge experiments pairing strains from across the broader phylogeny, closely related strains were able to inhibit each other [46]. The indolocarbazole 9-methoxyrebeccamycin (13) was produced by *Pseudonocardia* isolated from *A. dentigerum* and inhibited other *Pseudonocardia* strains isolated from the same region [47]. Recently, the thiopeptide GE37468 (14) was isolated from *Pseudonocardia* associated with *Trachymyrmex septentrionalis* ants and shown to inhibit other *Pseudonocardia* strains isolated from *Trachymyrmex* nests [48]. Another *A. dentigerum*-derived *Pseudonocardia* strain produced the angucyclines 6-deoxy-8-O-methylrabelomycin (15) and X-14881, and the glycosylated pseudonocardones. The

non-glycosylated compounds showed some antibacterial and antimalarial activity, but their role in ant nests is yet to be established [49]. Streptomycetes associated with fungus-growing ants also produce antibacterials, most notably actinomycin D (**16**) and actinomycin X<sub>2</sub> (**17**), which inhibited both *Pseudocardia* and other *Streptomyces* species and acted synergistically with **10a-d** compounds. Valinomycins were also produced by several *Streptomyces* spp.; however, these were not active against the ant-nest species tested [39].

### Ant defences

In marked contrast to the *Acromyrmex* leafcutters, *Atta* ants lack cuticular actinomycete cultures. Instead, they rely on phenylacetic acid secretions from their metapleural glands to fend off fungal pathogens. Although phenylacetic acid is also active against generalist fungal pathogens, *E. weberi* appears to be particularly vulnerable, with strains isolated from lower attine ant genera being more susceptible than those from higher attines [50]. The reason for the divergence in disease management strategies for these close relatives remains unclear, as do the comparative benefits of chemical *versus* ‘biological’ pest control. On the face of it, the capacity to recruit bacterial symbionts would appear to offer broader evolutionary adaptability; however, the limited evidence of natural *Escovopsis* outbreaks suggests that while they are more widespread in *Atta* colonies, they lead to colony collapse far less frequently than in *Acromyrmex* nests [50].

A range of aldehyde, ketone and alcohol volatiles detected in metapleural gland secretions of *Apterostigma*, *Acromyrmex* and *Atta* have been shown to have strong antibacterial and antifungal activity [51,52]. These compounds may constitute a further, more generalist line of defence in the nest.

### Defence recruitment

Understanding the role of the *Streptomyces* strains commonly isolated from attine cuticles is a crucial next step. One intriguing hypothesis proposes that the ants specifically recruit or ‘screen in’ antibiotic-producing actinomycetes such as *Streptomyces* species while keeping other bacteria out [12,53]. They achieve this by providing public resources to create a competitive environment attractive for bacterial colonisation, and then use their vertically transmitted *Pseudocardia* mutualist strain to create a demanding environment in which only antibiotic-producers can colonise and survive. These *Pseudocardia* strains are known to make broad-spectrum antibacterial molecules that inhibit most unicellular bacteria, but they do not inhibit *Streptomyces* species, which themselves make multiple antimicrobials and carry multiple antibiotic resistance

genes [29]. The result is an ant cuticle dominated by *Pseudocardia* and *Streptomyces* species, which make several antibacterial and antifungal compounds that are useful to the ants [54].

### Secret weapons

Genome sequencing of bacterial mutualists isolated from fungus-growing ants has shed light on their untapped biosynthetic potential. Analysis of *Pseudocardia* strains associated with *A. echinator*, which split into two phylotypes Ps1 (species name *P. octospinosus*) and Ps2 (*Pseudocardia echinator*), showed that Ps2 strains have the potential to produce new nystatin derivatives that arise from a novel 3-amino-5-hydroxybenzoic acid biosynthetic starter unit and encode several bacteriocins which may be involved in inter-species competition [29]. Recent population genomic analyses of nearly 50 *Pseudocardia* strains isolated from *Apterostigma* ants revealed 27 BGC families, including lassopeptides, siderophores, and terpenes [55,56].

Likewise, although *E. weberi* has a reduced genome size consistent with its parasitic lifestyle, genome sequencing reveals the biosynthetic capabilities of *Escovopsis* strains go well beyond what has been observed in the laboratory, thus far. In *E. weberi* isolated from *Atta cephalotes* 17 putative BGCs were identified [8]. A further five *E. weberi* strains isolated from *Acromyrmex* and *Atta* colonies contained 20–23 putative BGCs each, and in common with the previously sequenced strain, these mainly comprised terpene, type I polyketide synthase and non-ribosomal peptide synthetase BGCs [16].

### Conclusions

A multipartite web of mutualistic and antagonistic interactions exists between the symbionts in fungus-growing ant nests [14]. These ancient systems offer a gateway to a wealth of chemical diversity created by a 50 million-year-old arms race and provide tractable models for understanding the functions of specialized metabolites in nature. While the major players are well-established, recent work suggests that these represent only a fraction of the many microbes present in these colonies, and distinguishing symbionts from opportunistic environmental microbes will be important to develop a full understanding of the interactions at play. These symbiotic interactions are widespread in the plant and animal kingdoms and are all likely driven by chemical communication and chemical warfare. For example, a Kenyan fungus growing plant-ant system is already proving to be an exciting resource for the discovery of new chemical diversity [57,58]. Understanding the roles and regulation of microbial specialized

metabolites in their natural habitats is crucial if we are to unlock and discover the vast range of activities encoded by these organisms.

### Role of the funding source

This work was supported by the Biotechnology and Biological Sciences Research Council (BBSRC) via Institute Strategic Program Project BBS/E/J/00PR9791 to the John Innes Centre, BBSRC responsive mode grants BB/S009000/1 (to BW) and BB/S00811X/1 (to MIH) and Natural Environment Research Council grants NE/J01074X/1 and NE/M015033/1 (to MIH) and NE/M014657/1 (to BW).

### Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

### Acknowledgements

The authors would like to thank our colleagues Dr Sarah Worsley, Dr Neil Holmes and Dr Daniel Heine for their contributions to the work reviewed here. The SEM image of *Pseudonocardia* hyphae was taken by Dr Kim Findlay (JIC), the image of *Escovopsis weberi* was taken by Dr Claudio Greco (JIC), the images of the *Acromyrmex* ant in Figures 1 and 2 and of *Leucoagaricus gongylophorus* were taken by Andrew Davis (JIC), the image of *Streptomyces albidoflavus* S4 was taken by Dr Jörg Barke, and the image of the *Acromyrmex* ant in Figure 3 was taken by Dr Sarah Worsley.

### References

Papers of particular interest, published within the period of review, have been highlighted as:

- \* of special interest
- \*\* of outstanding interest

1. Schultz TR, Brady SG: **Major evolutionary transitions in ant agriculture.** *Proc Natl Acad Sci USA* 2008, **105**:5435–5440.
  2. Currie CR: **A community of ants, fungi, and bacteria: a multilateral approach to studying symbiosis.** *Annu Rev Microbiol* 2001, **55**:357–380.
  3. De Fine Licht HH, Boomsma JJ, Tunlid A: **Symbiotic adaptations in the fungal cultivar of leaf-cutting ants.** *Nat Commun* 2014, **5**:5675.
  4. Mueller UG, Kardish MR, Ishak HD, Wright AM, Solomon SE, Bruschi SM, Carlson AL, Bacci Jr M: **Phylogenetic patterns of ant–fungus associations indicate that farming strategies, not only a superior fungal cultivar, explain the ecological success of leafcutter ants.** *Mol Ecol* 2018, **27**:2414–2434.
  5. Poulsen M, Boomsma JJ: **Mutualistic fungi control crop diversity in fungus-growing ants.** *Science* 2005, **307**:741–744.
  6. Currie CR, Wong B, Stuart AE, Schultz TR, Rehner SA, Mueller UG, Sung G-H, Spatafora JW, Straus NA: **Ancient tripartite coevolution in the attine ant-microbe symbiosis.** *Science* 2003, **299**:386–388.
  7. Osti JF, Rodrigues A: ***Escovopsioides* as a fungal antagonist of the fungus cultivated by leafcutter ants.** *BMC Microbiol* 2018, **18**:130.
- This paper demonstrates that *Escovopsioides* represents another genera of the fungal antagonist in the nests of Leafcutter ants, but has morphological features distinct from *Escovopsis* genera. This expands our knowledge of fungal pathogens and suggests there are many more types of specialised pathogens waiting to be discovered.
8. Currie CR: **Prevalence and impact of a virulent parasite on a tripartite mutualism.** *Oecologia* 2001, **128**:99–106.
  9. de Man Tjb, Stajich JE, Kubicek CP, Teiling C, Chenthamara K, Atanasova L, Druzhinina IS, Levenkova N, Birnbaum SSL, Barribeau SM, *et al.*: **Small genome of the fungus *Escovopsis weberi*, a specialized disease agent of ant agriculture.** *Proc Natl Acad Sci USA* 2016, **113**:3567–3572.
- This paper reported the first genome sequence of an *Escovopsis* species, giving the first insight into its biosynthetic capacity and revealing that specialized metabolites are upregulated when it is co-cultured with the fungal cultivar.
10. Bonadies E, Wcislo W, Gálvez D, Hughes W, Fernández-Marín H: **Hygiene defense behaviors used by a fungus-growing ant depend on the fungal pathogen stages.** *Insects* 2019, **10**:130.
  11. Marsh SE, Poulsen M, Pinto-Tomás A, Currie CR: **Interaction between workers during a short time window is required for bacterial symbiont transmission in *Acromyrmex* leaf cutting ants.** *PLoS One* 2014, **9**, e103269.
  12. Scheuring I, Yu DW: **How to assemble a beneficial microbiome in three easy steps.** *Ecol Lett* 2012, **15**:1300–1307.
  13. Van Arnam EB, Currie CR, Clardy J: **Defense contracts: molecular protection in insect-microbe symbioses.** *Chem Soc Rev* 2018, **47**:1638–1651.
  14. Worsley SF, Innocent TM, Heine D, Murrell CJ, Yu DW, Wilkinson B, Hutchings MI, Boomsma JJ, Holmes NA: **Symbiotic partnerships and their chemical interactions in the leafcutter ants (Hymenoptera: formicidae).** *Myrmecol News* 2018, **27**: 59–47.
  15. Dhodary B, Schilg M, Wirth R, Spiteller D: **Secondary metabolites from *Escovopsis weberi* and their role in attacking the garden fungus of leaf-cutting ants.** *Chem Eur J* 2018, **24**: 4445–4452.
- This paper is one of the first to purify and characterise specialised metabolites from *Escovopsis* species, demonstrating the production of terpene indole alkaloids and cycloarthropsones.
16. Heine D, Holmes NA, Worsley SF, Santos ACA, Innocent TM, Scherlach K, Patrick EH, Yu DW, Murrell JC, Vieira PC, *et al.*: **Chemical warfare between leafcutter ant symbionts and a co-evolved pathogen.** *Nat Commun* 2018, **9**:2208.
- This paper is one of the first to purify and characterise specialised metabolites from *Escovopsis* species, demonstrating the production of compounds active against ants, mutualistic bacteria and the fungal cultivar. The genomes of five *Escovopsis* strains were also sequenced and deposited.
17. Gerardo NM, Mueller UG, Price SL, Currie CR: **Exploiting a mutualism: parasite specialization on cultivars within the fungus-growing ant symbiosis.** *Proc R Soc B* 2004, **271**: 1791–1798.
  18. Marfétán JA, Romero AI, Folgarait PJ: **Pathogenic interaction between *Escovopsis weberi* and *Leucoagaricus* sp.: mechanisms involved and virulence levels.** *Fungal Ecol* 2015, **17**: 52–61.
  19. Varanda-Haifig SS, Albarici TR, Nunes PH, Haifig I, Vieira PC, Rodrigues A: **Nature of the interactions between hypocrealean fungi and the mutualistic fungus of leaf-cutter ants.** *Antonie Leeuwenhoek* 2017, **110**:593–605.
- This paper was the first to demonstrate *Escovopsis* extracts had antibacterial activity.
20. Boya PCA, Fernández-Marín H, Mejía LC, Spatafora C, Dorrestein PC, Gutiérrez M: **Imaging mass spectrometry and MS/MS molecular networking reveals chemical interactions among cuticular bacteria and pathogenic fungi associated with fungus-growing ants.** *Sci Rep* 2017, **7**: 5604-5604.
- This paper was the first to propose *Escovopsis* species were capable of producing shearinine compounds.
21. Xu M, Gessner G, Groth I, Lange C, Christner A, Bruhn T, Deng Z, Li X, Heinemann SH, Grabley S, *et al.*: **Shearinines D–K, new indole triterpenoids from an endophytic *Penicillium* sp. (strain HK10459) with blocking activity on large-conductance calcium-activated potassium channels.** *Tetrahedron* 2007, **63**: 435–444.

22. Little AEF, Currie CR: **Symbiotic complexity: discovery of a fifth symbiont in the attine ant-microbe symbiosis.** *Biol Lett* 2007, **3**:501–504.
23. Attili-Angelis D, Duarte APM, Pagnocca FC, Nagamoto NS, de Vries M, Stielow JB, de Hoog GS: **Novel *Phialophora* species from leaf-cutting ants (tribe Attini).** *Fungal Divers* 2014, **65**: 65–75.
24. Duarte APM, Attili-Angelis D, Baron NC, Forti LC, Pagnocca FC: **Leaf-cutting ants: an unexpected microenvironment holding human opportunistic black fungi.** *Antonie Leeuwenhoek* 2014, **106**:465–473.
25. Little AEF, Currie CR: **Black yeast symbionts compromise the efficiency of antibiotic defenses in fungus-growing ants.** *Ecology* 2008, **89**:1216–1222.
26. Vasse M, Voglmayr H, Mayer V, Gueidan C, Nepel M, Moreno L, Hoog Sd, Selosse M-A, McKey D, Blatrix R: **A phylogenetic perspective on the association between ants (Hymenoptera: formicidae) and black yeasts (Ascomycota: Chaetothiales).** *Proc R Soc B* 2017, **284**:20162519.
27. Currie CR, Poulsen M, Mendenhall J, Boomsma JJ, Billen J: **Coevolved crypts and exocrine glands support mutualistic bacteria in fungus-growing ants.** *Science* 2006, **311**:81–83.
28. Mattoso TC, Moreira DDO, Samuels RI: **Symbiotic bacteria on the cuticle of the leaf-cutting ant *Acromyrmex subterraneus subterraneus* protect workers from attack by entomopathogenic fungi.** *Biol Lett* 2012, **8**:461–464.
29. Holmes NA, Innocent TM, Heine D, Bassam MA, Worsley SF, Trottmann F, Patrick EH, Yu DW, Murrell JC, Schiøtt M, *et al.*: **Genome analysis of two *Pseudonocardia* phylotypes associated with *Acromyrmex* leafcutter ants reveals their biosynthetic potential.** *Front Microbiol* 2016, **7**.
- This paper was the first to publish high-quality genome sequences of attine ant-associated *Pseudonocardia* strains and demonstrated that *Acromyrmex echinator* ants are always associated with one of two distinct *Pseudonocardia* species.
30. Currie CR, Bot ANM, Boomsma JJ: **Experimental evidence of a tripartite mutualism: bacteria protect ant fungus gardens from specialized parasites.** *Oikos* 2003, **101**:91–102.
31. Poulsen M, Cafaro MJ, Erhardt DP, Little AE, Gerardo NM, Tebbets B, Klein BS, Currie CR: **Variation in *Pseudonocardia* antibiotic defence helps govern parasite-induced morbidity in *Acromyrmex* leaf-cutting ants.** *Environ Microbiol Rep* 2010, **2**:534–540.
32. Oh D-C, Poulsen M, Currie CR, Clardy J: **Dentigerumycin: a bacterial mediator of an ant-fungus symbiosis.** *Nat Chem Biol* 2009, **5**:391–393.
33. Wyche TP, Ruzzini AC, Beemelmans C, Kim KH, Klassen JL, Cao S, Poulsen M, Bugni TS, Currie CR, Clardy J: **Linear peptides are the major products of a biosynthetic pathway that encodes for cyclic depsipeptides.** *Org Lett* 2017, **19**: 1772–1775.
34. Sit CS, Ruzzini AC, Van Arnem EB, Ramadhar TR, Currie CR, Clardy J: **Variable genetic architectures produce virtually identical molecules in bacterial symbionts of fungus-growing ants.** *Proc Natl Acad Sci USA* 2015, **112**:13150–13154.
35. Zotchev SB: **Polyene macrolide antibiotics and their applications in human therapy.** *Curr Med Chem* 2003, **10**: 211–223.
36. Barke J, Seipke RF, Gruschow S, Heavens D, Drou N, Bibb MJ, Goss RJM, Yu DW, Hutchings MI: **A mixed community of actinomycetes produce multiple antibiotics for the fungus farming ant *Acromyrmex octospinosus*.** *BMC Biol* 2010, **8**:109.
37. Van Arnem EB, Ruzzini AC, Sit CS, Horn H, Pinto-Tomás AA, Currie CR, Clardy J: **Selvamicin, an atypical antifungal polyene from two alternative genomic contexts.** *Proc Natl Acad Sci USA* 2016, **113**:12940–12945.
- This paper identified a new polyene macrolide antifungal from an *Apterostigma dentigerum*-associated *Pseudonocardia*, with an apparently novel mode of action, demonstrating the exciting capacity for novel chemistry in the fungus-growing ant systems.
38. Haeder S, Wirth R, Herz H, Spiteller D: **Candididin-producing *Streptomyces* support leaf-cutting ants to protect their fungus garden against the pathogenic fungus *Escovopsis*.** *Proc Natl Acad Sci USA* 2009, **106**:4742–4746.
39. Schoenian I, Spiteller M, Ghaste M, Wirth R, Herz H, Spiteller D: **Chemical basis of the synergism and antagonism in microbial communities in the nests of leaf-cutting ants.** *Proc Natl Acad Sci USA* 2011, **108**:1955–1960.
40. Seipke RF, Barke J, Brearley C, Hill L, Yu DW, Goss RJM, Hutchings MI: **A single *Streptomyces* symbiont makes multiple antifungals to support the fungus farming ant *Acromyrmex octospinosus*.** *PLoS One* 2011, **6**, e22028.
41. Ortega HE, Ferreira LLG, Melo WGP, Oliveira ALL, Ramos Alvarenga RF, Lopes NP, Bugni TS, Andricopulo AD, Pupo MT: **Antifungal compounds from *Streptomyces* associated with attine ants also inhibit *Leishmania donovani*.** *PLoS Neglected Trop Dis* 2019, **13**, e0007643.
- This paper was the first to report antiprotozoal activity for compounds isolated from the nests of fungus-growing ants.
42. Rodrigues A, Cable RN, Mueller UG, Bacci M, Pagnocca FC: **Antagonistic interactions between garden yeasts and microfungal garden pathogens of leaf-cutting ants.** *Antonie Leeuwenhoek* 2009, **96**:331–342.
43. Santos AV, Dillon RJ, Dillon VM, Reynolds SE, Samuels RI: **Occurrence of the antibiotic producing bacterium *Burkholderia* sp. in colonies of the leaf-cutting ant *Atta sexdens rubropilosa*.** *FEMS Microbiol Lett* 2004, **239**:319–323.
44. Wang Y, Mueller UG, Clardy J: **Antifungal diketopiperazines from symbiotic fungus of fungus-growing ant *Cyphomyrmex minutus*.** *J Chem Ecol* 1999, **25**:935–941.
45. Birnbaum SS, Gerardo NM: **Patterns of specificity of the pathogen *Escovopsis* across the fungus-growing ant symbiosis.** *Am Nat* 2016, **188**:52–65.
46. Poulsen M, Erhardt DP, Molinaro DJ, Lin T-L, Currie CR: **Antagonistic bacterial interactions help shape host-symbiont dynamics within the fungus-growing ant-microbe mutualism.** *PLoS One* 2007, **9**, e960.
47. Van Arnem EB, Ruzzini AC, Sit CS, Currie CR, Clardy J: **A rebeccamycin analog provides plasmid-encoded niche defense.** *J Am Chem Soc* 2015, **137**:14272–14274.
48. Chang PT, Rao K, Longo LO, Lawton ES, Scherer G, Van Arnem EB: **Thiopeptide defense by an ant's bacterial symbiont.** *J Nat Prod* 2020, **83**:725–729.
- This paper describes a thiopeptide antibiotic isolated from *Trachymyrmex septentrionalis* associated *Pseudonocardia*.
49. Carr G, Derbyshire ER, Caldera E, Currie CR, Clardy J: **Antibiotic and antimalarial quinones from fungus-growing ant-associated *Pseudonocardia* sp.** *J Nat Prod* 2012, **75**: 1806–1809.
50. Fernández-Marín H, Nash DR, Higginbotham S, Estrada C, Zweden JSv, d'Ettorre P, Wcislo WT, Boomsma JJ: **Functional role of phenylacetic acid from metapleural gland secretions in controlling fungal pathogens in evolutionarily derived leaf-cutting ants.** *Proc R Soc B* 2015, **282**:20150212.
51. de Lima Mendonça A, da Silva CE, de Mesquita FLT, da Silva Campos R, Do Nascimento RR, de Azevedo Ximenes ECP, Sant'Ana AEG: **Antimicrobial activities of components of the glandular secretions of leaf cutting ants of the genus *Atta*.** *Antonie Leeuwenhoek* 2009, **95**:295–303.
52. Hogan CT, Jones TH, Zhukova M, Sosa-Calvo J, Adams RMM: **Novel mandibular gland volatiles from *Apterostigma* ants.** *Biochem Systemat Ecol* 2017, **72**:56–62.
53. Barke J, Seipke RF, Yu DW, Hutchings MI: **A mutualistic microbiome: how do fungus-growing ants select their antibiotic-producing bacteria?** *Commun Integr Biol* 2011, **4**: 41–43.

54. Innocent T, Holmes N, Bassam MA, Schiøtt M, Scheuring I, Wilkinson B, Hutchings MI, Boomsma JJ, Yu DW: **Experimental demonstration that screening can enable the environmental recruitment of a defensive microbiome.** *bioRxiv* 2018:375634. This preprint gives new insights into how ants recruit their defensive microbiome.
55. Caldera EJ, Chevrette MG, McDonald BR, Currie CR: **Local adaptation of bacterial symbionts within a geographic mosaic of antibiotic coevolution.** *Appl Environ Microbiol* 2019, **85**. e01580-01519.
56. McDonald BR, Chevrette MG, Klassen JL, Horn HA, Caldera EJ, Wendt-Pienkowski E, Cafaro MJ, Ruzzini AC, Van Arnam EB, Weinstock GM, *et al.*: **Biogeography and microscale diversity shape the biosynthetic potential of fungus-growing ant-associated *Pseudonocardia*.** *bioRxiv* 2019:545640.
57. Qin Z, Munnoch JT, Devine R, Holmes NA, Seipke RF, Wilkinson KA, Wilkinson B, Hutchings MI: **Formicamycins, antibacterial polyketides produced by *Streptomyces formicae* isolated from African *Tetraponera* plant-ants.** *Chem Sci* 2017, **8**:3218–3227.
58. Vikeli E, Widdick DA, Batey SF, Heine D, Holmes NA, Bibb MJ, Martins DJ, Pierce NE, Hutchings MI, Wilkinson B: ***In situ* activation and heterologous production of a cryptic lantibiotic from a plant-ant derived *Saccharopolyspora* species.** *Appl Environ Microbiol* 2019. AEM.01876-01819.