

1
2
3
4 **TITLE**

5 **Prescription Of analgesia in Emergency Medicine (POEM) secondary analysis: an**
6 **observational multicentre comparison of pain relief provided to adults and children with**
7 **an isolated limb fracture and/or dislocation**
8
9

10 **Corresponding author**

11 Dr Sarah J Wilson,
12 Emergency Department, Wexham Park Hospital, Frimley Health NHS Foundation Trust,
13 Wexham Street, Slough, SL2 4HL

14 Sarah.wilson2@nhs.net

15 0300 6154050
16
17
18

19 **Authorship details**

20 Sarah Wilson¹, sarah.wilson2@nhs.net <https://orcid.org/0000-0003-3964-0809>

21 Jane Quinlan², jane.quinlan@ouh.nhs.uk <https://orcid.org/0000-0002-0212-4459>

22 Sally Beer³, sally.beer@ouh.nhs.uk <https://orcid.org/0000-0002-0988-3458>

23 Melanie Darwent³, Melanie.darwent@ouh.nhs.uk <https://orcid.org/0000-0003-0600-0020>

24 Jack R Dainty⁴, jack.dainty@uea.ac.uk <https://orcid.org/0000-0002-0056-1233>

25 James Robert Sheehan⁵ jrs24@doctors.org.uk <https://orcid.org/0000-0002-2885-8629>

26 Liza Keating⁶, liza.keating@royalberkshire.nhs.uk <https://orcid.org/0000-0001-9028-2520>
27
28

29
30 ¹ Emergency Department, Wexham Park Hospital, Frimley Health NHS Foundation Trust, Slough, UK

31 ² Nuffield Department of Anaesthesia, Oxford University Hospitals NHS Foundation Trust, Oxford, UK

32 ³ Emergency Department, John Radcliffe Hospital, Oxford University Hospitals NHS Foundation Trust, Oxford, UK

33 ⁴ Norwich Medical School, University of East Anglia, Norwich, UK

34 ⁵ Department of Anaesthesia, Addenbrookes Hospital, Cambridge University Hospitals NHS Foundation Trust, Cambridge, UK

35 ⁶ Emergency Department, Royal Berkshire Hospital NHS Foundation Trust, Reading, UK

36 **Word count, excluding title page, abstract, references, figures and tables**

37 **1168**
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **ABSTRACT**

4
5 **Background**

6 Acute pain is a common reason for Emergency Department (ED) attendance. Royal College
7 of Emergency Medicine (RCEM) pain management audits have shown national variation and
8 room for improvement. Previous evidence suggests that children receive less satisfactory
9 pain management than adults.

10
11
12 **Methods**

13 POEM (Prescription Of analgesia in Emergency Medicine) is a cross-sectional observational
14 study of consecutive patients presenting to 12 NHS Emergency Departments with an
15 isolated long bone fracture and/or dislocation, and was carried out between 2015 and 2017.
16 Using the recommendations in the RCEM Best Practice Guidelines, pain management in ED
17 was assessed for differences of age (adults vs children) and hospital type (children's vs all
18 patients).

19
20
21 **Results**

22 From the total 8346 patients, 38% were children (median age 8y). There was better
23 adherence to the RCEM guidance for children than adults (24% (766/3196) vs 11%
24 (579/5123)) for the combined outcome of timely assessment, pain score and appropriate
25 analgesia. In addition, children were significantly more likely than adults to receive analgesia
26 appropriate to the pain score (of those with a recorded pain score 67% (1168/1744) vs 52%
27 (1238/2361)). Children's hospitals performed much better across all reported outcomes
28 compared to general hospitals.

29
30
31 **Conclusions**

32 In contrast to previous studies, children with a limb fracture/dislocation are more likely than
33 adults to have a pain score documented and to receive appropriate analgesia. Unexpectedly,
34 children's EDs performed better than general EDs in relation to timely and appropriate
35 analgesia but the reasons for this are not apparent from the present study.
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **INTRODUCTION**

4 Good pain management correlates with positive patient experience and clinical outcomes
5 but, worldwide, ED pain management is often inadequate.[1] Similar to Emergency Medicine
6 organisations internationally, the Royal College of Emergency Medicine (RCEM) Best Practice
7 Guidelines 2014[2] (updated for children 2017[3]) describe standards for acute pain
8 management in the Emergency Department (ED). Multiple RCEM national audits[4] have
9 found wide variation in performance and concluded pain management could be improved.

10
11
12 Prior studies have demonstrated that patient age influences pain management after fracture
13 in the ED with children seemingly receiving inferior treatment.[5] In addition, the type of
14 hospital appears to modify the treatment of pain by clinicians.[6] The reasons for these
15 variations are unclear and need investigation.

16
17
18 This study examines the differences in ED pain management for adults and children
19 presenting with isolated limb fracture and/or dislocation across several UK NHS sites. The
20 findings are examined with the aim of improving recommendations for patient care.

21
22 **METHODS**

23 **Study design**

24
25 This is a pre-planned sub-analysis of a cross-sectional observational study of patients
26 presenting to 12 NHS EDs between 2015 and 2017. Five trauma centres and six trauma units
27 contributed to the final POEM study dataset. A list of participating sites can be found in the
28 online supplementary appendix.

29 The Berkshire Research Ethics Committee (REC 14/SC/0167) and the Confidential Advisory
30 Group (CAG 3-02(c)/2014) approved the study.

31 Inclusion/exclusion criteria, the study sample size calculation and data collection are
32 described in Sheehan et al.[7]

33
34
35 **Data analysis**

36 The primary outcome was the proportion of patients receiving pain management as per the
37 RCEM Best Practice Guidelines: a pain score recorded within twenty minutes of arrival in the
38 ED and analgesia provided appropriate to the pain score. The secondary outcome was the
39 influence of age and setting on pain management provided in the ED by comparing adults
40 and children (<18y). Chi-squared tests were used to examine associations between variables.
41 All analyses were performed using the R Statistics program (R Foundation for Statistical
42 Computing, Vienna, Austria).

43
44
45 **RESULTS**

46 **Participants**

47 Patient characteristics are described in **Table 1**.

48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. Patient characteristics

Note: all missing data <2% unless otherwise stated.

Characteristic	Adult	Children (all) (<18y)	p-value ^b	Children at children's hospitals	Children at mixed hospitals	p-value ^c
Number Analysed	5123	3196		1089	2107	
Age (Median (IQR))	64 (43, 80)	8 (5, 12)	<0.001	8 (4, 11)	9 (5, 13)	<0.001
IMD ^a (Median (IQR))	11.6 (2.0, 45.0)	14.1 (0.5, 79.0)	<0.001	33.2 (14.7,50.4)	10.9 (5.7,19.3)	<0.001
Gender			<0.001			0.895
Male	1878 (36.7%)	1915 (60.1%)		656 (60.2%)	1259 (59.8%)	
Female	3222 (62.9%)	1269 (39.9%)		431 (39.6%)	838 (39.8%)	
Ethnicity	642 missing (12.5%)	272 missing (8.5%)	<0.001	49 missing (4.5%)	223 missing (10.6%)	<0.001
White	4251 (83.0%)	2187 (68.4%)		618 (56.7%)	1569 (74.5%)	
Non-white	230 (4.5%)	737 (23.1%)		422 (38.8%)	315 (15.0%)	
Arrival mode			<0.001			0.272
Self presented	2553 (49.8%)	2783 (87.1%)		945 (86.8%)	1838 (87.2%)	
Ambulance (road and air)	2490 (48.6%)	390 (12.2%)		144 (13.2%)	246 (11.7%)	
Other	60 (1.6%)	17 (0.7%)		0 (0%)	17 (1.1%)	
Location after ED			<0.001			<0.001
Home	2977 (58%)	2541 (80%)		898 (84%)	1643 (79%)	
Hospital	2061 (40%)	604 (19%)		166 (16%)	438 (21%)	
Type of injury			<0.001			0.018
Fracture	4495 (88%)	3088 (97%)		1066 (98%)	2022 (96%)	
Dislocation	466 (9%)	71 (2%)		17 (2%)	54 (3%)	
Fracture/ Dislocation	159 (3%)	35 (1%)		6 (1%)	29 (1%)	
Injury location			<0.001			0.159
Upper limb injury	2906 (57%)	2795 (87%)		966 (89%)	1829 (87%)	
Lower limb injury	2205 (43%)	397 (12%)		123 (11%)	274 (13%)	

ED: emergency department; IMD: index of multiple deprivation.

^aIMD is a numerical score whereby a higher score represents greater deprivation.^bstatistical test for the difference between adults and children^cstatistical test for the difference between children at children's hospitals and children at mixed hospitals

A total of 8346 patients attended the EDs with a limb fracture and/or dislocation. Children comprised 38% (3196) of our population (median age 8 years). Children had a male:female

1
2
3 ratio of 60:40, the reverse of adults (37:63). There is little difference in the attendances by
4 day of the week between adults and children (online supplementary appendix). Children had
5 a greater proportion of upper limb injuries (87%) compared with adults (56%) (online
6 supplementary appendix).
7
8

9 **Pain assessment and management**

10 **Figure 1** summarises the primary outcome data for adults and children. Children were more
11 likely than adults to have an initial assessment (triage) within twenty minutes of arrival (63%
12 (1997/3155) vs 50% (2512/5031); $p<0.001$). Of those seen within twenty minutes of arrival,
13 55% (1093/1997) of children and 46% (1145/2512) of adults had a pain score recorded
14 ($p<0.001$). Irrespective of timings, children were more likely than adults to have a pain score
15 documented (55% (1756/3196) vs 47% (2401/5123); $p<0.001$).
16
17

18 Children were more likely than adults to receive analgesia appropriate to the pain score (of
19 those with a recorded pain score 67% (1168/1744) vs 52% (1238/2361); $p<0.001$). The
20 combined outcome of timely assessment and appropriate analgesia also shows better
21 adherence to RCEM guidance for children than adults (24% (766/3196) vs 11% (579/5123);
22 $p<0.001$).
23
24

25 Children were less likely to have a pain score reassessed than adults (7% (230/3196) vs 11%
26 (545/5123); $p<0.001$) except in the children's only hospitals (15% (163/1089); $p<0.001$).
27

28 Regardless of pain score, 66% (2112/3196) of children received pharmacological analgesia in
29 the ED compared with 53% (2727/5123) of adults ($p<0.001$). This proportion was even
30 greater (76% (826/1089)) in children's only hospitals.
31
32

33 Considering the whole patient journey (Table 2), 23.7% (759/3196) of children had 'self-
34 medicated' before ED arrival (vs adults 17.2% (882/5123)). The other marked difference was
35 that almost one third of adults were given a weak opioid, but only 1.5% of children, all of
36 whom attended mixed hospitals.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 2. Pain score and Analgesia throughout the patient journey

	Adults (n=5123, %)	Children (n=3196, %)	Children at children's hospitals (n=1089, %)	Children at mixed hospitals (n=2107, %)
Pain Score				
No pain score	2722 (53.1%)	1440 (45.1%)	283 (26.0%)	1157 (54.9%)
No pain (pain score 0)	188 (3.7%)	298 (9.3%)	203 (18.6%)	95 (4.5%)
Mild/moderate pain (Pain scores 1-3/4-6)	1726 (33.7%)	1078 (33.7%)	392 (36.0%)	686 (32.6%)
Severe pain (pain scores 7-10)	487 (9.5%)	380 (11.9%)	211 (19.4%)	169 (8.0%)
Analgesia				
Self-medication	882 (17.2%)	759 (23.7%)	226 (20.8%)	533 (25.3%)
Analgesia given by pre- hospital clinician	1735 (33.9%)	355 (11.1%)	120 (11.0%)	235 (11.2%)
Manipulation	1160 (22.6%)	136 (4.3%)	24 (2.2%)	112 (5.3%)
Sedation	548 (10.7%)	56 (1.8%)	12 (1.1%)	44 (2.1%)
Block	790 (15.4%)	21 (0.7%)	13 (1.2%)	8 (0.4%)
Analgesia given in ED	2727 (53.2%)	2112 (66.1%)	826 (75.8%)	1286 (61.0%)
Mild/moderate potency	1482 (28.9%)	1619 (50.6%)	650 (59.7%)	969 (46.0%)
Severe potency	1192 (23.3%)	478 (15.0%)	176 (16.2%)	302 (14.3%)
Non-opioid: paracetamol	1666 (32.5%)	1513 (47.3%)	611 (56.1%)	902 (42.8%)
Non-opioid: NSAID*	635 (12.4%)	1169 (36.6%)	469 (43.1%)	700 (33.2%)
Weak opioid#	861 (16.8%)	31 [§] (1.0%)	0 (0%)	31 (1.5%)
Strong opioid [¶]	1009 (19.7%)	547 (17.1%)	201 (18.5%)	346 (16.4%)
Missing	101 (2%)	32 (1.0%)	1 (0.1%)	31 (1.5%)

*Ibuprofen, diclofenac, ketorolac, naproxen

#codeine, dihydrocodeine, tramadol

[§]dihydrocodeine=1, codeine=29, tramadol=1[¶]morphine, hydromorphone, diamorphine, fentanyl, alfentanil, pethidine, oxycodone, methadone**DISCUSSION**

Children were more likely to have a pain score recorded within twenty minutes of ED arrival and to receive analgesia appropriate to that pain score than adults. This contrasts with previous literature which found that children fared worse than adults in terms of ED provision of analgesia for fractures.[5,6,8] Potential explanations are that children are frequently accompanied by an adult who can act as an advocate; there may be a different staff response to a child in pain; or perhaps children are less inhibited in displaying their pain which prompts staff to react and provide analgesia.

Much of the difference in our data was accounted for by the children's hospitals which provided the best adherence to the RCEM criteria. This is contrary to the findings of Cimpello et al [9] who found no difference between paediatric and general EM physicians in the provision of analgesia to children with extremity fractures.

1
2
3 The increased provision of analgesia for children when compared to adults extended
4 through the whole patient journey including provision of 'self-medication' or by pre-hospital
5 clinicians. This suggests that the difference is not limited only to the hospital environment.
6 Minimal use of weak opioids in children is probably a response to the UK guidance to avoid
7 codeine in children.[10]
8
9

10 There is an unacceptable proportion of patients who appear not to have been offered some
11 means of analgesia. Both POEM and the RCEM audit data demonstrate similar proportions
12 of children without a recorded pain score (45%). Interestingly, in previous analysis from
13 POEM [7] we showed that patients with a pain score recorded were also more likely to
14 receive analgesia. Repeating this analysis split into adults and children indicates that this is
15 almost entirely an effect in children rather than in adults. Over half of adults received ED
16 analgesia irrespective of whether a pain score was recorded. In children, 61.1% received ED
17 analgesia without a pain score and 70.1% received ED analgesia with a pain score,
18 demonstrating that ED analgesia provision was better in children and further improved by
19 recording a pain score. Reassessment of pain was poorly documented, in line with the RCEM
20 data. We found that children's hospitals reassessed pain most often (15%).
21
22

23 Future work should concentrate on learning from the children's hospitals to improve
24 provision of pain relief to all patients.
25
26

27 **Limitations**

28 We are limited by reliance on documentation. We were unable to reliably consider non-
29 pharmacological modalities of pain management such as ice or splints. We have not
30 considered factors such as volume of ED attendances for impact on the ability to provide
31 timely treatment. We have only considered patients with certain injuries and not all causes
32 of pain that present to EDs.

33 We are unable to determine why there are marked differences in pain management
34 between the children's hospitals and the remaining departments.
35
36

37 **CONCLUSION**

38 Children were statistically more likely than adults to have a pain score documented and to
39 receive appropriate analgesia. The two children's hospitals in our study outperformed
40 general EDs for provision of timely and appropriate analgesia to children. It is unclear why
41 this might be, and we recommend this should be explored further to improve pain
42 management for all patients.
43
44
45
46

47 **What is already known on the subject?**

48 There is mixed evidence for the effect of age on the provision of analgesia in the Emergency
49 Department.
50

51 **What this study adds:**

52 For patients with an isolated long bone fracture/dislocation, children receive better pain
53 management than adults in the ED. There is a difference in performance between children's
54 hospital EDs and general EDs.
55
56

57 **Figure 1: Primary outcomes and other metrics split by age and department type**

58
59
60

REFERENCES

1. Sampson FC, Goodacre SW, O’Cathain A. Interventions to improve the management of pain in emergency departments: systematic review and narrative synthesis. *Emerg Med J*. 31(e1):e9–18.
2. Royal College of Emergency Medicine. Management of Pain in Adults Best Practice Guideline 2014 [Internet]. 2014 [cited 2020 Mar 2]; Available from: [https://www.rcem.ac.uk/docs/College Guidelines/5w. Management of Pain in Adults \(Revised December 2014\).pdf](https://www.rcem.ac.uk/docs/College Guidelines/5w. Management of Pain in Adults (Revised December 2014).pdf)
3. Royal College of Emergency Medicine. Pain in Children Best Practice Guideline 2017 [Internet]. 2017 [cited 2020 Mar 2]; Available from: [https://www.rcem.ac.uk/docs/RCEM Guidance/RCEM Pain in Children - Best Practice Guidance \(REV Jul 2017\).pdf](https://www.rcem.ac.uk/docs/RCEM Guidance/RCEM Pain in Children - Best Practice Guidance (REV Jul 2017).pdf)
4. Royal College of Emergency Medicine. Pain in Children National Audit Report 2017-18 [Internet]. [cited 2020 Mar 2]; Available from: [https://www.rcem.ac.uk/docs/QI + Clinical Audit/Pain in Children 2017_18 National Report \(Oct 2018\).pdf](https://www.rcem.ac.uk/docs/QI + Clinical Audit/Pain in Children 2017_18 National Report (Oct 2018).pdf)
5. Brown JC, Klein EJ, Lewis CW, Johnston BD, Cummings P. Emergency department analgesia for fracture pain. *Ann Emerg Med*. 42(2):197–205.
6. Petrack EM, Christopher NC, Kriwinsky J. Pain management in the emergency department: Patterns of analgesic utilization. *Pediatrics*. 99(5):711–4.
7. Sheehan JR, Wilson S, Quinlan J, Beer S, Darwent M, Dainty JR et al. Prescription Of analgesia in Emergency Medicine (POEM): a multicentre observational survey of pain relief in patients presenting with an isolated limb fracture and/or dislocation. *Br J Pain*. :204946371985851. Available from: <http://journals.sagepub.com/doi/10.1177/2049463719858513>
8. Boccio E, Wie B, Pasternak S, Salvador-Kelly A, Ward MF, D’Amore J. The relationship between patient age and pain management of acute long-bone fracture in the ED. *Am J Emerg Med*. 32(12):1516–9.
9. Cimpello LB, Khine H, Avner JR. Practice Patterns of Pediatric Versus General Emergency Physicians for Pain Management of Fractures in Pediatric Patients. Vol. 20, *Pediatric Emergency Care*. p. 228–32.
10. Medicines and Healthcare products Regulatory Agency. Codeine for cough and cold:restricted use in children. *Drug Saf Updat*. 8(9):2.

1
2
3 **Footnotes**

4 **Acknowledgements**

5 We thank the investigators and participants at the participating sites for their involvement in
6 and support of the survey. Particular thanks to David Porter (TVSM CRN) and Carys Jones
7 (RBH) for providing ongoing support and encouragement for the study. Thanks also to the
8 PPI representatives and all those who have in some way contributed to the conception,
9 conduction and reporting of the POEM survey. After publication of the planned primary and
10 secondary analyses, the trial data can be shared upon reasonable request to the
11 corresponding author. The views expressed are those of the authors and not necessarily
12 those of the RCEM or the NIHR.

13
14
15 **Contributors**

16 LK and JQ conceived the survey. JS was the chief investigator. LK, JQ, MD, SW and Peter
17 Thomas were co-investigators on the protocol. Martyn Ezra drew up the initial statistical
18 strategy and JRD performed the statistical analysis. SB contributed to the entirety of the
19 project together with the other investigators. SW wrote the first draft of the manuscript; all
20 authors revised this draft. All authors read and approved the final version.

21
22
23 **Funding**

24 Initial funding was received from an RCEM research grant and the study was subsequently
25 adopted onto the NIHR portfolio.

26
27
28 **COI disclosure**

29 All authors have completed the ICMJE uniform disclosure form at
30 www.icmje.org/coi_disclosure.pdf and declare: JRD reports personal fees from Royal
31 Berkshire Hospital during this analysis of the study and personal fees from Reading
32 University during the conduct of the study. All remaining authors did not declare any
33 interests.

34
35
36 **Patient and Public Involvement (PPI) statement**

37 PPI representatives worked with us to refine the research question and protocol. At the end
38 of the study, the PPI representatives met with the study steering group to review the
39 findings. Their opinion was sought about the interpretation of the results. One of our PPI
40 representatives has read and reviewed the submitted and revised manuscripts.


41
42
43 **Collaborators**

44 Stuart Hartshorn, Katy Pettit, Karen Faulkner and Anna Smith (Birmingham Children's
45 Hospital, Birmingham's Women's and Children's NHS Foundation Trust); Rachel Wharton,
46 Sarah Moreton, Denise Maitland and Stephanie Jones (Dorset County Hospital NHS
47 Foundation Trust); Alison Walker and Rachel Worton (Harrogate and District NHS
48 Foundation Trust); Peter Thomas, Antoanela Colda, Charlotte White and Gill Ritchie (Milton
49 Keynes University Hospital NHS Foundation Trust); Melanie Darwent, Sally Beer, Alexis
50 Espinosa, Dominique Georgiou, Louise Findlay, Holly Coles, Jose Martinez, Alex Novak,
51 Amanda Budden, Soubera Rymell, Peter Chater-Lea, Sandy Farrow, Hannah Ward and Neil
52 Dawson (Oxford University Hospitals NHS Foundation Trust—John Radcliffe Hospital and
53 Horton General Hospital); James Sheehan, Liza Keating, Sarah Kempster, Katie Keating-
54 Fedders, Will Orchard, Christina Ambrose, Carys Jones, Justine Loh, Nicola Jacques, Gemma
55 Grimwood, Nyla Haque, Charlotte Philips, Victoria Murray, Carl Evans, Rachel Brooke, Lucie
56 Tesarova and Claire Burnett (Royal Berkshire Hospital NHS Foundation Trust); Alasdair Gray,
57 Mia Paderanga, Adam Lloyd, Kate Easterford, Emma Ward and Rachel O'Brien (Royal
58 Infirmary of Edinburgh); Stuart Gormley, Nick Weatherhill and Shammi Ramlakha (Sheffield
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Children’s NHS Foundation Trust); Anthony Kehoe, Jason Smith, Rosalyn Squire and Alison Jeffery (University Hospitals Plymouth NHS Trust); Sarah Wilson, Louise Chandler, Joana Da Rocha, Saheli Das, Louise Foster, Gail Warren, Thanuja Weerasinghe, Clare Gargini and Maria Cruz (Wexham Park Hospital, Frimley Health NHS Foundation Trust).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45



Supplementary Data**Appendix 1: Alphabetical list of participating sites**

Birmingham Children's Hospital, Birmingham Women's and Children's NHS Foundation Trust
 Dorset County Hospital NHS Foundation Trust
 Harrogate and District NHS Foundation Trust
 Milton Keynes University Hospital NHS Foundation Trust
 Oxford University Hospitals NHS Foundation Trust (John Radcliffe Hospital, Horton General Hospital)
 Royal Berkshire Hospital NHS Foundation Trust
 Royal Infirmary of Edinburgh
 Sheffield Children's NHS Foundation Trust
 University Hospitals Plymouth NHS Trust
 Wexham Park Hospital, Frimley Health NHS Foundation Trust

Appendix 2: Attendance by day of the week and time of day

	Adults	Children
Arrival in ED (day of week)	2 missing (0%)	1 missing (0%)
Monday	737 (14%)	436 (14%)
Tuesday	722 (14%)	436 (14%)
Wednesday	683 (13%)	409 (13%)
Thursday	694 (14%)	507 (16%)
Friday	650 (13%)	453 (14%)
Saturday	781 (15%)	433 (14%)
Sunday	854 (17%)	521 (16%)
Arrival in ED (time of day)	1 missing (0%)	1 missing (0%)
Midnight to 8am	593 (12%)	65 (2%)
8am to 4pm	2477 (48%)	1488 (47%)
4pm to midnight	2052 (40%)	1642 (51%)

Appendix 3: Bone injured

	Adults	Children
Bone	12 missing (0.2%)	4 missing (0.1%)
Humerus	634 (12%)	476 (15%)
Clavicle	291 (6%)	360 (11%)
Acromioclavicular joint	35 (1%)	5 (0%)
Glenohumeral joint (Shoulder)	301 (6%)	30 (1%)
Sternoclavicular joint	2 (0%)	0 (0%)
Elbow joint	39 (1%)	33 (1%)
Radius	1163 (23%)	1144 (36%)
Ulna	113 (2%)	78 (2%)
Wrist	0 (0%)	0 (0%)
Radius/Ulna	328 (6%)	669 (21%)
Femur	244 (5%)	63 (2%)
Hip joint	101 (2%)	0 (0%)
Knee joint	4 (0%)	2 (0%)
Tibia	167 (3%)	164 (5%)
Fibula	528 (10%)	83 (3%)
Tibia/Fibula	235 (5%)	84 (3%)
Ankle	67 (1%)	1 (0%)
Neck of femur	859 (17%)	0 (0%)