


Journal Pre-proof

Repositioning for pressure injury prevention in adults: An abridged Cochrane systematic review and meta-analysis.

Brigid M Gillespie , Rachel M Walker , Sharon L Latimer ,
Lukman Thalib , Jennifer A Whitty , Elizabeth McInnes ,
Ishtar Lockwood , Wendy P Chaboyer

PII: S0020-7489(21)00121-8
DOI: <https://doi.org/10.1016/j.ijnurstu.2021.103976>
Reference: NS 103976



To appear in: *International Journal of Nursing Studies*

Received date: 17 March 2021
Revised date: 27 April 2021
Accepted date: 1 May 2021

Please cite this article as: Brigid M Gillespie , Rachel M Walker , Sharon L Latimer , Lukman Thalib , Jennifer A Whitty , Elizabeth McInnes , Ishtar Lockwood , Wendy P Chaboyer , Repositioning for pressure injury prevention in adults: An abridged Cochrane systematic review and meta-analysis., *International Journal of Nursing Studies* (2021), doi: <https://doi.org/10.1016/j.ijnurstu.2021.103976>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Published by Elsevier Ltd.

TITLE

Repositioning for pressure injury prevention in adults: An abridged Cochrane systematic review and meta-analysis.

AUTHORS

Brigid M Gillespie^{1,2,3}, Rachel M Walker^{1,3,4}, Sharon L Latimer^{1,2,3}, Lukman Thalib¹, Jennifer A Whitty⁵, Elizabeth McInnes^{1,6}, Ishtar Lockwood¹, Wendy P Chaboyer^{1,3}

¹NHMRC Centre of Research Excellence in Wiser Wound Care, Menzies Health Institute Queensland

²Gold Coast University Hospital, Gold Coast Health, Southport, Australia.

³School of Nursing and Midwifery, Griffith University, Brisbane and Gold Coast, Australia.

⁴Division of Surgery, Princess Alexandra Hospital, Metro South Health, Brisbane, Australia.

⁵Health Economics Group, Norwich Medical School, Faculty of Medicine and Health Sciences, University of East Anglia, Norwich, UK.

⁶Nursing Research Institute, St Vincent's Health Australia Sydney, St Vincent's Hospital Melbourne & Australian Catholic University, Fitzroy, Melbourne, Australia

Contact address: Brigid M Gillespie, School of Nursing and Midwifery, Griffith University, Brisbane, Queensland, Australia. b.gillespie@griffith.edu.au

*This article is based on a Cochrane Review published in the Cochrane Database of Systematic Reviews (CDSR) 2020, Issue 6, DOI: 10.1002/14651858.CD009958.pub3 (see www.cochranelibrary.com for information). Cochrane Reviews are regularly updated as new evidence emerges and in response to feedback, and the CDSR should be consulted for the most recent version of the review.

Abstract

Background: A pressure injury is an area of localised damage to the skin and underlying tissues. Patient repositioning is an important prevention strategy, as those with limited mobility are at increased risk of developing pressure injury.

Objectives: To assess the clinical and cost-effectiveness of repositioning schedules on the prevention of pressure injury in adults.

Design: Systematic review and meta-analysis.

Data sources: The Cochrane Wounds Specialised Register; the Cochrane Central Register of Controlled Trials; MEDLINE (Ovid); Embase (Ovid) and Cumulative Index of Nursing and Allied Health Literature Plus (EBSCO) were searched in February 2019. No restrictions were applied to language or date of publication.

Review methods: Studies were eligible if they were randomised controlled trials including cluster trials, published or unpublished, and undertaken in any healthcare setting that assessed the clinical and/or cost effectiveness of repositioning schedules for prevention of pressure injury in adults. Methodological quality of the studies was independently assessed by three authors. Heterogeneity between studies was assessed using the I^2 statistic, and the pooled risk ratios along with their 95% confidence intervals were estimated using either fixed and random effects models, as indicated. Grading of Recommendations Assessment, Development and Evaluation was used to appraise the certainty of evidence.

Results: Eight eligible trials involving 3,941 participants published between 2004 and 2018 were identified. Trials compared either different repositioning frequencies or positioning regimens. Three trials (1074 participants) compared 2-hourly with 4-hourly repositioning (risk ratio 1.06, 95% confidence interval 0.80 to 1.41; $I^2 = 45\%$). Two other trials (252

participants) compared a 30-degree tilt with a 90-degree tilt (risk ratio 0.62, 95% confidence interval 0.10 to 3.97; $I^2 = 69\%$).

Only two trials included economic analyses, both amongst nursing home residents. One study estimated the costs of repositioning to be Canadian dollars \$11.05 and Canadian dollars \$16.74 less per resident per day for the 3-hourly or 4-hourly regimens, respectively, when compared to 2-hourly regimen. The second study reported 3-hourly repositioning using a 30-degree tilt to cost €46.50 (95% confidence interval €1.25 to €74.60) less per patient in nursing time compared with 6-hourly repositioning with a 90-degree lateral rotation.

Conclusion: It remains unclear which repositioning frequencies or positions are most effective in preventing pressure injury in adults. There is limited evidence to support the cost effectiveness of repositioning frequencies and positions.

Registration: Cochrane protocol published in 2012.

What is already known about the topic?

- ◁ Pressure injuries are areas of localised damage to the skin or underlying tissue, often over a bony prominence and generally caused by pressure and/or shear.
- ◁ Immobility is a major risk factor for the development of pressure injury and is an important component of risk assessment.
- ◁ Repositioning is an important strategy that redistributes pressure and is recommended in international pressure injury prevention clinical practice guidelines.

What this paper adds

- ◁ The body of evidence on the clinical and cost effectiveness of particular positions and repositioning frequencies for pressure injury prevention in adults is low quality.
- ◁ There is a paucity of evidence on the effectiveness of regular repositioning. Given the lack of evidence does not prove a lack of effect, repositioning remains an important pressure injury prevention strategy and recommended by most guidelines.
- ◁ Findings from the eight clinical trials and two economic analyses included in this review suggest the evidence to support the use of one particular repositioning frequency and position over another to prevent pressure injuries is low in quality and limited in amount. It remains unclear which position or frequency of repositioning is the most effective in reducing pressure injury development.

Keywords: Pressure ulcer, repositioning, randomised controlled trial, systematic review, meta-analysis, cost-effectiveness

1. Introduction

A pressure injury is an area of underlying tissue damage caused by shear or pressure for a prolonged period, often occurring over bony prominences ². While the terms pressure 'ulcer' or 'bed sore' are also used to describe pressure injuries, the term pressure 'injury' is used in this paper in accordance with recent international guidelines ². Hospital acquired pressure injuries are preventable adverse events and are associated with increased risk of mortality, significantly higher hospitalisation costs, and longer lengths of hospital stay for patients ³⁻⁵.

The results of a recent systematic review and meta-analysis undertaken to describe the global prevalence and incidence of pressure injuries indicated that pooled prevalence of hospital-acquired pressure injuries was 12.8% (95% confidence interval 11.8 to 13.9), with an incidence rate of 5.4 per 10,000 patient days ⁶. The most frequently occurring locations of pressure injury identified in this same systematic review were the sacrum, heels, and hips ⁶. Costs of pressure injury treatment varies between countries but is considerably higher in developed countries. For example, a cost analysis of 273 hospitals in the United Kingdom showed that pressure injuries cost £750 million ² per annum resulting in the loss of 26 healthy life years ⁷. Estimates based on Australian data suggest that pressure injury treatment costs \$1.8 billion Australian dollars annually ⁸, while in the United States, the national cost of hospital acquired pressure injuries has been estimated at US\$26.8 billion per annum ⁹. A large proportion of these healthcare costs relate to nursing time associated with the delivery of care ^{8 10}.

Clinical guidelines recommend repositioning for the prevention of pressure injuries in at risk patients ². Repositioning regimes involve frequency (e.g. 2-hourly), positioning using tilt and/or position (lateral, supine, prone), and are used to redistribute pressure between the

body and support surface⁵. Repositioning is believed to work as a prevention strategy by reducing the duration of pressure on tissues, decreasing tissue hypoxia and thus risk of pressure ulceration^{11 12}. However, recommendations for routine repositioning are largely based on findings from small observational studies conducted more than 25-years ago¹³.

A systematic review synthesising evidence for effectiveness of repositioning in adults was undertaken. The aim was to assess the clinical and cost-effectiveness of repositioning schedules on the prevention of pressure in adults regardless of risk in any healthcare setting.

2. Materials and methods

2.1. Design

This systematic review is an abridged version based on our 2020 Cochrane review¹⁴ (updated from the original review, published in 2014¹⁵; the protocol was published in 2012¹⁶). It followed the recommendations from Preferred Reporting Items for Systematic Review and Meta-Analysis (PRISMA) guidelines for reporting systematic reviews and meta analyses¹⁷.

2.2 Inclusion/exclusion criteria

Publications identified in the search of the five databases were combined and duplicates removed. Randomised controlled trials and cluster-controlled trials cluster-controlled trials were included, regardless of the cluster group (i.e. patient, nurse, hospital). Cross-over trials and quasi-randomised studies were excluded as were all other designs.

All studies were reviewed for eligibility against the population, intervention, comparison, outcomes (PICO) criteria. The population included adult patients admitted to any healthcare or long-term care facility without an existing pressure injury at baseline. Studies where the

only systematic differences were due to varying repositioning frequencies were included. So were the trials comparing *different positions* for repositioning, where the systematic difference between groups were due to such positionings. Studies comparing any repositioning regimen with standard practice, were also included.

The primary outcome measure was the proportion of participants with a new pressure injury (i.e. cumulative incidence) of any stage or location/body region, measured in the study as either primary or as a secondary outcome. Pressure injury stage was defined according to published criteria^{2 13}, or as defined by study authors. Secondary outcomes included health-related quality of life, procedural pain, patient satisfaction, and costs (e.g. costs of pressure injury prevention, related health practitioner time or visits, costs avoided by pressure injury prevention).

Comparative full and partial economic evaluations conducted within the framework of eligible randomised controlled trials, or cluster-controlled trials reporting information such as estimates of resource use or costs associated with repositioning and a comparator were included as part of the review of health economic evidence. Only health economics studies conducted alongside effectiveness studies were considered.

2.3 Search strategy

Electronic databases searched to identify relevant clinical trials included the Cochrane Wounds Specialised Register of Controlled Trials (; 2019); Ovid MEDLINE; Ovid Embase; EBSCO Cumulative Index of Nursing and Allied Health Literature Plus; and the National Health Service Economic Evaluation Database (2015). All searches were on 12 February 2019. The US National Institutes of Health Ongoing Trials Register, ClinicalTrials.gov, the World Health Organization (WHO) International Clinical Trials Registry Platform and the

European Clinical Trials Register were searched 10 March 2019. To identify economic studies, filters developed by the Centre for Reviews and Dissemination were applied to the Ovid MEDLINE, Ovid Embase, and EBSCO Cumulative Index of Nursing and Allied Health Literature Plus searches (CRD 2013).

The reference lists of included trials, relevant systematic reviews, meta-analyses, and health technology assessment reports were also searched to identify other potentially eligible trials or supplementary publications. Search filters were applied to databases, including the Cochrane Highly Sensitive Search Strategy for identifying randomised trials in MEDLINE; the Ovid Embase filter; and the trial filters which were applied to the Cumulative Index of Nursing and Allied Health Literature Plus searches. No restrictions were applied to language, date of publication, or study setting.

2.4. Study selection

Three review authors independently assessed all titles and abstracts of retrieved citations against the eligibility criteria. Full reports of all potentially relevant trials were retrieved for further assessment of eligibility. Where there were discrepancies, review authors discussed and made decisions by consensus.

2.5. Data collection process and extraction

Data were independently extracted from included studies by three review authors. A specifically designed data collection tool was used to extract information (e.g., author, title, journal title, year of publication, country; healthcare setting; eligibility criteria; sample size; intervention; primary and secondary outcome measures). If data were missing from reports, attempts were made to contact authors. Data were entered into Review Manager 5

software (Review Manager 2014) by one author and a data check for accuracy was performed by two review authors.

2.6. Risk of bias assessment

Two review authors independently assessed the risk of bias of eligible trials using the Cochrane tool for assessing risk of bias¹⁸. The domains randomisation, allocation concealment, blinding of outcome assessors, and incomplete outcome data were assessed as low risk of bias, high risk of bias, or unclear risk of bias. Overall high risk of bias was defined as if a trial was rated as 'high' for any one of the three key domains.

Assessment of risk of bias is presented in the 'Risk of bias' summary figure that details reviewers' judgements in a cross-tabulation of studies. Where authors did not report validity criteria, these trials were recorded as being at unclear risk of bias. During study selection, data extraction, and assessment of risk of bias, any disagreements between review authors were resolved by consensus or by referral to another review author.

2.7. Data analysis

For measures of treatment effect, effect estimates for dichotomous outcomes (e.g. relative proportions of patients developing pressure injuries during follow-up) are reported as risk ratios with 95% confidence intervals¹⁹. In the absence of clinical and methodological heterogeneity we conducted a fixed effects analysis. A random effect model was used when heterogeneity exceeded 50%. The Chi-squared and I^2 statistics were used to calculate heterogeneity and reported as the pooled estimate together with its 95% confidence interval. Where appropriate, the data are presented using forest plots. The decision to pool data depended on the availability of outcome data and the assessment of between-trial heterogeneity. Quantitative data were analysed using Review Manager 5²⁰.

Where possible, costs relating to implementing pressure injury prevention and treatment strategies (including complications or infections), health practitioner time or visits, duration or costs of hospital stay for pressure injury and associated adverse events were extracted. Where reported, indirect costs to society associated with pressure injury such as lost productivity were also extracted.

2.8. Summary of findings and assessment of the certainty of the evidence

The 'Summary of findings' tables provide key information about the certainty of the evidence, the magnitude of the intervention effects, the sum of the available data for the primary outcome and secondary outcomes²¹ and an overall grading of the body of evidence related to these outcomes using the Grading of Recommendations, Assessment, Development and Evaluations assessment^{15 16}. Grading of Recommendations, Assessment, Development and Evaluations assesses the certainty of a body of evidence as high, moderate, low, or very low, which represents the extent to which one can be confident that an estimate of effect is close to the true effect¹⁶. The level of certainty is downgraded according to risk of bias, precision of the effect estimate, consistency of individual study results, how directly the evidence answers the question(s) of interest, and risk of reporting and publication biases^{22 23}.

3. Results

3.1. Study characteristics

The study flow diagram is shown in Figure 1. This search yielded 463 intervention records in addition to 18 records from clinical trial registries, resulting in a total of 305 unique records after duplications removed. Fourteen full-text articles and 18 records from clinical trials registries were retrieved. Eight trials (three from the previous review¹) were included in this

review, representing a total of 3941 participants. Two of the eight included trials were cluster-controlled trials, and studies were conducted in Europe (n=4)²⁴⁻²⁷, North America (n=2)^{28,29}, and Asia (n=2)^{30,31}.

Journal Pre-proof




Figure 1. Study flow diagram for clinical studies (undertaken in February 2019)

Three of the eight trials assessed different combinations of repositioning frequencies (2-, 3-, 4- and 6-hourly) ^{24 25 28} and three compared different tilt positions (30°, 45°, or 90°) ^{26 27 31}.

The Pickham trial assessed whether the feedback provided from a wearable patient sensor would increase turning compliance using a 2-hourly turning regimen with 20° tilt ²⁹. Zhou 2014 compared a prone position with a standard supine position ³⁰. Further details regarding the characteristics of included studies are outlined in Table 1.

Electronic searches for this update yielded 237 economic evaluation records. Of these, 236 records were excluded because they did not meet the inclusion criteria. Two economic sub studies met the inclusion criteria ^{32 33} (Figure 2).

Journal Pre-proof




Figure 2 Study flow diagram for economic studies (undertaken in February 2019)

Table 1 Characteristics of included randomised controlled trials (listed in reverse chronological order)

Author, year	Setting and participants	Study design and intervention	Outcomes
Pickham 2018	<p>Location: Intensive care unit (n=2) in a large academic medical centre in California, USA.</p> <p>Number of participants: n=1312 (intervention: n=659; control: n=653).</p> <p>Inclusion criteria: Critically ill medical, surgical and trauma patients</p> <p>Exclusion criteria: Patients less than 18 years of age; patients with an issue preventing effective sensor adhesion (i.e. a sternal dressing) or known adhesive sensitivity; acuity precluding participation; patient refusal</p>	<p>Study design: pragmatic, investigator-initiated, open-label, single-site, randomised clinical trial.</p> <p>Aims: Assess the clinical effectiveness of a wearable patient sensor to improve care delivery and patient outcomes by increasing the total time with turning compliance and preventing pressure injuries in acutely ill patients.</p> <p>Group A (experimental): Optimal turning: all participants had a sensor applied. Participants received care from nurses who had access to a User Dashboard that provides visual advisories for patient turning, based on data obtained from a wearable patient sensor (Leaf Healthcare Inc). Turning regimen 2-hourly.</p> <p>Group B (control): All participants had a sensor applied. Participants received care from nurses who DID NOT have access to a User Dashboard that provides visual advisories for patient turning. Instead, these participants received standard care practices, patient turning initiated by nurses as necessary. Turning regimen 2-hourly.</p>	<p>Primary outcome: hospital acquired pressure injury</p> <p>Secondary outcomes: Total time with turning compliance.</p> <p>Time points: First 72 hours in ventilator-dependent participants.</p>

Ghezjeljeh 2017	<p>Location: intensive care unit in selected government hospitals in Tehran, Iran.</p> <p>Number of participants: n=120</p> <p>Inclusion criteria: Adults aged 18 years; no history of ventilator-acquired pneumonia; hospitalised in intensive care unit; undergoing mechanical ventilation support for 8 hours following hospitalisation; no spinal or unstable pelvic fractures.</p> <p>Exclusion criteria: Patient death; Remaining in selected positions for less than 6 hours over the last 24 hours; history of pressure injury with elevation of head of bed to 45°.</p>	<p>Study design: 3-group randomised clinical trial.</p> <p>Aims: Primary aim was to compare the effect 30° and 45° head of bed tilts on the incidence of ventilator-acquired pneumonia. Incidence of pressure injury was a secondary outcome.</p> <p>Group A: 30° HOB tilt (n = 40). Group B: 45° HOB tilt (n = 40). Group C: routine position (n = 40).</p> <p>Standard care across all groups: Other interventions relative to changing bed sheets, lifting patient rather than dragging, and 2-hourly position changes were performed for participants in all groups.</p>	<p>Primary outcome: unrelated to review outcomes.</p> <p>Secondary outcomes: "the mean probability of pressure injury within 3 days of the analysis of variance test showed that there was no statistically significant difference relative to the mean of the probability of pressure ulcer according to the Braden scale (P = 0.652). Furthermore, none of the patients in the groups suffered from pressure ulcer after 3 days".</p> <p>Time points: participants followed through for 3 days and their skin checked 2-hourly.</p>	High
Manzano 2014	<p>Location: Mixed ICUs (n=2) of a university hospital in southern Spain.</p> <p>Number of participants:</p>	<p>Study design: RCT using 2 groups.</p> <p>Aims: Compare the effectiveness of repositioning every 2 or 4 hours for preventing pressure injury (stage 2) development in intensive care unit</p>	<p>Primary outcome: Occurrence of a new pressure injury (stage 2) at any anatomic site between enrolment in the study and intensive care unit discharge.</p>	Low

	n=329. Inclusion criteria: Critically ill adults; no pressure injury at intensive care unit admission; received invasive mechanical ventilation for at least 24 hours between February 2009 and January 2011. Exclusion criteria: Pregnancy; < 18 years; not being on an alternating air-pressure mattress (due to lack of availability); weight greater than 140 kg or less than 45 kg (as per alternating air-pressure mattress specifications); refusal to consent; mechanical ventilation for more than 48 hours before enrolment in the study; and inclusion in a related trial.	patients under mechanical ventilation. Secondary aims: compare clinical outcomes relative to motility, intensive care unit and hospital length of stay, mechanical ventilation duration, adverse/safety events, and nursing workload. Group A: 2-hourly repositioning (n = 165). Group B: 4-hourly repositioning (n = 164). Standard care across all groups: All participants had the same alternating air-pressure mattress. Standard sedation and analgesia consisted of fentanyl plus propofol or midazolam. The weaning protocol included the daily interruption of sedatives and spontaneous awakening trials.	Secondary outcomes: unrelated to review outcomes. Time points: follow-up for 24 hours.	
Zhou 2014	Location: intensive care unit (n=1) in Beijing, China Number of participants: n=116	Study design: 2-armed randomised controlled trial with a 28-day follow-up period Aims: Examine the effects of prone positioning on the occurrence of pressure injury (secondary outcome).	Primary outcome: unrelated to review outcomes Secondary outcomes: occurrence of pressure injury Time points: daily for up to 72 hours	Unclear

	<p>Inclusion criteria: -</p> <p>Exclusion criteria: -</p> <p>unclear, details not available from translation.</p>	<p>Group A (intervention): Prone position for 10 hours, alternating 4 to 6 hours supine, and then repositioned in the prone position for another 10 hours.</p> <p>Group B (control): Supine position with standard care</p>		
Bergstrom 2013	<p>Location: Nursing homes in the US (n=20) and Canada (n=7)</p> <p>Number of participants: n=967</p> <p>Inclusion criteria: Nursing home residents; aged 65 years; no existing pressure injuries; Braden scale either moderate (13 to 14) or high (11 to 12); limited mobility (3 on Braden subscale of mobility).</p> <p>Exclusion criteria: individuals deemed not competent to provide consent.</p>	<p>Study design: randomised controlled trial</p> <p>Aims: Determine the effectiveness of 3 repositioning (turning) schedules (2-, 3-, 4-hourly) for prevention of pressure injury in nursing home residents</p> <p>Group A: repositioning every 2 hours/± 30 minutes of scheduled time: (n = 335)</p> <p>Group B: repositioning every 3 hours/± 30 minutes of scheduled time: (n = 333)</p> <p>Group C: repositioning every 4 hours/± 30 minutes of scheduled time: (n = 299)</p> <p>Standard care across all groups: all groups repositioned on high-density foam mattresses.</p>	<p>Primary outcome: pressure injuries on sites susceptible to pressure when lying in bed (coccyx or sacrum, trochanter, heel) weekly. Stage 1 pressure injury identified on 2 consecutive days excluded false positives caused by reactive hyperaemia.</p> <p>Secondary outcomes: none reported.</p> <p>Time points: weekly follow-up for 3 weeks (21 days).</p>	Unclear
Moore 2011	<p>Location: Long term aged facilities (n=12) in Ireland</p> <p>Number of participants:</p>	<p>Study design: 2-arm cluster-controlled trial with a 4-week (28-day) follow-up period</p>	<p>Primary outcome: incidence of all pressure injuries during a 28-day period</p>	High

n=213	Inclusion criteria:	Aims: Examine whether repositioning using 30° tilt and 3-hourly repositioning reduces the incidence of compared with usual care.	Secondary outcomes:	Economic outcomes:
	<ul style="list-style-type: none"> ◀ Inpatient in a long-term geriatric facility ◀ Over 65 years of age ◀ At risk of pressure injury development using the activity and mobility components of Braden scale ◀ No pressure injury at time of recruitment to study ◀ No medical condition that would preclude the use of repositioning ◀ Consent 	<p>Group A: 30° tilt (n = 99 participants randomised, 99 analysed)</p> <p>Group B: Usual care (n = 114 participants randomised, 99 analysed)</p> <p>Group C:</p> <p>Co-interventions: participants in both groups nursed as per planned care regarding nutritional regimens, toileting, changing of incontinence pads, preparation for feeding, and pressure redistribution devices on chairs. Repositioned every 2- to 3-hours during the day.</p>	<ul style="list-style-type: none"> ◀ Mean daily nurse time for repositioning ◀ Nurse time cost per patient ◀ Cost of patient free of pressure injury ◀ Projected annual cost 	Time points: weekly follow-up over 4 weeks
	Exclusion criteria:			
	<ul style="list-style-type: none"> ◀ Patients with existing pressure injury 			
Defloor 2005	Location: Wards (n=32) across nursing homes (n=11) in Flanders, Belgium	Study design: 5-armed cluster-controlled trial with a 4-week (28-day) follow-up period	Primary outcome: incidence of a pressure injury (any stage) during a 28-day period.	High
		Aims: Investigate the effect of 4	Secondary outcomes: unrelated to	

Number of participants: n=838	different preventative regimens involving either frequent turning (2- to 3-hourly) or use of pressure-reducing mattress in combination with less frequent turning (4- to 6-hourly).	review outcomes.		
Eligibility criteria:	<ul style="list-style-type: none"> ◀ Geriatric residents with a Braden score of < 17 or a Norton score of < 12 ◀ Informed consent of the patient/family ◀ No pressure injury at time of recruitment to study 	Time points: twice weekly for 4 weeks (28 days).		
Exclusion criteria:	<ul style="list-style-type: none"> • none stated, but total of 1114 people excluded. 	<p>Alternating turning positions: semi-Fowler's with feet elevated 30° alternating with 30° lateral rotation, pillow placement under back from shoulder on standard mattress.</p> <p>Specified sitting position: intervention group sitting periods were recorded but not standardised; participants sat on thick air cushions. Backrest tilt on chair, legs on footrest, but heels not supported. Cushion for back.</p> <p>Group 2 control: n=576</p>		
Young 2004	Location: medical ward of an acute general	Study design: randomised controlled trial (groupings for allocation not	Primary outcome: incidence of non-blanchable erythema during a 24-hour	High

hospital in Wales.	reported) with a 24-hour follow-up period.	period.
Number of participants: n=46	Aims: Examine the effects of the 30° tilt in reducing non-blanchable erythema	Secondary outcomes: not stated Time points: 1, at 24 hours
Inclusion criteria: <ul style="list-style-type: none"> • Elderly patients • At risk of developing a pressure injury using Waterlow score • Able to lie in 30° tilt position • Given informed consent • No existing pressure injury • Caucasian 	Group A (intervention): Repositioning using 30° tilt (left side, back, right side, back) 2- to 3-hourly overnight, 2- to 3-hourly during the day. Sacrum and heels free from contact with support surface. Group B: 90° lateral and supine positions 2- to 3-hourly overnight, 2- to 3-hourly during the day. Standard care across all groups: support mattress: low air loss mattress or alternating air pressure mattresses.	
Exclusion criteria: not stated		

Abbreviations:

US,

United

States

3.2. Risk of bias assessment

Figure 3 summarises risk of bias across included studies. All trials were assessed as being at high risk of bias in at least one domain ¹⁷⁻²⁴. The risk of bias was unclear for at least one domain for all but three studies ^{17 20 24}.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Bergstrom 2013	+	+	-	+	?	+	?
Defloor 2005	+	?	-	-	-	+	+
Ghezeljeh 2017	+	+	-	-	+	+	+
Manzano 2014	+	+	-	+	+	+	+
Moore 2011	+	+	-	-	+	+	?
Pickham 2018	+	+	-	+	+	+	?
Young 2004	+	+	-	+	-	+	+
Zhou 2014	?	?	-	?	+	+	+

Figure 3: Risk of bias summary for the 8 included studies
(Colour coding: Green=low risk; Yellow=unclear risk; Red=high risk)

3.3. Frequency of repositioning

Data from four trials with 2870 participants compared 2-hourly versus 3-hourly positioning^{24 25 28 29} were included. Data from the two trials were unable to be pooled due to high statistical heterogeneity ($I^2 = 77%$)^{25 28}. There were no differences in the risk of pressure injury for 2-hourly versus 3-hourly frequencies in either study (risk ratio 4.06, 95% confidence interval 0.87 to 18.98 and risk ratio 0.90, 95% confidence interval 0.69 to 1.16, respectively), and the certainty of evidence is low²⁸ to very low²⁵.

Figure 4 shows a forest plot including three trials comparing 2-hourly with 4-hourly repositioning (1074 participants) (fixed-effect; $I^2 = 45%$, pooled risk ratio 1.06, 95% confidence interval 0.80 to 1.41)^{24 25 28}. It is uncertain whether 2-hourly repositioning compared with 4-hourly repositioning used in conjunction with any support surface increases or decreases the incidence of pressure injury, and the certainty of evidence is very low due to high risk of bias.




Figure 4: Pressure injury occurrence (stages 1-4), 2-hour versus 4-hourly positioning on any support surface

Bergstrom et al.²⁸ compared repositioning regimens using 3-hourly (n = 209), and 4-hourly (n = 198) frequencies, with all participants being nursed on high-density foam mattresses. There was no significant difference in pressure injury incidence between with 3-hourly compared with 4-hourly repositioning regimens (risk ratio 0.20, 95% confidence interval 0.04 to 0.92), and the certainty of evidence is low.

Defloor et al.²⁵ compared the number of new pressure injuries of any stage in participants being nursed on viscoelastic foam mattresses receiving 4-hourly repositioning compared with those receiving 6-hourly repositioning²⁵. There was a reported 27% reduction associated with 4-hourly repositioning (risk ratio 0.73, 95% confidence interval 0.53 to 1.02), however the certainty of evidence was very low due to high risk of bias. Finally, Pickham et al.²⁹ compared a 2-hourly turning regimen using a 20° tilt with "standard care", and participants in the intervention group developed fewer pressure injuries compared to the control group (risk ratio 0.28, 95% confidence interval 0.10 to 0.75), with moderate certainty of evidence²⁹.

3.4 Different positions for positioning

Data from four trials with 495 participants were included that compared different positions for positioning^{26 27 30 31}. Figure 5 shows the forest plot results of two trials comparing 30° and 90° tilts (pooled risk ratio 0.62, 95% confidence interval 0.10 to 3.97)^{26 27}. Overall, there was no clear difference in occurrence of stage 1 or 2 pressure injuries (persistent erythema) between the degree of tilting and the certainty of evidence is very low due to high risk of bias.




Figure 5: Pressure injury occurrence (stages 1-4), 30° tilt overnight versus 90° tilt overnight

There were two other trials undertaken in intensive care unit settings that included 236 participants, however their results could not be pooled due to the differences in tilt and repositioning regimens. Ghezalje et al.³¹ compared the use of 30° 2-hourly head-of-bed tilt with a 45° 2-hourly head of bed tilt and "standard care" in three randomised groups³¹. For the outcome of pressure injury, the authors stated that none of the patients who were recruited in this study developed pressure injuries. The certainty of evidence is low. Zhou et al.³⁰ compared the effects of prone positioning (intervention) versus supine positioning on the development of pressure injury. The occurrence of stage 1 pressure injury was higher in intervention participants, but there was no clear difference in the risk of stage 2 pressure injury between the two groups (P > 0.05). However, the authors presented no numerical data and the certainty of evidence is low due to high risk of bias.

3.5. Economic outcomes

A within-trial cost-minimisation analysis undertaken alongside Bergstrom et al. compared the costs of 3-hourly (n = 326) and 4-hourly (n = 295) repositioning to a 2-hourly (n = 321) repositioning schedule³³. In Canadian dollars, the cost of repositioning was estimated to be \$11.05 lower per resident per day for the 3- hourly regimen or Canadian dollars \$16.74 lower per resident per day for the 4- hourly regimen, compared to the 2-hourly regimen. The estimates of economic benefit were from the perspective of the Ontario Ministry of Health and Long-Term Care and were driven primarily on the value of freed nursing time. The analysis assumed 2-, 3-, or 4-hourly repositioning was associated with a similar incidence of pressure injury based on the trial findings²⁸.

Moore et al.³² performed a cost-effectiveness analysis based on data from their cluster-controlled trial²⁶, comparing the nursing time cost of 3-hourly repositioning using a 30° tilt with standard care (6-hourly repositioning with a 90° lateral rotation) among nursing home residents. The authors estimated 3-hourly repositioning using a 30° tilt to cost €46.50 (95% confidence interval €1.25 to €74.60) less per patient in nursing time compared with 6-hourly repositioning with a 90-degree lateral rotation. Consequently, based on the trial findings and subsequent economic evaluation, the authors concluded that the 30° tilt 3-hourly regimen was less costly in terms of nurse time (as well as more effective in reducing the incidence of pressure injuries in their trial) than the standard care of 6-hourly repositioning with a 90° lateral rotation (Moore 2011²⁶; 2013³²).

4. Discussion

The objective of this updated review was to assess the clinical and cost effectiveness of repositioning regimens on the prevention of pressure injury in adults regardless of risk in any setting. This review included data from eight clinical trials and two economic analyses with a total of 3,941 participants. The Grading of Recommendations, Assessment, Development and Evaluations assessment judged the certainty of evidence for effectiveness of repositioning frequencies, and positions as low or very low certainty, with serious risk of bias related to lack of blinding and imprecision. Therefore, it was not possible to draw reliable conclusions as to whether one particular repositioning frequency or position is more effective in pressure injury prevention.

In this review, one study reported the cost of nursing time for repositioning to be lower when patients were more frequently positioned with a lower tilt regimen (i.e., more frequent turning required less nursing time³²). The result of the other economic sub study³³ suggested that less frequent repositioning used in combination with viscoelastic mattresses was more cost-effective primarily in terms of nursing time. Nonetheless, with only two within trial economic evaluations included in this review, both in nursing home residents, these results are at best inconclusive and unlikely to be generalisable. Therefore, understanding which repositioning regimen provides optimal value is an important area for further research.

4.1. Limitations of included studies

Overall, limitations of included studies diminished the completeness and applicability of evidence. Trials were conducted in developed nations²⁴⁻²⁷, and five were undertaken in

acute care settings^{24 27 29-31}, limiting the extent to which findings can be generalised to lower-income and non-acute care settings. All trials had small samples and thus were underpowered to detect treatment effects, the focus of interventions varied, and no study examined outcomes such as participant pain, quality of life, or patient satisfaction. Inconsistent follow-up periods and variation in the types of support surfaces used across the included trials also limits generalisability.

Assessment of risk of bias identified limitations relative to blinding. Difficulty in blinding outcome assessors to the intervention was particularly concerning and a serious limitation in three trials^{25 26 31}. The certainty of the evidence was assessed as low or very low for most of the included trials. Studies were downgraded due to small samples with consequent imprecision, lack of allocation concealment and/or blinding of personnel and outcome assessors. These results are consistent with others' assessment of the evidence for frequencies of repositioning and repositioning positions³⁶⁻³⁸. Findings from a recently published systematic review by Avsar et al³⁶ concluded that the evidence for turning and repositioning schedules remains inconclusive³⁶. While Asar et al.³⁶ similarly examined the effects of different repositioning regimens, there are some differences between their review and the current review: Avsar et al.³⁶ included before-and-after design studies and descriptive studies as well as randomised controlled trials and cluster-controlled trials, and outcomes of patient preferences rather than health-related quality of life outcomes. The current review considered data derived from only randomised controlled trials and cluster-controlled trials, and included health-related quality of life, procedural pain, and patient satisfaction.

4.2. Strengths and limitations of this review

We conducted a rigorous and comprehensive systematic literature search that was reproducible. This review was guided by clearly defined, prespecified procedures to prevent potential bias in the review process and all evidence that could be obtained in the review was considered, including one study that was not published in English ³⁰. Nevertheless, we may have missed trials published in journals that were outside our search strategy. We had planned to undertake a sensitivity analysis to test the robustness of the results based on those lost to follow-up, but, due to the low volume and quality of the evidence and the inability to draw any conclusions, a sensitivity analysis would not have aided understanding. That these meta-analyses included only two to three studies is a limitation of this review. Further, it is possible this review is subject to cultural bias; six of the included trials were conducted in developed nations, thereby limiting the generalisability of findings to lower-income settings.

4.3. Implications for clinical practice

There is currently insufficient evidence to recommend one repositioning regimen in preference to another. Repositioning per se is recommended in all clinical practice guidelines ^{2 39 40}, though implementation is probably variable and highly dependent on the available resources. More recent clinical practice guidelines no longer advocate repositioning patients every two hours ². Rather, guidelines recommend that the patient's level of activity and ability to reposition themselves should guide health professionals' decision-making regarding frequency and amount of assistance for repositioning ². This seems like a very common-sense approach and reflects more patient-centred, individualised care.

4.4. Implications for research

To address the methodological limitations identified in included trials, researchers must ensure transparency of research process and adhere to the Consolidated Standards of Reporting Trials statement for reporting randomised controlled trials⁴¹. To minimise the sources of bias, researchers need to ensure rigorous processes in research design and execution of allocation concealment, randomisation, blinding, and participant attrition⁴². For instance, having assessors who are blinded to the outcome. If cluster-controlled trials are used, researchers need to also consider the potential for bias relative to participant selection, baseline comparability, analysis, and loss of clusters¹⁸.

Further research is needed relative to: repositioning frequencies and optimal positioning; use of manual repositioning regimens and electronic repositioning aids; effects of repositioning in high-risk patient populations (e.g. spinal cord injury); effects of position sensors on repositioning regimens; use of pressure sensor technologies to map pressure in relation to different tilt angles during repositioning; use of repositioning monitors to quantify patient repositioning while in bed; economic costs (including incremental costs) of pressure injuries; and economic and social impacts of pressure injuries on patients' health-related quality of life using valid and reliable measures. However, a properly conducted trial will be challenging to plan, fund and deliver and will likely require a repositioning intervention that is complex. Thus, it is unlikely to be as simple as a 2-hourly versus 4-hourly repositioning regimen because patients' clinical conditions need to be considered.

5. Conclusions

These review findings suggest a lack of robust evaluations of repositioning frequency and positioning for pressure injury prevention and uncertainty about their clinical and cost

effectiveness. Nonetheless, we cannot conclude that these interventions are ineffective as all comparisons are underpowered. This lack of quality trials reflects the challenges in undertaking research focused on repositioning for pressure injury prevention. The growing body of evidence on the aetiology of pressure injuries helps explain the mechanism of action of repositioning as one strategy that makes theoretical sense.

Acknowledgements: This is an abridged version of a review published in the Cochrane Database of Systematic Reviews (CDSR) 2020, Issue 6, DOI: 10.1002/14651858.CD009958.pub3 (see www.cochranelibrary.com for information).

Conflict of interest

The authors declare they have no conflict.

Funding sources

No external funding.

References

1. Gillespie BM, Chaboyer WP, McInnes E, et al. Repositioning for pressure ulcer prevention in adults. *Cochrane Database of Syst Rev* 2014(4)
2. European Pressure Ulcer Advisory Panel, National Pressure Injury Advisory Panel, Pan Pacific Pressure Injury Alliance. Prevention and Treatment of Pressure Ulcers/Injuries: Clinical Practice Guideline: The International Guideline. In: (Ed.) EH, ed. 3rd ed, 2019:1-405.
3. Institute for Healthcare Improvement. Getting Started Kit: Prevent Pressure Ulcers Supplement for Rural Hospitals, 2008.
4. Moore Z. Patient safety and pressure ulcers. *European Wound Management Association Journal* 2013;13(1):63-64.
5. McInnes E, Jammali Blasi A, Bell Syer SE, et al. Support surfaces for pressure ulcer prevention. *Cochrane Database of Syst Rev* 2015(9)
6. Li Z, Lin F, Thalib L, et al. Global prevalence and incidence of pressure injuries in hospitalised adult patients: A systematic review and meta-analysis. *Int J Nurs Stud* 2020;105:103546.
7. Hauck KD, Wang S, Vincent C, et al. Healthy life-years lost and excess bed-days due to 6 patient safety incidents: empirical evidence from english hospitals. *Medical care* 2017;55(2):125.
8. Nguyen K-H, Chaboyer W, Whitty JA. Pressure injury in Australian public hospitals: a cost-of-illness study. *Aust Health Rev* 2015;39(3):329-36.
9. Padula WV, Delarmente BA. The national cost of hospital acquired pressure injuries in the United States. *International wound journal* 2019;16(3):634-40.
10. Bennett G, Dealey C, Posnett J. The cost of pressure ulcers in the UK. *Age Ageing* 2004;33(3):230-35.
11. Catania K, Huang C, James P, et al. PUPPI: the pressure ulcer prevention protocol interventions. *Am J Nurs* 2007;107(4):44-52.
12. Braden B, Bergstrom N. A conceptual schema for the study of the etiology of pressure sores. *Rehabil Nurs* 1987;12(1):8-16.
13. Walker RM, Gillespie BM, McInnes E, et al. Prevention and treatment of pressure injuries: A meta-synthesis of Cochrane Reviews. *J Tissue Viability* 2020
14. Gillespie BM, Walker RM, Latimer SL, et al. Repositioning for pressure injury prevention in adults. *Cochrane Database of Syst Rev* 2020(6)
15. Gillespie BM, Chaboyer WP, McInnes E, et al. Repositioning for pressure ulcer prevention in adults. *Cochrane Database Syst Rev* 2014(4):44. doi: 10.1002/14651858.CD009958.pub2
16. Gillespie BM, Chaboyer W, McInnes E, et al. Repositioning for pressure ulcer prevention in adults. Cochrane Review [Protocol]. *The Cochrane Library* 2012(7)
17. Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Int J Surg* 2010;8(5):336-41.
18. Higgins JP, Deeks JJ, Altman DG. Chapter 16: Special topics in statistics. In: Higgins JP, S G, eds. *Cochrane handbook for systematic reviews of interventions: The Cochrane Collaboration* 2011.
19. Deeks JJ. Issues in the selection of a summary statistic for meta analysis of clinical trials with binary outcomes. *Stat Med* 2002;21(11):1575-600.
20. The Cochrane Collaboration Review Manager 5 (RevMan 5) [program]. 5.3 version. Copenhagen: Nordic Cochrane Centre, The Cochrane Collaboration, 2014.
21. Schünemann HJ, Oxman AD, Vist GE, et al. Chapter 12: Interpreting results and drawing conclusions. In: Higgins JP, Green S, eds. *Cochrane Handbook for Systematic Reviews of Interventions Version 510: The Cochrane Collaboration* 2011.
22. Ryan R, Hill S. How to GRADE the quality of the evidence. Cochrane Consumers and Communication Group. 2016, 2019.
23. Schünemann HJ, Higgins JP, Vist G, et al. Chapter 14: Completing 'Summary of findings' tables and grading the certainty of the evidence. In: Higgins JP, Thomas J, Chandler J, et al., eds. *Cochrane Handbook for Systematic Reviews of Interventions version 61* 2020.

24. Manzano F, Colmenero M, Pérez-Pérez AM, et al. Comparison of two repositioning schedules for the prevention of pressure ulcers in patients on mechanical ventilation with alternating pressure air mattresses. *Intensive Care Med* 2014;40(11):1679-87.
25. Defloor T, De Bacquer D, Grypdonck MH. The effect of various combinations of turning and pressure reducing devices on the incidence of pressure ulcers. *Int J Nurs Stud* 2005;42(1):37-46.
26. Moore Z, Cowman S, Conroy RM. A randomised controlled clinical trial of repositioning, using the 30 tilt, for the prevention of pressure ulcers. *J Clin Nurs* 2011;20(17 18):2633-44.
27. Young T. The 30 tilt position vs the 90 lateral and supine positions in reducing the incidence of non-blanching erythema in a hospital inpatient population: a randomised controlled trial. *J Tissue Viability* 2004;14(3):88-96.
28. Bergstrom N, Horn SD, Rapp MP, et al. Turning for Ulcer Reduction: a multisite randomized clinical trial in nursing homes. *J Am Geriatr Soc* 2013;61(10):1705-13.
29. Pickham D, Berte N, Pihulic M, et al. Effect of a wearable patient sensor on care delivery for preventing pressure injuries in acutely ill adults: A pragmatic randomized clinical trial (LS-HAPI study). *Int J Nurs Stud* 2018;80:12-19.
30. Zhou X, Liu D, Long Y, et al. The effects of prone position ventilation combined with recruitment maneuvers on outcomes in patients with severe acute respiratory distress syndrome. *Zhonghua nei ke za zhi* 2014;53(6):437-41.
31. Ghezjeljeh TN, Kalhor L, Moghadam OM, et al. The Comparison of the Effect of the Head of Bed Elevation to 30 and 45 Degrees on the Incidence of Ventilator Associated Pneumonia and the Risk for Pressure Ulcers: A Controlled Randomized Clinical Trial. *Iran Red Crescent Med J* 2017;19(7)
32. Moore Z, Cowman S, Posnett J. An economic analysis of repositioning for the prevention of pressure ulcers. *J Clin Nurs* 2013;22(15-16):2354-60.
33. Paulden M, Bergstrom N, Horn SD, et al. Turning for ulcer reduction (TURN) study: an economic analysis. *Ont Health Technol Assess Ser* 2014;14(12):1.
34. Mulligan S, Prentice J, Scott L. WoundsWest wound prevalence survey 2011 state-wide overview report. *Perth: Western Australia: Ambulatory Care Services, Department of Health* 2011
35. Queensland Health. 2016 Queensland bedside audit: statewide inpatient report. Brisbane, Qld Australia, 2017:1-163.
36. Avsar P, Moore Z, Patton D, et al. Repositioning for preventing pressure ulcers: a systematic review and meta-analysis. *Journal of Wound Care* 2020;29(9):496-508.
37. Jocelyn Chew HS, Thiara E, Lopez V, et al. Turning frequency in adult bedridden patients to prevent hospital acquired pressure ulcer: A scoping review. *International Wound Journal* 2018;15(2):225-36.
38. Reddy M, Gill SS, Rochon PA. Preventing pressure ulcers: a systematic review. *Jama* 2006;296(8):974-84.
39. Beeckman D, Matheï C, Van Lancker A, et al. A national guideline for the treatment of pressure ulcers. *KCE Reports* 2013;203
40. Registered Nurses' Association of Ontario. Risk assessment and prevention of pressure ulcers: Guideline supplement. . Toronto, Ontario: Registered Nurses' Association of Ontario,, 2011.
41. Schulz KF, Altman DG, Moher D, et al. CONSORT 2010 statement: updated guidelines for reporting parallel group randomised trials. *Trials* 2010;11(1):32.
42. Polit DF, Gillespie BM. Intention to treat in randomized controlled trials: Recommendations for a total trial strategy. *Research in nursing & health* 2010;33(4):355-68.