

Article type : Original Article

# Allelic variants of the NLR protein Rpi-chc1 differentially recognise members of the *Phytophthora infestans* PexRD12/31 effector superfamily through the leucine-rich repeat domain

Daniel Monino-Lopez <sup>1</sup>, Maarten Nijenhuis <sup>1,2</sup>, Linda Kodde <sup>1</sup>, Sophien Kamoun <sup>3</sup>, Hamed Salehian <sup>1</sup>, Kyrilo Schentsnyi <sup>1,4</sup>, Remco Stam <sup>1,5</sup>, Anoma Lokossou <sup>1</sup>, Ahmed Abd-El-Haliem <sup>1,6</sup>, Richard GF Visser <sup>1</sup>, Jack H Vossen <sup>1\*</sup>

<sup>1</sup> Plant Breeding, Wageningen University & Research, Droevendaalsesteeg 1, 6708PB, Wageningen, The Netherlands.

<sup>2</sup> Current address: Agrico Research, Burchtweg 17, 8314PP, Bant, The Netherlands;

<sup>3</sup> The Sainsbury Laboratory, University of East Anglia, Norwich Research Park, Norwich, United Kingdom.

<sup>4</sup> Current address: Center for Plant Molecular Biology, Auf der Morgenstelle 32, 2076 Tübingen, Germany;

<sup>5</sup> Current address: Technical University Munich;

<sup>6</sup> Current address: Rijk Zwaan Breeding B.V., Burgemeester Crezéelaan 40, The Netherlands.

**Running title:** Rpi-chc1 variants recognize PexRD12/31 effectors

**Key words:** NLR cluster, Leucine rich repeat, *Phytophthora infestans*, Late blight resistance gene, RXLR effector, Solanum species, potato

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/tbj.15284](https://doi.org/10.1111/tbj.15284)

This article is protected by copyright. All rights reserved

\* Corresponding author:

Jack H. Vossen

+31 317 481 590

*jack.vossen@wur.nl*

## Summary

*Phytophthora infestans* is a pathogenic oomycete that causes the infamous potato late blight disease. Resistance (*R*) genes from diverse *Solanum* species encode intracellular receptors that trigger effective defence responses upon the recognition of cognate RXLR avirulence (*Avr*) effector proteins. To deploy these *R* genes in a durable fashion in agriculture, we need to understand the mechanism of effector recognition and the way the pathogen evades recognition. In this study, we cloned sixteen allelic variants of the *Rpi-chc1* gene from *Solanum chacoense* and other *Solanum* species, and identified the cognate *P. infestans* RXLR effectors. These tools were used to study effector recognition and co-evolution. Functional and non-functional alleles of *Rpi-chc1* encode Coiled-Coil Nucleotide Binding Leucine-Rich-Repeat (CNL) proteins, being the first described representatives of the CNL16 family. These alleles have distinct patterns of RXLR effector recognition. While Rpi-chc1.1 recognised multiple PexRD12 (*Avrchc1.1*) proteins, Rpi-chc1.2 recognised multiple PexRD31 (*Avrchc1.2*) proteins, both belonging to the PexRD12/31 effector superfamily. Domain swaps between Rpi-chc1.1 and Rpi-chc1.2 revealed that overlapping subdomains in the LRR were responsible for the difference in effector recognition. This study showed that Rpi-chc1.1 and Rpi-chc1.2, evolved to recognize distinct members of the same PexRD12/31 effector family via the LRR domain. The biased distribution of polymorphisms suggests that exchange of LRRs during host-pathogen co-evolution can lead to novel recognition specificities. These insights will help future strategies to breed for durable resistant varieties.

## Introduction

Potato (*Solanum tuberosum*) is the third largest food crop in terms of human consumption in the world after rice and wheat, with more than 368 million tonnes produced in 2018 (FAO, 2020; Devaux *et al.*, 2020). Potato late blight, caused by the oomycete *Phytophthora infestans* (*P. infestans*), is one of the most infamous potato diseases. During the mid-1840s, this pathogen caused the Great Irish Famine from which around one million people died (Callaway, 2013). Nowadays, losses from late blight are estimated to still reach 16% of the world production and the main disease management is based on biocide applications. Including yield losses and crop protection measures, late blight causes a global economic loss of € 5.2 billion per year (Haverkort *et al.*, 2016).

*P. infestans* is an oomycete with sexual and asexual life cycles, which exhibits a hemibiotrophic lifestyle on potato. Together with its large and fast evolving genome (estimated to be 240 Mb), its population diversity leads to the regular emergence of new aggressive and virulent strains (Haas *et al.*, 2009). The infection starts when a spore lands on the plant surface, germinates and forms a penetration structure called appressorium. Alternatively, spores can also enter through natural openings such as stomata. After passing the epidermis, hyphae spread intercellularly projecting haustorium structures into the mesophyll cells. These haustoria are specialised infection structures that secrete both apoplastic and cytoplasmic effectors to create an intimate association with the host cell and facilitate nutrient uptake (Fry, 2008). Effectors are pathogen molecules that interact with different host targets to suppress the host defence response and enable colonisation. The publication of the *P. infestans* T30-4 genome, revealed the presence of 563 effector genes encoding the conserved Arg–any amino acid–Leu–Arg (RXLR) peptide motif (Haas *et al.*, 2009). These effectors, rapidly evolve by gaining and losing repeat-rich domains through recombination with different paralogs, transposon movement, and point mutations (Goss, Press and Grünwald, 2013). During co-evolution, potato has evolved receptors to recognise some of these effectors and trigger an immune response.

Wild *Solanum* species are the main source of resistance (*R*) genes to *P. infestans* (*Rpi*). To date, over 20 *Rpi* genes have been characterised in different *Solanum* species, e.g. *R1*, *R2*, *R3a*, *R3b*, *R8*, *R9a* from *S. demissum*, *Rpi-blb1*, 2 from *S. bulbocastanum*, *Rpi-vnt1* from *S. venturii* and *Rpi-amr1* from *S. americanum* (Ballvora *et al.*, 2002; van der Vossen *et al.*, 2003, 2005; Huang *et al.*, 2005; Lokossou *et al.*, 2009; Pel *et al.*, 2009; Li *et al.*, 2011; Jo *et al.*, 2015; Vossen *et al.*, 2016; Witek *et al.*, 2021). All these receptors belong to the nucleotide-binding (NB)–leucine-rich repeat (NLR) type of receptors and contain a coiled-coil domain (CC) in their N-termini, referred to as CC-NB-LRR or CNL. The recognition of a specific effector or avirulence factor (Avr) leads to the activation of the effector triggered immunity (ETI) and the restriction of the pathogen growth (Jones, Vance and Dangl, 2016). ETI is mostly monogenic and therefore well suited and commonly deployed for resistance breeding and crop protection strategies. To keep up with the fast evolution of effectors, NLR genes are also very diverse and rapidly evolving. Gene duplications, recombinations, unequal crossing overs and transpositions have been proposed to provide the basis for the evolution of the NLR recognition spectrum (Leister, 2004; McDowell and Simon, 2006). This fast evolution can lead to the independent development of new receptors in different geographical locations that recognise the same effector. For instance, the recognition of the effector Avr2 from *P. infestans*, by the unrelated *R2* and *Rpi-mcq1* CNLs (Aguilera-Galvez *et al.*, 2018). *R2* is located on chromosome IV in the Mexican species *S. demissum*, while *Rpi-mcq1* is located on chromosome IX from a Peruvian accession of *S. mochiquense* (Smilde *et al.*, 2005; Foster *et al.*, 2009). More recently, *Rpi-amr1* alleles have been described to cause differential recognition of *Avramr1* homologs from several *P. infestans* isolates (Witek *et al.*, 2021). When the doubled-monoploid DM1-3 519 R44 potato genome was published, 755 NLR genes were identified (Jupe *et al.*, 2013). Many of them were found in clusters together with closely related paralogs. All of these clusters were formed in ancestral species and had sequence homology to syntenic genomic regions from other *Solanum* species harbouring late blight resistance genes. Thus, alleles from functional *Rpi* genes that do not provide resistance (*rpi*) can be found in all studied *Solanum* genomes.

Here, we studied *Solanum chacoense* (*S. chacoense*); a diploid wild potato relative from South America considered a source of resistance to *P. infestans* (Vossen *et al.*, 2009;

Karki, Jansky and Halterman, 2021). We identified two functionally distinct receptors, Rpi-chc1.1 and Rpi-chc1.2, which are allelic variants that recognise distinct *P. infestans* effectors from the same PexRD12/31 effector superfamily. Remarkably, only Rpi-chc1.1 is able to provide resistance against several *P. infestans* isolates. The expression and recognition of PexRD12 effectors was associated with Rpi-chc1.1 mediated resistance and, therefore, designated as Avrhc1.1 effectors. PexRD31 effectors were still expressed in several *P. infestans* isolates, but were rapidly downregulated during the interaction with potato. This, potentially explains the inability of Rpi-chc1.2 to provide late blight resistance. We postulate that Rpi-chc1.2 is a ubiquitous ancient *R* gene that was recently overcome and PexRD31 may have functioned as Avrhc1.2. An allele mining strategy revealed Rpi-chc1 orthologs in different wild *Solanum* accessions and potato cultivars that could be classified by their sequence and recognition spectrum of Avrhc1.1, Avrhc1.2, or non-functionality. Finally, using domain swaps, we found that the LRR domain harboured the recognition specificity of both Avrhc1.1 and Avrhc1.2. The specificities resided in overlapping LRR subdomains and could not be combined into one active protein using domain exchanges.

## Results

### Cloning and characterization of Rpi-chc1.1

The *S. chacoense* accession CHC543 from Bolivia is a previously described wild potato relative harbouring resistance to *P. infestans* (Vleeshouwers *et al.*, 2011a). To identify the genetic locus of resistance, the resistant seedling CHC543-5 was crossed with the susceptible seedling CHC544-5 to generate the F1 population 7650, consisting initially of 212 individuals. This population was challenged with *P. infestans* isolate 90128 in a detached leaf assay (DLA). A clear 1:1 segregation was observed, indicating the presence of a single dominant resistance gene which will henceforth be referred to as Rpi-chc1. CAPS markers from chromosome 10 were tested as this chromosome was known to harbour Rpi-ber from the related species *S. berthaultii* (Vossen *et al.*, 2013). The marker TG63 in chromosome 10 was indeed linked to the Rpi-chc1 resistance. Successive fine mapping in a recombinant population representing 2357 individuals was

performed using markers derived from RH89-39-16 BAC clones from chromosome 10 (PGSC) (The Potato Genome Sequencing Consortium, 2011; Sharma *et al.*, 2013). A narrow genetic window between markers RH106G03-T and RH97D21\_C21-4 was identified to contain *Rpi-chc1* (**Figure 1a**). To generate a physical map of the mapping interval, two Bacterial Artificial Chromosome (BAC) clones, B1 and B2, were selected from a BAC library that was derived from CHC543-5 genomic DNA. After sequencing the BAC clones, two NLR genes were identified in clone B1 and another six NLR in clone B2. Further fine-mapping revealed that only the last six were located within the mapping interval and only three (B2-1, B2-2 and B2-3) encoded complete NLR proteins (**Figure 1b**). The three candidates were subcloned including their native 5' and 3' regulatory elements, and complementation analyses were performed in *N. benthamiana*. After two days, the agroinfiltrated area was challenged with *P. infestans* 90128. *Rpi-blb1*, which was shown to provide resistance to *P. infestans*, was used as a positive control. The leaves agroinfiltrated with candidate B2-3 and *Rpi-blb1* showed severely compromised pathogen growth, while leaves with candidates B2-1 and B2-2 were completely susceptible to *P. infestans* 90128 (**Figure 1c**). This result suggested that B2-3 was the gene in CHC543-5 that provides resistance to *P. infestans*. To verify this result, the three candidates B2-1, B2-2 and B2-3 were stably transformed into the susceptible *S. tuberosum* cv. 'Desiree'. Indeed, only the events containing candidate B2-3 showed resistance to *P. infestans* (**Figure 1d**). Furthermore, three different single guide RNAs (sgRNAs) were designed to specifically target the LRR domain of candidate B2-3 (**Figure 2a**). The resistant CHC543-5 genotype was stably transformed with CRISPR-Cas9 and these sgRNAs. The transformation events were challenged with *P. infestans* 90128 and IPO-C isolates, and 48% of the transformants had become susceptible to both isolates (**Table S1; Figure 2b**), which suggested that the active late blight resistance gene was specifically and successfully mutated. Four transformants were genotyped and frameshift mutations were found only in the three susceptible transformants (**Figure 2c**). Therefore, we concluded that B2-3 was the gene from CHC543-5 that was causal for late blight resistance. Henceforth, we will refer to gene B2-3 as *Rpi-chc1.1* as it is the first *Rpi-chc1* allele that is identified in *S. chacoense*.

*Rpi-chc1.1* is a CC-NB-LRR or CNL and it belongs to the CNL-16 (Witek *et al.*, 2016) immune receptor family (**Figure 3a**). *Rpi-chc1.1* has one uninterrupted open reading frame of 3909 bp, which is predicted to be translated into 1303 amino acids (**Figure S2**). No introns were predicted. The CC domain contains the N terminal MADA motif, 4 predicted  $\alpha$ -helices and the typical hhGRExE, but the distinctive EDVID motif was less conserved (Rairdan *et al.*, 2008). The NB domain contains the characteristic Kinase 1a VYND, Kinase 2, and Kinase 3a motifs (Saraste, Sibbald and Wittinghofer, 1990; Campbell, 2003; Leipe, Koonin and Aravind, 2004; Pal, Chakrabarti and Basak, 2007; Wendler *et al.*, 2012). The ARC1 domain contains the RNBS-C, the Motif 3 and the GLPL motif. The ARC2 domain contains the Motif 2, the RNBS-D and two copies of the MHDL motif (Danot *et al.*, 2009; Reubold, Wohlgemuth and Eschenburg, 2011; Sukarta, Sloodweg and Goverse, 2016). The LRR domain consists of 29 imperfect repeats. Both LRR3 and LRR4 contain a central VLDL motif which is conserved in the third LRR of most functional NLRs (Warren *et al.*, 1998; Bendahmane *et al.*, 2002).

### Identification of *Rpi-chc1.1* allelic variants

In order to identify different *Rpi-chc1.1* allelic variants, we pursued an allele mining approach in several closely related resistant and susceptible *S. chacoense*, *S. berthaultii*, *S. tarijense*, and a *S. tuberosum* accessions. Homologous sequences were amplified using primers overlapping the start and stop codons of *Rpi-chc1.1*. The PCR fragments of the expected 3.9 kb size were cloned and sequenced, resulting in the identification of fifteen *Rpi-chc1.1*-like sequences. The mined *Rpi-chc1.1* variants contained between 1296 to 1303 amino acids (**Figure 3b**; **Figure S2**). From the selected diploid accessions one or two sequence variants were identified, suggesting that indeed *Rpi-chc1* alleles were mined rather than paralogs. In case only one variant was mined from an accession, this suggested that the second allele has significant sequence polymorphisms at (one of) the primer annealing sites. Phylogenetic analysis of the sequences showed strong sequence similarities among the alleles (94.6 – 100% identity). Even within this high identity range, the presence of four main clades was revealed (**Figure 3b**). In clade 1, the *Rpi-chc1.1* allele was found, together with three sequences from *S. berthaultii* that were nearly identical to each other, and a sequence from *S. tarijense*. From clade 1, together

with *Rpi-chc1.1*, we selected one sequence from *S. berthaultii* (94-2031) and the *S. tarijense* (TAR852-5) for complementation analysis. Transformation of the corresponding genes to susceptible Desiree plants showed that they provide resistance to *P. infestans* isolates 90128 and IPO-C, like *Rpi-chc1.1* (**Table S2**). We therefore concluded that clade 1 contains functional homologs of *Rpi-chc1.1*. The *S. tarijense* allele will be referred to as *Rpi-tar1.1*. The *S. berthaultii* allele will be referred to as *Rpi-ber1.1* which matches to the previously described *Rpi-ber* and *Rpi-ber1* genes that were derived from the same accession (PI473331) at similar genetic positions (Rauscher *et al.*, 2006; Tan *et al.*, 2010; Vossen *et al.*, 2013).

The allele mining in accession CHC543-5 resulted not only in the re-identification of the active *Rpi-chc1.1* but also in the identification of a presumed allelic variant, which we will refer to as *Rpi-chc1.2*. To test if *Rpi-chc1.1* and *Rpi-chc1.2* were indeed alleles of the same gene, we tested *Rpi-chc1.2* specific markers in the recombinant population 6750 (CHC543-5 x CHC544-5). We found a perfect repulsion between *Rpi-chc1.2* and *Rpi-chc1.1*, strongly suggesting that both genes are allelic variants (**Table S3**). Additionally, this analysis proved that *Rpi-chc1.2* does not cause resistance against *P. infestans* 90128, even though *Rpi-chc1.2* is expressed during infection (**Table S2**; **Figure S1f**). The *Rpi-chc1.2* protein sequence clusters in clade 2 together with four identical sequences from *S. berthaultii*. Close to clade 2, we can observe clade 3, which consisted of a *S. berthaultii*, a *S. tarijense* and a *S. tuberosum* allele from RH89-039-16, a diploid clone previously characterised as susceptible to *P. infestans* (Vleeshouwers *et al.*, 2011a). The clade 3 allele from *S. tarijense* contained an in frame stop codon, making it unlikely that this allele is producing an active resistance protein. Additionally, a fourth clade contained only *S. berthaultii* alleles. The allelic variants were numbered according to the clade in which they were found (i.e. *Rpi-ber1.1* from clade 1 and *Rpi-ber1.2* from clade 2, etc.) followed by an extension to indicate the genotype from which the allele was derived.

## ***Rpi-chc1.1* recognises the RXLR PexRD12 effector family from *P. infestans***

To understand the resistance mechanism of the *S. chacoense* CHC543-5 accession, we searched for the effector recognised by *Rpi-chc1.1*. A collection of ninety *P. infestans* extracellular (Pex) proteins in a PVX agroinfectious vector, of which 54 contained the RXLR-DEER motif (PexRD), was screened. Individual clones from the Pex collection were co-agroinfiltrated with *Rpi-chc1.1* in *N. benthamiana* leaves. As a positive control, we used a mix of the *R3a/Avr3a* *R* gene effector pair which is known to trigger a strong hypersensitive response (HR) in *N. benthamiana* leaves. Only two effectors from the Pex collection were able to trigger an *Rpi-chc1.1* dependent HR, PexRD12-1 and PexRD12-2 (PITG\_16233 and PITG\_16240, respectively) (**Figure 4a**). Neither the inactive paralogs B2-1 and B2-2 nor *R3a* produced an HR upon co-agroinfiltration with PexRD12. These results showed that PexRD12 is specifically recognised by *Rpi-chc1.1*. We could further confirm this finding using transgenic Desiree potato plants that were transformed with *Rpi-chc1.1*. About half of this transgenic population showed late blight resistance while the other half was susceptible, probably due to impaired transgene expression. Interestingly, the plants that showed late blight resistance also showed PexRD12 recognition, while the susceptible transgenic plants did not show any response upon PexRD12 agroinfiltration (**Table S4**). We sought for further evidence that PexRD12 was indeed causing avirulence on *Rpi-chc1.1* expressing plants. In a field trial with natural infection, we found isolates that were virulent on plants containing *Rpi-chc1.1*. The infected material was collected and used for gene expression analysis via RT-qPCR. The expression of PexRD12 was undetected in the *Rpi-chc1.1* resistance breaking isolates, while other effectors such as *Avrsto1* were normally expressed (Vleeshouwers *et al.*, 2008). Reciprocally, we found that *Rpi-sto1* breaking isolates still expressed PexRD12 normally (**Figure 4b**). Altogether, these results suggest that PexRD12 corresponds to *Avrchc1.1*.

### **The PexRD12/31 superfamily is a complex *P. infestans* RXLR effector family**

Using Blast analyses of the T30-4 proteome, we found 9 homologs of PexRD12 in the *P. infestans* T30-4 genome. Additionally, we found that PexRD12 proteins had strong

homology with 9 members of the PexRD31 family and two additional, more distantly related sequences (**Table S5**). These 20 effectors are encoded by clusters of paralogs mainly in three supercontigs (**Figure S3**) and will henceforth be referred to as the PexRD12/31 superfamily (see also Petre *et al.*, 2020). All PexRD12/31 effectors are small proteins that include a signal peptide in the N-terminus, an effector domain in the C-terminus, and the conserved RXLR and EER motifs in the centre; except for PITG\_16243 and PITG\_09577 which contain an RXXR-EER and RXXLR-EER motifs, respectively (**Figure 5a**).

The alignment of the protein sequences and the phylogenetic analysis of the PexRD12/31 superfamily members resulted in five main clades (**Figure 5a**). Two highly homologous clades can be distinguished to form the PexRD12 family, PexRD12-A1 and PexRD12-A2. The clade PexRD12-A2 also includes truncated versions which partly or completely miss the effector domain. In addition, two related clades constitute the PexRD31 family, PexRD31-B and PexRD31-C. Additionally, PITG\_16428 and PITG\_09577 were much less related and are together referred to as PexRD12/31 group D.

To determine the degree to which PexRD12/31 members are expressed *in planta*, as observed for other Avr effectors of *P. infestans* (Vleeshouwers *et al.*, 2011b; Rietman *et al.*, 2012), we tested their expression during infection with quantitative PCR on cDNA using clade A, B and C specific primers. The relative expression was calculated and normalised for the relative amount of *P. infestans*. Three different *P. infestans* isolates were evaluated at different time points after inoculation of different susceptible potato genotypes (**Figure S1a-d**). In all the tested genotypes, PexRD12 showed the highest relative expression. In two isolates a maximum expression was found from 4 to 24 hours after inoculation and expression remained high till after 48 hours in all four isolates. The PexRD31-B effectors were expressed in 2 isolates but were rapidly downregulated in the first hours after inoculation with hardly any expression left. The expression of PexRD31-C was mostly undetectable along the inoculation time course. Similar results were observed when the PexRD12/31 expression was analysed during the infection of the four different

*P. infestans* isolates EU\_13\_A2, Ec1, EU\_6\_A1 and US23 (**Figure S4**), using the data from the PenSeq dataset (Lin *et al.*, 2020).

### **Rpi-chc1.2 recognises the RXLR PexRD31 effector family from *P. infestans***

In order to describe the spectrum of effector recognition by different *Rpi-chc1* alleles, several representatives from each clade were selected and co-agroinfiltrated with different PexRD12/31 members in *N. benthamiana*. *Rpi-chc1.1\_543-5* and *Rpi-ber1.1\_94-2031-01* from clade 1, *Rpi-chc1.2\_543-5* and *Rpi-ber1.2\_493-7* from clade 2, *rpi-tub1-RH89-039-16* from clade 3 and *rpi-ber1.4\_561-2* from clade 4 were selected. As a representation from each of the clades of the PexRD12/31 effector superfamily, we selected: PITG\_16245 (PexRD12-A1), PITG\_20934 (PexRD12-A2), PITG\_16235 (PexRD31-B), and PITG\_23069 (PexRD31-C). The different *Rpi-chc1* alleles were co-agroinfiltrated with the PexRD12/31 effectors in *N. benthamiana* leaves. Three days after agroinfiltration, we observed that the members from clade 1, *Rpi-chc1.1* and *Rpi-ber1.1*, specifically recognised both PexRD12-A1 and PexRD12-A2 effectors (**Figure 5b**). This result showed that *Rpi-chc1.1* and *Rpi-ber1.1* recognise multiple members of the PexRD12 family, suggesting that *Avrchc1.1* is encoded by multiple redundant paralogs. On the other hand, *Rpi-chc1.2* and *Rpi-ber1.2* from clade 2, specifically recognised both PexRD31-B and PexRD31-C effectors (**Figure 5b**). This suggests that multiple PexRD31 paralogs correspond to *Avrchc1.2*. The selected alleles from clade 3 and 4, *rpi-tub1.3\_RH89-039-16* and *rpi-ber1.4\_561-2*, were not able to recognize any of the PexRD12/31 members (**Figure 5b**), showing that clade 3 and 4 encode functionally more distant receptors, and is in agreement with the known susceptibility of RH89-039-16 to *P. infestans* (Vleeshouwers *et al.*, 2011a).

### **The LRR domain of the *Rpi-chc1* variants determines the PexRD12/31 effector recognition specificity**

Since the allelic variants of *Rpi-chc1* could be divided into three activity groups (recognition of either Avrhc1.1, Avrhc1.2 or none), while having an amino acid identity up to 96%, they provide ideal tools to investigate the Rpi-chc1 mechanism of recognition. Therefore, we performed progressive exchanges of the different receptor domains. The chimeric receptors were co-agroinfiltrated with the PexRD12/31 effectors in *N. benthamiana* leaves to evaluate their recognition specificity. First, we selected Rpi-chc1.2 and rpi-tub1.3\_RH89-039-16 as representatives of clade B and clade C, respectively. When aligning the protein sequences, 54 single amino acid polymorphisms (SAPs) were found and most of them were located in the LRR domain (**Figure 6a**). As previously mentioned, Rpi-chc1.2 specifically recognises PexRD31-B and PexRD31-C, while rpi-tub1.3\_RH89-039-16 does not recognise any of the PexRD12/31 effectors. When the complete rpi-tub1.3\_RH89-039-16 LRR domain was exchanged for the Rpi-chc1.2 LRR, the chimeric receptor RH::C2\_2-29 was able to recognise both PexRD31-B and PexRD31-C. Reciprocally, the exchange of the Rpi-chc1.2 LRR for the rpi-tub1.3\_RH89-039-16 in C2::RH\_2-29, led to the inability to recognise any of the PexRD12/31 effectors. This result demonstrates the importance of the LRR domain during the Avrhc1.2 recognition. Additional domain exchanges were performed in order to identify the essential LRR repeats for the effector recognition. The required LRR repeats for the Avrhc1.2 recognition could be narrowed down with the construct RH::C2\_14-19 to nine amino acid polymorphisms (**Figure 6a**). Due to the absence of polymorphisms in the LRR repeats 14 and 15, we can conclude that the exchange of the amino acid polymorphisms present in LRRs 16 to 19 activate rpi-tub1.3\_RH89-039-16 to recognise Avrhc1.2. Interestingly, the majority of these nine amino acid polymorphisms are particularly situated in the solvent exposed domain (xxLxLxxxx) of every LRR repeat. The exchange of the tested solvent exposed residues led to the partial or complete loss of effector recognition, suggesting the contribution of an effector binding surface (**Figure 6b**).

To understand the difference in effector recognition specificity between Rpi-chc1.1 and Rpi-chc1.2, and to explore the possibility to combine both recognitions in one receptor, we performed a similar progressive domain exchange approach between Rpi-chc1.1 and Rpi-chc1.2 (**Figure 7**). The exchange of the LRR domain in the chimeric receptors C1::C2\_8-29 and C2::C1\_8-29 led to a shift in effector recognition, from Avrhc1.1 to

Avrchc1.2. Further exchanges revealed that the LRR repeats 14 to 23 from Rpi-chc1.2 led to an opposite effector recognition pattern as the chimeric receptor C1::C2\_14-23 was able to only recognise Avrhc1.2. With reciprocal domain exchanges of Rpi-chc1.1 into Rpi-chc1.2, we found that LRR repeats 8 to 29, led to Avrhc1.1 recognition. In an attempt to further reduce the length of the exchanged sequence, the recognition of Avrhc1.1 resulted in partial (C2::C1\_8-25 and C2::C1\_8-23) or complete (C2::C1\_14-25 and C2::C1\_14-23) loss of recognition. Especially, when comparing the receptors C2::C1\_8-29 and C2::C1\_8-25, already the modification of the last five SAPs led to the reduced recognition of Avrhc1.1. But, apparently not only the last LRR repeats are involved in the effector recognition. Also the first LRR repeats, from 8 to 14, are also important for Avrhc1.1 recognition as C2::C1\_8-25 was able to partially recognise Avrhc1.1, while C2::C1\_14-25 did not trigger any HR. We conclude that the LRR repeats 8 to 29 in Rpi-chc1.1 are important for the Avrhc1.1 recognition, which overlaps with LRR repeats 16-19 from Rpi-chc1.2 which were required for Avrhc1.2 recognition.

## Discussion

In this study, we identified *Rpi-chc1.1* and 15 additional allelic variants from *S. chacoense*, *S. berthaultii*, *S. tarijense* and *S. tuberosum*. Phylogenetic analysis of the encoded protein sequences revealed four clades. These four clades were not only supported by sequence similarity but also by differences in effector and *P. infestans* recognition. Clade 1 genes encode receptors that recognise PexRD12 effectors and includes the active orthologs *Rpi-chc1.1*, *Rpi-ber1.1* and *Rpi-tar1.1* (**Figure 3 and 5**). Clade 2 receptors could be distinguished by the recognition of the PexRD31 effectors (**Figure 5**). Receptors encoded by clades 3 and 4 do not recognise PexRD12/31 effectors and no other activity has been found. Interestingly, clade 3 alleles are also present in domesticated potato clones that are susceptible to late blight; e.g. RH89-039-16 (**Figure 3**) and the varieties Colomba and Altus (to be published elsewhere), implying that the encoded receptors are not able to effectively provide resistance against *P. infestans*.

*Rpi-ber1.1\_94-2031-01* was derived from the same accession as the previously described *Rpi-ber* (Rauscher *et al.*, 2006; Vossen *et al.*, 2009; Tan *et al.*, 2010), and *Rpi-ber1* genes (Park *et al.*, 2009). In these four studies, *Rpi-ber* and *Rpi-ber1* mapped close to marker TG63 but slightly different genetic positions were reported. The population from Park *et al.* was quite small and a single recombination event may have caused the deviating genetic distance. In the case of Tan *et al.*, a single mis-phenotyping could explain the mapping of *Rpi-ber* distal to TG63. We therefore assume that *Rpi-ber* and *Rpi-ber1* are the same genes and adopt the *Rpi-ber1* naming as it is more consistent with current nomenclature for late blight resistance genes. *Rpi-ber2*, as described by Park *et al.*, was derived from the same accession that was used in our allele mining studies (BER493). We could not find a clade 1 *Rpi-chc1* allele from the BER493 accessions, which supports the idea that a more distantly related CNL16 member may be present that lacks sufficient match to the primer sequences, explaining the *Rpi-ber2* map position distal from TG63.

The presence of *Rpi-chc1* alleles in *S. tarijense* and *S. berthaultii* suggests a functional common ancestor existed before their speciation. However, it must be noted that the geographic locations where the accessions were found are close to each other in Bolivia. Since *S. chacoense*, *S. tarijense* and *S. berthaultii* are closely related, the presence of functional *Rpi-chc1* alleles in the three species might be a result from a recent species intercrossing.

Sequence similarity among the studied allelic variants correlated with their functionality, deduced by their ability to provide late blight resistance and *P. infestans* effector recognition (**Figure 4 and 5**). This is not the first described case of *R* gene allelic variants across *Solanum* species. *Rpi-blb1*, *Rpi-sto1* and *Rpi-pta1*, from the Mexican species *S. bulbocastanum*, *S. stoloniferum* and *S. papita* are allelic variants that recognise the same IpiO or PexRD6 *P. infestans* effector (Vleeshouwers *et al.*, 2008). Among allelic variants of late blight resistance genes (i.e. *Rpi-blb3* and *Rpi-hjt1* that recognise Avr2 effectors), overlapping recognition specificities have been previously described (Champouret, 2010) but also more recently (Witek *et al.*, 2021). Moreover, highly similar, but non-allelic *R* genes from the same CNL cluster had different recognition specificities, i.e. *Rpi-vnt1*, *Rpi-*

mcq1, R9a, Ph-3 (Smilde *et al.*, 2005; Foster *et al.*, 2009; Zhang *et al.*, 2014; Jo *et al.*, 2015). In the current report, we describe for the first time that allelic variants of a late blight resistance gene show non-overlapping effector recognition specificities in the combinations that were tested. Remarkably, the recognized effectors belonged to the same effector family, which is a further refinement of our insight in host pathogen co-evolution.

When studying Rpi-chc1 protein domain structure, we identified most of the conserved CNL motifs. Remarkably, the MADA motif (Adachi *et al.*, 2019) was not located downstream of the starting methionine, but downstream of the second methionine in position 46 of the Rpi-chc1 protein. Further research is needed to show if either or both methionines are used as translational start codons. Interestingly, we recently cloned the functional late blight resistance gene from the late blight resistant variety Carolus (*Rpi-Carolus* gene, to be published elsewhere). Rpi-Carolus differed only at 7 amino acid positions from Rpi-ber1, but its N-terminus was shorter as a stop codon was present between the first two methionine codons. This strongly suggests that translation in Rpi-Carolus starts from the second methionine while retaining biological activity.

In contrast to the relatively conserved N termini of the proteins encoded by the *Rpi-chc1* alleles, most interallelic sequence variation localised to the Leucine Rich Repeat regions. Through domain interchange between the *rpi-tub1.3\_RH89-039-16* and a *Rpi-chc1.2* alleles and between Rpi-chc1.1 and Rpi-chc1.2, we discovered that the LRR domain defines recognition specificity (**Figure 6 and 7**). Polymorphisms in the LRR of some NLR receptors were previously shown to determine the effector recognition specificity (Dodds, Lawrence and Ellis, 2001; Shen *et al.*, 2003; Catanzariti *et al.*, 2010; Krasileva, Dahlbeck and Staskawicz, 2010; Ravensdale *et al.*, 2012). In one example, a domain exchange between Rx1 and Gpa2 converted the virus resistance into nematode resistance, and vice versa (Slootweg *et al.*, 2017). The recognition of both nematode and virus could not be combined into one chimeric receptor, as we also observed with Rpi-chc1.1 and Rpi-chc1.2. The reason for this is the overlap between the LRRs involved in recognition.

Most of the amino acids in Rpi-chc1.2 that were required for Avrhc1.2 recognition are located in the LRR solvent exposed motif (xxLxLxxxx) and modification of the tested solvent exposed amino acids led to the partial or complete loss of PexRD31 recognition (**Figure 6b**). The co-requirement of these solvent exposed amino acids suggests that they are involved in recognition of a particular epitope. This observation, combined with the observation of unequal distribution of SAPs, allow us to hypothesise that *Rpi-chc1* alleles evolved through insertion of a stretch of DNA into the LRR domain rather than through accumulation of independent mutation. A similar model of evolution was recently proposed for allelic variants of *Rpi-amr1* (Witek *et al.*, 2021). Such insertions may happen through unequal crossing-over with paralog sequences or through retro-transposition. Interestingly, the evolution of integrated domains in *R* genes has been postulated to be caused by an unknown recombination or transposon independent translocation mechanism (Bailey *et al.*, 2018). The same mechanism may be active in LRR exchange to evolve recognition of non-integrated domains like guardees or decoys (Kourelis and van der Hoorn, 2018) or direct effector recognition.

Interestingly, some of the PexRD31 family members have been previously identified as one of the most rapidly diversifying and fast evolving RXLR effectors in the T30-4 genome, with  $\omega$  values higher than 1.55 (Haas *et al.*, 2009). Additionally, several members of the PexRD12/31 superfamily have recently been characterised to target the host vesicle trafficking machinery by interacting with the vesicle associated membrane protein 72 (VAMP72) family (Petre *et al.*, 2020). Even though both PexRD12 and PexRD31 have the same or functionally similar host targets, they are differentially expressed during *P. infestans* infection. While PexRD12 is highly expressed in all the tested isolates, PexRD31 is expressed at low levels after contact with potato (**Figure S1 and S4**). This would also explain why *Rpi-chc1.2* is not able to provide resistance against *P. infestans*, since most of the isolates have low or undetectable expression levels of *Avrhc1.2* (**Figure S4**). Consequently, clade A (PexRD12) may have evolved to avoid detection by *Rpi-chc1.2* while retaining its targeting of the vesicle trafficking machinery.

Another step in the co-evolution between *Rpi-chc1.1* and the PexRD12/31 family was found by analysing the effector expression in plants expressing *Rpi-chc1.1*. The isolates

that overcome the *Rpi-chc1.1* resistance no longer express PexRD12, while the expression in untransformed Desiree plants was normal and comparable to the expression of *Avrsto1* (**Figure 4b**). Similarly, evasion of recognition through transcriptional suppression, was previously observed in plants expressing *Rpi-vnt1* infected with *P. infestans* (Pel, 2010). Once more, we confirmed the plasticity of the *P. infestans* effector secretion and the fast evolution capacity of some aggressive isolates to break down single Rpi resistances.

The introgression of single *R* genes is driving *P. infestans* to evolve and evade recognition. In order to durably deploy late blight resistance in agriculture, we need novel strategies informed by knowledge of disease resistance genes in varieties, their recognition specificities and the presence of the cognate effectors in the pathogen populations. Virulence information from the field must be rapidly translated to decision support systems (DSS) for the risk prediction and calculation of biocide spraying intervals. Additionally, DSS can be used to determine *R* gene composition of (novel) varieties to be deployed in the next season. To meet these requirements, novel breeding strategies are needed to rapidly tailor the *R* gene contents of the potato varieties to the pathogen populations. In current breeding schemes, it takes 10-15 years to select a late blight resistant potato variety. Moreover, susceptible varieties with dominant market shares will not be easy to replace. A system of varieties with flexible *R* gene content is needed. In other crops this has been accomplished through F1 hybrid varieties. In potato, this route has only recently been opened (Su *et al.*, 2020) and no hybrid potato varieties have reached the market yet. Proof of principle for flexible late blight resistance varieties produced through cisgenesis was provided several years ago (Haverkort *et al.*, 2016). Unfortunately, the EU legislation does not distinguish between cisgenic and transgenic products, making it impossible to market cisgenic varieties. In other parts of the world similar improvement technologies are being deployed to obtain late blight resistance (Habig *et al.*, 2018; Ghislain *et al.*, 2019). It is promising that many different national authorities now apply, or consider separate oversight regulations for events that have been enriched with “cisgenes” or “genes with a history of safe use”. Knowledge as obtained in this study is essential to pursue such enrichment strategies, that can be achieved through transformation of cisgenes, or through novel gene editing tools. We now

know how inactive resistance genes from susceptible varieties could be repaired by replacing minimal fragments with the corresponding fragments of alleles from wild relatives. This would provide an unprecedented accuracy and speed which is not present in introgression breeding.

## Methods

### Plant materials and growth conditions

The wild *Solanum* species used in this study are listed in **Table S6** (Tan *et al.*, 2010; Vleeshouwers *et al.*, 2011a). The potato plants were maintained *in vitro* on MS20 at 24°C under 16/8h day/night regime (Domazakis *et al.*, 2017). The 7650 F1 population was generated by crossing *S. chachoense* (CHC543-5) x *S. chacoense* (CHC544-5). *S. tuberosum* cv. 'Desiree' was used for stable transformations of the different *Rpi-chc1.1* candidates. Four week old *Nicotiana benthamiana* leaves were used for agroinfiltration. The agroinfiltrated plants were kept in climate regulated greenhouse compartments of Unifarm (Wageningen University & Research) at 20-25°C and under 16/8h day/night regime.

### BAC clone isolation and sequencing

The procedure has been described in patent US9551007B2. Briefly: Two different BAC libraries were produced using partial digestion of CHC543-5 genomic DNA with HindIII. Fragments larger than 100 kb were ligated into pBeloBAC or pCC1BAC arms (Epicenter). The BAC clones were collected and stored as bacterial pools of approximately 700 to 1000 white colonies. BAC pools were screened with selected markers and individual clones were identified using colony PCR. The ends of positive individual BACs were sequenced for the purpose of fine mapping RH106G03T and RH137D14\_C37-7-4. The complete inserts were sequenced using shotgun sequencing of 2 kb library fragments generated by partial digestion with EcoR1 by Macrogen, Inc (Seoul, South Korea). Assembly of the sequences resulted in contigs as indicated in **Figure 1** (Genbank accession number MW383255).

## Cloning of *Rpi-chc1* allelic variants and chimeric constructs

The *Rpi-chc1* allelic variants were amplified using genomic DNA from the different wild *Solanum* species using PCR primers as described in **Table S7** and DNA polymerase with proofreading activity. The fragments were cloned into pGEM-T easy vector (Promega) for sequencing. Genbank submission numbers as in **Table S6**. The *Rpi-ber1.1* and *Rpi-tar1.1* genes were amplified using primers in the promotor and terminator. The resulting PCR fragments were cloned into pBINPLUS-PASSA (Jo *et al.*, 2016) and were expressed in transgenic Desiree plants under the control of their native regulatory elements. For transient expression analyses, the coding sequences of the allelic variants were cloned under the *Rpi-chc1.1* regulatory elements (900 bp promotor and 400 bp terminator) into pDEST using a multisite gateway protocol. *Escherichia coli* strain DH10 $\beta$  was transformed with the gateway reaction products and clones with the correct insert were selected. *Agrobacterium tumefaciens* AGL1+VirG was used for transient and stable transformations of *N. benthamiana* leaves and *S. tuberosum* cv. 'Desiree'.

The chimeric constructs were cloned using the Golden Gate modular cloning principle. As acceptor vector, we used a Golden Gate compatible version of pBINPLUS (McBride & Summerfelt, 1990), PBINPLUS-GG (Vossenbergh *et al.*, 2019). The final acceptor vector was constructed to contain 800 bp *Rpi-chc1.1* promoter::CDS::1000 bp *Rpi-ber* terminator (**Figure S5**). The different PCR fragments were amplified using the Phusion High-Fidelity PCR Kit (Thermo SCIENTIFIC) and primers with BsaI sites as overhang (**Table S7**) and purified using the DNA Clean&Concentrator Kit (ZYMO RESEARCH). PCR fragments and the acceptor vector were incubated in Buffer G (Thermo SCIENTIFIC) with ATP 1 mM for thirty cycles of 37°C for 5 min + 16°C for 5 min. Additionally, we performed a final step at 37°C for 10 min, to digest the plasmids wrongly assembled, and 65°C for 20 min, to heat inactivate the BsaI enzyme.

## Hypersensitive cell death assays

Transient expression of the different receptors and PexRD12/31 effectors were performed in four weeks old *N. benthamiana* leaves. R3a/Avr3a was used as a positive

control. All the constructs were agroinfiltrated at an OD<sub>600</sub> of 0.5. Each construct was agroinfiltrated twice on two leaves of four plants in at least two independent experiments. Cell death responses were observed after 3-4 days post inoculation.

### **Phylogenetic analysis of Rpi-chc1.1 homologs and the PexRD12/31 superfamily**

The sequences of the PexRD12/31 effectors were retrieved from the *P. infestans* T30-4 genome (Haas *et al.*, 2009). Twenty family members were found to form the PexRD12/31 superfamily. The coding sequences of the *Rpi-chc1* variants as obtained in this study, were aligned using MUSCLE and a neighbour joining tree was calculated using Megalign from the DNASTAR package. The closest homolog of *Rpi-chc1.1* from the DM reference genome (SoltuDM10G021850.1) was used as an outgroup.

The protein sequences of PexRD12/31 effectors were aligned using Clustal OMEGA and manually edited in MEGAX (Sievers *et al.*, 2011; Kumar *et al.*, 2018). The phylogenetic relationship was inferred using the Maximum Likelihood method based on the JTT matrix-based model in MEGAX with 1000 bootstraps (Jones, Taylor and Thornton, 1992). The tree with the highest log likelihood was shown. The two more distant effectors PITG\_16428 and PITG\_09577, served as an outgroup.

### ***P. infestans* isolates and Detach Leaf Assay (DLA)**

The *P. infestans* isolates used in this study (90128, IPO-C and NL08645) were retrieved from our in-house collection. Isolates were grown at 15°C on solid rye medium in the dark (Caten and Jinks, 1968). After two weeks, sporulating mycelium was flooded with 20 mL of ice-cold water, adjusted to 70 zoospores/μL and incubated at 4°C for 2-3 hours. After the incubation, the detached leaves were inoculated with 10 μL of the zoospore suspension on the abaxial side of the leaves. Detached leaves were inserted into wet floral foam. For each biological replicate the three leaflets from four leaves from two independent plants were used. Twelve spots on each leaf were inoculated with the zoospore suspension and closed in a plastic bag, to maintain high humidity. The leaves were kept in a climate cell at 18°C for 5 days. Disease resistance was scored on a scale

from 1 to 10 for each leaflet. 10=no symptoms; 9=HR no larger than the inoculum droplet; 8=HR lesion of up to 0,5 cm diameter; 7=diffuse lesions up to 1 cm diameter, no sporulation, no water soaking; 5= lesions larger than 1 cm sometimes with water soaking, no sporulation; 4=large water soaked lesions with sporulation only visible through binoculars; 2= large lesions with macroscopically visible sporulation on one side of the leaflet; 1= large lesions with macroscopically visible sporulation on both sides of the leaflet.

### **Relative effector and *R* gene expression**

The *P. infestans* effectors used in this study are listed in **Table S5**. The different genotypes were inoculated with the different *P. infestans* isolates and samples were collected after 0, 3, 8, 24, 48, 72, 96 and 120 hours. Infected plant material with the different *P. infestans* isolates was collected and RNA was isolated using RNA Purification Kit (QIAGEN). The isolated RNA was converted into cDNA using the QuantiTect Reverse Transcription Kit (QIAGEN). The primers used in this study are listed in **Table S7**. The expression of the different effectors in the infected material was evaluated using RT-qPCR SYBR Green (Bio-Rad). The samples were heated to 95°C for 2 min. Then 40 cycles of 15 sec at 95°C, 30 sec at 60°C and 30 sec at 72°C. Fluorescence was measured after each cycle. After the final amplification cycle a melting curve was calculated. Relative gene expression was calculated using the  $2^{-\Delta\Delta CT}$  method (Livak and Schmittgen, 2001). The normalised gene expression was obtained by dividing the relative gene expression by the relative *P. infestans* elongation factor 2 gene (*ef2*) expression.

### **sgRNA and CRISPR-Cas9 construct design**

The CRISPOR web tool (<http://crispor.org>) was used to design the sgRNAs with lower off-target and higher on-target potentials (Concordet and Haeussler, 2018).

A Modular Cloning (MoClo) system based on the Golden Gate cloning technology was used to assemble the different sgRNAs and binary vectors as previously described for tomato mutagenesis (Engler, Kandzia and Marillonnet, 2008; Weber *et al.*, 2011). Briefly, each sgRNA was fused to the *Arabidopsis thaliana* U6-26 promoter as *AtU6*-

26::gRNA. The Level 1 constructs *pICH47732-pNOS::NPTII::tOCS*, *pICH47742-p2x35S::hCas9::tNOS* and the linker *pICH41780* were used to build the Level 2 vector *pICSL4723* (Werner *et al.*, 2012). The primers used for cloning the gRNAs are listed in **Table S7**.

## **Data availability statement**

Supplementary figures and tables are available as supporting information through the TPJ website. All described sequences have been submitted to GenBank. Accession numbers are listed in Table S6.

## **Acknowledgements**

The first part of this research was performed in the DuRPh program, funded by the Ministry of Agriculture, Nature and Food Quality in the Netherlands. The second part of the research was funded by the Ministry of Infrastructure and Water Management in the Netherlands through the NWO-TTW program Biotechnology and Safety (projectnumber 15815). We thank Jan de Boer and Adillah Tan for advice in selecting BAC clones and marker sequences from chr10. Evert Jacobsen and Clemens van der Wiel are thanked for their advice about regulation and safety aspects of plant biotechnology. Sidrat Abdullah is thanked for testing late blight resistance of transgenic potato plants harbouring *Rpi-chc1* allelic variants. Last but not least, we thank Marjan Bergervoet, Gert van Arkel, Koen Pelgrom, Dirk Jan Huigen and Isolde Pereira for plant transformations, molecular biology assistance and plant maintenance.

## **Author contributions**

JV planned and designed the research; DML, MN and LK performed the majority of the experiments; SK provided the Pex-RD set; DML and JV wrote the manuscript; RV proofread the manuscript and provided the essential research environment. SA, HS and KS contributed by mapping, cloning and characterization of *Rpi-chc1* allelic variants. RS,

AL and AAH contributed through the identification of Avrhc1 and their differential recognition specificities by *Rpi-chc1* allelic variants.

## Conflict of interest statement

Some results presented in this manuscript have been included in a patent application.

## Short supporting material legends

**Figure S1:** Effector and *R* gene expression in potato leaves inoculated with *P. infestans*.

**Figure S2:** Rpi-chc1.1 protein domain organization.

**Figure S3:** Localization of *PexRD12/31* effectors in the *P. infestans* T30-4 contigs.

**Figure S4:** *PexRD12/31* effector expression in potato leaves infected with *P. infestans* from PenSeq dataset.

**Figure S5:** pBINPLUS-PASSA-GG vector map.

**Table S1:** CRISPR-Cas9 targeting of *Rpi-chc1.1* allele in *S. chacoense* resistant accession CHC543-5.

**Table S2:** Late blight resistance assessment of different *Rpi-chc1* alleles.

**Table S3:** Segregation of markers and late blight resistance of *Rpi-chc1.1* and *Rpi-chc1.2*.

**Table S4:** Functional expression of *Rpi-chc1.1* in Desiree transgenic events correlates with responsiveness to *PexRD12*.

**Table S5:** *P. infestans* effectors used in this study.

**Table S6:** Accession numbers of *Solanum* genotypes and *Rpi-chc1* sequences.

**Table S7:** Primers used in this study.

## References

- Adachi, H. *et al.* (2019) 'An N-terminal motif in NLR immune receptors is functionally conserved across distantly related plant species', *eLife*, 8, p. e49956. doi: 10.7554/eLife.49956.
- Aguilera-Galvez, C. *et al.* (2018) 'Two different R gene loci co-evolved with Avr2 of *Phytophthora infestans* and confer distinct resistance specificities in potato', *Studies in Mycology*, 89, pp. 105–115. doi: 10.1016/j.simyco.2018.01.002.
- Bailey, P. C. *et al.* (2018) 'Dominant integration locus drives continuous diversification of plant immune receptors with exogenous domain fusions', *Genome Biology*, 19(1), p. 23. doi: 10.1186/s13059-018-1392-6.
- Ballvora, A. *et al.* (2002) 'The R1 gene for potato resistance to late blight (*Phytophthora infestans*) belongs to the leucine zipper/NBS/LRR class of plant resistance genes', *The Plant Journal*, 30(3), pp. 361–371. doi: 10.1046/j.1365-313X.2001.01292.x.
- Bendahmane, A. *et al.* (2002) 'Constitutive gain-of-function mutants in a nucleotide binding site-leucine rich repeat protein encoded at the Rx locus of potato', *The Plant Journal*, 32(2), pp. 195–204. doi: 10.1046/j.1365-313X.2002.01413.x.
- Callaway, E. (2013) 'Pathogen genome tracks Irish potato famine back to its roots', *Nature*, p. nature.2013.13021. doi: 10.1038/nature.2013.13021.
- Campbell, T. A. (2003) 'Investigation of variations in NBS motifs in alfalfa (*Medicago sativa*), *M. edgeworthii*, and *M. ruthenica*', *Canadian Journal of Plant Science*, 83(2), pp. 371–376. doi: 10.4141/P02-002.
- Catanzariti, A.-M. *et al.* (2010) 'The AvrM Effector from Flax Rust Has a Structured C-Terminal Domain and Interacts Directly with the M Resistance Protein', *Molecular Plant-Microbe Interactions*®, 23(1), pp. 49–57. doi: 10.1094/MPMI-23-1-0049.

Caten, C. E. and Jinks, J. L. (1968) 'Spontaneous variability of single isolates of *Phytophthora infestans*. I. Cultural variation', *Canadian Journal of Botany*, 46(4), pp. 329–348. doi: 10.1139/b68-055.

Champouret, N. (2010) *Functional genomics of Phytophthora infestans effectors and Solanum resistance genes*. Ph.D. Dissertation. Wageningen University. Available at: <https://edepot.wur.nl/138174>.

Concordet, J.-P. and Haeussler, M. (2018) 'CRISPOR: intuitive guide selection for CRISPR/Cas9 genome editing experiments and screens', *Nucleic Acids Research*, 46(W1), pp. W242–W245. doi: 10.1093/nar/gky354.

Danot, O. *et al.* (2009) 'Wheel of Life, Wheel of Death: A Mechanistic Insight into Signaling by STAND Proteins', *Structure*, 17(2), pp. 172–182. doi: 10.1016/j.str.2009.01.001.

Devaux, A. *et al.* (2020) 'Global Food Security, Contributions from Sustainable Potato Agri-Food Systems', in Campos, H. and Ortiz, O. (eds) *The Potato Crop*. Cham: Springer International Publishing, pp. 3–35. doi: 10.1007/978-3-030-28683-5\_1.

Dodds, P. N., Lawrence, G. J. and Ellis, J. G. (2001) 'Six Amino Acid Changes Confined to the Leucine-Rich Repeat  $\beta$ -Strand/ $\beta$ -Turn Motif Determine the Difference between the P and P2 Rust Resistance Specificities in Flax', *The Plant Cell*, 13(1), pp. 163–178. doi: 10.1105/tpc.13.1.163.

Domazakis, E. *et al.* (2017) 'Effectoromics-Based Identification of Cell Surface Receptors in Potato', in Shan, L. and He, P. (eds) *Plant Pattern Recognition Receptors*. New York, NY: Springer New York (Methods in Molecular Biology), pp. 337–353. doi: 10.1007/978-1-4939-6859-6\_29.

Engler, C., Kandzia, R. and Marillonnet, S. (2008) 'A One Pot, One Step, Precision Cloning Method with High Throughput Capability', *PLoS ONE*. Edited by H. A. El-Shemy, 3(11), p. e3647. doi: 10.1371/journal.pone.0003647.

Foster, S. J. *et al.* (2009) 'Rpi-vnt1.1 , a Tm-2 Homolog from *Solanum venturii* , Confers Resistance to Potato Late Blight', *Molecular Plant-Microbe Interactions®*, 22(5), pp. 589–600. doi: 10.1094/MPMI-22-5-0589.

Fry, W. (2008) 'Phytophthora infestans: the plant (and R gene) destroyer', *Molecular Plant Pathology*, 9(3), pp. 385–402. doi: 10.1111/j.1364-3703.2007.00465.x.

Ghislain, M. *et al.* (2019) 'Stacking three late blight resistance genes from wild species directly into African highland potato varieties confers complete field resistance to local blight races', *Plant Biotechnology Journal*, 17(6), pp. 1119–1129. doi: 10.1111/pbi.13042.

Goss, E. M., Press, C. M. and Grünwald, N. J. (2013) 'Evolution of RXLR-Class Effectors in the Oomycete Plant Pathogen *Phytophthora ramorum*', *PLoS ONE*. Edited by A. Palsson, 8(11), p. e79347. doi: 10.1371/journal.pone.0079347.

Haas, B. J. *et al.* (2009) 'Genome sequence and analysis of the Irish potato famine pathogen *Phytophthora infestans*', *Nature*, 461(7262), pp. 393–398. doi: 10.1038/nature08358.

Habig, J. W. *et al.* (2018) 'Food safety evaluation for R-proteins introduced by biotechnology: A case study of VNT1 in late blight protected potatoes', *Regulatory Toxicology and Pharmacology*, 95, pp. 66–74. doi: 10.1016/j.yrtph.2018.03.008.

Haverkort, A. J. *et al.* (2016) 'Durable Late Blight Resistance in Potato Through Dynamic Varieties Obtained by Cisgenesis: Scientific and Societal Advances in the DuRPh Project', *Potato Research*, 59(1), pp. 35–66. doi: 10.1007/s11540-015-9312-6.

Huang, S. *et al.* (2004) 'The R3 Resistance to *Phytophthora infestans* in Potato is Conferred by Two Closely Linked R Genes with Distinct Specificities', *Molecular Plant-Microbe Interactions*®, 17(4), pp. 428–435. doi: 10.1094/MPMI.2004.17.4.428.

Huang, S. *et al.* (2005) 'Comparative genomics enabled the isolation of the R3a late blight resistance gene in potato: Cloning the potato late blight R3a gene by synteny', *The Plant Journal*, 42(2), pp. 251–261. doi: 10.1111/j.1365-3113X.2005.02365.x.

Jo, K.-R. *et al.* (2015) 'Characterisation of the late blight resistance in potato differential MaR9 reveals a qualitative resistance gene, R9a, residing in a cluster of Tm-2 2 homologs on chromosome IX', *Theoretical and Applied Genetics*, 128(5), pp. 931–941. doi: 10.1007/s00122-015-2480-6.

- Jo, K.-R. *et al.* (2016) 'Problematic Crops: 1. Potatoes: Towards Sustainable Potato Late Blight Resistance by Cisgenic *R* Gene Pyramiding', in Collinge, D. B. (ed.) *Plant Pathogen Resistance Biotechnology*. Hoboken, NJ: John Wiley & Sons, Inc, pp. 171–191. doi: 10.1002/9781118867716.ch9.
- Jones, D. T., Taylor, W. R. and Thornton, J. M. (1992) 'The rapid generation of mutation data matrices from protein sequences', *Bioinformatics*, 8(3), pp. 275–282. doi: 10.1093/bioinformatics/8.3.275.
- Jones, J. D. G., Vance, R. E. and Dangl, J. L. (2016) 'Intracellular innate immune surveillance devices in plants and animals', *Science*, 354(6316), pp. aaf6395–aaf6395. doi: 10.1126/science.aaf6395.
- Jupe, F. *et al.* (2013) 'Resistance gene enrichment sequencing (RenSeq) enables reannotation of the NB-LRR gene family from sequenced plant genomes and rapid mapping of resistance loci in segregating populations', *The Plant Journal*, 76(3), pp. 530–544. doi: 10.1111/tpj.12307.
- Karki, H. S., Jansky, S. H. and Halterman, D. A. (2021) 'Screening of Wild Potatoes Identifies New Sources of Late Blight Resistance', *Plant Disease*, 105(2), pp. 368–376. doi: 10.1094/PDIS-06-20-1367-RE.
- Kourelis, J. and van der Hoorn, R. A. L. (2018) 'Defended to the Nines: 25 Years of Resistance Gene Cloning Identifies Nine Mechanisms for R Protein Function', *The Plant Cell*, 30(2), pp. 285–299. doi: 10.1105/tpc.17.00579.
- Krasileva, K. V., Dahlbeck, D. and Staskawicz, B. J. (2010) 'Activation of an Arabidopsis Resistance Protein Is Specified by the in Planta Association of Its Leucine-Rich Repeat Domain with the Cognate Oomycete Effector', *The Plant Cell*, 22(7), pp. 2444–2458. doi: 10.1105/tpc.110.075358.
- Kumar, S. *et al.* (2018) 'MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms', *Molecular Biology and Evolution*. Edited by F. U. Battistuzzi, 35(6), pp. 1547–1549. doi: 10.1093/molbev/msy096.

Leipe, D. D., Koonin, E. V. and Aravind, L. (2004) 'STAND, a Class of P-Loop NTPases Including Animal and Plant Regulators of Programmed Cell Death: Multiple, Complex Domain Architectures, Unusual Phyletic Patterns, and Evolution by Horizontal Gene Transfer', *Journal of Molecular Biology*, 343(1), pp. 1–28. doi: 10.1016/j.jmb.2004.08.023.

Leister, D. (2004) 'Tandem and segmental gene duplication and recombination in the evolution of plant disease resistance genes', *Trends in Genetics*, 20(3), pp. 116–122. doi: 10.1016/j.tig.2004.01.007.

Li, G. *et al.* (2011) 'Cloning and Characterization of R3b; Members of the R3 Superfamily of Late Blight Resistance Genes Show Sequence and Functional Divergence', *Molecular Plant-Microbe Interactions*®, 24(10), pp. 1132–1142. doi: 10.1094/MPMI-11-10-0276.

Lin, X. *et al.* (2020) 'Identification of Avramr1 from *Phytophthora infestans* using long read and cDNA pathogen-enrichment sequencing (PenSeq)', *Molecular Plant Pathology*, 21(11), pp. 1502–1512. doi: 10.1111/mpp.12987.

Livak, K. J. and Schmittgen, T. D. (2001) 'Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the  $2^{-\Delta\Delta CT}$  Method', *Methods*, 25(4), pp. 402–408. doi: 10.1006/meth.2001.1262.

Lokossou, A. A. *et al.* (2009) 'Exploiting Knowledge of R/Avr Genes to Rapidly Clone a New LZ-NBS-LRR Family of Late Blight Resistance Genes from Potato Linkage Group IV', *Molecular Plant-Microbe Interactions*®, 22(6), pp. 630–641. doi: 10.1094/MPMI-22-6-0630.

McBride, K. E. and Summerfelt, K. R. (1990) 'Improved binary vectors for *Agrobacterium*-mediated plant transformation', *Plant Molecular Biology*, 14(2), pp. 269–276. doi: 10.1007/BF00018567.

Mcdowell, J. M. and Simon, S. A. (2006) 'Recent insights into R gene evolution', *Molecular Plant Pathology*, 7(5), pp. 437–448. doi: 10.1111/j.1364-3703.2006.00342.x.

Pal, A., Chakrabarti, A. and Basak, J. (2007) 'New motifs within the NB-ARC domain of R proteins: Probable mechanisms of integration of geminiviral signatures within the host species of

Fabaceae family and implications in conferring disease resistance', *Journal of Theoretical Biology*, 246(3), pp. 564–573. doi: 10.1016/j.jtbi.2007.01.013.

Park, T.-H. *et al.* (2009) 'Two distinct potato late blight resistance genes from *Solanum berthaultii* are located on chromosome 10', *Euphytica*, 165(2), pp. 269–278. doi: 10.1007/s10681-008-9784-4.

Pel, M. A. *et al.* (2009) 'Mapping and Cloning of Late Blight Resistance Genes from *Solanum venturii* Using an Interspecific Candidate Gene Approach', *Molecular Plant-Microbe Interactions*®, 22(5), pp. 601–615. doi: 10.1094/MPMI-22-5-0601.

Pel, M. A. (2010) *Mapping, isolation and characterization of genes responsible for late blight resistance in potato*. Wageningen University.

Petre, B. *et al.* (2020) *Host-interactor screens of Phytophthora infestans RXLR proteins reveal vesicle trafficking as a major effector-targeted process*. preprint. Plant Biology. doi: 10.1101/2020.09.24.308585.

Rairdan, G. J. *et al.* (2008) 'The Coiled-Coil and Nucleotide Binding Domains of the Potato Rx Disease Resistance Protein Function in Pathogen Recognition and Signaling', *The Plant Cell*, 20(3), pp. 739–751. doi: 10.1105/tpc.107.056036.

Rauscher, G. M. *et al.* (2006) 'Characterization and mapping of Rpi-ber , a novel potato late blight resistance gene from *Solanum berthaultii*', *Theoretical and Applied Genetics*, 112(4), pp. 674–687. doi: 10.1007/s00122-005-0171-4.

Ravensdale, M. *et al.* (2012) 'Intramolecular Interaction Influences Binding of the Flax L5 and L6 Resistance Proteins to their AvrL567 Ligands', *PLoS Pathogens*. Edited by J.-R. Xu, 8(11), p. e1003004. doi: 10.1371/journal.ppat.1003004.

Reubold, T. F., Wohlgemuth, S. and Eschenburg, S. (2011) 'Crystal Structure of Full-Length Apaf-1: How the Death Signal Is Relayed in the Mitochondrial Pathway of Apoptosis', *Structure*, 19(8), pp. 1074–1083. doi: 10.1016/j.str.2011.05.013.

Rietman, H. *et al.* (2012) 'Qualitative and Quantitative Late Blight Resistance in the Potato Cultivar Sarpio Mira Is Determined by the Perception of Five Distinct RXLR Effectors', *Molecular Plant-Microbe Interactions*®, 25(7), pp. 910–919. doi: 10.1094/MPMI-01-12-0010-R.

Saraste, M., Sibbald, P. R. and Wittinghofer, A. (1990) 'The P-loop — a common motif in ATP- and GTP-binding proteins', *Trends in Biochemical Sciences*, 15(11), pp. 430–434. doi: 10.1016/0968-0004(90)90281-F.

Sharma, S. K. *et al.* (2013) 'Construction of Reference Chromosome-Scale Pseudomolecules for Potato: Integrating the Potato Genome with Genetic and Physical Maps', *G3&#58;Genes|Genomes|Genetics*, 3(11), pp. 2031–2047. doi: 10.1534/g3.113.007153.

Shen, Q.-H. *et al.* (2003) 'Recognition Specificity and RAR1/SGT1 Dependence in Barley Mla Disease Resistance Genes to the Powdery Mildew Fungus', *The Plant Cell*, 15(3), pp. 732–744. doi: 10.1105/tpc.009258.

Sievers, F. *et al.* (2011) 'Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega', *Molecular Systems Biology*, 7(1), p. 539. doi: 10.1038/msb.2011.75.

Slootweg, E. *et al.* (2017) 'Sequence Exchange between Homologous NB-LRR Genes Converts Virus Resistance into Nematode Resistance, and Vice Versa', *Plant Physiology*, 175(1), pp. 498–510. doi: 10.1104/pp.17.00485.

Smilde, W. D. *et al.* (2005) 'Solanum mochiquense chromosome IX carries a novel late blight resistance gene Rpi-moc1', *Theoretical and Applied Genetics*, 110(2), pp. 252–258. doi: 10.1007/s00122-004-1820-8.

Song, J. *et al.* (2003) 'Gene RB cloned from Solanum bulbocastanum confers broad spectrum resistance to potato late blight', *Proceedings of the National Academy of Sciences*, 100(16), pp. 9128–9133. doi: 10.1073/pnas.1533501100.

Su, Y. *et al.* (2020) 'Introgression of Genes for Resistance against Phytophthora infestans in Diploid Potato', *American Journal of Potato Research*, 97(1), pp. 33–42. doi: 10.1007/s12230-019-09741-8.

Sukarta, O. C. A., Slootweg, E. J. and Goverse, A. (2016) 'Structure-informed insights for NLR functioning in plant immunity', *Seminars in Cell & Developmental Biology*, 56, pp. 134–149. doi: 10.1016/j.semcdb.2016.05.012.

Tan, M. Y. A. *et al.* (2010) 'The effect of pyramiding *Phytophthora infestans* resistance genes R Pi-mcd1 and R Pi-ber in potato', *Theoretical and Applied Genetics*, 121(1), pp. 117–125. doi: 10.1007/s00122-010-1295-8.

The Potato Genome Sequencing Consortium (2011) 'Genome sequence and analysis of the tuber crop potato', *Nature*, 475(7355), pp. 189–195. doi: 10.1038/nature10158.

Vleeshouwers, V. G. *et al.* (2011) 'SolRgene: an online database to explore disease resistance genes in tuber-bearing *Solanum* species', *BMC Plant Biology*, 11(1), p. 116. doi: 10.1186/1471-2229-11-116.

Vleeshouwers, V. G. A. A. *et al.* (1999) 'A Laboratory Assay for *Phytophthora infestans* Resistance in Various *Solanum* Species Reflects the Field Situation', *European Journal of Plant Pathology*, 105(3), pp. 241–250. doi: 10.1023/A:1008710700363.

Vleeshouwers, V. G. A. A. *et al.* (2008) 'Effector Genomics Accelerates Discovery and Functional Profiling of Potato Disease Resistance and *Phytophthora infestans* Avirulence Genes', *PLoS ONE*. Edited by H. A. El-Shemy, 3(8), p. e2875. doi: 10.1371/journal.pone.0002875.

Vleeshouwers, V. G. A. A. *et al.* (2011) 'Understanding and Exploiting Late Blight Resistance in the Age of Effectors', *Annual Review of Phytopathology*, 49(1), pp. 507–531. doi: 10.1146/annurev-phyto-072910-095326.

van der Vossen, E. A. G. *et al.* (2005) 'The Rpi-blb2 gene from *Solanum bulbocastanum* is an Mi-1 gene homolog conferring broad-spectrum late blight resistance in potato: Isolation of the late blight resistance gene Rpi-blb2', *The Plant Journal*, 44(2), pp. 208–222. doi: 10.1111/j.1365-313X.2005.02527.x.

van der Vossen, E. *et al.* (2003) 'An ancient R gene from the wild potato species *Solanum bulbocastanum* confers broad-spectrum resistance to *Phytophthora infestans* in cultivated potato and tomato', *The Plant Journal*, 36(6), pp. 867–882. doi: 10.1046/j.1365-313X.2003.01934.x.

Vossen, J. H. *et al.* (2009) 'Cloning and exploitation of a functional R gene from *Solanum chacoense*'.

Vossen, J. H. *et al.* (2013) 'Novel applications of motif-directed profiling to identify disease resistance genes in plants', *Plant Methods*, 9(1), p. 37. doi: 10.1186/1746-4811-9-37.

Vossen, J. H. *et al.* (2016) 'The *Solanum demissum* R8 late blight resistance gene is an Sw-5 homologue that has been deployed worldwide in late blight resistant varieties', *Theoretical and Applied Genetics*, 129(9), pp. 1785–1796. doi: 10.1007/s00122-016-2740-0.

Warren, R. F. *et al.* (1998) 'A Mutation within the Leucine-Rich Repeat Domain of the Arabidopsis Disease Resistance Gene RPS5 Partially Suppresses Multiple Bacterial and Downy Mildew Resistance Genes', *The Plant Cell*, 10(9), pp. 1439–1452. doi: 10.1105/tpc.10.9.1439.

Weber, E. *et al.* (2011) 'A Modular Cloning System for Standardized Assembly of Multigene Constructs', *PLoS ONE*. Edited by J. Peccoud, 6(2), p. e16765. doi: 10.1371/journal.pone.0016765.

Wendler, P. *et al.* (2012) 'Structure and function of the AAA+ nucleotide binding pocket', *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research*, 1823(1), pp. 2–14. doi: 10.1016/j.bbamcr.2011.06.014.

Werner, S. *et al.* (2012) 'Fast track assembly of multigene constructs using Golden Gate cloning and the MoClo system', *Bioengineered*, 3(1), pp. 38–43. doi: 10.4161/bbug.3.1.18223.

Witek, K. *et al.* (2016) 'Accelerated cloning of a potato late blight–resistance gene using RenSeq and SMRT sequencing', *Nature Biotechnology*, 34(6), pp. 656–660. doi: 10.1038/nbt.3540.

Witek, K. *et al.* (2020) *A complex resistance locus in Solanum americanum recognizes a conserved Phytophthora effector*. preprint. Plant Biology. doi: 10.1101/2020.05.15.095497.

Witek, K. *et al.* (2021) 'A complex resistance locus in *Solanum americanum* recognizes a conserved *Phytophthora* effector', *Nature Plants*, 7(2), pp. 198–208. doi: 10.1038/s41477-021-00854-9.

Zhang, C. *et al.* (2014) 'The Ph-3 gene from *Solanum pimpinellifolium* encodes CC-NBS-LRR protein conferring resistance to *Phytophthora infestans*', *Theoretical and Applied Genetics*, 127(6), pp. 1353–1364. doi: 10.1007/s00122-014-2303-1.

## Figure legends

### Figure 1. Map based cloning of *Rpi-chc1.1*.

(A) Genetic map of *P. infestans* (isolate 90128) resistance from CHC543-5. The number between the markers represents the number of recombinants found in a population derived from 2357 seedlings. Markers starting with RH were derived from BAC end sequences generated by PGSC. Marker B2-S represents the BAC end marker from clone 2. The black horizontal line represents the interval of *Rpi-chc1.1*. (B) Two BAC clones were isolated to generate the physical map. Annotation revealed the presence of NB-LRR genes, genes with or without predicted function and transposable elements. Three complete NB-LRR (B2-1, B2-2 and B2-3) genes between flanking markers RH106G03T and B2-S were selected as candidates. (C) The three candidates were expressed through agroinfiltration in *N. benthamiana* leaves. An empty vector (EV) and *Rpi-blb1* were used as negative and positive controls, respectively. Only candidate B2-3 was able to compromise the growth of *P. infestans* isolate 90128. (D) The three candidates were stably transformed into the potato variety Desiree. After inoculation with isolate 90128, only the candidate B2-3 was able to provide resistance. Untransformed Desiree and Desiree plants stably transformed with *Rpi-blb1* were used as negative and positive controls, respectively.

### Figure 2. CRISPR-Cas9 induced frameshifts in B2-3 candidate (*Rpi-chc1.1*) leads to *Phytophthora* susceptibility.

(A) Three different sgRNAs were designed to target the LRR domain of candidate B2-3. (B) A construct with Cas9 and all three sgRNAs was stably transformed into the resistant genotype CHC543-5. Transformation events were inoculated with *P. infestans* 90128 and IPO-C isolates. 48% of the transformation events became susceptible to both isolates

(Table S1). (C) One resistant (#1) and three susceptible events (#21, 24, 26) were genotyped together with the untransformed plant CHC543-5. The three susceptible events contained frameshift mutations, while the tested resistant event had no mutations in the B2-3 candidate.

**Figure 3. *Rpi-chc1* allele mining.**

(A) *Rpi-chc1.1* allele belongs to the immune receptor family. Different motifs were found in the different CNL receptor domains. The LRR domain consists of 29 imperfect repeats. (B) Sixteen *Rpi-chc1.1*-like sequences were cloned from eleven different diploid *Solanum* accessions. The phylogenetic analysis of the DNA sequences led to the identification of four clades. The branch lengths represent the percentage of phylogenetic distance. In the protein alignment mismatches are highlighted in red, gaps are indicated by dashes (-), while insertions relative to *Rpi-chc1.1* (anchor) are indicated by a blue bracket.

**Figure 4. The RXLR effector PexRD12 corresponds to Avrhc1.1.**

(A) The three *Rpi-chc1.1* candidates were co-agroinfiltrated with the Pex effector collection in *N. benthamiana* leaves to screen for Avrhc1.1. *Rpi-chc1.1* induces cell death when co-expressed with both PITG\_16233 and PITG\_16240, from the PexRD12 family. *R3a* and *Avr3a* were used as negative controls, a mix of *R3a* and *Avr3a* was used as a positive control. (B) Relative *Avrhc1.1* and *Avrsto1* effector expression was measured in infected plant material from a field trial with natural *P. infestans* infection (2013, Wageningen). Untransformed Desiree plants, and Desiree plants transformed with *Rpi-chc1.1* (Desiree:*Rpi-chc1.1*), or *Rpi-sto1* (Desiree:*Rpi-sto1*) were used in this study. Relative expression data was obtained by dividing the relative effector gene expression by the *P. infestans* elongation factor 2 gene (*ef2*) expression. Three independent samples were included in the RT-qPCR experiment. Stars represent statistical difference in a two sample T-test,  $p < 0.008$ .

**Figure 5. *Rpi-chc1* alleles show non-overlapping recognition of the PexRD12/31 effector superfamily.**

(A) Twenty members of the PexRD12/31 superfamily were found in *P. infestans* isolate T30-4. In the amino acid sequence, we can distinguish a signal peptide in the N-terminus, the conserved RXLR-EER motifs in the center, and the effector domain in the C-terminus. Some PexRD12/31 family members differed at the nucleotide level but were identical at the protein level (PITG\_16245 = PITG\_16418; PITG\_16233 = PITG\_16240; PITG\_20934 = PITG\_20936; PITG\_16409 = PITG\_16424). The phylogenetic analysis of the complete protein sequences led to the identification of five clades. This analysis was performed in MEGA X by using the Maximum Likelihood method based on the JTT matrix-based model. The tree with the highest log likelihood (-766) is shown. The bootstrapping values, which indicates the percentage of trees that had the particular branch, are shown in each branch. In the protein alignment: blue, hydrophobic residues (A, I, L, M, F, W, V and C); red, positively charged residues (K and R); magenta, negatively charged residues (E and D); green, polar residues (N, Q, S and T); pink, cysteines residue (C); orange, glycines residue (G); yellow, prolines residue (P); cyan, aromatic residues (H and Y); and white, unconserved residues or gaps. (B) Different *Rpi-chc1* allelic variants were co-agroinfiltrated in *N. benthamiana* with a member from each PexRD12/31 clade. While variants from clade 1 recognize both PexRD12 A1 and A2 clades, *Rpi-chc1* variants from clade 2 recognize PexRD31 B and C. Receptors from clade 3 and 4 do not recognise any PexRD12/31 effector. A mix of *R3a* and *Avr3a* was used as a positive control.


**Figure 6.** Domain exchanges between *Rpi-chc1.2* and *rpi-tub1.3\_RH89-039-16*.

(A) The positions of SAPs and the corresponding protein domains are indicated on top. *Rpi-chc1.2* and *rpi-tub1.3\_RH89-039-16*, are represented as light blue and yellow bars, respectively. Below, the domain exchanges are shown. The chimeric constructs were co-agroinfiltrated with the PexRD12/31 effectors in *N. benthamiana* leaves. After 4 days, the HR that was visible and recorded. Experiments were repeated three times with 12 inoculation sites each time. A representative leaf for the response of each chimeric construct is shown in the right panel. GUS was used as a negative control. It is concluded that the exchange of the complete LRR domain led to recognition of PexRD31. With the final construct, RH::C2\_14-19, the exchange of only nine amino acids led to the activation of the *rpi-tub1.3\_RH89-036-16* protein. (B) Seven new modified receptors were derived

from RH::C2\_14-19 in order to pinpoint the amino acids involved in the Rpi-chc1.2 recognition specificity. SAPs present in Rpi-chc1.2 are highlighted in blue font. Most of the SAPs are located in the solvent exposed xxLxLxxxx motif of the LRR 16-19. The chimeric constructs were co-agroinfiltrated with the PexRD12/31 effectors in *N. benthamiana* leaves. A representative leaf for the response of each chimeric construct is shown in the right panel. Experiments were repeated three times with 12 inoculation sites each time. GUS was used as a negative control. The modification of the tested residues of the Rpi-chc1.2 solvent exposed specific amino acids (blue) for the corresponding amino acid present in rpi-tub1.3\_RH89-039-16 (yellow), led to the partial or complete loss of effector recognition.


**Figure 7.** The effector recognition specificity could be exchanged between Rpi-chc1.1 and Rpi-chc1.2.

The alignment of Rpi-chc1.1 and Rpi-chc1.2 shows that all the 41 amino acids polymorphisms (red bars) are located in the LRR domain. The chimeric constructs were co-agroinfiltrated with the PexRD12/31 effectors in *N. benthamiana* leaves. A representative leaf for the response of each chimeric construct is shown in the right panel. Experiments were repeated three times with 12 inoculation sites each time. GUS was used as a negative control. In the construct C1::C2\_14-23, we could see that the LRR 16-19 reappear again as determining for the PexRD31 recognition. The required domain exchanges of the Rpi-chc1.1 LRR are more complex and encompasses almost the complete LRR.




**Figure 1.** Map based cloning of *Rpi-chc1.1*.

(A) Genetic map of *P. infestans* (isolate 90128) resistance from CHC543-5. The number between the markers represents the number of recombinants found in a population derived from 2357 seedlings. Markers starting with RH were derived from BAC end sequences generated by PGSC. Marker B2-S represents the BAC end marker from clone 2. The black horizontal line represents the interval of *Rpi-chc1.1*. (B) Two BAC clones were isolated to generate the physical map. Annotation revealed the presence of NB-LRR genes, genes with or without predicted function and transposable elements. Three complete NB-LRR (B2-1, B2-2 and B2-3) genes between flanking markers RH106G03T and B2-S were selected as candidates. (C) The three candidates were expressed through agroinfiltration in *N. benthamiana* leaves. An empty vector (EV) and *Rpi-blb1* were used as negative and positive controls, respectively. Only candidate B2-3 was able to compromise the growth of *P. infestans* isolate 90128. (D) The three candidates were stably transformed into the potato variety Desiree. After inoculation with isolate 90128, only the candidate B2-3 was able to provide resistance. Untransformed Desiree and Desiree plants stably transformed with *Rpi-blb1* were used as negative and positive controls, respectively.




**Figure 2.** CRISPR-Cas9 induced frameshifts in B2-3 candidate (Rpi-chc1.1) leads to Phytophthora susceptibility.

(A) Three different sgRNAs were designed to target the LRR domain of candidate B2-3. (B) A construct with Cas9 and all three sgRNAs was stably transformed into the resistant genotype CHC543-5. Transformation events were inoculated with *P. infestans* 90128 and IPO-C isolates. 48% of the transformation events became susceptible to both isolates (Table S1). (C) One resistant (#1) and three susceptible events (#21, 24, 26) were genotyped together with the untransformed plant CHC543-5. The three susceptible events contained frameshift mutations, while the tested resistant event had no mutations in the B2-3 candidate.




**Figure 3.** Rpi-chc1 allele mining.

(A) Rpi-chc1.1 allele belongs to the immune receptor family. Different motifs were found in the different CNL receptor domains. The LRR domain consists of 29 imperfect repeats. (B) Sixteen Rpi-chc1.1-like sequences were cloned from eleven different diploid *Solanum* accessions. The phylogenetic analysis of the DNA sequences led to the identification of four clades. The branch lengths represent the percentage of phylogenetic distance. In the protein alignment mismatches are highlighted in red, gaps are indicated by dashes (-), while insertions relative to Rpi-chc1.1 (anchor) are indicated by a blue bracket.




**Figure 4.** The RXLR effector PexRD12 corresponds to Avrhc1.1.

(A) The three Rpi-chc1.1 candidates were co-agroinfiltrated with the Pex effector collection in *N. benthamiana* leaves to screen for Avrhc1.1. Rpi-chc1.1 induces cell death when co-expressed with both PITG\_16233 and PITG\_16240, from the PexRD12 family. R3a and Avr3a were used as negative controls, a mix of R3a and Avr3a was used as a positive control. (B) Relative Avrhc1.1 and Avrsto1 effector expression was measured in infected plant material from a field trial with natural *P. infestans* infection (2013, Wageningen). Untransformed Desiree plants, and Desiree plants transformed with Rpi-chc1.1 (Desiree:Rpi-chc1.1), or Rpi-sto1 (Desiree:Rpi-sto1) were used in this study. Relative expression data was obtained by dividing the relative effector gene expression by the *P. infestans* elongation factor 2 gene (*ef2*) expression. Three independent samples were included in the RT-qPCR experiment. Stars represent statistical difference in a two sample T-test,  $p < 0.008$ .




**Figure 5.** Rpi-chc1 alleles show non-overlapping recognition of the PexRD12/31 effector superfamily.

(A) Twenty members of the PexRD12/31 superfamily were found in *P. infestans* isolate T30-4. In the amino acid sequence, we can distinguish a signal peptide in the N-terminus, the conserved RXLR-EER motifs in the center, and the effector domain in the C-terminus. Some PexRD12/31 family members differed at the nucleotide level but were identical at the protein level (PITG\_16245 = PITG\_16418; PITG\_16233 = PITG\_16240; PITG\_20934 = PITG\_20936; PITG\_16409 = PITG\_16424). The phylogenetic analysis of the complete protein sequences led to the identification of five clades. This analysis was performed in MEGA X by using the Maximum Likelihood method based on the JTT matrix-based model. The tree with the highest log likelihood (-766) is shown. The bootstrapping values, which indicates the percentage of trees that had the particular branch, are shown in each branch. In the protein alignment: blue, hydrophobic residues (A, I, L, M, F, W, V and C); red, positively charged residues (K and R); magenta, negatively charged residues (E and D); green, polar residues (N, Q, S and T); pink, cysteines residue (C); orange, glycines residue (G); yellow, prolines residue (P); cyan, aromatic residues (H and Y); and white, unconserved residues or gaps. (B) Different Rpi-chc1 allelic variants were co-agroinfiltrated in *N. benthamiana* with a member from each PexRD12/31 clade. While variants from clade 1 recognize both PexRD12 A1 and A2 clades, Rpi-chc1 variants from clade 2 recognize PexRD31 B and C. Receptors from clade 3 and 4 do not recognise any PexRD12/31 effector. A mix of R3a and Avr3a was used as a positive control.



**Figure 6.** Domain exchanges between Rpi-chc1.2 and rpi-tub1.3\_RH89-039-16. (A) The positions of SAPs and the corresponding protein domains are indicated on top. Rpi-chc1.2 and rpi-tub1.3\_RH89-039-16, are represented as light blue and yellow bars, respectively. Below, the domain exchanges are shown. The chimeric constructs were co-agroinfiltrated with the PexRD12/31 effectors in *N. benthamiana* leaves. After 4 days, the HR that was visible and recorded. Experiments were repeated three times with 12 inoculation sites each time. A representative leaf for the response of each chimeric construct is shown in the right panel. GUS was used as a negative control. It is concluded that the exchange of the complete LRR domain led to recognition of PexRD31. With the final construct, RH::C2\_14-19, the exchange of only nine amino acids led to the activation of the rpi-tub1.3\_RH89-036-16 protein. (B) Seven new modified receptors were derived from RH::C2\_14-19 in order to pinpoint the amino acids involved in the Rpi-chc1.2 recognition specificity. SAPs present in Rpi-chc1.2 are highlighted in blue font. Most of the SAPs are located in the solvent exposed xxLxLxxxx motif of the LRR 16-19. The chimeric constructs were co-agroinfiltrated with the PexRD12/31 effectors in *N. benthamiana* leaves. A representative leaf for the response of each chimeric construct is shown in the right panel. Experiments were repeated three times with 12 inoculation sites each time. GUS was used as a negative control. The modification of the tested residues of the Rpi-chc1.2 solvent exposed specific amino acids (blue) for the corresponding amino acid present in rpi-tub1.3\_RH89-039-16 (yellow), led to the partial or complete loss of effector recognition.



**Figure 7.** The effector recognition specificity could be exchanged between Rpi-chc1.1 and Rpi-chc1.2.

The alignment of Rpi-chc1.1 and Rpi-chc1.2 shows that all the 41 amino acids polymorphisms (red bars) are located in the LRR domain. The chimeric constructs were co-agroinfiltrated with the PexRD12/31 effectors in *N. benthamiana* leaves. A representative leaf for the response of each chimeric construct is shown in the right panel. Experiments were repeated three times with 12 inoculation sites each time. GUS was used as a negative control. In the construct C1::C2\_14-23, we could see that the LRR 16-19 reappear again as determining for the PexRD31 recognition. The required domain exchanges of the Rpi-chc1.1 LRR are more complex and encompasses almost the complete LRR.