

**Oxidation of iodide to iodate by cultures of marine ammonia-oxidising bacteria**

Claire Hughes<sup>1\*</sup>, Eleanor Barton<sup>1\*</sup>, Helmke Hepach<sup>1</sup>, Rosie Chance<sup>2</sup>, Matt Pickering<sup>1</sup>, Karen Hogg<sup>3</sup>, Andreas Pommerening-Röser<sup>4</sup>, Martin R. Wadley<sup>5</sup>, David P. Stevens<sup>5</sup> and Tim D. Jickells<sup>6</sup>

<sup>1</sup> Department of Environment and Geography, University of York, Wentworth Way, Heslington, York, YO10 5NG, UK

<sup>2</sup> Wolfson Atmospheric Chemistry Laboratory, Department of Chemistry, University of York, Heslington, York, YO10 5DD, UK

<sup>3</sup> Bioscience Technology Facility, Department of Biology, University of York, Wentworth Way, York, YO10 5DD, UK

<sup>4</sup> University of Hamburg, Mikrobiologie & Biotechnologie, Ohnhorststr. 18, D-22609 Hamburg, Germany

<sup>5</sup> School of Mathematics, University of East Anglia, Norwich Research Park, Norwich, NR4 7TJ, UK

<sup>6</sup> School of Environmental Sciences, University of East Anglia, Norwich Research Park, Norwich, NR4 7TJ, UK

\*authors contributed equally to the manuscript

Corresponding author: Rosie Chance (rosie.chance@york.ac.uk)

**Highlights:**

- Oxidation of iodide to iodate by marine nitrifying bacteria demonstrated for first time
- Laboratory cultures of ammonium oxidising bacteria produced iodate from iodide substrate
- Nitrification used to parameterise iodide sink in global marine iodine cycling model
- Changes in nitrification may increase sea surface iodide, impacting atmospheric chemistry

## Abstract


Reaction with iodide ( $I^-$ ) at the sea surface is an important sink for atmospheric ozone, and causes sea-air emission of reactive iodine which in turn drives further ozone destruction. To incorporate this process into chemical transport models, improved understanding of the factors controlling marine iodine speciation, and especially sea-surface iodide concentrations, is needed. The oxidation of  $I^-$  to iodate ( $IO_3^-$ ) is the main sink for oceanic  $I^-$ , but the mechanism for this remains unknown. We demonstrate for the first time that marine nitrifying bacteria mediate  $I^-$  oxidation to  $IO_3^-$ . A significant increase in  $IO_3^-$  concentrations compared to media-only controls was observed in cultures of the ammonia-oxidising bacteria *Nitrosomonas* sp. (Nm51) and *Nitrosococcus oceanus* (Nc10) supplied with 9-10 mM  $I^-$ , indicating  $I^-$  oxidation to  $IO_3^-$ . Cell-normalised production rates were  $15.69 (\pm 4.71)$  fmol  $IO_3^-$  cell $^{-1}$  d $^{-1}$  for *Nitrosomonas* sp., and  $11.96 (\pm 6.96)$  fmol  $IO_3^-$  cell $^{-1}$  d $^{-1}$  for *Nitrosococcus oceanus*, and molar ratios of iodate-to-nitrite production were  $9.2 \pm 4.1$  and  $1.88 \pm 0.91$  respectively. Preliminary experiments on nitrite-oxidising bacteria showed no evidence of  $I^-$  to  $IO_3^-$  oxidation. If the link between ammonia and  $I^-$  oxidation observed here is representative, our ocean iodine cycling model predicts that future changes in marine nitrification could alter global sea surface  $I^-$  fields with potential implications for atmospheric chemistry and air quality.

## Introduction

Iodine plays an important role in catalytic ozone destruction and new particle formation in the troposphere, thereby impacting the oxidative capacity of the atmosphere (Sherwen *et al.*, 2016) and the Earth's radiation balance (O'Dowd *et al.*, 2002). Sea-to-air iodine transfer is known to be the main source of iodine to the atmosphere (Carpenter, 2003; Sherwen *et al.*, 2016). Reactive inorganic iodine ( $I_2$ , HOI) emissions resulting from the reaction of gas-phase ozone with sea surface iodide ( $I^-$ ) is now thought to be the dominant mechanism mediating sea-air iodine emissions (Carpenter *et al.*, 2013). The strength of the surface reactive iodine flux is related to sea surface  $I^-$  concentrations (Carpenter *et al.*, 2013) so knowledge of ocean  $I^-$  distributions is required in order to estimate the significance of this process. Furthermore, a detailed understanding of the processes controlling inorganic iodine speciation is needed to allow us to develop predictive capacity regarding sea surface  $I^-$ , ozone-deposition rates and sea-air emission of reactive iodine.

Total inorganic iodine is found at 400-500 nM in seawater and predominantly exists as iodate ( $IO_3^-$ ) and  $I^-$  (Chance *et al.*, 2014) with inter-conversion between these two species alongside physical mixing being the main causes of spatial and temporal variability in sea surface  $I^-$ . Iodate is the thermodynamically stable form and the dominant form in the deep ocean. The existence of relatively higher levels of  $I^-$  in the euphotic zone (reviewed by Chance *et al.*, 2014) has led to the suggestion that  $IO_3^-$  reduction to  $I^-$  is linked to primary productivity. This theory has been supported by observations of  $I^-$  production in cultures of a wide range of marine phytoplankton (e.g. Chance *et al.*, 2007; Bluhm *et al.*, 2010; Hepach *et al.*, 2020) and some field studies (Chance *et al.*, 2010). The mechanism of biogenic iodate reduction to iodide is not yet known, but may be related to senescence processes (Bluhm *et al.*, 2010; Hepach *et al.*, 2020; Carrano *et al.*, 2020). Reduction of  $IO_3^-$  to  $I^-$  by phytoplankton nitrate reductase enzymes (Hung *et al.*, 2005), or macroalgal cell surface reductases (Carrano *et al.*, 2020), has also been suggested but neither has been confirmed as a significant route of conversion.

Oxidation of  $\Gamma^-$  back to  $\text{IO}_3^-$  is the dominant sink for  $\Gamma^-$ , but is a relatively slow reaction with rate estimates ranging from  $\sim 4$  to  $670 \text{ nM yr}^{-1}$  (Chance *et al.*, 2014; Hardisty *et al.*, 2020). The rates and processes involved in  $\Gamma^-$  to  $\text{IO}_3^-$  oxidation are associated with large uncertainty (Truesdale *et al.*, 2001; Amachi *et al.*, 2008), and the mechanisms involved remain undefined. This uncertainty has been suggested to be one of the factors hindering the development of mathematical models of iodine transformations in the global oceans (Truesdale *et al.*, 2001). Abiotic oxidation of  $\Gamma^-$  back to  $\text{IO}_3^-$  in the ocean (e.g. by oxygen, hydroxyl radicals, hydrogen peroxide and ozone) is thought to occur so slowly as to be insignificant (e.g. Wong, 1991), and so  $\Gamma^-$  oxidation to  $\text{IO}_3^-$  is also thought to be associated with marine microbiological activity.  $\Gamma^-$  oxidation to  $\text{I}_2$  has been observed in bacterial isolates obtained from a range of environments including seawater aquaria (Gozlan *et al.*, 1968), natural gas brines (Iino *et al.*, 2016) and seawater/marine mud (Fuse *et al.*, 2003). Additionally, based on field observations, a number of studies (Truesdale *et al.*, 2001; Žic *et al.*, 2013) have proposed that  $\Gamma^-$  oxidation to  $\text{IO}_3^-$  is linked to nitrification in marine systems. Nitrification is the two-stage biological transformation of ammonia ( $\text{NH}_3$ ) to nitrate ( $\text{NO}_3^-$ ) (Equations 1 and 2; Koops & Pommerening-Röser, 2001) mediated by chemoautotrophic ammonia-oxidising bacteria (AOB), and nitrite-oxidising bacteria (NOB). Previously thought to only occur outside of the euphotic zone, nitrification is now known to occur throughout the oceanic water-column (reviewed by Yool *et al.*, 2007).



A link between  $\Gamma^-$  oxidation/  $\text{IO}_3^-$  production and nitrification is yet to be confirmed but, if established, would suggest that  $\Gamma^-$  oxidation to  $\text{IO}_3^-$  is widespread throughout the world's oceans (Yool *et al.*, 2007).

The primary aim of this study was to establish whether  $\text{I}^-$  oxidation to  $\text{IO}_3^-$  is associated with marine nitrification. Our objectives were to determine if  $\text{IO}_3^-$  production occurs in cultures of marine ammonia- and nitrite-oxidising bacteria supplied with  $\text{I}^-$ , determine the relative rates of  $\text{IO}_3^-$  production and nitrification and explore the possible implications of the findings.

## Methods

### *Cultures*

Stock bacterial cultures were taken from the existing culture collections of the authors. Two marine AOB cultures, *Nitrosomonas* sp. Nm51 (C-15) and *Nitrococcus oceanus* Nc10 (C-107, ATCC 19707) were investigated for  $\text{IO}_3^-$  production in the presence of  $\text{I}^-$  as the only iodine source. These strains were originally isolated from seawater in the south Pacific and the north Atlantic respectively (Watson and Mandel, 1971). Cultures were grown in the dark in a water bath at 25 °C in autoclaved ESAW artificial seawater mixture (Berges *et al.*, 2001) made up using distilled water. The ESAW media was supplemented with 7-8 mM ammonium chloride and potassium phosphate. We also conducted preliminary tests on three active marine NOB: *Nitrospira marina* Nb-295 (isolated from Gulf of Maine, Watson *et al.*, 1976); *Nitrospina gracilis* 3/211 (isolated from the south Atlantic, Watson and Waterbury, 1971); *Nitrococcus mobilis* Nb-231 (ATCC 25380, isolated from Galapagos seawater, Watson and Waterbury, 1971). However we saw no evidence of  $\text{IO}_3^-$  production in any of the NOB cultures studied and these results are not discussed further. Handling of cultures was done at all times in a biosafety cabinet using sterile equipment.

### *Experimental Set Up*

For the AOB experiments triplicate cultures were incubated alongside triplicate media-only controls for periods of 8-12 days. The experiments were kept as short as possible to avoid significant changes in pH in the bulk media which would impact inorganic iodine speciation. Hence experiments were only run until an increase in nitrite across two time-points was observed. Samples were taken at regular intervals of between 1 to 6 days for pH measurement, cell counts and determination of  $\text{NO}_2^-$ ,  $\text{IO}_3^-$ ,  $\text{I}^-$  and  $\text{NH}_4^+/\text{NH}_3$  concentrations. In all cases,  $\text{I}^-$  (Aristar) was added to be at similar concentrations with the  $\text{NH}_4^+$  required in the growth media. The levels of  $\text{I}^-$  are much higher than those encountered in the oceans (global ocean median=77 nM  $\text{I}^-$  [interquartile range 28-140 nM], Chance *et al.*, 2014) but were chosen to be similar to the levels of  $\text{NH}_4^+$ . This is because in the marine environment nitrifiers would be exposed to similar ratio of  $\text{NH}_4^+$  and  $\text{I}^-$ . For example, Rees *et al.* (2006) show that  $\text{NH}_4^+/\text{NH}_3$  occurs at concentrations ranging from 60-300 nM in the Atlantic between 60°N to 50°S.

### ***pH***

A spectrophotometric method using a Lambda 25 UV/Vis spectrophotometer (Perkin-Elmer) and m-cresol purple dye (Dickson *et al.*, 2007) with measurements at 730, 578 and 434 nm was used to determine pH in the cultures and media-only controls. Salinity, needed for the pH calculation, was calculated from conductivity measured using a calibrated Hanna Instruments hand-held probe.

### ***Cell counts***

Immediately after sampling, 4 mL of the culture was fixed with 15  $\mu\text{L}$  of 50% glutaraldehyde (Alfa Aesar), flash frozen in liquid nitrogen and placed in a -80 °C freezer for later determination of cell density. Cell counts were made using a Beckman Coulter Cytoflex S flow cytometer (flow rate of 10  $\mu\text{L min}^{-1}$ ) within 2 months of collection. DAPI (Sigma; 2  $\mu\text{g mL}^{-1}$ ) stained samples were excited by a

laser at 405 nm and the emitted fluorescence detected using an avalanche photodiode detector with a reflective band pass filter 450/45. The flow cytometer thresholds were set using the 405 nm laser side scatter and the DAPI fluorescence signals.

### ***Nitrite concentration***

$\text{NO}_2^-$  was measured in 0.45  $\mu\text{m}$  (Millex) filtered samples using a spectrophotometric method (Lambda 25 UV/Vis spectrophotometer, Perkin-Elmer) developed by Norwitz & Keliher (1984). The method involves diazotizing nitrite with sulfanilamide (Fisher, analytical reagent grade) and coupling with N-1-naphthylethylenediamine dihydrochloride (Fisher, analytical reagent grade) to form a coloured azo dye which is measured spectrophotometrically at 540 nm. The method was calibrated using  $\text{NaNO}_2$  standards (Fisher, analytical reagent grade) prepared in the ESAW-based media.


### ***Iodate Concentration***

$\text{IO}_3^-$  concentrations were measured in 0.45  $\mu\text{m}$  (Millex) filtered samples using a manual version of the spectrophotometric (Lambda 25 UV/Vis spectrophotometer) method detailed in Truesdale & Spencer, 1974 and Jickells *et al.*, 1988. Absorbance was measured at 350 nm. Strictly, this method determines all oxidised (0 to +5 oxidation state) forms of inorganic iodine, but in seawater derived media this is predominantly  $\text{IO}_3^-$ , and so will be referred to as  $\text{IO}_3^-$  iodate hereafter. The method was calibrated using potassium iodate (Aristar) standard solutions made up in ESAW.

Some validation and modification to the method was required due to the nature of our experimental set-up. Chapman & Liss (1977) show that  $\text{NO}_2^-$  can interfere with spectrophotometric  $\text{IO}_3^-$  measurements (using sulfamic acid) at ambient seawater concentrations with a 15% error. Clearly significant interference would be an issue for our experiments where  $\text{NO}_2^-$  was being produced so we ran tests. We found that the presence of  $\text{NO}_2^-$  up to 10  $\mu\text{M}$  had negligible impact on  $\text{IO}_3^-$



measurements (between 0.1-50  $\mu\text{M}$ ). We did however identify that the high starting concentration of  $\text{I}^-$  ( $\sim 10 \mu\text{M}$ ) in the culture media was problematic. The iodate analysis method comprises two steps: the first involves an initial absorbance reading after the addition of sulfamic acid; the second involves the addition of excess  $\text{I}^-$ . Under acidic conditions  $\text{I}^-$  reacts with  $\text{IO}_3^-$  to form  $\text{I}_2$  (equation 3a) which reacts with excess  $\text{I}^-$  to form the coloured ion  $\text{I}_3^-$  (equation 3b) that can be measured spectrophotometrically.



The difference between the first and second absorbance readings is then used to calibrate the method. In the case of our experiments the media already contained excess  $\text{I}^-$  so the formation of  $\text{I}_2$  and  $\text{I}_3^-$  was initiated as soon as the acid was added in the first step. Hence we calibrated the method based on a single absorbance reading obtained after acid and then additional  $\text{I}^-$  was added. Calibrations and standard checks revealed this approach did not have any impact on the quality of the data.

### ***Ammonium Concentration***

$\text{NH}_4^+$  concentrations were measured in 0.45  $\mu\text{m}$  (Millex) filtered samples with a Seal Analytical Autoanalyser 3 according to method G-109-93 rev. 10 (Seal Analytical) using sodium salicylate, dichloro-isocyanuric acid and citrate buffer. The method was calibrated using standards ranging from 0-2 mg/L prepared from dilutions of a 1000 mg/L ammonium standard solution (Merck).

### ***Iodide Concentration***

$\text{I}^-$  concentrations were determined using a Dionex ICS-2000 ion chromatograph equipped with an EGC III KOH elugen cartridge, AG18 (2 x 50 mm) guard column, AS18 (2 x 250 mm) analytical

column, ASRS 300 (2 mm) suppressor, DS6 heated conductivity cell and AS40 autosampler.

Samples were diluted 100-fold with 18 M $\Omega$  deionised water for analysis and 5  $\mu$ L was injected onto the ion chromatograph. Aqueous potassium hydroxide was used as the eluent at a flow rate of 0.25 mL min<sup>-1</sup> with a gradient program starting from an initial concentration of 2 mM hydroxide (hold 1 min) to 20 mM at 18 min then to 41 mM at 19 min (hold 2 min) before returning to 2 mM. The I<sup>-</sup> retention time was 19 min. The instrument was calibrated with matrix-matched standards ranging from 0-100 nM (I<sup>-</sup>), prepared from dilutions of a 1000 mg/L iodide standard solution (Fisher Scientific) with 18 M $\Omega$  deionised water and containing a final concentration of 1% ESAW.

### **Data Analysis**

As in Guerrero and Jones (1996), the NH<sub>4</sub><sup>+</sup> oxidation rate is defined here as the rate of increase in NO<sub>2</sub><sup>-</sup>. Similarly, we define the rate of I<sup>-</sup> oxidation as the rate of increase in IO<sub>3</sub><sup>-</sup>. This is appropriate as no other iodine species were supplied to the cultures and conversion between I<sup>-</sup> and IO<sub>3</sub><sup>-</sup> is known to be the main cause of variability in inorganic iodine speciation (Bluhm *et al.*, 2010; Chance *et al.*, 2014). Average NO<sub>2</sub><sup>-</sup> and IO<sub>3</sub><sup>-</sup> production rates were calculated for each replicate culture using Equation 4.


$$\text{Production Rate (nM day}^{-1}\text{)} = \frac{(C_{\text{end}} - C_0)}{t} \quad (4)$$

where C<sub>0</sub> and C<sub>end</sub> are the NO<sub>2</sub><sup>-</sup> or IO<sub>3</sub><sup>-</sup> concentrations observed at the start and end of the experiment and t is the experimental duration in days. Cell-normalised rates were calculated by dividing these rates by the final cell density observed in each AOB culture and are hence likely to be minimum values.

## Results

### Cell counts and pH


Increases in cell density were observed in all replicates of *Nitrosomonas* sp. and *Nitrosococcus oceani* between the start and end of the experiment indicating growth (Figure 1). Average initial cell density in the *Nitrosomonas* sp. cultures was 21,767 ( $\pm 4,046$ ) cells mL<sup>-1</sup> and this increased to 150,983 ( $\pm 7,585$ ) cells mL<sup>-1</sup> by the end of the experiment (8 days). For *Nitrosococcus oceani* start and end (12 days) cell densities were 16,947 ( $\pm 3,098$ ) and 71,430 ( $\pm 9,062$ ) cells mL<sup>-1</sup>, respectively. Average pH levels in the culture experiments calculated from measurements at each time point (data not shown) were 7.69 ( $\pm 0.07$ ) for *Nitrosomonas* sp. and 7.41 ( $\pm 0.12$ ) for *Nitrosococcus* sp. These pH levels are consistent with those found in the media-only controls (7.64 $\pm$ 0.07 for *Nitrosomonas* sp; 7.64 $\pm$ 0.15 for *Nitrosococcus oceani*).



**Figure 1.** Average cell number in the *Nitrosomonas* sp. (grey bars) and *Nitrosococcus oceani* (white bars) cultures used in this study at the start (T<sub>0</sub>) and end (T<sub>end</sub>; 8 days for *Nitrosomonas* sp. and 12 days for *Nitrosococcus oceani*) of each experiment. Error bars are standard deviations from three replicate cultures.

### *Iodine and nitrogen speciation*

Figure 2 shows that significant increases in the concentrations of  $\text{IO}_3^-$  (compared to media-only controls) were observed alongside  $\text{NO}_2^-$  production in both AOB cultures studied. In *Nitrosomonas* sp. (Figure 2ai and 2bi) there was a steady increase in  $\text{IO}_3^-$  concentrations throughout the experiment reaching a maximum of 19,921 ( $\pm 4,754$ ) nM by the end of the experiment (day 8). In contrast  $\text{NO}_2^-$  concentrations reached a maximum of 2,360 ( $\pm 386$ ) nM by day 6 and remained at around that level until the end of the experiment. In *Nitrosococcus oceani* (Figure 2aii and 2bii)  $\text{IO}_3^-$  concentrations increased rapidly during the initial stages of the experiment reaching 23,943 ( $\pm 8,568$ ) nM by day 6.  $\text{IO}_3^-$  concentrations at the end of the experiment (day 12) were 16,365 ( $\pm 7,603$ ) nM.  $\text{NO}_2^-$  concentrations increased gradually throughout the experiment reaching 5,547 ( $\pm 1,251$ ) nM by day 12. There was larger variability in  $\text{IO}_3^-$  concentrations between replicates for *Nitrosococcus oceani* but despite this a clear increase in all replicates was observed.




**Figure 2.** Changes in iodate (a) and nitrite (b) concentrations in cultures (closed symbols) and media-only controls (open symbols) for two cultures of ammonia-oxidising bacteria: i) *Nitrosomonas* sp.; and, ii) *Nitrosococcus oceanii* supplied with 9-10 mM iodide and 7-8 mM  $\text{NH}_4^+$ . Error bars show the standard deviation of three replicate cultures.

Average production rates of  $\text{IO}_3^-$  and  $\text{NO}_2^-$  are presented in Table 1. In *Nitrosomonas* sp. average rates ( $\pm$ standard deviation) were 2,348 ( $\pm$ 593) nM  $\text{IO}_3^-$  day $^{-1}$  and 298 ( $\pm$ 141) nM  $\text{NO}_2^-$  day $^{-1}$ . In *Nitrosococcus oceanii* average rates were 897 ( $\pm$ 640) nM  $\text{IO}_3^-$  day $^{-1}$  and 445 ( $\pm$ 99) nM  $\text{NO}_2^-$  day $^{-1}$ . Minimum cell-normalised rates (based on the final cell density observed in each culture) were 15.69 ( $\pm$ 4.71) fmol  $\text{IO}_3^-$  cell $^{-1}$  day $^{-1}$  and 1.96 ( $\pm$ 0.88) fmol  $\text{NO}_2^-$  cell $^{-1}$  day $^{-1}$  for *Nitrosomonas* sp., and 11.96 ( $\pm$ 6.96) fmol  $\text{IO}_3^-$  cell $^{-1}$  day $^{-1}$  and 6.19 ( $\pm$ 0.56) fmol  $\text{NO}_2^-$  cell $^{-1}$  day $^{-1}$  for *Nitrosococcus oceanii*. Molar ratios of iodate-to-nitrite production were 9.2 $\pm$ 4.0 for *Nitrosomonas* sp. and 1.88 $\pm$ 0.91 for *Nitrosococcus oceanii*.

**Table 1.** Nitrite and iodate production rates ( $\pm$  standard deviations) observed in cultures of the ammonia-oxidising bacteria *Nitrosomonas* sp. and *Nitrosococcus oceanus*. Cell-normalised values are a minimum as they are calculated using maximum cell densities.

Culture	Nitrite		Iodate	
	nM day <sup>-1</sup>	fmol cell <sup>-1</sup> day <sup>-1</sup>	nM day <sup>-1</sup>	fmol cell <sup>-1</sup> day <sup>-1</sup>
<i>Nitrosomonas</i> sp.	298 ( $\pm$ 141)	1.96 ( $\pm$ 0.88)	2,348 ( $\pm$ 593)	15.69 ( $\pm$ 4.71)
<i>Nitrosococcus oceanus</i>	445 ( $\pm$ 99)	6.19 ( $\pm$ 0.56)	897 ( $\pm$ 640)	11.96 ( $\pm$ 6.96)

Figure 3 shows that, within error, a decline in  $\Gamma$  or  $\text{NH}_4^+$  concentrations was not observed during either of the AOB experiments. Average starting  $\Gamma$  or  $\text{NH}_4^+$  concentrations in *Nitrosomonas* sp. were 9.8 ( $\pm$ 0.2) mM and 7.6 ( $\pm$ 0.1) mM respectively. At the end of the experiment these values were 10.2 ( $\pm$ 0.3) mM  $\Gamma$  and 7.7 ( $\pm$ 0.1) mM  $\text{NH}_4^+$ . For *Nitrosococcus oceanus* the start and end concentrations were 9.8 ( $\pm$ 0.3) and 9.4 ( $\pm$ 0.1) mM for  $\Gamma$  and 7.8 ( $\pm$ 0.2) and 7.7 ( $\pm$ 0.1) mM for  $\text{NH}_4^+$ . This result was expected as the average standard deviations associated with the observed concentrations of  $\Gamma$  or  $\text{NH}_4^+$  (i.e. 0.1 to 0.3 mM) are at least an order of magnitude higher than the maximum levels of  $\text{IO}_3^-$  and  $\text{NO}_2^-$  observed in the culture experiments, i.e. very little of the initial stock of  $\text{NO}_2^-$  or  $\text{NH}_4^+$  was oxidised during the experiments.



**Figure 3.** Start and end concentrations of a) iodide and b) ammonia in cultures of *Nitrosomonas* sp. (grey bars) and *Nitrosococcus oceani* (white bars). Error bars show the standard deviation of three replicate cultures.

## Discussion

### *Iodate production by ammonia-oxidising bacteria*

Our results confirm that  $\text{IO}_3^-$  production occurs in cultures of the ammonia-oxidising bacteria *Nitrosomonas* sp. and *Nitrosococcus oceani* supplied with  $\text{I}^-$ , but not in cultures of nitrite oxidising bacteria. Coincident increases in  $\text{NO}_2^-$  (Figure 2) show that both cultures were actively oxidising ammonia throughout the experiments at rates of  $1.96 \pm 0.68 \text{ fmol NO}_2^- \text{ cell}^{-1} \text{ day}^{-1}$  for *Nitrosomonas* sp. and  $6.19 \pm 0.56 \text{ fmol NO}_2^- \text{ cell}^{-1} \text{ day}^{-1}$  for *Nitrosococcus oceani*. Whilst these cell-normalised oxidation rates are of the same order as those reported in the literature (e.g.  $6\text{--}20 \text{ fmol NO}_2^- \text{ cell}^{-1} \text{ day}^{-1}$ ; Ward *et al.*, 1987; 1989) they are at the lower end. This is consistent with the approach taken here to calculate the rates by normalising to the final (highest) cell densities. It is also worth noting that the cultures were at an early stage of growth and had relatively low cell densities during the experiment. This was done to avoid significant changes in pH in the bulk media which would impact inorganic iodine speciation (Section 3.2). The observation of an increase in  $\text{IO}_3^-$  concentrations alongside active biological ammonia oxidation supports previous studies (e.g. Truesdale *et al.*, 2001; Zic *et al.*, 2013) which have shown that high aqueous concentrations of  $\text{IO}_3^-$  are found in regions of enhanced nitrification, and provides the first direct confirmation of a biological basis for at least one mechanism of iodide oxidation.

Whilst we did not set out to establish the mechanism for  $\text{I}^-$  to  $\text{IO}_3^-$  oxidation by marine nitrifiers, some speculations can be made. As  $\text{I}^-$  oxidation to  $\text{IO}_3^-$  requires the transfer of six electrons, it may

occur in a series of one- or two- electron transfer steps. Initially,  $\Gamma^-$  may be oxidised to molecular iodine ( $\Gamma^- \rightarrow \text{I}_2$ ), a reaction which is thermodynamically unfavourable at the pH of seawater (Luther *et al.*, 1995). Further oxidation to  $\text{IO}_3^-$  by disproportionation ( $\text{I}_2 \rightarrow \text{HOI} \rightarrow \text{IO}_3^-$ ) can occur spontaneously, but in seawater is subject to competition with reduction of  $\text{I}_2$  by organic matter (Truesdale & Moore, 1992; Truesdale *et al.*, 1995). It is not known whether the ammonia-oxidisers mediate just the first stage of  $\Gamma^-$  oxidation, with the observed  $\text{IO}_3^-$  production due to subsequent spontaneous reactions in the culture media, or if they are involved in driving the complete conversion of  $\Gamma^-$  to  $\text{IO}_3^-$ . However, bacteria which just oxidise  $\Gamma^-$  to  $\text{I}_2$  have been isolated from seawater aquaria (Gozlan, 1968),  $\Gamma^-$ -rich natural gas brine waters (Amachi *et al.*, 2005) and marine environmental samples (Fuse *et al.*, 2003; Amachi *et al.*, 2005).

The observed  $\text{IO}_3^-$  production is either linked to the nitrification process itself or associated with other metabolic activities of the AOB studied. Truesdale *et al.* (2001) has proposed that  $\Gamma^-$  oxidation to  $\text{IO}_3^-$  would be energetically advantageous for chemoautotrophic AOB. In that case the key enzymes used to obtain energy during the oxidation of  $\text{NH}_4^+$  to  $\text{NO}_2^-$  (ammonia monooxygenase [AMO] and hydroxylamine oxidoreductase [HAO]) could also have the potential to use  $\Gamma^-$  as a substrate. The observed  $\text{IO}_3^-/\text{NO}_2^-$  molar production rates ( $9.2 \pm 4.0$  for *Nitrosomonas* sp. and  $2.3 \pm 1.1$  for *Nitrosococcus oceanus*) are intriguing. If AMO/HAO are involved, this suggests that the enzymes have higher affinities for  $\Gamma^-$  than  $\text{NH}_4^+/\text{NH}_2\text{OH}$  given the similar concentrations of  $\Gamma^-$  and  $\text{NH}_4^+$  used in the experiments. Other enzymes that have been implicated in  $\Gamma^-$  oxidation include the chloroperoxidases (Thomas & Hager, 1968) but we do not know if they occur in AOB. The exact metabolic pathway driving the observed  $\text{IO}_3^-$  production and its controls (i.e. substrate concentrations, light intensity) will need to be determined in future work. To establish if such further experimentation is warranted we need to explore whether the link between nitrification and  $\Gamma^-$  oxidation is likely to be an important part of inorganic iodine cycling in seawater.



### *Implications for inorganic iodine speciation in the oceans*

Our culture studies suggest that the molar rate of  $\Gamma^-$  oxidation ( $\text{IO}_3^-$  production) is ~2-9 times higher than that for ammonia oxidation (nitrification). Note that although ammonium and iodide concentrations were much higher in the experimental media than in the oceans, the concentration ratio of these species was comparable to that found naturally. Ammonia oxidation rates in seawater range from below detection to  $10^2 \text{ nM day}^{-1}$  (Table 2). Literature estimates of the rate of  $\Gamma^-$  oxidation in the marine environment range from ~4 to  $670 \text{ nM year}^{-1}$  or  $0.01$  to  $1.84 \text{ nM day}^{-1}$  (reviewed in Chance *et al.*, 2014). If the oxidation molar ratios observed in this study (~2-9) are representative, predicted rates of  $\Gamma^-$  oxidation are in-line (i.e. 2-9 times higher) with the lower end of observed ammonia oxidation rates (Table 2).

**Table 2.** Ammonia-oxidation rates measured in a range of ocean regions.


Study	Location	Rate ( $\text{nM day}^{-1}$ )
Newell <i>et al.</i> (2011)	Arabian Sea, Indian Ocean	undetected to 21.6
Smith <i>et al.</i> (2015)	Northeast Pacific	< 0.01 to 90
Peng <i>et al.</i> (2015)	Eastern tropical north Pacific	< 1 to 8.6
Newell <i>et al.</i> (2013)	Subtropical Atlantic, Sargasso Sea (BATS)	< 2
Lam <i>et al.</i> (2007)	Black Sea	7-24
Beman <i>et al.</i> (2012)	Gulf of California, eastern tropical north Pacific	0-348

Truesdale *et al.* (2001) derive likely  $\Gamma^-$  oxidation (or  $\text{IO}_3^-$  production) rates for the near surface Black Sea using an iodine budget and this allows us to examine the potential importance of the link between nitrification and  $\Gamma^-$  oxidation on a local scale. They predict a minimum  $\Gamma^-$  oxidation flux of  $3.89 \times 10^{-4} \text{ mol I m}^{-2} \text{ year}^{-1}$  which is an average of  $0.02 \text{ nM day}^{-1}$  at a mixed-layer depth (MLD) of 50 m or  $0.11 \text{ nM day}^{-1}$  at an MLD of 10 m. Lam *et al.* (2007) report an AOB abundance of  $\leq 1,400$  cells  $\text{mL}^{-1}$  in the Black Sea. If we apply a cell density of 1,400 AOB cells  $\text{mL}^{-1}$  to the average cell-normalised rates of  $\text{IO}_3^-$  production observed in this study (Table 1) we derive  $\Gamma^-$  oxidation rates of

~20 nM d<sup>-1</sup>. This is clearly much higher than the rates suggested in Truesdale *et al.* (2001). This discrepancy could be explained in a number of ways. Firstly, Lam *et al.* (2007) state that net nitrification only takes place within a narrow depth range of the Black Sea water column (i.e. between 71 and 81 m) and, the  $\Gamma$  oxidation values derived in Truesdale *et al.* (2001) are minimum values. It is also possible that the AOB studied here have a higher capacity for  $\Gamma$  oxidation (per unit ammonia-oxidised) than other ammonia-oxidisers or that our culture conditions (e.g. substrate availability) promoted higher  $\Gamma$  oxidation rates than would be observed in marine systems. For example, ammonia-oxidising Archaea (AOA), which can outnumber known bacterial ammonia oxidisers by orders of magnitudes in environments such as the marine water-column (reviewed by Schleper & Nicol, 2010), may have a very different capacity for  $\Gamma$  oxidation compared to the AOB studied here. Further studies are needed to establish the relationship between ammonia- and  $\Gamma$  oxidation in the marine environment.

#### ***Potential implications for future oceanic inorganic iodine distributions***

Environmental factors which are known to be currently undergoing change in the oceans (e.g. oxygen, light, pH, temperature) have all been found to impact rates and patterns of marine nitrification (reviewed by Pajares and Ramos, 2019). Whilst there remains some uncertainty about the future magnitude and, in some cases, sign of the response, some of the expected future changes in marine nitrification are large. For example, whilst some studies have seen no impact on specific marine nitrifiers (e.g. Qin *et al.*, 2014), Beman *et al.* (2011) suggest that expected rates of acidification could cause a decline in ammonia oxidation by up to 44% within the next few decades. It is hence worth exploring how possible future changes in marine nitrification could impact ocean iodine cycling.



**Figure 4.** Modelled changes in surface  $\text{I}^-$  concentration (nM) resulting from a) +10%, b) -10%, changes in the rates of nitrification. Negative percent values indicate a decline in the rate of nitrification and *vice-versa*. Negative values on the scale bar indicate a decrease in  $\text{I}^-$  concentrations and *vice versa*.

In order to explore the possible impact of future changes in marine nitrification rates on sea surface iodine fields we used the ocean cycling model described in Wadley et al. (2020). Within the model iodide production is driven by primary productivity, and  $\text{I}^-$  oxidation to  $\text{IO}_3^-$  linked to nitrification in the mixed layer. Nitrogen fluxes and the spatial distribution of mixed layer ammonia oxidation are derived from a global biogeochemical cycling model (Yool et al., 2007).  $\text{I}^-$  is oxidised to  $\text{IO}_3^-$  in association with the ammonia oxidation, with the same I:N:C ratio as associated with iodide production (Truesdale et al., 2001; Long et al., 2015). The model does not use any of the rates derived in the current study as these are based on results from only 2 AOB species cultured at high substrate concentrations. Model outputs (Figure 4) show that even with small (+/- 10%) changes in ammonia oxidation there is a clear alteration to sea surface  $\text{I}^-$  fields. Sea surface  $\text{I}^-$  concentrations increase as ammonium oxidation rates decrease and *vice-versa*. For example, the ocean cycling model suggests there could be an average global increase of 0.13 nM  $\text{I}^-$  per 1% decrease in nitrification. The outputs suggest that the change in the iodine fields is spatially variable and will increase as the perturbation to ammonia oxidation increases. For example, at the 44% decline in

nitrification predicted by Beman *et al.* (2011) the model predicts there will be a 25% increase (+30 nM) in sea surface  $\Gamma^-$  in the sub-tropical gyres. Carpenter *et al.* (2013) show that  $\text{I}_2$  emissions due to ozone deposition increase near linearly with  $\Gamma^-$  concentration. Hence, the predicted changes to sea surface  $\Gamma^-$  fields under future ocean acidification could have a major impact on ozone deposition to the sea surface, atmospheric chemistry and resulting sea-air iodine emissions.

### 5.3. Conclusions

This study has shown that  $\Gamma^-$  oxidation to  $\text{IO}_3^-$  occurs in cultures of ammonia oxidising (nitrifying) bacteria, but not nitrite oxidising bacteria. Our calculations suggest that  $\Gamma^-$  oxidation by AOB could be an important control on inorganic iodine speciation in seawater, but to confirm this further study is needed on a wider range of ammonia-oxidiser including ammonia oxidising archaea (AOA). Simulations from our iodine cycling model suggest that changes in nitrification rate, such as those predicted to occur under acidification (Beman *et al.*, 2011), could have an important impact on sea surface  $\Gamma^-$  fields. A future change in marine nitrification could alter sea surface  $\Gamma^-$  fields. In turn, this could lead to a change in ozone deposition to the sea surface and sea-air iodine emissions with potentially major implications for atmospheric chemistry and air quality.

### Acknowledgements

We thank Eva Spieck (University of Hamburg, Germany) for supplying the nitrite oxidising bacteria (NOB) cultures used in our preliminary experiments.

## Author contributions

Claire Hughes: Conceptualisation, Methodology, Formal Analysis, Investigation, Writing – Original Draft, Writing – Review & Editing, Visualisation, Supervision, Project Administration, Funding Acquisition. Eleanor Barton: Formal Analysis, Investigation, Writing – Original Draft. Helmke Hepach: Methodology, Validation, Investigation, Resources. Rosie Chance: Conceptualisation, Methodology, Resources, Data Curation, Writing – Review & Editing, Project Administration, Funding Acquisition. Matt Pickering: Methodology, Validation, Resources. Karen Hogg: Methodology, Resources. Andreas Pommerening-Röser: Methodology, Resources. Martin R. Wadley: Conceptualisation, Methodology, Software, Validation, Formal Analysis, Investigation, Writing – Original Draft, Visualisation. David J. Stevens: Conceptualisation, Methodology, Resources, Supervision, Funding Acquisition. Tim D. Jickells: Conceptualisation, Methodology, Supervision, Funding Acquisition.

**Data Statement:** The data presented in this manuscript is available from the British Oceanographic Data Centre (BODC), doi:10.5285/aaa4dab8-7c9d-0cca-e053-6c86abc00b1d

**Funding:** This work was supported by the Natural Environment Research Council [grant no. NE/N01054X/1]; and the University of York Laidlaw Scholarship [work undertaken by Eleanor Barton].

**Declaration of interest:** The authors have no financial conflicts of interest with the research in this paper.

Journal Pre-proof

## References

- Amachi, S., 2008. Microbial Contribution to Global Iodine Cycling: Volatilization, Accumulation, Reduction, Oxidation, and Sorption of Iodine. *Microbes Environ.*, 23(4): 269.10.1264/jsme2.ME08548
- Amachi, S., Muramatsu, Y., Akiyama, Y., Miyazaki, K., Yoshiki, S., Hanada, S., Kamagata, Y., Ban-nai, T., Shinoyama, H. and Fujii, T., 2005. Isolation of iodide-oxidizing bacteria from iodide-rich natural gas brines and seawaters. *Microb. Ecol.* 49(4): 547
- Beman, J.M., Chow, C.-E., King, A.L., Feng, Y., Fuhrman, J.A., Andersson, A., Bates, N.R., Popp, B.N. and Hutchins, D.A., 2011. Global declines in oceanic nitrification rates as a consequence of ocean acidification. *Proceedings of the National Academy of Sciences*, 108(1): 208.10.1073/pnas.1011053108
- Beman, J.M., Popp, B. N. & Alford, S. E., 2012. Quantification of ammonia oxidation rates and ammonia-oxidizing archaea and bacteria at high resolution in the Gulf of California and eastern tropical North Pacific Ocean. *Limnology and Oceanography*, 57, 712-726
- Bluhm, K., Croot, P., Wuttig, K. and Lochte, K., 2010. Transformation of iodate to iodide in marine phytoplankton driven by cell senescence. *Aquat. Biol.*, 11(1): 1.10.3354/ab00284
- Carpenter, L.J., 2003. Iodine in the marine boundary layer. *Chem. Rev.*, 103(12): 4953.10.1021/cr02064-65
- Carpenter, L.J., MacDonald, S.M., Shaw, M.D., Kumar, R., Saunders, R.W., Parthipan, R., Wilson, J. and Plane, J.M.C., 2013. Atmospheric iodine levels influenced by sea surface emissions of inorganic iodine. *Nature Geoscience*, 6(2): 108.10.1038/ngeo1687
- Carrano, M.W., Yarimizu, K., Gonzales, J.L., Cruz-López, R., Edwards, M.S., Tymon, T.M., Küpper, F.C. and Carrano, C.J. 2020. The influence of marine algae on iodine speciation in the coastal ocean. *Algae*, 35 (2): 167-176. 10.4490/algae.2020.35.5.25

- Chance, R., Baker, A.R., Carpenter, L. and Jickells, T.D., 2014. The distribution of iodide at the sea surface. *Environmental Science-Processes & Impacts*, 16(8): 1841.10.1039/c4em00139g
- Chance, R., Malin, G., Jickells, T. and Baker, A.R., 2007. Reduction of iodate to iodide by cold water diatom cultures. *Mar. Chem.*, 105(1-2): 169.10.1016/j.marchem.2006.06.008
- Chance, R., Weston, K., Baker, A.R., Hughes, C., Malin, G., Carpenter, L., Meredith, M.P., Clarke, A., Jickells, T.D., Mann, P. and Rossetti, H., 2010. Seasonal and interannual variation of dissolved iodine speciation at a coastal Antarctic site. *Mar. Chem.*, 118(3-4): 171.10.1016/j.marchem.2009.11.009
- Chapman, P. and Liss, P.S., 1977. Effect of Nitrite on Spectrophotometric Determination of Iodate in Seawater. *Mar. Chem.*, 5(3): 243
- Dickson, A. G., Sabine, C. L. & Christian, J. R. (Eds.), 2007. Guide to Best Practices for Ocean CO<sub>2</sub> Measurements. PICES Special Publication 5. 161 pp.
- Fuse, H., Inoue, H., Murakami, K., Takimura, C. and Yamaoka, Y., 2003. Production of free and organic iodine by *Roseovarius* spp. *FEMS Microbiol. Lett.*, 229(2): 189.10.1016/s0378-1097(03)00839-5
- Gozlan, R.S., 1968. Isolation of Iodine-Producing Bacteria from Aquaria. *Antonie Van Leeuwenhoek Journal of Microbiology*, 34(2): 226
- Hardisty DS, Horner TJ, Wankel SD, Blusztajn J, and Nielsen SG. 2020. Experimental observations of marine iodide oxidation using a novel sparge-interface MC-ICP-MS technique. *Chem Geol*, 532:119360. 10.1016/j.chemgeo.2019.119360.
- Hepach H, Hughes C, Hogg K, Collings S, and Chance R., 2020. Senescence as the main driver of iodide release from a diverse range of marine phytoplankton. *Biogeosciences*, 17:2453–71. <https://doi.org/10.5194/bg-17-2453-2020>.
- Hung, C.C., Wong, G.T.F. and Dunstan, W.M., 2005. Iodate reduction activity in nitrate reductase extracts from marine phytoplankton. *Bull. Mar. Sci.*, 76(1): 61



- Iino, T., Ohkuma, M., Kamagata, Y. and Amachi, S., 2016. *Iodidimonas muriae* gen. nov., sp. nov., an aerobic iodide-oxidizing bacterium isolated from brine of a natural gas and iodine recovery facility, and proposals of *Iodidimonadaceae* fam. nov., *Iodidimonadales* ord. nov., *Emcibacteraceae* fam. nov. and *Emcibacterales* ord. nov. Int. J. Syst. Evol. Microbiol., 66(12): 5016. <https://doi.org/10.1099/ijsem.0.001462>
- Jickells, T.D., Boyd, S.S. and Knap, A.H., 1988. Iodine cycling in the Sargasso Sea and the Bermuda Inshore waters. Mar. Chem., 24(1): 61.10.1016/0304-4203(88)90006-0
- Koops, H.-P. and Pommerening-Röser, A., 2001. Distribution and ecophysiology of the nitrifying bacteria emphasizing cultured species. FEMS Microbiol. Ecol., 37(1): 1.10.1111/j.1574-6941.2001.tb00847.x
- Lam, P., Jensen, M.M., Lavik, G., McGinnis, D.F., Müller, P., Schubert, C.J., Amann, R., Thamdrup, B. and Kuypers, M.M.M., 2007. Linking crenarchaeal and bacterial nitrification to anammox in the Black Sea. Proceedings of the National Academy of Sciences, 104(17): 7104.10.1073/pnas.0611081104
- Li, H.P., Daniel, B., Creeley, D., Grandbois, R., Zhang, S.J., Xu, C., Ho, Y.F., Schwehr, K.A., Kaplan, D.I., Santschi, P.H., Hansel, C.M. and Yeager, C.M., 2014. Superoxide Production by a Manganese-Oxidizing Bacterium Facilitates Iodide Oxidation. Appl. Environ. Microbiol., 80(9): 2695.10.1128/aem.00400-14
- Long, A., Dang, A., Xiao, H. and Yu, X., 2015. The Summer Distribution of Dissolved Inorganic Iodine along 18°N in the South China Sea. Journal of Marine Science: Research & Development, 5: 169.10.4172/2155-9910.1000169
- Newell, S.E., Babbitt, A.R., Jayakumar, A. and Ward, B.B., 2011. Ammonia oxidation rates and nitrification in the Arabian Sea. Global Biogeochemical Cycles, 25(4).10.1029/2010GB003940

- Newell, S.E., Fawcett, S.E. and Ward, B.B., 2013. Depth distribution of ammonia oxidation rates and ammonia-oxidizer community composition in the Sargasso Sea. *Limnology and Oceanography*, 58(4): 1491.10.4319/lo.2013.58.4.1491
- Norwitz, G. and Keliher, P.N., 1984. Spectrophotometric determination of nitrite with composite reagents containing sulphanilamide, sulphanilic acid or 4-nitroaniline as the diazotisable aromatic amine and N-(1-naphthyl)ethylenediamine as the coupling agent. *Analyst*, 109(10): 1281.10.1039/AN9840901281
- O'Dowd, C.D., Jimenez, J.L., Bahreini, R., Flagan, R.C., Seinfeld, J.H., Hameri, K., Pirjola, L., Kulmala, M., Jennings, S.G. and Hoffmann, T., 2002. Marine aerosol formation from biogenic iodine emissions. *Nature*, 417(6889): 632
- Pajores, S. and R. Ramos (2019) Processes and microorganisms involved in the marine nitrogen cycle: knowledge and gaps, *Frontiers in Marine Science*, 6, 10.3389/fmars.2019.00739
- Peng, X., Fuchsman, C.A., Jayakumar, A., Cleveland, S., Martens-Habbena, W., Devol, A.H. and Ward, B.B., 2015. Ammonia and nitrite oxidation in the Eastern Tropical North Pacific. *Global Biogeochemical Cycles*, 29(12): 2034.10.1002/2015GB005278
- Qin, W., Amin, S.A., Martens-Habbena, W., Walker, C.B., Urakawa, H., Devol, A.H., Ingalls, A.E., Moffett, J.W., Armbrust, F.W. and Stahl, D.A., 2014. Marine ammonia-oxidizing archaeal isolates display obligate mixotrophy and wide ecotypic variation. *Proc. Natl. Acad. Sci. U. S. A.*, 111(34): 12504.10.1073/pnas.1324115111
- Schleper, C. and Nicol, G.W., 2010. Ammonia-oxidising archaea--physiology, ecology and evolution. *Adv. Microb. Physiol.*, 57: 1.10.1016/b978-0-12-381045-8.00001-1
- Schulz, K.G., Barcelos e Ramos, J., Zeebe, R.E. and Riebesell, U., 2009. CO<sub>2</sub> perturbation experiments: similarities and differences between dissolved inorganic carbon and total alkalinity manipulations. *Biogeosciences*, 6(10): 2145.10.5194/bg-6-2145-2009

- Sherwen, T., Evans, M.J., Carpenter, L.J., Andrews, S.J., Lidster, R.T., Dix, B., Koenig, T.K., Sinreich, R., Ortega, I., Volkamer, R., Saiz-Lopez, A., Prados-Roman, C., Mahajan, A.S. and Ordóñez, C., 2016. Iodine's impact on tropospheric oxidants: a global model study in GEOS-Chem. *Atmospheric Chemistry and Physics*, 16(2): 1161.10.5194/acp-16-1161-2016
- Smith, J.M., Damashek, J., Chavez, F.P. and Francis, C.A., 2016. Factors influencing nitrification rates and the abundance and transcriptional activity of ammonia-oxidizing microorganisms in the dark northeast Pacific Ocean. *Limnology and Oceanography*, 61(2): 596.10.1002/lno.10235
- Thomas, J.A. and Hager, L.P., 1968. The peroxidation of molecular iodine to iodate by chloroperoxidase. *Biochimica Biophysica Research Communications*, 32: 770
- Truesdale, V.W. and Luther, G.W., 1995. Molecular iodine reduction by natural and model organic substances in seawater. *Aquatic Geochemistry*, 9(1): 89
- Truesdale, V.W. and Moore, R.M., 1992. Further-Studies on the Chemical-Reduction of Molecular-Iodine Added to Seawater. *Mar. Chem.*, 40(3-4): 199
- Truesdale, V.W. and Spencer, C.P., 1974. Studies on the determination of inorganic iodine in seawater. *Mar. Chem.*, 2(1): 33.[https://doi.org/10.1016/0304-4203\(74\)90004-8](https://doi.org/10.1016/0304-4203(74)90004-8)
- Truesdale, V.W., Watts, S.F. and Wendell, A.R., 2001. On the possibility of iodide oxidation in the near-surface of the Black Sea and its implications to iodine in the general ocean. *Deep-Sea Research Part I-Oceanographic Research Papers*, 48(11): 2397.10.1016/s0967-0637(01)00021-8
- Wadley, M.R., Stevens, D.P., Jickells, T.D., Hughes, C., Chance, R., Hepach, H., Tinel, L. and Carpenter, L.J., 2020. A Global Model for Iodine Speciation in the Upper Ocean. *Global Biogeochemical Cycles*, e2019GB006467.10.1029/2019GB006467

- Ward, B.B., 1987. Nitrogen transformations in the Southern California Bight. Deep Sea Research Part A. Oceanographic Research Papers, 34(5): 785.[https://doi.org/10.1016/0198-0149\(87\)90037-9](https://doi.org/10.1016/0198-0149(87)90037-9)
- Ward, B.B., Glover, H.E. and Lipschultz, F., 1989. Chemoautotrophic activity and nitrification in the oxygen minimum zone off Peru. Deep Sea Research Part A. Oceanographic Research Papers, 36(7): 1031.[https://doi.org/10.1016/0198-0149\(89\)90076-9](https://doi.org/10.1016/0198-0149(89)90076-9)
- Watson, S.W., Bock, E., Valois, F.W., Waterbury, J.B. and Schlosser, U., 1986. *Nitrospira marina* gen. nov. sp. nov.: a chemolithotrophic nitrite-oxidizing bacterium. *Arch. Microbiol.* 144: 1–7. <https://doi.org/10.1007/BF00454941>
- Watson S. W. and Mandel, M., 1971. Comparison of the morphology and deoxyribonucleic acid composition of 27 strains of nitrifying bacteria. *J. Bacteriol.*, 107 (2): 563-569
- Watson, S.W. and Waterbury, J.B., 1971. Characteristics of two marine nitrite oxidizing bacteria, *Nitrospina gracilis* nov. gen. nov. sp. and *Nitrococcus mobilis* nov. gen. nov. sp. *Archiv. Mikrobiol.* 77: 203–230 <https://doi.org/10.1007/BF00408114>
- Wong, G.T.F., 1991. The marine geochemistry of iodine. *Reviews in Aquatic Sciences*, 4(1): 45
- Yool, A., Martin, A.P., Fernandez, C. and Clark, D.R., 2007. The significance of nitrification for oceanic new production. *Nature*, 447(7147): 999.[10.1038/nature05885](https://doi.org/10.1038/nature05885)
- Zic, V., Caric, M. and Ciglenecki, I., 2013. The impact of natural water column mixing on iodine and nutrient speciation in a eutrophic anchialine pond (Rogoznica Lake, Croatia). *Estuarine Coastal and Shelf Science*, 133: 260.[10.1016/j.ecss.2013.09.008](https://doi.org/10.1016/j.ecss.2013.09.008)

## Figure Captions

**Figure 1.** Average cell number in the *Nitrosomonas* sp. (grey bars) and *Nitrosococcus oceani* (white bars) cultures used in this study at the start ( $T_0$ ) and end ( $T_{end}$ ; 8 days for *Nitrosomonas* sp. and 12 days for *Nitrosococcus oceani*) of each experiment. Error bars are standard deviations from three replicate cultures.

**Figure 2.** Changes in iodate (a) and nitrite (b) concentrations in cultures (closed symbols) and media-only controls (open symbols) for two cultures of ammonia-oxidising bacteria: i) *Nitrosomonas* sp.; and, ii) *Nitrosococcus oceani* supplied with 9-10 mM iodide and 7-8 mM  $\text{NH}_4^+$ . Error bars show the standard deviation of three replicate cultures.

**Figure 3.** Start and end concentrations of a) iodide and b) ammonia in cultures of *Nitrosomonas* sp. (grey bars) and *Nitrosococcus oceani* (white bars). Error bars show the standard deviation of three replicate cultures.

**Figure 4.** Modelled changes in surface  $\Gamma$  concentration (nM) resulting from a) +10%, b) -10%, changes in the rates of nitrification. Negative percent values indicate a decline in the rate of nitrification and *vice-versa*. Negative values on the scale bar indicate a decrease in  $\Gamma$  concentrations and *vice versa*.

**Highlights:**

- Oxidation of iodide to iodate by marine nitrifying bacteria demonstrated for first time
- Laboratory cultures of ammonium oxidising bacteria produced iodate from iodide substrate
- Nitrification used to parameterise iodide sink in global marine iodine cycling model
- Changes in nitrification may increase sea surface iodide, impacting atmospheric chemistry

Journal Pre-proof






Figure 1


a,i *Nitrosomonas* sp.,  $\text{IO}_3^-$



a,ii *Nitrosococcus oceanus*,  $\text{IO}_3^-$



b,i *Nitrosomonas* sp.,  $\text{NO}_2^-$



b,ii *Nitrosococcus oceanus*,  $\text{NO}_2^-$




Figure 2








Figure 3

Nitrification + 10%



Nitrification - 10%





Figure 4