




Integrated Care in the
Community: The Case
of the *Programa Maior Cuidado* (Older Adult
Care Programme) in Belo
Horizonte-Minas Gerais, BRA

INTEGRATED CARE CASE

JANAÍNA DE SOUZA AREDES (D)
JENNY BILLINGS (D)
KARLA CRISTINA GIACOMIN (D)
PETER LLOYD-SHERLOCK (D)
JOSÉLIA OLIVEIRA ARAÚJO FIRMO (D)

*Author affiliations can be found in the back matter of this article



ABSTRACT

Internationally, there is a large body of scientific evidence concerning the benefits of integrating health and social care to ensure that frail older people living in the community receive the assistance they need to maintain independence. In the Brazilian city of Belo Horizonte, located in the state of Minas Gerais, an integrated care intervention has been developed: the *Programa Maior Cuidado – Older Adult Care Programme* (PMC). This programme represents a pioneering example in Brazil of the provision of carers for highly vulnerable older people, through integrated action between public health and social service agencies. This paper draws on the first phase of a mixed method evaluation of PMC, including data from documentary sources, focus groups, empirical observation and expert workshops, to examine the processes that led to the establishment of programme. The origins of the PMC are discussed and its operational processes, with a particular emphasis on integrated activities and the roles of different actors. The paper situates PMC within comparable international experiences of integrated provision for older people and considers how it has been affected by unique context and challenging of a middle-income country.

CORRESPONDING AUTHOR:

Janaína de Souza Aredes

Centre for Studies in Public Health and Aging, René Rachou Institute, Fiocruz, Minas Gerais, BRA

janainaaredes@gmail.com

KEYWORDS:

integrated care; older vulnerable people; health and social integration; Brazil

TO CITE THIS ARTICLE:

Aredes JS, Billings J, Giacomin KC, Lloyd-Sherlock P, Firmo JOA. Integrated Care in the Community: The Case of the *Programa Maior Cuidado* (Older Adult Care Programme) in Belo Horizonte-Minas Gerais, BRA. *International Journal of Integrated Care*, 2021; 21(2): 28, 1–12. DOI: https://doi.org/10.5334/ijic.5619

INTRODUCTION

Rapid population ageing represents a significant challenge for most low and middle- income countries. By 2015, 62% of people aged 70 or more were living in less developed regions, and this is projected to reach 76% in 2050 [1]. As in high-income countries, an ageing population requires policies to promote integrated health and social care, to support independent living at home and reduce unnecessary hospital admissions [2]. Numerous studies refer to the potential benefits of integrating health and social care, including more efficient use of health services and improved health outcomes for older people and their carers. These integrated interventions can take many forms, but mainly focus on collaboration between health and social assistance professionals, and with family carers [3, 4, 5].

Brazil currently has the 6th largest population of older people in the world, with approximately 30 million people aged 60 or older - 15% of the country's total population. By 2030, this number will surpass the number of children and adolescents and by 2050 it will reach 64 million people, 30% of Brazilian population [6]. Over the next 25 years, the share of people aged 70 or older will treble, to reach 16.4% - more than current levels in more developed regions [7]. Brazil is ageing in a context of deep economic crisis, and profound social and gender inequality, all of which present challenges to public policies. Austerity policies, similar to those in Europe, are being implemented, which may further compromise the provision of care to the most vulnerable population groups [8]. All of the above have been exacerbated by the Covid-19 pandemic, which has hit Brazil hard, with the majority of deaths occurring among people aged 60 or more [9].

As in many countries, Brazil has separate administrative systems for health services and for social care: the Unified Health System (Sistema Único de Saúde – SUS in Portuguese) established in 1990 and the Unified System for Social Assistance (Sistema Único de Assistência Social – SUAS in Portuguese) established in 2004.

There are national mechanisms to support coordination and learning between local governments and responsibility for this is mainly decentralised to the municipal level. These mechanisms focus on community-based primary health care, involving interdisciplinary Family Health Teams (*Equipe de Saúde da Família* – ESF in Portuguese) of doctors, nurses, community health agents (*Agente Comunitário de Saúde* – ACS in Portuguese), social workers, nutritionists, physiotherapist and psychologists [10]. Each team offers a comprehensive set of primary health care services to defined communities of 3000 to 4000 inhabitants. These ESFs are responsible for referral, coordinating different SUS health services and operating as a bridge between the health system and local communities [10]. The local institutional hub for

this system is the Health Centre (*Centro de Saúde* – CS in Portuguese), which provides primary health care to designated geographical populations [10].

SUAS has a somewhat wider set of roles than social welfare agencies in many high-income countries, including the management of national cash benefit schemes and the provision of social services. It has a similar decentralised structure to that of the SUS, with national coordination and local hubs for community engagement. Local hubs known as Social Assistance Reference Centres (Centro de Referência em Assistência Social – CRAS in Portuguese) perform a similar role to SUS Health Centres (CS), offering a wide range of services for people of all ages, with a particular emphasis on protecting and strengthening relationships between family members and guaranteeing human rights [11]. Unlike CS, however, it does not operate in all parts of Brazil: its coverage is restricted to areas defined as highly vulnerable and at social risk, such as precarious urban neighbourhoods.

In Brazil, even before the Covid-19 pandemic, there was growing awareness of the need to manage pressures on the SUS in the context of population ageing [12]. Rates of health service use and many preventable hospital admissions of older people [3] are higher for older Brazilians than for the population in general [13]. Different studies report that unnecessary hospital admissions and protracted lengths of hospital stay are in part due to a lack of family support or social care in the community at home [14, 15, 16]. Unnecessary hospital stays expose older people to other health risks and contravene the preferences of many older people to remain in their own homes as much as possible [16, 17]. To date, Brazil has made little progress to integrate health and social care services for older people at the national level. However, the decentralised structures of SUS and SUAS allow local governments some scope to develop their own policy interventions to promote an integrated approach. The most notable example of this is in the city of Belo Horizonte, which has been running an innovative intersectoral scheme, Programa Maior Cuidado (PMC -Older Adult Care Programme), for nearly ten years.

Implemented in 2011 as a partnership between Belo Horizonte's municipal departments of health and social assistance, PMC represents a novel approach for managing social care for highly vulnerable older people in Brazil [18]. PMC originated from discussions held by an Intersectoral Working Group formed in 2010, composed of representatives of different agencies (Education; Health; Social Assistance; Citizenship; Culture, Sports and Leisure) and the Municipal Council of Older People's Rights. This group aimed to develop strategies for supporting families with frail older people in situations of social vulnerability – as a consequence of weak family/social ties or limited opportunities for inclusion in the community, generating situations of risk, exclusion and social isolation [18, 19].

This paper draws on the first phase of a evaluation of PMC, including data from documentary sources, focus groups and empirical observation, to examine the processes that led to the establishment of PMC. We also compare how PMC functions in theory to the actual experiences of different health and social assistance professionals involved in its day to day operation. This pays particular attention to the degree PMC has promoted enhanced service integration. It also considers the specific setting in which this integrated programme has been developed. The vast majority of literature on comparable interventions refer to high-income countries [3]. Brazil, and more specifically, the city of Belo Horizonte represent a distinct context. With a population of around 2.5 million people, Belo Horizonte is the sixth largest city in Brazil and, as elsewhere in the country, is characterised by high levels of poverty, social exclusion and inequality [20]. This is most evident in the city's most deprived neighbourhoods, which are where PMC operates. As such, a key contribution of this study is the unique insight it provides into processes of service integration for older people in a highly challenging middle-income country setting.

STUDY DESIGN

This is an exploratory study, [21] seeking to map the organisation and functioning of PMC and to reveal the nature of relationships between key stakeholders and actors in different participating agencies. It is part of a wider evaluation project, "Improving the effectiveness and efficiency of Health and social care services for vulnerable Older Brazilians - IHOB" (2018-2021) approved by the René Rachou Research Ethics Committee, Fiocruz-MG (CAAE: 96033418.9.0000.5091). The research team is multidisciplinary, including social scientists, geriatricians, a nurse, health services managers and public policy makers with interests in ageing. The main objective of the larger study is to provide evidence to support policies to reduce unnecessary stays of older people in hospitals and long-term care facilities. These include admissions for conditions that could be treated outside inpatient settings, admissions for conditions that could reasonably be prevented by adequate community-based health and social care, and delayed discharge due to a lack of homebased support [2].

The project used a multi-method [22] approach and has two phases (*Figure 1*). This paper draws entirely on Phase 1, as Phase 2 has been delayed as a result of the Covid-19 pandemic. Our data sources include a systematic review of international studies of comparable interventions [3], documentary analysis of PMC's internal records and new data obtained from focus groups, non-participant observation and expert workshops. Phase 1 focussed on mapping and analysing the development

and operation of PMC. Phase 2 (ongoing) focusses on outcomes and impacts.

DATA ANALYSIS

The data obtained in Phase 1 were transcribed into Microsoft Word®. We applied three stages of content analysis: pre-analysis (data organization); exploration of the material (definition of analytical categories); and treatment of results, inferences and interpretation (reflective analysis) [23]. This included data coding and thematic analysis and was independently reviewed by two members of the research team in order to reach consensus on the final categories.

Although the specific components of the research design for Phase 1 addressed different objectives, there were important complementarities between them. For example, the document analysis revealed a number of gaps in written guidance about how PMC should operate, including inter-agency collaboration, guidelines and protocols. This in itself is a relevant finding. Consequently, the focus groups served to both fill in some knowledge gaps about how PMC operated in theory, as well as to explore how this compared to daily realities.

RESULTS

PMC IN THEORY: OBJECTIVES AND OPERATION

Plans for PMC were first developed in the Department of Health. However, this Department was concerned about the costs of employing paid carers and looked for financial support from the other municipal agency with related interests: Social Assistance. PMC then emerged as a partnership between these two departments, with Social Assistance bearing the full costs of the carers (Document analysis, 2019).

The main objectives of the PMC are presented in Figure 2. PMC aims to provide a paid daily carer to semidependent and dependent older people identified as clinically and socially vulnerable [18]. The carers only work during normal working hours, but the time spent varies according to the complexity of the case. They are hired through a contract between the municipality and a non-governmental Civil Society Organisation (Organização da Sociedade Civil - OSC in Portuguese), responsible for ensuring their training and qualifications, under the supervision of the Departments of Health and Social Assistance. It is also the role of the OSC to select the caregivers (127 in total for the 9 administrative regions), in addition to 02 administrative assistants and 02 general coordinators – who work at the headquarters of the OSC (Document analysis, 2019).

PMC's management is described as intersectoral, with joint oversight at the central level of the Secretariats of Health and Social Assistance, as well as at the local

Phases	Research steps							
^a Phase 1 - Exploratory	Systematic Review [3] Objective: to synthesise existing literature on interventions addressing avoidable displacement from home for older people with multimorbidity or frailty. The review focused on home- based interventions by multidisciplinary teams.	Fifteen studies were included in the review. Four themes were identified and discussed: (1) types of intervention (2) composition of teams, (3) intervention effectiveness and (4) types of outcomes. The review identified son potential interventions and relevant topics to be addressed in order to develop effective and sustainab interventions to reduce the avoidable displacement from home of older people. However the review was not at to identify robust impact evidence, either in terms of quantity or quality from the studies presented. As such, the available evidence is not sufficiently robust to inform policy or interventions for reducing avoidable displacement from home. This finding reflects the complexity of these interventions and a lack of systematic data collection.						
Study [21] Data collection:	Document analysis Objective: initial, exploratory analysis of PMC.	Collection of documentary data on PMC. Review of existing records of people included in PMC from it implementation until the end of 2018 (including follow-up correction of gaps and inconsistencies in records, base on complementary data sources).						
between September 2018 to May 2019	Focus groups in all health districts (n= 9) ^b Objective: map PMC's structure and organization from the perspective of professionals.	All professionals working in PMC were invited to participate. The meetings took place at the CS or CRAS an lasted an average of 84 minutes. Groups included an average of 6 to 10 professionals*, including local practitioner (nurses, social workers, nutritionists, physiotherapist and psychologists) and municipal-level manager (geriatricians and senior social workers). Topics were: PMC's objectives; registration and monitoring of participants; PMC's organization; factors that had facilitated/impeded PMC's operation (with particular reference to articulation between Health and Social Assistance); and links to higher levels of service provision.						
	Observation of meetings (n=6) Objective: Engagement with local teams; explore day-to-day dynamics.	Non-participation observation of managerial and operational meetings involving local and municipal-level health and social assistance staff, applying an appropriate, validated methodology [21].						
Phase 2 Data	Quantitative analysis	- Hospitalization rates among PMC participants Quantitative analysis of possible effects of PMC on the use of health services.						
collection: in progress	In-depth case study	- In-depth semi-structured interviews with family members, professionals, caregivers and older people included in PMC. Observational data from meetings and working groups.						

Figure 1 Study design.

a: In this stage, three Expert Workshop were also held. Objective: networks and dissemination of the first results to public policy makers.

b: Belo Horizonte, capital of the state of Minas Gerais, is organized into 9 administrative regions (health districts). PMC operates in all administrative regions.

c: In two focus groups there was also participation of Community Health Agents (Agente Comunitário de Saúde - ACS in Portuguese).

Data recording: Field notes (focus groups and observation of meetings), digitization and filing of the documents collected (document analysis). Organization of a database with the systematization of the main documents and records identified.

Source: Elaborated by the authors.




Figure 2 PMC in theory: objectives, criteria and organisational structure.

Abbreviations: CRAS: Social Assistance Reference Center (*Centro de Referência em Assistência Social* – CRAS in Portuguese); CS: Health Centre (*Centro de Saúde* – CS in Portuguese); GT: Working Group (*Grupo de Trabalho* – GT in Portuguese); Pré-GT: Pre-Working Group (*Pré-Grupo de Trabalho* – Pré-GT in Portuguese); OSC: Civil Society Organisation (*Organização da Sociedade Civil* – OSC in Portuguese); ESF: Family Health Teams (*Equipe de Saúde da Família* – ESF in Portuguese).

Source: Elaborated by the authors, research data. Document analysis, 2019.

level through joint activities involving the CS and CRAS teams (*Figure 2*). Each CRAS has staff members with specific responsibility for PMC. At the central level, programme supervisors have overall responsibility for the management of caregivers and for care quality assurance. At the local level, PMC is supported by a close collaboration between families, health professionals, social assistance professionals and PMC carers (Document analysis, 2019).

At the time of Phase 1, PMC covered 62 (40.8%) of the Belo Horizonte's 152 CS and 26 (76.5%) of its 34 CRAS (its coverage has since increased). However, the geographical territories of CS and CRAS do not always coincide. Some areas covered by a CS do not have a CRAS, and in other areas the same CS corresponds with several CRAS (Document analysis, 2019).

The stages for admission into and monitoring of PMC users are summarised in *Figure 2*. Pre-inclusion consists of identification of potential users, by a CS or a CRAS or referrals from families, hospitals or neighbours. The criteria for *inclusion* are provided in a set of survey forms which evaluate the older person's health, functional and social status. After staff from the CS and CRAS complete these forms, applications are prioritised according to the income and social vulnerability of each family, as well as the clinical status and level of functional dependence of the older person. If identified as eligible, the CRAS will meet the family and older person to discuss the scheme and obtain their consent for participation (Document analysis, 2019).

Monitoring activities include a monthly pre-Working Group meeting and the main Working Group meeting. The pre-meeting involves the paid carers, the OSC supervisor and the CRAS team. They agree on practical issues, such as shifts and rotations between caregivers, and discuss the difficulties faced by caregivers related to the care itself or other environmental risks. The main Working Group meeting runs immediately after the pre-meeting, whose participants are joined by health staff (nurses, nutritionists, physiotherapist) and where complex cases of very ill people, problems of familiar violence and discharges from PMC are discussed. Monitoring activities also include monthly and quarterly spreadsheets completed by health staff and sent to the Municipal Office for Health. Older people enrolled in PMC can remain in it for an initial period of up to two years, which can be extended if needed (Document analysis, 2019).

PMC IN PRACTICE: PRELIMINARY EVIDENCE

One of the objectives of the documentary analysis was to collect records for all people included in PMC since its implementation. These include information on age, gender, degree of care dependence and (if leaving PMC) the reason for exiting. This permits some limited descriptive quantitative analysis.

Table 1 presents data derived from PMC records for the numbers of older people included in the programme over time, by age and degree of dependence. In each year following 2011 (when PMC began), around a third of people enrolled in PMC (range 29.3% in 2014 to 45.9% in 2018) left the programme and were replaced by new members.

As revealed in the documentary analysis, since the implementation of the PMC (in 2011) until the year 2018, 1,980 elderly people were admitted to the Programme. In

YEAR	N	GENDER		DEGRE DEPEN		RATE OF RENEWAL	
		М ^ь	F°	Dd	SDe		
		%	%	%	%	— %	
2011	549	29.51	70.49	45.7	54.3	-	
2012	214	28.97	71.03	47.7	52.3	39.00	
2013	207	29.47	70.53	48.3	51.7	37.70	
2014	161	27.95	72.05	37.3	62.7	29.30	
2015	208	32.69	67.31	41.8	58.2	37.90	
2016	177	31.07	68.93	39.5	60.5	32.20	
2017	212	32.08	67.92	41	59	38.60	
2018	252	36.11	63.89	34.9	65.1	45.90	
Average	-	30.98	69.02	42.03	57.98	37.2	

Table 1 People included in PMC (by sex and degree of dependence) and rate of renewal (2011–2018).

Note: a: Total elderly people included in the PMC since its implementation: 1,980 elderly people.

Abbreviations: b: Male; c: Female; d: Dependent; e: Semi-dependent.

Source: Elaborated by the authors based on research data. Document analysis, 2019.

that same period the average volume of resources invested is around 890,000 dollars a year, which includes the hiring of the caregivers and other professionals necessary for the management of human resources and administrative staff (Field notes – Observation of meetings, 2019).

Table 2 presents data for people who left PMC, along with the main reason for leaving. The most common reason was the death of the older person (45.04% of cases). This was unsurprising given the health and functional status of the target population. The second most frequent reason (26.27%) was moving away from the local health district and hence losing local registration with PMC. In 10.94% of cases the main reason given was "institutionalization" (placement into a long-term care facility) and 7.22% of departures from PMC were due to the "rehabilitation" of the older person (so that they no longer needed assistance from PMC).

During the focus groups professionals affirmed that PMC appears to be appreciated and valued by all those involved, namely the users, professionals and family members. It was possible to identify three main thematic categories that highlighted the potential of the programme, namely: support for the family; prevention of self-neglect and prevention of clinical problems, hospitalisations and institutionalisation. *Figure 3* provides quotes from participants and recorded observations to illustrate these thematic categories. Although there was a strong consensus from study participants that PMC "mainly works well" on a day-to-day basis, a number of issues and frustrations were also identified. These were framed in terms of areas for improvement, rather than grounds for discontinuing PMC. These concerns are summarised in *Figure 3*.

Staff questioned the Rigidity of the inclusion criteria - age and territory (Figure 3). Potential inclusion in PMC was restricted to people aged 60 or more living in defined administrative zones served by specific CRASs. This could potentially exclude some vulnerable families and some CSs staff referred to the frustration of identifying older people as potentially eligible for inclusion in PMC, only to discover that they did not live within a neighbourhood linked to a CRAS. Staff recognised that these rigid criteria were mainly applied to manage high levels of demand for inclusion in PMC, which greatly exceeded their resourcing capacity. They saw this as part of a wider problem of under-resourced community services, observing that both the CSs and CRASs are small, understaffed, poorly maintained and unable to meet the needs of their communities (Field notes - Focus groups, 2019). As such, rather than an inherent criticism of PMC itself, these concerns reflected a wider frustration with the weakly resourced setting within which it operated.

A separate set of concerns related to the many challenges facing PMC carers who were working with vulnerable families in highly deprived neighbourhoods – Challenges facing PMC carers (Figure 3). This could expose PMC carers to threatening situations and generate a range of tensions and misunderstandings. These included challenges in establishing which family members had the main care responsibility for the older person and in distinguishing between the roles of PMC carers and those of health care professionals. There were sometimes tensions between the theoretical premise that families would cooperate closely with PMC carers and the complexities of dealing with

REASON	MALE	MALE			FEMALE				TOTAL	
	Da	%	SDb	%	D	%	SD	%	N	%
Change of residence	16	16.33	22	15.07	45	21.74	86	19.72	233	26.27
Family resumes care	21	21.43	22	15.07	56	27.05	97	22.25	196	22.10
Family opt out	24	24.49	26	17.81	49	23.67	69	15.83	168	18.94
Institutionalization	11	11.22	16	10.96	22	10.63	48	11.01	97	10.94
Older person opts out	6	6.12	21	14.38	11	5.31	57	13.07	95	10.71
Risk situation for the caregiver ^c	14	14.29	21	14.38	18	8.70	40	9.17	93	10.48
Rehabilitation of the elderly	5	5.10	17	11.64	6	2.90	36	8.26	64	7.22
No data ^d	1	1.02	1	0.68	-	-	3	0.69	5	0.56
Subtotal (all reasons above)	98	36.98	146	60.58	207	45.59	436	66.67	887	54.96
Death	167	63.02	95	39.42	247	54.41	218	33.33	727	45.04
Total	265	100.0	241	100.0	454	100.0	654	100.0	1614	100.0

Table 2 Reason for discharge/interruption of participation in PMC by sex and degree of dependence (2011-2018).

Source: elaborated by the authors based on research data. Document analysis, 2019.

^a: Dependent; ^b: Semi-dependent; ^c: Lack of security in the territory of operation; ^d: No reason provided.

Ti	hematic categories	Quotes							
POTENTIALITIES	Support for the family	"We have many cases of family insufficiency. Family members who cannot read, write, administer medications. The [PMC] carer comes to he families. There is an older person who has a family, but at the same time he does not there is the case of an older person with 10 children, but of takes care of him. When the [PMC] carer started working at the house, he even helped the health of the son of this older man who was overwork [PMC] carer even helps in the quality of life of the family caregiver". (Social Worker - Focus groups, 2019)							
	Prevention of self- neglect	"The programme allows us to see before and after care. It not only prolongs the life of older people, but brings the older people back to life". (Nurse - Focus groups, 2019) "Here we had a remarkable case of a lady known as "garbage hair". She stored all kinds of garbage in her hair - cigarette butts, candy wrappers, food packaging, etc. After the work of the programme's carer, the older woman began to bathe frequently and to have more energy. It's the kind of case we tell							
	Prevention of clinical problems, hospitalisation and institutionalisation	everyone" (), (Nurse - Focus groups, 2019) "The issue of self-neglect in relation to care is very common among older people. This makes pre-existing diseases worse and even other disease sappear. The presence of the [PMC] carer is very important in this regard. He signals and intervenes before the problem gets worse. There are older people who would have passed away". (Nurse - Focus groups, 2019) "We had the case of a very frail older woman who lives with her husband who is also frail. She was not able to self-care. Prior to the programme, she had been hospitalised several times for dehydration. The programme avoids these unnecessary hospitalisations because the [PMC] carer intervenes in these cases. They notify the Health Centre that takes the necessary steps without even having to send the older person to a hospital. In the case of this lady, without the programme, she would have died long ago". (Social Worker - Focus groups, 2019) "The existence of the programme saves a lot for the State. In addition, we are investing in those who have great need, needy people. The presence of the [PMC] carer positively influences the progress of their disease". (Social Worker - Focus groups, 2019). "Older people create a closeness, a bond with the [PMC] carer. With this, the older people provide information to the professional who is not comfortable telling the doctor, for example. This helps a lot in care, especially in the Health Centre". (Nurse - Focus groups, 2019) "There are cases of older people who improve a lot after the [PMC] carer comes in and we even think about discharging them. However, most of the time when older people leave the programme, their condition gets worse, because they often live alone and do not have the support of the family. In these cases, we choose to decrease the amount of hours spent with this older person instead of taking him out of the programme". (Nurse - Field Diary, 2019)							
	Rigidity of the inclusion criteria (age and territory)	"How can we adjust (the criterion) to that 57/58 years old patient who is in the same frail condition as an older person over 60 years old?". (Social Worker -Focus groups, 2019) "We had a case where we were at the door of the older person's house with everything prepared for inclusion. Even his neighbour across the road was a member of the PMC. When we arrived at the door of the older person's house, the driver came and said: 'but this house is not part of the CRAS area'. We said: 'What do you mean?! The man across the street is!' Then we had to go back. It was sad because he was a very needy older person, and everything was ready, filling in the instruments was all that was missing.' (Nurse - Focus groups, 2019) "There are more serious patients who are placed outside the CRAS area. There was an older man who lived in a shantytown and when we went to visit him he had several bug infestations. There were even animals under his bed. I remember that a drug dealer who lived near his house, the day I was there he asked me: 'Do you want me to shoot him in the forehead? I can do it now and get it over with'. The dealer, in fact, was compassionate about that situation, he wanted to give that gentleman a chance. He ended up being institutionalised". (Social Worker - Focus groups, 2019)							
LIMITS		"It must be made clear that the [PMC] carer can also work in areas with high vulnerability. Health works anywhere. Why can't the PMC carer also? Health is universal, but assistance is restricted". (Social Worker - Focus groups, 2019)							
	Challenges facing PMC carers	"There is a [PMC] carer who doesn't even know the older man's son. We have cases of extremely negligent family members. There is still a lack of understanding of the family in relation to the real role and function of the PMC carer. We received cases of family members asking us why the PMS carer did not do household chores in the older person's home, like: 'it was full of pots in the kitchen to wash and he didn't wash. I had to make lunch and he didn't My father wanted to eat pork and the [PMC] carer didn't want to do it'. This is still common". (Nurse - Focus groups, 2019) "We cannot ignore the history of the older people. There are cases of older people who in the past raped their daughters, were alcoholics, aggressive. He was not a good father, a good husband. The family has no obligation to take on this care. It is her right not to want to be involved with that person anymore and that is beyond our control". (Social Worker - Focus groups, 2019) "Older people who live alone is a complex issue, as one of the criteria of the PMC is the frail older person who has a family caregiver/ friend/neighbour who can provide support. However, there are cases of older people very frail without family members. Do we put this older person in the PMC or not? It is an issue that needs to be discussed: assess whether older people who live alone is part of the PMC profile. There are cases in which the [PMC] carer arrives at the house and the older person is lying on the floor he fell because he had no support in the carer's absence". (Nurse - Focus groups, 2019) "We recently had a family member with a severe mental disorder in an area of high social vulnerability and violence. In a crisis situation for this family member, the [PMC] carer was prevented from leaving the residence (the family member locked him in the house). There are cases of family members involved in violence and use of illicit drugs. There was also the case of an infestation of chicken lice in the older person's home. This caused allergies in two followed							
	Lack of institutional quality assurance in the programme	"It is necessary to look more at [PMC] carers. It is a very exhausting service. There are situations in homes where the [PMC] carer is not able to use the bathroom during care and no place to eat your snack". (Social Worker - Focus groups, 2019) "We did not receive any guidance on how to record and monitor older people in the programme. I know that some were hospitalised and records I have are for my own use. In fact, there were adways these two spreadsheets: the monthly and the quarterly, but the guidance was always forwarded to the personal email of the [technical reference of the Municipal Health Department]. I always filled out the spreadsheets and forwarded them to her". (Nutrse - Focus groups, 2019) "It is necessary to review and simplify the content of these monitoring worksheets and the care plan itself. The information needed is very complex. There are many documents to complete and it is not always the same professional who fills them in. There is also a need for guidance and alignment between CRAS and the Health Centre regarding the organisation and systematisation of these documents". (Social Worker - Focus groups, 2019) "We do not have a document that contains all of the programme's guidelines and each team's own duties. I think there is a need to strengthen the integration between the teams and refine the relationshins between the professionals". (Social Worker - Focus groups, 2019)							

Figure 3 Potential and limits of the PMC – professionals' perspective.

Source: Testimonials collected during the meetings focus groups. Field notes, 2019.

highly vulnerable individuals. The capacity of family members to work alongside PMC carers to support older relatives was sometimes limited due to overload, a lack of education and problematic personal relations between relatives (Field notes – Focus groups, 2019). Again, rather than an inherent weakness of PMC, these concerns reflected the hugely challenging contexts in which it operated.

A third set of concerns were more integral to PMC itself – Lack of institutional quality assurance in the programme (Figure 3). These refer to the limited institutional embeddedness of the programme and challenges in cross-agency collaboration. These concerns were supported by our documentary review which found few official materials defining PMC's formal institutional status, management guidelines or collaboration protocols. Instead, there was evidence that relationships between different agencies and ways of working together, both at

the community and municipal levels, had been allowed to develop organically as the programme evolved. This may have permitted some flexibility through common sense learning-by-doing, but had impeded the formalisation of these arrangements. The absence of comprehensive written protocols occurred in a context of high staff turnover at both the municipal and community levels, generating confusion about roles and responsibilities (Field notes – Focus groups, 2019). According to some focus group participants, CS health teams did not always meet PMC expectations due to the many demands on their time and a lack of specific protocols.

A direct consequence of the limited formalisation of PMC was inconsistent record keeping and a lack of information sharing across different agencies (Field notes – Focus groups, 2019). The application of monitoring instruments (monthly and quarterly spreadsheets) was inconsistent across different areas.

There were gaps in the PMC data systems operated by both the departments of Health and Social Assistance and they were mutually incompatible, thus preventing data sharing (Field notes – Focus groups and Document analysis, 2019). Separately from this, PMC carers were required to write up daily case notes and compile these into monthly reports for the Civil Society Organisation which had originally recruited and trained them. These records were not shared with the rest of the Health and Social Assistance teams and most professionals involved in PMC did not understand the purpose of this exercise or the meaning of the data (Field notes - Focus groups, 2019).

In sum, our data revealed that, although PMC was almost universally viewed as highly effective, the programme was not without flaws and areas of weakness. These strengths and limitations, summarised in Figure 4, mainly reflected PMC's somewhat ad hoc evolution and the challenges of working with vulnerable families. Having operated along somewhat informal lines for several years, the view of most study participants was that the time had come to take PMC "to the next level", with clearer and more consistent understandings. Following the presentation of these findings to the Departments of Health and Social Assistance, steps have been taken to new develop simpler and more systematic records, as well as enhanced protocols for data sharing across agencies. A new official document sets out the main elements of PMC, as well as the specific roles and responsibilities of different agencies [24]. This is the first document to set out this information and, as such, represents a key step towards formalising

PMC's institutional status. More generally, the evident strengths of PMC revealed in our study led to the expansion of the programme to all previously unserved CRAS in Belo Horizonte and the piloting of similar schemes in other cities.

DISCUSSION

This study presents results from the first phase of evaluation of an innovative integrated health and social care programme for older people in Brazil. These data must be analysed with reference to the wider national context. Although Brazil has an extensive legal framework for supporting older people, public policies are still not being fully implemented. Brazil has mainly focussed on providing contributory pensions and meanstested cash benefits for older people [25]. However, these income transfers are not enough to address the care needs of a population that is ageing in difficult social and economic conditions. There is little coordination between federal departments responsible for health and social care. This disconnect feeds down to lower tiers of government, including at the district level where separate infrastructures are based around CS and CRAS whose territories in some cases do not coincide.

It should be noted that the Program should serve a much larger number of people when compared to the total elderly population in the region studied. Currently the city of Belo Horizonte has about 380,000 elderly people, of whom it is estimated that 8% would be incapacitated in basic activities of daily living [26] and 75% totally dependent on the Unified Health System.



LIMITS POTENTIAL STRENTHS

- Institutional quality assurance;
- Weak integration between Social Assistance and Health policies;
- Incompatible information systems;
- Absence of evaluation and monitoring.

- Sustainability: continuity during different political administrations;
- Positive evaluation of the staff of the two policies;
- Compliance with basic objectives

IMPLICATIONS OF THE PRELIMINARY RESEARCH

- Programme strengthening and institutionality:
 - Publication of the first inter-sectorial decree of co-responsibility between Health and Social Assistance policies;
 - Improvement of instruments (Instrument for Inclusion of Older People in PMC, Care Plan for Dependent Older People and Multidimensional Evaluation Protocol);
 - Expansion of the Program to all CRAS in the city of Belo Horizonte;
 - Organisation of periodic meetings between Health and Social Assistance at the central management level of the Program.
- Expansion to the municipalities of Contagem (Minas Gerais) and Fortaleza (Ceará).



Figure 4 Limits and potential strengths of PMC.

Abbreviations: CRAS: Social Assistance Reference Center (Centro de Referência em Assistência Social - CRAS in Portuguese).

Source: Elaborated by the authors based Observation of meetings. Field notes, 2019.

When it comes to comparing the theoretical aims, objectives and ambitions of PMC with the practicalities of implementation, there are a number of contrasting issues. Just as in other countries, the experience of PMC demonstrates that effective intersectoral cooperation is a major challenge, especially in contexts of high clinical and social vulnerability. Notwithstanding positive statements by stakeholders on the scheme, the problems identified were similar to other studies of integrated health and social care interventions in highincome countries and referred to difficulties due to the complexities of existing professional communication pathways and incompatibility of systems [27, 28]. The inclusion of other professionals such as mental health teams and community doctors is also seen as desirable, but often unfeasible due to competing priorities and resource deficits [29]. Some barriers to effective collaboration are more specific to counties like Brazil. For example, Brazil has very high rates of staff turnover in health and welfare agencies, among other things, due to temporary contracting and a tendency for new governments to replace staff with their "own people" [30].

There are clear discrepancies between theoretical expectations about collaboration with families (which to some degree reflected Latin American cultural norms and ideals of family care-giving) and the challenging realities of struggling families in highly-deprived communities [31]. Laws requiring families to care for older people need to consider their real capacities to do so [32]. By offering family carers respite and support, PMC goes some way towards recognising these challenges, but is not always able to resolve them. Nevertheless, PMC shows the value of interventions that see family support, along with formal health and social care provision, as complementary parts of an integrated support system for frail older people [17].

Despite the challenges outlined, PMC represents a pioneering programme that has been able to establish a meaningful partnership between local departments of health and social assistance and to sustain this partnership over nearly a decade. These positive perceptions contributed to decisions to sustain PMC over two different municipal administration. This is unusual in Brazil, where programmes of this nature rarely outlast a single electoral cycle [33]. They also contributed to a decision, in 2019, to extend PMC into new neighbourhoods, to strengthen its information systems and to formalise its institutional design. These developments reflect the appetite of policymakers in Brazil and other middle-income counties to identify more effective approaches to meet the health and social care needs of their fast-growing older populations [34, 35]. PMC can also be understood as a social innovation, promoting Ageing in Place through adequate health and social support [17, 36, 37].

LIMITATIONS AND STRENGTHS OF THE STUDY

These results analyse the structure and operation of PMC, but do not explicitly discuss its effects or impacts. Clearly, the two are linked, inasmuch as effective operation and intersectoral coordination are conducive to more positive impacts, and general comments about PMC from a range of participants and stakeholders were favourable.

A strength of our research design is its use of multiple methods: document review, focus group discussions and non-participant observation. This permitted an element of validation and consistency checking across data, although some of the qualitative data may reflect the subjective biases and preferences of participants [38, 39]. The second phase of the study will develop a more substantial set of data from different sources, including the perspectives of older people and families participating in PMC.

CONCLUSION

The insufficient provision of adequate services facing the new demographic condition affects the daily routine of users, the community and the Health and Social Assistance systems. Intersectoriality is another challenge, especially in contexts of high clinical and social vulnerability.

The Older Adult Care Programme is a beacon of integration between the two social protection public policies in the Brazil. This research allowed to identify the potential of the Programme, promoting its strengthening and institutionalization. Notwithstanding the challenges PMC has faced and its sometimes *ad hoc* operation, overall comments about the programme from a range of participants and stakeholders were strongly favourable.

However, some gaps were identified in the implementation of the Program, which corroborates the need to systematically monitor its execution, not to point out errors and flaws, but to improve processes that guarantee its effectiveness. The preliminary findings from this study also provided a clearer description of the challenges and considered the extent to which they reflect specific problems with the PMC itself or are the result of other bigger issues, such as a lack of resources.

Although the results of this study are limited to a level of local action, it is observed that supplement the complex health needs with limited resources is a global challenge.

REVIEWERS

Dr. Julie MacInnes, Research Fellow, Integrated Care, ICAP Programme Lead, Centre for Health Services Studies, University of Kent, Canterbury, UK.

One anonymous reviewer.

FUNDING INFORMATION

This study is funded by a partnership between the UK's Medical Research Council/Newton Fund (Grant Number: MR/R024219/1) and the Confederation of Foundations for Brazilian Research Support, with the participation of FAPEMIG (Grant Number: APQ-01141-18) and FUNCAP/CE (Grant Number: 0133-00003.01.00/18).

COMPETING INTERESTS

The authors have no competing interests to declare.

AUTHOR AFFILIATIONS

Janaína de Souza Aredes o orcid.org/0000-0002-4147-2405 Centre for Studies in Public Health and Aging, René Rachou Institute, Fiocruz, Minas Gerais, BRA

Jenny Billings orcid.org/0000-0002-5660-9478
Centre for Health Services Studies, University of Kent, Canterbury, Kent, UK

Karla Cristina Giacomin Dorcid.org/0000-0002-9510-6953 Belo Horizonte Municipal Health Secretary, Minas Gerais, BRA

Peter Lloyd-Sherlock D orcid.org/0000-0002-1367-0703
University of East Anglia, School of International Development,
Norwich, UK

Josélia Oliveira Araújo Firmo o orcid.org/0000-0001-5330-476X
Centre for Studies in Public Health and Aging, René Rachou
Institute, Fiocruz, Minas Gerais, BRA

PUBLISHER'S NOTE

This paper underwent peer review using the *Cross-Publisher COVID-19 Rapid Review Initiative*.

REFERENCES

- United Nations. Department of Economic and Social Affairs; 2017. Population Division. World Population Prospects The 2017 Revision Volume I: Comprehensive Tables. Available from: https://population.un.org/wpp/ DataSources/.
- Lloyd-Sherlock P, Billings J, Giacomin K, Aredes J, Bastos J, Camarano A, et al. Meeting the complex challenge of health and social care provision for rapidly-ageing populations. Introducing the concept of "avoidable displacement from home". Reports in Public Health [Cadernos de Saúde Pública], 2020; 36(3): 1–12. DOI: https://doi.org/10.1590/0102-311x00162819 [in Portuguese]
- 3. **Sempé L, Billings J, Lloyd-Sherlock P.** Multidisciplinary interventions for reducing the avoidable displacement from home of frail older people: a systematic review.

- BMJ Open, 2019; 9(11). DOI: https://doi.org/10.1136/bmjopen-2019-030687
- 4. **The Kings Fund.** A year of integrated care systems: reviewing the journey so far. 2018. Available from: https://www.kingsfund.org.uk/publications/year-integrated-care-systems
- Stranges PM, Marshall VD, Walker PC, Hall KE, Griffith DK, Remington T. A multidisciplinary intervention for reducing readmissions among older adults in a patient-centered medical home. The American journal of managed care, 2015; 21(2): 106–113. Available from: https://www.ajmc. com/view/a-multidisciplinary-intervention-for-reducingreadmissions-among-older-adults-in-a-patient-centeredmedical-home
- Alcântara AO, Camarano AA, Giacomin KC. Política nacional do idoso: velhas e novas questões. Rio de Janeiro: Ipea; 2016.
- United Nations. World Population Prospects 2019.
 Available from: https://population.un.org/wpp/Download/ Standard/Population/
- Santos IS, Vieira FS. The Right to healthcare and fiscal austerity: the Brazilian case from an international perspective. [Direito à saúde e austeridade fiscal: o caso brasileiro em perspectiva internacional]. Rev Ciência & Saúde Coletiva, 2018; 23(7): 2303–2314. DOI: https://doi.org/10.1590/1413-81232018237.09192018 [in Portuguese]
- 9. Barbosa IR, Galvão MHR, Souza TA, Gomes SM, Medeiros AA, Lima KC. Incidence of and mortality from COVID-19 in the older Brazilian population and its relationship with contextual indicators: an ecological study. [Incidência e mortalidade por COVID-19 na população idosa brasileira e sua relação com indicadores contextuais: um estudo ecológico]. Rev Bras Geriatr Gerontol, 2020; 23(1). Available from: https://www.rbgg.com.br/edicoes/v23n1/RBGG%20 v23n1%20PORT_2020-0171.pdf [in Portuguese]. DOI: https://doi.org/10.1590/1981-22562020023.200171
- Macinko J, Harris MJ. Brazil's family health strategy delivering community-based primary care in a universal health system. New England Journal of Medicine, 2015; 372(23): 2177–2181. DOI: https://doi.org/10.1056/ NEJMp1501140
- 11. Borges MCM. The elderly person in the single social assistance system. [A pessoa idosa no sistema único de assistência social]. In: Berzins MV, Borges MC. Public policies for an aging country [Políticas públicas para um país que envelhece]. São Paulo: Martini, 2012; 99–124. [in Portuguese]
- 12. **Paim J, Travassos C, Almeida C, Bahia L, Macinko J.** The Brazilian health system: history, advances, and challenges. *Lancet*, 2011; 377: 1778–97. DOI: https://doi.org/10.1016/S0140-6736(11)60054-8
- 13. Stopa SR, Malta DC, Monteiro CN, Szwarcwald CL, Chester MG, Cesar LG. Use of and access to health services in Brazil, 2013 National Health Survey [Acesso e uso de serviços de saúde pela população brasileira, Pesquisa Nacional de Saúde 2013]. Rev Saúde Pública, 2017; 51(1:3s). DOI:

https://doi.org/10.1590/s1518-8787.2017051000074 [in Portuguese]

- 14. Romero DE, Marques A, Barbosa AC, Sabino R.

 Hospitalizations of elderly people for long-term care in SUS hospitals in Rio de Janeiro: an analysis of their characteristics and the fragility of social care networks [Internações de idosos por cuidados prolongados em hospitais do SUS no Rio de Janeiro: uma análise de suas características e da fragilidade das redes sociais de cuidado]. In: Camarano A, editor. Long-term care for the elderly: a new social risk to be taken? [Cuidados de longa duração para a população idosa: um novo risco social a ser assumido?] Rio de Janeiro: Ipea; 2010. [in Portuguese]
- 15. Fialho CB, Lima-Costa MF, Giacomin KC, Loyola Filho AI. Functional capacity and use of health services by the elderly in the Metropolitan Region of Belo Horizonte, Minas Gerais, Brazil: a population-based study [Capacidade funcional e uso de serviços de saúde por idosos da Região Metropolitana de Belo Horizonte, Minas Gerais, Brasil: um estudo de base populacional]. Cad Saúde Pública, 2014; 30(3): 599–610. DOI: https://doi.org/10.1590/0102-311X00090913 [in Portuguese]
- 16. Silva AMM, Mambrini JVM, Peixoto SV, Malta DC, Lima-Costa MF. Use of health services by Brazilian older adults with and without functional limitation [Uso de serviços de saúde por idosos brasileiros com e sem limitação funcional]. Rev de Saúde Pública, 2017; 51(5s). DOI: https://doi.org/10.1590/s1518-8787.2017051000243 [in Portuguese]
- World Health Organization (WHO). World Report on Ageing and Health. In Geneva: World Health Organisation;
 2015
- 18. **Sartini CM, Correia AM.** Greater Care Program: qualifying and humanizing care [Programa Maior Cuidado: qualificando e humanizando o cuidado]. *Pensar/BH Política Social*, 2012; 31: 10–13. [in Portuguese]
- 19. Barbosa DCA. Public Policies and Population Aging: analysis of the Greater Care Program in Belo Horizonte [Políticas Públicas e Envelhecimento Populacional: análise do Programa Maior Cuidado em Belo Horizonte]. Masters dissertation in Public Administration [Mestrado em Admnistração Pública] Escola de Governo Professor Paulo Neves de Carvalho da Fundação João Pinheiro, Belo Horizonte; 2019. Available from: http://tede.fjp.mg.gov.br/handle/tede/434 [in Portuguese].
- Brazilian Institute of Geography and Statistics [Instituto Brasileiro de Geografia e Estatística]. Demographic censos, 2010 [Censo demográfico]. Available from: https://cidades. ibge.gov.br/brasil/mg/belo-horizonte/pesquisa/23/25359 [in Portuguese].
- 21. **Quivy R, Campenhoudt LV.** Manuel de recherche en sciences sociales. Paris: Dunod; 1995.
- Fetters MD, Curry LA, Creswell JW. Achieving integration in mixed methods designs principles and practices.
 Health Service Research, 2013; 48(6): 2134–2156. DOI: https://doi.org/10.1111/1475-6773.12117

- 23. **Bardin L.** L'analyse de contenu. Paris: Presses Universitaires de France; 2013. DOI: https://doi.org/10.3917/puf.bard.2013.01
- 24. Belo Horizonte City Hall [Prefeitura Municipal de Belo Horizonte]. Ordinance nº 007, of November 29, 2019. Establishes the principles and guidelines for the management and operation of the Older Adult Care Programme and institutes management and dialogue instances and takes other measures [Estabelece os princípios e diretrizes para gestão e operacionalização do Programa Maior Cuidado e institui instâncias de gestão e de interlocução e dá outras providências]. Official Gazette of the Municipality. 03 Dec 2019. [in Portuguese]
- 25. Alcântara AO, Camarano AA, Giacomin KC. National policy for the elderly: old and new issues [Política nacional do idoso: velhas e novas questões]. Rio de Janeiro: Instituto de Pesquisa Econômica Aplicada; 2016. [in Portuguese]
- 26. Bernardes GM, Mambrini JVDM, Lima-Costa MF, Peixoto SV. Multimorbidity profile associated with disability among the elderly living in the Metropolitan Region of Belo Horizonte, Brazil [Perfil de multimorbidade associado à deficiência em idosos residentes na Região Metropolitana de Belo Horizonte, Brasil]. Cienc saúde coletiva, 2019; 24(5): 1853–1864. DOI: https://doi.org/10.1590/1413-81232018245.17192017 [in Portuguese]
- Hutchinson T, Young J, Forsyth D. National pilot audit of intermediate care. Clinical Medicine, 2011; 11(2): 146–149.
 DOI: https://doi.org/10.7861/clinmedicine.11-2-146
- MacInnes J, Jaswel S, Mikelyte R, Billings J. Implementing an integrated acute response service: professional perceptions of intermediate care. *Journal of Integrated* Care, 2020. DOI: https://doi.org/10.1108/JICA-03-2020-0016
- 29. De Bruin SR, Billings J, Stoop A, Lette M, Ambugo EA, Gadsby E, et al. Different contexts, similar challenges. SUSTAIN's experiences with improving integrated care in Europe. International Journal of Integrated Care, 2020; 20(2): 17. DOI: https://doi.org/10.5334/ijic.5492
- 30. Medeiros CRG, Junqueira AGW, Schwingel G, Carreno I, Jungles LAP, Saldanha OMFL. Nurses and doctors turnover: an impasse in the implementation of the Family Health Strategy [A rotatividade de enfermeiros e médicos: um impasse na implementação da Estratégia de Saúde da Família]. Cienc saúde coletiva, 2010; 15(s1): 1521–1531. DOI: https://doi.org/10.1590/S1413-81232010000700064 [in Portuguese]
- 31. Gascón S, Redondo N. Quality of long-term services for dependent elderly people [Calidad de los servicios de largo plazo para personas adultas mayores con dependencia]. Santiago: EuroSocial CEPAL; 2014 (Colección Estudios, 7). [in Spanish]
- 32. Giacomin K, Duarte Y, Camarano A, Nunes D, Fernandes D. Care and functional disabilities in daily activities ELSI-Brazil [Cuidado e limitações funcionais em atividades cotidianas ELSI-Brasil]. Rev Saude Publica, 2018; 52(2:9s). DOI: https://doi.org/10.11606/s1518-8787.2018052000650 [in Portuguese]

- Colombo LA, Alves AT. Public policies and government agendas: main theories [Políticas públicas e agendas governamentais: principais teorias]. Rev Profanações, 2020; 7(esp): 173–184. [in Portuguese].
- 34. **Southby K, Gamsu M.** Factors affecting general practice collaboration with voluntary and community sector organisations. *Health & Social Care in the Community*, 2018; 26(3): 360–369. DOI: https://doi.org/10.1111/hsc.12538
- 35. You E, Dunt D, Doyle C. What is the role of a case manager in community aged care? A qualitative study in Australia. Health & Social Care in the Community, 2016; 24(4): 495–506. DOI: https://doi.org/10.1111/hsc.12238
- 36. **Wiles JL, Leibing A, Guberman N, Reeve J, Allen RE.**The meaning of "aging in place" to older people. *The*

- Gerontologist, 2012; 52(3): 357–366. DOI: https://doi.org/10.1093/geront/gnr098
- 37. **Gulliford MC, Jack RH, Adams G, Ukoumunne OC.**Availability and structure of primary medical care services and population health and health care indicators in England.

 BMC Health Services Research, 2004; 4(12). DOI: https://doi.org/10.1186/1472-6963-4-12
- 38. **Nooteboom B.** Learning by interaction: Absorptive capacity, cognitive distance and governance. *Journal of Management and Governance*, 2000; 4: 69–92. DOI: https://doi.org/10.1023/A:1009941416749
- 39. **Amunarriz G, Alcalde-Heras H.** Towards an Integrated Care Organisation from a CEO Perspective. *International Journal of Integrated Care*, 2020; 20(3): 10. DOI: https://doi.org/10.5334/ijic.5559

TO CITE THIS ARTICLE:

Aredes JS, Billings J, Giacomin KC, Lloyd-Sherlock P, Firmo JOA. Integrated Care in the Community: The Case of the *Programa Maior Cuidado* (Older Adult Care Programme) in Belo Horizonte-Minas Gerais, BRA. *International Journal of Integrated Care*, 2021; 21(2): 28, 1–12. DOI: https://doi.org/10.5334/ijic.5619

Submitted: 29 September 2020 Accepted: 16 March 2021 Published: 21 June 2021

COPYRIGHT:

© 2021 The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC-BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See http://creativecommons.org/licenses/by/4.0/.

International Journal of Integrated Care is a peer-reviewed open access journal published by Ubiquity Press.

