

RESEARCH

Open Access



Smoking cessation message testing to inform app-based interventions for young adults – an online experiment

Josef Hamoud¹, Janardan Devkota², Timothy Regan³, Amanda Luken², Joseph Waring², Jasmin Jiuying Han², Felix Naughton⁴, Roger Vilardaga⁵, Jonathan Bricker^{6,7}, Carl Latkin⁸, Meghan Moran⁸ and Johannes Thru^{2,9,10*}

Abstract

Background To improve the efficacy of digital smoking cessation interventions for young adults, intervention messages need to be acceptable and appropriate for this population. The current study compared ratings of smoking cessation and urge reduction messages based on Cognitive Behavioral Therapy (distraction themed) and Acceptance and Commitment Therapy (acceptance themed) in young adults who smoke.

Methods A total of 124 intervention messages were rated by an online Qualtrics panel of $N=301$ diverse young adults who currently smoked tobacco cigarettes (Age $M=26.6$ years; 54.8% male; 51.5% racial/ethnic minority; 16.9% sexual or gender minority (SGM); 62.5% daily smoking). Each participant rated 10 randomly selected messages (3,010 total message ratings; 24.3 ratings per message) on 5-point scales (higher scores representing more favorable ratings) evaluating quality of content, quality of design, perceived support for coping with smoking urges, and perceived support for quitting smoking. Mixed models examined associations between message category (distraction vs. acceptance), participant level predictors (sociodemographic variables, readiness and motivation to quit, daily smoking, psychological flexibility), and message ratings.

Results Overall ratings ranged from $M=3.61$ ($SD=1.25$) on support for coping with urges to $M=3.90$ ($SD=1.03$) on content, with no differences between distraction and acceptance messages. Male participants gave more favorable ratings on the dimensions of support for coping ($p<0.01$) and support for quitting ($p<0.01$). Participants identifying as SGM gave lower ratings for message design ($p<0.05$). Participants with a graduate degree gave higher ratings on support for coping with urges and support for quitting (both $p<0.05$). Higher motivation to quit was associated with more favorable scores across all dimensions (all $p<0.01$). Those smoking daily rated messages as less helpful for coping with urges ($p<0.01$) and quitting smoking ($p<0.05$) compared to those smoking non-daily. Few interactions were found between message category distraction vs. acceptance and participant characteristics.

Conclusions Distraction and acceptance messages received similar ratings among young adults who smoke cigarettes. Message revisions may be needed to increase appeal to women, SGM, those with lower education, and

*Correspondence:
Johannes Thru
jthru@jhu.edu

Full list of author information is available at the end of the article



© The Author(s) 2025. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

those less motivated to quit. Messages will be refined and used in an ongoing micro-randomized trial to investigate their real-time impact on smoking urges and behaviors.

Keywords Smoking cessation, Messages, Mobile health, Communication, Evaluation

Introduction

Cigarette smoking accounts for close to half a million deaths in the US each year, and is the leading health risk behavior for increased morbidity and mortality [1, 2]. While smoking prevalence has generally decreased over recent decades [3], smoking rates among young adults remain high, and evidence suggests many people who smoke initiate smoking in young adulthood, making this demographic an important group for new smoking cessation interventions [4–9]. More public health research is needed to understand how to reduce smoking disparities as well as high smoking rates among young adults [10].

Low socioeconomic status (SES) is an important risk factor for cigarette smoking and reduced success in quitting. Increased smoking rates among members of racial/ethnic minority groups [3, 11] and adults of low SES [3, 12] are well established. For example, 21.3% of adults with an annual household income <\$35k smoke cigarettes compared to 13.7% of adults in the general US population [3]. Similar disparities exist based on education level, such as low education level correlating with increased smoking prevalence among all racial/ethnic groups [13]. Some of these disparities along lines of race/ethnicity and SES [14, 15] can already be observed during the developmental phase of young adulthood [16], and may endure into later life. Moreover recent evidence suggests disparities among adolescent and young adult members of sexual or gender minority groups (SGM) compared to their heterosexual counterparts, putting them at an increased risk for tobacco use [17, 18]. In sum, young adults, who are also often exposed to extensive digital marketing of nicotine products [19, 20], are priority populations for early smoking cessation interventions and sociodemographic risk factors need to be considered in the design of these interventions.

While digital and mobile interventions are in demand, their effective implementation and evaluation in real life settings is lacking [21]. For example, smartphone apps for smoking cessation, a subtype of digital interventions, have become increasingly popular among people who smoke who want to quit. These smartphone apps typically provide tools and resources to help users quit smoking through features like personalized quit plans, progress tracking, urge management, motivational messages, and access to support communities, often enhanced with gamification (elements of game playing, e.g., points, competitions) or reminders that sustain user engagement [22]. As of 2020, English-language smoking cessation apps have been downloaded 33 million times

[23]. The low cost and convenience of smartphone-delivered interventions make them an appealing option for many people who smoke, especially those disproportionately affected by the harms of smoking, including those with low income or education, and racial minority subgroups [24]. In addition, the widespread availability of smartphones—96% of US adults 18–29 years old owned one as of 2022—has helped extend the reach of smoking cessation apps to a large number of young adults who may not otherwise have access to cessation services [25]. Thus, smartphone-based interventions hold great promise for reducing the high burden and disparities in smoking-related health risk and mortality. Although many cessation apps do not follow evidence-based guidelines [26–28], some have now been developed using established evidence-based therapies [22]. However, their efficacy is still a major point of investigation in current research [29–32].

Two key psychotherapy approaches that inform digital smoking cessation interventions and apps are Cognitive Behavioral Therapy (CBT) [33, 34] and Acceptance and Commitment Therapy (ACT) [35, 36]. CBT is focused on changing maladaptive thought and behavior patterns, and has been established as an effective therapy for the treatment of a variety of mental health and substance use disorders, including smoking cessation treatment [37–40], for in-person as well as digital delivery. Distraction, or shifting attention and awareness to something else when feeling preoccupied with maladaptive or counterproductive thoughts, is one of multiple skills taught in CBT. Thus, distraction may be conceptualized as a coping strategy in CBT, and may be particularly relevant for just-in-time mobile interventions that aim to interrupt habitual smoking behaviors in the moment. Evidence suggests that distraction strategies can help people who smoke cope with smoking urges; [41] however, more research is needed to investigate perceptions of distraction-themed CBT messages before they can be successfully implemented in digital interventions. In contrast to distraction-themed CBT intervention messages, messages based on a contemporary “third wave” form of CBT called Acceptance and Commitment Therapy (ACT), encourage individuals to pay attention to and build a different relationship with their internal urge-related experiences. For example, one skill emphasized in ACT is acceptance, or being willing to feel and not act on urges to smoke cigarettes, and this concept is called “psychological flexibility” [42]. ACT for smoking cessation has been investigated in previous randomized trials [43–45].

Smoking cessation messages modeled around not only traditional CBT but also ACT theory could provide a wider array of evidence-based and potentially effective digital interventions; however, their perception among diverse groups of young adults needs to be established before efficacy testing.

In summary, to improve the development and efficacy of novel digital smoking cessation interventions for young adults, including those from diverse backgrounds, intervention messages must meet their needs. While the rating of intervention messages may not align with their actual effectiveness, low rated messages certainly preclude and limit their effectiveness. Therefore, both perceived and actual effectiveness are critical requirements of these digital interventions. The current study used an online panel of young adults who smoke interested in quitting smoking to investigate message ratings of both CBT and ACT-themed intervention messages on different dimensions related to content, design, and perceived message effectiveness to cope with smoking urges and to quit smoking.

Methods

Procedure

A total of 124 intervention messages were developed by the research team. Intervention messages came from several previous studies that tested CBT- and ACT-based smartphone interventions [23, 46, 47], were refined

internally, and combined with image content from free stock photo websites (Pexels, Unsplash) [48, 49]. Broadly, distraction messages focused on action-oriented strategies, prompting individuals to redirect their attention away from cravings through specific tasks or behaviors. By promoting active engagement and creative problem-solving, distraction messages aimed at helping individuals break habitual responses to smoking triggers. In contrast, acceptance messages emphasized mindfulness, self-awareness, and reframing cravings as transient and external experiences, encouraging individuals to observe urges without judgment and focus on the present moment. These acceptance messages aimed to empower users by fostering a sense of control through non-resistance and cognitive reframing. See Fig. 1 for intervention message examples. All messages are available on the study OSF page. Study procedures were approved by the Institutional Review Board of the Johns Hopkins Bloomberg School of Public Health.

For the current study, these 124 intervention messages were rated by an online Qualtrics panel of $N=301$ diverse young adults (18–30 years) who endorsed current cigarette smoking. Each participant rated 10 randomly selected messages (3,010 total message ratings; 24.3 ratings per message) on dimensions of content, design, perceived support for smoking urge reduction, and perceived support for quitting smoking. The sample size was selected based on recommendations in the existing

Distraction message	Acceptance message
	
Keep your mind occupied if you get an urge. Do a mental puzzle, make your next shopping list, read a book, write a poem...you might be surprised how well occupying your mind can get rid of urges.	Take a minute to notice what the urge feels like in your body without smoking a cigarette. Notice what is going on. See what happens. Does the urge pass?

Fig. 1 Intervention message examples for distraction and acceptance messages, including image and text content

literature, that suggested 25 evaluations per message to strike a balance between accuracy and efficiency [50].

Participants

We used Qualtrics panels, an online panel provider for market research, which is increasingly used in health and tobacco research [51, 52], to customize the study sample and facilitate the distribution as well as administration of the survey. The US-based panel included participants who had smoked 100 or more cigarettes in their lifetime, currently smoked cigarettes every day or some days (as non-daily smoking is common among young adults) [8, 9], were between 18 and 30 years old, and were currently trying to quit smoking or intended to quit in the next month. Quotas were set by the research team to ensure diversity of the sample, aiming for approximately 50% male and female sex, 20% Hispanic/Latino ethnicity, and 40% non-White race. These quotas were based on the US Census [53].

Qualtrics partners with over 20 online sample providers and most samples come from traditional, actively managed, double-opt-in market research panels. Potential participants are randomly selected from sites where users are likely to qualify. Participants' names, addresses and dates of birth are third-party validated (e.g., TrueSample, RelevantID, Verity, etc.). Invitations to participate in the survey are sent through email or on the survey platform with a generic message, a hyperlink to the survey, and the compensation offered. Invitations do not include specific study details to avoid self-selection bias. Since participants are recruited from a variety of sources, incentives vary (e.g., airline points, retail shopping points, cash, or gift cards), but the specific incentive a participant will receive is explicitly stated in the email before participants proceed to the survey link. Digital informed consent was obtained before participants completed the survey.

Qualtrics provides a series of data quality checks, including checking for duplicates, IP addresses, speeding (completion of survey in less than half of median time), bot detection, and straightlining (providing the same answers on 3 or more matrix tables in a survey), among others, before delivering final datasets.

Measures

Participants completed a survey assessing sociodemographic data, including age, sex, SGM status (Response options: Straight; Gay or lesbian; Bisexual; Pansexual; Queer; Transgender, transsexual, or gender non-conforming; Other – specify; recoded to binary variable SGM yes/SGM no), race/ethnicity, education levels, as well as current smoking behavior. For the purposes of reporting message ratings in tables, age was dichotomized into categories 18–25 vs. 26–30, while mixed

models included the continuous scale. Smoking behavior measures assessed daily smoking, time to first cigarette (within thirty minutes vs. after 30 min) [54], readiness to quit smoking (currently trying to quit; will quit in the next month), and motivation to quit smoking (five-point Likert scale). For the purposes of reporting message ratings in tables, this variable was dichotomized into categories 1–3 (not at all to moderately motivated – low motivation) vs. 4–5 (very or extremely motivated – high motivation), while mixed models included the continuous scale. Psychological flexibility was assessed using the Acceptance and Action Questionnaire – version two (AAQ-II) [55], with higher scores indicating more inflexibility. The AAQ-II measures psychological flexibility, which includes the ability to accept negative experiences, unwanted thoughts, and feelings while remaining engaged in present-moment living and pursuing actions aligned with personal values. Psychological flexibility may be a moderator of message rating of distraction and acceptance messages. Again, for the purposes of reporting message ratings in tables, this variable was dichotomized into categories 7–27 (flexible) vs. 28–49 (inflexible), while mixed models included the continuous scale. A cutoff of 28 or higher has been associated with mental distress in previous work [55].

Message ratings were assessed across four dimensions: (1) perceived quality of *content* (“How would you rate the content (that is, the words and meaning) of this message?”) and (2) perceived quality of *design* (“How would you rate the design (that is, how the message looks) of this message?”), both assessed on a Likert scale from 1 (Very poor) to 5 (Very good). Additional dimensions included: (3) perceived message support for *coping* with smoking urges (“How helpful would this message be to support you in coping with a smoking urge or craving?”), and (4) perceived message support for *quitting* smoking (“How helpful would this message be to support you in quitting or reducing smoking?”), both assessed on a Likert scale from 1 (Not at all helpful) to 5 (Extremely helpful).

Statistical analyses

In descriptive analyses, we examined message ratings by message category (acceptance vs. distraction) and participant characteristics [50]. We used mixed models to examine the relationship between the message category (CBT/distraction vs. ACT/acceptance) and participant level predictors (sociodemographic factors; daily smoking; time to first cigarette; readiness and motivation to quit; psychological flexibility) of message ratings. The model included both fixed and random effects to account for the hierarchical structure of the data, where message ratings were nested within participants. Fixed effects were specified for message category (CBT/distraction

vs. ACT/acceptance) and participant characteristics, estimating their average effects across the sample. To account for variability in baseline levels of message ratings across participants, we included a random intercept for participants. The initial mixed models included only main effects. Additional mixed models were estimated to investigate cross-level interaction effects between message category (CBT/distraction vs. ACT/acceptance) and participant level predictors. Separate models were estimated for each interaction effect and controlled for all other predictors.

Results

Sample description

Participant ($N=301$) mean age was 26.57 ($SD=2.94$) years and 54.8% of participants were male. 16.9% of participants identified as SGM (Table 1). A total of 48.5% identified as Non-Hispanic White, 21.9% as Non-Hispanic Black, and 22.6% as Hispanic. The majority (65.9%)

Table 1 Participant characteristics

	%; M (SD)
Age, mean (SD)	26.57 (2.94)
Age categories	
18–25	31.89
26–30	68.11
Sex	
Female	45.18
Male	54.82
Sexual or gender minority	
No	83.06
Yes	16.94
Race/Ethnicity	
Non-Hispanic White	48.50
Non-Hispanic Black	21.93
Hispanic	22.59
Other or multi-racial	6.98
Education	
High school or less	34.22
Some college	30.56
Bachelor's degree	23.26
Graduate degree	11.96
Daily smoking	62.46
Time to first cigarette (30 min or less)	60.80
Readiness to quit smoking	
Currently trying to quit	73.75
Will quit in the next month	26.25
Motivation to quit smoking, mean (SD)	3.82 (0.93)
Motivation to quit smoking categories	
Low (1–3)	32.56
High (4–5)	67.44
Psychological flexibility (AAQ-II), mean (SD)	23.44 (8.22)
Psychological flexibility (AAQ-II) categories	
Flexible (7–27)	65.45
Inflexible (28–49)	34.55

had at least some college education. Regarding smoking behavior, 62.5% of participants reported smoking daily, with 60.8% reporting smoking within thirty minutes of waking up, and 73.7% reported currently trying to quit smoking. The mean rating for motivation to quit smoking was 3.82 ($SD=0.93$, range 1–5). Psychological flexibility (AAQ-II) mean score was 23.44 ($SD=8.22$, range 7–39), with 34.6% reporting scores between 28 and 49, indicating moderate levels of inflexibility.

Average message ratings

Overall average message ratings were $M=3.90$ ($SD=1.03$) on content, $M=3.84$ ($SD=1.06$) on design, $M=3.61$ ($SD=1.25$) on support for coping, and $M=3.62$ ($SD=1.29$) on support for quitting smoking, with little to no differences between distraction and acceptance messages (Fig. 2).

Males, participants aged between 26 and 30 years of age, Hispanic participants and participants with at least a bachelor's degree gave relatively higher scores on average on most dimensions (Table 2). Participants reporting daily smoking, time to first cigarette at thirty minutes or less, high motivation to quit smoking, or more flexibility gave relatively higher scores across all dimensions compared to their counterparts. Conversely, participants identifying as a SGM or currently trying to quit in the next month averaged lower scores.

Mixed models

Mixed models predicting message scores based on message and participant characteristics

Mixed models examined main effects of message theme and participant characteristics (Table 3). Distraction messages were rated higher than acceptance messages on the dimension of perceived quality of design ($p<0.05$) and there were no other message category main effects on any of the rating dimensions. Male participants were significantly more likely to report favorable ratings on the dimensions of support for coping ($p<0.01$) and support for quitting ($p<0.01$). Participants identifying as SGM gave significantly lower ratings for message design ($p<0.05$), compared to their non-SGM counterparts. Participants with a graduate degree gave a higher rating on the dimensions of support for coping and support for quitting (both $p<0.05$), compared to those with high school education or less. High motivation to quit was associated with significantly more favorable message scores across all dimensions (all $p<0.01$). Participants who smoked daily rated messages as less helpful to support coping with smoking urges ($p<0.01$) and support for quitting smoking ($p<0.05$), compared to participants who smoked non-daily.




Fig. 2 Acceptance and distraction messages and mean rating scores for each message on dimensions content, design, support for coping with smoking urges, and support for quitting smoking (median; interquartile range (IQR) Q1 to Q3; whiskers are lower and upper adjacent values, respectively)

Mixed models predicting message scores with interactions between message and participant characteristics

Few significant interactions were observed between participant characteristics and ratings of distraction- vs. acceptance themed messages.

Participants who identified as SGM gave significantly lower ratings on distraction-themed messages on the dimension of perceived message support for coping with urges compared to their non-SGM counterparts ($b = -0.187$, $SE = 0.092$, $z = -2.02$, $p = 0.043$). Regarding racial and ethnic background, compared to Non-Hispanic White participants, Non-Hispanic Black ($b = -0.169$, $SE = 0.081$, $z = 2.09$, $p = 0.037$) and Hispanic participants ($b = 0.198$, $SE = 0.079$, $z = 2.50$, $p = 0.013$) gave significantly higher ratings for distraction-themed messages on the dimension of perceived quality of design. Compared to Non-Hispanic White participants, those of other racial or multiracial background gave more favorable ratings for distraction-themed messages on the dimension of perceived message support for coping ($b = 0.339$, $SE = 0.136$, $z = 2.48$, $p = 0.013$). The only significant interaction between participant education level and message theme was observed among participants with a bachelor's degree, who gave significantly lower ratings for distraction-themed messages on the dimensions of perceived quality of design ($b = -0.180$, $SE = 0.084$, $z = -2.15$, $p = 0.032$), perceived message support for coping ($b = -0.219$, $SE = 0.09$, $z = -2.39$, $p = 0.017$), and perceived message support for quitting ($b = -0.207$, $SE = 0.090$, $z = -2.28$, $p = 0.022$), compared to participants with high school education or less.

Discussion

The current study recruited an online panel of young adults who smoked and were interested in quitting smoking to evaluate CBT/distraction and ACT/acceptance-themed intervention messages on different dimensions related to content, design, and support for coping with smoking urges, and support for quitting smoking. Our results suggest that CBT/distraction and ACT/acceptance-themed messages were equally well received by our sample of young adults who smoke, reaching similar average ratings across all dimensions. These findings are in line with previous studies comparing the effectiveness of both therapy strategies [56, 57]. Overall, our findings are in accordance with previous work that have established the feasibility of both CBT and ACT smartphone-based interventions for smoking cessation [37–41, 43–45], and suggest that both strategies of smoking cessation intervention are similarly well received among young adults who smoke. Although messages of acceptance may be perceived as counterintuitive and different from messaging to avoid urges to smoke based on CBT approaches, our results suggest that the framing and presentation of the acceptance messaging were just as effective as CBT distraction messaging.

Results of the current study also highlighted sex differences in message ratings and suggest that male participants rated messages more positively than female participants on the dimensions of support for coping with smoking urges and for quitting smoking. These findings align with prior research indicating that biological and psychosocial factors influence smoking behaviors

Table 2 Average ratings for acceptance ($N = 1,575$ ratings) and distraction messages ($N = 1,434$ ratings) on dimensions content, design, coping, and quitting

Message theme	Content score (M, SD)		Design score (M, SD)		Coping score (M, SD)		Quitting score (M, SD)	
	Acceptance	Distraction	Acceptance	Distraction	Acceptance	Distraction	Acceptance	Distraction
Overall	3.90 (1.03)	3.90 (1.03)	3.81 (1.07)	3.88 (1.04)	3.60 (1.26)	3.62 (1.25)	3.60 (1.30)	3.63 (1.27)
Age								
18–25	3.84 (1.09)	3.87 (1.11)	3.70 (1.12)	3.81 (1.06)	3.53 (1.25)	3.55 (1.26)	3.50 (1.33)	3.57 (1.25)
26–30	3.93 (1.00)	3.92 (1.00)	3.86 (1.04)	3.91 (1.03)	3.63 (1.26)	3.65 (1.24)	3.65 (1.29)	3.66 (1.28)
Sex								
Female	3.79 (1.07)	3.74 (1.11)	3.67 (1.12)	3.71 (1.13)	3.35 (1.35)	3.31 (1.35)	3.35 (1.39)	3.30 (1.40)
Male	4.00 (1.00)	4.04 (0.95)	3.93 (1.01)	4.02 (0.94)	3.81 (1.14)	3.87 (1.09)	3.81 (1.19)	3.90 (1.08)
SGM								
No	3.95 (1.02)	3.98 (1.00)	3.87 (1.04)	3.96 (1.00)	3.66 (1.24)	3.72 (1.22)	3.68 (1.29)	3.74 (1.23)
Yes	3.69 (1.08)	3.52 (1.14)	3.51 (1.13)	3.48 (1.14)	3.33 (1.29)	3.10 (1.27)	3.24 (1.32)	3.09 (1.32)
Race/Ethnicity								
Non-Hispanic White	3.88 (0.99)	3.89 (1.00)	3.86 (1.03)	3.86 (1.03)	3.63 (1.19)	3.59 (1.21)	3.63 (1.23)	3.63 (1.21)
Non-Hispanic Black	3.82 (1.09)	3.78 (1.09)	3.71 (1.10)	3.78 (1.04)	3.48 (1.39)	3.46 (1.36)	3.45 (1.43)	3.43 (1.41)
Hispanic	4.08 (1.01)	4.00 (1.04)	3.85 (1.07)	4.00 (1.04)	3.81 (1.20)	3.84 (1.15)	3.83 (1.25)	3.87 (1.17)
Other or multi-racial	3.78 (1.17)	4.05 (1.05)	3.67 (1.14)	3.95 (1.07)	3.21 (1.36)	3.55 (1.37)	3.20 (1.37)	3.41 (1.43)
Education								
High school or less	3.82 (1.09)	3.80 (1.05)	3.69 (1.09)	3.81 (1.01)	3.38 (1.34)	3.44 (1.25)	3.35 (1.39)	3.43 (1.29)
Some college	3.76 (1.09)	3.72 (1.11)	3.63 (1.12)	3.69 (1.09)	3.44 (1.32)	3.43 (1.32)	3.42 (1.32)	3.40 (1.33)
Bachelor's degree	4.11 (0.88)	4.17 (0.90)	4.04 (0.98)	4.10 (0.97)	3.87 (1.07)	3.92 (1.14)	3.89 (1.16)	3.97 (1.13)
Graduate degree	4.11 (0.86)	4.13 (0.91)	4.17 (0.86)	4.14 (1.02)	4.12 (0.90)	4.03 (1.00)	4.20 (0.89)	4.15 (0.99)
Daily smoking								
Every day	3.98 (0.99)	3.93 (1.00)	3.90 (1.03)	3.92 (1.01)	3.75 (1.18)	3.70 (1.20)	3.75 (1.23)	3.72 (1.21)
Some days	3.78 (1.10)	3.85 (1.09)	3.66 (1.12)	3.82 (1.10)	3.35 (1.34)	3.48 (1.32)	3.36 (1.39)	3.49 (1.35)
Time to first cigarette								
After 30 min	3.86 (1.05)	3.86 (1.07)	3.74 (1.11)	3.82 (1.09)	3.46 (1.33)	3.52 (1.28)	3.46 (1.37)	3.49 (1.33)
Within 30 min	3.93 (1.02)	3.94 (1.01)	3.85 (1.04)	3.93 (1.00)	3.69 (1.21)	3.69 (1.22)	3.70 (1.25)	3.73 (1.22)
Readiness to quit smoking								
Currently trying to quit	3.86 (1.05)	3.84 (1.06)	3.78 (1.10)	3.84 (1.08)	3.54 (1.30)	3.54 (1.17)	3.54 (1.34)	3.54 (1.30)
Will quit in the next month	4.03 (0.96)	4.09 (0.94)	3.90 (0.96)	4.01 (0.92)	3.78 (1.12)	3.84 (1.14)	3.79 (1.17)	3.87 (1.13)
Motivation to quit smoking								
Low (1–3)	3.58 (1.09)	3.64 (1.07)	3.54 (1.13)	3.74 (1.06)	3.19 (1.29)	3.29 (1.27)	3.14 (1.35)	3.31 (1.29)
High (4–5)	4.06 (0.97)	4.03 (0.99)	3.94 (1.01)	3.95 (1.02)	3.80 (1.19)	3.78 (1.20)	3.82 (1.22)	3.79 (1.23)
Psychological flexibility (AAQ-II)								
Flexible (7–27)	3.88 (1.00)	3.90 (1.00)	3.80 (1.05)	3.87 (1.03)	3.58 (1.25)	3.60 (1.24)	3.58 (1.28)	3.63 (1.26)
Inflexible (28–49)	3.94 (1.09)	3.91 (1.09)	3.82 (1.11)	3.91 (1.07)	3.64 (1.27)	3.65 (1.26)	3.64 (1.35)	3.64 (1.29)

and cessation outcomes differently by sex and gender identity. While women are more likely to adhere to smoking interventions, they experience increased difficulty in cessation [58–60, 71]. Thus, our research provides further evidence for the need of appropriate interventions in the form of tailored message design and content to facilitate coping with smoking urges and support quitting among female young adults who smoke.

Compared to non-SGM participants, those identifying as SGM rated messages lower on the design dimension. Since intervention messages were not tailored to specific subgroups of young adults who smoke cigarettes, including those identifying as SGM, these findings may indicate the need for customized intervention messages

to culturally fit this group [61]. There is some evidence that culturally tailored smoking cessation interventions may work better for SGM young adults who smoke compared to non-tailored interventions [62]. However, other research has indicated that while SGM individuals who smoke may prefer smoking cessation interventions that are culturally tailored, untailored interventions show similar effects for SGM people who smoke compared to heterosexual individuals [63]. Future work is needed to determine the efficacy of digital smoking cessation interventions for SGM adults who smoke cigarettes and to investigate the need for cultural tailoring of intervention content.

Table 3 Mixed models with main effects predicting message rating scores on dimensions content, design, coping, and quitting by message and participant / rater characteristics

	Unstan- dardized coefficient (b)	Stan- dard error (SE)	T-value	p-value	Unstan- dardized coefficient (b)	Stan- dard error (SE)	T-value	p- value
	Content				Design			
Message category: distraction (ref. acceptance)	0.006	(0.030)	0.2	0.843	0.065	(0.031)	2.1	0.037
Age	0.020	(0.013)	1.5	0.128	0.026	(0.013)	1.9	0.054
Male sex (ref. female)	0.113	(0.081)	1.4	0.164	0.148	(0.082)	1.8	0.073
SGM (ref. non-minority)	-0.189	(0.106)	-1.8	0.074	-0.242*	(0.107)	-2.3	0.024
Non-Hispanic Black (ref. NH White)	-0.093	(0.097)	-1.0	0.341	-0.123	(0.099)	-1.3	0.210
Hispanic (ref. NH White)	0.110	(0.101)	1.1	0.274	0.063	(0.102)	0.6	0.540
Other or multi-racial (ref. NH White)	0.042	(0.151)	0.3	0.781	-0.004	(0.153)	0.0	0.978
Some college (ref. high school or less)	-0.173	(0.094)	-1.8	0.065	-0.179	(0.095)	-1.9	0.061
Bachelor's degree (ref. high school or less)	0.095	(0.104)	0.9	0.365	0.111	(0.106)	1.1	0.293
Graduate degree (ref. high school or less)	0.070	(0.131)	0.5	0.595	0.182	(0.133)	1.4	0.171
Daily smoking (ref. some days)	-0.153	(0.087)	-1.8	0.078	-0.148	(0.088)	-1.7	0.093
Time to first cigarette within 30 min (ref. after 30 min)	-0.074	(0.087)	-0.9	0.394	-0.066	(0.088)	-0.7	0.455
Readiness to quit smoking within the next month (ref. currently trying to quit)	0.082	(0.091)	0.9	0.367	0.036	(0.092)	0.4	0.694
Motivation to quit smoking	0.227***	(0.042)	5.5	0.000	0.151***	(0.042)	3.6	0.000
Psychological flexibility (AAQ-II)	0.000	(0.005)	0.0	0.971	0.000	(0.005)	0.1	0.925
	Coping				Quitting			
Message category: distraction (ref. acceptance)	0.024	(0.034)	0.7	0.485	0.038	(0.034)	1.1	0.268
Age	0.026	(0.017)	1.5	0.126	0.024	(0.018)	1.4	0.170
Male sex (ref. female)	0.330**	(0.103)	3.2	0.001	0.308**	(0.109)	2.8	0.005
SGM (ref. non-minority)	-0.165	(0.135)	-1.2	0.221	-0.211	(0.142)	-1.5	0.136
Non-Hispanic Black (ref. NH White)	-0.102	(0.124)	0.8	0.408	-0.147	(0.130)	-1.1	0.259
Hispanic (ref. NH White)	0.184	(0.128)	1.4	0.151	0.188	(0.135)	1.4	0.164
Other or multi-racial (ref. NH White)	-0.143	(0.192)	-0.7	0.458	-0.223	(0.203)	-1.1	0.272
Some college (ref. high school or less)	-0.110	(0.119)	-0.9	0.354	-0.117	(0.126)	-0.9	0.353
Bachelor's degree (ref. high school or less)	0.182	(0.133)	1.4	0.170	0.224	(0.140)	1.6	0.109
Graduate degree (ref. high school or less)	0.348*	(0.167)	2.1	0.037	0.456*	(0.176)	2.6	0.010
Daily smoking (ref. some days)	-0.295**	(0.110)	-2.7	0.007	-0.272*	(0.116)	-2.3	0.019
Time to first cigarette within 30 min (ref. after 30 min)	-0.079	(0.110)	0.7	0.476	-0.044	(0.116)	-0.4	0.704
Readiness to quit smoking within the next month (ref. currently trying to quit)	0.070	(0.115)	0.6	0.546	0.082	(0.122)	0.7	0.498
Motivation to quit smoking	0.223***	(0.053)	4.2	0.000	0.233***	(0.056)	4.2	0.000
Psychological flexibility (AAQ-II)	0.000	(0.006)	0.0	0.986	-0.002	(0.006)	-0.3	0.801

Note: NH = Non-Hispanic; AAQ-II = Acceptance and Action Questionnaire – version two

* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$

Regarding effects of education, participants with a graduate degree rated messages significantly more favorable on support for coping with smoking urges and support for quitting smoking compared to participants with a high school degree or less. These findings agree with well-known smoking disparities by education and socioeconomic status [64, 65]. Moreover, previous research has demonstrated that among people who smoke, those with less education engage less frequently with traditional smoking cessation aids, despite facing greater exposure to harmful products [3, 11, 12]. Based on these findings, the development and testing of smoking cessation interventions that are acceptable among people who

smoke with low educational attainment and meet their needs should be prioritized. Specifically, future interventions should make sure to engage this population to ensure messages are understandable and appealing, and to make any necessary adaptations.

Our findings showed that participants with a high motivation to quit gave higher message ratings on all dimensions. Conversely, those smoking daily rated messages as less helpful for coping with smoking urges and for quitting. Previous studies have already established the importance of motivation to quit for the effectiveness of smoking cessation interventions [66–68], which is also reflected in our results.

Future work may need to improve these smoking intervention messages to not only appeal to those who are already highly motivated to quit, but also to those who report indicators of high nicotine dependence or who may be less motivated to quit. For example, content based on motivational interviewing has been shown to generate good engagement among young adults who smoke and have low motivation to quit in social media delivered interventions [69, 70]. Overall, communication research finds that tailored messaging can support smoking cessation and health behavior change more broadly [71–74], and additional tailoring of the messages used in the current study may be needed to improve their efficacy and impact.

Limitations

The current study has several limitations. While the online panel we used in the current study allowed us to include various quotas to ensure our participants had diversity regarding sex and race/ethnicity, the study sample may not be representative for all young adults who smoke in the US. For example, our panel did not include a quota for education. Study participants were recruited from different sources with varying incentives, which may have impacted their motivation to diligently complete study procedures. The current study compared different categories of messages (e.g., distraction vs. acceptance) and did not investigate additional intervention message features (e.g., image content) that may have impacted message ratings. Despite evidence from anti-smoking campaigns suggesting that perceived effectiveness correlates with the actual effectiveness of intervention messages to some extent [75], it is unclear to what degree results of this intervention will be translated into message efficacy to support coping with smoking urges and quitting smoking. In addition, message ratings were aggregated which may have concealed within-individual or within-message variability or consistency. Lastly, messages only addressed cigarette smoking and additional modifications may be needed to address other nicotine and tobacco products popular among young people, which include electronic cigarettes [76] and oral nicotine pouches [77, 78].

Conclusions

Distraction and acceptance messages to support coping with smoking urges and smoking cessation received similar ratings among young adults who smoked cigarettes. Message revisions may be needed to increase appeal to women, SGM, those with lower education, and those less motivated to quit smoking. As a next step, intervention messages are tested for efficacy to reduce smoking urges and support smoking cessation in a micro-randomized trial to investigate their real-time impact on behavioral

outcomes. This trial is currently in progress and results of the current study helped to select a subset of intervention messages for inclusion. Finally, building on the findings of our current study, future research should also consider collecting and evaluating qualitative feedback and focus-group discussions of intervention messages to further refine and improve digital smoking cessation interventions for priority populations.

Abbreviations

AAQ-II	Acceptance and Action Questionnaire – version two
ACT	Acceptance and commitment therapy
CBT	Cognitive behavioral therapy
IQR	Interquartile range
M	Mean
NH	Non-Hispanic
OSF	Open Science Framework
SD	Standard Deviation
SES	Socioeconomic status
SGM	Sexual or gender minority
US	United States

Acknowledgements

The opinions and assertions expressed herein are those of the author(s) and do not reflect the official policy or position of the Uniformed Services University of the Health Sciences or the Department of Defense.

Author contributions

JH: Writing - Original Draft; Writing - Review & Editing. JD: Data Curation; Formal analysis; Investigation; Writing - Review & Editing. TR: Writing - Review & Editing. AL: Investigation; Writing - Review & Editing. JW: Investigation; Writing - Review & Editing. JH: Writing - Review & Editing. FN: Resources; Writing - Review & Editing. RV: Resources; Writing - Review & Editing. JB: Resources; Writing - Review & Editing. CL: Conceptualization; Methodology; Writing - Review & Editing. MM: Conceptualization; Methodology; Writing - Review & Editing. JT: Conceptualization; Formal analysis; Funding acquisition; Methodology; Supervision; Writing - Original Draft; Writing - Review & Editing.

Funding

This work was supported by the National Cancer Institute (NCI; R01 CA246590) and the National Institute on Drug Abuse (NIDA; T32 DA007292).

Data availability

No datasets were generated or analysed during the current study.

Declarations

Ethics approval and consent to participate

This study adhered to the Declaration of Helsinki and was approved by the Institutional Review Board of the Johns Hopkins Bloomberg School of Public Health. Digital informed consent was obtained before participants completed the survey.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Author details

¹Faculty of Medicine, Department of Medical Statistics, University of Göttingen, Göttingen, Germany

²Department of Mental Health, Johns Hopkins Bloomberg School of Public Health, Baltimore, USA

³Department of Medical and Clinical Psychology, School of Medicine, Uniformed Services University of the Health Sciences, Bethesda, USA

⁴Addiction Research Group, University of East Anglia, Norwich, UK

⁵Department of Implementation Science, Wake Forest University School of Medicine, Winston-Salem, USA

⁶Fred Hutchinson Cancer Center, Seattle, USA

⁷Department of Psychology, University of Washington, Seattle, USA

⁸Department of Health, Behavior and Society, Johns Hopkins Bloomberg School of Public Health, Baltimore, USA

⁹Sidney Kimmel Comprehensive Cancer Center at Johns Hopkins, Baltimore, USA

¹⁰Centre for Alcohol Policy Research, La Trobe University, Melbourne, Australia

Received: 24 December 2024 / Accepted: 30 April 2025

Published online: 20 May 2025

References

- Department of Health U, Services H. The Health Consequences of Smoking—50 Years of Progress: A Report of the Surgeon General. www.cdc.gov/tobacco.
- Threl J, Riehm KE, Cohen JE, Caleb Alexander G, Vernick JS, Mojtahab R. Tobacco control policies and smoking cessation treatment utilization: A moderated mediation analysis. *PLoS ONE*. 2021;16(8 August). <https://doi.org/10.1371/journal.pone.0241512>.
- Babb S, Malarcher A, Schauer G, Asman K, Jamal A. Quitting smoking among Adults — United States, 2000–2015. *MMWR Morb Mortal Wkly Rep*. 2017;65(52):1457–64. <https://doi.org/10.15585/mmwr.mm6552a1>.
- Ghadenik AE, Gauvin L, Frohlich KL. Smoking in young adults: A study of 4-Year smoking behavior patterns and residential presence of features facilitating smoking using data from the interdisciplinary study of inequalities in smoking cohort. *Nicotine Tob Res*. 2020;22(11):1997–2005. <https://doi.org/10.1093/ntr/ntaa035>.
- Barrington-Trimis JL, Braymiller JL, Unger JB, et al. Trends in the age of cigarette smoking initiation among young adults in the US from 2002 to 2018. *JAMA Netw Open*. 2020;3(10):E2019022. <https://doi.org/10.1001/jamanetworkopen.2020.19022>.
- Watkins SL, Threl J, Max W, Ling PM. Cold Turkey and hot vapes?? A National study of young adult cigarette cessation strategies. *Nicotine Tob Res*. 2020;22(5):638–46. <https://doi.org/10.1093/ntr/nty270>.
- Watkins SL, Threl J, Max W, Ling PM. Real-world effectiveness of smoking cessation strategies for young and older adults: findings from a nationally representative cohort. *Nicotine Tob Res*. 2020;22(9):1560–8. <https://doi.org/10.1093/ntr/ntz223>.
- Hooshmand S, Rodriguez EJ, Pérez-Stable EJ. *Nicotine Tob Res*. 2025;27(3):558–60. <https://doi.org/10.1093/ntr/ntae226>. How Much Longer Will We Ignore Nondaily Cigarette Smoking?.
- Doran N, Myers M, Luczak S, Trim R, Strong D, Tully L. Non-daily cigarette smoking: stability and transition to abstinence in young adults. *Nicotine Tob Res*. 2023;25(1):151–8. <https://doi.org/10.1093/ntr/ntac189>.
- Centers for Disease Control and Prevention. Tobacco-Related Disparities. Accessed January 27, 2020. <https://www.cdc.gov/tobacco/disparities/index.htm>
- Drope J, Liber AC, Cahn Z, et al. Who's still smoking? Disparities in adult cigarette smoking prevalence in the United States. *CA Cancer J Clin*. 2018;68(2):106–15. <https://doi.org/10.3322/caac.21444>.
- Hiscock R, Bauld L, Amos A, Fidler JA, Munafò M. Socioeconomic status and smoking: a review. *Ann NY Acad Sci*. 2012;1248(1):107–23. <https://doi.org/10.1111/j.1749-6632.2011.06202.x>.
- Nguyen-Grozavu FT, Pierce JP, Sakuma KLK, et al. Widening disparities in cigarette smoking by race/ethnicity across education level in the United States. *Prev Med (Baltim)*. 2020;139:106220. <https://doi.org/10.1016/j.jypmed.2020.106220>.
- Carlson S, Widome R, Fabian L, Luo X, Forster J. Barriers to quitting smoking among young adults: the role of socioeconomic status. *Am J Health Promotion*. 2018;32(2):294–300. <https://doi.org/10.1177/0890117117696350>.
- Stewart DW, Cano MÁ, Correa-Fernández V, et al. Lower health literacy predicts smoking relapse among racially/ethnically diverse smokers with low socioeconomic status. *BMC Public Health*. 2014;14(1). <https://doi.org/10.1186/1471-2458-14-716>.
- Villanti AC, Johnson AL, Rath JM. Beyond education and income: identifying novel socioeconomic correlates of cigarette use in U.S. Young adults. *Prev Med (Baltim)*. 2017;104:63–70. <https://doi.org/10.1016/j.jypmed.2017.06.019>.
- Harlow AF, Liu F, Young LE, et al. Sexual and gender identity disparities in nicotine and tobacco use susceptibility and prevalence: disaggregating emerging identities among adolescents from California, USA. *Nicotine Tob Res*. 2024;26(2):203–11. <https://doi.org/10.1093/ntr/ntad131>.
- Vogel EA, Humfleet GL, Meacham M, Prochaska JJ, Ramo DE. Sexual and gender minority young adults' smoking characteristics: assessing differences by sexual orientation and gender identity. *Addict Behav*. 2019;95:98–102. <https://doi.org/10.1016/j.addbeh.2019.03.005>.
- Jackson KM, Janssen T, Gabrielli J. Media/Marketing influences on adolescent and young adult substance abuse. *Curr Addict Rep*. 2018;5(2):146–57. <https://doi.org/10.1007/s40429-018-0199-6>.
- Cruz TB, McConnell R, Low BW, et al. Tobacco marketing and subsequent use of cigarettes, e-cigarettes, and Hookah in adolescents. *Nicotine Tob Res*. 2019;21(7):926–32. <https://doi.org/10.1093/ntr/nty107>.
- White JS, Salem MK, Toussaert S, et al. Developing a game (Inner Dragon) within a leading smartphone app for smoking cessation: design and feasibility evaluation study. *JMIR Serious Games*. 2023;11. <https://doi.org/10.2196/46602>.
- Bold KW, Garrison KA, DeLucia A, et al. Smartphone apps for smoking cessation: systematic framework for app review and analysis. *J Med Internet Res*. 2023;25. <https://doi.org/10.2196/45183>.
- Bricker JB, Watson NL, Mull KE, Sullivan BM, Heffner JL. Efficacy of smartphone applications for smoking cessation: A randomized clinical trial. *JAMA Intern Med*. 2020;180(11):1472–80. <https://doi.org/10.1001/jamainternmed.2020.4055>.
- Henley SJ, Thomas CC, Sharapova SR, et al. Vital signs: disparities in Tobacco-Related Cancer incidence and Mortality — United States, 2004–2013. *MMWR Morb Mortal Wkly Rep*. 2016;65(44):1212–8. <https://doi.org/10.15585/mmwr.mm6544a3>.
- Faverio M. Share of those 65 and older who are tech users has grown in the past decade. 2021. Accessed August 20, 2021. <https://www.pewresearch.org/short-reads/2022/01/13/share-of-those-65-and-older-who-are-tech-users-has-grown-in-the-past-decade/>
- Whittaker R, McRobbie H, Bullen C, Borland R, Rodgers A, Gu Y. Mobile phone-based interventions for smoking cessation. *Cochrane database of systematic reviews*. John Wiley & Sons, Ltd; 2012. <https://doi.org/10.1002/14651858.cd006611.pub3>.
- Abroms LC, Padmanabhan N, Thaweethai L, Phillips T. iPhone apps for smoking cessation: A content analysis. *Am J Prev Med*. 2011;40(3):279–85. <https://doi.org/10.1016/j.amepre.2010.10.032>.
- Abroms LC, Lee Westmaas J, Bontemps-Jones J, Ramani R, Mellerson J. A content analysis of popular smartphone apps for smoking cessation. *Am J Prev Med*. 2013;45(6):732–6. <https://doi.org/10.1016/j.amepre.2013.07.008>.
- Zhou X, Wei X, Cheng A, et al. Mobile Phone–Based interventions for smoking cessation among young people: systematic review and Meta-Analysis. *JMIR Mhealth Uhealth*. 2023;11:e48253. <https://doi.org/10.2196/48253>.
- Micalizzi L, Mattingly DT, Hart JL, Jensen JK, Mahabee-Gittens EM, Garrison KA. Smartphone apps targeting youth tobacco use prevention and cessation: an assessment of credibility and quality. *Curr Addict Rep*. 2023;10(4):649–63. <https://doi.org/10.1007/s40429-023-00524-0>.
- Chen J, Chu J, Marsh S, Shi T, Bullen C. Smartphone App-Based interventions to support smoking cessation in smokers with mental health conditions: A systematic review. *Psych*. 2023;5(4):1077–100. <https://doi.org/10.3390/psych5040072>.
- Rupp A, Rietzler S, Di Lellis MA, Weiland T, Tschirner C, Kreuter M. Digital smoking cessation with a comprehensive Guideline-Based App—Results of a nationwide, multicentric, parallel, randomized controlled trial in Germany. *Nicotine Tob Res*. 2024;26(7):895–902. <https://doi.org/10.1093/ntr/ntae009>.
- Force USPST. Interventions for tobacco smoking cessation in adults, including pregnant persons: US preventive services task force recommendation statement. *JAMA*. 2021;325(3):265–79. <https://doi.org/10.1001/jama.2020.25019>.
- Perkins KA, Conklin CA, Levine MD. Cognitive-Behavioral therapy for smoking cessation. *Routledge*; 2008. <https://doi.org/10.4324/9780203844533>.
- Kwan YK, Lau Y, Ang WW, Lau ST. Immediate, Short-term, Medium-term, and Long-term effects of acceptance and commitment therapy for smoking cessation: A systematic review and Meta-Analysis. *Nicotine Tob Res*. 2024;26(1):12–22. <https://doi.org/10.1093/ntr/ntad145>.
- McCallion EA, Zvolensky MJ. Acceptance and commitment therapy (ACT) for smoking cessation: a synthesis. *Curr Opin Psychol*. 2015;2:47–51. <https://doi.org/10.1016/j.copsyc.2015.02.005>.
- Tudor-Sfetcu C, Rabee R, Najim M, et al. Evaluation of two mobile health apps in the context of smoking cessation: qualitative study of cognitive behavioral

- therapy (CBT) versus NON-CBT-based digital solutions. *JMIR Mhealth Uhealth*. 2018;6(4). <https://doi.org/10.2196/mhealth.9405>.
38. Hofmann SG, Asnaani A, Vonk IJ, Sawyer AT, Fang A. The efficacy of cognitive behavioral therapy: A review of meta-analyses. *Cognit Ther Res*. 2012;36(5):427–40. <https://doi.org/10.1007/s10608-012-9476-1>.
39. Magill M, Ray LA. Cognitive-Behavioral Treatment With Adult Alcohol and Illicit Drug Users: A Meta-Analysis of Randomized Controlled Trials*. 2009.
40. Watts S, Mackenzie A, Thomas C, et al. CBT for depression: A pilot RCT comparing mobile phone vs. computer. *BMC Psychiatry*. 2013;13. <https://doi.org/10.1186/1471-244X-13-49>.
41. Ploderer B, Smith W, Pearce J, Borland R. A mobile app offering distractions and tips to Cope with cigarette craving: A qualitative study. *JMIR Mhealth Uhealth*. 2014;2(2). <https://doi.org/10.2196/mhealth.3209>.
42. Zhang CQ, Leeming E, Smith P, Chung PK, Hagger MS, Hayes SC. Acceptance and commitment therapy for health behavior change: A contextually-driven approach. *Front Psychol*. 2018;8(JAN). <https://doi.org/10.3389/fpsyg.2017.02350>.
43. Singh S, Starkey NJ, Sargisson RJ. Using SmartQuit*, an acceptance and commitment therapy smartphone application, to reduce smoking intake. *Digit Health*. 2017;3:205520761772953. <https://doi.org/10.1177/2055207617729535>.
44. O'Connor M, Whelan R, Bricker J, Mchugh L. Randomized controlled trial of a smartphone application as an adjunct to acceptance and commitment therapy for smoking cessation.; 2019. www.sciencedirect.com/locate/bt
45. Bricker JB, Mull KE, Kientz JA, et al. Randomized, controlled pilot trial of a smartphone app for smoking cessation using acceptance and commitment therapy. *Drug Alcohol Depend*. 2014;143(1):87–94. <https://doi.org/10.1016/j.drugalcdep.2014.07.006>.
46. Vilardaga R, Rizo J, Palenski PE, Mannelli P, Oliver JA, McClernon FJ. Pilot randomized controlled trial of a novel smoking cessation app designed for individuals with co-occurring tobacco use disorder and serious mental illness. *Nicotine Tob Res*. 2020;22(9):1533–42. <https://doi.org/10.1093/ntr/ntz202>.
47. Naughton F, Hope A, Siegle-Brown C et al. A smoking cessation smartphone app that delivers real-time 'context aware' behavioural support: the quit sense feasibility RCT. *Public Health Res Published Online April 2024*:1–99. <https://doi.org/10.3310/kqyt5412>
48. Unsplash. Accessed December 31, 2021. <https://unsplash.comhttps://unsplash.com>
49. Pexels. Accessed December 31, 2021. <https://www.pexels.com>
50. Kim M, Cappella JN. An efficient message evaluation protocol: two empirical analyses on positional effects and optimal sample size. *J Health Commun*. 2019;24(10):761–9. <https://doi.org/10.1080/10810730.2019.1668090>.
51. Olson CK, Sherwood N, Berkane K, Gilligan K, McKinney WJ. Perceptions, intentions, and actual use of a consumer nicotine gum. *Harm Reduct J*. 2023;20(1). <https://doi.org/10.1186/s12954-023-00864-0>.
52. Patev AJ, Maldonado GT, Sawyer-Mays AN, Cobb CO, Breland A. Smoking discrimination moderates the relationship between smoking stigma and cigarettes per day among sexual and gender minority individuals. *Subst Use Misuse*:1–8. <https://doi.org/10.1080/10826084.2024.2422952>
53. United. States Census.
54. Baker T, Piper M, McCarthy D, et al. Time to first cigarette in the morning as an index of ability to quit smoking: implications for nicotine dependence. *Nicotine Tob Res*. 2007;9(SUPPL 4):555–70. <https://doi.org/10.1080/14622200701673480>.
55. Bond FW, Hayes SC, Baer RA, et al. Preliminary psychometric properties of the acceptance and action Questionnaire-II: A revised measure of psychological inflexibility and experiential avoidance. *Behav Ther*. 2011;42(4):676–88. <https://doi.org/10.1016/j.beth.2011.03.007>.
56. Bricker JB, Bush T, Zbikowski SM, Mercer LD, Heffner JL. Randomized trial of telephone-delivered acceptance and commitment therapy versus cognitive behavioral therapy for smoking cessation: A pilot study. *Nicotine Tob Res*. 2014;16(11):1446–54. <https://doi.org/10.1093/ntr/ntu102>.
57. McClure JB, Bricker J, Mull K, Heffner JL. Comparative effectiveness of group-delivered acceptance and commitment therapy versus cognitive behavioral therapy for smoking cessation: A randomized controlled trial. *Nicotine Tob Res*. 2020;22(3):354–62. <https://doi.org/10.1093/ntr/nty268>.
58. Smith PH, Bessette AJ, Weinberger AH, Sheffer CE, McKee SA. Sex/gender differences in smoking cessation: A review. *Prev Med (Baltim)*. 2016;92:135–40. <https://doi.org/10.1016/j.ypmed.2016.07.013>.
59. O'Connell N, Burke E, Dobbie F, et al. The effectiveness of smoking cessation interventions for socio-economically disadvantaged women: a systematic review and meta-analysis. *Syst Rev*. 2022;11(1). <https://doi.org/10.1186/s13643-022-01922-7>.
60. Bauld L, Ferguson J, McEwen A, Hiscock R. Evaluation of a drop-in rolling-group model of support to stop smoking. *Addiction*. 2012;107(9):1687–95. <https://doi.org/10.1111/j.1360-0443.2012.03861.x>.
61. Leinberger-Jabari A, Golob MM, Lindson N, Hartmann-Boyce J. Effectiveness of culturally tailoring smoking cessation interventions for reducing or quitting combustible tobacco: A systematic review and meta-analyses. *Addiction*. 2024;119(4):629–48. <https://doi.org/10.1111/add.16400>.
62. Vogel EA, Ramo DE, Meacham MC, Prochaska JJ, Delucchi KL, Humfleet GL. The put it out project (POP) Facebook intervention for young sexual and gender minority smokers: outcomes of a pilot, randomized, controlled trial. *Nicotine Tob Res*. 2020;22(9):1614–21. <https://doi.org/10.1093/ntr/ntz184>.
63. Lee JGL, Matthews AK, McCullen CA, Melvin CL. Promotion of tobacco use cessation for lesbian, gay, bisexual, and transgender people: A systematic review. *Am J Prev Med*. 2014;47(6):823–31. <https://doi.org/10.1016/j.amepre.2014.07.051>.
64. Maralani V. Understanding the links between education and smoking. *Soc Sci Res*. 2014;48:20–34. <https://doi.org/10.1016/j.ssresearch.2014.05.007>.
65. Kock L, Brown J, Hiscock R, Tattan-Birch H, Smith C, Shahab L. Individual-level behavioural smoking cessation interventions tailored for disadvantaged socioeconomic position: a systematic review and meta-regression. *Lancet Public Health*. 2019;4(12):e628–44. [https://doi.org/10.1016/S2468-2667\(19\)30220-8](https://doi.org/10.1016/S2468-2667(19)30220-8).
66. Rajani NB, Mastellos N, Filippidis FT. Self-Efficacy and motivation to quit of smokers seeking to quit: quantitative assessment of smoking cessation mobile apps. *JMIR Mhealth Uhealth*. 2021;9(4):e25030. <https://doi.org/10.2196/25030>.
67. Roberts NJ, Kerr SM, Smith SMS. Behavioral interventions associated with smoking cessation in the treatment of tobacco use. *Health Serv Insights*. 2013;6. <https://doi.org/10.4137/HSI.S11092>.
68. Elshatarat RA, Yacoub MI, Khraim FM, Saleh ZT, Afaneh TR. Self-efficacy in treating tobacco use: A review article. *Proceedings of Singapore Healthcare*. 2016;25(4):243–248. <https://doi.org/10.1177/2010105816667137>
69. Thrul J, Meacham MC, Tice C, Kelly O, Ramo DE. Live counselor contact in a Facebook intervention predicts smoking cessation outcomes. *Psychol Addict Behav*. 2020;34(2):360–9. <https://doi.org/10.1037/adb0000541>.
70. Thrul J, Klein AB, Ramo DE. Smoking cessation intervention on Facebook: which content generates the best engagement? *J Med Internet Res*. 2015;17(11). <https://doi.org/10.2196/jmir.4575>.
71. Altendorf Ma, Hoving C, Van Weert JCM, Smit ES. Effectiveness of message frame-tailoring in a web-based smoking cessation program: randomized controlled trial. *J Med Internet Res*. 2020;22(4). <https://doi.org/10.2196/17251>.
72. Noar SM, Benac CN, Harris MS. Does tailoring matter? Meta-analytic review of tailored print health behavior change interventions. *Psychol Bull*. 2007;133(4):673–93. <https://doi.org/10.1037/0033-2909.133.4.673>.
73. Tripp HL, Strickland JC, Mercincavage M, Audrain-McGovern J, Donny EC, Strasser AA. Tailored cigarette warning messages: how individualized loss aversion and delay discounting rates can influence perceived message effectiveness. *Int J Environ Res Public Health*. 2021;18(19). <https://doi.org/10.3390/ijerph181910492>.
74. Tversky A, Kahneman D. The Framing of Decisions and the Psychology of Choice.
75. Noar SM, Barker J, Bell T, Yzer M. Does perceived message effectiveness predict the actual effectiveness of tobacco education messages? A systematic review and Meta-Analysis. *Health Commun*. 2020;35(2):148–57. <https://doi.org/10.1080/10410236.2018.1547675>.
76. Sanford BT, Brownstein NC, Baker NL, et al. Shift from smoking cigarettes to vaping nicotine in young adults. *JAMA Intern Med*. 2024;184(11):106–8. <https://doi.org/10.1001/jamainternmed.2023.5239>.
77. Dowd AN, Thrul J, Czaplicki L, Kennedy RD, Moran MB, Spindle TR. A Cross-Sectional survey on oral nicotine pouches: characterizing Use-Motives, topography, dependence levels, and adverse events. *Nicotine Tob Res*. 2024;26(2):245–9. <https://doi.org/10.1093/ntr/ntad179>.

78. Zamarripa CA, Dowd AN, Elder HJ et al. A comprehensive review on oral nicotine pouches: available scientific evidence and future research needs. *Exp Clin Psychopharmacol* Published Online Dec 2, 2024.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.